



HAL
open science

Complete partial metric spaces have partially metrizable computational models-CMMSE2010

Oscar Valero, Salvador Romaguera, Pedro Tirado

► **To cite this version:**

Oscar Valero, Salvador Romaguera, Pedro Tirado. Complete partial metric spaces have partially metrizable computational models-CMMSE2010. *International Journal of Computer Mathematics*, 2011, pp.1. 10.1080/00207160.2011.559229 . hal-00710044


HAL Id: hal-00710044

<https://hal.science/hal-00710044>

Submitted on 20 Jun 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Complete partial metric spaces have partially metrizable computational models-CMMSE2010

Journal:	<i>International Journal of Computer Mathematics</i>
Manuscript ID:	GCOM-2010-0802-A
Manuscript Type:	Original Article
Date Submitted by the Author:	13-Oct-2010
Complete List of Authors:	VALERO, OSCAR; Universidad de las Islas Baleares, Departamento de Ciencias Matematicas e Informatica Romaguera, Salvador; Instituto Universitario de Matemática Pura y Aplicada Tirado, Pedro; Instituto Universitario de Matemática Pura y Aplicada
Keywords:	computational model, complete partial metric, domain, formal ball, The Scott topology

SCHOLARONE™
Manuscripts

RESEARCH ARTICLE

Complete partial metric spaces have partially metrizable
computational modelsS. Romaguera^{a*}, P. Tirado^{b*} and O. Valero^{c*}^asromague@mat.upv.es, Instituto Universitario de Matemática Pura y Aplicada,
Universidad Politécnica de Valencia, 46071 Valencia, Spain;^bpedtipe@mat.upv.es, Instituto Universitario de Matemática Pura y Aplicada,
Universidad Politécnica de Valencia, 46071 Valencia, Spain^co.valero@uib.es, Departamento de Ciencias Matemáticas e Informática, Universidad de
las Islas Baleares, Palma de Mallorca, Spain

(Received 00 Month 200x; in final form 00 Month 200x)

We show that the domain of formal balls of a complete partial metric space (X, p) can be endowed with a complete partial metric that extends p and induces the Scott topology. This result, that generalizes well-known constructions of Edalat and Heckmann (Theoret. Comput. Sci. 1998) and Heckmann (Appl. Cat. Struct. 1998) for metric spaces and improves a recent result of Romaguera and Valero (Math. Struct. Comput. Sci. 2009), motivates a notion of a partially metrizable computational model which allows us to characterize those topological spaces that admit a compatible complete partial metric via this model.

Keywords: computational model, complete partial metric, domain, formal ball, the Scott topology

AMS Subject Classification: MCS Codes: 68Q01, 68Q05, 54E50, 06A06, 06B35.

1. Introduction

Motivated by the fact that metric spaces and domain theory constitute fundamental mathematical tools in computer science, several authors have investigated the problem of obtaining links between them and constructing, with this approach, suitable models in the theory of computation [2, 3, 5, 7, 8, 11–13, 15, 19, 23, etc]. In particular, Edalat and Heckmann [2] presented a very nice and explicit construction of a computational model for (complete) metric spaces by means of the notion of a formal ball. They proved, among other results, that the poset of formal balls of a metric space (X, d) is a domain if and only if (X, d) is complete and that this poset is an ω -domain if and only if (X, d) is complete and separable (previously, Lawson characterized in [11] separable completely metrizable spaces in terms of ω -domains). Later on, Heckmann [7] constructed a partial metric on the (continuous) poset of formal balls of a metric space (X, d) which extends the metric d and induces the Scott topology (similar results were obtained by Rutten [19] with a different approach). Heckmann's construction is very interesting because the use of a distance function on the formal balls provides a “quantitative” computational model, which suggested, by one hand, the study of computational models from a

* The authors acknowledge the support of the Spanish Ministry of Science and Innovation, under grant MTM2009-12872-C02-01

quantitative point of view and, on the other hand, the study of partial metrics and related structures by means of domain theory (see [1, 9, 14, 16–18, 20–23, etc] for recent contributions in these directions).

Here, and improving a recent construction by Romaguera and Valero [17], we shall prove that the domain of formal balls of a complete partial metric space (X, p) can be endowed with a complete partial metric that, among other things, extends the partial metric p and induces the Scott topology. Motivated by this result we shall define the notion of a partially metrizable computational model and then we shall show that the topological spaces having a partially metrizable computational model are exactly those admitting a compatible complete partial metric.

2. Background

The letters \mathbb{N} , \mathbb{R} and \mathbb{R}^+ will denote the set of positive integer numbers, the set of real numbers and the set of non-negative real numbers, respectively.

Our basic reference for domain theory is [6], for general topology is [4] and for quasi-metric spaces it is [10].

Let us recall that a partially ordered set, or poset for short, is a set L equipped with a partial order \sqsubseteq ; it will be denoted by (L, \sqsubseteq) or simply by L if no confusion arises.

A subset D of a poset L is directed provided that it is nonempty and any pair of elements of D has an upper bound in D . The least upper bound of a subset D of L is denoted by $\bigsqcup D$ if it exists.

A poset L is said to be directed complete, and is called a dcpo, if every directed subset of L has a least upper bound.

An element x of L is said to be maximal if the condition $x \sqsubseteq y$ implies $x = y$. The set of all maximal points of L will be denoted by $\text{Max}((L, \sqsubseteq))$ or simply by $\text{Max}(L)$ if no confusion arises.

Let L be a poset and $x, y \in L$; we say that x is way below y , in symbols $x \ll y$, if for each directed subset D of L for which $\bigsqcup D$ exists, the relation $y \sqsubseteq \bigsqcup D$ implies the existence of some $z \in D$ with $x \sqsubseteq z$.

A poset L is called continuous if for each $x \in L$, the set $\Downarrow x = \{u \in L : u \ll x\}$ is directed and $x = \bigsqcup(\Downarrow x)$.

A continuous poset which is also a dcpo is called a continuous domain or, simply, a domain.

A subset B of a poset L is a basis for L if for each $x \in L$, the set $\Downarrow x_B = \{u \in B : u \ll x\}$ is directed and $x = \bigsqcup(\Downarrow x_B)$.

Recall that a poset has a basis if and only if it is continuous. Therefore, a dcpo has a basis if and only if it is a domain.

A dcpo having a countable basis is said to be an ω -continuous domain, or simply an ω -domain if no confusion arises.

The Scott topology $\sigma(L)$ of a poset (L, \sqsubseteq) is constructed as follows: A subset U of L is open with respect to $\sigma(L)$ provided that: (i) $U = \uparrow U$, where $\uparrow U = \{y \in L : x \sqsubseteq y \text{ for some } x \in U\}$; (ii) for each directed subset D of L with $\bigsqcup D \in U$, it follows that $D \cap U \neq \emptyset$.

If D is a subset of L , we denote by $\sigma(L)|_D$ the restriction of $\sigma(L)$ to D .

If (L, \sqsubseteq) is a continuous poset, then the sets $\uparrow x$, $x \in L$, form an open base for $\sigma(L)$, where $\uparrow x = \{y \in L : x \ll y\}$ (see [6, Proposition II-1.10]).

The lower (or weak) topology of a poset L is the one that has as a subbase the collection of sets of the form $L \setminus \uparrow x$, $x \in L$, and is denoted by $\omega(L)$. Let us recall

1 that the supremum topology of $\sigma(L)$ and $\omega(L)$ is the Lawson topology of L , which
 2 is denoted by $\lambda(L)$. If D is a subset of L , we denote by $\lambda(L)|_D$ the restriction of
 3 $\lambda(L)$ to D .

4
 5 If (X, τ) is a T_0 topological space, then the binary relation \leq_τ defined on X by
 6 $x \leq_\tau y \Leftrightarrow x \in \text{cl}\{y\}$, is a partial order on X called the specialization order (see,
 7 for instance, p. 42 of [6]).

8 Following the modern terminology, by a quasi-metric on a set X we mean a
 9 function $d : X \times X \rightarrow \mathbb{R}^+$ such that for all $x, y, z \in X$: (i) $x = y \Leftrightarrow d(x, y) =$
 10 $d(y, x) = 0$; (ii) $d(x, z) \leq d(x, y) + d(y, z)$.

11 A quasi-metric space is a pair (X, d) such that X is a set and d is a quasi-metric
 12 on X .

13 Each quasi-metric d on X induces a T_0 topology τ_d on X which has as a base the
 14 family of open balls $\{B_d(x, r) : x \in X, r > 0\}$, where $B_d(x, \varepsilon) = \{y \in X : d(x, y) <$
 15 $\varepsilon\}$ for all $x \in X$ and $\varepsilon > 0$.

16 Note that if (X, d) is a quasi-metric space, then the binary relation \leq_d defined
 17 on X by $x \leq_d y \Leftrightarrow d(x, y) = 0$, is a partial order on X , which coincides with the
 18 specialization order of (X, τ_d) .

19 Given a quasi-metric d on X , then the function d^{-1} defined by $d^{-1}(x, y) = d(y, x)$,
 20 is also a quasi-metric on X , called the conjugate of d , and the function d^s defined
 21 by $d^s(x, y) = \max\{d(x, y), d^{-1}(x, y)\}$ is a metric on X .

22 On the other hand, the notion of a partial metric space, and its equivalent
 23 weightable quasi-metric space, was introduced by Matthews in [13] as a part of
 24 the study of denotational semantics of dataflow networks.

25 Let us recall that a partial metric on a set X is a function $p : X \times X \rightarrow \mathbb{R}^+$
 26 such that for all $x, y, z \in X$: (i) $x = y \Leftrightarrow p(x, x) = p(x, y) = p(y, y)$; (ii) $p(x, x) \leq$
 27 $p(x, y)$; (iii) $p(x, y) = p(y, x)$; (iv) $p(x, z) \leq p(x, y) + p(y, z) - p(y, y)$.

28 If p is a partial metric on X we denote by $\ker p$ the subset of X consisting of all
 29 points $x \in X$ such that $p(x, x) = 0$. Obviously, every metric d on X is a partial
 30 metric with $\ker d = X$.

31 A partial metric space is a pair (X, p) such that X is a set and p is a partial
 32 metric on X .

33 Each partial metric p on X induces a T_0 -topology τ_p on X which has as a base
 34 the family of open p -balls $\{B_p(x, \varepsilon) : x \in X, \varepsilon > 0\}$, where $B_p(x, \varepsilon) = \{y \in X :$
 35 $p(x, y) < \varepsilon + p(x, x)\}$ for all $x \in X$ and $\varepsilon > 0$.

36 We say that a topological space (X, τ) admits a compatible partial metric if there
 37 is a partial metric p on X such that $\tau_p = \tau$.

38 Matthews observed in [13] that each partial metric p on a set X induces a quasi-
 39 metric d_p on X given by $d_p(x, y) = p(x, y) - p(x, x)$ for all $x, y \in X$. Moreover
 40 $\tau_p = \tau_{d_p}$.

41 If (X, p) is a partial metric space, then the binary relation \leq_p on X given by
 42 $x \leq_p y \Leftrightarrow p(x, y) = p(x, x)$, is a partial order on X . Hence (X, \leq_p) is a poset
 43 ([7, 13]). Note that in this case one has $\leq_p = \leq_{d_p}$.

44 In Definition 5.3 of [13], Matthews also introduced the notion of a complete
 45 partial metric and stated, with a slight different terminology, that a partial metric
 46 p on a set X is complete if and only if the metric $(d_p)^s$ is complete on X .

47 We will say that a partial metric space (X, p) is complete if p is a complete partial
 48 metric on X .

3. Partially metrizable computational models

Recall [2] that if (X, d) is a metric space, then $\mathbf{B}X = X \times \mathbb{R}^+$ is said to be the set of formal balls of (X, d) , and the relation \sqsubseteq_d defined on $\mathbf{B}X$ by $(x, r) \sqsubseteq_d (y, s) \Leftrightarrow d(x, y) \leq r - s$, is a partial order on X . The pair $(\mathbf{B}X, \sqsubseteq_d)$ is called the poset of formal balls of (X, d) .

Then, Edalat and Heckmann proved that for any metric space (X, d) , the poset of formal balls is continuous, and the mapping $i : X \rightarrow \mathbf{B}X$, given by $i(x) = (x, 0)$, is a homeomorphism between (X, τ_d) and $(\text{Max}(\mathbf{B}X), \sigma(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)})$, and $\sigma(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)} = \lambda(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)}$.

As we mentioned above, Heckmann constructed in [7] a partial metric on the set $\mathbf{B}X$ of formal balls of a metric space (X, d) , that extends the metric d and such that both the partial order \sqsubseteq_d and the Scott topology are induced by it.

The partial metric P on $\mathbf{B}X$, given below, is a slight modification of Heckmann's original construction (see Section 4 of [17]):

$$P((x, r), (y, s)) = \max\{d(x, y), |r - s|\} + r + s,$$

for all $(x, r), (y, s) \in \mathbf{B}X$.

In Theorem 1 below we extend Heckmann's construction to (complete) partial metric spaces which also provides an improvement of Theorem 4.1 of [17].

To this end, we first recall some pertinent concepts and facts.

The poset of formal balls of a quasi-metric space (X, d) is the poset $(\mathbf{B}X, \sqsubseteq_d)$, where both $\mathbf{B}X$ and the partial order \sqsubseteq_d are defined exactly as in the metric case (see [1, 17], where quasi-metric and partial metric versions of several results of Edalat and Heckmann were obtained. In particular, it was proved in Theorem 3.1 of [17] that a partial metric space (X, p) is complete if and only if $(\mathbf{B}X, \sqsubseteq_{d_p})$ is a domain).

According to Waszkiewicz [22], a weak partial metric on a set X is a function $p : X \times X \rightarrow \mathbb{R}^+$ such that for all $x, y, z \in X$: (i) $x = y \Leftrightarrow p(x, x) = p(x, y) = p(y, y)$; (ii) $p(x, y) = p(y, x)$; (iii) $p(x, z) \leq p(x, y) + p(y, z) - p(y, y)$.

Heckmann [7] showed that every weak partial metric p on a set X induces a partial metric p' on X given by $p'(x, y) = \max\{p(x, y), p(x, x), p(y, y)\}$ for all $x, y \in X$.

Now let (X, p) be a partial metric space. Construct a function $P : \mathbf{B}X \times \mathbf{B}X \rightarrow \mathbb{R}^+$ by

$$P((x, r), (y, s)) = p(x, y) + r + s,$$

for all $(x, r), (y, s) \in \mathbf{B}X$.

The proof of the following result is straightforward (compare with the first paragraph of 3.2 in [7]).

Proposition 1. *For each partial metric space (X, p) , the function P constructed above is a weak partial metric on $\mathbf{B}X$.*

Therefore, the function $P' : \mathbf{B}X \times \mathbf{B}X \rightarrow \mathbb{R}^+$ defined by

$$P'((x, r), (y, s)) = \max\{P((x, r), (y, s)), P((x, r), (x, r)), P((y, s), (y, s))\},$$

for all $(x, r), (y, s) \in \mathbf{B}X$, is a partial metric on $\mathbf{B}X$.

Note that, in fact,

$$P'((x, r), (y, s)) = \max\{p(x, y) + r + s, p(x, x) + 2r, p(y, y) + 2s\},$$

for all $(x, r), (y, s) \in \mathbf{B}X$.

Now we are in position of establishing our main result.

Theorem 1. *Let (X, p) be partial metric space and let P' be the partial metric on $\mathbf{B}X$ defined above. Then*

(a) *The mapping $i : X \rightarrow \mathbf{B}X$, given by $i(x) = (x, 0)$, is an isometry between (X, p) and a $\tau_{(d_{P'})^{-1}}$ -closed subspace of $(\mathbf{B}X, P')$.*

(b) *$\leq_{P'} = \sqsubseteq_{d_p}$, and $i(\text{Max}(X, \leq_p)) = \text{Max}(\mathbf{B}X)$.*

(c) *If (X, p) is complete, then $\tau_{P'} = \sigma(\mathbf{B}X)$.*

(d) *If (X, p) is complete and $\ker p = \text{Max}(X, \leq_p)$, then $\sigma(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)} = \lambda(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)}$.*

(e) *$(\mathbf{B}X, P')$ is complete if and only if (X, p) is complete.*

Proof. (a) Let $x, y \in X$. Then

$$P'(i(x), i(y)) = P'((x, 0), (y, 0)) = \max\{p(x, y), p(x, x), p(y, y)\} = p(x, y).$$

So i is an isometry from (X, p) into $(\mathbf{B}X, P')$.

In order to show that $i(X)$ is $\tau_{(d_{P'})^{-1}}$ -closed in $(\mathbf{B}X, P')$, let $(y, s) \in \mathbf{B}X$ such that there is a sequence $(x_n)_{n \in \mathbb{N}}$ in X with $d_{P'}((x_n, 0), (y, s)) < 1/n$ for all $n \in \mathbb{N}$. It follows that $p(x_n, y) + s - p(x_n, x_n) < 1/n$, so $s < p(x_n, x_n) - p(x_n, y) + 1/n \leq 1/n$ for all $n \in \mathbb{N}$, and hence $s = 0$. We conclude that $(y, s) \in i(X)$ and consequently $i(X)$ is $\tau_{(d_{P'})^{-1}}$ -closed in $(\mathbf{B}X, P')$.

(b) For each $(x, r), (y, s) \in \mathbf{B}X$ we have

$$\begin{aligned} (x, r) \leq_{P'} (y, s) &\Leftrightarrow P'((x, r), (y, s)) = P'((x, r), (x, r)) \\ &\Leftrightarrow p(x, y) + r + s \leq p(x, x) + 2r \\ &\Leftrightarrow p(x, y) - p(x, x) \leq r - s \Leftrightarrow (x, r) \sqsubseteq_{d_p} (y, s). \end{aligned}$$

So $\leq_{P'} = \sqsubseteq_{d_p}$.

On the other hand, it is clear (compare Theorem 5.1 of [17]) that $\text{Max}(\mathbf{B}X) = \{(x, 0) : x \in \text{Max}(X, \leq_p)\}$. Hence $i(\text{Max}(X, \leq_p)) = \text{Max}(\mathbf{B}X)$.

(c) If (X, p) is complete, we have that $(\mathbf{B}X, \sqsubseteq_{d_p})$ is a domain by Theorem 3.1 of [17], and hence the sets $\uparrow(x, r), (x, r) \in \mathbf{B}X$, form a base for $\sigma(\mathbf{B}X)$. Moreover, by Corollary 3.1 of [17], we have that $(x, r) \ll (y, s) \Leftrightarrow d_p(x, y) < r - s$, for all $(x, r), (y, s) \in \mathbf{B}X$.

It is easy to check that for each $(x, r) \in \mathbf{B}X$ and each $\varepsilon > 0$, one obtains

$$\uparrow(x, r + \frac{\varepsilon}{2}) \subseteq B_{P'}((x, r), \varepsilon).$$

Consequently $\tau_{P'} \subseteq \sigma(\mathbf{B}X)$.

Now let $(z, t) \in \mathbf{B}X$ such that $(x, r) \in \uparrow(z, t)$. Then, there is $\varepsilon > 0$ such that $d_p(z, x) < t - r - \varepsilon$. Then, for $(y, s) \in B_{P'}((x, r), \varepsilon)$, we have

$$d_p(z, y) \leq d_p(z, x) + d_p(x, y) < (t - r - \varepsilon) + (r - s + \varepsilon) = t - s,$$

and thus $(y, s) \in \uparrow(z, t)$. Therefore $\sigma(\mathbf{B}X) \subseteq \tau_{P'}$.

(d) It suffices to show that $\omega(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)} \subseteq \sigma(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)}$. Let $(x, r) \in \text{Max}(\mathbf{B}X)$ and $(z, t) \in \mathbf{B}X$ such that $(x, r) \in \mathbf{B}X \setminus \uparrow(z, t)$. By (b), $x \in \text{Max}(X, \leq_p)$ and $r = 0$. Then $p(x, x) = 0$ by our hypothesis. Since $d_p(z, x) > t$ there exists $\varepsilon > 0$ such that $d_p(z, x) > t + \varepsilon$, so, $B_{P'}((x, 0), \varepsilon) \in \sigma(\mathbf{B}X)$ by (c). Then, for each $(y, 0) \in B_{P'}((x, 0), \varepsilon) \cap \text{Max}(\mathbf{B}X)$, we have $p(y, y) = 0$ and thus

$$t + \varepsilon < d_p(z, x) \leq d_p(z, y) + d_p(y, x) = d_p(z, y) + d_p(x, y) < d_p(z, y) + \varepsilon.$$

Hence $d_p(z, y) > t$, i.e., $(y, 0) \in \mathbf{B}X \setminus \uparrow(z, t)$. We conclude that $\omega(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)} \subseteq \sigma(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)}$, so $\sigma(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)} = \lambda(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)}$.

(e) The proof is similar to the one given in Theorem 4.1 (e) of [17], so we omit some details.

Suppose that $(\mathbf{B}X, P')$ is complete and let $(x_n)_{n \in \mathbb{N}}$ be a Cauchy sequence in the metric space $(X, (d_p)^s)$. Then $((x_n, 0))_n$ is a Cauchy sequence in the complete metric space $(\mathbf{B}X, (d_{P'})^s)$, so there is $(z, t) \in \mathbf{B}X$ such that $((x_n, 0))_n$ converges to (z, t) in $(\mathbf{B}X, (d_{P'})^s)$. So, in particular, we obtain that for each $\varepsilon > 0$, $p(z, z) + 2r - p(z, z) < \varepsilon$, and thus $r = 0$. Then, it immediately follows that $(x_n)_n$ converges to z in $(X, (d_p)^s)$. We conclude that p is a complete partial metric on X .

Conversely, suppose that (X, p) is complete and let $((x_n, r_n))_{n \in \mathbb{N}}$ be a Cauchy sequence in $(\mathbf{B}X, (d_{P'})^s)$. Then, for each $\varepsilon > 0$ there is $n_\varepsilon \in \mathbb{N}$ such that $p(x_n, x_m) - p(x_n, x_n) + r_m - r_n < \varepsilon$ and $p(x_n, x_m) - p(x_m, x_m) + r_n - r_m < \varepsilon$ for all $n, m \geq n_\varepsilon$. Hence the sequence $(r_n)_{n \in \mathbb{N}}$ is bounded, so there exist a subsequence $(r_{n_k})_{k \in \mathbb{N}}$ of $(r_n)_{n \in \mathbb{N}}$ and an $r \in \mathbb{R}^+$ such that $\lim_k r_{n_k} = r$. It immediately follows that $(x_{n_k})_{k \in \mathbb{N}}$ is a Cauchy sequence in $(X, (d_p)^s)$. Then, there is $x \in X$ such that $(x_{n_k})_{k \in \mathbb{N}}$ converges to x in $(X, (d_p)^s)$. Finally, it is easy to deduce that $((x_n, r_n))_{n \in \mathbb{N}}$ converges to (x, r) in $(\mathbf{B}X, (d_{P'})^s)$. This concludes the proof.

Remark 1. If (X, p) is partial metric space such that τ_p is a T_1 -topology, then $\text{Max}(X, \leq_p) = X$, so, by Theorem 1(b), $i(X) = \text{Max}(\mathbf{B}X)$ in this case.

The following example shows that condition “ $\ker p = \text{Max}(X, \leq_p)$ ” can not be omitted in Theorem 1 (d).

Example 1. Let $X = \{0, \infty\} \cup \mathbb{N} \setminus \{1\}$, and let $p : X \times X \rightarrow \mathbb{R}^+$ defined as $p(0, 0) = 0$, $p(0, \infty) = 2$, $p(\infty, \infty) = 0$, $p(n, n) = 1/n$, $p(0, n) = 1 + 1/n$, and $p(n, \infty) = 1$ for all $n \in \mathbb{N} \setminus \{1\}$, and $p(n, m) = 1 + 1/n + 1/m$ for all $n, m \in \mathbb{N} \setminus \{1\}$ with $n \neq m$.

It is routine to check that p is a partial metric on X , and it is complete because every Cauchy sequence in $(X, (d_p)^s)$ is eventually constant.

On the other hand, $\text{Max}(X, \leq_p) = X$, so $i(X) = \text{Max}(\mathbf{B}X)$ by Theorem 1 (b). However $\ker p = \{\infty\}$.

We show that $\sigma(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)} \neq \lambda(\mathbf{B}X)|_{\text{Max}(\mathbf{B}X)}$. Indeed, we have $(0, 0) \in \mathbf{B}X \setminus \uparrow(\infty, 1)$ because $d_p(\infty, 0) = p(\infty, 0) - p(\infty, \infty) = 2$. Nevertheless, for each $n \in \mathbb{N}$ we have

$$(n + 1, 0) \in B_{P'}((0, 0), 1/n) \cap \uparrow(\infty, 1),$$

because $P'((0, 0), (n + 1, 0)) = 1 + 1/(n + 1) < 1 + 1/n = P'((0, 0), (0, 0)) + 1/n$, and $d_p(\infty, n + 1) = 1$ for all $n \in \mathbb{N}$.

1 Since by Theorem 1 (c), $\{B_{P'}((0,0), 1/n) : n \in \mathbb{N}\}$ is a $\sigma(\mathbf{B}X)$ -local base at
2 $(0,0)$, we conclude that $\lambda(\mathbf{B}X)_{|\text{Max}(\mathbf{B}X)}$ is strictly finer than $\sigma(\mathbf{B}X)_{|\text{Max}(\mathbf{B}X)}$.

3
4 The computational model suggested by conditions (1)-(3) in p. 59 of [22] joint
5 with Definition 5.1 of [17] and Theorem 1 above, motivate the following notions.

6
7 **Definition 1.** A partially metrizable computational model is a triple (L, \sqsubseteq, P)
8 such that (L, \sqsubseteq) is a domain and P is a complete partial metric on L that induces
9 the Scott topology.

10
11 **Definition 2.** A topological space (X, τ) has a partially metrizable computa-
12 tional model if there is a partially metrizable computational model (L, \sqsubseteq, P) and
13 an homeomorphism ϕ between (X, τ) and a $\tau_{(d_P)^{-1}}$ -closed subset of (L, P) such
14 that $\phi(\text{Max}(X, \leq_\tau)) = \text{Max}(L)$.

15
16 We conclude the paper with a characterization of those topological spaces that
17 admit a compatible complete partial metric via the model proposed in Definition 1.

18
19 **Theorem 2.** *A topological space has a partially metrizable computational model
20 if and only if it admits a compatible complete partial metric.*

21
22 *Proof.* Suppose that the topological space (X, τ) has a partially metrizable computa-
23 tional model (L, \sqsubseteq, P) . Let ϕ an homeomorphism between (X, τ) and a $\tau_{(d_P)^{-1}}$ -
24 closed subset of (L, P) such that $\phi(\text{Max}(X, \leq_\tau)) = \text{Max}(L)$. For each $x, y \in X$
25 put $p(x, y) = P(\phi(x), \phi(y))$. Then p is a partial metric on X such that $\tau_p = \tau$.
26 Moreover p is complete: Indeed, let $(x_n)_{n \in \mathbb{N}}$ be a Cauchy sequence in the metric
27 space $(X, (d_p)^s)$. Then $(\phi(x_n))_n$ is a Cauchy sequence in the complete metric space
28 $(L, (d_P)^s)$, so there is $z \in L$ such that $(\phi(x_n))_n$ converges to z in $(L, (d_P)^s)$, and
29 hence in $(L, (d_P)^{-1})$. Since $\phi(X)$ is $\tau_{(d_P)^{-1}}$ -closed in L , it follows that $z \in \phi(X)$,
30 and consequently $(x_n)_n$ converges to $\phi^{-1}(z)$ in $(X, (d_p)^s)$. We conclude that p is a
31 complete partial metric on X .

32
33 Conversely, if (X, τ) admits a compatible complete partial metric p , then
34 $(\mathbf{B}X, \sqsubseteq_{d_p})$ is a domain. So, by Theorem 1, $(\mathbf{B}X, \sqsubseteq_{d_p}, P')$ is a partially metrizable
35 computational model for (X, τ) .

36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

References

- [1] M. Ali-Akbari, B. Honari, M. Pourmahdian, M.M. Rezaei, The space of formal balls and models of quasi-metric spaces, *Math. Struct. Comput. Sci.* 19 (2009), 337-355.
- [2] A. Edalat, R. Heckmann, A computational model for metric spaces, *Theoret. Comput. Sci.* 193 (1998), 53-73.
- [3] A. Edalat, Ph. Sünderhauf, Computable Banach spaces via domain theory, *Theoret. Comput. Sci.* 219 (1999), 169-184.
- [4] R. Engelking, *General Topology*, Monografie Mat. Tom 60, PWN-Polish Scientific Publishers, Warszawa, 1977.
- [5] B. Flagg, R. Kopperman, Computational models for ultrametric spaces, *Electronic Notes Theoret. Comput. Sci.* 6 (1997), 151-159.
- [6] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, D.S. Scott, *Continuous Lattices and Domains*, Encyclopedia of Mathematics and its Applications vol. 93. Cambridge Univ. Press, 2003.
- [7] R. Heckmann, Approximation of metric spaces by partial metric spaces, *Appl.Cat. Struct.* 7 (1999), 71-83.
- [8] R. Kopperman, H.P. Künzi, P. Waszkiewicz, Bounded complete models of topological spaces, *Topology Appl.* 139 (2004), 285-297.
- [9] M. Krötzsch, Generalized ultrametric spaces in quantitative domain theory, *Theoret. Comput. Sci.* 368 (2006), 30-49.
- [10] H.P.A. Künzi, Nonsymmetric distances and their associated topologies: About the origins of basic

- 1 ideas in the area of asymmetric topology, in: C.E. Aull, R. Lowen (Eds.), Handbook of the History
2 of General Topology, vol. 3, Kluwer, Dordrecht, 2001, pp. 853-968.
- 3 [11] J. Lawson, Spaces of maximal points, *Math. Struct. Comput. Sci.* 7 (1997), 543-556.
- 4 [12] K. Martin, Domain theoretic models of topological spaces, *Electronic Notes Theoret. Comput. Sci.*
5 13 (1998), 173-181.
- 6 [13] S.G. Matthews, Partial metric topology, in: Proceedings 8th Summer Conference on General Topology
7 and Applications, *Ann. New York Acad. Sci.* 728 (1994), 183-197.
- 8 [14] J. Rodríguez-López, S. Romaguera, O. Valero, Denotational semantics for programming languages,
9 balanced quasi-metrics and fixed points, *Internat. J. Comput. Math.* 85 (2008), 623-630.
- 10 [15] S. Romaguera, On Computational Models for the Hyperspace, *Advances in Mathematics Research*,
11 volume 8, Nova Science Publishers, New York, 2009, pp. 277-294.
- 12 [16] S. Romaguera, O. Valero, A quasi-metric computational model from modular functions on monoids,
13 *Internat. J. Comput. Math.* 86 (2009), 1668-1677.
- 14 [17] S. Romaguera, O. Valero, A quantitative computational model for complete partial metric spaces via
15 formal balls, *Math. Struct. Comput. Sci.* 19 (2009), 541-563.
- 16 [18] S. Romaguera, O. Valero, Domain theoretic characterisations of quasi-metric completeness in terms
17 of formal balls, *Math. Struct. Comput. Sci.* 20 (2010), 453-472.
- 18 [19] J.J.M.M. Rutten, Weighted colimits and formal balls in generalized metric spaces, *Topology Appl.* 89
19 (1998), 179-202.
- 20 [20] M. Schellekens, A characterization of partial metrizability. Domains are quantifiable, *Theoret. Com-*
21 *put. Sci.* 305 (2003), 409-432.
- 22 [21] M.B.Smyth, The constructive maximal point space and partial metrizability, *Ann. Pure Appl. Logic*
23 137 (2006), 360-379.
- 24 [22] P. Waszkiewicz, Quantitative continuous domains, *Appl. Cat. Struct.* 11 (2003), 41-67.
- 25 [23] P. Waszkiewicz, Partial metrisability of continuous posets, *Math. Struct. Comput. Sci.* 16 (2006),
26 359-372.
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60