

The dynamics of Salmonella occurrence in commercial laying hen flocks throughout a laying period

Jochen Schulz, Sebastiaan van Hoorebeke, Birthe Hald, Jörg Hartung, Filip van Immerseel, Ines Radtke, Susanne Kabell, Jeroen Dewulf

▶ To cite this version:

Jochen Schulz, Sebastiaan van Hoorebeke, Birthe Hald, Jörg Hartung, Filip van Immerseel, et al.. The dynamics of Salmonella occurrence in commercial laying hen flocks throughout a laying period. Avian Pathology, 2011, 40 (03), pp.243-248. 10.1080/03079457.2010.544290 . hal-00710047

HAL Id: hal-00710047 https://hal.science/hal-00710047

Submitted on 20 Jun 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The dynamics of Salmonella occurrence in commercial laying hen flocks throughout a laying period

Journal:	Avian Pathology
Manuscript ID:	CAVP-2010-0158.R1
Manuscript Type:	Review
Date Submitted by the Author:	17-Nov-2010
Complete List of Authors:	Schulz, Jochen; University of Veterinary Medicine Hannover, Foundation, Germany, Institute for Animal Hygiene, Animal Welfare and Farm Animal Behaviour Van Hoorebeke, Sebastiaan; Ghent University, Obstetrics, Reproduction and Herd Health Hald, Birthe; Technical University of Denmark, National Veterinary Institute, Department of poultry, fish and fur animals Hartung, Jörg; University of Veterinary Medicine Hannover, Foundation, Germany, Institute for Animal Hygiene, Animal Welfare and Farm Animal Behaviour Van Immerseel, Filip; Ghent University, Pathology, Bacteriology and Avian Diseases Radtke, Ines; University of Veterinary Medicine Hannover, Foundation, Germany, Institute for Animal Hygiene, Animal Welfare and Farm Animal Behaviour Van Immerseel, Filip; Ghent University, Pathology, Bacteriology and Avian Diseases Radtke, Ines; University of Veterinary Medicine Hannover, Foundation, Germany, Institute for Animal Hygiene, Animal Welfare and Farm Animal Behaviour Kabell, Susanne; Food and Veterinary Agency, Dept. for Fish- and Animal Health Dewulf, Jeroen; Ghent University, Obstetrics, Reproduction and Herd Health
Keywords:	Salmonella, occurence, laying hen flocks, laying period
	1



Cavp-2010-0158.R1

The dynamics of *Salmonella* occurrence in commercial laying hen flocks throughout a laying period

J. Schulz¹*, S. Van Hoorebeke², B. Hald⁴, J. Hartung¹, F. Van Immerseel³, I. Radtke¹, S. Kabell⁵ and J. Dewulf²

 ¹ Institute for Animal Hygiene, Animal Welfare and Farm Animal Behaviour, University of Veterinary Medicine Hannover, Foundation, Büntewg 17p, 30559 Hannover, Germany
 ² Department of Reproduction, Obstetrics and Herd Health, Veterinary Epidemiology Unit, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium
 ³ Department of Pathology, Bacteriology and Avian Diseases, Veterinary Epidemiology Unit, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium
 ⁴ National Veterinary Institute, Department of poultry, fish and fur animals, Technical University of Denmark, Hangøvej 2, 8200 Århus, Denmark
 ⁵ Food and Veterinary Agency, Dept. for Fish- and Animal Health, 100 Tórshavn, Faroe Islands, Denmark

Short title

Salmonella occurrence in laying hens

Corresponding author: Dr. Jochen Schulz, Tel.: 0049-511-953-8546; Fax.: 0049-511-953-

8588; E-mail: Jochen.Schulz@tiho-hannover.de

Received: 17 November 2010

Abstract

Contaminated eggs and egg products have been recognised since many years as an important source of Salmonella infections in humans in the European Union and in the United States. Longitudinal studies can help to increase our knowledge about the dynamics of the occurrence of *Salmonella* in the course of a laying period. The total of 41 laying hen flocks, 18 in Belgium, 6 in Denmark and 17 in Germany were followed during an entire laying period. Samples taken from the empty cleaned and disinfected poultry houses were all negative for Salmonella. After hens arrived on the farms five pooled faecal samples, one pooled dust sample and 40 cloacal swabs (Belgium and Germany) or 40 swabs from fresh droppings (Denmark) were taken four times from 18 flocks, three times from 21 flocks and two times from 2 flocks in the course of the laying period. Ten flocks (two Belgian and eight German flocks) were tested up to three times positive for Salmonella. Forty three out of 50 positive samples contained Salmonella Enteritidis phage type 4 (29 isolates) or phage type 8 (14 isolates). The probability of subsequent Salmonella positive findings increased significantly in *Salmonella* positive flocks (p < 0.05, odds ratio = 6.4). However, the probability of finding Salmonella did not depend on the time of sampling in the laying period or the season.

Introduction

Salmonellosis belongs to the most frequently reported zoonotic food-borne diseases worldwide. It is known since many years that contaminated eggs and egg products are important vehicles for the transmission of these organisms to the consumer in the European Union and in the United States (Patrick et al., 2004; European Food Safety Authority, 2009). Salmonella can be found either on the surface of the eggshell or in the egg content. The eggshell can become contaminated as a result of infection of the oviduct or by colonization of the intestine of a hen (De Reu et al., 2008). Also the presence of Salmonella in faeces can lead to eggshell contamination (Gast & Beard, 1990). Salmonella in the egg content is resulting from an infection of the reproductive organ or is a consequence of the migration of these bacteria through the egg shell and membranes (Gantois et al., 2009). The most common Salmonella serovar in eggs and egg products is Salmonella Enteritidis which is also the most frequently reported serovar in Salmonella infections in humans in the European Union and in the United states (Braden, 2006; European Food Safety Authority, 2009). Case-control studies and outbreak investigations have demonstrated the association between human cases and the presence of Salmonella Enteritidis in laying hen flocks (Altekruse et al., 1993; Rampling, 1993; Trepka et al., 1999).

Many studies have investigated the transmission routes and identified risk factors for *Salmonella* infections in laying hen flocks (Davies & Breslin, 2003; Gradel *et al.*, 2004; Mollenhorst, 2005; Humphrey, 2006; Carrique-Mas *et al.*, 2008; Namata *et al.*, 2008; Van Hoorebeke *et al.*, 2010). However, studies over longer time periods describing the dynamics of the within flock *Salmonella* prevalence during a production period are still rare. This may be caused by high costs of such complex field studies and simply by the difficulty to convince

farmers to take part in long lasting investigations (Wales *et al.*, 2007). Therefore, in the framework of the EU project SAFEHOUSE a longitudinal study was carried out in laying hen flocks on different farms in Belgium (BE), Denmark (DK) and Germany (DE). The flocks were sampled several times in the course of one laying period in order to describe the dynamics of *Salmonella* occurrence throughout a production period under field conditions.

Materials and Methods

Sampled flocks. In total 41 flocks were included in the study: 18 flocks from BE (1 enriched cage system, 4 battery cage, 5 free range, 4 barn and 4 organic systems), 6 flocks from DK (1 free range system, 3 battery cage and 2 barn system) and 17 flocks from DE (4 battery cage, 5 free range, 5 barn and 3 small colony keeping systems). A flock was defined according to regulation (EC) 2160/2003 of the European Parliament and of the Council (Anon, 2003). That means all hens of the same health status kept on the same premises or in the same enclosure and constituting a single epidemiological unit. In case of housed hens this includes all birds sharing the same airspace. Only flocks larger than 1000 birds were included. On farms with several flocks (overall 24 farms and 21 of them with flocks of different ages) one flock was selected randomly for sampling. The housing systems were categorised according to the EU directive 2002/4/EC and council regulations (EC) No 1804/1999 (Anon., 1999; Anon., 2002). An exception was the German small colony keeping system which is defined by the German authorities (German Federal Ministry of Justice, 2006). This system resembles the EU furnished cage but offers the hens more space and extra perches at different heights.

Periods of sampling. Field studies were performed from February 2007 until July 2008 in BE and DE. In DK flocks were sampled from October 2007 until March 2009. First samples were taken from the empty and disinfected animal houses one week before the hens arrived at the farm. The age at placement varied between 15 and 22 weeks. In three flocks no samples were taken from empty houses because hens of different ages were housed in the same premise and the animal houses were never empty. In total 139 samplings were carried out at different times during one egg-laying period. Flocks were sampled three or four times at different intervals with the exception of two flocks which were sampled only twice. Between the different countries sampling intervals varied because of different logistics. The laying periods were divided in six 10-week intervals for data analysis. Table 1 shows the number of flocks sampled in each country during these intervals.

Number and sample type taken on-farm. One pooled dust sample, five nest-box samples or 5 cage samples were taken in the empty and disinfected houses. The dust was collected from 20 different locations in the animal house. The dusty material was transferred by a gloved hand or sterile brush into a sterile bag. The minimum amount was 25 g dust per pooled sample. If no suitable surfaces with dust were present in the cleaned houses, surfaces from 20 locations were swabbed with sterile swabs, which were placed in Ames medium (Oxoid, Deutschland GmbH). Swabs and Ames medium were also used to swab 5 randomly selected nest-boxes except in battery cages which do not provide nest-boxes. In the latter case the grids of 5 randomly selected cages were swabbed.

From each laying hen flock 5 nest-box samples or 5 cage samples and 1 pooled dust sample were taken per visit as described above. Furthermore 5 pooled faecal samples and individual cloacal swabs from 40 randomly selected laying hens (in BE and DE) were sampled at each farm visit. In Denmark, the 40 swabs were obtained from 40 randomly

selected fresh faecal droppings because farmers did not allow sampling their hens. Depending on the housing system, the pooled faecal samples were taken as follows:

- Cage systems and small colony keeping systems: 5 samples of mixed fresh faeces were either taken from dropping belts, scrapers or deep pits depending on the type of cage (conventional battery cage, furnished cage, small colony keeping system). Each pooled faecal sample consisted of approximately 250 g.
- Barn, free-range and organic systems: for each of the 5 pooled faecal samples 60 piles of fresh faeces were collected from the floor and the slats. Each pooled faecal sample consisted of approximately 250 g.

The pooled faecal samples were put in a sterile box. New gloves were used for collecting each pooled faecal sample.

Cloacal swabs were taken by inserting a cotton-tipped swab approximately 5 cm into the cloaca, taking care to avoid contact with the surrounding feathers and skin. Afterwards, the swabs were stored in tubes containing Ames medium. A sample size of 40 swabs allows detecting a within-flock prevalence of 7% at a level of confidence of 95% in a flock with more than 1000 hens (WinEpiscope 2.0). All samples were placed in leak-proof bags and transported in container boxes to the laboratory under ambient temperatures. The samples were analysed either immediately after arrival at the laboratory or stored at 4°C for analysis within the next 24 hours.

Bacteriological analysis of samples. All samples were analyzed according to ISO 6579:2002/Amd 1:2007 Annex D: Detection of *Salmonella* spp. in animal faeces and in environmental samples from the primary production stage. From each of the pooled faecal and mixed dust sample, 25 g was added to 225 ml of buffered peptone water (BPW) (Oxoid, Basingstoke, Hampshire, UK). All cloacal swabs, swabs from nest-boxes and from dusty

surfaces were individually placed in 9 ml of BPW. Pooled faecal and mixed dust samples were mixed in a stomacher bag for 1 minute. All samples were incubated for 18 ± 2 h at $37 \pm$ 1 °C. Three droplets of each pre-enrichment culture were dropped on the surface of a modified semi-solid Rappaport-Vassiliadis (MSRV) (Difco; Becton Dickinson) agar plate containing 0.01 g l⁻¹ novobiocine. The plates were incubated for $2 \times 24 \pm 3$ h at 41.5 ± 1 °C. Material of suspected white cultures from the borderlines of the growth zones were transferred on Brilliant Green Agar (BGA; Oxoid) and Xylose Lysine Deoxycholate agar (XLD; Oxoid), followed by incubation for 24 ± 2 h at 37 ± 1 °C. Presumed *Salmonella* colonies on BGA and XLD were biochemically confirmed using urea agar, triple sugar iron agar and lysine-decarboxylase broth. Serotyping of *Salmonella*-isolates according to the White-Kauffmann-Le Minor scheme (Grimont & Weill, 2007) was performed in the national Reference Laboratories for *Salmonella* in each participating country. Phage typing of *S.* Enteritidis was carried out as described by Ward *et al.* (1987).

Data processing and analysis. In order to get reliable information on the sampled flocks (e.g. number and age of hens in a flock) and the farms (e.g. total number of flocks and hens, age of different flocks, cleaning measures and disinfection procedures) a questionnaire was developed and filled in by interviewing the farmer at one of the farm visits as described by Van Hoorebeke *et al.* (2010). Answers in the questionnaires were coded and transferred into an excel spread sheet (Microsoft Office Excel 2003, Microsoft Cooperation). The variable "period of sampling" was divided into six 10-weeks intervals. To define the category "season", the meteorological seasons for Europe (spring begins on the first of March, summer on the first of June, autumn on the first of September and winter on the first of December) were chosen. The effect of the period of sampling and the season on the within flock prevalence was analysed using logistic regression as was the effect of a previous sample result

within the same production round on the next sample result. For statistical analysis the software SPSS 17.0 (SPSS Inc., Chicago, IL) was used.

Results

Salmonella detection. In none of the animal houses members of Salmonella were found after cleaning and disinfecting prior to stocking. Ten out of the 41 sampled flocks were tested once, twice or three times positive for *Salmonella* in the course of one laying period. Two Belgian and eight German flocks were Salmonella-positive across five different housing systems. Table 2 gives an overview of the kind of positive samplings, the number of positive samples, sampling intervals and detected serovars (including S. Enteritidis phage types) in the positive tested flocks. Six of these flocks (no. 7, 8, 25, 26, 31 and 41) were housed in all-in and all-out systems, two flocks (no. 27 and 32) originate from farms with hens of different ages in different houses and two flocks (no. 33 and 40) were kept in multiage systems. Two flocks (no. 25 and 41) hosted different serovars. The most prevalent isolate was S. Enteritidis found in 44 (88%) of the 50 positive tested samples (29 isolates of phage type 4, 14 isolates of phage type 8, phage type of 1 isolate could not be identified). The serovars S. Infantis, S. London and one unidentified serotype from serogroup B were isolated from 6 samples (12%). Table 2 shows that Salmonella were detected 18 times in 10 flocks by using three kinds of sampling techniques. Already 17 positive samplings in 10 flocks were obtained by environmental samplings (pooled faecal and pooled dust samples). The number of positive flocks (pooled faecal = 7, pooled dust = 6, cloacal swabs = 3) and samplings (pooled faecal = 12, pooled dust = 10, cloacal swabs = 5) decreases when only a single sampling technique is taken into account.

Statistical analysis. Observed *Salmonella* prevalence throughout the production period is given in Figure 1. Although differences are obvious between the different sampling intervals, these were not statistically significant. No significant effect of the season was found either. However, the risk to find *Salmonella* in a flock increased significantly (p < 0.01) (odds ratio = 6.4, 95% confidence interval: 2.051 – 19.745) when *Salmonella* was found in the previous sampling of the same flock.

Discussion

The occurrence of *Salmonella* was observed in Belgian and German laying hen flocks but not in Danish flocks what can be explained by the relatively small number of Danish flocks included into the study and the well-known low *Salmonella* prevalence in Danish laying hen flocks (Wegener *et al.*, 2003; European Food Safety Authority, 2006). Therefore swab samples taken from fresh droppings instead of cloacal swabs in Denmark had most likely no considerable impact on the overall results.

Ten out of the 41 sampled flocks were tested one, two or three times *Salmonella*positive in the course of the investigated laying period by using cloacal swabs, dust samples and samples of faeces. Taking cloacal swabs was less useful to detect *Salmonella*. This was probably influenced by the relatively low within flock prevalence of shedding hens (< 7%) even in a *Salmonella* contaminated environment. Sampling the environment was rather successful and especially the combination of faeces and dust samplings seems to be more effective than using a single sampling technique. These findings support the recommendation to sample both faeces and dust to increase the sensitivity of *Salmonella* detection in laying

hen houses (Carrique-Mas & Davies, 2008). Although non-shedding hens with infected organs cannot be detected with the applied methods (Van Hoorebeke *et al.*, 2009), the use of more sensitive environmental samplings can help identify flocks at risk of producing *S*. Enteritidis-contaminated eggs (Henzler *et al.*, 1998).

A Salmonella positive finding in a flock significantly increased the probability of subsequent positive results in the same flock. In these cases generally the same serovars and phage types were detected. Especially the same phage types in environmental samples and in cloacal swabs of the same flock demonstrate the close relationship between the infected animals and the environment which seems to play an important role as a reservoir. Furthermore, in most of the positive tested flocks the infection was already observed in the early stage of the laying period. Although the samplings of the cleaned and disinfected laying hen houses were negative, undetected residual contamination may have contributed to an initial infection of hens. Especially in cage and multi-age systems, which represent seven (70%) of the positive tested flocks in this study, an effective cleaning and disinfection is difficult to perform in practice and may even fail (Davies & Breslin, 2004; Wales et al., 2007). Another source for infections in the early laying period may be living vectors such as rodents and arthropods. This was not specifically investigated in this study. However, it is known that they can contribute to the horizontal transmission of infectious agents including Salmonella (Gast et al., 1998; Carrique-Mas et al., 2008). These vectors may also be responsible for infections even on farms with all-in all-out management because they can stay on the farm after depopulation and re-enter into the new flocks (Hald et al., 1998). Beside horizontal transmission also vertical transmission may have contributed to the Salmonella presence in the investigated flocks. Salmonella infections in parent breeding flocks are described as an existing problem in Germany (European Food Safety Authority, 2009). Although vaccination is compulsory in German parent breeding flocks, this route of

11

transmission should not be neglected because even vaccinated hens can become infected with *Salmonella* Enteritidis and may transmit this organism to their descendants (Davies & Breslin, 2004).

The presented results show no significant differences concerning the Salmonella prevalence in the different sampling intervals. These findings contradict somehow other studies which described an increase of infected flocks with increasing age of the hens. A higher susceptibility of older hens to infections, molting and an accumulation of Salmonella in the environment were suggested as factors which lead to an increase of Salmonella positive flocks and an increase of Salmonella positive samples in ageing flocks respectively (Schlosser et al., 1999; Garber et al., 2003; Van de Giessen et al., 2006; Wales et al., 2007). In the present study induced molting was not practiced on the sampled farms and no higher numbers of Salmonella positive faecal, dust and cloacal swab samples were observed towards the end of the laying period. These findings may partly be influenced by cleaning procedures and the removal of faeces during the production cycle (Wales et al., 2007). Hence, differences in the farm management and in the design of studies make it difficult to compare the results of this study and previous investigations. However, an early Salmonella contamination in the laying period did apparently not result in an increased number of shedding hens. In contrast to other opinions, it seems that an increasing number of shedding hens towards the end of a laying period does not hold true for infected flocks in general. This may be typical in particular for early infected flocks due to the decline of shedding hens within a few weeks after infection (De Vylder et al., 2009) and should be considered in future longitudinal studies by taking samples in the very early stage of lay (in the first three weeks after housing).

Seasonal effects may also influence the detection of *Salmonella* in laying hen flocks in longitudinal studies (Wales *et al.*, 2007). Therefore the season was included in the logistic regression model but no significant seasonal effects were observed. The discussion about

seasonal effects is controversial and it seems that not the changing climatic conditions itself but rather their consequences for the management or the presence of living vectors are influencing *Salmonella* detection and transmission (Carrique-Mas *et al.*, 2008; Van Hoorebeke *et al.*, 2010).

The dynamics of *Salmonella* infection in laying hen flocks is very complex. It is strongly depending on the farm management (e.g. cleaning and disinfection procedures), the immune status of the birds, the presence of living vectors as carriers and an accumulation of *Salmonella* in the flock environment. This study indicates that early infections of laying hen flocks with the important human pathogen *S*. Enteritidis (PT4 and PT8) can lead to a continuous infection of hens and to a persistent contamination of their environment. This may increase also the risk off egg contaminations. Therefore more stringent action should be taken both to eliminate *Salmonella* from breeding and production flocks by applying comprehensive bio-security measures. Special regards should be paid to avoid *Salmonella* introduction in laying hen flocks at any stage. This will be a considerable contribution to decrease the risk for the consumer.

Acknowledgements

The study was funded by the European Community (Project FOOD-CT-2006-035547). The authors would like to thank all farmers participating in the study for very good cooperation.

References

- Altekruse, S.F., Koehler, J., Hickmanbrenner, F., Tauxe, R.V. & Ferris, K. (1993). A comparison of *Salmonella* Enteritidis phage types from egg-associated outbreaks and implicated laying flocks. *Epidemiology and Infection*, 110, 17-22.
- Anon. (1999). Council regulation (EC) No 1804/1999 of 19 July 1999 supplementing
 Regulation (EEC) No 2092/91 on organic production of agricultural products and
 indications referring thereto on agricultural products and foodstuffs to include livestock
 production. *Official Journal of the European Community, L 222, 1-28.*
- Anon. (2002). Comission Directive 2002/4/EC of 30 January 2002 on the registration of establishment keeping laying hens covered by Council Directive 1999/74/EC. Official Journal of the European Communit, L 30, 44-46.
- Anon. (2003). Regulation (EC) No 2160/2003 of the European Parliament and of the Council of November 2003 on the control of *Salmonella* and other specific foodborne zoonotic agents. *Official Journal of the European Communit,L 325,* 1-15.
- Braden, C.R. (2006). *Salmonella* enterica serotype Enteritidis and eggs: A national epidemic in the United States. *Clinical Infectious Diseases*, *43*, 512-517.
- Carrique-Mas, J.J. & Davies, R.H. (2008). Salmonella Enteritidis in commercial layer flocks in Europe: Legislative background, on-farm sampling and main challenges. *Brazilian Journal of Poultry Science*, *10*, 1-9.
- Carrique-Mas, J.J., Breslin, M., Snow, L., Mclaren, I., Sayers, A.R. & Davies, R.H. (2008). Persistence and clearance of different *Salmonella* serovars in buildings housing laying hens. *Epidemiology and Infection*, 137, 837-846.

- Davies R. & Breslin, M. (2003). Observations on Salmonella contamination of commercial laying farms before and after cleaning and disinfection. Veterinary Record, 152, 283-287.
- Davies R. & Breslin, M. (2004). Observation on Salmonella contamination of eggs from infected commercial laying hen flocks where vaccination for Salmonella enterica serovar Enteritidis had been used. Avian Pathology, 33, 133-144.
- De Reu, K., Messens, W., Heyndrickx, M., Rodenburg, T.B., Uyttendaele, M. & Herman, L.
 (2008). Bacterial contamination of table eggs and the influence of housing system.
 World's Poultry Science Journal, 64, 5-19.
- De Vylder, J., Van Hoorebeke, S., Ducatelle, R., Pasmans, F., Haesebrouck, F., Dewulf, J.
 & Van Immerseel, F. (2009). Effect of the housing system on shedding and colonization of gut and internal organs of laying hens with *Salmonella* Enteritidis. *Poultry Science*, 88, 2491–2495.
- European Food Safety Authority (2006). Preliminary report on the analysis of the baseline study on the prevalence of *Salmonella* in Laying Hen Flocks of *Gallus Gallus. The EFSA Journal*, 81, 1-71.
- European Food Safety Authority (2009). The Community summary report on trends and sources of zoonoses and zoonotic agents in the European Union in 2007. *The EFSA Journal*, 223, 310 pp.
- Gantois, I., Ducatelle, R., Pasmans, F., Haesebrouck, F., Gast, R.K., Humphrey, T.J. & Van Immreseel, F. (2009). Mechanisms of egg contamination by *Salmonella* Enteritidis.
 FEMS *Microbiological Reviews*, *33*, 718-738
- Garber, L., Schmeltzer, M., Fedorka-Cray, P., Ladely, S. & Ferris, K. (2003). *Salmonella* enterica Serotype enteritidis in table egg layer house environments and in mice in US layer houses and associated risk factors. *Avian Diseases*, 47, 134-142.

- Gast, R.K., & Beard, C.W. (1990). Isolation of *Salmonella* Enterititidis from internal organs of experimentally infected hens. *Avian Diseases*, *34*, 991-993.
- Gast, R.K., Mitchell, B.W. & Holt, P. S. (1998). Airborne transmission of Salmonella Enteritidis infection between groups of chicks in controlled-environment isolation cabinets. Avian Diseases, 42, 315-320.
- German Federal Ministry of Justice (2006). Tierschutz-Nutztierhaltungsverordnung. Neugefasst durch Bek. v. 22.8.2006 I 2043; geändert durch V v. 1.10.2009 I 3223. *http://bundesrecht.juris.de/tierschnutztv/*
- Gradel, K.O., Jorgensen, J.C., Andersen, J.S., & Corry, J.E.L. (2004). Monitoring the efficacy of steam and formaldehyde treatment of naturally *Salmonella*-infected layer houses. *Journal of Applied Microbiology*, 96, 613-622.
- Grimont, P.A.D., & Weill, F.-X. (2007). Antigenic Formulae of the *Salmonella* Serovars, (ninth ed.) Paris: WHO Collaborating Centre for Reference and Research on *Salmonella*, Institut Pasteur. Available at:

http://www.pasteur.fr/sante/clre/cadrecnr/salmoms/WKLM_En.pdf.

- Hald, B., A. Olsen, A. & Madsen, M. (1998). *Typhaea stercorea* (Coleoptera: Mycetophagidae), a carrier of *Salmonella enterica* serovar Infantis in a Danish broiler house. *Journal of Economic Entomology*, *91*, 660-664.
- Henzler, D.J., Kradel, D.C. & Sischo, W.M. (1998). Management and environmental risk factors for *Salmonella* enteritidis contamination of eggs. *American Journal of Veterinary Research*, 59, 824-9.
- Humphrey T. (2006). Are happy chickens safer chickens? Poultry welfare and disease susceptibility. *British Poultry Science*, 47, 379-391.

- Mollenhorst H., Van Woudenbergh, C.J., Bokkers, E.G.M. & De Boer, I.J.M. (2005). Risk factors for *Salmonella* Enteritidis infections in laying hens. *Poultry Science*, *84*, 1308-1313.
- Namata H., Méroc, E., Aerts, M., Faes, C., Abrahantes, J.C., Imberechts, H. & Mintiens, K. (2008). Salmonella in Belgian laying hens: an identification of risk factors. Preventive Veterinary Medicine, 83, 323-336.
- Patrick, M.E., Adcock, P.M., Gomez, T.M., Altekruse, S.F., Holland, B.H., Tauxe, R.V. & Swerdlow, D. L. (2004). Salmonella Enteritidis Infections, United States, 1985-1999.
 Emerging Infectious Diseases, 10, 1-7.
- Rampling, A. (1993). Salmonella Enteritidis five years on. Lancet, 342, 317-318.
- Schlosser, W.D., Henzler, D.J., Mason, J., Kradel, D., Shipman, L., Trock, S., Hurd, S.H., Hogue, A.T., Sischo, W. & Ebel, E.D. (1999). The *Salmonella enterica* serovar Enteritidis pilot project. In A.M. Saeed, R.K. Gast & M.E. Potter (Eds.). Salmonella enterica *serovar Enteritidis in humans and animals: Epidemiology, pathogenesis, and control* (pp. 353-365). Iowa State University Press. Ames, IA.
- Trepka, M. J., Archer, J.R., Altekruse, S.F., Proctor, M.E. & Davis, J.P. (1999). An increase in sporadic and outbreak-associated *Salmonella* Enteritidis Infections in Wisconsins:
 The role of eggs. *The Journal of Infectious Diseases*, 180, 1214-1219.
- Van de Giessen, A.W., Boutwknegt, M., Dam-Deisz, W.D.C., Van Pelt, W., Wannet, W.J.B.
 & Visser, G. (2006). Surveillance of Salmonella spp. and Campylobacter spp. in poultry production flocks in the Netherlands. *Epidemiology and Infection*, 134, 1266-1275.
- Van Hoorebeke, S., Van Immerseel, F., Schulz, J., Hartung, J., Harisberger, M., Barco, L.,
 Ricci, A., Theodoropoulos, G., Xylouri, E., De Vylder, J., Ducatelle, R., Haesebrouck,
 F., Pasmans, F., De Kruif, A. & Dewulf, J. (2010). Determination of the within and
 between flock prevalence and identification of risk factors for *Salmonella* infections in

laying hen flocks housed in conventional and alternative systems. *Preventive Veterinary Medicine*, *94*, 94-100.

- Van Hoorebeke S., Van Immerseel, F., De Vylder, J., Ducatelle, R., Haesebrouck, F.,
 Pasmans, F., De Kruif, A. & Dewulf, J. (2009). Faecal sampling underestimates the actual prevalence of *Salmonella* in laying hen flocks. *Zoonoses and Public Health*, 56, 471-476.
- Wales A., Breslin, M., Carter, B., Sayers, R. & Davies R. (2007). A longitudinal study of environmental *Salmonella* contamination in caged and free-range layer flocks. *Avian Pathology*, 36, 187-197.
- Ward, L.R., De Sa, J.D.H. & Rowe, B. (1987): A phage-typing scheme for Salmonella enteritidis. Epidemiology and Infection, 99, 291-294.
- Wegener, H.C., Hald, T., Lo Fo Wong, D., Madsen, M., Korsgaard, H., Bager, F., Gerner-Smidt, P. & Mølbak, K. (2003). Salmonella Control Programs in Denmark. Emerging Infectious Diseases, 9, 774-780.

Figure Legends

Figure 1. Salmonella prevalence detected in six different sampling intervals throughout a laying period

Tuble 1. Completed samplings daming a laying period										
No. of flocks,	week of laying period									
country	-1 (empty house)	0 - 10	11 - 20	21 - 30	31 - 40	41 - 50	51 - 60			
2, DK	Х		Х		Х		Х			
2, DK	Х			Х	Х		Х			
1, DK	X		Х	Х		Х				
1, DK	X		Х		Х					
18, BE	X	Х		Х	Х		Х			
17, DE	X		Х	Х		Х				

Table 1. Completed samplings during a laying period

Table 2. Sampling intervals, negative samplings (0), number and kind of positive samples (p.f., p.d., c.s.) and detected serovars (phage types) of the positive tested flocks

1 3							
Flock no., housing				week of laying period			
system and country	-1 (empty	0 - 10	11 - 20	21 - 30	31 - 40	41 - 50	51 - 60
	house)						
7. barn, BE	0	0	- 1	0	0	_	1 p.f. (S.E. PT4)
8. enriched cage, BE	0	0	_	0	1 p.d. (S.E., nt)	-	0
25. battery cage, DE	0	-	4 p.f. (S.E. PT4; S.I.)	0	-	1 p.d. (S.E. PT4)	-
26. battery cage, DE	0	_	0	0	-	1 p.f. (S.E. PT4)	-
27. battery cage, DE	0	_	3 p.f. (S.E. PT4)	2 c.s. (S.E. PT4)	-	2 p.f. (S.E. PT4)	-
31. barn, DE	0	-	2 p.f., 1 p.d., 1 c.s.	1 p.f., 2 c.s. (S.E.	-	3 p.f., 1 p.d., 1 c.s.	-
			(S.E. PT8)	PT8)		(S.E. PT8)	
32. barn, DE	0	_	0	1 p.d. (S.L.)	_	0	-
33. barn, DE	-	-	5 p.f., 1 p.d., 7 c.s.	2 p.f., 1 p.d. (S.E.		-	-
			(S.E. PT4)	PT4)			
40. small colony, DE	_	_	2 p.f. (S.E. PT4)	0	_	0	-
41. small colony, DE	0	_	1 p.d. (<i>S</i> . group B)	1 p.d. (S. group B)	_	1 p.f., 1 p.d. (S.E.	-
						PT8)	

p.f. = pooled faecal sample; p.d. = pooled dust sample; c.s. = cloacal swab sample; - = no sample was taken in this interval; nt = not typable; S. E. = Salmonella Enteritidis;

S.I. = *Salmonella* Infantis; *S.L.* = *Salmonella* London; PT = phage type



Salmonella prevalence detected in six different sampling intervals throughout a laying period