



## On the toughness of the corona of two graphs

Rocio M. Casablanca, Ana Diáñez, Pedro García-Vázquez

### ► To cite this version:

Rocio M. Casablanca, Ana Diáñez, Pedro García-Vázquez. On the toughness of the corona of two graphs. International Journal of Computer Mathematics, 2011, pp.1. 10.1080/00207160.2011.564277 . hal-00710050


**HAL Id: hal-00710050**

**<https://hal.science/hal-00710050>**

Submitted on 20 Jun 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



### On the toughness of the corona of two graphs

Journal:	<i>International Journal of Computer Mathematics</i>
Manuscript ID:	GCOM-2010-0689-A.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	17-Dec-2010
Complete List of Authors:	Casablanca, Rocio; Universidad de Sevilla, Matematica Aplicada I Diánez, Ana; Universidad de Sevilla, Matemática Aplicada I García-Vázquez, Pedro; Universidad de Sevilla, Matemática Aplicada I
Keywords:	toughness, networks, vulnerability parameter, cutset, connectivity

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**List of corrections**

- (1) Referee 1: Page 4, lines 25-27 and lines 46-48;  $|S| - |S_j|$ ; Referee 2: The authors seem to have set  $|S_j| = n$ , but the explanation is missing. This is corrected in blue color in page 4.
- (2) Referee 1: Page 6, line 48; Then  $S$  is a cutset of  $G \circ H$  since.... This is corrected in blue color in page 6.
- (3) Referee 1: Page 8, line 17:  $S_0 \neq \emptyset$  by Remark 2.1 and  $k \geq 1$ , if..... This is corrected in blue color in page 8.
- (4) Referee 2: Page 9, line 4. "loss" This is corrected in blue color in page 9.
- (5) Referee 2: Page 9, line 6. I think "We may also suppose" sounds better This is corrected in blue color in page 9.
- (6) Referee 1: Page 9, line 25:  $\omega(G \circ H - S) = |S_0|$  (this mistake is also noted by Referee 2), line 29:  $\omega(G \circ H - S) = 1 + |S_0|$ , line 55: there exists an integer... This is corrected in blue color in page 9.
- (7) Referee 2: Page 9, line 11: "Proposition 3.1. Thus," This is corrected in blue color in page 9.
- (8) Referee 2: Page 9, line 12:  $\omega(G \circ H - S_0) \geq 1 \implies \omega_0 \geq 1$  and  $\omega(G \circ H - S_0) = 1 \implies \omega_0 = 1$ ; Page 9, line 13:  $\omega(G \circ H - S_0) \geq 2 \implies \omega_0 \geq 2$ . This is clarified in blue color in page 9.
- (9) Referee 1: Page 10, line 7: " $|S_0| > |S|$ " is not correct, I suggest to delete " $|S_0| > |S|$ " and the word "then", write like "Since ..., we have".; the same mistake is noted by Referee 2. This is corrected in blue color in page 10.
- (10) Referee 1: Page 11, line 6, delete "then", as I know "Since A, then B" is not correct in English. This is corrected in blue color in page 11.
- (11) Referee 1: Page 11, in Table 1, it seems to me that "si" should be "if", please check; the same mistake is noted by Referee 2. This is corrected in blue color in Table 1 of page 11.
- (12) Referee 2: Page 12, line 13: "focus" This is corrected in blue color in page 13.
- (13) Referee 1: There is a paper: A. Mamut and E. Vumar, Vertex vulnerability parameters of Kronecker products of complete graphs, Inform. Process. Lett.106(2008)258-262, I suggest to mention it since it is related to the present paper and this will demonstrate that the present paper deals with a "hot topic". This paper has been added as reference [21].
- (14) Editor: update your Bibliography The Bibliography has been updated by adding references [1,17,19,20,21,22,27].
- (15) We have revised the paper and some words and/or expressions have been modified and marked in red color.

# On the toughness of the corona of two graphs

R. M. Casablanca, A. Diánez, P. García-Vázquez\*

*Departamento de Matemática Aplicada I*

Universidad de Sevilla, Sevilla, Spain

## Abstract

The toughness of a noncomplete graph  $G = (V, E)$  is defined as  $\tau(G) = \min\{|S|/\omega(G - S)\}$ , where the minimum is taken over all cutsets  $S$  of vertices of  $G$  and  $\omega(G - S)$  denotes the number of components of the resultant graph  $G - S$  by deletion of  $S$ . The corona of two graphs  $G$  and  $H$ , written as  $G \circ H$ , is the graph obtained by taking one copy of  $G$  and  $|V(G)|$  copies of  $H$ , and then joining the  $i$ th vertex of  $G$  to every vertex in the  $i$ th copy of  $H$ . In this paper, we investigate the toughness of this kind of graphs and obtain the exact value for the corona of two graphs belonging to some families as paths, cycles, stars, wheels or complete graphs.

**Keywords.** Connectivity, toughness, networks, vulnerability.

## 1 Introduction

Throughout this paper, all the graphs are simple, that is, without loops and multiple edges. Notations and terminology not explicitly given here can be found in the book by Chartrand and Lesniak [9].

Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . The graph  $G$  is called *connected* if every pair of vertices is joined by a path. A *cutset* in a graph  $G$  is a subset  $S \subset V(G)$  of vertices of  $G$  such that  $G - S$  is not connected. The existence of a cutset is always guaranteed in every graph different from a complete graph  $K_n$ . The index of

---

\* rociomc@us.es anadianez@us.es pgvazquez@us.es

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*connectivity* of  $G$ , denoted by  $\kappa(G)$ , is defined as the minimum cardinality over all cutsets of  $G$ , if  $G$  is a non complete graph, or  $|V(G)| - 1$ , otherwise.

There are several measures of vulnerability of a network (see for instance [6, 18, 21, 23]). The vulnerability parameters one generally encounters are the indices of connectivity and edge-connectivity. These two parameters give the minimum cost to disrupt the network, but they do not take into account what remains after destruction. To measure the vulnerability of networks more properly, some vulnerability parameters have been introduced and studied. Among them are toughness [3, 17, 19, 20, 22], integrity [2], neighbor-integrity [1, 26], scattering number [27], tenacity [24] and several variants of connectivity and edge-connectivity called conditional connectivity [5, 7, 11, 12, 16], each of which measures not only the difficulty of breaking down the network but also the effect of the damage. In general, for most of the aforementioned parameters, the corresponding computing problem is *NP*-hard. So it is of interest to give the formulae or algorithms for computing these parameters for special classes of graphs. For our purpose, we deal with the notion of *toughness*, introduced by Chvátal [10], which pays special attention to the relationship between the cardinality of the rupture set in the network and the number of components after rupture. The parameter is defined as

$$\tau(G) = \min \{|S|/\omega(G - S) : S \subseteq J(G)\},$$

where  $J(G) = \{S \subset V(G) : S \text{ is a cutset of } G \text{ or } G - S \text{ is an isolated vertex}\}$ , and  $\omega(G - S)$  denotes the number of components in the resultant graph  $G - S$  by removing  $S$ .

Since this parameter was introduced, a lot of research has been done, mainly relating toughness conditions to the existence of cycle structures. Historically, most of the research was based on a number of conjectures in [10]. Some of most interesting results are addressed in [4, 8, 13]. However, exact values of  $\tau(G)$  are known only for a few families of graphs  $G$ , as paths and cycles [25], the cartesian product of two complete graphs [10] and of paths and/or cycles [15], and the composition of two graphs, one of them being a path, a cycle or a complete bipartite graph [15].

In this paper we focus on the toughness of the corona of two graphs. If for each vertex  $x$  in a graph  $G$ , we introduce a new vertex  $x'$  and join  $x$  and  $x'$  by an edge, the resulting graph is called the *corona* of  $G$ . The operation of adding one vertex for each vertex of  $G$  and connecting them by an edge can be generalized as follows. The corona of any two graphs  $G$  and  $H$ , denoted by  $G \circ H$ , is the graph obtained by taking one copy of  $G$  and  $|V(G)|$  copies of  $H$ , and then joining the  $i$ th vertex of  $G$  to every vertex in the  $i$ th copy

of  $H$ . Observe that the particular case in which  $H = K_1$ , the graph  $G \circ K_1$  is called the corona of  $G$ . This notion was introduced in [14].

There exist several kinds of interconnection networks whose structure can be modeled in terms of the corona of two predetermined networks. This is the case, for instance, of electric networks distributed in a big city where each transformer must guarantee the energy supply of its catchment area. In order to optimize resources, the distribution of transformers is made by dividing the city in catchment areas of the same entity. Thus, in terms of Graph Theory, the structure to be analyzed consists of a network transformers, modeled by a graph  $G$ , where each transformer is connected with its catchment area, modeled by the graph  $H$ . The resultant graph is the corona  $G \circ H$  of  $G$  and  $H$ . In the maintenance of electric networks is relevant to avoid the disruption of the energy supply, but when the failure of some nodes produces the rupture of the network, the greater the number of fragments in which the network has been divided, the greater the cost of reconstruction.

The relationship between the cardinality of a cutset of a graph  $G$  and the remaining component after disruption is analyzed by the notion of *toughness*, defined above. So our aim in this work is to determine the toughness of the corona  $G \circ H$  of two connected graphs  $G$  and  $H$  in terms of known parameters of them. As a consequence, we will deduce the exact value of the corona of some families of graphs involving paths, cycles, stars, wheels or complete graphs.

## 2 Notations and remarks

Let  $G, H$  be two connected graphs on  $m$  and  $n$  vertices, respectively. Let us set  $V(G) = \{v_1, \dots, v_m\}$  and denote by  $H_i$  the copy of  $H$  that is joined to vertex  $v_i$  of  $G$  in  $G \circ H$ . Thus, every cutset  $S$  of  $G \circ H$  will henceforth expressed as  $S = S_0 \cup \bigcup_{i=1}^m S_i$ , where  $S_0 \subseteq V(G)$  and  $S_i \subseteq V(H_i)$ , for  $i = 1, \dots, m$ . We denote by  $\omega_0 = \omega(G - S_0)$ ,  $\omega_i = \omega(H_i - S_i)$ ,  $i = 1, \dots, m$ , that is, the number of component of  $G - S_0$  and  $H_i - S_i$ ,  $i = 1, \dots, m$ , respectively.

A cutset of  $G \circ H$  such that  $|S|/\omega(G \circ H - S) = \tau(G \circ H)$  will be called a  $\tau$ -cut of  $G \circ H$ . Let us see some remarks on the  $\tau$ -cut of the corona of two graphs.

**Remark 2.1** If  $S = S_0 \cup \bigcup_{i=1}^m S_i$  is a cutset of the corona  $G \circ H$  of two connected graphs

$G, H$ , then  $S_0 \neq \emptyset$ .

**Proof.** If  $S_0 = \emptyset$  then every vertex of  $G \circ H - S$  either is in  $V(G)$  or is adjacent to one vertex of  $G$ , hence,  $G \circ H - S$  is connected, against the fact that  $S$  is a cutset of  $G \circ H$ .

■

**Remark 2.2** Let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of the corona  $G \circ H$  of two connected graphs  $G, H$ . If  $v_j \in S_0$  then either  $S_j = \emptyset$  or  $S_j$  is a cutset of  $H_j$ .

**Proof.** Let  $v_j \in S_0$  and suppose by contradiction that  $S_j \neq \emptyset$  is not a cutset of  $H_j$ . Let us consider the set  $S^* = S \setminus S_j$ . Observe that either  $H_j - S_j$  is a component of  $G \circ H - S$  or  $S_j = V(H_j)$  and  $H_j$  is a component of  $G \circ H - S^*$ . Thus,  $\omega(G \circ H - S^*) \geq \omega(G \circ H - S)$  and therefore,

$$\frac{|S^*|}{\omega(G \circ H - S^*)} \leq \frac{|S| - |S_j|}{\omega(G \circ H - S)} < \frac{|S|}{\omega(G \circ H - S)} = \tau(G \circ H - S),$$

which contradicts the hypothesis that  $S$  is a  $\tau$ -cut of  $G \circ H$ . Then either  $S_j = \emptyset$  or  $S_j$  is a cutset of  $H_j$ . ■

**Remark 2.3** Let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of the corona  $G \circ H$  of two connected graphs  $G, H$ . If  $v_j \notin S_0$  then  $S_j = \emptyset$ .

**Proof.** Let  $v_j \notin S_0$  and suppose by contradiction that  $S_j \neq \emptyset$ . Let us consider the set  $S^* = S \setminus S_j$ . Observe that either  $H_j - S_j$  belongs to the component of  $G \circ H - S$  that contains vertex  $v_j$  or  $S_j = V(H_j)$  and  $H_j$  belongs to the component of  $G \circ H - S^*$  that contains vertex  $v_j$ . Thus,  $\omega(G \circ H - S^*) = \omega(G \circ H - S)$  and therefore,

$$\frac{|S^*|}{\omega(G \circ H - S^*)} = \frac{|S| - |S_j|}{\omega(G \circ H - S)} < \frac{|S|}{\omega(G \circ H - S)} = \tau(G \circ H - S),$$

which is again a contradiction with the fact that  $S$  is a  $\tau$ -cut of  $G \circ H$ . Then  $S_j = \emptyset$ . ■

Let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of  $G \circ H$ . From now on, we may assume without loss of generality that the vertices of the set  $V(G) = \{v_1, \dots, v_m\}$  are ordered so that  $|S_1| \geq \dots \geq |S_m|$ . Let  $k \in \{1, \dots, m\}$  be the maximum integer such that  $S_i \neq \emptyset$  for all  $i = 1, \dots, k$ . Then, as an immediate consequence of Remark 2.1, Remark 2.2 and Remark 2.3, it follows that  $|S| = |S_0| + \sum_{i=1}^k |S_i|$  and  $\omega(G \circ H - S) = \omega_0 + \sum_{i=1}^k \omega_i + |S_0| - k$ .

### 3 Main results

Let  $G, H$  be two connected graphs on  $m$  and  $n$  vertices, respectively. Our purpose is to determine the toughness of the corona  $G \circ H$  of  $G$  and  $H$ . To begin with, given a  $\tau$ -cut of  $G \circ H$ , the first question that we must answer is whether every copy of graph  $H$  can be disconnected in the same way. The following lemma provides an answer to this question.

**Lemma 3.1** *Let  $G, H$  be two connected graphs of order  $m$  and  $n$ , respectively, and let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of  $G \circ H$  of minimum cardinality. If  $S_i \neq \emptyset, S_j \neq \emptyset$ , for  $i, j = 1, \dots, m$  with  $i \neq j$ , then  $|S_i| = |S_j|$  and  $\omega_i = \omega_j$ .*

**Proof.** Let us consider the vertex set  $V(G) = \{v_1, \dots, v_m\}$  ordered so that  $|S_1| \geq \dots \geq |S_m|$ , and let  $k \in \{1, \dots, m\}$  be the maximum integer such that  $S_i \neq \emptyset$  for all  $i = 1, \dots, k$ . Thus,  $|S| = |S_0| + \sum_{i=1}^k |S_i|$ . Since  $S$  is a  $\tau$ -cut of  $G \circ H$ , we have

$$\tau(G \circ H) = \frac{|S_0| + \sum_{i=1}^k |S_i|}{\omega_0 + \sum_{i=1}^k \omega_i + |S_0| - k} \leq \frac{|S_0| + k|S_\ell|}{\omega_0 + k\omega_\ell + |S_0| - k}, \quad \text{for every } \ell = 1, \dots, k, \quad (1)$$

yielding that

$$\begin{aligned} & \left( |S_0| + \sum_{i=1}^k |S_i| \right) k\omega_\ell + (\omega_0 + |S_0| - k) \sum_{i=1}^k |S_i| \\ & \leq |S_0| \sum_{i=1}^k \omega_i + \left( \omega_0 + \sum_{i=1}^k \omega_i + |S_0| - k \right) k|S_\ell|, \quad \text{for } \ell = 1, \dots, k. \end{aligned} \quad (2)$$

By taking summation in (2) we deduce that

$$\begin{aligned} & \left( |S_0| + \sum_{i=1}^k |S_i| \right) k \sum_{\ell=1}^k \omega_\ell + k(\omega_0 + |S_0| - k) \sum_{i=1}^k |S_i| \\ & \leq k|S_0| \sum_{i=1}^k \omega_i + \left( \omega_0 + \sum_{i=1}^k \omega_i + |S_0| - k \right) k \sum_{\ell=1}^k |S_\ell| \\ & = \left( |S_0| + \sum_{\ell=1}^k |S_\ell| \right) k \sum_{i=1}^k \omega_i + k(\omega_0 + |S_0| - k) \sum_{\ell=1}^k |S_\ell|, \end{aligned}$$

which implies that all the inequalities of (2) become equalities, and therefore, all the inequalities of (1) become equalities. Thus,

$$\tau(G \circ H) = \frac{|S_0| + k|S_i|}{\omega_0 + k\omega_i + |S_0| - k} = \frac{|S_0| + k|S_j|}{\omega_0 + k\omega_j + |S_0| - k}, \quad \text{for all } i, j = 1, \dots, k, \quad (3)$$



which means that the set  $S^* = S_0 \cup \bigcup_{i=1}^k S_i^*$ , where  $S_i^* = S_k$ , for all  $i = 1, \dots, k$ , is also a  $\tau$ -cut. Hence,

$$|S| = |S_0| + \sum_{i=1}^k |S_i| \geq |S_0| + k|S_k| = |S^*|,$$

leading to  $|S_1| = \dots = |S_k|$  because  $S$  has minimum cardinality. Moreover, given any two subsets  $S_i, S_j$ , with  $i, j \in \{1, \dots, k\}$  and  $i \neq j$ , from (3) it is clear that  $\omega_i = \omega_j$ . Then the result holds. ■

Given a  $\tau$ -cut  $S = S_0 \cup \bigcup_{i=1}^m S_i$  of  $G \circ H$  with minimum cardinality, by Lemma 3.1 we may assume without loss of generality that for each  $i = 1, \dots, m$ , either  $S_i = \emptyset$  or  $S_i = S_H$ , for some  $S_H \subset V(H)$ . Furthermore, it follows that either  $\omega(H_i - S_i) = 1$  (if  $S_i = \emptyset$ ) or  $\omega(H_i - S_i) = \omega(H - S_H)$  (if  $S_i = S_H$ ).

To upper bound the index of toughness of  $G \circ H$ , it is enough to find a cutset  $S$  of  $G \circ H$  and compute  $|S|/\omega(G \circ H - S)$ . There are some alternatives in the choice of such a cutset, as the following proposition shows.

**Proposition 3.1** *Let  $G, H$  be two connected graphs of order  $m$  and  $n$ , respectively. Let  $S_H \subset V(H)$  be any cutset of  $H$  of cardinality  $|S_H| = p$  and denote by  $q = \omega(H - S_H)$ . Then*

$$\tau(G \circ H) \leq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)}, \frac{1+p}{1+q}, \frac{1+p}{\frac{1}{\tau(G)} + q} \right\}.$$

**Proof.** First, let  $v_j$  be any vertex of  $V(G)$  and let us consider the set  $S = \{v_j\}$  in  $G \circ H$ . Then  $S$  is a cutset and  $G \circ H$  since  $v_j$  separates the copy  $H_j$  of  $H$  from  $G \circ H - (\{v_j\} \cup V(H_j))$ . Furthermore,  $G \circ H - S$  has at least two components, i.e.,  $\omega(G \circ H - S) = 1 + \omega(G \circ H - (\{v_j\} \cup V(H_j))) \geq 2$ , yielding that  $\tau(G \circ H) \leq \frac{|S|}{\omega(G \circ H - S)} \leq \frac{1}{2}$ .

Second, let  $S \subset V(G)$  be a  $\tau$ -cut of  $G$ . Then  $S$  is a cutset of  $G \circ H$  and  $\omega(G \circ H - S) = \omega(G - S) + |S|$  and therefore,

$$\tau(G \circ H) \leq \frac{|S|}{\omega(G \circ H - S)} = \frac{|S|}{\omega(G - S) + |S|} = \frac{\frac{|S|}{\omega(G - S)}}{1 + \frac{|S|}{\omega(G - S)}} = \frac{\tau(G)}{1 + \tau(G)}.$$

Third, let  $S_H \subset V(H)$  be any cutset of  $H$  of cardinality  $|S_H| = p$  and denote by  $q = \omega(H - S_H)$ . Take any vertex  $v_j \in V(G)$  and set  $S_j = S_H \subset V(H_j)$ . Let us

consider the vertex set  $S = \{v_j\} \cup S_j$  and observe that  $S$  is a cutset of  $G \circ H$ . Indeed,  $\omega(G \circ H - S) = \omega(G - v_j) + \omega(H_j - S_j) \geq 1 + \omega(H_j - S_j)$ . Thus, it follows that

$$\tau(G \circ H) \leq \frac{|S|}{\omega(G \circ H - S)} \leq \frac{1 + |S_j|}{1 + \omega(H_j - S_j)} = \frac{1 + p}{1 + q}.$$

Finally, take any cutset  $S_H \subset V(H)$  of  $H$  of cardinality  $|S_H| = p$  and denote by  $q = \omega(H - S_H)$ . Let  $S_0 = \{u_1, \dots, u_{|S_0|}\} \subset V(G)$  be a  $\tau$ -cut of  $G$  and denote by  $H_i$  the copy of  $H$  joined to vertex  $u_i$  in  $G \circ H$ , for  $i = 1, \dots, |S_0|$ . Let us consider the vertex set  $S = S_0 \cup \bigcup_{i=1}^{|S_0|} S_i$ , where  $S_i = S_H$ , for every  $i = 1, \dots, |S_0|$ . Clearly  $S$  is a cutset of  $G \circ H$  and  $\omega(G \circ H - S) = \omega(G - S_0) + |S_0|\omega(H - S_H)$ . Hence,

$$\begin{aligned} \tau(G \circ H) &\leq \frac{|S|}{\omega(G \circ H - S)} = \frac{|S_0| + |S_0||S_H|}{\omega(G - S_0) + |S_0|\omega(H - S_H)} \\ &= \frac{|S_0|(1 + p)}{\omega(G - S_0) + |S_0|q} \\ &= \frac{\tau(G)(1 + p)}{1 + \tau(G)q} \\ &= \frac{1 + p}{1/\tau(G) + q}. \end{aligned}$$

Thus,  $\tau(G \circ H) \leq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)}, \frac{1 + p}{1 + q}, \frac{1 + p}{\frac{1}{\tau(G)} + q} \right\}$  and the result holds. ■

The next result gives a necessary condition for a  $\tau$ -cut of  $G \circ H$  to contain vertices of some copy  $H_i$ .

**Lemma 3.2** *Let  $G, H$  be two connected graphs of order  $m$  and  $n$ , respectively, and let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of  $G \circ H$  of minimum cardinality. If  $S_j \neq \emptyset$  for some  $j = 1, \dots, m$ , then  $|S_j|/\omega(H_j - S_j) < 1/2$ .*

**Proof.** From Lemma 3.1 there exists a vertex set  $S_H \subset V(H)$  such that either  $S_i = \emptyset$  or  $S_i = S_H$ , for every  $i = 1, \dots, m$ . So without loss of generality we may assume that there is an integer  $k \in \{1, \dots, m\}$  such that  $S = S_0 \cup \bigcup_{i=1}^k S_H$ ; that is,  $S_i = S_H$  if  $i \in \{1, \dots, k\}$  and  $S_i = \emptyset$  otherwise. Therefore, it is enough to us to prove that  $|S_H|/\omega(H - S_H) < 1/2$ . To clarify expressions, denote by  $\omega_0 = \omega(G - S_0)$  and  $\omega_H = \omega(H - S_H)$ . By applying Remark 2.1, we know that  $S_0 \neq \emptyset$ , and from Remark 2.2 and Remark 2.3 it follows that  $k \leq |S_0|$ . Thus,  $|S| = |S_0| + k|S_H|$  and  $\omega(G \circ H - S) = \omega_0 + k\omega_H + |S_0| - k$ . By applying Proposition 3.1 we know that  $\tau(G \circ H) \leq 1/2$ , which implies that

$$\frac{|S|}{\omega(G \circ H - S)} = \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0| - k} \leq \frac{1}{2},$$

leading to

$$\frac{|S_H|}{\omega_H} \leq \frac{1}{2} + \frac{\omega_0 - (|S_0| + k)}{2k\omega_H}. \quad (4)$$

Since  $S_0 \neq \emptyset$  by Remark 2.1 and  $k \geq 1$ , if  $S_0$  is not a cutset of  $G$  then  $\omega_0 \leq 1$  (i.e.,  $\omega_0 = 0$  if  $S_0 = V(G)$ , and  $\omega_0 = 1$  otherwise). Hence, applying inequality  $\omega_0 - (|S_0| + k) < 0$  in (4), we have  $\frac{|S_H|}{\omega_H} < \frac{1}{2}$ . Thus, suppose that  $S_0 \subset V(G)$  is a cutset of  $G$ .

First assume that  $|S_0|/\omega_0 \geq 1$ . This means that  $\omega_0 - (|S_0| + k) < \omega_0 - |S_0| \leq 0$ , yielding in (4) that  $\frac{|S_H|}{\omega_H} < \frac{1}{2}$ .

Second assume that  $|S_0|/\omega_0 < 1$ . Since  $S_0$  is a cutset of  $G$  then it is also a cutset of  $G \circ H$  and  $\omega(G \circ H - S_0) = \omega_0 + |S_0|$ . Therefore, by using that  $S$  is a  $\tau$ -cut of  $G \circ H$ , it follows that

$$\frac{|S_0|}{\omega_0 + |S_0|} \geq \tau(G \circ H) = \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0| - k} > \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0|}. \quad (5)$$

Combining the first and the last members of (5) we deduce that

$$\frac{|S_H|}{\omega_H} < \frac{|S_0|}{\omega_0 + |S_0|} = \frac{\frac{|S_0|}{\omega_0}}{1 + \frac{|S_0|}{\omega_0}} < \frac{1}{2},$$

because  $|S_0|/\omega_0 < 1$ . This concludes the proof. ■

From these previous results it follows the next theorem where the toughness of the corona  $G \circ H$  of two connected graphs is determined in terms of some parameters of  $G$  and  $H$ .

**Theorem 3.1** *Let  $G, H$  be two connected graphs of order  $m$  and  $n$ , respectively. Then the following assertions hold:*

(i) *If  $\tau(G) \geq 1$  and  $\tau(H) \geq 1/2$ , then  $\tau(G \circ H) = \frac{1}{2}$ .*

(ii) *If  $\tau(G) < 1$  and  $\tau(H) \geq 1/2$ , then  $\tau(G \circ H) = \frac{\tau(G)}{1 + \tau(G)}$ .*

(iii) *If  $\tau(G) \geq 1$  and  $\tau(H) < 1/2$ , then  $\tau(G \circ H) = \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega(H - S_H)} \right\}$ .*

(iv) *If  $\tau(G) < 1$  and  $\tau(H) < 1/2$ , then*

$$\tau(G \circ H) = \min \left\{ \frac{\tau(G)}{1 + \tau(G)}, \min_{S_H \in J(H)} \frac{1 + |S_H|}{\frac{1}{\tau(G)} + \omega(H - S_H)} \right\}.$$

**Proof.** Let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of  $G \circ H$ . Without loss of generality we may assume that  $V(G) = \{v_1, \dots, v_m\}$ , where the vertices are numbered so that  $|S_i| \geq |S_{i+1}|$ , for all  $i = 1, \dots, m-1$ . We may also suppose that  $S$  has minimum cardinality over all the  $\tau$ -cuts of  $G \circ H$ .

First, assume that  $\tau(H) \geq 1/2$ . Then by applying Lemma 3.2 we deduce that  $S_i = \emptyset$ , for all  $i = 1, \dots, m$ , hence,  $S = S_0$ . This implies that  $\tau(G \circ H) = \frac{|S|}{\omega(G \circ H - S)} = \frac{|S_0|}{\omega_0 + |S_0|}$ . Note that  $S_0 \neq V(G)$ , because otherwise,  $\omega_0 = 0$ ; that is,  $\omega(G \circ H - S_0) = |S_0|$  and therefore,  $\tau(G \circ H) = 1$ , which is a contradiction with Proposition 3.1. Thus,  $S_0 \subset V(G)$ , which means that  $\omega_0 \geq 1$ , yielding that  $\omega(G \circ H - S_0) \geq 1 + |S_0|$ . If  $S_0$  is not a cutset of  $G$  then  $\omega_0 = 1$ , and therefore,  $\tau(G \circ H) = \frac{|S_0|}{\omega_0 + |S_0|} = \frac{|S_0|}{1 + |S_0|} \geq \frac{1}{2}$ . If  $S_0$  is a cutset of  $G$  then  $\omega_0 \geq 2$  and therefore,

$$\tau(G \circ H) = \frac{|S_0|}{\omega_0 + |S_0|} = \frac{\frac{|S_0|}{\omega_0}}{1 + \frac{|S_0|}{\omega_0}} \geq \frac{\tau(G)}{1 + \tau(G)}.$$

Hence,  $\tau(G \circ H) \geq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)} \right\}$ . Moreover, by Proposition 3.1 we have  $\tau(G \circ H) \leq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)} \right\}$ , leading to

$$\tau(G \circ H) = \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)} \right\} = \begin{cases} \frac{1}{2}, & \text{if } \tau(G) \geq 1 \\ \frac{\tau(G)}{1 + \tau(G)}, & \text{if } \tau(G) < 1, \end{cases}$$

which proves items (i) and (ii).

Second, assume that  $\tau(H) < 1/2$ . If  $S_1 = \emptyset$  then  $S_i = \emptyset$  for every  $i = 1, \dots, m$ , and reasoning as above, we prove that

$$\tau(G \circ H) \geq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)} \right\} = \begin{cases} \frac{1}{2}, & \text{if } \tau(G) \geq 1 \\ \frac{\tau(G)}{1 + \tau(G)}, & \text{if } \tau(G) < 1. \end{cases} \quad (6)$$

Thus, suppose that  $S_1 \neq \emptyset$ , then by Lemma 3.1 we may assume that there exists an integer  $k \in \{1, \dots, m\}$  and a nonempty vertex set  $S_H \subset V(H)$  such that  $S_i = S_H$  if  $i \leq k$ , and  $S_i = \emptyset$  otherwise. Further, from Lemma 3.2, it follows that  $|S_H|/\omega(H - S_H) < 1/2$ . Again to clarify expressions, denote by  $\omega_H = \omega(H - S_H)$ . Notice that  $k \leq |S_0|$  by Remark 2.2 and Remark 2.3, and therefore,  $\tau(G \circ H) = \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0| - k}$ . Since  $S_0$  is also a cutset of  $G \circ H$  and  $S$  is a  $\tau$ -cut of  $G \circ H$  of minimum cardinality, we have

$$\frac{|S_0|}{\omega_0 + |S_0|} > \tau(G \circ H) = \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0| - k},$$

yielding that

$$|S_H|(\omega_0 + |S_0|) - |S_0|(\omega_H - 1) < 0. \quad (7)$$

The function  $f(k) = \frac{|S_0|+k|S_H|}{\omega_0+k\omega_H+|S_0|-k}$  has derivate

$$\frac{df}{dk} = \frac{|S_H|(\omega_0 + |S_0|) - |S_0|(\omega_H - 1)}{(\omega_0 + k\omega_H + |S_0| - k)^2},$$

and by (7), we deduce that  $f(k)$  is decreasing in  $k$ . Hence,

$$\tau(G \circ H) = f(k) \geq f(|S_0|) = \frac{|S_0|(1 + |S_H|)}{\omega_0 + |S_0|\omega_H}. \quad (8)$$

If  $S_0$  is not a cutset of  $G$  then  $\omega_0 \leq 1$  ( $\omega_0 = 0$  if  $S_0 = V(G)$ , and  $\omega_0 = 1$  otherwise), and from (8) it follows that

$$\begin{aligned} \tau(G \circ H) &\geq \frac{|S_0|(1 + |S_H|)}{1 + |S_0|\omega_H} = \frac{1 + |S_H|}{1/|S_0| + \omega_H} \\ &\geq \frac{1 + |S_H|}{1 + \omega_H} \\ &\geq \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega_H} \right\}. \end{aligned} \quad (9)$$

If  $S_0$  is a cutset of  $G$  then  $|S_0|/\omega_0 \geq \tau(G)$  and therefore, from (8) it follows that

$$\begin{aligned} \tau(G \circ H) &\geq \frac{|S_0|(1 + |S_H|)}{\omega_0 + |S_0|\omega_H} = \frac{1 + |S_H|}{\omega_0/|S_0| + \omega_H} \\ &\geq \frac{1 + |S_H|}{1/\tau(G) + \omega_H} \\ &\geq \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1/\tau(G) + \omega_H} \right\}. \end{aligned} \quad (10)$$

(iii) Suppose that  $\tau(G) \geq 1$ . Then combining (6), (9) and (10), we deduce that

$$\begin{aligned} \tau(G \circ H) &\geq \min \left\{ \frac{1}{2}, \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega_H} \right\}, \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1/\tau(G) + \omega_H} \right\} \right\} \\ &= \min \left\{ \frac{1}{2}, \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega_H} \right\} \right\}. \end{aligned}$$

Since  $\tau(H) < 1/2$ , there exists a cutset  $S_H \subset V(H)$  such that  $|S_H|/\omega_H < 1/2$ , which implies that  $2|S_H| + 1 \leq \omega_H$ . Then

$$\frac{1 + |S_H|}{1 + \omega_H} \leq \frac{\omega_H - |S_H|}{1 + \omega_H} = \frac{1 + \omega_H - (1 + |S_H|)}{1 + \omega_H} = 1 - \frac{1 + |S_H|}{1 + \omega_H},$$

which means that  $\frac{1 + |S_H|}{1 + \omega_H} \leq 1/2$  and therefore,

$$\tau(G \circ H) = \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega_H} \right\}.$$

(iv) Now suppose that  $\tau(G) < 1$ , then from (6), (9) and (10) it follows that

$$\tau(G \circ H) \geq \min \left\{ \frac{\tau(G)}{1 + \tau(G)}, \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1/\tau(G) + \omega_H} \right\} \right\}.$$

■

$\circ$	$S_n$	$P_n$	$C_n$	$W_{1,n}$	$K_n$
$S_m$	$\begin{cases} \frac{1}{m}, & \text{if } m > n - 2, \\ \frac{2}{m+n-2}, & \text{if } m \leq n - 2. \end{cases}$	$\frac{1}{m}$	$\frac{1}{m}$	$\frac{1}{m}$	$\frac{1}{m}$
$P_m$	$\begin{cases} \frac{1}{3}, & \text{if } n < 5, \\ \frac{2}{n+1}, & \text{if } n \geq 5. \end{cases}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$
$C_m$	$\frac{2}{n}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
$W_{1,m}$	$\frac{2}{n}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
$K_m$	$\frac{2}{n}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$

Table 1: The toughness of the corona of some families of graphs.

As a consequence of Theorem 3.1, the toughness of the corona of some families of graphs can be derived. Let  $n \geq 3$  be an integer. Let us denote by  $P_n$  and  $C_n$  the path and the cycle with  $n$  vertices, respectively; by  $S_n$  the complete bipartite graph  $K_{1,n-1}$ ; by  $W_{1,n}$  the wheel with  $n + 1$  vertices; and by  $K_n$  the complete graph of order  $n$ . As a consequence of Theorem 3.1, the toughness of the corona of two graphs, one of them being a complete graph is deduced. Further, in Table 1 we can find the toughness of the corona of two graphs belonging to some of these families: stars, paths, cycles, wheels and complete graphs.

**Corollary 3.1** *Let  $m \geq 3$ ,  $n \geq 3$  be two integers and let  $G, H$  be two connected graphs. Then the following assertions hold:*

$$(i) \quad \tau(G \circ K_n) = \begin{cases} \frac{1}{2}, & \text{if } \tau(G) \geq 1, \\ \frac{\tau(G)}{1+\tau(G)}, & \text{if } \tau(G) < 1. \end{cases}$$

$$(ii) \tau(K_m \circ H) = \begin{cases} \frac{1}{2}, & \text{if } \tau(H) \geq 1/2, \\ \min_{S_H \subset J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega(H - S_H)} \right\}, & \text{if } \tau(H) < 1/2. \end{cases}$$

### 3.1 Conclusions

To measure the vulnerability of networks more properly, there exist some parameters which focus not only the difficulty of breaking down the network but also the effect of the damage. In this work we have studied the toughness of the corona of two graphs. This family models the structure of interconnection networks in several fields, where the reliability is one of most important features. We have determined the value of the toughness of this kind of graphs.

### Acknowledgment

We want to deeply thank the anonymous referees who helped us to improve this paper. This research was supported by the Ministry of Education and Science, Spain, and the European Regional Development Fund (ERDF) under project MTM2008-06620-C03-02/MTM; Catalanian Government 1298 SGR2009. And by the Andalusian Government under project P06-FQM-01649.

### References

- [1] V. Aytaç, *Vulnerability in graphs: the neighbour-integrity of line graphs*, Int. J. Comput. Math. 82(1) (2005), pp. 35–40.
- [2] C.A. Barefoot, R. Entringer, and H. Swart, *Vulnerability in graphs—a comparative survey*, J. Combin. Math. Combin. Comput. 1 (1987), pp. 13–22.
- [3] D. Bauer, H. Broersma, and E. Schmeichel, *Toughness in graphs—a survey*, Graphs Combin. 22(1) (2006), pp. 1–35.
- [4] D. Bauer, S.L. Hakimi, and E. Schmeichel, *Recognizing tough graphs is NP-hard*, Discrete Appl. Math. 28 (1990), pp. 191–195.
- [5] F.T. Boesch, *Synthesis of reliable networks—A survey*, IEEE Trans. Reliability 35 (1986), pp. 240–246.

- [6] F. T. Boesch, A. Satyanarayana, and C. L. Suffel, *A Survey of Some Network Reliability Analysis and Synthesis Results*, Networks 54(2) (2009), pp. 99–107.
- [7] F.T. Boesch and R. Tindell, *Circulants and their connectivities*, J. Graph Theory 8(4) (1984), pp. 487–499.
- [8] H.J. Broersma, E. Engbers, and H. Trommel, *Various results on the toughness of graphs*, Networks 33 (1999), pp. 233–238.
- [9] G. Chartrand and L. Lesniak, *Graphs and Digraphs*, Chapman and Hall/CRC, 2005.
- [10] V. Chvátal, *Tough graphs and Hamiltonian circuits*, Discrete Math. 5 (1973), pp. 215–228.
- [11] J. Fàbrega and M.A. Fiol, *On the extraconnectivity of graphs*, Discrete Math 155 (1996), pp. 49–57.
- [12] J. Fàbrega and M.A. Fiol, *Extraconnectivity of graphs with large girth*, Discrete Math 127 (1994), pp. 163–170.
- [13] R. Faudree, R. Gould, M. Jacobson, L. Lesniak, and A. Saito, *Toughness, degrees and 2-factors*, Discrete Math. 286 (2004), pp. 245–249.
- [14] R. Frucht and F. Harary, *On the corona of two graphs*, Aequationes Math. 4 (1970), pp. 322–325.
- [15] W.D. Goddard and H.C. Swart, *On the toughness of a graph*, Quaestiones Math. 13 (1990), pp. 217–232.
- [16] A. Hellwig and L. Volkmann, *Maximally edge-connected and vertex-connected graphs and digraphs: a survey*, Discrete Math. 308(15) (2008), pp. 3265–3296.
- [17] P. Katerinis and T. Wang, *Toughness of graphs and 2-factors with given properties*, Ars Combin. 95 (2010), pp. 161–177.
- [18] Y. Li, S. Zhang, and Q. Zhang, *Vulnerability parameters of split graphs*, Int. J. Comput. Math. 85(1) (2008), pp. 19–23.
- [19] S. Liu, *On toughness and fractional  $(g, f, n)$ -critical graphs*, Inform. Process. Lett. 110(10) (2010), pp. 378–382.



- [20] S. Liu and J. Cai, *Toughness and existence of fractional  $(g, f)$ -factors in graphs*, Ars Combin. 93 (2009), pp. 305–311.
- [21] A. Mamut and E. Vumar, *Vertex vulnerability parameters of Kronecker products of complete graphs*, Inform. Process. Lett. 106 (2008), pp. 258–262.
- [22] M. Metsidik and E. Vumar, *Toughness and matching extension in  $P_3$ -dominated graphs*, Graphs Combin. 26(3) (2010), pp. 425–432.
- [23] M. Metsidik and E. Vumar, *Edge vulnerability parameters of bisplit graphs*, Comput. Math. Appl. 56(7) (2008), pp. 1741–1747.
- [24] D. Moazzami and B. Salehian, *Some results related to the tenacity and existence of  $k$ -trees*, Discrete Appl. Math. 157(8) (2009), pp. 1794–1798.
- [25] R. E. Pippert, *On the toughness of a graph*, Lecture Notes in Math. 303 (1972), pp. 225–233.
- [26] Z. Wei and S. Zhang, *Vertex-neighbour-integrity of composition graphs of paths and cycles*, Int. J. Comput. Math. 85(5) (2008), pp. 727–733.
- [27] S. Zhang, X. Li, and X. Han, *Computing the scattering number of graphs*, Int. J. Comput. Math. 79(2) (2002), pp. 179–187.

# On the toughness of the corona of two graphs

R. M. Casablanca, A. Diánez, P. García-Vázquez\*

*Departamento de Matemática Aplicada I*

Universidad de Sevilla, Sevilla, Spain

## Abstract

The toughness of a noncomplete graph  $G = (V, E)$  is defined as  $\tau(G) = \min\{|S|/\omega(G - S)\}$ , where the minimum is taken over all cutsets  $S$  of vertices of  $G$  and  $\omega(G - S)$  denotes the number of components of the resultant graph  $G - S$  by deletion of  $S$ . The corona of two graphs  $G$  and  $H$ , written as  $G \circ H$ , is the graph obtained by taking one copy of  $G$  and  $|V(G)|$  copies of  $H$ , and then joining the  $i$ th vertex of  $G$  to every vertex in the  $i$ th copy of  $H$ . In this paper, we investigate the toughness of this kind of graphs and obtain the exact value for the corona of two graphs belonging to some families as paths, cycles, stars, wheels or complete graphs.

**Keywords.** Connectivity, toughness, networks, vulnerability.

## 1 Introduction

Throughout this paper, all the graphs are simple, that is, without loops and multiple edges. Notations and terminology not explicitly given here can be found in the book by Chartrand and Lesniak [9].

Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . The graph  $G$  is called *connected* if every pair of vertices is joined by a path. A *cutset* in a graph  $G$  is a subset  $S \subset V(G)$  of vertices of  $G$  such that  $G - S$  is not connected. The existence of a cutset is always guaranteed in every graph different from a complete graph  $K_n$ . The index of

---

\* rociomc@us.es anadianez@us.es pgvazquez@us.es

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*connectivity* of  $G$ , denoted by  $\kappa(G)$ , is defined as the minimum cardinality over all cutsets of  $G$ , if  $G$  is a non complete graph, or  $|V(G)| - 1$ , otherwise.

There are several measures of vulnerability of a network (see for instance [6, 18, 21, 23]). The vulnerability parameters one generally encounters are the indices of connectivity and edge-connectivity. These two parameters give the minimum cost to disrupt the network, but they do not take into account what remains after destruction. To measure the vulnerability of networks more properly, some vulnerability parameters have been introduced and studied. Among them are toughness [3, 17, 19, 20, 22], integrity [2], neighbor-integrity [1, 26], scattering number [27], tenacity [24] and several variants of connectivity and edge-connectivity called conditional connectivity [5, 7, 11, 12, 16], each of which measures not only the difficulty of breaking down the network but also the effect of the damage. In general, for most of the aforementioned parameters, the corresponding computing problem is *NP*-hard. So it is of interest to give the formulae or algorithms for computing these parameters for special classes of graphs. For our purpose, we deal with the notion of *toughness*, introduced by Chvátal [10], which pays special attention to the relationship between the cardinality of the rupture set in the network and the number of components after rupture. The parameter is defined as

$$\tau(G) = \min \{|S|/\omega(G - S) : S \subseteq J(G)\},$$

where  $J(G) = \{S \subset V(G) : S \text{ is a cutset of } G \text{ or } G - S \text{ is an isolated vertex}\}$ , and  $\omega(G - S)$  denotes the number of components in the resultant graph  $G - S$  by removing  $S$ .

Since this parameter was introduced, a lot of research has been done, mainly relating toughness conditions to the existence of cycle structures. Historically, most of the research was based on a number of conjectures in [10]. Some of most interesting results are addressed in [4, 8, 13]. However, exact values of  $\tau(G)$  are known only for a few families of graphs  $G$ , as paths and cycles [25], the cartesian product of two complete graphs [10] and of paths and/or cycles [15], and the composition of two graphs, one of them being a path, a cycle or a complete bipartite graph [15].

In this paper we focus on the toughness of the corona of two graphs. If for each vertex  $x$  in a graph  $G$ , we introduce a new vertex  $x'$  and join  $x$  and  $x'$  by an edge, the resulting graph is called the *corona* of  $G$ . The operation of adding one vertex for each vertex of  $G$  and connecting them by an edge can be generalized as follows. The corona of any two graphs  $G$  and  $H$ , denoted by  $G \circ H$ , is the graph obtained by taking one copy of  $G$  and  $|V(G)|$  copies of  $H$ , and then joining the  $i$ th vertex of  $G$  to every vertex in the  $i$ th copy

of  $H$ . Observe that the particular case in which  $H = K_1$ , the graph  $G \circ K_1$  is called the corona of  $G$ . This notion was introduced in [14].

There exist several kinds of interconnection networks whose structure can be modeled in terms of the corona of two predetermined networks. This is the case, for instance, of electric networks distributed in a big city where each transformer must guarantee the energy supply of its catchment area. In order to optimize resources, the distribution of transformers is made by dividing the city in catchment areas of the same entity. Thus, in terms of Graph Theory, the structure to be analyzed consists of a network transformers, modeled by a graph  $G$ , where each transformer is connected with its catchment area, modeled by the graph  $H$ . The resultant graph is the corona  $G \circ H$  of  $G$  and  $H$ . In the maintenance of electric networks is relevant to avoid the disruption of the energy supply, but when the failure of some nodes produces the rupture of the network, the greater the number of fragments in which the network has been divided, the greater the cost of reconstruction.

The relationship between the cardinality of a cutset of a graph  $G$  and the remaining component after disruption is analyzed by the notion of *toughness*, defined above. So our aim in this work is to determine the toughness of the corona  $G \circ H$  of two connected graphs  $G$  and  $H$  in terms of known parameters of them. As a consequence, we will deduce the exact value of the corona of some families of graphs involving paths, cycles, stars, wheels or complete graphs.

## 2 Notations and remarks

Let  $G, H$  be two connected graphs on  $m$  and  $n$  vertices, respectively. Let us set  $V(G) = \{v_1, \dots, v_m\}$  and denote by  $H_i$  the copy of  $H$  that is joined to vertex  $v_i$  of  $G$  in  $G \circ H$ . Thus, every cutset  $S$  of  $G \circ H$  will henceforth expressed as  $S = S_0 \cup \bigcup_{i=1}^m S_i$ , where  $S_0 \subseteq V(G)$  and  $S_i \subseteq V(H_i)$ , for  $i = 1, \dots, m$ . We denote by  $\omega_0 = \omega(G - S_0)$ ,  $\omega_i = \omega(H_i - S_i)$ ,  $i = 1, \dots, m$ , that is, the number of component of  $G - S_0$  and  $H_i - S_i$ ,  $i = 1, \dots, m$ , respectively.

A cutset of  $G \circ H$  such that  $|S|/\omega(G \circ H - S) = \tau(G \circ H)$  will be called a  $\tau$ -cut of  $G \circ H$ . Let us see some remarks on the  $\tau$ -cut of the corona of two graphs.

**Remark 2.1** If  $S = S_0 \cup \bigcup_{i=1}^m S_i$  is a cutset of the corona  $G \circ H$  of two connected graphs

$G, H$ , then  $S_0 \neq \emptyset$ .

**Proof.** If  $S_0 = \emptyset$  then every vertex of  $G \circ H - S$  either is in  $V(G)$  or is adjacent to one vertex of  $G$ , hence,  $G \circ H - S$  is connected, against the fact that  $S$  is a cutset of  $G \circ H$ .

■

**Remark 2.2** Let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of the corona  $G \circ H$  of two connected graphs  $G, H$ . If  $v_j \in S_0$  then either  $S_j = \emptyset$  or  $S_j$  is a cutset of  $H_j$ .

**Proof.** Let  $v_j \in S_0$  and suppose by contradiction that  $S_j \neq \emptyset$  is not a cutset of  $H_j$ . Let us consider the set  $S^* = S \setminus S_j$ . Observe that either  $H_j - S_j$  is a component of  $G \circ H - S$  or  $S_j = V(H_j)$  and  $H_j$  is a component of  $G \circ H - S^*$ . Thus,  $\omega(G \circ H - S^*) \geq \omega(G \circ H - S)$  and therefore,

$$\frac{|S^*|}{\omega(G \circ H - S^*)} \leq \frac{|S| - |S_j|}{\omega(G \circ H - S)} < \frac{|S|}{\omega(G \circ H - S)} = \tau(G \circ H - S),$$

which contradicts the hypothesis that  $S$  is a  $\tau$ -cut of  $G \circ H$ . Then either  $S_j = \emptyset$  or  $S_j$  is a cutset of  $H_j$ . ■

**Remark 2.3** Let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of the corona  $G \circ H$  of two connected graphs  $G, H$ . If  $v_j \notin S_0$  then  $S_j = \emptyset$ .

**Proof.** Let  $v_j \notin S_0$  and suppose by contradiction that  $S_j \neq \emptyset$ . Let us consider the set  $S^* = S \setminus S_j$ . Observe that either  $H_j - S_j$  belongs to the component of  $G \circ H - S$  that contains vertex  $v_j$  or  $S_j = V(H_j)$  and  $H_j$  belongs to the component of  $G \circ H - S^*$  that contains vertex  $v_j$ . Thus,  $\omega(G \circ H - S^*) = \omega(G \circ H - S)$  and therefore,

$$\frac{|S^*|}{\omega(G \circ H - S^*)} = \frac{|S| - |S_j|}{\omega(G \circ H - S)} < \frac{|S|}{\omega(G \circ H - S)} = \tau(G \circ H - S),$$

which is again a contradiction with the fact that  $S$  is a  $\tau$ -cut of  $G \circ H$ . Then  $S_j = \emptyset$ . ■

Let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of  $G \circ H$ . From now on, we may assume without loss of generality that the vertices of the set  $V(G) = \{v_1, \dots, v_m\}$  are ordered so that  $|S_1| \geq \dots \geq |S_m|$ . Let  $k \in \{1, \dots, m\}$  be the maximum integer such that  $S_i \neq \emptyset$  for all  $i = 1, \dots, k$ . Then, as an immediate consequence of Remark 2.1, Remark 2.2 and Remark 2.3, it follows that  $|S| = |S_0| + \sum_{i=1}^k |S_i|$  and  $\omega(G \circ H - S) = \omega_0 + \sum_{i=1}^k \omega_i + |S_0| - k$ .

### 3 Main results

Let  $G, H$  be two connected graphs on  $m$  and  $n$  vertices, respectively. Our purpose is to determine the toughness of the corona  $G \circ H$  of  $G$  and  $H$ . To begin with, given a  $\tau$ -cut of  $G \circ H$ , the first question that we must answer is whether every copy of graph  $H$  can be disconnected in the same way. The following lemma provides an answer to this question.

**Lemma 3.1** *Let  $G, H$  be two connected graphs of order  $m$  and  $n$ , respectively, and let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of  $G \circ H$  of minimum cardinality. If  $S_i \neq \emptyset, S_j \neq \emptyset$ , for  $i, j = 1, \dots, m$  with  $i \neq j$ , then  $|S_i| = |S_j|$  and  $\omega_i = \omega_j$ .*

**Proof.** Let us consider the vertex set  $V(G) = \{v_1, \dots, v_m\}$  ordered so that  $|S_1| \geq \dots \geq |S_m|$ , and let  $k \in \{1, \dots, m\}$  be the maximum integer such that  $S_i \neq \emptyset$  for all  $i = 1, \dots, k$ . Thus,  $|S| = |S_0| + \sum_{i=1}^k |S_i|$ . Since  $S$  is a  $\tau$ -cut of  $G \circ H$ , we have

$$\tau(G \circ H) = \frac{|S_0| + \sum_{i=1}^k |S_i|}{\omega_0 + \sum_{i=1}^k \omega_i + |S_0| - k} \leq \frac{|S_0| + k|S_\ell|}{\omega_0 + k\omega_\ell + |S_0| - k}, \quad \text{for every } \ell = 1, \dots, k, \quad (1)$$

yielding that

$$\begin{aligned} & \left( |S_0| + \sum_{i=1}^k |S_i| \right) k\omega_\ell + (\omega_0 + |S_0| - k) \sum_{i=1}^k |S_i| \\ & \leq |S_0| \sum_{i=1}^k \omega_i + \left( \omega_0 + \sum_{i=1}^k \omega_i + |S_0| - k \right) k|S_\ell|, \quad \text{for } \ell = 1, \dots, k. \end{aligned} \quad (2)$$

By taking summation in (2) we deduce that

$$\begin{aligned} & \left( |S_0| + \sum_{i=1}^k |S_i| \right) k \sum_{\ell=1}^k \omega_\ell + k(\omega_0 + |S_0| - k) \sum_{i=1}^k |S_i| \\ & \leq k|S_0| \sum_{i=1}^k \omega_i + \left( \omega_0 + \sum_{i=1}^k \omega_i + |S_0| - k \right) k \sum_{\ell=1}^k |S_\ell| \\ & = \left( |S_0| + \sum_{\ell=1}^k |S_\ell| \right) k \sum_{i=1}^k \omega_i + k(\omega_0 + |S_0| - k) \sum_{\ell=1}^k |S_\ell|, \end{aligned}$$

which implies that all the inequalities of (2) become equalities, and therefore, all the inequalities of (1) become equalities. Thus,

$$\tau(G \circ H) = \frac{|S_0| + k|S_i|}{\omega_0 + k\omega_i + |S_0| - k} = \frac{|S_0| + k|S_j|}{\omega_0 + k\omega_j + |S_0| - k}, \quad \text{for all } i, j = 1, \dots, k, \quad (3)$$

which means that the set  $S^* = S_0 \cup \bigcup_{i=1}^k S_i^*$ , where  $S_i^* = S_k$ , for all  $i = 1, \dots, k$ , is also a  $\tau$ -cut. Hence,

$$|S| = |S_0| + \sum_{i=1}^k |S_i| \geq |S_0| + k|S_k| = |S^*|,$$

leading to  $|S_1| = \dots = |S_k|$  because  $S$  has minimum cardinality. Moreover, given any two subsets  $S_i, S_j$ , with  $i, j \in \{1, \dots, k\}$  and  $i \neq j$ , from (3) it is clear that  $\omega_i = \omega_j$ . Then the result holds. ■

Given a  $\tau$ -cut  $S = S_0 \cup \bigcup_{i=1}^m S_i$  of  $G \circ H$  with minimum cardinality, by Lemma 3.1 we may assume without loss of generality that for each  $i = 1, \dots, m$ , either  $S_i = \emptyset$  or  $S_i = S_H$ , for some  $S_H \subset V(H)$ . Furthermore, it follows that either  $\omega(H_i - S_i) = 1$  (if  $S_i = \emptyset$ ) or  $\omega(H_i - S_i) = \omega(H - S_H)$  (if  $S_i = S_H$ ).

To upper bound the index of toughness of  $G \circ H$ , it is enough to find a cutset  $S$  of  $G \circ H$  and compute  $|S|/\omega(G \circ H - S)$ . There are some alternatives in the choice of such a cutset, as the following proposition shows.

**Proposition 3.1** *Let  $G, H$  be two connected graphs of order  $m$  and  $n$ , respectively. Let  $S_H \subset V(H)$  be any cutset of  $H$  of cardinality  $|S_H| = p$  and denote by  $q = \omega(H - S_H)$ . Then*

$$\tau(G \circ H) \leq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)}, \frac{1+p}{1+q}, \frac{1+p}{\frac{1}{\tau(G)} + q} \right\}.$$

**Proof.** First, let  $v_j$  be any vertex of  $V(G)$  and let us consider the set  $S = \{v_j\}$  in  $G \circ H$ . Then  $S$  is a cutset of  $G \circ H$  since  $v_j$  separates the copy  $H_j$  of  $H$  from  $G \circ H - (\{v_j\} \cup V(H_j))$ . Furthermore,  $G \circ H - S$  has at least two components, i.e.,  $\omega(G \circ H - S) = 1 + \omega(G \circ H - (\{v_j\} \cup V(H_j))) \geq 2$ , yielding that  $\tau(G \circ H) \leq \frac{|S|}{\omega(G \circ H - S)} \leq \frac{1}{2}$ .

Second, let  $S \subset V(G)$  be a  $\tau$ -cut of  $G$ . Then  $S$  is a cutset of  $G \circ H$  and  $\omega(G \circ H - S) = \omega(G - S) + |S|$  and therefore,

$$\tau(G \circ H) \leq \frac{|S|}{\omega(G \circ H - S)} = \frac{|S|}{\omega(G - S) + |S|} = \frac{\frac{|S|}{\omega(G - S)}}{1 + \frac{|S|}{\omega(G - S)}} = \frac{\tau(G)}{1 + \tau(G)}.$$

Third, let  $S_H \subset V(H)$  be any cutset of  $H$  of cardinality  $|S_H| = p$  and denote by  $q = \omega(H - S_H)$ . Take any vertex  $v_j \in V(G)$  and set  $S_j = S_H \subset V(H_j)$ . Let us



consider the vertex set  $S = \{v_j\} \cup S_j$  and observe that  $S$  is a cutset of  $G \circ H$ . Indeed,  $\omega(G \circ H - S) = \omega(G - v_j) + \omega(H_j - S_j) \geq 1 + \omega(H_j - S_j)$ . Thus, it follows that

$$\tau(G \circ H) \leq \frac{|S|}{\omega(G \circ H - S)} \leq \frac{1 + |S_j|}{1 + \omega(H_j - S_j)} = \frac{1 + p}{1 + q}.$$

Finally, take any cutset  $S_H \subset V(H)$  of  $H$  of cardinality  $|S_H| = p$  and denote by  $q = \omega(H - S_H)$ . Let  $S_0 = \{u_1, \dots, u_{|S_0|}\} \subset V(G)$  be a  $\tau$ -cut of  $G$  and denote by  $H_i$  the copy of  $H$  joined to vertex  $u_i$  in  $G \circ H$ , for  $i = 1, \dots, |S_0|$ . Let us consider the vertex set  $S = S_0 \cup \bigcup_{i=1}^{|S_0|} S_i$ , where  $S_i = S_H$ , for every  $i = 1, \dots, |S_0|$ . Clearly  $S$  is a cutset of  $G \circ H$  and  $\omega(G \circ H - S) = \omega(G - S_0) + |S_0|\omega(H - S_H)$ . Hence,

$$\begin{aligned} \tau(G \circ H) &\leq \frac{|S|}{\omega(G \circ H - S)} = \frac{|S_0| + |S_0||S_H|}{\omega(G - S_0) + |S_0|\omega(H - S_H)} \\ &= \frac{|S_0|(1 + p)}{\omega(G - S_0) + |S_0|q} \\ &= \frac{\tau(G)(1 + p)}{1 + \tau(G)q} \\ &= \frac{1 + p}{1/\tau(G) + q}. \end{aligned}$$

Thus,  $\tau(G \circ H) \leq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)}, \frac{1 + p}{1 + q}, \frac{1 + p}{\frac{1}{\tau(G)} + q} \right\}$  and the result holds. ■

The next result gives a necessary condition for a  $\tau$ -cut of  $G \circ H$  to contain vertices of some copy  $H_i$ .

**Lemma 3.2** *Let  $G, H$  be two connected graphs of order  $m$  and  $n$ , respectively, and let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of  $G \circ H$  of minimum cardinality. If  $S_j \neq \emptyset$  for some  $j = 1, \dots, m$ , then  $|S_j|/\omega(H_j - S_j) < 1/2$ .*

**Proof.** From Lemma 3.1 there exists a vertex set  $S_H \subset V(H)$  such that either  $S_i = \emptyset$  or  $S_i = S_H$ , for every  $i = 1, \dots, m$ . So without loss of generality we may assume that there is an integer  $k \in \{1, \dots, m\}$  such that  $S = S_0 \cup \bigcup_{i=1}^k S_H$ ; that is,  $S_i = S_H$  if  $i \in \{1, \dots, k\}$  and  $S_i = \emptyset$  otherwise. Therefore, it is enough to us to prove that  $|S_H|/\omega(H - S_H) < 1/2$ . To clarify expressions, denote by  $\omega_0 = \omega(G - S_0)$  and  $\omega_H = \omega(H - S_H)$ . By applying Remark 2.1, we know that  $S_0 \neq \emptyset$ , and from Remark 2.2 and Remark 2.3 it follows that  $k \leq |S_0|$ . Thus,  $|S| = |S_0| + k|S_H|$  and  $\omega(G \circ H - S) = \omega_0 + k\omega_H + |S_0| - k$ . By applying Proposition 3.1 we know that  $\tau(G \circ H) \leq 1/2$ , which implies that

$$\frac{|S|}{\omega(G \circ H - S)} = \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0| - k} \leq \frac{1}{2},$$



leading to

$$\frac{|S_H|}{\omega_H} \leq \frac{1}{2} + \frac{\omega_0 - (|S_0| + k)}{2k\omega_H}. \quad (4)$$

Since  $S_0 \neq \emptyset$  by Remark 2.1 and  $k \geq 1$ , if  $S_0$  is not a cutset of  $G$  then  $\omega_0 \leq 1$  (i.e.,  $\omega_0 = 0$  if  $S_0 = V(G)$ , and  $\omega_0 = 1$  otherwise). Hence, applying inequality  $\omega_0 - (|S_0| + k) < 0$  in (4), we have  $\frac{|S_H|}{\omega_H} < \frac{1}{2}$ . Thus, suppose that  $S_0 \subset V(G)$  is a cutset of  $G$ .

First assume that  $|S_0|/\omega_0 \geq 1$ . This means that  $\omega_0 - (|S_0| + k) < \omega_0 - |S_0| \leq 0$ , yielding in (4) that  $\frac{|S_H|}{\omega_H} < \frac{1}{2}$ .

Second assume that  $|S_0|/\omega_0 < 1$ . Since  $S_0$  is a cutset of  $G$  then it is also a cutset of  $G \circ H$  and  $\omega(G \circ H - S_0) = \omega_0 + |S_0|$ . Therefore, by using that  $S$  is a  $\tau$ -cut of  $G \circ H$ , it follows that

$$\frac{|S_0|}{\omega_0 + |S_0|} \geq \tau(G \circ H) = \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0| - k} > \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0|}. \quad (5)$$

Combining the first and the last members of (5) we deduce that

$$\frac{|S_H|}{\omega_H} < \frac{|S_0|}{\omega_0 + |S_0|} = \frac{\frac{|S_0|}{\omega_0}}{1 + \frac{|S_0|}{\omega_0}} < \frac{1}{2},$$

because  $|S_0|/\omega_0 < 1$ . This concludes the proof. ■

From these previous results it follows the next theorem where the toughness of the corona  $G \circ H$  of two connected graphs is determined in terms of some parameters of  $G$  and  $H$ .

**Theorem 3.1** *Let  $G, H$  be two connected graphs of order  $m$  and  $n$ , respectively. Then the following assertions hold:*

(i) *If  $\tau(G) \geq 1$  and  $\tau(H) \geq 1/2$ , then  $\tau(G \circ H) = \frac{1}{2}$ .*

(ii) *If  $\tau(G) < 1$  and  $\tau(H) \geq 1/2$ , then  $\tau(G \circ H) = \frac{\tau(G)}{1 + \tau(G)}$ .*

(iii) *If  $\tau(G) \geq 1$  and  $\tau(H) < 1/2$ , then  $\tau(G \circ H) = \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega(H - S_H)} \right\}$ .*

(iv) *If  $\tau(G) < 1$  and  $\tau(H) < 1/2$ , then*

$$\tau(G \circ H) = \min \left\{ \frac{\tau(G)}{1 + \tau(G)}, \min_{S_H \in J(H)} \frac{1 + |S_H|}{\frac{1}{\tau(G)} + \omega(H - S_H)} \right\}.$$

**Proof.** Let  $S = S_0 \cup \bigcup_{i=1}^m S_i$  be a  $\tau$ -cut of  $G \circ H$ . Without loss of generality we may assume that  $V(G) = \{v_1, \dots, v_m\}$ , where the vertices are numbered so that  $|S_i| \geq |S_{i+1}|$ , for all  $i = 1, \dots, m-1$ . We may also suppose that  $S$  has minimum cardinality over all the  $\tau$ -cuts of  $G \circ H$ .

First, assume that  $\tau(H) \geq 1/2$ . Then by applying Lemma 3.2 we deduce that  $S_i = \emptyset$ , for all  $i = 1, \dots, m$ , hence,  $S = S_0$ . This implies that  $\tau(G \circ H) = \frac{|S|}{\omega(G \circ H - S)} = \frac{|S_0|}{\omega_0 + |S_0|}$ . Note that  $S_0 \neq V(G)$ , because otherwise,  $\omega_0 = 0$ ; that is,  $\omega(G \circ H - S_0) = |S_0|$  and therefore,  $\tau(G \circ H) = 1$ , which is a contradiction with Proposition 3.1. Thus,  $S_0 \subset V(G)$ , which means that  $\omega_0 \geq 1$ , yielding that  $\omega(G \circ H - S_0) \geq 1 + |S_0|$ . If  $S_0$  is not a cutset of  $G$  then  $\omega_0 = 1$ , and therefore,  $\tau(G \circ H) = \frac{|S_0|}{\omega_0 + |S_0|} = \frac{|S_0|}{1 + |S_0|} \geq \frac{1}{2}$ . If  $S_0$  is a cutset of  $G$  then  $\omega_0 \geq 2$  and therefore,

$$\tau(G \circ H) = \frac{|S_0|}{\omega_0 + |S_0|} = \frac{\frac{|S_0|}{\omega_0}}{1 + \frac{|S_0|}{\omega_0}} \geq \frac{\tau(G)}{1 + \tau(G)}.$$

Hence,  $\tau(G \circ H) \geq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)} \right\}$ . Moreover, by Proposition 3.1 we have  $\tau(G \circ H) \leq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)} \right\}$ , leading to

$$\tau(G \circ H) = \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)} \right\} = \begin{cases} \frac{1}{2}, & \text{if } \tau(G) \geq 1 \\ \frac{\tau(G)}{1 + \tau(G)}, & \text{if } \tau(G) < 1, \end{cases}$$

which proves items (i) and (ii).

Second, assume that  $\tau(H) < 1/2$ . If  $S_1 = \emptyset$  then  $S_i = \emptyset$  for every  $i = 1, \dots, m$ , and reasoning as above, we prove that

$$\tau(G \circ H) \geq \min \left\{ \frac{1}{2}, \frac{\tau(G)}{1 + \tau(G)} \right\} = \begin{cases} \frac{1}{2}, & \text{if } \tau(G) \geq 1 \\ \frac{\tau(G)}{1 + \tau(G)}, & \text{if } \tau(G) < 1. \end{cases} \quad (6)$$

Thus, suppose that  $S_1 \neq \emptyset$ , then by Lemma 3.1 we may assume that there exists an integer  $k \in \{1, \dots, m\}$  and a nonempty vertex set  $S_H \subset V(H)$  such that  $S_i = S_H$  if  $i \leq k$ , and  $S_i = \emptyset$  otherwise. Further, from Lemma 3.2, it follows that  $|S_H|/\omega(H - S_H) < 1/2$ . Again to clarify expressions, denote by  $\omega_H = \omega(H - S_H)$ . Notice that  $k \leq |S_0|$  by Remark 2.2 and Remark 2.3, and therefore,  $\tau(G \circ H) = \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0| - k}$ . Since  $S_0$  is also a cutset of  $G \circ H$  and  $S$  is a  $\tau$ -cut of  $G \circ H$  of minimum cardinality, we have

$$\frac{|S_0|}{\omega_0 + |S_0|} > \tau(G \circ H) = \frac{|S_0| + k|S_H|}{\omega_0 + k\omega_H + |S_0| - k},$$

yielding that

$$|S_H|(\omega_0 + |S_0|) - |S_0|(\omega_H - 1) < 0. \quad (7)$$

The function  $f(k) = \frac{|S_0|+k|S_H|}{\omega_0+k\omega_H+|S_0|-k}$  has derivate

$$\frac{df}{dk} = \frac{|S_H|(\omega_0 + |S_0|) - |S_0|(\omega_H - 1)}{(\omega_0 + k\omega_H + |S_0| - k)^2},$$

and by (7), we deduce that  $f(k)$  is decreasing in  $k$ . Hence,

$$\tau(G \circ H) = f(k) \geq f(|S_0|) = \frac{|S_0|(1 + |S_H|)}{\omega_0 + |S_0|\omega_H}. \quad (8)$$

If  $S_0$  is not a cutset of  $G$  then  $\omega_0 \leq 1$  ( $\omega_0 = 0$  if  $S_0 = V(G)$ , and  $\omega_0 = 1$  otherwise), and from (8) it follows that

$$\begin{aligned} \tau(G \circ H) &\geq \frac{|S_0|(1 + |S_H|)}{1 + |S_0|\omega_H} = \frac{1 + |S_H|}{1/|S_0| + \omega_H} \\ &\geq \frac{1 + |S_H|}{1 + \omega_H} \\ &\geq \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega_H} \right\}. \end{aligned} \quad (9)$$

If  $S_0$  is a cutset of  $G$  then  $|S_0|/\omega_0 \geq \tau(G)$  and therefore, from (8) it follows that

$$\begin{aligned} \tau(G \circ H) &\geq \frac{|S_0|(1 + |S_H|)}{\omega_0 + |S_0|\omega_H} = \frac{1 + |S_H|}{\omega_0/|S_0| + \omega_H} \\ &\geq \frac{1 + |S_H|}{1/\tau(G) + \omega_H} \\ &\geq \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1/\tau(G) + \omega_H} \right\}. \end{aligned} \quad (10)$$

(iii) Suppose that  $\tau(G) \geq 1$ . Then combining (6), (9) and (10), we deduce that

$$\begin{aligned} \tau(G \circ H) &\geq \min \left\{ \frac{1}{2}, \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega_H} \right\}, \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1/\tau(G) + \omega_H} \right\} \right\} \\ &= \min \left\{ \frac{1}{2}, \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega_H} \right\} \right\}. \end{aligned}$$

Since  $\tau(H) < 1/2$ , there exists a cutset  $S_H \subset V(H)$  such that  $|S_H|/\omega_H < 1/2$ , which implies that  $2|S_H| + 1 \leq \omega_H$ . Then

$$\frac{1 + |S_H|}{1 + \omega_H} \leq \frac{\omega_H - |S_H|}{1 + \omega_H} = \frac{1 + \omega_H - (1 + |S_H|)}{1 + \omega_H} = 1 - \frac{1 + |S_H|}{1 + \omega_H},$$

which means that  $\frac{1 + |S_H|}{1 + \omega_H} \leq 1/2$  and therefore,

$$\tau(G \circ H) = \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega_H} \right\}.$$

(iv) Now suppose that  $\tau(G) < 1$ , then from (6), (9) and (10) it follows that

$$\tau(G \circ H) \geq \min \left\{ \frac{\tau(G)}{1 + \tau(G)}, \min_{S_H \in J(H)} \left\{ \frac{1 + |S_H|}{1/\tau(G) + \omega_H} \right\} \right\}.$$

■

$\circ$	$S_n$	$P_n$	$C_n$	$W_{1,n}$	$K_n$
$S_m$	$\begin{cases} \frac{1}{m}, & \text{if } m > n - 2, \\ \frac{2}{m+n-2}, & \text{if } m \leq n - 2. \end{cases}$	$\frac{1}{m}$	$\frac{1}{m}$	$\frac{1}{m}$	$\frac{1}{m}$
$P_m$	$\begin{cases} \frac{1}{3}, & \text{if } n < 5, \\ \frac{2}{n+1}, & \text{if } n \geq 5. \end{cases}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$
$C_m$	$\frac{2}{n}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
$W_{1,m}$	$\frac{2}{n}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
$K_m$	$\frac{2}{n}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$

Table 1: The toughness of the corona of some families of graphs.

As a consequence of Theorem 3.1, the toughness of the corona of some families of graphs can be derived. Let  $n \geq 3$  be an integer. Let us denote by  $P_n$  and  $C_n$  the path and the cycle with  $n$  vertices, respectively; by  $S_n$  the complete bipartite graph  $K_{1,n-1}$ ; by  $W_{1,n}$  the wheel with  $n + 1$  vertices; and by  $K_n$  the complete graph of order  $n$ . As a consequence of Theorem 3.1, the toughness of the corona of two graphs, one of them being a complete graph is deduced. Further, in Table 1 we can find the toughness of the corona of two graphs belonging to some of these families: stars, paths, cycles, wheels and complete graphs.

**Corollary 3.1** *Let  $m \geq 3$ ,  $n \geq 3$  be two integers and let  $G$ ,  $H$  be two connected graphs. Then the following assertions hold:*

$$(i) \quad \tau(G \circ K_n) = \begin{cases} \frac{1}{2}, & \text{if } \tau(G) \geq 1, \\ \frac{\tau(G)}{1+\tau(G)}, & \text{if } \tau(G) < 1. \end{cases}$$

$$(ii) \tau(K_m \circ H) = \begin{cases} \frac{1}{2}, & \text{if } \tau(H) \geq 1/2, \\ \min_{S_H \subset J(H)} \left\{ \frac{1 + |S_H|}{1 + \omega(H - S_H)} \right\}, & \text{if } \tau(H) < 1/2. \end{cases}$$

### 3.1 Conclusions

To measure the vulnerability of networks more properly, there exist some parameters which focus not only the difficulty of breaking down the network but also the effect of the damage. In this work we have studied the toughness of the corona of two graphs. This family models the structure of interconnection networks in several fields, where the reliability is one of most important features. We have determined the value of the toughness of this kind of graphs.

### Acknowledgment

We want to deeply thank the anonymous referees who helped us to improve this paper. This research was supported by the Ministry of Education and Science, Spain, and the European Regional Development Fund (ERDF) under project MTM2008-06620-C03-02/MTM; Catalanian Government 1298 SGR2009. And by the Andalusian Government under project P06-FQM-01649.

### References

- [1] V. Ayta, *Vulnerability in graphs: the neighbour-integrity of line graphs*, Int. J. Comput. Math. 82(1) (2005), pp. 35–40.
- [2] C.A. Barefoot, R. Entringer, and H. Swart, *Vulnerability in graphs—a comparative survey*, J. Combin. Math. Combin. Comput. 1 (1987), pp. 13–22.
- [3] D. Bauer, H. Broersma, and E. Schmeichel, *Toughness in graphs—a survey*, Graphs Combin. 22(1) (2006), pp. 1–35.
- [4] D. Bauer, S.L. Hakimi, and E. Schmeichel, *Recognizing tough graphs is NP-hard*, Discrete Appl. Math. 28 (1990), pp. 191–195.
- [5] F.T. Boesch, *Synthesis of reliable networks—A survey*, IEEE Trans. Reliability 35 (1986), pp. 240–246.

- [6] F. T. Boesch, A. Satyanarayana, and C. L. Suffel, *A Survey of Some Network Reliability Analysis and Synthesis Results*, Networks 54(2) (2009), pp. 99–107.
- [7] F.T. Boesch and R. Tindell, *Circulants and their connectivities*, J. Graph Theory 8(4) (1984), pp. 487–499.
- [8] H.J. Broersma, E. Engbers, and H. Trommel, *Various results on the toughness of graphs*, Networks 33 (1999), pp. 233–238.
- [9] G. Chartrand and L. Lesniak, *Graphs and Digraphs*, Chapman and Hall/CRC, 2005.
- [10] V. Chvátal, *Tough graphs and Hamiltonian circuits*, Discrete Math. 5 (1973), pp. 215–228.
- [11] J. Fàbrega and M.A. Fiol, *On the extraconnectivity of graphs*, Discrete Math 155 (1996), pp. 49–57.
- [12] J. Fàbrega and M.A. Fiol, *Extraconnectivity of graphs with large girth*, Discrete Math 127 (1994), pp. 163–170.
- [13] R. Faudree, R. Gould, M. Jacobson, L. Lesniak, and A. Saito, *Toughness, degrees and 2-factors*, Discrete Math. 286 (2004), pp. 245–249.
- [14] R. Frucht and F. Harary, *On the corona of two graphs*, Aequationes Math. 4 (1970), pp. 322–325.
- [15] W.D. Goddard and H.C. Swart, *On the toughness of a graph*, Quaestiones Math. 13 (1990), pp. 217–232.
- [16] A. Hellwig and L. Volkmann, *Maximally edge-connected and vertex-connected graphs and digraphs: a survey*, Discrete Math. 308(15) (2008), pp. 3265–3296.
- [17] P. Katerinis and T. Wang, *Toughness of graphs and 2-factors with given properties*, Ars Combin. 95 (2010), pp. 161–177.
- [18] Y. Li, S. Zhang, and Q. Zhang, *Vulnerability parameters of split graphs*, Int. J. Comput. Math. 85(1) (2008), pp. 19–23.
- [19] S. Liu, *On toughness and fractional  $(g, f, n)$ -critical graphs*, Inform. Process. Lett. 110(10) (2010), pp. 378–382.

- [20] S. Liu and J. Cai, *Toughness and existence of fractional  $(g, f)$ -factors in graphs*, Ars Combin. 93 (2009), pp. 305–311.
- [21] A. Mamut and E. Vumar, *Vertex vulnerability parameters of Kronecker products of complete graphs*, Inform. Process. Lett. 106 (2008), pp. 258–262.
- [22] M. Metsidik and E. Vumar, *Toughness and matching extension in  $P_3$ -dominated graphs*, Graphs Combin. 26(3) (2010), pp. 425–432.
- [23] M. Metsidik and E. Vumar, *Edge vulnerability parameters of bisplit graphs*, Comput. Math. Appl. 56(7) (2008), pp. 1741–1747.
- [24] D. Moazzami and B. Salehian, *Some results related to the tenacity and existence of  $k$ -trees*, Discrete Appl. Math. 157(8) (2009), pp. 1794–1798.
- [25] R. E. Pippert, *On the toughness of a graph*, Lecture Notes in Math. 303 (1972), pp. 225–233.
- [26] Z. Wei and S. Zhang, *Vertex-neighbour-integrity of composition graphs of paths and cycles*, Int. J. Comput. Math. 85(5) (2008), pp. 727–733.
- [27] S. Zhang, X. Li, and X. Han, *Computing the scattering number of graphs*, Int. J. Comput. Math. 79(2) (2002), pp. 179–187.