



HAL
open science

Infectious bronchitis virus variants - History, current situation and control measures

Sjaak de Wit, Jane K A Cook, Harold M.J.F. van Der Heijden

► To cite this version:

Sjaak de Wit, Jane K A Cook, Harold M.J.F. van Der Heijden. Infectious bronchitis virus variants - History, current situation and control measures. *Avian Pathology*, 2011, 40 (03), pp.223-235. 10.1080/03079457.2011.566260 . hal-00710052

HAL Id: hal-00710052

<https://hal.science/hal-00710052>

Submitted on 20 Jun 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Infectious bronchitis virus variants – History, current situation and control measures

Journal:	<i>Avian Pathology</i>
Manuscript ID:	CAVP-2010-0187.R1
Manuscript Type:	Review
Date Submitted by the Author:	06-Feb-2011
Complete List of Authors:	de Wit, Sjaak; GD Deventer, Laboratory Cook, Jane Van der Heijden, Harold; GD, R&D
Keywords:	IBV , history, control, variants

SCHOLARONE™
Manuscripts

Cavp-2010-0187.R1

Review article

Infectious bronchitis virus variants: a review of the history, current situation and control measures

J.J. (Sjaak) de Wit^{1,*}, Jane K.A. Cook² and Harold M.J.F. van der Heijden¹

¹ GD (Animal Health Service), P.O.B. 9, 7400 AA, Deventer, The Netherlands

² 138 Hartford Road, Huntingdon, Cambs. PE29 1XQ, UK

NOTE: FIG 1 and TABLE 1 are included in manuscript Figs, 2,3,4, & 5 are separate

Short title: IBV variants: a review

Part of the contents of this paper were published in the Proceedings of the Workshop on Infectious Bronchitis (IB) in the Brazilian Poultry Industry. *Brazilian Journal of Poultry Science*, (2010). 12, 97-106.

* to whom correspondence should be addressed. Tel: + 31 570 660178. Email:

j.d.wit@gddeventer.com

Received: 6 February 2010

For Peer Review Only

Abstract

The history, current situation and control measures for infectious bronchitis virus (IBV) variants are reviewed. A large number of IBV variants exist worldwide; some being unique to a particular area, others having a more general distribution. The possible reasons why some strains spread readily over major parts of the world, whereas other strains stay more localised are discussed. The advantages and disadvantages of strain classification by protectotyping, serotyping and genotyping are discussed in relation to *in vivo* protection. The different vaccination strategies are also considered.

Introduction

Infectious bronchitis virus (IBV) is ubiquitous in most parts of the world where poultry are reared and is able to spread very rapidly in non-protected birds. It is shed via both the respiratory tract and the faeces and can persist in the birds and the intestinal tract for several weeks or months. Outdoors, survival of IBV for 56 days in litter has been reported (Cavanagh & Gelb, 2008). Although strict biosecurity and working with a one-age system are essential control measures, normally vaccination is required to increase the resistance of the chickens against challenge with IBV strains. This is made more difficult to achieve because IBV exists in the form of many different antigenic or genotypic types, commonly referred to as variants. Being a coronavirus and therefore a single stranded RNA virus, IBV has an enormous capacity to change both by spontaneous mutation and by genetic recombination (Cavanagh & Gelb, 2008). When these events occur with IBV both are most likely to result in the emergence of new variants if they occur in the hypervariable regions of the spike gene. Whilst many new variants are unable to replicate or survive for only a short time, a few emerge which become of economic importance either world wide or in restricted geographic areas (see below).

This paper presents an overview of the history and current situation of IBV worldwide, including the measures necessary to control the many IB variants found throughout the world.

History of infectious bronchitis variants. For many years it was widely believed that the first variants of IBV occurred in the early 1950s when Jungherr *et al.* (1956) in the USA showed that the Connecticut (Conn) isolate of 1951 neither cross-neutralised nor cross-protected with the original Massachusetts (Mass) isolate from the early 1940s. However, a retrospective study (Jia *et al.*, 2002), using monoclonal antibodies (mAbs) and molecular

analysis of part of the S1 subunit of the spike glycoprotein (S) gene, identified non-Mass IBVs among isolates made in the USA as early as the 1940s. Although some of these are found elsewhere, most countries are now known to have their own indigenous variants as well, and this paper will consider the current situation worldwide.

USA. In the USA a number of different infectious bronchitis (IB) variants had been identified by the early 1970s, generally on the basis of serological analysis (Hitchner *et al.*, 1966; Hopkins, 1974; Cowen & Hitchner, 1975; Johnson & Marquardt, 1976). Hofstad (1961; 1981) showed a poor level of cross-protection between several of them. New variants continued to be identified and poor cross-protection shown (Gelb *et al.*, 1981). In a survey using a reverse transcriptase polymerase chain reaction (RT-PCR) for typing, Jackwood *et al.* (2005) identified 82 different IB variants among 1523 submissions to their laboratory over an 11-year period. As the number of variants identified in USA increased, a few were shown to be both widely distributed and of major economic importance. Possibly the most significant of these was the variant known as Arkansas [Ark] (Fields, 1973); its importance quickly being recognised by the development of live-attenuated vaccines from this strain (Gelb & Cloud, 1983). Vaccines developed from Mass, Conn and Ark IBVs are widely used in the USA, either singly or in combination (for example a combined Mass-Conn commercial product) and these vaccines have also proved beneficial in providing adequate cross-protection against other IB variants including California 99 (Alvarado *et al.*, 2003), thereby indicating the possibility of cross-protection between IBVs. The Ark serotype continued to be of economic importance in the main poultry producing areas, including Georgia and Delmarva (Gelb *et al.*, 1983). By the mid 1990s, the availability of molecular diagnostic methods led to the identification of subtypes or “quasispecies” within the Ark serotype (Nix *et al.*, 2000; Jackwood *et al.*, 2005). However, suggestions that the currently available Ark

vaccines were providing incomplete protection against more recent isolates of this serotype were discounted by Sander *et al.* (1997) who demonstrated that poor protection was more likely to be due to poor vaccine application than to antigenic drift in the virus in the field. This finding, supported by the recent work of Jackwood *et al.* (2009), confirms the importance of careful application of IB vaccines.

Another IB variant of importance in the USA, Delaware (DE) 072, first reported in the north east of the country in 1992 (Gelb *et al.*, 1997), was found to show little genomic relatedness in the S1 region of the S-gene to other US variants, but interestingly, to be closely related to the Dutch variant, D1466 (Lee & Jackwood, 2001b) (see below). DE072 increased in incidence throughout the next decade, also causing major disease problems in vaccinated flocks in Georgia (Lee *et al.*, 2001). It has apparently undergone both genetic drift and recombination (Lee & Jackwood, 2000, 2001a), such that DE072 vaccines provide only poor protection against field strains, and molecular analysis of field isolates has led to the recognition of a new IB variant, designated Georgia 98 (GA98) (Lee *et al.*, 2001). A homologous vaccine was developed from the Georgia 98 (GA98) strain that provided adequate protection after two applications against a homologous challenge but also against a DE072 challenge (Jackwood *et al.*, 2003). Recently, a new IBV variant designated GA08, for which existing vaccines were not protective, was detected in the same region (Jackwood *et al.*, 2010). Genetic analyses, serotyping and cross-protection trials with strains from California have now shown that different parts of the USA can have their own IBV variants and have confirmed that new ones continue to emerge (Jackwood *et al.*, 2007).

Latin America. IBV had appeared in Latin America by the 1950s and the first reported isolate was of the Mass serotype in Brazil (Hipólito, 1957), although isolation of a variant (Ark) was not reported in that country until some 10 years later (Branden & Da Silva, 1986).

In a study carried out in the mid 1990s, IBV isolates of at least five different antigenic types were found in commercial chickens of all types throughout Brazil, but mainly in the major poultry producing area of the south (Di Fabio *et al.*, 2000). Several different genotypes have now been identified in Brazil by analysis of either the S1 (Montassier *et al.*, 2006; Villarreal *et al.*, 2007a; Villarreal *et al.*, 2007b; Montassier *et al.*, 2008) or the nucleoprotein (N) gene (Abreu *et al.*, 2006), but protection studies have not been performed. More recently it has been shown (Villarreal *et al.*, 2010) that both a unique cluster of IBV strains (subdivided into three subclusters) as well as the 4/91 genotype currently coexist in that country. It is important to remember that in Brazil, as in many other parts of the world, the only live attenuated IB vaccines licensed for use are of the Mass serotype and that protection studies (Cook *et al.*, 1999; Di Fabio *et al.*, 2000) showed that the currently available Mass vaccine provided inadequate protection against some of the Brazilian variants.

Hidalgo *et al.* (1976) reported the first isolation of IB (Mass serotype) in Chile in 1975 and variants were reported for the first time there some 10 years later (Hidalgo *et al.*, 1986). By the mid 1980s, IB was reported as a serious problem in commercial chicken flocks and novel variants as well as the Mass and Conn serotypes were isolated from broiler and layer flocks and again, Mass vaccines were found to protect poorly against challenge with these variants (Cubillos *et al.*, 1991). Elsewhere in Latin America a variant IBV was isolated from commercial chickens in Honduras in 1997 and poor protection against Mass vaccines was demonstrated (Cook *et al.*, 1999). Following the isolation of the Ark serotype in Mexico in the early 1990s (Quiroz *et al.*, 1993), the use of molecular methods subsequently identified IBV variants unique to that country in commercial chickens (Escorcia *et al.*, 2000; Callison *et al.*, 2001; Gelb *et al.*, 2001). They were shown by neutralisation tests to be different from Mass or Conn serotypes, but *in vivo* protection studies were not performed. Rather surprisingly, IBV variants do not appear to have been reported in Argentina until very

recently when Rimondi *et al.* (2009), using only molecular techniques, detected three unique genotype clusters (in addition to Mass and Conn); one of which was closely related to isolates from Brazil. Similar techniques had been used a few years earlier to identify for the first time a genetically unique IB variant in Colombia (Alvarado *et al.*, 2005).

Europe. Until the late 1970s, it was believed that only IBVs of the Mass serotype were important causes of disease in Europe. Then Dawson & Gough (1971) reported the detection of IBV variants in UK and workers at the Doorn Institute in the Netherlands isolated IBVs belonging to at least four different IBV serotypes associated with disease outbreaks in Mass-vaccinated commercial flocks (Davelaar *et al.*, 1984). These IBVs belonged to novel serotypes: D207 (also known as D274), D212 (better known as D1466), D3896 and D3128 and experimental data showed that existing IB vaccines protected poorly against them. Vaccines were developed using some of these variants, with several still in use today. The development in the UK of techniques which made use of tracheal organ cultures (TOCs), for both the isolation and serotyping of IBVs (Cook, 1984; Cook & Huggins, 1986) resulted in increased interest in the detection of new IB variants. Many new ones were isolated, not only in UK (Gough *et al.*, 1996) but also in other European countries including France (Picault *et al.*, 1986), Belgium (Meulemans *et al.*, 1987), Italy (Capua *et al.*, 1994; Zanella *et al.*, 2000; Zanella *et al.*, 2003), Poland (Minta *et al.*, 1998) and Spain (Dolz *et al.*, 2006, 2008).

Many, perhaps the majority, of these variants were detected for only a brief period and were probably of limited importance. However, occasionally a variant that caused major disease outbreaks did occur. One example of this is the IB variant B1648, mainly associated with renal problems in vaccinated flocks in Belgium and neighbouring countries for a relatively short period in the 1990s (Lambrechts *et al.*, 1993; Pensaert & Lambrechts, 1994). However, of major importance internationally was the variant called variously 4/91, 793B

and CR88 [named here 4/91] (Gough *et al.*, 1992; Parsons *et al.*, 1992; Le Gros, 1998) which emerged in the 1990s, associated with major welfare and economic problems in apparently well-vaccinated flocks. IB 4/91 quickly spread to many parts of the world and necessitated the development of live-attenuated IB vaccines to control it. This virus continues to be a major concern in poultry of all ages in many parts of the world, but interestingly has not been reported in the USA.

More recently, the use of RT-PCR technology has led to an increase in the detection of novel IB variants in Europe (Worthington *et al.*, 2008). However, it is important to remember that detecting an IB variant by molecular methods does not necessarily mean that the virus concerned is causing major disease problems. This can be illustrated by the IB variant Italian 02 (It-02). This virus appears to be easily detected by RT-PCR (Jones *et al.*, 2005; Dolz *et al.*, 2006) but is difficult to isolate and its association with widespread disease outbreaks remains to be established. On the other hand, the variant QX IBV is easy to detect by RT-PCR, readily isolated and is associated with major disease outbreaks in many areas. This virus, first reported associated with proventriculitis in China in the late 1990s (Yu *et al.*, 2001), spread throughout Russia (Bochkov *et al.*, 2006), and then appeared in much of Europe (Beato *et al.*, 2005; Landman *et al.*, 2005; Benyeda *et al.*, 2009). It is associated with nephritis and respiratory distress in broilers and with the so-called “false layer syndrome” and drops in egg production in breeders and layers. It is therefore causing major economic problems in IB-vaccinated flocks. Although no homologous vaccine was available until very recently, some success is reported in controlling its effects using currently available IB vaccines (De Wit *et al.*, 2006; Worthington & Jones, 2006; Terregino *et al.*, 2008; De Wit & Van de Sande, 2009; De Wit *et al.*, 2009).

Africa. IBV associated with swollen head syndrome and causing severe problems throughout southern Africa was isolated in the early 1980s (Morley & Thomson, 1984), confirmed as a variant and shown to be poorly protected against by Mass vaccines (Cook *et al.*, 1999). The only other known incidence of IBV variants in sub-Saharan Africa is the recent report by Ducatez *et al* (2009) who detected a novel IBV in Nigeria and Niger, that was antigenically and genetically distinct from other known IBVs. However, no association with disease was demonstrated and there is no information on the ability of currently available vaccines to protect against it.

IBV variants have been recognised in Egypt since the 1950s (Sheble *et al.*, 1986; Eid, 1998) with the isolation of a variant shown by neutralisation tests to be closely related to the Dutch variant D3128. Subsequently variants related to Mass, other European IBVs and one related to an Israeli variant have been identified by genome analysis in that country (Abdel-Moneim *et al.*, 2006). Bourogaa *et al.* (2009) used both molecular methods and cross neutralization tests to identify IBV isolates from Tunisia as variants which are closely related to ones found in Europe. In the early 1980s the unusual enterotropic variant, known as IB “G” was isolated in Morocco (El-Houadfi & Jones, 1985; El-Houadfi *et al.*, 1986). Interestingly, S1 sequence data have shown that IBV G and 4/91 are very closely related, possibly with a common origin (Jones *et al.*, 2004). The suggestion was made that parts of Africa, where variants have been little studied, might be a reservoir for such viruses, although the increasing number of variants being reported in other countries, such as Brazil and China, indicate that several such reservoirs might exist. More recently, El Bouqdaoui *et al.* (2005) used RT-PCR and RFLP techniques to study outbreaks of nephritis associated with IB in Morocco, where Mass and 4/91 vaccines have been used since 1960 and 2000 respectively. Three novel genotypes were identified, against which Mass vaccines provided poor protection.

Middle East. Variant IBVs have been recognised in Israel since at least the mid 1990s (Meir *et al.*, 1998; Callison *et al.*, 2001; Meir *et al.*, 2004) on the basis of both virus neutralisation tests and molecular techniques, and protection studies have shown that Mass vaccines provide inadequate protection against some of these novel variants. In Jordan, the use of RT-PCR enabled European IBV variants D274 and 4/91 to be detected (Roussan *et al.*, 2008), but since the primers were designed to detect only these specific variants, it is possible that others are present in that country. Similar methods have been used to identify 4/91 in Iran (Seyfi Abad Shapouri *et al.*, 2004).

India and Pakistan. Antibodies to several “American” (Muneer *et al.*, 1987a) and “European” (Ahmed *et al.*, 2007) IBV variants have been demonstrated in Pakistan, but virological studies have still to be performed. An IBV isolated in India in the early 2000s from cases of nephritis was reported to have a unique S1 sequence, indicating it to be different from other known IBVs (Bayry *et al.*, 2005).

Asia. In Asia, many studies have been performed in different countries in recent years. In Malaysia where IBV was first isolated in 1967, variants have been present since at least 1979 (reported by Lohr, 1988). More recently, molecular epidemiological studies of IBVs isolated in Malaysia and Singapore showed that, whilst some were of the Mass serotype (probably identical to the H120 vaccine), others were similar to IBVs reported from China and Taiwan (Yu *et al.*, 2001b). These studies led the authors to suggest that IBV variants have existed in Asia for some time. This finding was substantiated by Zulperi *et al.* (2009) who used sequence and phylogenetic analysis to study two variants isolated in Malaysia, 10 years apart. One was similar to several Chinese variants whilst the other was characterised as unique to Malaysia, but no protection studies were performed. IB has been a problem in Thailand since

the 1950s, despite the use of many different IB vaccines and a recent molecular study by Pohuang *et al.* (2009) has identified two groups of IBV variants in Thailand by phylogenetic analysis of the SI gene. Group I appeared to be unique to that country, whilst Group II showed a close relationship to Chinese IBVs, including variant A2 (see below).

IBV variants have been associated with disease outbreaks in Korea since at least the mid 1980s (Song *et al.*, 1998). Initially Mass vaccines were successful in controlling disease, but since 1990 IB outbreaks have been experienced in well vaccinated flocks with an increased incidence of renal problems. Song *et al.* (1998) classified 40 IBV isolates into Mass plus four local genotypes, one of which was not only the predominant type, but in pathogenicity studies, caused 50% mortality in SPF chicks inoculated at day-old. More recent studies (Lee *et al.*, 2004) have extended this work and reported further genetic diversity amongst Korean IBV variants isolated from diseased flocks; some of which are indigenous to Korea, whilst others share genetic relationships to IBV variants from other countries in the region (Lee *et al.*, 2008). It is suggested that Korean IBVs are continually evolving (Jang *et al.*, 2007).

Doi *et al.* (1982) studied the antigenic relationship of eight Japanese IBV isolates obtained between 1960 to 1974 and concluded that there were many distinct serotypes of IBV in Japan. Mase *et al.* (2004) carried out a detailed analysis of Japanese IBV variants by looking at the N-terminus of the S1 glycoprotein and identified three major genetic groups not found in other countries. One group, present in Japan since at least the 1960s, may be found only in Japan, whilst the other two, which were more recent, are related to Chinese and Taiwanese variants (see below). These groups were distinct from those found in Europe or the USA, although the 4/91 serotype has been isolated in Japan (Mase *et al.*, 2008; Shimazaki *et al.*, 2008). The origin of these variants is not clear, but Shieh *et al.* (2004), reported the close relationship of isolates from Taiwan and Japan to IBVs found in Australia

and the USA, and suggested that the Asian variants are recombinants; their S gene being derived from Australian variants and the N genes from US strains. More recently, analysis of the hypervariable region of the S1 subunit of Japanese field isolates has demonstrated the presence of a novel IBV variant (Mase *et al.*, 2010).

IBV variants have been recognised in Taiwan since at least the mid 1960s and two distinct lineages have been identified, as well as Mass and IBVs related to ones reported in neighbouring countries (Wang & Tsai, 1996; Wang & Huang, 2000; Huang *et al.*, 2004). However, recent data have suggested that the currently dominant IB variant in Taiwan may have arisen as a result of recombination in the 5' end of the N gene between a local and a foreign IBV (Kuo *et al.*, 2010). The failure of Mass vaccines to provide adequate protection has led to the development of vaccines from indigenous strains (Huang & Wang, 2006).

For many years little was known of the situation regarding IBV variants in China, but the fact that Mass type vaccines were used successfully suggests that variants may not have been a problem before the 1980s. However, by the mid 1990s this was clearly no longer the case and in the last decade many published reports have revealed the diversity of IBVs causing disease in that country. By means of *in vivo* studies and antigenic typing using mAbs and cross-neutralisation tests, Wu *et al.* (1998) identified highly pathogenic variant IBVs in China associated with both respiratory disease and nephritis and showed that H120 vaccine provided poor protection against challenge with these isolates. Possibly the most significant IBV variant worldwide to have emerged from China is the QX variant reported by Yu *et al.* (2001) in association with proventriculitis (see above). Another IBV genotype Q1 has also been associated with proventriculitis (Yu *et al.*, 2001a). However, it remains unclear whether the problems with proventriculitis that were associated with the detection of QX or Q1 strains were really caused by the IBV infection or by another factor because local replication of the IBV strain in the proventriculus was not shown. In a recent study of 26 IBVs isolated

between 1985 and 2008 from a variety of disease conditions in the Guangxi region of China, Wei *et al.* (2009) identified four clusters on the basis of RT-PCR analysis of the N gene. They were grouped into seven serotypes by neutralisation tests, but there was poor correlation between the results of genotyping and serotyping. In an analysis of the genome of 26 IBV variants isolated from kidneys, proventriculus and oviduct in different areas of China between 1995 and 2004, Liu *et al.* (2006) identified Mass type IBVs plus five genotypes apparently found only in China and co-circulating there. One of these (genotype A2) was subsequently shown to be the dominant indigenous type in China at that time (Liu *et al.*, 2009b), although more recently based on analysis of S1 gene sequences, Zou *et al.* (2010) have reported that a different genotype, designated LX4, is now dominant. Others showed close relationships with either Korean or Taiwanese IBV variants, and one was closely related to an Australian isolate (Liu *et al.*, 2006). Cuiping *et al.* (2007) identified the 4/91 variant in China, along with the Australian T strain, as well as one variant indigenous to China. Thus the diversity of IBV variants in China is now well established, some being restricted to and co-circulating in that country, whilst others show similarities with IBV variants identified in other countries in the region.

Russia. Molecular characterization using a part of the S gene of 91 IBV strains isolated between 1998 and 2002 from chickens in Russia showed the complexity of the IBV situation in that country (Bochkov *et al.*, 2006). The major group of isolates (38 viruses) was of the Mass genotype, circulating in Russia since the early 1970s; a second group of 22 strains were of the known European genotypes D274, 4/91, B1648, 624/I and It-02. Two further isolates from very distant geographic locations in Russia (the far east and European parts) clustered together with Chinese strains of the QX IBV genotype. The remaining 27 Russian isolates were divided in 11 novel genotypes.

Australia and New Zealand. In Australia, where IBV has always evolved independently from the rest of the world due to its geographical isolation (Ignjatovic *et al.*, 2006), many different IBV variants have been isolated and characterised since the early 1960s (Cumming, 1963) and *in vivo* protection studies performed with these variants (Klieve & Cumming, 1988). Using both mAbs directed against the major IBV proteins and sequencing studies, several distinct lineages have now been recognised (Ignjatovic *et al.*, 1997; Ignjatovic *et al.*, 2006), all different from those found elsewhere. Ignjatovic *et al.* (2002) in a study comparing the pathogenicity of 25 Australian IBV strains isolated between the 1960s and 1990s, found that of 12 IBVs isolated between 1961 and 1976, nine were nephropathogenic, whilst only three of 13 strains isolated between 1981 and 1994 were associated with nephritis. They suggested that this indicated a change in the prevalent IBV strains from highly nephropathogenic (1960s to 1970s) to respiratory (1980s to early 1990s). Their work also indicated the emergence in the late 1980s of respiratory strains with altered tissue tropism in Australia. It does not appear to be clear as yet whether or not similar changes in pathogenicity have occurred in other parts of the world.

Interestingly, in New Zealand, IB problems were uncommon before the 1970s, when IBV variants were first reported (Lohr, 1988). It was initially believed, on the basis of cross neutralisation tests, that IBV variants had evolved independently of those reported in Australia or the USA and at least four different variants were identified (Lohr, 1976, 1977). However, sequencing of the S1 gene has recently revealed genetic relationships between these early IBV isolates and ones made since 2000, and phylogenetic analysis has also showed that they are more closely related to Australian than to European or North American ones (McFarlane & Verma, 2008).

Spreading of the variants. The above account highlights the large number of IBV variants that exist worldwide; some being unique to a particular area, others having a more general distribution. The reason why some strains spread readily over major parts of the world, whereas others remain more local is unknown. Variants of major importance in major parts of the world, such as 4/91 or QX that have spread over Asia, Europe and Africa in a short period have not been reported in, for example, the USA or Australia. On the other hand, the major strain of the USA, the Arkansas strain has hardly been reported outside the USA. A pathogenic strain like D1466 that has been in some countries in Western Europe for three decades now has hardly been reported outside Western Europe, although it is very difficult to achieve a sufficient level of protection (Cook *et al.*, 1999) against this strain, which would make it easier for the strain to spread to other areas. It seems likely that geographical isolation and control measures employed in countries may play a part in preventing entry of IBV variants. The recent discoveries of IBV and IBV-like strains in other species of birds than the chicken (Cavanagh, 2005), such as geese, ducks, and pigeons might also play a role in the spread of IBV strains over the world. Specific IBV strains that were able to infect another bird species, especially if it is a migratory bird, would spread more easily over long distances than a strain that could not replicate in that bird species. An unknown IBV-like virus that might be common in a migratory bird could infect the poultry industry in different parts of the world, leaving us with the mystery of how this new virus spread so fast within the poultry industry. Although the role of the wild birds in the world of IBV is largely unknown and speculative, it is certainly an area that deserves more attention and research.

Current situation. As discussed, it is becoming clearer that many countries have to deal with many types of IBV. At the moment genotyping is by far the most used system and has largely

replaced serotyping and protectotyping. Does this create a problem? The preferred typing system depends on the goal (for example selection of vaccination programmes, or epidemiological studies), available techniques, experience, field situation and costs.

Classification systems are divided into two major groups: functional tests, that regard the biological function of a virus, and non-functional tests that examine the viral genome (De Wit, 2000). Typing by functional tests results in protectotypes and antigenic types (serotypes and epitope-types). Tests that look at the genome result in genotypes.

Protectotypes. With protectotyping, the complete immune response of a chicken against an IBV strain is measured. For the field, grouping of IBV strains into protectotypes is the most important system from a practical point of view, because it provides direct information about the efficacy of a vaccine. Strains that induce protection against each other in chickens belong to the same protectotype. However, protectotyping is laborious and expensive and requires both SPF chickens and high level facilities for performing vaccination-challenge studies.

Serotypes. Serotyping is based on the reaction between an IBV strain and chicken-induced IBV serotype-specific antibodies. Two strains (A and B) are considered to be of the same serotype when two-way heterologous neutralisation titres (antiserum A with virus B, and antiserum B with virus A) differ less than 20-fold from the homologous titres (antiserum A with virus A, antiserum B with virus B) in both directions (Hesslink, 1991). Serotyping becomes less practical as more IBV types are detected in a certain area since every serotype needs its own neutralization test, and for new strains that appear to be different, an antiserum has to be raised in SPF birds. As mentioned before, more and more countries have to deal with an increasing number of variants, which decreases the practicability of serotyping.

Genotypes. Grouping of strains based on genetic characterisation of (a part of) the genome results in genotypes. Methods include sequencing, detection of genotype-specific parts of the genome by RT-PCR, or determination of the position of enzyme cleavage sites (De Wit, 2000). Genomic information is objective and provides essential information for epidemiological studies (Figure 1). Most used for genotyping is the part of the genome that codes for the S1 subunit of the spike glycoprotein, which is the major inducer of protective immunity and carries most of the virus neutralizing epitopes, including serotype-specific epitopes, which are usually conformation-dependent (Mockett *et al.*, 1984; Cavanagh & Davis, 1986; Koch *et al.*, 1990; Cavanagh *et al.*, 1992).

Relationship between genotype, serotype and protectotype. A complicating factor with regard to genotyping of IBV is that a change of only a small percent of the amino acids in the S1 protein can result in a change of serotype (Cavanagh *et al.*, 1992) due to a change in a virus neutralizing epitopes, whereas other larger percentages of mutations at other parts of S1 might not result in a relevant change in antigenicity of the virus. On the other hand, IB viruses of different serotypes and genotypes not only have different epitopes, but also share common epitopes which are of importance in cross-immunity (Cavanagh *et al.*, 1992; Cavanagh *et al.*, 1997) and cell-mediated immune responses (Boots *et al.*, 1992; Ignjatovic & Galli, 1995). These features of the IB virus result in a disadvantage of genotyping for use in the field, as direct translation of information about usually only a part of the genome (generally a part of the S gene) of an IBV strain to biological function or antigenicity of the virus is not possible or is not without risk. Despite these limitations, there are reports that the S1 gene sequence comparison (part of 700 nucleotides (nt)) is a better predictor of challenge of immunity in chickens than serotyping by virus neutralization (Ladman *et al.*, 2006). Whether this is a general rule is unknown, as only a small number of strains and vaccines

have been compared, and also different parts of the S1 gene were used for the comparison of the homology. But in general, a lower homology in sequence of the S1 subunit of two strains (e.g. a vaccine and a field strain) means a greater chance that relevant mutations have occurred, which might result in a lower cross-protection. An analyses of several papers reporting the level of homology of the S1 gene or a part of it and level of cross-protection (Cavanagh *et al.*, 1997; Cook *et al.*, 2001; Meir *et al.*, 2004; Gelb *et al.*, 2005; Abdel-Moneim *et al.*, 2006; Ladman *et al.*, 2006; Liu *et al.*, 2009a) show that, in general, there is a higher chance of a good level of cross-protection between strains with a high level of homology than between strains with a low homology (Figure 2). However, these data also show that the relationship is not very strong. Some strains, which differ by only a few percent from other strains in the sequenced part of the genome showed a significant drop in cross-protection (Meir *et al.*, 2004; Abdel-Moneim *et al.*, 2006), whereas there was a high level of cross-protection against other strains with a much lower homology (Meir *et al.*, 2004). Figure 2 also shows the wide variation in level of cross-protection that is detected for strains with the same level of homology in comparison with the strains that are used as vaccine. Another limitation that needs to be considered is the different size and location of the S1 region of the S gene that different laboratories and research groups use for their analyses of the level of homology (De Wit *et al.*, 2010a). Several laboratories sequence a part of S1 that includes the highly variable regions (HVR) 1 and 2; others do not. Analysis of different parts of S1 can result in different levels of homology leading to different conclusions regarding the relationship between virus strains. An example is shown in Table 1 and Figures 3, 4, and 5. For this analysis the sequences of four IBV strains of different genotypes and serotypes, as published by Adznar *et al.* (1997), were used. The homology of the four strains (7/91 (genotype 4/91), H120 (Mass serotype), D274 and D1466) was compared for nucleotides 1 to 1600 of the S1 region of the S gene. The extent of nucleotide identity between the strains was

expressed on nucleotide level as follows: score 100% when all four strains had the same nucleobase (in the copy DNA); score 75% when three out of four strains had the same nucleobase; score 50% when two strains had the same nucleobase and the two other strains had another nucleobase in common; score 25% when two strains had the same nucleobase and the two other strains each had another different nucleobase; score 0% when all strains had a different nucleobase at that position of the genome. In this comparison, a missing nucleotide (by deletion) was considered to be a different nucleobase. Table 1 shows the overview of the comparison. When windows of 10 nt were used for the comparison of the four strains, the average homology of the four strains could have been 42.5% but also 100% based on the location of the 10 nt that have been used for the comparison. When windows of 100 nt were used for the comparison of the four strains, the average homology of the four strains could have been 68.0% but also 86.8% based on the location of the 100 nt that were used for the comparison. When windows of 400 nt were used for the comparison, the average homology of the four strains varied from 73.2% to 81.6% based of the location of the 400 nt that were used for the comparison.

When the IBV strain tested is the result of a recombination event between different IB genotypes, examination of different parts of S1 can mean the detection of a different genotype (Wang *et al.*, 1993; Jia *et al.*, 1995; Dolz *et al.*, 2008). The detected homology with other strains is then very dependent on the part of the genome that is being used for the comparisons of the homology. Using larger parts of S1 for the comparison of strains results in a decreased risk of finding high levels of homology between strains where in fact this is not really the case. In the field situation there is another limitation on the predictive value of a vaccine against the field challenge strain based on the homology in the S1 gene. In many countries, IBV vaccines are used that are a combination of two strains, or the vaccination programme contains two vaccinations with two (or more) different IBV strains. The level of

cross-protection then depends on the efficacy of the combination of strains, which makes it impossible to determine a level of homology between the field strain and the “vaccine strain”.

This leads to the conclusion that genotyping is an excellent tool for epidemiological studies (Figure 1), and is a convenient, practical tool for typing that can be used best as a means of screening to select potentially important strains. In situations where there is suspicion in the field that the genotype of recent isolates does not provide accurate information about the true antigenic nature of these IBV isolates, then conventional testing (serotyping) and especially *in vivo* protection studies are required.

Control measures. IBV is ubiquitous in most parts of the world where poultry are reared and is able to spread very rapidly in non-protected birds (De Wit *et al.*, 1998). It is shed via both the respiratory tract and the faeces and can persist in the birds, intestinal tract and faeces for several weeks or months. Although strict biosecurity and working with a one-age system are essential control measures, normally vaccination is an essential tool to increase the resistance of the chickens against challenge with IBV strains (Cook, 2008).

For vaccination of chickens against IBV, both live attenuated and inactivated (usually oil-adjuvanted) vaccines are used. Live vaccines are especially used in young birds to achieve early protection against challenge and also for priming of future layers and breeders that will be boosted with the inactivated vaccines. In areas with an increased level of field challenge, live attenuated vaccines are also used periodically during the laying period with the intention of keeping the local protection of the respiratory tract at a high level.

Most used in the world are vaccines of the Mass serotype. In several parts of the world, Mass vaccines are the only vaccines allowed, but elsewhere vaccines of one or more other serotypes are permitted. Vaccines of a certain serotype or genotype are normally able to

protect the well vaccinated chicken against a homologous challenge. Often there is a partial protection against strains of other protectotypes, serotypes or genotypes that can vary from high to low (Figure 2 and reviewed for the H-strain by Bijlenga *et al.* (2004). The magnitude and duration of the response to vaccination is dependent on many factors, including age of the chick, levels of maternal immunity, immunogenicity of the vaccine, method of vaccine application, virulence of the field strain challenge, interval between vaccination and challenge and immunocompetency of the host. Chickens vaccinated under optimal conditions may have immunity lasting many months and for broilers, this may be life-long (Bijlenga *et al.*, 2004).

It has been shown that vaccination with two antigenically distinct live-attenuated vaccines such as Mass and 4/91, can result in a broad cross-protection against many different IBV types (Cook *et al.*, 1999; Terregino *et al.*, 2008). The cross-protection was broader when these vaccines were applied with a 2 week interval than when the vaccines were combined on the same day.

Results of challenge studies and field work have shown that vaccination with a bivalent vaccine containing the Mass and Ark strains (Gelb *et al.*, 1989; Gelb *et al.*, 1991; Gelb *et al.*, 2005) provided, on average, a higher level of cross-protection against certain heterologous field strains than other combinations of vaccines such as Mass together with Conn or with JMK. However, the Mass and Ark vaccine did not provide significant protection against challenge with another strain (Ladman *et al.*, 2002). It is unknown whether the separate application of these two IBV strains would have resulted in a higher (or lower) level of cross-protection against the same heterologous challenges.

A well vaccinated chicken is protected against challenge with a virulent homologous IBV strain. This means that this well vaccinated chicken is also protected against an early revaccination with a homologous vaccine (Davelaar & Kouwenhoven, 1980). Despite this,

revaccination of young birds, especially broilers, using a vaccine of the same serotype as the first vaccine, has proven to be beneficial under field conditions. This is an indication that the quality of the first vaccination might need careful attention. Whatever live vaccine is used, the application is a very critical step. IBV virus is a sensitive virus that can be inactivated easily (Cavanagh & Gelb, 2008) which may result in inadequate efficacy of the vaccination under field conditions (Jackwood *et al.*, 2009; De Wit *et al.*, 2010b). The vaccine may be applied by eye- or nasal drop, spray or drinking water routes and it is essential that a high percentage of the birds receive a required dose of the vaccine in the right tissue. Inadequate 'take' of the vaccine may result in no or a decreased level of protection, delayed protection, prolonged presence/circulation of the vaccine virus in the flock resulting in an increased risk of infections with *E.coli* or other bacteria (Goren, 1978; Hopkins & Yoder, 1984; Smith *et al.*, 1985; Cook *et al.*, 1986; Matthijs *et al.*, 2003) and even increase of virulence of the virus (Hopkins & Yoder, 1986).

In order to achieve an increased level of protection during the laying period of commercial layers and parent stock, the use of inactivated IBV vaccines after a priming with live attenuated IBV vaccines has been shown to be effective against homologous Mass challenges (Gough *et al.*, 1977; Box *et al.*, 1980; Timms & Bracewell, 1983; Box & Ellis, 1985; Box *et al.*, 1988). The efficacy of increasing the level of protection against heterologous challenges in the laying period has rarely been reported (De Wit *et al.*, 2009), although birds that had been vaccinated twice with a live Mass type vaccine and boosted with a killed oil-emulsion vaccine containing a Mass strain showed no protection against challenge with a strain of the Ark type (Muneer *et al.*, 1987b).

Finally, it must be emphasised that whilst high quality live-attenuated and inactivated IB vaccines are available for use worldwide, two points require careful consideration by those responsible for the welfare of commercial chicken flocks. Firstly, the

importance of careful application of the vaccines cannot be overstated. Secondly, there may be a need to amend vaccine programmes, using vaccines licensed for use in a particular area, in order to optimise protection against the IB variants currently circulating in that area.

Conclusion

The problems of how to type the large number of IBV variants that are now found worldwide, and how to relate the findings to the best vaccination strategy for protection, are clearly complex. Whilst genotyping has advantages of ease and speed, the different sizes and locations of the part of the S1 region of the S gene that are used by different groups for the comparison of strains makes interpretation of results very difficult and this review emphasises the need for a standardised method of performing genotyping. Another major point that needs to be remembered is that only the vaccinated chicken decides whether genetic or antigenic differences as shown by genotyping or serotyping are relevant for the level of cross-protection in that bird.

References

- Abdel-Moneim, A.S., El-Kady, M.F., Ladman, B.S. & Gelb, J. (2006). S1 gene sequence analysis of a nephropathogenic strain of avian infectious bronchitis virus in Egypt. *Virology Journal*, 3, 78.
- Abreu, J.T., Resende, J.S., Flatschart, R.B., Folgueras-Flatschart, A.V., Mendes, A.C., Martins, N.R., Silva, C.B., Ferreira, B.M. & Resende, M. (2006). Molecular analysis of Brazilian infectious bronchitis field isolates by reverse transcription-polymerase

chain reaction, restriction fragment length polymorphism, and partial sequencing of the N gene. *Avian Diseases*, 50, 494-501.

Adzhar, A., Gough, R.E., Haydon, D., Shaw, K., Britton, P. & Cavanagh, D. (1997).

Molecular analysis of the 793/B serotype of infectious bronchitis virus in Great Britain. *Avian Pathology*, 26, 625-640.

Ahmed, Z., Naeem, K. & Hameed, A. (2007). Detection and seroprevalence of infectious bronchitis virus strains in commercial poultry in Pakistan. *Poultry Science*, 86, 1329-1335.

Alvarado, I.R., Villegas, P., El-Attrache, J. & Brown, T.P. (2003). Evaluation of the protection conferred by commercial vaccines against the California 99 isolate of infectious bronchitis virus. *Avian Diseases*, 47, 1298-1304.

Alvarado, I.R., Villegas, P., Mossos, N. & Jackwood, M.W. (2005). Molecular characterization of avian infectious bronchitis virus strains isolated in Colombia during 2003. *Avian Diseases*, 49, 494-499.

Ammayappan, A., Upadhyay, C., Gelb, J. & Vakharia, V.N. (2008). Complete genomic sequence analysis of infectious bronchitis virus Ark DPI strain and its evolution by recombination. *Virology Journal*, 5, 157.

Bayry, J., Goudar, M.S., Nighot, P.K., Kshirsagar, S.G., Ladman, B.S., Gelb, J., Ghalsasi, G.R. & Kolte, G.N. (2005). Emergence of a nephropathogenic avian infectious bronchitis virus with a novel genotype in India. *Journal of Clinical Microbiology*, 43, 916-918.

Beato, M.S., De Battisti, C., Terregino, C., Drago, A., Capua, I. & Ortali, G. (2005).

Evidence of circulation of a Chinese strain of infectious bronchitis virus (QXIBV) in Italy. *Veterinary Record*, 156, 720.

- Benyeda, Z., Mato, T., Suveges, T., Szabo, E., Kardi, V., Abonyi-Toth, Z., Rusvai, M. & Palya, V. (2009). Comparison of the pathogenicity of QX-like, M41 and 793/B infectious bronchitis strains from different pathological conditions. *Avian Pathology*, 38, 449-456.
- Bijlenga, G., Cook, J.K.A., Gelb, J. & de Wit, J.J. (2004). Development and use of the H strain of avian infectious bronchitis virus from the Netherlands as a vaccine: a review. *Avian Pathology*, 33, 550-557.
- Bochkov, Y.A., Batchenko, G.V., Shcherbakova, L.O., Borisov, A.V. & Drygin, V.V. (2006). Molecular epizootiology of avian infectious bronchitis in Russia. *Avian Pathology*, 35, 379-393.
- Boots, A.M., Benaissa-Trouw, B.J., Hesselink, W., Rijke, E., Schrier, C. & Hensen, E.J. (1992). Induction of anti-viral immune responses by immunization with recombinant-DNA encoded avian coronavirus nucleocapsid protein. *Vaccine*, 10, 119-124.
- Bourogaa, H., Miled, K., Gribaa, L., El Behi, I. & Ghram, A. (2009). Characterization of new variants of avian infectious bronchitis virus in Tunisia. *Avian Diseases*, 53, 426-433.
- Box, P.G., Beresford, A.V. & Roberts, B. (1980). Protection of laying hens against infectious bronchitis with inactivated emulsion vaccines. *Veterinary Record*, 106, 264-268.
- Box, P.G. & Ellis, K.R. (1985). Infectious bronchitis in laying hens: interference with response to emulsion vaccine by attenuated live vaccine. *Avian Pathology*, 14, 9-22.
- Box, P.G., Holmes, H.C., Finney, P.M. & Froymann, R. (1988). Infectious bronchitis in laying hens: the relationship between haemagglutination inhibition antibody levels and resistance to experimental challenge. *Avian Pathology*, 17, 349-361.
- Branden, R.C. & Da Silva, E.N. (1986). Ocurrência de “nuevos” serotipos de bronquitis infecciosa en Brasil. In: P. Villegas (Ed.). Proceedings of VI Seminario Internacional de Patologia aviar Georgia USA.

- Callison, S.A., Jackwood, M.W. & Hilt, D.A. (2001). Molecular characterization of infectious bronchitis virus isolates foreign to the United States and comparison with United States isolates. *Avian Diseases*, 45, 492-499.
- Capua, I., Gough, R.E., Mancini, M., Casaccia, C. & Weiss, C. (1994). A 'novel' infectious bronchitis strain infecting broiler chickens in Italy. *Zentralblatt für Veterinärmedizin. Reihe .*, 41, 83-89.
- Cavanagh, D. (2005). Coronaviruses in poultry and other birds. *Avian Pathology*, 34, 439-448.
- Cavanagh, D. & Davis, P.J. (1986). Coronavirus IBV: removal of spike glycopolyptide S1 by urea abolishes infectivity and haemagglutination but not attachment to cells. *Journal of General Virology*, 67, 1443-1448.
- Cavanagh, D., Davis, P.J., Cook, J.K.A., Li, D., Kant, A. & Koch, G. (1992). Location of the amino acid differences in the S1 spike glycoprotein subunit of closely related serotypes of infectious bronchitis virus. *Avian Pathology*, 21, 33-43.
- Cavanagh, D., Ellis, M.M. & Cook, J.K.A. (1997). Relationship between sequence variation in the S1 spike protein of infectious bronchitis virus and the extent of cross-protection in vivo. *Avian Pathology*, 26, 63-74.
- Cavanagh, D. & Gelb, J. (2008). Infectious Bronchitis. In Y. M. Saif, Fadly, A.M., Glisson, J.R., McDougald, L.R., Nolan, L.K. and Swayne, D.E. (Eds.), *Diseases of Poultry* (12th edn, pp. 117-135).
- Cook, J.K.A. (1984). The classification of new serotypes of infectious bronchitis virus isolated from poultry flocks in Britain between 1981 and 1983. *Avian Pathology*, 13, 733-741.
- Cook, J.K.A. (2008). Coronaviridae. In M. Pattison, P.F. McMullin, J.M. Bradbury and D.J. Alexander (Eds), *Poultry Diseases* (6th edn, pp. 340-349): Saunders Elsevier.

- Cook, J.K.A., Chesher, J., Baxendale, W., Greenwood, N., Huggins, M.B. & Orbell, S.J. (2001). Protection of chickens against renal damage caused by a nephropathogenic infectious bronchitis virus. *Avian Pathology*, 30, 423-426.
- Cook, J.K.A. & Huggins, M.B. (1986). Newly isolated serotypes of infectious bronchitis virus: their role in disease. *Avian Pathology*, 15, 129-138.
- Cook, J.K.A., Orbell, S.J., Woods, M.A. & Huggins, M.B. (1999). Breadth of protection of the respiratory tract provided by different live-attenuated infectious bronchitis vaccines against challenge with infectious bronchitis viruses of heterologous serotypes. *Avian Pathology*, 28, 477-485.
- Cook, J.K.A., Smith, H.W. & Huggins, M.B. (1986). Infectious bronchitis immunity: its study in chickens experimentally infected with mixtures of infectious bronchitis virus and *Escherichia coli*. *Journal of General Virology*, 67, 1427-1434.
- Cowen, B.S. & Hitchner, S.B. (1975). Serotyping of avian infectious bronchitis viruses by the virus-neutralization test. *Avian Diseases*, 19, 583-595.
- Cubillos, A., Ulloa, J., Cubillos, V. & Cook, J.K.A. (1991). Characterisation of strains of infectious bronchitis virus isolated in Chile. *Avian Pathology*, 20, 85-99.
- Cuiping, X., Jixun, Z. & Xudong, Z. (2007). Isolation and identification of four isolates of infectious bronchitis strains in China and analysis of their SI protein gene. *Veterinary Microbiology*, 122, 61-71.
- Cumming, R.B. (1963). Infectious avian nephrosis (uraemia) in Australia. *Australian Veterinary Journal*, 39, 145-147.
- Davelaar, F.G. & Kouwenhoven, B. (1980). Vaccination of 1-day-old broilers against infectious bronchitis by eye drop application or coarse droplet spray and the effect of revaccination by spray. *Avian Pathology*, 9, 499-510.

- Davelaar, F.G., Kouwenhoven, B. & Burger, A.G. (1984). Occurrence and significance of infectious bronchitis virus variant strains in egg and broiler production in the Netherlands. *Veterinary Quarterly*, 6, 114-120.
- Dawson, P.S. & Gough, R.E. (1971). Antigenic variation in strains of avian infectious bronchitis virus. *Archiv für die gesamte Virusforschung*, 34, 32-39.
- De Wit, J.J. (2000). Detection of infectious bronchitis virus. *Avian Pathology*, 29, 71-93.
- De Wit, J.J., Cook, J.K.A. & van der Heijden, H.M.J.F. (2010a). Infectious bronchitis virus in Asia, Africa, Australia and Latin America - history, current situation and control measures. *Brazilian Journal of Poultry Science*, 12, 97-106.
- De Wit, J.J., de Jong, M.C., Pijpers, A. & Verheijden, J.H. (1998). Transmission of infectious bronchitis virus within vaccinated and unvaccinated groups of chickens. *Avian Pathology*, 27, 464-471.
- De Wit, J.J., Nieuwenhuizen, J. & Fabri, T.H.F. (2006). Protection by maternally derived antibodies and vaccination at day of hatch against early challenge with IBV serotype D388. In: U. Heffels-Redmann & E.F. Kaleta (Eds.), *Proceedings of the Vth International Symposium on Corona- and Pneumovirus Infections* (pp. 314-318). Rauschholzhausen, Germany.
- De Wit, J.J., Swart, W.A.J.M. & Fabri, T.H.F. (2010b). The efficacy of infectious bronchitis virus vaccinations in the field: association between the α -IBV IgM response, protection and vaccine application parameters. *Avian Pathology*, 39, 123-132.
- De Wit, J.J. & Van de Sande, H. (2009). Efficacy of combined vaccines at day of hatch against a D388 challenge in SPF and commercial chickens. In: U. Heffels-Redmann & E.F. Kaleta (Eds.) *Proceedings of the VIth International Symposium on Corona- and Pneumoviruses and Complicating Pathogens* (pp. 177-182). Rauschholzhausen, Germany.

- De Wit, J.J., Van de Sande, H. & Prandini, F. (2009). Enhanced efficacy of the use of a monovalent infectious bronchitis virus inactivated vaccine in layers primed with H120 and 793B live IBV vaccines to increase the protection against challenge with 3 European serotypes of IBV. In: U. Heffels-Redmann & E.F. Kaleta (.), *Proceedings of the Vth. International Symposium on Corona- and Pneumoviruses and Complicating Pathogens* (pp. 159-167). Rauschholzhausen, Germany.
- Di Fabio, J., Rossini, L.I., Orbell, S.J., Paul, G., Huggins, M.B., Malo, A., Silva, B.G. & Cook, J.K.A. (2000). Characterization of infectious bronchitis viruses isolated from outbreaks of disease in commercial flocks in Brazil. *Avian Diseases*, *44*, 582-589.
- Doi, M., Yamakami, T., Koimaru, H., Yoshimura, M., Masu, S., Shirai, J. & Kawamura, H. (1982). Serotypes of avian infectious bronchitis virus isolates from field cases in Japan. *Avian Diseases*, *26*, 946-956.
- Dolz, R., Pujols, J., Ordonez, G., Porta, R. & Majo, N. (2006). Antigenic and molecular characterization of isolates of the Italy 02 infectious bronchitis virus genotype. *Avian Pathology*, *35*, 77-85.
- Dolz, R., Pujols, J., Ordonez, G., Porta, R. & Majo, N. (2008). Molecular epidemiology and evolution of avian infectious bronchitis virus in Spain over a fourteen-year period. *Virology*, *374*, 50-59.
- Ducatez, M.F., Martin, A.M., Owoade, A.A., Olatoye, I.O., Alkali, B.R., Maikano, I., Snoeck, C.J., Sausy, A., Cordioli, P. & Muller, C.P. (2009). Characterization of a new genotype and serotype of infectious bronchitis virus in Western Africa. *Journal of General Virology*, *90*, 2679-2685.
- Eid, A.A.M. (1998). Infectious bronchitis virus infection in Egypt. In: E.F. Kaleta & U. Heffels-Redmann (Eds.), *Proceedings of the International Symposium on Infectious*

- Bronchitis and Pneumovirus Infections in Poultry* (pp. 145-156). Rauschholzhausen, Germany.
- El-Bouqdaoui, M., Mhand, R. A., Bouayoune, H. & Ennaji, M.M. (2005). Genetic grouping of nephropathogenic avian infectious bronchitis virus isolated in Morocco. *International Journal of Poultry Science*, 4, 721-727.
- El-Houadfi, M. & Jones, R.C. (1985). Isolation of avian infectious bronchitis viruses in Morocco including an enterotropic variant. *Veterinary Record*, 116, 445.
- El-Houadfi, M., Jones, R.C., Cook, J.K.A. & Ambali, A.G. (1986). The isolation and characterisation of six avian infectious bronchitis viruses isolated in Morocco. *Avian Pathology*, 15, 93-105.
- Escorcía, M., Jackwood, M.W., Lucio, B., Petrone, V.M., Lopez, C., Fehervari, T. & Tellez, G. (2000). Characterization of Mexican strains of avian infectious bronchitis isolated during 1997. *Avian Diseases*, 44, 944-947.
- Fields, D.B. (1973). Arkansas 99, a new infectious bronchitis serotype. *Avian Diseases*, 17, 659-661.
- Gelb, J. & Cloud, S.S. (1983). Effect of serial embryo passage of an Arkansas-type avian infectious bronchitis virus isolate on clinical response, virus recovery, and immunity. *Avian Diseases*, 27, 679-687.
- Gelb, J., Keeler, C.L., Jr., Nix, W.A., Rosenberger, J.K. & Cloud, S.S. (1997). Antigenic and S-1 genomic characterization of the Delaware variant serotype of infectious bronchitis virus. *Avian Diseases*, 41, 661-669.
- Gelb, J., Ladman, B.S., Tamayo, M., Gonzalez, M. & Sivanandan, V. (2001). Novel infectious bronchitis virus S1 genotypes in Mexico 1998-1999. *Avian Diseases*, 45, 1060-1063.

- Gelb, J., Leary, J.H. & Rosenberger, J.K. (1983). Prevalence of Arkansas-type infectious bronchitis virus in Delmarva peninsula chickens. *Avian Diseases*, 27, 667-678.
- Gelb, J., Perkins, B.E., Rosenberger, J.K. & Allen, P.H. (1981). Serologic and cross-protection studies with several infectious bronchitis virus isolates from Delmarva-reared broiler chickens. *Avian Diseases*, 25, 655-666.
- Gelb, J., Rosenberger, J.K., Fries, P.A., Cloud, S.S., Odor, E.M., Dohms, J.E. & Jaeger, J.S. (1989). Protection afforded infectious bronchitis virus-vaccinated sentinel chickens raised in a commercial environment. *Avian Diseases*, 33, 764-769.
- Gelb, J., Weisman, Y., Ladman, B.S. & Meir, R. (2005). S1 gene characteristics and efficacy of vaccination against infectious bronchitis virus field isolates from the United States and Israel (1996 to 2000). *Avian Pathology*, 34, 194-203.
- Gelb, J., Wolff, J.B. & Moran, C.A. (1991). Variant serotypes of infectious bronchitis virus isolated from commercial layer and broiler chickens. *Avian Diseases*, 35, 82-87.
- Goren, E. (1978). Observations on experimental infection of chicks with *Escherichia coli*. *Avian Pathology*, 7, 213-224.
- Gough, R.E., Allan, W.H. & Nedelciu, D. (1977). Immune response to monovalent and bivalent Newcastle disease and infectious bronchitis inactivated vaccines. *Avian Pathology*, 6, 131-142.
- Gough, R.E., Cox, W.J., Gutierrez, E., MacKenzie, G., Wood, A.M. & Dagless, M.D. (1996). Isolation of 'variant' strains of infectious bronchitis virus from vaccinated chickens in Great Britain. *Veterinary Record*, 139, 552.
- Gough, R.E., Randall, C.J., Dagless, M., Alexander, D.J., Cox, W.J. & Pearson, D. (1992). A 'new' strain of infectious bronchitis virus infecting domestic fowl in Great Britain. *Veterinary Record*, 130, 493-494.

- Hesselink, W.G. (1991). Serotyping avian infectious bronchitis virus: selection of a unified method. In: E.F. Kaleta & U. Heffels-Redmann (Eds.), *Proceedings of the 2nd International Symposium on Infectious Bronchitis* (pp. 87-97). Rauischholzhausen, Germany.
- Hidalgo, H., Gallardo, R. & Rosende, S. (1976). Isolation of infectious bronchitis virus from broiler chickens in Chile. *Avian Diseases*, 20, 601-603.
- Hidalgo, H., Gallardo, R. & Toro, H. (1986). Antigenic and pathogenic properties of three isolates of infectious bronchitis virus obtained from vaccinated chickens. *Journal of Veterinary Medicine*, B, 33, 26-35.
- Hipólito, O. (1957). Isolamento e identificação do vírus da bronquite infecciosa das galinhas no Brasil. *Arquivo Escola Veterinária Universidade de Minas Gerais*, 10, 131-151.
- Hitchner, S.B., Winterfield, R.W. & Appleton, G.S. (1966). Infectious bronchitis virus types - incidence in the United States. *Avian Diseases*, 10, 98-102.
- Hofstad, M.S. (1961). Antigenic and immunological studies on several isolates of avian infectious bronchitis virus. *Avian Diseases*, 5, 102-107.
- Hofstad, M.S. (1981). Cross-immunity in chickens using seven isolates of avian infectious bronchitis virus. *Avian Diseases*, 25, 650-654.
- Hopkins, S.R. (1974). Serological comparisons of strains of infectious bronchitis virus using plaque-purified isolants. *Avian Diseases*, 18, 231-239.
- Hopkins, S.R. & Yoder, H.W., Jr. (1984). Increased incidence of airsacculitis in broilers infected with mycoplasma synoviae and chicken-passaged infectious bronchitis vaccine virus. *Avian Diseases*, 28, 386-396.
- Hopkins, S.R. & Yoder, H.W., Jr. (1986). Reversion to virulence of chicken-passaged infectious bronchitis vaccine virus. *Avian Diseases*, 30, 221-223.

- Huang, Y.P., Lee, H.C., Cheng, M.C. & Wang, C.H. (2004). S1 and N gene analysis of avian infectious bronchitis viruses in Taiwan. *Avian Diseases*, 48, 581-589.
- Huang, Y.P. & Wang, C.H. (2006). Development of attenuated vaccines from Taiwanese infectious bronchitis virus strains. *Vaccine*, 24, 785-791.
- Ignjatovic, J., Ashton, D.F., Reece, R., Scott, P. & Hooper, P. (2002). Pathogenicity of Australian strains of avian infectious bronchitis virus. *Journal of Comparative Pathology*, 126, 115-123.
- Ignjatovic, J. & Galli, U. (1995). Immune responses to structural proteins of avian infectious bronchitis virus. *Avian Pathology*, 24, 313-332.
- Ignjatovic, J., Gould, G. & Sapats, S. (2006). Isolation of a variant infectious bronchitis virus in Australia that further illustrates diversity among emerging strains. *Archives of Virology*, 151, 1567-1585.
- Ignjatovic, J., Sapats, S.I. & Ashton, F. (1997). A long-term study of Australian infectious bronchitis viruses indicates a major antigenic change in recently isolated strains. *Avian Pathology*, 26, 535-552.
- Jackwood, M.W., Hilt, D.A. & Brown, T.P. (2003). Attenuation, safety, and efficacy of an infectious bronchitis virus GA98 serotype vaccine. *Avian Diseases*, 47, 627-632.
- Jackwood, M.W., Hilt, D.A., Lee, C.W., Kwon, H.M., Callison, S.A., Moore, K.M., Moscoso, H., Sellers, H. & Thayer, S. (2005). Data from 11 years of molecular typing infectious bronchitis virus field isolates. *Avian Diseases*, 49, 614-618.
- Jackwood, M.W., Hilt, D.A., McCall, A.W., Polizzi, C.N., McKinley, E.T. & Williams, S.M. (2009). Infectious bronchitis virus field vaccination coverage and persistence of Arkansas-type viruses in commercial broilers. *Avian Diseases*, 53, 175-183.

- Jackwood, M.W., Hilt, D.A., Sellers, H.S., Williams, S.M. & Lasher, H.N. (2010). Rapid heat-treatment attenuation of infectious bronchitis virus. *Avian Pathology*, 39, 227-233.
- Jackwood, M.W., Hilt, D.A., Williams, S.M., Woolcock, P., Cardona, C. & O'Connor, R. (2007). Molecular and serologic characterization, pathogenicity, and protection studies with infectious bronchitis virus field isolates from California. *Avian Diseases*, 51, 527-533.
- Jang, J.H., Sung, H.W., Song, C.S. & Kwon, H.M. (2007). Sequence analysis of the S1 glycoprotein gene of infectious bronchitis viruses: identification of a novel phylogenetic group in Korea. *Journal of Veterinary Science*, 8, 401-407.
- Jia, W., Karaca, K., Parrish, C.R. & Naqi, S.A. (1995). A novel variant of avian infectious bronchitis virus resulting from recombination among three different strains. *Archives of Virology*, 140, 259-271.
- Jia, W., Mondal, S.P. & Naqi, S.A. (2002). Genetic and antigenic diversity in avian infectious bronchitis virus isolates of the 1940s. *Avian Diseases*, 46, 437-441.
- Johnson, R.B. & Marquardt, W.W. (1976). Strains of infectious bronchitis virus on the Delmarva peninsula and in Arkansas. *Avian Diseases*, 20, 382-386.
- Jones, R.C., Savage, C.E., Naylor, C.J., Cook, J.K.A. & El-Houadfi, M. (2004). A possible North African progenitor of the major European infectious bronchitis variant (793B, 4/91, CR88). In: E.F. Kaleta & U. Heffels-Redmann (Eds.), *Proceedings of the IV International Symposium on Avian Corona- and Pneumovirus Infections* (pp. 105-111). Rauschholzhausen, Germany.
- Jones, R.C., Worthington, K.J. & Gough, R.E. (2005). Detection of the Italy O2 strain of infectious bronchitis virus in the UK. *Veterinary Record*, 156, 260.

- Jungherr, E.L., Chomiak, T.W. & Luginbuhl, R.E. (1956). Immunologic differences in strains of infectious bronchitis virus. *Proceedings of 60th Annual Meeting of the United States Livestock Sanitary Association* (pp. 203-209). Chicago, USA.
- Klieve, A.V. & Cumming, R.B. (1988). Immunity and cross-protection to nephritis produced by Australian infectious bronchitis viruses used as vaccines. *Avian Pathology*, 17, 829-839.
- Koch, G., Hartog, L., Kant, A. & van Roozelaar, D.J. (1990). Antigenic domains on the peplomer protein of avian infectious bronchitis virus: correlation with biological functions. *Journal of General Virology*, 71, 1929-1935.
- Kuo, S.M., Wang, C.H., Hou, M.H., Huang, Y.P., Kao, H.W. & Su, H.L. (2010). Evolution of infectious bronchitis virus in Taiwan: characterisation of RNA recombination in the nucleocapsid gene. *Veterinary Microbiology*, 144, 293-302.
- Ladman, B.S., Loupos, A.B. & Gelb, J. (2006). Infectious bronchitis virus S1 gene sequence comparison is a better predictor of challenge of immunity in chickens than serotyping by virus neutralization. *Avian Pathology*, 35, 127-133.
- Ladman, B.S., Pope, C.R., Ziegler, A.F., Swieczkowski, T., Callahan, C.J., Davison, S. & Gelb, J. (2002). Protection of chickens after live and inactivated virus vaccination against challenge with nephropathogenic infectious bronchitis virus PA/Wolgemuth/98. *Avian Diseases*, 46, 938-944.
- Lambrechts, C., Pensaert, M. & Ducatelle, R. (1993). Challenge experiments to evaluate cross-protection induced at the trachea and kidney level by vaccine strains and Belgian nephropathogenic isolates of avian infectious bronchitis virus. *Avian Pathology*, 22, 577-590.
- Landman, W.J.M., Dwars, R.M. & De Wit, J.J. (2005). High incidence of false layers in (re)production hens supposedly attributed to a juvenile infectious bronchitis infection.

XIVth International Congress of the World Veterinary Poultry Association (p. 369).
Turkey, Istanbul.

- Le Gros, F.X. (1998). Serotyping studies on recent IBV isolates from France and various regions of the world. In: E.F. Kaleta & U. Heffels-Redmann (Eds), *Proceedings of the International Symposium on Infectious Bronchitis and Pneumovirus Infections in Poultry* (pp. 205-209). Rauschholzhausen, Germany.
- Lee, C.W., Hilt, D.A. & Jackwood, M.W. (2001). Identification and analysis of the Georgia 98 serotype, a new serotype of infectious bronchitis virus. *Avian Diseases*, 45, 164-172.
- Lee, C.W. & Jackwood, M.W. (2000). Evidence of genetic diversity generated by recombination among avian coronavirus IBV. *Archives of Virology*, 145, 2135-2148.
- Lee, C.W. & Jackwood, M.W. (2001a). Origin and evolution of Georgia 98 (GA98), a new serotype of avian infectious bronchitis virus. *Virus Research*, 80, 33-39.
- Lee, C.W. & Jackwood, M.W. (2001b). Spike gene analysis of the DE072 strain of infectious bronchitis virus: origin and evolution. *Virus Genes*, 22, 85-91.
- Lee, E.K., Jeon, W.J., Lee, Y.J., Jeong, O.M., Choi, J.G., Kwon, J.H. & Choi, K.S. (2008). Genetic diversity of avian infectious bronchitis virus isolates in Korea between 2003 and 2006. *Avian Diseases*, 52, 332-337.
- Lee, S.K., Sung, H.W. & Kwon, H.M. (2004). S1 glycoprotein gene analysis of infectious bronchitis viruses isolated in Korea. *Archives of Virology*, 149, 481-494.
- Liu, S., Zhang, X., Wang, Y., Li, C., Liu, Q., Han, Z., Zhang, Q., Kong, X. & Tong, G. (2009a). Evaluation of the protection conferred by commercial vaccines and attenuated heterologous isolates in China against the CK/CH/LDL/97I strain of infectious bronchitis coronavirus. *Veterinary Journal*, 179, 130-136.

- Liu, S.W., Zhang, Q.X., Chen, J.D., Han, Z.X., Liu, X., Feng, L., Shao, Y.H., Rong, J.G., Kong, X.G. & Tong, G.Z. (2006). Genetic diversity of avian infectious bronchitis coronavirus strains isolated in China between 1995 and 2004. *Archives of Virology*, *151*, 1133-1148.
- Liu, X.L., Su, J.L., Zhao, J.X. & Zhang, G.Z. (2009b). Complete genome sequence analysis of a predominant infectious bronchitis virus (IBV) strain in China. *Virus Genes*, *38*, 56-65.
- Lohr, J.E. (1976). Serologic differences between strains of infectious bronchitis virus from New Zealand, Australia, and the United States. *Avian Diseases*, *20*, 478-482.
- Lohr, J.E. (1977). Studies on avian infectious bronchitis virus in New Zealand. I. Serotypes. *New Zealand Veterinary Journal*, *25*, 48-51.
- Lohr, J.E. (1988). Infectious bronchitis in New Zealand, Asia, East Europe. In: E.F. Kaleta & U. Heffels-Redmann (Eds), *Proceedings of the 1st International Symposium on Infectious Bronchitis* (pp. 70-75). Rauschholzhausen, Germany.
- Mase, M., Inoue, T., Yamaguchi, S. & Imada, T. (2008). Existence of avian infectious bronchitis virus with a European-prevalent 4/91 genotype in Japan. *Journal of Veterinary Medical Science*, *70*, 1341-1344.
- Mase, M., Kawanishi, N., Ootani, Y., Murayama, K., Karino, A., Inoue, T. & Kawakami, J. (2010). A novel genotype of avian infectious bronchitis virus isolated in Japan in 2009. *Journal of Veterinary Medical Science*, *72*, 1265-1268.
- Mase, M., Tsukamoto, K., Imai, K. & Yamaguchi, S. (2004). Phylogenetic analysis of avian infectious bronchitis virus strains isolated in Japan. *Archives of Virology*, *149*, 2069-2078.
- Matthijs, M.G., van Eck, J.H., Landman, W.J. & Stegeman, J.A. (2003). Ability of Massachusetts-type infectious bronchitis virus to increase colibacillosis susceptibility

- in commercial broilers: a comparison between vaccine and virulent field virus. *Avian Pathology*, 32, 473-481.
- McFarlane, R. & Verma, R. (2008). Sequence analysis of the gene coding for the S1 glycoprotein of infectious bronchitis virus (IBV) strains from New Zealand. *Virus Genes*, 37, 351-357.
- Meir, R., Malkinson, M. & Weisman, Y. (1998). Characterization of IBV isolates in Israel using RT-PCR and RFLP. In: E.F. Kaleta & U. Heffels-Redmann (Eds), *Proceedings of the International Symposium on Infectious Bronchitis and Pneumovirus Infections in Poultry* (pp. 229-234). Rauschholzhausen, Germany.
- Meir, R., Rosenblut, E., Perl, S., Kass, N., Ayali, G., Perk, S. & Hemsani, E. (2004). Identification of a novel nephropathogenic infectious bronchitis virus in Israel. *Avian Diseases*, 48, 635-641.
- Meulemans, G., Carlier, M. C., Gonze, M., Petit, P. & Vandebroek, M. (1987). Incidence, characterisation and prophylaxis of nephropathogenic avian infectious bronchitis viruses. *Veterinary Record*, 120, 205-206.
- Minta, Z., Bugajek, P., Karpinska, E., Gough, R., Cavanagh, D., Mawditt, K., Cox, W. J., Bartnicka, B. & Wiercinski, J. (1998). Isolation of new strains of IBV from broiler chickens in Poland. In: E.F. Kaleta & U. Heffels-Redmann (Eds), *International Symposium on Infectious Bronchitis and Pneumovirus Infections in Poultry* (pp. 180-188). Rauschholzhausen, Germany.
- Mockett, A.P.A., Cavanagh, D. & Brown, T.D. (1984). Monoclonal antibodies to the S1 spike and membrane proteins of avian infectious bronchitis coronavirus strain Massachusetts M41. *Journal of General Virology*, 65, 2281-2286.

- Montassier, M.F.S., Brentano, L., Montassier, H.J. & Richtzenhain, L.J. (2008). Genetic grouping of avian infectious bronchitis virus isolated in Brazil based on RT-PCR/RFLP analysis of the S1 gene. *Pesquisa Veterinaria Brasileira*, 28, 190-194.
- Montassier, M.F.S., Brentano, L., Richtzenhain, L.J. & Montassier, H.J. (2006). Genetic diversity on S1 glycoprotein of avian infectious bronchitis virus strains isolated in Brazil between 1988–2000. In: U. Heffels-Redmann & E.F. Kaleta (Eds), *Proceedings of the Vth International Symposium on Corona- and Pneumovirus Infections* (pp. 119-131). Rauschholzhausen, Germany.
- Morley, A.J. & Thomson, D.K. (1984). Swollen-head syndrome in broiler chickens. *Avian Diseases*, 28, 238-243.
- Muneer, M.A., Newman, J.A., Goyal, S. M. & Ajmal, M. (1987a). Antibodies to avian infectious bronchitis virus in Pakistani chickens. *Poultry Science*, 66, 765-767.
- Muneer, M.A., Newman, J.A., Halvorson, D.A., Sivanandan, V. & Coon, C.N. (1987b). Effects of avian infectious bronchitis virus (Arkansas strain) on vaccinated laying chickens. *Avian Diseases*, 31, 820-828.
- Nix, W.A., Troeber, D.S., Kingham, B.F., Keeler, C.L. & Gelb, J. (2000). Emergence of subtype strains of the Arkansas serotype of infectious bronchitis virus in Delmarva broiler chickens. *Avian Diseases*, 44, 568-581.
- Parsons, D., Ellis, M.M., Cavanagh, D. & Cook, J. K.A. (1992). Characterisation of an infectious bronchitis virus isolated from vaccinated broiler breeder flocks. *Veterinary Record*, 131, 408-411.
- Pensaert, M. & Lambrechts, C. (1994). Vaccination of chickens against a Belgian nephropathogenic strain of infectious bronchitis virus B1648 using attenuated homologous and heterologous strains. *Avian Pathology*, 23, 631-641.

- Picault, J.P., Drouin, P., Guittet, M., Bennejean, G., Protais, J., L'Hospitalier, R., Gillet, J.P., Lamande, J. & Bachelier, A.L. (1986). Isolation, characterisation and preliminary cross-protection studies with a new pathogenic avian infectious bronchitis virus (strain PL-84084). *Avian Pathology*, 15, 367-383.
- Pohuang, T., Chansiripornchai, N., Tawatsin, A. & Sasipreeyajan, J. (2009). Detection and molecular characterization of infectious bronchitis virus isolated from recent outbreaks in broiler flocks in Thailand. *Journal of Veterinary Science*, 10, 219-223.
- Quiroz, M.A., Retana, A. & Tamayo, M. (1993). Determinacion de la presencia del serotipo Arkansas a partir de aislamientos del virus de bronquitos infecciosa aviar en Mexico. *IV Jornada Medico Avicola* (191-198). Mexico.
- Rimondi, A., Craig, M.I., Vagnozzi, A., Konig, G., Delamer, M. & Pereda, A. (2009). Molecular characterization of avian infectious bronchitis virus strains from outbreaks in Argentina (2001-2008). *Avian Pathology*, 38, 149-153.
- Roussan, D.A., Totanji, W.S. & Khawaldeh, G.Y. (2008). Molecular subtype of infectious bronchitis virus in broiler flocks in Jordan. *Poultry Science*, 87, 661-664.
- Sander, J.E., Jackwood, M.W. & Rowland, G.N. (1997). Protection by a commercial Arkansas-type infectious bronchitis virus vaccine against a field isolate of the same serotype. *Avian Diseases*, 41, 964-967.
- Seyfi Abad Shapouri, M.R., Mayahi, M., Assasi, K. & Charkhkar, S. (2004). A survey of the prevalence of infectious bronchitis virus type 4/91 in Iran. *Acta Veterinaria Hungarica*, 52, 163-166.
- Sheble, A., Sabry, M.Z., Davelaar, F.G., Burger, A.G., Khafagy, A.R., Moustafa, M.M. & Henna, M. (1986). Present status of infectious bronchitis in Egypt. *Journal of the Egyptian Veterinary Medical Association*, 46, 393-411.

- Shieh, H.K., Shien, J.H., Chou, H.Y., Shimizu, Y., Chen, J.N. & Chang, P.C. (2004). Complete nucleotide sequences of S1 and N genes of infectious bronchitis virus isolated in Japan and Taiwan. *The Journal of Veterinary Medical Science*, 66, 555-558.
- Shimazaki, Y., Horiuchi, T., Harada, M., Tanimura, C., Seki, Y., Kuroda, Y., Yagyu, K., Nakamura, S. & Suzuki, S. (2008). Isolation of 4/91 type of infectious bronchitis virus as a new variant in Japan and efficacy of vaccination against 4/91 type field isolate. *Avian Diseases*, 52, 618-622.
- Smith, H.W., Cook, J.K.A. & Parsell, Z.E. (1985). The experimental infection of chickens with mixtures of infectious bronchitis virus and *Escherichia coli*. *Journal of General Virology*, 66, 777-786.
- Song, C.S., Lee, Y.J., Kim, J.H., Sung, H.W., Lee, C.W., Izumiya, Y., Miyazawa, T., Jang, H.K. & Mikami, T. (1998). Epidemiological classification of infectious bronchitis virus isolated in Korea between 1986 and 1997. *Avian Pathology*, 27, 409-416.
- Terregino, C., Toffan, A., Beato, M. S., De Nardi, R., Vascellari, M., Meini, A., Ortali, G., Mancin, M. & Capua, I. (2008). Pathogenicity of a QX strain of infectious bronchitis virus in specific pathogen free and commercial broiler chickens, and evaluation of protection induced by a vaccination programme based on the Ma5 and 4/91 serotypes. *Avian Pathology*, 37, 487-493.
- Timms, L.M. & Bracewell, C.D. (1983). Cell mediated and humoral immune response of chickens to inactivated oil-emulsion infectious bronchitis vaccine. *Research in Veterinary Science*, 34, 224-230.
- Villarreal, L.Y., Brandao, P.E., Chacon, J.L., Assayag, M.S., Maiorka, P.C., Raffi, P., Saidenberg, A.B., Jones, R.C. & Ferreira, A.J. (2007a). Orchitis in roosters with

- reduced fertility associated with avian infectious bronchitis virus and avian metapneumovirus infections. *Avian Diseases*, 51, 900-904.
- Villarreal, L.Y., Brandao, P.E., Chacon, J.L., Saidenberg, A.B., Assayag, M.S., Jones, R.C. & Ferreira, A.J. (2007b). Molecular characterization of infectious bronchitis virus strains isolated from the enteric contents of Brazilian laying hens and broilers. *Avian Diseases*, 51, 974-978.
- Villarreal, L.Y., Sandri, T.L., Souza, S.P., Richtzenhain, L.J., de Wit, J.J. & Brandao, P.E. (2010). Molecular epidemiology of avian infectious bronchitis in Brazil from 2007 to 2008 in breeders, broilers, and layers. *Avian Diseases*, 54, 894-898.
- Wang, C.H. & Huang, Y.C. (2000). Relationship between serotypes and genotypes based on the hypervariable region of the S1 gene of infectious bronchitis virus. *Archives of Virology*, 145, 291-300.
- Wang, C.H. & Tsai, C.T. (1996). Genetic grouping for the isolates of avian infectious bronchitis virus in Taiwan. *Archives of Virology*, 141, 1677-1688.
- Wang, L., Junker, D. & Collisson, E.W. (1993). Evidence of natural recombination within the S1 gene of infectious bronchitis virus. *Virology*, 192, 710-716.
- Wei, P.L.M., Wei, Z.J., Wang, X.Y., Mo, M.L. & Chen, Q.Y. (2009). Genotyping and serotyping of Guangxi IBV isolates during 1985-2008. In: E.F. Kaleta & U. Heffels-Redmann (Eds), *Proceedings of the VIth International Symposium on Avian Corona- and Pneumoviruses and Complicating Pathogens* (pp. 59-66). Rauschholzhausen, Germany.
- Worthington, K.J., Currie, R.J. & Jones, R.C. (2008). A reverse transcriptase-polymerase chain reaction survey of infectious bronchitis virus genotypes in Western Europe from 2002 to 2006. *Avian Pathology*, 37, 247-257.

- Worthington, K.J. & Jones, R.C. (2006). An update on the European RT-PCR IBV survey and recent findings on a novel IBV genotype. In: U. Heffels-Redmann & E.F. Kaleta (Eds), *Proceedings of the Vth International Symposium on Corona- and Pneumovirus Infections* (pp. 176-188). Rauischholzhausen, Germany.
- Wu, Z.Q., Yang, Q.W., Fu, C., Zhao, X.Y. & Ignjatovic, J. (1998). Antigenic and immunogenic characterization of infectious bronchitis virus strains isolated in China between 1986 and 1995. *Avian Pathology*, 27, 578-585.
- Yu, L., Jiang, Y., Low, S., Wang, Z., Nam, S.J., Liu, W. & Kwangac, J. (2001a). Characterization of three infectious bronchitis virus isolates from China associated with proventriculus in vaccinated chickens. *Avian Diseases*, 45, 416-424.
- Yu, L., Wang, Z., Jiang, Y., Low, S. & Kwang, J. (2001b). Molecular epidemiology of infectious bronchitis virus isolates from China and Southeast Asia. *Avian Diseases*, 45, 201-209.
- Zanella, A., Coaro, R., Fabris, G., Marchi, R. & Lavazza, A. (2000). Avian infectious bronchitis virus: isolation of an apparently new variant in Italy. *Veterinary Record*, 146, 191-193.
- Zanella, A., Lavazza, A., Marchi, R., Moreno Martin, A. & Paganelli, F. (2003). Avian infectious bronchitis: characterization of new isolates from Italy. *Avian Diseases*, 47, 180-185.
- Zou, N.L., Zhao, F.F., Wang, Y.P., Liu, P., Cao, S.J., Wen, X.T. & Huang, Y. (2010). Genetic analysis revealed LX4 genotype strains of avian infectious bronchitis virus became predominant in recent years in Sichuan area, China. *Virus Genes*, 41, 202-209.

Zulperi, Z.M., Omar, A.R. & Arshad, S.S. (2009). Sequence and phylogenetic analysis of S1, S2, M, and N genes of infectious bronchitis virus isolates from Malaysia. *Virus Genes*, 38, 383-391.

For Peer Review Only

Figure 1.

Phylogenetic tree of a selection of IBV variants from different parts of the world showing the wide diversity of IBV. The tree is based on comparison of the partial S gene (between nucleotides 20447 and 20924 (numbering compared to the genome of Ark DPI ((Ammayappan et al., 2008), coding for a part of the S1 glycoprotein including the highly variable regions 1 and 2). The phylogenetic tree analysis was conducted by neighbour-joining method using bootstrap analysis (100 replications).




Table 1. Average level of genetic homology for 4 IBV strains using different nt window sizes across the entire S1 gene of different genotypes/serotypes (Adznan et al, 1997)

Number of nucleotides per window that were used to calculate the homology	Number of windows	Lowest homology ^a	Highest homology	Difference in homology between highest and lowest windows (%)
per 1	1600	0	100	100
per 10 ^b	160	42.5	100	57.5
per 50	32	64.5	91.0	26.5
per 100	16	68.0	86.8	18.2
per 200	8	69.6	83.9	14.3
per 400	4	73.2	81.6	8.4
per 800 ^c	2	76.8	80.2	3.4
All 1600	1	78.5*		

^aThe extent of nucleotide identity between the strains was expressed on nucleotide level as follows: score 100% when all four strains had the same nucleobase (in the copy DNA); score 75% when three out of four strains had the same nucleobase; score 50% when two strains had the same nucleobase and the two other strains had another nucleobase in common; score 25% when two strains had the same nucleobase and the two other strains each had another different nucleobase; score 0% when all strains had a different nucleobase at that position of the genome. In this comparison, a missing nucleotide (by deletion) was considered to be a different nucleobase.

^bHomology in nucleotides 1-10, 11-20, 21-30, to 1591-1600.

^cHomology in nucleotides 1-800 and 801-1600.

For Peer Review Only




Figure 2. Correlation between the level of homology in the S1 region of the S gene of IBV strains and the level of cross-protection between these strains as reported in seven publications.




Figure 3. Average level of homology between four strains of IBV in 160 different windows of 10 nucleotides of nucleotides 1-1600 of the S1 region on the S gene of different genotypes/serotypes (Adznar et al., 1997).




Figure 4. Average level of homology between four strains of IBV in 16 different windows of 100 nucleotides of nucleotides 1-1600 of the S1 region on the S gene of different genotypes/serotypes (Adznar et al., 1997).




Figure 5. Average level of homology between four strains of IBV in four different windows of 400 nucleotides of nucleotides 1-1600 of the S1 region on the S gene of different genotypes/serotypes (Adznaret al., 1997).