



HAL
open science

Double-frequency noise generation by surface gravity waves in finite depth: gravity, acoustic and seismic modes

Fabrice Ardhuin, Thomas H. C. Herbers

► To cite this version:

Fabrice Ardhuin, Thomas H. C. Herbers. Double-frequency noise generation by surface gravity waves in finite depth: gravity, acoustic and seismic modes. 2012. hal-00711326v1

HAL Id: hal-00711326

<https://hal.science/hal-00711326v1>

Preprint submitted on 23 Jun 2012 (v1), last revised 29 Oct 2012 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Double-frequency noise generation by surface gravity waves in finite depth: gravity, acoustic and seismic modes

FABRICE ARDHUIN¹ AND T. H. C. HERBERS²

¹Ifremer, Laboratoire d'Océanographie Spatiale, Plouzané, France

²Department of Oceanography, Naval Postgraduate School, Monterey, California, 93943, USA

(Received ?; revised ?; accepted ?. - To be entered by editorial office)

Oceanic observations, even in very deep water, and atmospheric pressure or seismic records, from anywhere on Earth, contain noise with dominant periods between 3 and 10 seconds, that can be related to ocean waves. This noise has been explained by a non-linear wave-wave interaction mechanism, and takes the form of surface gravity waves, acoustic or seismic waves. Previous theoretical works on seismic noise focused on surface (Rayleigh) waves, while acoustic studies have generally considered an unbounded ocean, and atmospheric noise theory has been limited to vertical sound propagation. Here we show how directional wave spectra, through the same theory, produce seismic waves (including Rayleigh and body waves) and associated acoustic-gravity waves in the ocean. In the limit of slow phase speeds, the theory gives the known and well-verified gravity wave result, previously derived for an incompressible ocean. This consistent approach makes it possible to account for noise generation by finite depth surface gravity waves, which is particularly relevant for the noise periods larger than 10 s, and requires the consideration of the direct wave-induced pressure at the ocean bottom, a contribution previously overlooked in the context of seismic noise. For acoustic and seismic modes, that are much faster than the ocean waves, the noise level of each mode is given by a mode-specific amplification of the same wave-induced pressure field near the zero wavenumber.

Key words: surface gravity waves

1. Introduction

Ocean waves generate noise in a wide range of frequencies. The upper end of the spectrum, at least for $f > 100$ Hz, is dominated by wave breaking and associated bubbles (Knudsen *et al.* 1948), whereas the lower frequency part, nominally $f < 2$ Hz is mostly expected to be caused by the nonlinearity of the hydrodynamic equations, on which we focus here. The general sound generation by fluid flows was described theoretically by Lighthill (1952). Longuet-Higgins (1950) showed how seismic waves can be generated by the same process, with noise radiating along the Earth's crust in the form of Rayleigh waves. That theory was extended to random waves by Hasselmann (1963), and later cast in the more general framework of wave-wave interactions (Hasselmann 1966). Interest in seismic noise has risen sharply over the last few years with the enforcement of the Comprehensive Nuclear Test Ban Treaty, and since Shapiro *et al.* (2005) have demonstrated that seismic noise correlation could provide a unique monitoring method for the properties of the solid Earth. Recent numerical models based on the Longuet-Higgins–Hasselmann (LH-H) theory for Rayleigh wave generation have shown good agreement

of modeled seismic noise spectra with observations (Kedar *et al.* 2008; Ardhuin *et al.* 2011). It is still unclear whether most of the uncertainties on the modeled noise level can be attributed to errors in the seismic sources, associated with a poorly controlled directional distribution of the wave spectrum, or to errors in the seismic propagation. Also, many observations have focused on body waves for which no complete theory has been proposed to date. Vinnik (1973) did propose a theory for compressional (P) waves, but he did not consider the important effect of the water layer.

The level of acoustic noise has also been investigated as a potential source of information on the poorly known directional spectrum of short gravity waves (Tyler *et al.* 1974). Farrell & Munk (2008, 2010) and Duennebieer *et al.* (2012) showed a large variability of the spectral level in the frequency range 0.1–50 Hz that is clearly related to the sea state. Their interpretation of the data, following previous studies of underwater noise (e.g. Hughes 1976; Lloyd 1981; Kibblewhite & Ewans 1985), is based on sound generated by waves in an unbounded ocean. Although this simplified approach is reasonable for high frequency noise, the neglected reflection from the seafloor and subsequent reverberations in the bounded ocean may strongly amplify the lower frequency resonant modes.

Given the renewal of interest in seismic and acoustic noise, we found it appropriate to revisit Hasselmann’s theory. We thus illustrate, correct, and add a few missing aspects. These corrections include important terms for intermediate and shallow water that have not been considered before for the compressible conditions, although they were verified in the incompressible limit (Herbers & Guza 1994). The compressible equations of motion are used to derive a consistent solution in terms of gravity, acoustic and seismic modes, including both surface Rayleigh waves and compressional (P) or shear (S) body waves. This solution for body waves has not been given before.

Building on the seismo-acoustic paradigm proposed by Arrowsmith *et al.* (2010), the basic idea of the present paper is that all modes of motion can be excited by ocean surface gravity waves (OSGW) of any frequency. For a given pair of interacting frequencies f and f' , the frequency of the generated noise is $f + f'$, and the different types of waves are only distinguished by their phase velocity, or equivalently by their horizontal wavenumber K which is the norm of the vector sum of the wave numbers, $\mathbf{K} = \mathbf{k} + \mathbf{k}'$ of the interacting OSGW, as shown in figure 1. This type of wave-wave interaction is one of the lowest order interactions (Hasselmann 1966).

In the physical space, one such interaction excites waves with horizontal wavelength $L_h = 2\pi/K$, and different vertical patterns in the atmosphere, ocean and crust, due to the very different speeds for compression waves in these three media. Figure 2 shows how the same forcing can give almost vertical-propagating waves in the atmosphere, waves that propagate almost horizontally in the ocean, and evanescent waves in the crust. In practice, the broad-banded wave spectrum results in a superposition of all possible waves, propagating or evanescent, that radiate in all directions. For wavelengths much larger than those of the interacting OSGW and for a spatially homogeneous wave field, the excited waves have a spectrum similar to that excited by a point force exerted on the sea surface (Longuet-Higgins 1950). Table 1 points directly to the main theoretical results of the present paper.

2. Water waves theory and noise sources

Here we give only a brief derivation of the solution, which is a straightforward compressible extension to the solution given by Hasselmann (1962). Water column motions are expanded in powers of the sea surface slope with a linear motion for which compressible effects may be neglected (Longuet-Higgins 1950), and a second order motion




FIGURE 1. (a) Typical distribution of speeds: α_w is for sound waves in the ocean, β and α_c are shear and compression speeds in the upper crust. These velocities separate the horizontal phase speeds in four domains. From slow to fast, these are the acoustic-gravity (AG) domain, the Rayleigh (R) wave domain, and two body wave domains (S only, and P and S together) (b) For any fixed frequency, the four domains correspond to 4 concentric regions in the wavenumber plane. For three selected frequencies in the infragravity, dominant and high frequency ranges of the forcing wave field, the limiting wavelengths between the four domains are indicated. One example of interaction (black vectors) is shown with two gravity wave modes that interact to generate a Rayleigh waves.




FIGURE 2. Left, schematic of second-order pressure field, in colours, associated with double-frequency noise, forced by the interaction of a single pair of directionally opposing monochromatic wave trains on the sea surface that form periodic groups with wave number K . The individual waves in the (near standing) group advance very slowly, at the OSWG phase speed. However, in this particular case, the group as a whole travels slightly faster than the sound speed in water, and slower than the crust shear and compression velocity. As a result, the crust elastic waves are evanescent while ocean and atmospheric waves propagate. The number of waves in the group was reduced for visibility, and for the same reason the amplitudes of the sea surface and bottom elevations have been exaggerated as well as the pressure fluctuation in the atmosphere relative to those in the ocean. Right, schematic of ocean waves with a relatively broad spectrum, giving rise to the interaction of all possible pairs of wave trains and noise radiation in all directions.

TABLE 1. Summary of theoretical expressions for the gravity, acoustic and seismic noise fields.

	infinite water depth	finite water depth h
Surface forcing field	eq. (2.26)	eq. (2.25)
Bottom forcing field	n.a.	eq. (3.7)
gravity-like modes	eq. (3.11)	not given
acoustic spectrum	eq. (3.10)	eq. (4.23)
microbaroms	eq. (3.19)	eq. (3.19)
seismic source (R)	n.a.	eq. (4.32)
seismic source (P)	n.a.	eq. (4.39)
seismic source (S)	n.a.	eq. (4.41)
seismic spectrum (R)	n.a.	eq. (4.37)
seismic spectrum (P)	n.a.	eq. (4.46)

with pressure p_2 and velocity potential ϕ_2 . It is for this second order motion that we are extending previous results. Following again Longuet-Higgins (1950) we will see that ϕ_2 approximately obeys the same linear acoustic wave equation as ϕ_1 , so that the forcing of the double-frequency noise is only due to boundary conditions. In contrast to Hasselmann (1963), we consider here the general finite depth expression for the surface pressure forcing $\widehat{p}_{2,\text{surf}}$, and in the boundary condition at the ocean bottom the additional forcing $\widehat{p}_{2,\text{bot}}$ that accounts for the Bernoulli effect of the near-bed orbital wave motion. Compared to the incompressible second order solution of Hasselmann (1962), the differences comes from compressibility of the water and the elasticity of the bottom.

2.1. Equations of motion

We decompose the water density into a mean value ρ_w and a perturbation $\rho \ll \rho_w$. Neglecting stratification effects due to temperature and salinity, the fluctuation in pressure p and water density ρ are related by an equation of state, which involves the speed of sound in water α_w (Lighthill 1978, eq. 32).

$$\frac{dp}{dt} = \alpha_w^2 \frac{d\rho}{dt}, \quad (2.1)$$

We assume the oceanic motion to be irrotational so that the velocity field is given by the gradient of the velocity potential ϕ . Because of the importance of the apparent gravity acceleration g , which defines the vertical axis, we separate horizontal and vertical components using vectors and gradient operators in the horizontal plane, e.g. $\mathbf{u} = \nabla\phi = (\partial\phi/\partial x, \partial\phi/\partial y)$ and $w = \partial\phi/\partial z$. The conservation of mass of sea water is

$$\frac{d\rho}{dt} = -\rho_w \nabla^2 \phi - \rho_w \frac{\partial^2 \phi}{\partial z^2}. \quad (2.2)$$

Equations (2.2)–(2.1) can be combined to eliminate ρ ,

$$\frac{dp}{dt} = -\rho_w \alpha_w^2 \left[\nabla^2 + \frac{\partial^2}{\partial z^2} \right] \phi \quad (2.3)$$

From eq. (2.1), the water density is only a function of the pressure. Hence the two unknowns p and ϕ are also related by the momentum conservation equation, which can be cast in the form of Bernoulli's equation (see e.g. Lamb 1932, section 20),

$$\frac{\partial\phi}{\partial t} = -\frac{1}{2} \left[|\nabla\phi|^2 + \left(\frac{\partial\phi}{\partial z} \right)^2 \right] - \frac{p}{\rho_w} - gz + C(t). \quad (2.4)$$

The boundary conditions at the surface $z = \zeta$ are given by the continuity of pressure and vertical velocity

$$p = p_a \quad (2.5)$$

$$\frac{\partial \phi}{\partial z} = \nabla \phi \cdot \nabla \zeta + \frac{\partial \zeta}{\partial t} \quad \text{at } z = \zeta, \quad (2.6)$$

with the atmospheric pressure p_a . This expression is translated to the mean sea level $z = 0$ with a Taylor expansion for ϕ

$$\frac{\partial \zeta}{\partial t} - \frac{\partial \phi}{\partial z} \simeq -\nabla \phi \cdot \nabla \zeta + \zeta \frac{\partial^2 \phi}{\partial z^2} \quad \text{at } z = 0. \quad (2.7)$$

Following Longuet-Higgins (1950), we shall now expand the solution in powers of the surface slope, with the sea surface elevation as ζ_1 , associated with linear waves, and a non-linear correction ζ_2 such that at the sea surface $|\zeta_2| \ll |\zeta_1|$ (see also e.g. Hasselmann 1962),

$$\zeta = \zeta_1 + \zeta_2, \quad \phi = \phi_1 + \phi_2. \quad (2.8a,b)$$

2.2. Linear solution

Compressibility effects in the linear solution are negligible for our purpose (see Longuet-Higgins 1950, eq. 123), so that we may use

$$\zeta_1 = \sum_{\mathbf{k},s} Z_{1,\mathbf{k}}^s e^{i(\mathbf{k}\cdot\mathbf{x}-s\sigma t)}, \quad \phi_1 \simeq \sum_{\mathbf{k},s} -i \frac{sg \cosh(kz + kh)}{\sigma \cosh(kh)} Z_{1,\mathbf{k}}^s e^{i(\mathbf{k}\cdot\mathbf{x}-s\sigma t)} \quad (2.9a,b)$$

where k is the norm of the horizontal wavenumber vector \mathbf{k} , s is a sign index equal to -1 or 1, so that $s = 1$ corresponds to waves propagating in the direction of the vector \mathbf{k} , and $s = -1$ corresponds to the opposite direction. The radian frequency σ is given by the dispersion relation for linear waves (de Laplace 1776) ,

$$\sigma = \sqrt{gk \tanh(kh)}, \quad (2.10)$$

giving the group speed

$$C_g = \frac{\partial \sigma}{\partial k} = \frac{\sigma}{k} \left[\frac{1}{2} + \frac{kh}{\sinh(2kh)} \right]. \quad (2.11)$$

2.3. The negligible near-surface forcing

Eliminating p between (2.3) and (2.4), we obtain the acoustic wave equation

$$\alpha_w^2 \left[\nabla^2 + \frac{\partial^2}{\partial z^2} \right] \phi = \frac{d}{dt} \left[\frac{\partial \phi}{\partial t} + \frac{1}{2} \left(|\nabla \phi|^2 + \left(\frac{\partial \phi}{\partial z} \right)^2 \right) \right], \quad (2.12)$$

where the gravity term has been removed by our approximation of a constant mean density. That term is usually neglected eventually in the solution (Stoneley 1926).

Our eq. (2.12) corresponds to eq. (130) in Longuet-Higgins (1950), who showed that the non-linear terms yield a contribution (his F terms) that is of the order of $g/(2\alpha_w^2 k)$ times the other terms. Even for very long (infragravity) waves with a wavelength of 50 km, this factor is 0.02, so that we shall neglect this source of wave forcing. This leads to a linear wave equation for the second order velocity potential

$$\frac{\partial^2 \phi_2}{\partial t^2} \simeq \alpha_w^2 \left[\nabla^2 + \frac{\partial^2}{\partial z^2} \right] \phi_2, \quad \text{for } -h \leq z \leq 0 \quad (2.13)$$

2.4. Surface forcing

Because the compressibility only affects the mass conservation equation, it does not modify the kinematic and dynamic boundary conditions. These are given by Hasselmann (1962). The unknown ζ is eliminated from the linear terms by adding $\partial(2.4)/\partial t$ evaluated with (2.5), and $g \times$ (2.7). This combination of kinetic and dynamic boundary condition gives an equation for the velocity potential to second order in the wave slope, valid at $z = 0$. Keeping the lowest order non-linear terms we have,

$$\left(\frac{\partial^2}{\partial t^2} + g\frac{\partial}{\partial z}\right)\phi_2 = -g \left[\nabla\phi_1 \cdot \nabla\zeta_1 + \zeta_1 \frac{\partial^2\phi_1}{\partial z^2} \right] - \frac{1}{2} \frac{\partial}{\partial t} \left[|\nabla\phi_1|^2 + \left(\frac{\partial\phi_1}{\partial z}\right)^2 + 2\zeta_1 \frac{\partial^2\phi_1}{\partial t\partial z} \right]. \quad (2.14)$$

We now give the explicit form of the right hand side using eqs. (2.9). As noted by Hasselmann (1963), we may rewrite (2.14) as

$$\left(\frac{\partial^2}{\partial t^2} + g\frac{\partial}{\partial z}\right)\phi_2 = -\frac{1}{\rho_w} \frac{\partial \hat{p}_{2,\text{surf}}}{\partial t}. \quad (2.15)$$

Namely, our problem is equivalent to the effect of a pressure field $\hat{p}_{2,\text{surf}}$ applied at $z = 0$. Using the linear solution (2.9),

$$\hat{p}_{2,\text{surf}} = \rho_w \sum_{\mathbf{k}, s, \mathbf{k}', s'} D_z(\mathbf{k}, s, \mathbf{k}', s') Z_{1,\mathbf{k}}^s Z_{1,\mathbf{k}'}^{s'} e^{i\Theta(\mathbf{k}, \mathbf{k}', s, s')} \quad (2.16)$$

with the phase function of interacting wave trains defined by

$$\Theta(\mathbf{k}, \mathbf{k}', s, s') = [(\mathbf{k} + \mathbf{k}') \cdot \mathbf{x} - (s\sigma + s'\sigma')t] \quad (2.17)$$

and coupling coefficient $D_z(\mathbf{k}, s, \mathbf{k}', s')$ given by eq. (A 1).

The shape of this pressure pattern is well understood when the full sum in eq. (2.16) is reduced to only two interacting deep water wave trains with amplitudes a and a' and slightly different frequencies σ and σ' , traveling in opposite directions 3. For $kh \gg 1$ we may use $D_z(\mathbf{k}, 1, -\mathbf{k}, 1) = -2\sigma^2$ and $D_z(\mathbf{k}, 1, \mathbf{k}, 1) = 0$. Defining $K = k - k'$, the equivalent surface pressure is given by the sum interaction,

$$\hat{p}_2 = -2\rho_w\sigma^2aa' \{\cos[Kx + 2\sigma t]\}. \quad (2.18)$$

This expression generalizes the result given by Longuet-Higgins (1950) for equal wave periods. The corresponding difference interaction yields a short (wave number $2k$) infragravity wave that will not be considered here (see e.g. Hasselmann 1963).

For wave directions nearly opposite, instead of exactly opposite, this first term can propagate in any horizontal direction, given by the direction of \mathbf{K} .

2.5. Noise spectrum and finite depth effects

Both LH50 and H63 have used the deep-water approximation

$$\hat{p}_{2,\text{surf}} \simeq \rho_w \left[|\nabla\phi|^2 + \left(\frac{\partial\phi}{\partial z}\right)^2 \right] \quad \text{valid for } kh \gg 1, \quad (2.19)$$

instead of the more complex but more general form given by eq. (2.16). We will illustrate the important differences between these two expressions for $kh < 1$, by considering the forcing of very long components. Indeed, sound waves in the ocean have velocities in excess of 1.4 km s^{-1} . As a result, regardless of the vertical wavenumber l , the horizontal




FIGURE 3. Generation of supersonic wave groups (in blue) by the superposition of two opposing deep water monochromatic wave trains of nearly equal periods T and T' . Each curve shows the surface elevation of the individual wave trains (black and red) or their combination (blue). The length of the group is 12 km, and propagates in the same direction as the wave train with the shortest wavelength. If the wave trains do not face exactly, then the group propagates in the direction of the vector $\mathbf{K} = \mathbf{k} + \mathbf{k}'$. The red and black dots are fixed on wave crests of each train, and move 100 times slower than the blue group.

wavenumber vector is relatively small $K = \sqrt{(\omega_s^2/\alpha_w^2 - l^2)} < \omega_s/\alpha_w$. Thus the acoustically noisy wave interactions verify $K \ll k$, which gives $\mathbf{k}' \simeq -\mathbf{k}$, and $f \simeq f'$ with $f_s \simeq 2f$. We may thus focus on the estimation of the spectrum of $\hat{p}_{2,\text{surf}}$ at $\mathbf{K} \simeq 0$.

For this purpose we introduce the spectral density of this surface pressure in the three spectral dimensions (K_x, K_y, f_s) . Using the the Fourier amplitude $\hat{p}_2(\mathbf{K}, f_s)$ of the forcing pressure \hat{p}_2 , with wavenumber vector \mathbf{K} and frequency f_s ,

$$F_{p2,\text{surf}}(\mathbf{K}, f_s) = 2 \lim_{|\mathbf{K}| \rightarrow 0, df_s \rightarrow 0} \frac{|\hat{p}_{2,\text{surf}}(\mathbf{K}, f_s)|^2}{dK_x dK_y df_s}, \quad (2.20)$$

The factor 2 in the expression makes this a single-sided spectrum, with non-zero values only for $f_s \geq 0$. This spectral density is in a three-dimensional spectral space, with S.I. units of $\text{Pa}^2 \text{m}^2 \text{s}$, it is denoted F_{p3D} in Ardhuin *et al.* (2011).

Using (2.16) the surface pressure spectrum can be expressed in terms of quadratic products of the (linear) sea surface elevation spectrum

$$E(k_x, k_y) = 2 \lim_{|d\mathbf{k}| \rightarrow 0} \frac{|Z_{1,\mathbf{k}}^+|^2}{dk_x dk_y} \quad (2.21)$$

with a coupling coefficient from (A 1) that simplifies for $K \simeq 0$ to

$$D_z(\mathbf{k}, 1, -\mathbf{k}, 1) = -2\sigma^2 \left[1 + \frac{1}{4 \sinh^2(kh)} \right]. \quad (2.22)$$

To transform the spectra to frequency-direction space we use the Jacobian transformation

$$E(f, \theta) = \frac{2\pi k}{C_g} E(k_x, k_y). \quad (2.23)$$

We now introduce the directional distribution M such that $E(f, \theta) = E(f)M(f, \theta)$, and

we define the directional integral

$$I(f) = \int_0^\pi M(f, \theta)M(f, \theta + \pi)d\theta. \quad (2.24)$$

With these notations we finally obtain

$$\begin{aligned} F_{p2,\text{surf}}(\mathbf{K} \simeq 0, f_s) &\simeq \rho_w^2 D_z^2 \int E(f, \theta)E(f, \theta + \pi) \frac{C_g^2 dk_x dk_y}{k^2 4\pi^2 df_s} \\ &\simeq \rho_w^2 g^2 f_s E^2(f) I(f) \tanh^2(kh) \left[1 + \frac{2kh}{\sinh(2kh)} \right] \left[1 + \frac{1}{4 \sinh^2(kh)} \right]^2 \end{aligned} \quad (2.25)$$

In deep water ($kh \gg 1$), the equivalent surface pressure is $\widehat{p}_{2,\text{surf}} \simeq \rho_w(u_1^2 + w_1^2)$, and its spectrum (Hasselmann 1963; Ardhuin *et al.* 2011)

$$F_{p2,\text{surf}}(\mathbf{K} \simeq 0, f_s) = \rho_w^2 g^2 f_s E^2(f) I(f). \quad (2.26)$$

As shown in figure 4, this spectral density of surface pressure is, in finite depth, up to four times smaller than the deep water approximation obtained from eq. (2.19) and used by Webb (2007) and Tanimoto (2007). This is because the surface pressure actually combines two terms. One is $\rho_w(u_1^2 + w_1^2)/2$, from the momentum conservation equation – which here takes the form of the Bernoulli equation (2.4) – and the other comes from the non-linearity of the surface boundary condition (2.7). As kh goes to zero, both this second term and w_1 become negligible compared to the $\rho_w u_1^2/2$ term from eq. (2.4).

This effect is not important for the dominant microseismic peak, generated by waves of period about 10 s, but it may be important for the much longer waves, known as hum, driven by free infragravity waves. However, in that case, one should also consider the direct action of the wave-induced pressure on the bottom.

2.6. Additional bottom forcing

In the limit $kh \rightarrow 0$, the nonlinearity of the kinematic boundary condition (2.7) becomes negligible compared to the $\rho_w u_1^2/2$ term from eq. (2.4), so that the combined surface boundary condition (2.14) now only expresses the momentum conservation: the water is accelerated horizontally by the advection due to the orbital velocity of the first order waves, while the pressure is zero at the sea surface. However, $kh \rightarrow 0$ makes both the orbital velocity u_1 and the second order potential ϕ_2 uniform over the depth. Because we have the same momentum balance at the sea surface and bottom, we thus have the same pressure field, which is uniformly zero. Therefore noise only exists as velocity in the water column, without an associated pressure signal.

For any value of kh , the coupling coefficient given by eq. (2.22) differs from the full second order coefficient for the bottom pressure (e.g. eq. 4 in Herbers & Guza 1991), which also involves the Bernoulli head (the bracket in eq. 2.4). However, that extra term is also relevant to the generation of seismic noise due to the bottom boundary condition that couples the solid crust to the water column. Indeed, the second-order pressure perturbation at the bottom writes,

$$p_2(-h) = -\rho_w \frac{\partial \phi_2}{\partial t} + \widehat{p}_{2,\text{bot}}, \quad (2.27)$$

where the Bernoulli head contribution to the pressure can be expressed from the first




FIGURE 4. Compared to deep water, the source of seismic noise power in finite water depth is amplified by a coefficient $G(kh) = \tanh^2(kh) [1 + 2kh/\sinh(2kh)]$, when accounting for both the surface and bottom forcing. This is very different from the approximation that considers the surface forcing only (leading to eq. 2.25), even more so when using its deep water approximation (eq. 2.19) which gives a factor 4 difference for $kh \ll 1$.

order wave amplitudes,

$$\hat{p}_{2,\text{bot}} = \rho_w \sum_{\mathbf{k}, s, \mathbf{k}', s'} D_{pb}(\mathbf{k}, s, \mathbf{k}', s', z = -h) Z_{1,\mathbf{k}}^s Z_{1,\mathbf{k}'}^{s'} e^{i\Theta(\mathbf{k}, \mathbf{k}', s, s')}, \quad (2.28)$$

with a coupling coefficient D_{pb} given by eq. (A 2).

We may interpret the bottom pressure (2.27) as the sum of the surface forcing $\hat{p}_{2,\text{surf}}$ transmitted to the bottom by ϕ_2 , and a direct effect of the Bernoulli head at the bottom which is an additional forcing $\hat{p}_{2,\text{bot}}$ that partly cancels $\hat{p}_{2,\text{surf}}$.

We shall see in the next section that the forcing term for seismic noise is $\hat{p}_{2,\text{surf}} + \cos(lh)\hat{p}_{2,\text{bot}}$, with $l \leq K \ll k$ the vertical wavenumber in the water. For shallow water gravity waves, $kh \ll 1$ and thus $\cos(lh) \simeq 1$ so that the effective forcing term becomes $\hat{p}_{2,\text{surf}} + \hat{p}_{2,\text{bot}}$, which equals the bottom pressure in the incompressible limit. The shallow water asymptote of the spectrum of this total forcing term is very different from the surface pressure only. Compared to eq. (2.25), the $[1 + 0.25/\sinh^2(kh)]^2$ factor now replaced by 1. For $kh \ll 1$, this is a factor $(kh)^4/16$ smaller, as shown in figure 4. This asymptote is relevant for the hum, the noise with periods larger than 30 s, which is believed to be driven by infragravity waves (Webb 2007). The source of this hum is

attenuated by several orders of magnitude on the continental shelves and not amplified according to the deep water approximation given by eq. (2.19).

3. From surface pressure spectrum to noise spectra

3.1. Dispersion relation and modes

The problem of noise generation by waves has been reduced to that of noise generation by an equivalent surface pressure field $\widehat{p}_{2,\text{surf}}(x, y, t)$, with the possible addition of a bottom pressure field $\widehat{p}_{2,\text{bot}}(x, y, t)$ for finite depths. From a statistical point of view, this equivalent surface pressure is fully represented by its spectrum, $F_{p2,\text{surf}}(\mathbf{K}, f_s)$ where $\mathbf{K} = \mathbf{k} + \mathbf{k}'$ is the sum of the interacting wavenumbers and $f_s = f \pm f'$ is the sum or difference of the interacting frequencies. Subjected to this surface forcing, our linearized wave equation (2.13) will have linear solutions. In particular, any homogeneous or evanescent solution will take the following form

$$\phi_2 \propto \exp[i(K_x x + K_y y + lz - \omega_s t)], \quad (3.1)$$

which, replaced in the wave equation, gives the dispersion relation

$$\{-\omega_s^2 + \alpha_w^2 [K^2 + l^2]\} = 0, \quad (3.2)$$

where $\omega_s = 2\pi f_s$. Both ω_s and $K = |\mathbf{K}|$ are imposed by the forcing, so that the magnitude of the complex vertical wavenumber l is given by

$$l = K \sqrt{\frac{\omega_s^2}{K^2 \alpha_w^2} - 1}, \quad (3.3)$$

which yields

$$\phi_2 = \left(C e^{il(z+h)} + D e^{-il(z+h)} \right) e^{i\Theta}, \quad \Theta = (K_x x + K_y y - \omega_s t) \quad (3.4a,b)$$

where C and D are the bottom amplitudes of the upward and downward propagating waves, determined by the surface and bottom boundary conditions.

There are two classes of solutions. Those for which $\omega_s/K \leq \alpha_w$ and thus l is imaginary: these are 'gravity-acoustic' modes with an amplitude that decays exponentially from the surface. For the shortest components, we have $\omega_s/K \ll \alpha_w$ and thus $l = ik$, corresponding to the incompressible limit in which only gravity is important. The other class of solutions, for which $\omega_s/K \geq \alpha_w$ and l is real, are the acoustic modes that propagate along both vertical and horizontal dimensions, as illustrated in figure 5.

The equations of elasticity in the crust yield the same wave equation, one for compressional and another for shear motions, with α_w replaced by the crusts compressional wave speed α_c and shear wave speed β_s , respectively (e.g. Aki & Richards 2002). Since we have $\alpha_w < \beta_s < \alpha_c$, we can distinguish four different regimes depending on the value of the horizontal phase speed ω_s/K , relative to these three velocities (fig. 1).

In the family of gravity-acoustic modes, there is no resonance in the forcing, namely there are no free waves with $\mathbf{K} = \mathbf{k} + \mathbf{k}'$ and $f_s = f \pm f'$ and thus the motion is locked with the forcing wave groups, as verified by Herbers & Guza (1994). The acoustic modes propagate obliquely to the vertical away from the surface, and interact with the ocean bottom (figure 5). The seismic Rayleigh waves modes propagate horizontally, combining an acoustic-like motion in the water and evanescent elastic waves in the crust, illustrated by figure 2. The coupling with the bottom motion selects a few resonant modes that dominate the solution, with an energy that grows linearly with the propagation distance




FIGURE 5. Schematic of wave groups and forced acoustic and seismic wave motion. For a given wave group period T , the horizontal wavelength $L = 2\pi/K$ can be larger than the acoustic wavelength in the water due to the oblique incidence of the sound waves. The superposition of two obliquely propagating sound waves (arrows) forms a mode pattern that propagates on the horizontal at a supersonic speed. The acoustic wavelength is $L_a = 2\pi/k_a = 2\pi/\sqrt{K^2 + l^2} = \alpha_w/T$. Both vertical and horizontal wavelengths are larger. For readability, the wave and bottom amplitude is not to scale, and also we have reduced the number of waves in the group from 104 to 10. Other than that, the angles are respected. The configuration shown here corresponds to the conditions for maximum amplification of mode 1 (see below), with a vertical wavelength to water depth ratio of 0.75. For a water depth $h = 4400$ m depth, this corresponds to $L = 7.7$ km and a seismic frequency $f_s = 0.29$ Hz.

(Hasselmann 1963). For wavenumbers that allow a vertical propagation in the crust, we obtain body waves. These waves can be compression or shear (P or S) waves, and these two types occur in overlapping ranges of K .

3.2. Acoustic noise in an unbounded ocean

In order to illustrate the different types of solutions, it is interesting to express the solution for an infinitely deep ocean, in which sound waves are radiated from the surface only. The velocity field and associated pressure fluctuations are

$$\phi_2 = \frac{1}{\rho_w} \int \frac{i\omega_s \hat{p}_2(\mathbf{K}, f_s)}{-\omega_s^2 + igl} e^{i[-lz + \Theta(\mathbf{k}, \mathbf{k}', s, s')]} d\mathbf{K} df_s \quad (3.5)$$

$$p_2 = \int \frac{\hat{p}_2(\mathbf{K}, f_s)}{1 - igl/\omega_s^2} e^{i[-lz + \Theta(\mathbf{k}, \mathbf{k}', s, s')]} d\mathbf{K} df_s \quad (3.6)$$

where p_2 has been obtained using the linearized version of eq. (2.4). The measurable pressure signal is the sum of the linear pressure p_1 , p_2 given by eq. (3.6), and the Bernoulli correction $p_{2,B}$ given by

$$p_{2,B}(z) = \rho_w \sum_{\mathbf{k}, s, \mathbf{k}', s'} D_{pb}(\mathbf{k}, s, \mathbf{k}', s', z) Z_{1,\mathbf{k}}^s Z_{1,\mathbf{k}'}^{s'} e^{i\Theta(\mathbf{k}, \mathbf{k}', s, s')}. \quad (3.7)$$

We shall neglect $g|l|/\omega_s^2$, which is bounded by the ratio between the deep water gravity and sound speeds, which is less than 0.1 for wave periods less than 180 s. We can also express the velocity potential as a sum of propagating (acoustic, l real) and evanescent (acoustic-gravity, l imaginary) modes,

$$\phi_2 = \phi_{2,p} + \phi_{2,e}. \quad (3.8)$$

We get the frequency spectrum of the propagating modes by summing on only the inner regions of the wavenumber space (labelled P+S, S and R in figure 1),

$$F_{p2,p}(f_s) = \int_{K < \omega_s/\alpha_w} F_{p2,\text{surf}}(\mathbf{K}, f_s) d\mathbf{K}. \quad (3.9)$$

For this range of wavenumbers $|k - k'| < K < \omega_s/\alpha_w$, so that $2f|f - f'| < g/(4\pi^2)|k - k'|$ gives $|f - f'| < g/(2\pi\alpha_w)$ which is close to = 0.001 Hz. Typical ocean wave spectra have a relative frequency half-width σ_f/f that is between 0.03 for swells and 0.07 for wind-seas (Hasselmann *et al.* 1973), so that $E(f) \simeq E(f')$ is a good approximation for typical wave frequencies, larger than 0.05 Hz.

The wave spectrum is thus broad enough for us to use $\mathbf{K} = 0$, so that the spectral density $F_{p2,\text{surf}}$, given by eq. (2.26), can be taken out of the integral in eq. (3.9). The acoustic spectrum simplifies as

$$F_{p2,p}(f_s) = \frac{\pi\omega_s^2}{\alpha_w^2} \rho_w^2 g^2 f_s E^2(f) I(f). \quad (3.10)$$

This is identical to the expression given by Lloyd (1981).

3.3. Gravity noise in an infinitely deep ocean

The pressure associated with acoustic-gravity modes is the other part of the integral in (3.9), for $K > \omega_s/\alpha_w$. The imaginary wave number l gives a vertical attenuation of the power spectrum by a factor $e^{-2|l|z}$. With that attenuation we may, for large enough depths, consider that only modes with $K \ll k$ contribute to the result, so that we may take $F_{p2,\text{surf}}(\mathbf{K}, f_s) \simeq F_{p2,\text{surf}}(\mathbf{K} = 0, f_s)$, and take it out of the integrand. This is only valid up to a maximum wavenumber $K_{\text{max}} = \epsilon k$. For numerical applications we have used $\epsilon = 0.2$.

With this approximation we have,

$$\begin{aligned} F_{p2,e}(f_s, z) &= F_{p2,\text{surf}}(\mathbf{K} = 0, f_s) 2\pi \int_{\omega_s/\alpha_w}^{K_{\text{max}}} K e^{2|l|z} dK \\ &= F_{p2,\text{surf}}(\mathbf{K} = 0, f_s) 2\pi \int_0^{K_{\text{max}}} |l| e^{2|l|z} d|l| \\ &= \frac{\pi}{2z^2} \rho_w^2 g^2 f_s [1 - e^{2zK_{\text{max}}}] E^2(f) I(f) \end{aligned} \quad (3.11)$$

A previous investigation by Cox & Jacobs (1989) included an extra factor $(1 + zK_{\text{max}})$ in front of the exponential term $e^{2zK_{\text{max}}}$, because they neglected compressibility effects. That term, however, becomes negligible very early as the depth increases. Hence, their observations at depth 100 to 290 m, in more than 4000 m of water, are not affected by this small compressibility correction.

As shown by figure 6, the oceanic pressure signals can be dominated by linear gravity waves down to depths of a few hundred meters. When looking at the double frequency band, linear waves may only dominate in the top 100 m. At these frequencies, the most important contribution between about 100 to 500 m, provided that $E^2(f)I(f)$ is large enough. Propagating modes should dominate only beyond about 1000 m in the case of an infinitely deep ocean, or only 300 m, when accounting for their amplification by the bottom elasticity, assuming a typical tenfold amplification[†]. All these numbers will be

[†] This amplification depends not only on the impedance ratio of the water and crust, which defines the amplification coefficients c_j derived below, but also on the seismic attenuation coefficient Q , which is discussed in section 4. Realistic calculations following Ardhuin *et al.* (2011)




FIGURE 6. Example of (a) directional wave spectrum and (b) resulting profiles of the different contributions to the pressure fluctuations in the ocean, assuming infinite water depth. The ratio of double frequency to linear wave contributions depends on the amplitude of the waves and on the directional spectral shape, because all double frequency contributions are proportional to $E^2(f)I(f)$. Here we have used a numerically estimated directional spectrum that corresponds to the source of a loud noise event recorded at the ocean bottom seismometer H2O, on 31 May 2002, with the encounter of swell from a North Pacific storm, with swell from Hurricane Alma (This event is analyzed in detail by Obrebski *et al.* 2012).

reduced in the case of surface gravity waves with shorter periods than the 15 s swell taken for the example in figure 6.

3.4. Atmospheric noise source: microbaroms

The source of noise in the atmosphere can also be derived with the same formalism, as an alternative to the Green functions used by Waxler & Gilbert (2006). Indeed, we may consider the atmospheric motion to be irrotational, so that the equations of motion are identical in the atmosphere and in an infinitely deep ocean, with the only difference that the atmospheric density is ρ_a and the atmospheric sound speed is α_a . The second-order velocity potential takes the form,

$$\phi_{2,a} \propto \exp[i(K_x x + K_y y + l_a z - \omega_s t)] \quad \text{for } z > 0, \quad (3.12)$$

with

$$l_a = \sqrt{\frac{\omega_s^2}{\alpha_a^2} - K^2}. \quad (3.13)$$

Because $\rho_w / \rho_a \simeq 1000$, the air motion has only a small $O(\rho_w / \rho_a)$ local influence on the water motion, so that the solutions derived above for the water motion remain valid in the presence of air. The air motion, with a velocity potential ϕ_a also obeying eq. (2.13) is fully determined from the water motion via the kinematic boundary conditions on the air and water-sides of the interface (2.7),

$$\frac{\partial \phi_a}{\partial z} - \frac{\partial \phi}{\partial z} \simeq \nabla(\phi_a - \phi) \cdot \nabla \zeta - \zeta \frac{\partial^2(\phi_a - \phi)}{\partial^2 z} \quad \text{at } z = 0. \quad (3.14)$$

typically give a factor 10 to 20 amplification of the sound in the water column due to the bottom elasticity.

From the first order potential in the air (e.g. Waxler & Gilbert 2006)

$$\phi_1 = \sum_{\mathbf{k}, s} i s \frac{g}{\sigma} Z_{1, \mathbf{k}}^s e^{-kz} e^{i(\mathbf{k} \cdot \mathbf{x} - s\sigma)} \quad (3.15)$$

we obtain the second order potential,

$$\frac{\partial \phi_{2,a}}{\partial z} = \frac{\partial \phi_2}{\partial z} + \sum_{\mathbf{k}, s, \mathbf{k}', s'} D_{za}(\mathbf{k}, s, \mathbf{k}', s', z) Z_{1, \mathbf{k}}^s Z_{1, \mathbf{k}'}^{s'} e^{i\Theta(\mathbf{k}, \mathbf{k}', s, s')} \quad (3.16)$$

and a new coupling coefficient

$$D_{za}(\mathbf{k}, s, \mathbf{k}', s', z) = -\frac{2isg}{\sigma} (kk' + \mathbf{k} \cdot \mathbf{k}'). \quad (3.17)$$

We note that for $\mathbf{k}' = -\mathbf{k}$, $D_{za} = 0$, so that the long-wavelength motion with $K \ll k$ simplifies to

$$\frac{\partial \phi_{2a}}{\partial z} \simeq \frac{\partial \phi_2}{\partial z} \quad \text{at } z = 0. \quad (3.18)$$

consistent with the result given by Posmentier (1967) for the interaction of monochromatic wave trains, and in disagreement with a factor 8 correction proposed by Arendt & Fritts (2000).

This gives a pressure spectrum for the propagating atmospheric waves,

$$F_{p2,ap}(f_s) = \int_{K < K_{\max}} \frac{\rho_a^2 l^2}{\rho_w^2 l_a^2} F_{p2,\text{surf}}(\mathbf{K}, f_s) d\mathbf{K} = R(K_{\max}) \frac{\pi \omega_s^2}{\alpha_w^2} \rho_a^2 g^2 f_s E^2(f) I(f). \quad (3.19)$$

In order to avoid the singularity for $l_a = 0$, and atmospheric ducting effects not represented here, we take $K_{\max} = \omega_s / (2\alpha_a)$ which restricts the acoustic propagation directions to be within 30 degrees from the vertical. In that case we have $R(K_{\max}) = 0.5$ instead of $R(K_{\max}) = 0.25$ with the vertical propagation approximation of Waxler & Gilbert (2006), which replaces the l^2/l_a^2 factor in the integral by its value α_a^2/α_w^2 for $K = 0$. Other than that, our expression is consistent with their low Mach number asymptote, i.e. $\sigma/k \ll \alpha_a$ (Waxler & Gilbert 2006, eq. 61). The present theory also allows the estimation of the evanescent wave components given by wavenumbers $K > \omega_s/\alpha_a$.

4. Noise in a finite depth ocean

For large depths compared to the ocean surface gravity wave wavelength, $kh \gg 1$, the finite depth has little effect on the evanescent modes except for a doubling of the motion amplitude near the bottom, as the vertical profiles of the form $\exp(Kz)$ are replaced by $\cosh(Kz + Kh)/\cosh(Kh)$. This is similar to the finite depth effect on linear wave motions. However, the propagating modes radiated by the surface will now undergo multiple reflections at the bottom and sea surface, as shown in figure 5. The oceanic acoustic field is tightly coupled to elastic waves in the crust through these reflections.

One of the greatest complications induced by the presence of a bottom is the heterogeneity of the sediment and rock layers below the water column. The natural layering of the crust has a strong influence on the sound reflection and the nature of the seismic modes (e.g. Latham & Sutton 1966; Abramovici 1968). These effects will not be considered here, and we follow exactly the theoretical setting of Hasselmann (1963).

4.1. Elastic wave theory

For simplicity, we assume here that the ocean bottom is a uniform and semi-infinite solid, with constant density ρ_s and compression and shear wave velocities α_c and β given from the Lamé coefficients λ and μ ,

$$\alpha_c^2 = \frac{\lambda + 2\mu}{\rho_s}, \quad \beta^2 = \frac{\mu}{\rho_s}. \quad (4.1a,b)$$

Assuming that the crust is a Poisson solid, $\mu = \lambda$ and $\alpha_c = \sqrt{3}\beta$.

The free wave problem in these conditions was solved by Stoneley (1926). Here we consider the forced problem treated by Hasselmann (1963) with a forcing by a pressure field $\widehat{p}_{2,\text{surf}}$ at the sea surface, but now generalized to an additional wave-induced bottom pressure $\widehat{p}_{2,\text{bot}}$.

The equations of motion in the water column are unchanged from the previous section, but they are now coupled to the elastic motions of the crust. Crustal motions can be separated in an irrotational part with a velocity potential ϕ_c and a rotational part with a stream function ψ , both solution to Laplace's equation. With a wave source at the surface, a horizontal propagating wave of phase $\Theta = Kx - \omega_s t$, implies that ϕ and ψ are either decaying or propagating downwards. They must therefore take the following form,

$$\phi_c = Ae^{m(z+h)} e^{i\Theta}, \quad \psi = Be^{n(z+h)} e^{i\Theta}. \quad (4.2a,b)$$

Both vertical wavenumbers m and n are given by the Fourier transform of eq. (2.13),

$$m = \sqrt{K^2 - \frac{\omega_s^2}{\alpha_c^2}}, \quad n = \sqrt{K^2 - \frac{\omega_s^2}{\beta^2}} \quad (4.3a,b)$$

where the sound speed in water has been replaced by the compression and shear velocities. For $K > \omega_s/\beta_s$, m and n are real and both compression and shear waves are evanescent. For $\omega_s/\alpha_c < K < \beta_s$ the compression wave is evanescent but there is a shear (S) wave that propagates through the crust.

The constants A and B have dimensions of m^2/s and are determined by the boundary conditions at the ocean bottom.

Horizontal and vertical ground displacements are given by the real parts of

$$\xi_x = \left(KAe^{m(z+h)} + i n B e^{n(z+h)} \right) e^{i\Theta} / \omega_s \quad (4.4)$$

$$\xi_z = \left(-i m A e^{m(z+h)} + K B e^{n(z+h)} \right) e^{i\Theta} / \omega_s. \quad (4.5)$$

Hooke's law of elasticity gives us

$$\tau_{zz} = \lambda \left(\frac{\partial \xi_x}{\partial x} + \frac{\partial \xi_z}{\partial z} \right) + 2\mu \frac{\partial \xi_z}{\partial z}, \quad \tau_{xz} = \mu \left(\frac{\partial \xi_x}{\partial z} + \frac{\partial \xi_z}{\partial x} \right) \quad (4.6a,b)$$

The zero tangential stress on the ocean bottom $\tau_{xz}(z = -h) = 0$ yields the following relationship between A and B , which is typical of Rayleigh waves,

$$B = \frac{2iKm}{n^2 + K^2} A. \quad (4.7)$$

As a result, on top of the two unknowns that are the bottom amplitudes C and D of the upward and downward velocity potential in the water (they appear in eq. 3.4), we have only one other unknown, the compression wave amplitude A .

The three equations that relate C , D and A are: the combined kinematic and dynamic

boundary condition (2.15), and the bottom continuity of normal velocity

$$\frac{\partial \phi_2}{\partial z} = \frac{\partial \phi_c}{\partial z} + \frac{\partial \psi}{\partial x} \quad \text{at } z = -h \quad (4.8)$$

and normal stress,

$$-\tau_{zz}(-h) = p(-h) = \rho_w \frac{\partial \phi_2(-h)}{\partial t} + \widehat{p}_{2,\text{bot}}. \quad (4.9)$$

For waves in intermediate or shallow water, i.e. $kh < \pi$, the Bernoulli term $\widehat{p}_{2,\text{bot}}$ that was not considered by Hasselmann (1963) should be included. We thus obtain the linear system of equations

$$(-\omega_s^2 \rho_w + gil) e^{ilh} C + (-\omega_s^2 \rho_w - gil) e^{-ilh} D = \frac{i\omega_s}{\rho_w} \widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s) \quad (4.10)$$

$$qA - ilC \quad + ilD \quad = 0 \quad (4.11)$$

$$rA - i\omega_s \rho_w C \quad - i\omega_s \rho_w D \quad = -\widehat{p}_{2,\text{bot}}(\mathbf{K}, f_s) \quad (4.12)$$

with

$$q = \frac{m\omega_s^2}{\omega_s^2 - 2K^2\beta^2} \quad (4.13)$$

$$\begin{aligned} r &= \frac{i}{\omega_s} \rho_s \left[-\frac{4\beta^4 K^2 mn}{\omega_s^2 - 2K^2\beta^2} + (\omega_s^2 - 2K^2\beta^2) \right] \\ &= \frac{i}{\omega_s} \left[-\rho_s m^2 \alpha_c^2 + \lambda k^2 + 4\mu \frac{K^2 mn}{n^2 + K^2} \right] \end{aligned} \quad (4.14)$$

Since we are in the range where $l < \omega_s/\alpha_w$ we may neglect again $g|l|/\omega_s^2$, which is less than 0.1 for OSGW periods less than 180 s. We then rewrite these equations in matrix form, it writes,

$$M[A, \quad C, \quad D]^T = [-\widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s), \quad 0, \quad -\widehat{p}_{2,\text{bot}}(\mathbf{K}, f_s)]^T \quad (4.15)$$

with

$$M = \begin{bmatrix} 0 & i\rho_w \omega_s e^{ilh} & i\rho_w \omega_s e^{-ilh} \\ q & -il & il \\ r & -i\omega_s \rho_w & -i\omega_s \rho_w \end{bmatrix} \quad (4.16)$$

The general solution of eq. (4.15) is the sum of one particular forced solution and the general solution of the homogeneous system, without the right hand side forcing, i.e. the free waves. This combination of free and forced waves is further determined by the initial conditions.

The forced solution is readily expressed using the determinant of the system

$$\det(M) = -\rho_w \omega_s (l e^{ilh} r + \omega_s \rho_w e^{-ilh} q - l e^{-ilh} r - \omega_s \rho_w e^{ilh} q) \quad (4.17)$$

$$= \frac{2\rho_w}{(\omega_s^2 - 2K^2\beta^2)} \left\{ l \rho_s \cos(lh) \left[4\beta^4 K^2 mn - (\omega_s^2 - 2K^2\beta^2)^2 \right] - \rho_w m \sin(lh) \omega_s^4 \right\}, \quad (4.18)$$

in the form

$$A = 2l\rho_w \omega_s \frac{\widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s) + \cos(lh)\widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s)}{\det(M)} \quad (4.19a)$$

$$C = \frac{-i(lr + q\omega_s \rho_w)\widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s) + qe^{-ilh}\widehat{p}_{2,\text{bot}}(\mathbf{K}, f_s)}{\det(M)} \quad (4.19b)$$

double-frequency noise generation by surface gravity waves waves 17

$$D = i(q\omega_s\rho_w - lr) \frac{(q\omega_s\rho_w - lr)\widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s) - qe^{-ilh}\widehat{p}_{2,\text{bot}}(\mathbf{K}, f_s)}{\det(M)} \quad (4.19c)$$

For $kh \gg 1$ we may neglect $\widehat{p}_{2,\text{bot}}$ and,

$$C = \frac{\widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s)}{\rho_w\omega_s} \frac{ir' - q'}{2r' \cos(lh) - 2q' \sin(lh)} \quad (4.20a)$$

$$D = \frac{\widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s)}{\rho_w\omega_s} \frac{ir' + q'}{2r' \cos(lh) - 2q' \sin(lh)} \quad (4.20b)$$

with

$$q' = \rho_w\omega_s q, \quad r' = ilr. \quad (4.21a,b)$$

4.2. Acoustic noise in a bounded ocean

Taking typical values of the water and crust density and sound speeds gives $r'/q' > 70$ for the free modes that are significantly generated by the waves (i.e. $c_j > 0.1$, as defined below by eq. 4.34). We can thus consider that $q'/r' \ll 1$, which gives $C \simeq D \simeq \widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s)/[2\rho_w\omega_s(\cos(lh) + q'/r')]$. The velocity potential and pressure in the water are given by,

$$\phi_2 \simeq i\widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s) \frac{\cos[l(z+h) - q'/r']}{\rho_w\omega_s(\cos(lh) - q'/r')} e^{i\Theta}, \quad p_2 \simeq i\widehat{p}_{2,\text{surf}}(\mathbf{K}, f_s) \frac{\cos[l(z+h)] - q'/r'}{(\cos(lh) - q'/r')} e^{i\Theta}. \quad (4.22a,b)$$

The small but finite factor q'/r' ensures that the solution remains finite as a small fraction of the acoustic energy is radiated into the crust, otherwise the acoustic energy would accumulate in the water column. The pressure oscillation are thus maximum for resonant frequencies such that the ratio of the water depth and vertical wavelength $lh/(2\pi)$ is $1/4, 3/4, 5/4 \dots$. We note that the vertical wavelength $2\pi/l$ is always greater than the acoustic wavelength $2\pi/\sqrt{K^2 + l^2} = f_s\alpha$. As a result, the resonant frequencies are shifted to higher values compared to the vertical resonant condition that is given by $f_s h \alpha_w = 1/4 \dots$

We may now integrate the pressure spectrum for all acoustic wavenumbers to find, again, the frequency spectrum,

$$F_{p2,ap}(f_s) = 2\pi\rho_w^2 g^2 f_s E^2(f) I(f) \int_0^{\omega_s/\alpha_w} \left[\frac{\cos(lz + lh) - q'/r'}{\cos(lh) - q'/r'} \right]^2 K dK. \quad (4.23)$$

We now have a depth dependence of the sound spectrum. At the surface it is equal to the widely used unbounded ocean value (Lloyd 1981), but in the water column it can be strongly amplified at depths where $\cos(lh)$ approaches 0, which, as we shall see, includes the near-bottom region in the cases where the noise generation is strongest.

4.3. Rayleigh waves

In order to simplify the algebra, we consider in this section waves in deep water and neglect bottom forcing. As result, this section brings no new results compared to Hasselmann (1963), it only gives more explicitly some intermediate results.

For a fixed frequency ω_s , there is at least one wavenumber K for which $\det(M) = 0$. This condition defines the dispersion relation of the Rayleigh modes (Stoney 1926),

$$\tan(lh) = \frac{l\rho_s}{m\rho_w} \times \frac{4\beta^4 K^2 mn - (\omega_s^2 - 2K^2\beta^2)^2}{\omega_s^4} \quad (4.24)$$




FIGURE 7. Wavenumbers of seismic and acoustic modes for a fixed frequency, which are the solutions of eq (4.24), for which the right hand side (RHS) equals the real part of $\tan(lh)$. Here we used $h = 5000$ m, $\alpha_w = 1500$ m/s, $\beta = 3200$ m/s, $\rho_w = 1000$ kg/m³ and $\rho_s = 2500$ kg/m³. This graph corresponds to a seismic frequency $f_s = 0.263$ Hz, for which $f_s h / \alpha_w = 0.88$. Here the determinant has three real roots, one acoustic mode K_a for which $l = 0$ (horizontal propagation) and two seismic modes K_0 et K_1 . The number of these roots increases with the frequency. One new solution appears every time $2\pi f_s$ becomes larger than $\omega_{s,cj}$ defined by eq. (4.29). For reference we also give the wavenumber for which the right hand side of eq. (4.24) is zero, which is the Rayleigh wavenumber K_R in the absence of the ocean layer ($h = 0$).

with the fundamental mode corresponding to the largest value K . Figure 7 illustrates the presence of different values for K .

The phase speeds of the Rayleigh modes vary continuously from the shear wave velocity β , in the limit $n = 0$ where the shear waves transition from propagating to evanescent, to the sound speed in water α_w , in the limit $l = 0$ where the acoustic modes become evanescent in the water (figure 8.a). This variation of the phase speed has an inflexion point close to the phase speed of Rayleigh waves without the water layer, corresponding to a maximum in group speed (figure 8.b).

We may now use eq. (4.19) to obtain the ground displacement amplitude

$$\delta \equiv \xi_z(-h) = G(K_x, K_y, \omega_s) \widehat{p}_{2,\text{surf}}(K_x, K_y, \omega_s) \quad (4.25)$$

as a function of the amplitude of the sea surface pressure, with the transfer function

$$G = \frac{2i\rho_w l m(K^2 - n^2)}{(n^2 + K^2) \det(M)}. \quad (4.26)$$

An example of this transfer function is shown in figure 9, in which the dispersion relation of the Rayleigh modes appear as the narrow red bands around the singularities of G .

To evaluate the complete family of Rayleigh wave solutions with dispersion relations $\omega_{s,j}(K)$, where j is the mode number, we need to examine the nature of these singularities.




FIGURE 8. Phase (top) and group (bottom) speeds of Rayleigh waves as a function of the dimensionless water depth. Calculations used $\alpha_w = 1470 \text{ m s}^{-1}$, $\beta = 3000 \text{ m s}^{-1}$, $\rho_w = 1000$ and $\rho_s = 2500 \text{ kg m}^{-3}$. Values without the water layer are also indicated with the blue line. In that case Rayleigh waves are not dispersive.

We re-write the determinant of the system as,

$$\det(M) = \frac{2il\rho_s\rho_w \cos(lh)}{(\omega_s^2 - 2K^2\beta^2)} \{ (4\beta^2 k^2) (mn + \omega_s^2 - \beta^2 K^2) - \omega_s^4 (1 + \tan(lh)\rho_w m / \rho_s l) \}. \quad (4.27)$$

For most frequencies, the singularities are simple, allowing a Taylor expansion of G in this form

$$G(K, \omega_s) \simeq \frac{G'(K_j(\omega_s))}{\omega_s^2 - \omega_{s,j}^2(K)}. \quad (4.28)$$

However, for each mode j , there is a critical frequency

$$\omega_{s,cj} = \frac{1}{h} \left[\arctan\left(-\frac{l}{m}\rho_s\rho_w\right) + j\pi \right] \quad (4.29)$$




FIGURE 9. Magnitude of the transfer function $G(K, f_s)$ illustrating the singularities along the dispersion relations of the free Rayleigh modes with wavenumbers $K = K_n(f_s)$. The oblique dashed lines correspond to phase speeds equal to α_c , β , and α_w , and separate the four domains body waves (P+S), mixed body and evanescent waves (S), Rayleigh waves (R) and acoustic-gravity modes (A-G).

for which both $\tan(lh)\rho_w m/\rho_s l = 1$, and $\omega_s^2 - K^2\beta^2 = 0$. These critical frequencies $\omega_{s,cj}$ exist for modes $j > 0$. These frequencies are those for which new Rayleigh modes appear, in a way similar to the Love waves discussed by Aki & Richards (2002, figure 7.3). The singularities at $(K, \omega_s) = (\omega_{s,cj}/\beta, \omega_{s,cj})$ are not simple and for these we have $G(\omega_s, k) \simeq G'(K_j(\omega_s))/\sqrt{K^2 - K_j^2(\omega_s)}$.

Solutions for the vertical displacement at the top of the crust have a spectral amplitude which is linearly related to the equivalent surface pressure amplitude,

$$\delta(K_x, K_y, \omega_s) = G(K_x, K_y, \omega_s)\hat{p}_{2,\text{surf}}(K_x, K_y, \omega_s). \quad (4.30)$$

For the simple singularities of G , i.e., away from the head waves, we may write $G(K, \omega_s) = G'(K, \omega_{s,j})/(\omega_s^2 - \omega_{s,j}^2) + O(\omega_s - \omega_{s,j})$. Taking initial conditions $\delta(t=0) = 0$ and $\partial\delta/\partial t = 0$ gives the full solution (Hasselmann 1962, eq. 3.2) of the j^{th} Rayleigh mode response

$$\delta(K_x, K_y, \omega_s) = G'(K, \omega_{s,j})e^{iKx} \left[\frac{1}{\omega_{s,j}^2 - \omega_s^2} e^{-i\omega_s t} - \frac{1}{2\omega_{s,j}} \left(\frac{e^{i\omega_{s,j}t}}{\omega_{s,j} - \omega_s} + \frac{e^{i\omega_{s,j}t}}{\omega_{s,j} + \omega_s} \right) \right], \quad (4.31)$$

as a function of K , where $\omega_{s,j}(K)$ is frequency of the j^{th} Rayleigh mode. For a forcing that varies slowly on the scale of the seismic period $2\pi/\omega_s$, and provided that the forcing spectrum is continuous in spectral space, this gives the rate of change of the ground displacement spectrum given by Hasselmann (1963),

$$\frac{\partial F_\delta(K_x, K_y)}{\partial t} = S_{DF}(K_x, K_y) = \frac{\pi |G'|^2}{2\omega_{s,j}^2} F_{p2,\text{surf}}(K_x, K_y, \omega_s), \quad (4.32)$$

The other discrete singularities, at $(K, \omega_s) = (\omega_{s,cj}/\beta, \omega_{s,cj})$, are associated with 'conical' or 'head' waves, for which the vertical wavenumber is $n = 0$, and that propagate along the ocean-crust interface. These horizontally propagating shear waves are generated with an evanescent compression wave. The singularity is integrable over the two spectral dimensions ω_s and K .

Although it looks like only the resonant forcing contributes to the solution, it is in fact the near-resonant forcing ($\omega_s \simeq \omega_{s,j}$) that builds up the seismic noise, because the exact resonant terms have a zero measure in spectral space. Indeed, a purely resonant forcing would give an amplitude that increases linearly with time, and an energy that increases like t^2 . The linear growth of energy in time can be interpreted as an effect of the narrowing of the frequency bandwidth in which the interaction is significant. This is a general property of wave-wave interactions (see also Hasselmann 1966).

We express the source of seismic noise with the rate of increase of the variance of δ per unit of propagation distance. It is

$$S_{DF}(\omega_s) = \frac{K(s)S_{DF}(K_x, K_y)}{V^2} = \frac{4\pi^2 f_s c_j^2}{\beta^5 \rho_s^2} F_{p2,\text{surf}}(K_x, K_y, \omega_s) \quad (4.33)$$

where V is the group speed of the seismic waves, and c_j is a dimensionless coefficient that depends on $\omega_s h / \alpha_w$ and the seismic mode index j , shown in figure 10,

$$c_j^2 = \frac{\beta^5 \rho_s^2 K_j \pi |G'_j|^2}{V_j^2 2\pi \omega_s 2\omega_s^2}. \quad (4.34)$$

We note that a missing 2π in eq. (5) of Arduin *et al.* (2011) has been corrected here.

A very rough simplification can be obtained by taking V and K independent of j and h . Then $S_{DF}(f_s)$ can be taken as a sum of all Rayleigh modes in the form

$$S_{DF}(f_s) = \frac{4\pi^2 f_s \tilde{C}}{\beta^5 \rho_s^2} F_p(\mathbf{K} \simeq 0, f_s). \quad (4.35)$$

where \tilde{C} is a dimensionless coefficient that is the sum of contributions c_j^2 from all Rayleigh modes,

$$\tilde{C} = \sum_{i=0}^{\infty} c_j^2. \quad (4.36)$$

Values of c_j are obtained from eq. (4.35), and shown in figure (10).

We may propagate these sources of seismic waves in a vertically symmetric earth model, neglecting all three-dimensional propagation effects, and parameterizing seismic wave scattering and dissipation with a uniform quality factor Q . Under these assumptions, the spectral density of the vertical ground displacement at $z = -h$ is

$$F_\delta(\lambda, \phi, f_s) = \int_{-\pi/2}^{\pi/2} \int_0^{2\pi} \frac{S_{DF}(f_s)}{R_E \sin \Delta} e^{-2\pi f_s \Delta R_E / (UQ)} (R_E^2 \sin \phi' d\lambda' d\phi') \quad (4.37)$$

with R_E the Earth radius, and U the seismic group velocity. The term $(R_E^2 \sin \phi' d\lambda' d\phi')$ is the Earth surface area element. The denominator $(R_E \sin \Delta)$ is the geometrical spreading factor for wave energy that follows geodesics on the sphere (e.g. Kanamori & Given 1981), replacing the distance $(R_E \Delta)$ used in flat Earth models (e.g. Hasselmann 1963).

The Rayleigh waves thus generated propagate away like free modes, those that exist without the local forcing. For these free modes with a monochromatic ground displacement $\delta(x, y, t)$ the surface pressure is constant so that $C = -D$ and the velocity potential




FIGURE 10. Dimensionless coefficients c_j that amplify the wave-induced pressure into ground displacement. The maxima of c_j correspond to quarter-wavelength resonance typical of organ pipes, except that here the sound waves propagate obliquely in the water column, which is why the maxima are at values of $f_s h / \alpha_w$ which are not exactly at $1/4, 3/4, \dots$, but shifted to higher frequencies by a factor $l / \sqrt{K^2 + l^2}$. The amplitude of the peaks depends on the impedance ratio of the sea water and crust. Hence the peak amplitude increases with $\rho_s \beta / (\rho_w \alpha_w)$. For example, $\beta = 2800 \text{ m s}^{-1}$, gives a maximum of 0.88 for c_0 instead of 1.03 here with $\beta = 3000 \text{ m s}^{-1}$.

and pressure in the water take the form,

$$\phi_2(z) = \frac{\omega_s}{l \cos(lh)} \sin(lz) \delta(x, y, t) \quad (4.38a,b)$$

$$p_2(z) = i \rho_w \frac{\omega_s^2}{l \cos(lh)} \sin(lz) \delta(x, y, t). \quad (4.38a,b)$$

For varying water depths, and on land, one may assume that the seismic energy is propagated along a ray and apply the refraction coefficient given by Hasselmann (1963).

4.4. Seismic P and S waves

Unlike the Rayleigh waves that grow resonantly in time owing to the trapping in the ocean/crust waveguide, the P and S waves, for which the vertical wavenumbers m or n are complex, radiate into the earth's interior, and their energy level is given directly by its value at the source. In particular, for $K < \omega_s / \alpha_c$ we have propagating P waves with a velocity potential amplitude A given by eq. (4.19). The frequency spectrum of vertical ground displacements at ($z = -h$) can be evaluated directly with (4.30) because G has no singularity in this range of wavenumbers,

$$F_{\delta,P}(f_s) = f_s E^2(f) I(f) \frac{\rho_w g^2}{\rho_s^2 \beta_s^4} c_P^2 \quad (4.39)$$

with a non-dimensional coefficient c_P ,

$$c_P^2 = 2\pi \int_0^{\omega_s / \alpha_c} \frac{4l^2 m^2 \rho_s^2 \beta_s^4}{\omega_s^2 \det^2(M)} K dK. \quad (4.40)$$




FIGURE 11. Non-dimensional coefficients c_P and c_S that amplify the wave-induced pressure into ground displacement associated with P and S waves.

A similar expression can be written for S waves, and both are illustrated in figure 11,

$$F_{\delta,S}(f_s) = f_s E^2(f) I(f) \frac{\rho_w^2 g^2}{\rho_s^2 \beta_s^4} c_S^2. \quad (4.41)$$

However, in the range of wavenumbers where S waves exist, $k < \omega_s/\beta$, there can also be evanescent P waves, and the system can approach the singularity for $\omega_s = \omega_{s,cj}$ and $k = \omega_s/\beta$. We thus evaluated numerically the coefficient

$$c_S^2 = 2\pi \int_0^{\omega_s/\beta} \frac{4l^2 m^2 k^2 \rho_s^2 \beta_s^4}{\omega_s^2 (n^2 + k^2) \det^2(M)} K dK. \quad (4.42)$$

Due to the typically three times stronger attenuation of S waves compared to P waves in the Earth mantle (e.g. Anderson & Hart 1978; Pasyanos *et al.* 2009), we will now focus on P waves only, which should dominate in the far field of the noise source.

For the estimation of the spectrum recorded outside of a source area, it is more convenient to express the local seismic source as a function of the horizontal propagation angle θ , and the vertical take-off angle φ . For P waves, this gives,

$$F_{\delta,P}(f_s, \theta, \varphi) = f_s E^2(f) I(f) \frac{1}{\rho_s^2 \beta_s^4} c_{P,\varphi}^2 \sin \varphi \quad (4.43)$$

with the non-dimensional coefficient $c_{P,\varphi}$ defined by

$$c_{P,\varphi}^2 = \frac{4l^2 m^2 \rho_s^2 \beta_s^4}{\omega_s \alpha_c \det^2(M)} \frac{\partial K}{\partial \varphi}, \quad (4.44)$$

which is the normalized source per unit solid angle Ω , so that the average over the half space of down-going directions Ω^- is

$$c_P^2 = \int_0^{2\pi} \int_0^{\pi/2} c_{P,\varphi}^2 \sin \varphi d\varphi = \int_{\Omega^-} c_{P,\varphi}^2 d\Omega \quad (4.45)$$

as defined by eq. (4.40).




FIGURE 12. Dimensionless coefficients $c_{P,\varphi}$ that amplify the wave-induced pressure into ground displacement. The maximum for a zero take-off angle corresponds to vertically propagating compression waves, and the compression waves that propagate along the crust have a zero amplitude.

It is noteworthy that the distribution of the P -wave energy with the take-off angle is very close to the one given by a small disk pushing at the top of a uniform half space, as given by Miller & Pursey (1955) and used by Vinnik (1973), although it also varies with the non-dimensional water depth $f_s h / \alpha_w$. The only missing item in the work by Vinnik (1973) is the very strong amplification of the motion for resonant frequencies associated with the water layer. Due to the large impedance contrast at the sea-crust interface the relative amplification of P waves is one order of magnitude stronger than for Rayleigh waves, and we thus expect a much stronger locking of the strong seismic noise sources on the isobaths corresponding to a maximum amplification.

4.5. Observable P wave spectra

We will now finish our review by estimating the ground motion due to P and Rayleigh waves as a function of the distance from the source. For a seismic station or hydrophone in the ocean, the incoming energy at the receiver will be strongly reflected at $z = -h$ but again amplified by multiple bottom and surface reflections in the water column. However, this P -wave signal from remote sources is likely to be dwarfed by locally generated noise.

For a land-based station, we may consider that incoming P waves arriving directly from the the source area. There are many other seismic wave phases that have undergone multiple reflections at the surface, these are called PP , PPP ..., or at any inner boundary of the Earth, like PkP phases that have reflected off the mantle-core interface (Aki & Richards 2002). These can be treated exactly like the direct P phase.

On arrival at the receiver, these waves are totally reflected at the Earth surface at $z = 0$, which doubles the ground motion, so that the ground displacement is given by the integral of four times the incoming spectral densities over the directions θ and φ , which can be replaced by a an integral over the source positions (λ, ϕ) . The transformation from the ray parameters (θ, φ) to the geographical coordinate can be obtained approximately for any type of seismic wave using travel time tables (e.g. Snoke 2009), which also provide the travel time τ .




FIGURE 13. Schematic of rays for seismic P waves radiated from a point source (red dashed lines) with a directional distribution $c_{P,\varphi}$ (solid red line), and ensemble of rays received at a given station from an extended noise source (yellow half-banana).

We may now express the ground displacement due to P waves at the observing station of coordinates (λ_o, ϕ_o) , as a function of the same quantity at the location of sources, as given by eq. (4.43)

$$F_{\delta,P}(\lambda_o, \phi_o, f_s) = \int_{\Omega^+} 4F_{\delta,P}(\lambda, \phi, f_s, \theta, \varphi) e^{-2\pi f_s \tau(\lambda, \phi)/Q} d\Omega, \quad (4.46)$$

with $d\Omega$ an element of solid angle that corresponds to the ensemble of rays arriving from an Earth surface element around the sources located at (λ, ϕ) . This sum may also be transformed as an integral over the ocean surface by properly mapping Ω to (λ, ϕ) . The elementary solid angle $d\Omega$ is zero for the so-called shadow zones, the regions for which there is no P -wave ray that connects to the observing station. For a single phase of seismic waves, this ensemble of rays has a half-banana shape, as illustrated on figure 13.

From our calculations, we expect that P waves will dominate the signal at large distances from the source. The exact location where P -wave levels overtakes Rayleigh waves depends on the relative attenuation of the two types of waves. With a realistic $Q = 2000$ for the P waves, and $Q = 400$ for the Rayleigh waves, figure 14 shows that it occurs at an epicentric angle of 40° , which is a distance of 4400 km, consistent with the observations reported by Vinnik (1973) using Kazakhstan array data.




FIGURE 14. Estimates of the rms vertical ground displacement associated to Rayleigh or P waves, as a function of the epicentric angle Δ , for a source of intensity $\int F_{p2,surf}(\mathbf{K} = 0, f)df = 4.2 \times 10^4$ hPa² m² over a 330 by 330 km square, assuming an attenuation factor $Q = 2000$ for the P waves (Pasyanos *et al.* 2009), with travel times given by the ak135 reference Earth model (Snoke 2009).

5. Conclusions

We have shown how the same physical process, the interaction of ocean surface gravity wave (OSGW) trains, can produce a wide variety of noises, in the atmosphere, ocean and Earth's crust, that can be classified according to their horizontal phase speed. The slowest noises in the ocean are acoustic-gravity waves that dominate pressure records at depths less than about one tenth of the acoustic wavelength. These acoustic-gravity waves cannot exist in the absence of OSGWs and are thus confined to the region of active wave forcing. Intermediate phase speeds correspond to Rayleigh waves that contain most of the energy of the seismic modes for distances less than about 4000 km from the source. Our derivations have also been the occasion to correct previously published asymptotic behaviour for very long period noise ($T > 30$ s). In particular we find that the sources of this long period noise are attenuated on the continental shelves, consistent with previous theory for forced gravity wave motion.

The common source of all these noises should allow a verification of the source magnitude for seismic waves by near-surface measurements of pressure which is dominated by acoustic-gravity modes. In particular, the direct modelling of the acoustic-gravity modes can be compared to pressure measurements in depths less than a few hundreds of meters. Unlike the analysis of seismic noise (e.g. Ardhuin *et al.* 2011), which suffers from poorly known seismic propagation and attenuation factors, the acoustic-gravity wave attenuation over the water column can be accurately predicted and thus pressure measurements in the upper ocean may provide a more quantitative verification of numerically modeled directional surface wave properties. In particular, as proposed by Cooper & Longuet-Higgins (1951), pressure measurements may provide a precise estimate of coastal reflection or wave scattering by currents, sea ice or other effects. Noise records from land-based or bottom-mounted seismometers are more ambiguous because they integrate sources over a large area. Also, as discussed by Hasselmann (1963), Abramovici (1968), and Latham

& Sutton (1966), the variations in water depths and horizontal and vertical variations of properties in the Earth's crust can significantly modify noise properties in both the water column and the crust.

F.A. is funded by ERC grant #240009 "IOWAGA" with additional from the U.S. National Ocean Partnership Program, under grant N00014-10-1-0383. T. C. H. is supported by the U.S. Office of Naval Research Littoral Geosciences & Optics Program and the U.S. National Oceanographic Partnership Program (NOPP). Discussions with S. Webb and L. Marié on the theoretical aspects are gratefully acknowledged.

Appendix A. Coupling coefficients

Using the coupling coefficient D given by Hasselmann (1962, eq. 4.3) for the velocity potentials, our coupling coefficient for the elevation amplitudes is

$$\begin{aligned} D_z(\mathbf{k}, s, \mathbf{k}', s') &= -\frac{g^2 D(\mathbf{k}, s, \mathbf{k}', s')}{i s \sigma s' \sigma' (s \sigma + s' \sigma')} \\ &= \frac{g^2}{s \sigma s' \sigma'} \left\{ \left[\mathbf{k} \cdot \mathbf{k}' - \frac{\sigma^2 \sigma'^2}{g^2} \right] + \frac{0.5}{(s \sigma + s' \sigma')} \left(\frac{s \sigma k'^2}{\cosh^2(k'h)} + \frac{s' \sigma' k^2}{\cosh^2(kh)} \right) \right\} \end{aligned} \quad (\text{A } 1)$$

In the bottom pressure, the additional term arising from the orbital velocity has a coupling coefficient

$$D_{pb}(\mathbf{k}, s, \mathbf{k}', s', z) = g^2 \frac{k k' \sinh[k(z+h)] \sinh[k'(z+h)] - \mathbf{k} \cdot \mathbf{k}' \cosh[k(z+h)] \cosh[k'(z+h)]}{2 s \sigma s' \sigma' \cosh(kh) \cosh(k'h)}. \quad (\text{A } 2)$$

The relationship with the coupling coefficient C given by Herbers & Guza (1991, their eq. 4) for the bottom pressure, expressed in meters of water, is given by solving eq. (2.15) for ϕ_2 , and then rewriting Bernoulli's equation (2.4), as

$$\frac{p_2}{\rho_w} = \frac{\partial \phi_2}{\partial t} - \frac{1}{2} \left[|\nabla \phi_1|^2 + \left(\frac{\partial \phi_1}{\partial z} \right)^2 \right] \quad (\text{A } 3)$$

This gives, for $z = -h$,

$$C = -\frac{D_z(s \sigma + s' \sigma')^2}{g [g K \tanh(Kh) - (s \sigma + s' \sigma')^2]} + \frac{D_{pb}(z = -h)}{g}. \quad (\text{A } 4)$$

REFERENCES

- ABRAMOVICI, FLAVIAN 1968 Diagnostic diagrams and transfer functions for oceanic wave-guides. *Bull. Seism. Soc. Am.* **58** (1), 426–456.
- AKI, KEIITI & RICHARDS, PAUL G. 2002 *Quantitative seismology*. Sausalito, California: University Science Books, second edition.
- ANDERSON, DON L. & HART, R. S. 1978 Q of the Earth. *J. Geophys. Res.* **83** (B12), 5869–5882.
- ARDHUIN, FABRICE, STUTZMANN, ELEONORE, SCHIMMEL, MARTIN & MANGENEY, ANNE 2011 Ocean wave sources of seismic noise. *J. Geophys. Res.* **116**, C09004.
- ARENDET, STEVE & FRITTS, DAVID C. 2000 Acoustic radiation by ocean surface waves. *J. Fluid Mech.* **415**, 1–21.
- ARROWSMITH, STEPHEN J., JOHNSON, JEFFREY B., DROB, DOUGLAS P., & HEDLIN, MICHAEL

- A. H. 2010 The seismoacoustic wavefield: a new paradigm in studying geophysical phenomena. *Rev. of Geophys.* **48**, RG4003.
- COOPER, R. I. B. & LONGUET-HIGGINS, MICHAEL S. 1951 An experimental study of the pressure variations in standing water waves. *Proc. Roy. Soc. Lond. A* **206**, 426–435.
- COX, CHARLES S. & JACOBS, DAVID C. 1989 Cartesian diver observations of double frequency pressure fluctuations in the upper levels of the ocean. *Geophys. Res. Lett.* **16** (8), 807–810.
- DUENNEBIER, FRED K., LUKAS, ROGER, NOSAL, EVA-MARIE, AUCAN, JÉRÔME & WELLER, ROBERT A. 2012 Wind, waves, and acoustic background levels at station aloha. *J. Geophys. Res.* **117**, C03017.
- FARRELL, W. E. & MUNK, WALTER 2008 What do deep sea pressure fluctuations tell about short surface waves? *Geophys. Res. Lett.* **35** (7), L19605.
- FARRELL, W. E. & MUNK, WALTER 2010 Booms and busts in the deep. *J. Phys. Oceanogr.* **40** (9), 2159–2169.
- HASSELMANN, K. 1962 On the non-linear energy transfer in a gravity wave spectrum, part 1: general theory. *J. Fluid Mech.* **12**, 481–501.
- HASSELMANN, K. 1963 A statistical analysis of the generation of microseisms. *Rev. of Geophys.* **1** (2), 177–210.
- HASSELMANN, K. 1966 Feynman diagrams and interaction rules of wave-wave scattering processes. *Rev. of Geophys.* **4** (1), 1–32.
- HASSELMANN, K., BARNETT, T. P., BOUWS, E., CARLSON, H., CARTWRIGHT, D. E., ENKE, K., EWING, J. A., GIENAPP, H., HASSELMANN, D. E., KRUSEMAN, P., MEERBURG, A., MÜLLER, P., OLBERS, D. J., RICHTER, K., SELL, W. & WALDEN, H. 1973 Measurements of wind-wave growth and swell decay during the Joint North Sea Wave Project. *Deut. Hydrogr. Z.* **8** (12), 1–95, suppl. A.
- HERBERS, T. H. C. & GUZA, R. T. 1991 Wind-wave nonlinearity observed at the sea floor. part I: forced-wave energy. *J. Phys. Oceanogr.* **21**, 1740–1761.
- HERBERS, T. H. C. & GUZA, R. T. 1994 Nonlinear wave interactions and high-frequency seafloor pressure. *J. Geophys. Res.* **99**, 10035–10048.
- HUGHES, B. 1976 Estimates of underwater sound (and infrasound) produced by nonlinearly interacting ocean waves. *J. Acoust. Soc. Amer.* **60** (5), 1032–1039.
- KANAMORI, HIRO & GIVEN, JEFFERY W. 1981 Use of long-period surface waves for rapid determination of earthquake-source parameters. *Phys. Earth and Planet. Int.* **27** (1), 8–31.
- KEDAR, SHARON, LONGUET-HIGGINS, MICHAEL, GRAHAM, FRANK WEBB NICHOLAS, CLAYTON, ROBERT & JONES, CATHLEEN 2008 The origin of deep ocean microseisms in the north Atlantic ocean. *Proc. Roy. Soc. Lond. A* pp. 1–35.
- KIBBLEWHITE, A. C. & EWANS, K. C. 1985 Wave-wave interactions, microseisms, and infrasonic ambient noise in the ocean. *J. Acoust. Soc. Amer.* **76** (3), 981–994.
- KNUDSEN, V. O., ALFORD, R. S. & EMLING, J. W. 1948 Underwater ambient noise. *J. Mar. Res.* **7**, 410–429.
- LAMB, HORACE 1932 *Hydrodynamics*, 6th edn. Cambridge, England: Cambridge University Press.
- DE LAPLACE, PIERRE SIMON 1776 Suite des recherches sur plusieurs points du système du monde (XXV–XXVII). *Mém. Présentés Acad. R. Sci. Inst. France* pp. 542–552.
- LATHAM, GARY V. & SUTTON, GEORGE H. 1966 Seismic measurements on the ocean floor. *J. Geophys. Res.* **71** (10), 2545–2573.
- LIGHTHILL, JAMES 1978 *Waves in fluids*. Cambridge, United Kingdom: Cambridge University Press, 504 p.
- LIGHTHILL, J. M. 1952 On sound generated aerodynamically. *Proc. Roy. Soc. Lond. A* **211**, 564–587.
- LLOYD, STUART P. 1981 Underwater sound from surface waves according to the Lighthill-Ribner theory. *J. Acoust. Soc. Amer.* **69** (2), 425–435.
- LONGUET-HIGGINS, MICHAEL S. 1950 A theory of the origin of microseisms. *Phil. Trans. Roy. Soc. London A* **243**, 1–35.
- MILLER, G. F. & PURSEY, H. 1955 On the partition of energy between elastic waves in a semi-infinite solid. *Proc. Roy. Soc. Lond. A* **233** (1192), 55–69.
- OBREBSKI, MATHIAS, ARDHUIN, FABRICE, STUTZMANN, ELEONORE & SCHIMMEL, MARTIN 2012

- How moderate sea states can generate loud seismic noise in the deep ocean. *Geophys. Res. Lett.* **39**, L11601, in press.
- PASYANOS, MICHAEL E., WALTER, WILLARD R. & MATZEL, ERIC M. 2009 A simultaneous multiphase approach to determine p -wave and s -wave attenuation of the crust and upper mantle. *Bull. Seismol. Soc. Am.* **99**, 3314–3325.
- POSMENTIER, E. 1967 A theory of microbaroms **13**, 487–501.
- SHAPIRO, NIKOLAI M., CAMPILLO, MICHEL, STEHLY, LAURENT & RITZWOLLER, MICHAEL H. 2005 High-resolution surface-wave tomography from ambient seismic noise. *Science* **307**, 1615–1617.
- SNOKE, J. A. 2009 Traveltime tables for iasp91 and ak135. *Seismological Research Letters* **88** (2), 260–262.
- STONELEY, R. 1926 The effect of the ocean on Rayleigh waves. *Mon. Not. Roy. Astron. Soc. Geophys. Suppl.* **1**, 349–356.
- TANIMOTO, TOSHIRO 2007 Excitation of normal modes by non-linear interaction of ocean waves. *Geophys. J. Int.* **168**, 571–582.
- TYLER, G. L., TEAGUE, C. C., STEWART, R. H., PETERSON, A. M., MUNK, W. H. & JOY, J. W. 1974 Wave directional spectra from synthetic aperture observations of radio scatter. *Deep Sea Res.* **21**, 989–1016.
- VINNIK, L. P. 1973 Sources of microseismic P waves. *Pure Appl. Geophys.* **103** (1), 282–289.
- WAXLER, ROGER & GILBERT, KENNETH E. 2006 The radiation of atmospheric microbaroms by ocean waves. *J. Acoust. Soc. Amer.* **119**, 2651–2664.
- WEBB, SPAHR C. 2007 The earth's 'hum' is driven by ocean waves over the continental shelves. *Nature* **445**, 754–756.