



**HAL**  
open science

## No-regret Dynamics and Fictitious Play

Yannick Viossat, Andriy Zapechelnyuk

► **To cite this version:**

Yannick Viossat, Andriy Zapechelnyuk. No-regret Dynamics and Fictitious Play. 2012. hal-00713871v1

**HAL Id: hal-00713871**

**<https://hal.science/hal-00713871v1>**

Preprint submitted on 2 Jul 2012 (v1), last revised 7 Sep 2012 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# No-regret Dynamics and Fictitious Play\*

Yannick Viossat<sup>†</sup>      Andriy Zapechelnyuk<sup>‡</sup>

July 3, 2012

## Abstract

Potential based no-regret dynamics are shown to be related to fictitious play. This allows to give alternative and sometimes much shorter proofs of known results on convergence of no-regret dynamics to the set of Nash equilibria.

*Keywords:* Regret minimization, no-regret strategy, fictitious play, best reply dynamics, Nash equilibrium, Hannan set, curb set

*JEL classification numbers:* C73, D81, D83

---

\*The authors thank William Sandholm whose comments led to a substantial improvement of the paper, as well as Josef Hofbauer, Mathieu Faure, Sergiu Hart, Alexander Matros, Karl Schlag, Eilon Solan and Sylvain Sorin for helpful comments and suggestions.

<sup>†</sup>CEREMADE, Université Paris-Dauphine, Place du Maréchal de Lattre de Tassigny, F-75775 Paris, France. *E-mail:* viossat  $\alpha\tau$  ceremade.dauphine.fr

<sup>‡</sup>School of Economics and Finance, Queen Mary, University of London, Mile End Road, London E1 4NS, UK. *E-mail:* a.zapechelnyuk  $\alpha\tau$  qmul.ac.uk

# 1 Introduction

*No-regret strategies* are simple adaptive learning rules that recently received a lot of attention in the literature.<sup>1</sup> In a repeated game, a player has a *regret* for an action if, loosely speaking, she could have obtained a greater average payoff had she played that action more often in the past. In the course of the game, the player reinforces actions that she regrets not having played enough, for instance, by choosing next action with probability proportional to the regret for that action, as in Hart and Mas-Colell's (2000) *regret matching* rule. A player's objective is to have no regrets in the long run, *no matter what the opponent does*. A behavior rule of a player which fulfills this objective is called a *no-regret strategy*. Existence of no-regret strategies is known since Hannan (1957); wide classes of no-regret strategies are identified by Hart and Mas-Colell (2001) and Cesa-Bianchi and Lugosi (2003).<sup>2</sup>

A *no-regret dynamics* is a stochastic process that describes trajectories of the average correlated play of players and that emerges when every player follows a no-regret strategy (different players may play different strategies). By definition, it converges to the Hannan set (the set of all correlated actions that satisfy the no-regret condition first stated by Hannan (1957)).<sup>3</sup> The Hannan set, however, is typically large. It contains, and generally properly contains, the set of correlated equilibria of the game. We show that it may even contain correlated actions that put positive weight *only* on strictly dominated actions. Thus convergence of the average play to the Hannan set often provides very little information about what the players will actually play, as it does not even imply exclusion of strictly dominated actions.

In this paper we show that no-regret dynamics are intimately linked to the classical fictitious play process (Brown, 1951). Drawing on Monderer et al. (1997), we first show that contrary to the standard, discrete-time version, continuous fictitious play leads to no-

---

<sup>1</sup>These rules have been used to investigate convergence to equilibria in the context of learning in games (Fudenberg and Levine (1995), Freund and Schapire (1999), Hart and Mas-Colell (2000, 2001, 2003)), for combining different forecasts (Foster and Vohra, 1993, 1999; for an overview of the forecast combination literature see Timmerman, 2006, and Clemen and Winkler, 2007) and for combining opinions, which is also of interest to management science (Larrick and Soll, 2006). In finance this method has been used to derive bounds on the prices of financial instruments (DeMarzo et al., 2006, Chen and Vaughan, 2010). This method can be applied to various tasks in computer science, such as job scheduling (Mansour, 2010) and routing (Blum et al., 2006) (for a survey of applicable problems in computer science see Irani and Karlin, 1996).

<sup>2</sup>This paper deals with the simplest notion of regret known as *unconditional* (or *external*) regret (Fudenberg and Levine, 1995; Hart and Mas-Colell, 2001, 2003). For more sophisticated regret notions, see Hart and Mas-Colell (2000), Lehrer (2003), and Cesa-Bianchi and Lugosi (2006).

<sup>3</sup>The Hannan set of a game is also known as the set of *weak correlated equilibria* (Moulin and Vial, 1978) or *coarse correlated equilibria* (Young, 2004, Ch.3).

regret. We then show that, for a large class of no-regret dynamics, the period-by-period behavior of each player converges to an approximate best reply to the average correlated play of the other players. For two-player finite games, the theory of perturbed differential inclusions (Benaïm et al., 2005, 2006) allows us to relate formally the asymptotic behavior of no-regret dynamics and of continuous fictitious play (or its time-rescaled version, the best-reply dynamics (Gilboa and Matsui, 1991)). In classes of games in which the behavior of continuous fictitious play is well known, this provides substantial information on the asymptotic behavior of no-regret dynamics. In particular, we recover most known convergence properties of no-regret dynamics. Our results do not just allow to find new and sometimes much shorter proofs of convergence of no-regret dynamics towards the set of Nash equilibria in some classes of games, such as dominance solvable game or potential games. They also allow to relate the asymptotic behavior of no-regret dynamics and continuous fictitious play in case of divergence, as in the famous Shapley game (Shapley, 1964).

These results extend only partially to  $n$ -player games (though they fully extend to  $n$ -player games with linear incentives (Selten, 1995)). The issue is that in  $n$ -player games no-regret dynamics turn out to be related with the correlated version of continuous fictitious play, in which the players play a best-reply to the *correlated* past play of the others. This version of fictitious play is defined through a correspondence which is not convex valued. This creates technical difficulties, because the theory of perturbed differential inclusions is not developed for non convex valued correspondences.

A different way to analyze no-regret dynamics is to show that that some sets attract nearby solution trajectories. We show that strict Nash equilibria and, more generally, the intersection of the Hannan set and the sets that are *closed under rational behavior (curb)*<sup>4</sup> are attracting for no-regret dynamics, in a sense to be defined in Section 4.

The remainder of the article is organized as follows. The next section introduces no-regret dynamics. Section 3 studies the links between no-regret dynamics and fictitious play. Section 4 shows that the intersection of the Hannan set and curb sets is attracting for no-regret dynamics. Section 5 shows similar results for the continuous-time version and the expected version of no-regret dynamics. Finally, the Appendix contains the proofs of the main results and a counterexample.

---

<sup>4</sup>A product set of action profiles is called *closed under rational behavior (curb)* if it contains all best replies of each player whenever she believes that no actions outside this set are being played by the other players (Basu and Weibull, 1991).

## 2 Preliminaries

Consider a bimatrix game  $\Gamma = (A_i, u_i)_{i=1,2}$ , where  $A_i$  is the set of actions of player  $i$  and  $u_i : A \rightarrow \mathbb{R}$  is her payoff function, with  $A = A_1 \times A_2$  (we use the term action for the base game, and strategy for the repeated game). For any finite set  $B$ , denote by  $\Delta(B)$  the set of probability distributions over  $B$ . A mixed action of player  $i$  is an element of  $\Delta(A_i)$ . A correlated action  $z$  is a probability distribution over the set of pure action profiles, i.e.  $z \in \Delta(A)$ . Given such a  $z$ , let  $z_i \in \Delta(A_i)$  and  $z_{-i} \in \Delta(A_{-i})$  denote its marginals for player  $i$  and her opponent, respectively. Thus,  $z_i(a_i) = \sum_{a_{-i} \in A_{-i}} z(a_i, a_{-i})$ . Throughout,  $-i$  refers to  $i$ 's opponent. As usual, let  $u_i(z) = \sum_{a \in A} u_i(a)z(a)$  and  $u_i(k, z_{-i}) = \sum_{a_{-i} \in A_{-i}} z_{-i}(a_{-i})u_i(k, a_{-i})$  for  $k \in A_i$ .

The game is played repeatedly in discrete time periods  $t \in \mathbb{N}^* = \{1, 2, \dots\}$ . In every period  $t$  each player  $i$  chooses an action  $a_i(t) \in A_i$  and receives the payoff  $u_i(a(t))$  where  $a(t) = (a_1(t), a_2(t))$ . Denote by  $h(t) = (a(1), a(2), \dots, a(t))$  the history of play up to  $t$ , and let  $\mathcal{H}$  be the set of all finite histories (including the empty history). A strategy of player  $i$  is a function  $q_i : \mathcal{H} \rightarrow \Delta(A_i)$  that stipulates to play in every period  $t = 1, 2, \dots$  a mixed action  $q_i(t) \equiv q_i(h(t-1))$  as a function of the history before  $t$ . The weight that this mixed action puts on action  $k \in A_i$  is denoted by  $q_{i,k}(t)$ .

The *average correlated play* up to period  $t$  is  $z(t) = \frac{1}{t} \sum_{\tau=1}^t a(\tau)$ , where we identify  $a(\tau)$  with the corresponding vertex of  $\Delta(A)$ . Since  $z(t) = \frac{1}{t} [a(t) + (t-1)z(t-1)]$ , it follows that for all  $t > 1$ ,

$$z(t) - z(t-1) = \frac{1}{t} (a(t) - z(t-1)). \quad (1)$$

For a correlated action  $z$ , the *regret* of player  $i$  for action  $k$  is defined as  $R_{i,k}(z) = u_i(k, z_{-i}) - u_i(z)$ , and her maximal regret as  $R_{i,\max}(z) = \max_{k \in A_i} R_{i,k}(z)$ . Typically we deal with the regret based on the average correlated play,  $z(t)$ , up to every period  $t$ . In this case the regret of player  $i$  for action  $k \in A_i$  is equal to the difference between the average payoff she would have obtained by always playing  $k$  (assuming that her opponent's play remains the same) and her average realized payoff:

$$R_{i,k}(z(t)) = u_i(k, z_{-i}(t)) - u_i(z(t)) = \frac{1}{t} \sum_{\tau=1}^t [u_i(k, a_{-i}(\tau)) - u_i(a(\tau))].$$

To simplify notations, we will often write  $R_{i,k}(t)$  for  $R_{i,k}(z(t))$  and  $R_{i,\max}(t)$  for  $R_{i,\max}(z(t))$ .

Player  $i$  has no asymptotic regret if her average realized payoff is asymptotically no

less than her best-reply payoff against the empirical distribution of her opponent:

$$\limsup_{t \rightarrow \infty} R_{i,\max}(t) \leq 0. \quad (2)$$

A strategy of player  $i$  is a *no-regret strategy* if for any strategy of the other player, inequality (2) holds almost surely. This property is also called *Hannan consistency* (Hart and Mas-Colell, 2001) or *universal consistency* (Fudenberg and Levine, 1995).

It is well known in the literature starting from Hannan (1957) that there exist simple no-regret strategies. Hart and Mas-Colell (2001) describe a wide class of *potential based* no-regret strategies. A twice differentiable, convex function  $P_i : \mathbb{R}^{A_i} \rightarrow \mathbb{R}$  is called a *potential* if it satisfies the following conditions:

**(R1)**  $P_i(\cdot) \geq 0$ , and  $P_i(x) = 0$  for all  $x \in \mathbb{R}_-^{A_i}$ ;

**(R2)**  $\nabla P_i(\cdot) \geq 0$ , and  $\nabla P_i(x) \cdot x > 0$  for all  $x \notin \mathbb{R}_-^{A_i}$ ;

**(R3)** if  $x \notin \mathbb{R}_-^{A_i}$  and  $x_k \leq 0$ , then  $\nabla_k P_i(x) = 0$ ,

where  $\nabla_k$  denote the partial derivative with respect to  $x_i(k)$ . The potential  $P_i$  can be viewed as a generalized distance function between a vector  $x \in \mathbb{R}_{A_i}$  and the nonpositive orthant  $\mathbb{R}_-^{A_i}$ . Let  $R_i(t) = (R_{i,k}(t))_{k \in A_i}$  denote player  $i$ 's regret vector.

**Proposition 1.** *Let  $P_i$  satisfy (R1)–(R3) and let strategy  $q_i$  satisfy*

$$q_{i,k}(t+1) = \frac{\nabla_k P_i(R_i(t))}{\sum_{s \in A_i} \nabla_s P_i(R_i(t))}, \quad \forall k \in A_i, \quad (Q1)$$

*whenever  $R_{i,\max}(t) > 0$ . Then  $q_i$  is a no-regret strategy.*

**Proof.** This holds by Theorem 3.3 of Hart and Mas-Colell (2001), whose conditions (R1) and (R2) are satisfied by our conditions (R1)–(Q1) and (R2), respectively. ■

By condition (R3),  $q_{i,k}(t+1) > 0$  only if  $R_{i,k}(t) > 0$ , or equivalently,  $u_i(k, z_{-i}(t)) > u_i(z(t))$ . Thus, condition (R3) is a *better reply property* that stipulates to assign a positive probability only on better reply actions to the opponent's empirical distribution of play (“better” with respect to the realized payoff).

A standard example of no-regret strategy satisfying the above conditions is obtained by letting  $P_i$  be the  $l_{\mathbf{p}}$ -norm on  $\mathbb{R}_+^{A_i}$ , i.e.  $P_i(x) = (\sum_{k \in A_i} [x_k]_+^{\mathbf{p}})^{1/\mathbf{p}}$  with  $1 < \mathbf{p} < \infty$ . The resulting strategy  $q_i$  is called the  $l_{\mathbf{p}}$ -norm strategy (Hart and Mas-Colell, 2001; Cesa-

Bianchi and Lugosi, 2003). It is defined by

$$q_{i,k}(t+1) = \frac{[R_{i,k}(t)]_+^{\mathbf{p}-1}}{\sum_{s \in A_i} [R_{i,s}(t)]_+^{\mathbf{p}-1}}, \quad \forall k \in A_i,$$

whenever  $R_{i,\max}(t) > 0$ . The  $l_2$ -norm strategy is the *regret-matching strategy* (Hart and Mas-Colell, 2000), that stipulates to play an action in the next period with probability proportional to the regret for that action. For large  $\mathbf{p}$ , the  $l_{\mathbf{p}}$ -norm strategies approximate fictitious play.

We say that the average correlated play  $z(t)$  follows a *no-regret dynamics* if both players use (possibly different) no-regret strategies. A trajectory  $(z(t))_{1 \leq t \leq +\infty}$  of a no-regret dynamics is thus a solution of (1) where  $a(t)$  is a realization of  $(q_1(t), q_2(t))$  and  $q_1, q_2$  are no-regret strategies. We focus on the class  $\mathcal{R}$  of no-regret dynamics such that:

(i) the no-regret strategies  $q_1, q_2$  of the players are potential-based: they satisfy (Q1) for some potentials  $P_1, P_2$  satisfying (R1)-(R3);

(ii) if a player has no-regret then he takes some constant pure action: for each  $i = 1, 2$ , there exists  $c \in A_i$  such that

$$q_i(t+1) = c \quad \text{whenever } R_{i,\max}(t) \leq 0. \quad (\text{Q2})$$

Our results are valid for a somewhat wider class of no-regret dynamics. What we really need, beside a no-regret dynamics, is that from some period  $t_0$  on:

(i') if a player has positive regret for some actions, then she plays one of these actions.

(ii') If a player never has any positive regret, then she plays an  $\varepsilon(t)$ -best-reply to the empirical distribution of her opponent, where  $\varepsilon(t) = \varepsilon(h(t)) \rightarrow 0$  almost surely.

**Remark 1.** Property (i') follows from (Q1). It implies that if  $R_{i,\max}(t) > 0$  in some period  $t$ , then  $R_{i,\max}(t') > 0$  for all  $t' > t$ . Indeed, when an action  $k$  with positive regret is played, the sign of  $R_{i,k}(t)$  does not change, hence the maximal regret remains positive (Hart and Mas-Colell, 2001, Proposition 4.3).

**Remark 2.** Assumption (Q2) is a simple way of ensuring (ii'), and in addition, that if  $R_{i,\max}(t) \leq 0$  for all  $t$ , then  $R_{i,\max}(t) \rightarrow 0$  as  $t \rightarrow +\infty$ .<sup>5</sup> Indeed, if  $R_{i,\max}(t) \leq 0$  for all  $t > t_0$  then by (Q2), for all  $t > t_0$ ,  $tR_{i,c}(t) = t_0R_{i,c}(t_0)$ , hence  $R_{i,c}(t) \rightarrow 0$ . It follows that  $R_{i,\max}(t) \rightarrow 0$  and that for all  $t > t_0$ , player  $i$  plays an  $\varepsilon(t)$ -best reply with  $\varepsilon(t) := \max_{k \in A_i} u_i(k, z_{-i}(t)) - u_i(c, z_{-i}(t)) = R_{i,\max}(t) - R_{i,c}(t) \rightarrow 0$ . For a discussion of other possible assumptions, see Hart and Mas-Colell (2003), Appendix A.

<sup>5</sup>This additional property is needed for Corollary 1 below, but for our main results (ii') suffices.

Note that there are no-regret dynamics that do not satisfy (i'). For instance, stochastic fictitious play with a noise parameter that varies with time and approaches zero at an appropriate rate (see, e.g., Benaïm and Faure, 2011).

Define the *Hannan set*  $H$  of the stage game  $\Gamma$  as the set of all correlated actions of the players where each player has no regret:

$$H = \left\{ z \in \Delta(A) \mid \max_{k \in A_i} u_i(k, z_{-i}) \leq u_i(z) \text{ for each } i = 1, 2 \right\}.$$

The *reduced Hannan set*  $H_R$  is the subset of  $H$  in which at least one regret is exactly zero for each player:

$$H_R = \left\{ z \in \Delta(A) \mid \max_{k \in A_i} u_i(k, z_{-i}) = u_i(z) \text{ for each } i = 1, 2 \right\}.$$

The next property of no-regret dynamics is straightforward by the definition of no-regret strategy and Remark 2 (see, e.g., Hart and Mas-Colell, 2003, Corollary 3.2).

**Corollary 1.** *For every no-regret dynamics in class  $\mathcal{R}$ , the trajectories converge almost surely to the reduced Hannan set.*

Convergence of the average play  $z(t)$  to set  $H_R$  does not imply its convergence to any particular point in  $H_R$ . Moreover, even if  $z(t)$  converges to a point, this point need not be a Nash equilibrium.

## 3 Fictitious play and no-regret dynamics

### 3.1 Fictitious play

In *discrete fictitious play*, in every period  $t$  after the initial one, player  $i$  plays a best reply  $q_i(t)$  to the average past play of her opponent  $x_{-i}(t-1) := \frac{1}{t-1} \sum_{\tau=1}^{t-1} q_{-i}(\tau)$ . The latter is called the *belief* of player  $i$  on her opponent's next move. Formally, for any  $x = (x_1, x_2)$  in  $\Delta(A_1) \times \Delta(A_2)$ , denote by  $BR_i(x_{-i})$  player  $i$ 's set of best replies to  $x_{-i}$ :

$$BR_i(x_{-i}) := \left\{ \sigma_i \in \Delta(A_i) \mid u_i(\sigma_i, x_{-i}) \geq \max_{k \in A_i} u_i(k, x_{-i}) \right\}, \quad i = 1, 2. \quad (3)$$



Let  $BR(x) = BR_1(x_2) \times BR_2(x_1)$ . A discrete-time trajectory  $(x(t))_{t=1}^{\infty}$  on  $\Delta(A_1) \times \Delta(A_2)$  is a solution of *discrete fictitious play* (DFP) if for every  $t > 1$

$$x(t) - x(t-1) = \frac{1}{t} (q(t) - x(t-1)). \quad (4)$$

where  $q(t) = (q_1(t), q_2(t))$  is a pure best reply action profile, i.e.,  $q(t) \in A \cap BR(x(t-1))$ .

Analogously, an absolutely continuous function  $x : [1, \infty) \rightarrow \Delta(A_1) \times \Delta(A_2)$  is a solution of *continuous fictitious play* (CFP) if for almost all  $t \geq 1$ ,  $x(t)$  is differentiable and

$$\dot{x}(t) = \frac{1}{t} (q(t) - x(t)), \quad (5)$$

where  $q(t) \in BR_i(x(t))$  is now a profile of *mixed* actions. This may be written as the differential inclusion:

$$\dot{x} \in \frac{1}{t} (BR(x) - x). \quad (6)$$

The average correlated play satisfies  $z(t) := \frac{1}{t} \left( z(1) + \int_1^t q(\tau) d\tau \right)$  for some initial condition  $z(1)$  such that  $z_i(1) = x_i(1)$ ,  $i = 1, 2$ . Thus,

$$\dot{z}(t) = \frac{1}{t} (q(t) - z(t)), \quad (7)$$

where for almost all  $t$ ,  $q_i$  is a best-reply to  $z_{-i}$ .<sup>6</sup>

In discrete or continuous fictitious play, the marginals  $z_1(t)$ ,  $z_2(t)$  of the average past play are equal to the beliefs  $x_1(t)$ ,  $x_2(t)$ . By analogy, if  $z(t)$  is the average past play generated by a no-regret dynamics, it is convenient to call  $z_{-i}(t)$  the belief of player  $i$  on her opponent's next move. This illuminates a crucial difference between fictitious play and no-regret dynamics in class  $\mathcal{R}$ : under fictitious play, a player chooses a *best reply* to her belief, whereas under no-regret dynamics, she chooses a *better reply* ("better" with respect to her average realized payoff).

---

<sup>6</sup> This definition of CFP guarantees that solutions exist in all games and for all initial conditions, and that by the change of time scale  $y(t) = x(e^t)$ , CFP corresponds to *best-reply dynamics* (Gilboa and Matsui, 1991; Matsui, 1992) defined by  $\dot{y} \in BR(y) - y$ . Another definition of CFP (e.g., Monderer et al., 1997, p. 445; Berger, 2007, pp. 252–253) considers only trajectories that are piecewise linear, such that  $q_i(t)$  is always a pure strategy, and that the times at which  $q(t)$  changes have no finite accumulation point. This restricted definition is easier to handle, but in many games such trajectories do not exist.

### 3.2 Continuous fictitious play leads to no regret

It is well known that discrete fictitious play does not lead to *no regret* (Young, 1993; Hart and Mas-Colell, 2001). Consider the following example:

	$L$	$R$
$L$	$1, \sqrt{2}$	$0, 0$
$R$	$0, 0$	$\sqrt{2}, 1$

Fig. 1

Because  $\sqrt{2}$  is irrational,  $L$  and  $R$  cannot both be best-replies to the empirical past play of the other player. Thus, any DFP process is entirely determined by its first move. Assume that the first move is off the diagonal, say  $(L, R)$ . Due to the symmetry of the game and the absence of ties, both players always switch to another action simultaneously. Therefore the play is locked off the diagonal and the maximal regret is at least  $\sqrt{2}/(1 + \sqrt{2})$  at any stage. This holds in the mixed extension of the game, since at any stage the players have a unique, pure best reply.

Since the continuous fictitious play process is a continuous-time version of DFP, intuitively, it should not lead to *no regret* either. The following result — a generalization of Theorem D of Monderer et al. (1997) — shows that this intuition is misleading.

**Proposition 2.** *Under any solution of continuous fictitious play, the average correlated play converges to the reduced Hannan set.*

This discrepancy between DFP and CFP may be explained as follows. Playing an action with positive regret decreases the regret for this action. In CFP, when an action is played it remains a best reply, hence it is associated with maximal regret for a small time increment.<sup>7</sup> Since the regret for this action decreases, so does the maximal regret. In contrast, in DFP, an action played at stage  $t$  has maximal regret at stage  $t$ , but not necessarily at stage  $t + 1$ . Thus the fact that the regret for this action decreases does not entail that the maximal regret does.

**Proof of Proposition 2.** For comparison with Hart and Mas-Colell (2003, Theorem 3.1), rescale time (let  $\tilde{t} = \exp t$ ) so that (7) become  $\dot{z} = q - z$ . For any mixed action  $\sigma_i \in \Delta(A_i)$ , let

$$R_{i,\sigma_i(t)} := \sum_{k \in A_i} \sigma_i(k) R_{i,k}(t) = u_i(\sigma_i, z_{-i}(t)) - u_i(z(t))$$

---

<sup>7</sup>This only holds up to a second order term (see the proof below), but this suffices.

Let  $v_i(t) = R_{i,\max}(t)$ . For almost all  $t$ , the function  $v_i$  is differentiable and the following holds: if  $k \in \text{Supp}(q_i(t))$ , then for any integer  $n \in \mathbb{N}^*$ , there exists  $t_n \in ]t, t+1/n[$  such that  $k$  is a best-reply to  $z_{-i}(t_n)$  (otherwise,  $\dot{z}_{i,k} = -z_{i,k}$  on  $]t, t+1/n[$ , hence  $z_i(t+s) = z_i(t)e^{-s}$  for all  $s \in ]0, 1/n[$ ; this implies  $q_{i,k} = 0$ , a contradiction). Therefore,  $v_i(t_n) = R_{i,k}(t_n)$ . Moreover,  $v_i(t) = R_{i,k}(t)$ . It follows that for any  $k \in \text{Supp}(q_i(t))$ ,  $\dot{v}_i = \dot{R}_{i,k}(t)$ , hence  $\dot{v}_i = \sum_k q_{i,k} \dot{R}_{i,k} = \dot{R}_{i,q_i}$ . Furthermore:

$$\dot{R}_{i,q_i} = u_i(q_i, \dot{z}_{-i}) - u_i(\dot{z}) = u_i(q_i, q_{-i} - z_{-i}) - u_i(q - z) = -[u_i(q_i, z_{-i}) - u_i(z)] = -R_{i,q_i} = -v_i.$$

Thus,  $\dot{v}_i = -v_i$ . Therefore,  $v_i(t)$  converges to zero for all  $i = 1, 2$ , hence  $z(t) \rightarrow H_r$ . ■

**Remark 3.** In the proof, we did not use that  $q_{-i}$  is a best-reply to  $z_i$ . This shows that the fact that CFP leads to no-regret is a unilateral property. That is, if a player's behavior evolves according to CFP, then she has no asymptotic regret, independently of her opponent's behavior (see also Monderer et al., 1997, p. 445).

**Remark 4.** CFP and the best-reply dynamics converge to the set of Nash equilibria in finite zero-sum games (Hofbauer and Sorin, 2006). The usual proof is to show that the ‘‘duality gap’’  $W(x) = \max_{k \in A_1} u_1(k, x_2) - \min_{s \in A_2} u_1(x_1, s)$  converges to zero. This follows from the above proof, since in a two-player zero sum game  $W(x(t)) = R_{1,\max}(z(t)) + R_{2,\max}(z(t))$ , where  $x$  is a solution of CFP and  $z$  the associated correlated play.

### 3.3 No-regret dynamics is perturbed CFP

In the previous subsection we showed that CFP leads to no regret. Conversely, we now show that any no-regret dynamics in class  $\mathcal{R}$  (as defined in Section 2) is closely related to CFP. We first explain the intuition. The crucial observation is the following.

**Lemma 1.** *Assume that the maximal regret is less than  $\varepsilon$ . Then any action with positive regret is an  $\varepsilon$ -best reply to the average play of the opponent.*

**Proof.** If player  $i$  has positive regret for action  $a_i$  at some  $z \in \Delta(A)$ , then  $u_i(z) - u_i(a_i, z_{-i}) < 0$ . But by assumption  $\max_{k \in A_i} u_i(k, z_{-i}) - u_i(z) \leq \varepsilon$ . Therefore,  $\max_{k \in A_i} u_i(k, z_{-i}) - u_i(a_i, z_{-i}) < \varepsilon$ , and  $a_i$  is an  $\varepsilon$ -best reply to  $z_{-i}$ . ■

Since no-regret dynamics in class  $\mathcal{R}$  only pick actions with positive regret, they only pick  $\varepsilon$ -best replies to the average play of the others, where  $\varepsilon$  is the maximal regret. Since this maximal regret approaches zero almost surely, eventually only almost-exact

best replies are picked. This provides the intuition why no-regret dynamics and fictitious play may exhibit similar asymptotic behavior. Finding a precise link, however, is not obvious. For instance, there could exist actions that are  $\varepsilon_t$ -best replies in each period  $t$ , with  $\varepsilon_t \rightarrow 0$ , but never exact best replies. Thus a limit play of no-regret dynamics may include such actions, but this cannot happen under fictitious play.

	$L$	$R$
$L$	$1, 0$	$0, \sqrt{2}$
$R$	$0, 1$	$\sqrt{2}, 0$
$C$	$\eta, 0$	$\eta, 0$

Fig. 2

Consider the example shown on Fig. 2. Let  $\eta = \sqrt{2}/(1 + \sqrt{2})$ . It is easy to verify that action  $C$  is player 1's best reply to player 2's mixed action  $x_2$  if and only if  $x_2 = (\eta, 1 - \eta)$ . Let us first consider DFP. Since  $\eta$  is an irrational number, after every finite history of play,  $C \notin BR_1(x_2(t))$ ; consequently DFP never picks  $C$  (except, possibly, at the initial period).<sup>8</sup> However, it may be shown that under any DFP trajectory, the average play  $x_2(t)$  of player 2 converges to  $(\eta, 1 - \eta)$ , to which  $C$  is a best-reply. It follows that  $C$  is an  $\varepsilon_t$ -best reply to  $x_2(t)$  for some sequence  $\varepsilon_t \rightarrow 0$ . Thus a no-regret dynamics with the same trajectory of the marginal play of player 2 *might* choose action  $C$  a positive fraction of time in the long run.

This example does not apply to CFP, as in this case  $x_2(t)$  need not be a rational number; and as we show below, the asymptotic behavior of no-regret dynamics and CFP can be formally related using the theory of perturbed differential inclusions (Benaïm et al., 2005, 2006).

Before stating a precise result, we need some definitions. A set  $L \subset \Delta(A_1) \times \Delta(A_2)$  is *invariant* under CFP if for every initial point  $x \in L$  there exists a solution  $x(\cdot)$  of CFP, defined for all  $t > 0$  (not only  $t \geq 1$ ) and such that  $x(1) = x$  and  $x(t) \in L$  for all  $t > 0$ . A nonempty compact invariant set is an *attractor* if it attracts uniformly all trajectories starting in its neighborhood. An invariant set  $L$  is *attractor-free* if no proper subset of  $L$  is an attractor for the dynamics restricted to  $L$ . A nonempty compact set  $L$  is *internally chain transitive* (ICT) if every pair of points in  $L$  can be connected by finitely many arbitrarily long pieces of orbits of the best-reply dynamics lying completely

---

<sup>8</sup>Starting with an arbitrary belief  $x_2(1)$  would not help since  $C$  is a best-reply only when  $x_2(t) = (\eta, 1 - \eta)$ , which happens at most once.

within  $L$  with arbitrarily small jumps between them.<sup>9</sup> Every ICT set is invariant and attractor free (Benaïm et al., 2006, Property 2). The *limit set of the beliefs* of a trajectory  $z(t)$  on  $\Delta(A_1 \times A_2)$  is the set of all accumulation points of its marginals  $(z_1(t), z_2(t)) \in \Delta(A_1) \times \Delta(A_2)$  as  $t \rightarrow \infty$ .

**Theorem 1.** *For every no-regret dynamics in class  $\mathcal{R}$ , the limit set of the beliefs is almost surely internally chain transitive for continuous fictitious play.*<sup>10</sup>

The proof is deferred to the Appendix. A consequence of this theorem is the following:

**Corollary 2.** *Let  $\mathcal{A}$  be the global attractor (i.e. the maximal invariant set) of CFP. For any no-regret dynamics in class  $\mathcal{R}$ , the limit set of the beliefs is almost surely a subset of  $\mathcal{A}$ .*

Note the similarity with Propositions 5.1 and 5.2 of Hofbauer et al. (2009), who study the links between the time-average of the replicator dynamics and CFP.

### 3.4 Applications of Theorem 1 and comments.

Theorem 1 allows us to find alternative and sometimes much shorter proofs of most known properties of no-regret dynamics. Note that some applications of Theorem 1 (points (a), (b) and (c) below) lead to the same conclusions about no-regret dynamics as those about the time average of the replicator dynamics described in Hofbauer et al. (2009, p. 267, points (2), (3) and (4)).

Below, we write that no-regret dynamics converge to some set  $E$  if the limit set of the beliefs is almost surely a subset of  $E$ .

**(a)** For any game which is best-reply equivalent to a two-person zero sum game, the global attractor of CFP is the set of Nash equilibria (Hofbauer and Sorin, 2006). Hence all no-regret dynamics in class  $\mathcal{R}$  converge to the set of Nash equilibria. Actually, in zero-sum games, if the correlated action  $z$  is in the Hannan set (recall that this is the set of correlated actions that satisfy *no-regret* for all players), then  $(z_1, z_2)$  is a Nash equilibrium. Consequently, in zero-sum games all dynamics that lead to no regret (not

---

<sup>9</sup>For the formal definitions of *attractor* and *attractor-free set* see Benaïm et al. (2006, p. 675); for the definition of *ICT* see Benaïm et al. (2005, p. 337). Note that the definition of invariance in Benaïm et al. (2005, 2006) applies to the best-reply dynamics, so an appropriate time rescaling must be used to apply it to CFP (see footnote 6). This explains that their definition considers solutions defined for all  $t \in \mathbb{R}$  while ours considers solutions defined for all  $t > 0$ .

<sup>10</sup>In the statement of Theorem 1, CFP can be replaced by the best-reply dynamics since they clearly have the same ICT sets (see footnote 6).

only those in class  $\mathcal{R}$ ) converge to the set of Nash equilibria. This holds more generally for *stable bimatrix games* (Hofbauer and Sandholm, 2009), because these are rescaled zero-sum games in the sense of Hofbauer and Sigmund (1998), as is easily shown and was known to Josef Hofbauer (private communication).

**(b)** For games with strictly dominated strategies, the global attractor of CFP is contained in the face of the simplex with no weight on these strategies. Hence all no-regret dynamics in class  $\mathcal{R}$  converge to this face. Similarly, these dynamics converge to the unique Nash equilibrium in strictly dominance solvable games.

	$A$	$B$	$C$
$A$	2	1	-4
$B$	1	0	-1
$C$	-4	-1	-2

(i)

	$A$	$A^-$	$B$	$B^-$
$A$	1	1	0	0
$A^-$	$1 - \varepsilon$	$1 - \varepsilon$	$-\varepsilon$	$-\varepsilon$
$B$	0	0	1	1
$B^-$	$-\varepsilon$	$-\varepsilon$	$1 - \varepsilon$	$1 - \varepsilon$

(ii)

Fig. 3

Contrary to (a), this need not be true for all dynamics that lead to no regret. Indeed, convergence to the Hannan set or even to the reduced Hannan set does not guarantee elimination of strictly dominated strategies. Consider, for instance, the games shown on Fig. 3. Both games are symmetric, so we indicate only the payoffs of the row player. Game (i) is an identical interest game which is strictly dominance solvable; yet the correlated action putting probabilities  $1/3$  on each diagonal square is in the reduced Hannan set. For  $\varepsilon = 0$ , game (ii) is a coordination game with duplicate strategies. For  $\varepsilon > 0$ , the duplicates  $A^-$ ,  $B^-$  are penalized and become strictly dominated. Thus, the correlated action putting probability  $1/2$  on  $(A^-, A^-)$  and  $1/2$  on  $(B^-, B^-)$  puts only weight on strictly dominated actions. Yet, for  $\varepsilon \leq 1/2$ , it belongs to the Hannan set.<sup>11</sup>

**(c)** In weighted potential games, all internally chain transitive sets of CFP are (subsets of) connected components of Nash equilibria on which the payoffs are constant (see Benaïm et al., 2005, Theorem 5.5 and Remark 5.6). Hence by Theorem 1, all no-regret dynamics in class  $\mathcal{R}$  converge to such components. Note that the original proof is much longer (Hart and Mas-Colell, 2003, Appendix A).

**(d)** If the beliefs  $(z_1(t), z_2(t))$  of a no-regret dynamics converge to the set of Nash equilibria, then the average realized payoff converges to the set of Nash equilibrium payoffs.

<sup>11</sup>See also the game of Moulin and Vial (1978, p. 205), where the third strategy of player 1 is strictly dominated but has a positive marginal probability under some correlated actions in the Hannan set.

To see why this is true, let  $\hat{z} \in \Delta(A)$  be a limit point of  $\{z(t)\}$  and let the marginals  $(\hat{z}_1, \hat{z}_2) \in \Delta(A_1) \times \Delta(A_2)$  constitute a Nash equilibrium. By Corollary 1 the maximal regret converges to zero, so for every  $i = 1, 2$

$$u_i(\hat{z}) = \max_{k \in A_i} u_i(k, \hat{z}_{-i}) = u_i(\hat{z}_i, \hat{z}_{-i}).$$

This result illuminates an important difference between no-regret dynamics and discrete fictitious play. It is well known that under DFP, if the beliefs of the players converge to a Nash equilibrium, their average realized payoffs *need not* approach the set of Nash equilibrium payoffs (as in the example illustrated by Fig. 1), whereas under no-regret dynamics it is always the case.

(e) Consider the  $3 \times 3$  game of Fig. 4 due to Shapley (1964), the historical counterexample to the convergence of fictitious play. This game has a unique equilibrium, in which both

	<i>A</i>	<i>B</i>	<i>C</i>
<i>A</i>	0, 0	1, 0	0, 1
<i>B</i>	0, 1	0, 0	1, 0
<i>C</i>	1, 0	0, 1	0, 0

Fig. 4

players randomize uniformly. Though this equilibrium attracts some solutions of continuous fictitious play (e.g. all those that start and remain symmetric), almost all solutions converge to an hexagon, the so-called Shapley polygon (Shapley, 1964; Gaunersdorfer and Hofbauer, 1995; Sparrow et al., 2008). It may be shown that the only ICT sets are the Nash equilibrium and the Shapley polygon. Consequently, the limit set of any no-regret dynamics in class  $\mathcal{R}$  is almost surely one of these two sets.

(f) In a number of classes of games, convergence of discrete fictitious play to the set of Nash equilibria has been established, but analogous results for continuous fictitious play are lacking. Thus we cannot use Theorem 1. These classes of games include generic  $2 \times n$  games (Berger, 2005), generic ordinal potential games, quasi-supermodular games<sup>12</sup> with diminishing returns (Berger, 2007) and some other special classes (see, e.g., Sparrow et al., 2008, p. 260). For ordinal potential games and quasi-supermodular games with diminishing returns, Berger (2007) proves convergence to the set of Nash equilibria of *some* solutions of continuous fictitious play as defined by (6) (see our footnote 6). This

---

<sup>12</sup>Also known as games of strategic complementarities (e.g., Tirole, 1988).

is not enough to apply the results of Benaïm et al. (2005). The same problem arises in Krishna and Sjöström (1998). Actually, as explained below, convergence of CFP to the set of Nash equilibria would not suffice to use Theorem 1: we would need some additional structure, such as a Lyapunov function, to get more information on the ICT sets.

**(g)** Consider a bimatrix game in which all solutions of CFP converge to the set of Nash equilibria. Because the definition of attractor requires a uniform attraction, this does not imply that the set of Nash equilibria is an attractor. Neither does it imply that all ICT sets are contained in the set of Nash equilibria, as shown in Appendix A.2. Therefore, we cannot deduce from Theorem 1 that no-regret dynamics in class  $\mathcal{R}$  converge to the set of Nash equilibria; whether this is always the case remains an open question.

**(h)** We show in Section 5 that Theorem 1 also applies, and under weaker assumptions, to the continuous-time version and to the expected version of no-regret dynamics in class  $\mathcal{R}$ . As apparent from the proof, the existence of a potential is not essential: for a good choice of behavior when there are no regrets, Theorem 1 holds for any no-regret dynamics such that a player always chooses an action with positive regret whenever he has some. It also applies to no-regret dynamics that do not have this property, such as the exponential weight algorithm (see Remark 6 at the end of Appendix A.1).

**(i)** Let us now comment on extensions of our results to  $n$ -player games. The definition of no-regret dynamics, as well as Proposition 1, extend to the  $n$ -player setting straightforwardly (e.g., Hart and Mas-Colell, 2001). The appropriate extension of CFP is *correlated CFP* where at each time  $t$  every player chooses a best reply action to the *correlated* past average play of the others. Specifically, an absolutely continuous function  $z : [1, \infty) \rightarrow \Delta(A)$  is a solution of correlated CFP if it is almost everywhere differentiable and satisfies

$$\dot{z}(t) \in \frac{1}{t}(\overline{BR}(z(t)) - z(t)),$$

where the correlated best-reply correspondence  $\overline{BR} : \Delta(A) \rightrightarrows \Delta(A)$  is defined by  $BR(z) = \times_{i=1}^n \overline{BR}_i(z_{-i})$  where  $\overline{BR}_i(z_{-i})$  is the set of mixed best replies of player  $i$  to the correlated action  $z_{-i} \in \Delta(A_{-i})$ .

In  $n$ -player games with linear incentives (Selten, 1995), also known as polymatrix games (Yanovskaya, 1968), the correlated and independent best-reply correspondences coincide; that is, for any correlated action  $z \in \Delta(A)$ ,  $\overline{BR}(z) = BR((z_1, \dots, z_n))$  where  $(z_1, \dots, z_n)$  is the vector of marginals of  $z$ , and  $BR$  the standard (independent) best-reply correspondence. For such games, our results extend easily. However, this is not the case in general. The main problem is that the correlated best reply correspondence is not



convex valued; that is,  $\overline{BR}(z)$  is not in general a convex subset of  $\Delta(A)$ .<sup>13</sup> This creates two issues:

(i) Existence of solutions of correlated CFP is not guaranteed by the classical results on differential inclusions we are aware of (e.g., Aubin and Cellina, 1984).

(ii) The theory of perturbed differential inclusions (Benaïm et al., 2005) does not apply to non-convex valued correspondences.

The first issue can be solved by building piecewise linear solutions of correlated CFP following the same ideas as for two-player games.<sup>14</sup> Moreover, due to Remark 3, Proposition 2 extends to the  $n$ -player setting. It then asserts that correlated CFP leads to no regrets. Lemma 1 also extends: it asserts that if the maximal regret of player  $i$  is less than  $\varepsilon$ , then she plays only  $\varepsilon$ -best reply actions to the *correlated* average play of the opponents. It follows that, analogously to two-player games, interpolated trajectories of no-regret dynamics are almost surely perturbed solutions of correlated CFP/ However, we cannot proceed to an analog of Theorem 1 because of the second issue. Thus, whether there is a formal relation between no-regret dynamics and correlated CFP in  $n$ -player games remains an open question. Similarly, the results of Hofbauer et al. (2009) on the links between the time-average of the replicator dynamics and CFP are restricted to bimatrix games (or games with linear incentives).

## 4 Curb sets

Theorem 1 does not answer whether *attracting sets* of CFP have an analogous property under no-regret dynamics.

A set  $\mathcal{C} \subset \Delta(A)$  is *eventually attracting* under a no-regret dynamic process if with any given probability it captures all no-regret trajectories originated from a small enough neighborhood of  $\mathcal{C}$  at all distant enough periods. Formally,  $\mathcal{C}$  is eventually attracting if for every  $\pi > 0$  there exists  $\varepsilon > 0$  and a period  $T$  such that: for every  $t_0 \geq T$ , if  $z(t_0)$  is in

---

<sup>13</sup>This is due to the fact that elements of  $\overline{BR}(z)$  are independent distributions and that the average of two independent distributions need not be an independent distribution.

<sup>14</sup>Assuming that  $z(t)$  is well defined, call  $G_r(t)$  the game in which the players are reduced to their best-replies to  $z(t)$ . Start with some initial condition  $z(t_0)$ . Then point to a Nash equilibrium of  $G_r(t_0)$  (i.e. fix  $b \in NE(G_r(T_0))$  and choose  $q(t) = b$ ) till the first time,  $t_1$ , when, for some player  $i$ , a strategy which was not a best-reply to  $z(t_0)$  is a best-reply to  $z(t_1)$ . Then iterate. If the times  $t_n$  accumulate towards some time  $t^*$ , then use the fact that  $z(t)$  must have a limit when  $t \rightarrow t^*$  (because  $z(t)$  is Lipschitz). Call it  $z(t^*)$  and restart from  $z(t^*)$ . Note that there might in principle be a countable infinity of such accumulation points  $t^*$ , and that they might themselves accumulate in some point  $t^{**}$ , but then define  $z^{**}$  as before and restart from there, etc. The largest (forward time) interval on which such a solution can be built is both open and closed in  $[t_0, +\infty[$  and is thus equal to  $[t_0, +\infty[$ .

an  $\varepsilon$ -neighborhood of  $\mathcal{C}$ , then  $z(t)$  converges to the set  $\mathcal{C}$  with probability at least  $1 - \pi$ .<sup>15</sup>

For this section it is convenient to replace assumption (Q2) by the following one:

(Q2')

If a player's maximal regret is nonpositive, then she plays a best-reply to the empirical distribution of her opponent.

This is not essential, since the interesting histories are those where both players have positive regrets, in which case (Q2) plays no role.<sup>16</sup>

A strict Nash equilibrium is eventually attracting. Indeed, if  $z(t_0)$  is close enough to a strict Nash equilibrium  $a = (a_1, a_2)$ , then for each player  $i$ , action  $a_i$  is the unique best reply and there is a negative regret for any action other than  $a_i$ . Since by (R3) only actions with positive regret can be chosen, and by (Q2') only best-reply actions can be chosen if *all* regrets are nonpositive, action  $a_i$  will be played by each player  $i$  in the following period, and so on.

Let us now consider a standard generalization of strict Nash equilibria. For each  $i = 1, 2$ , let  $B_i \subset A_i$ . With a slight abuse of notation, denote by  $\Delta(B_i)$  the set of probability measures on  $A_i$  with support on  $B_i$  only. The product set  $B = B_1 \times B_2$  is *closed under rational behavior (curb)* (Basu and Weibull (1991)) if

$$BR_i(y_{-i}) \subset \Delta(B_i) \text{ whenever } y_{-i} \in \Delta(B_{-i}), \quad i = 1, 2.$$

That is, the set  $B$  is curb if the players' pure best reply profiles are contained in  $B$  whenever they believe that no actions outside of  $B$  should be played.

Curb sets are known to be attracting under CFP (e.g., Balkenborg et al., 2011, Lemma 7). However, they need not be attracting under no-regret dynamics in class  $\mathcal{R}$ . Indeed, even if the support of  $z(t_0)$  is contained in some curb set  $B$ , there may be positive regrets for actions outside of  $B$ , since  $B$  need not be closed under *better* replies. However, we show that the intersection of the Hannan set and of the set of correlated actions with support on a curb set is eventually attracting.

Formally, let  $B = B_1 \times B_2$  be a curb set. Let  $\Delta_B(A)$  denote the set of correlated actions with support on  $B$  only. Let  $H_B = H \cap \Delta_B(A)$ .

---

<sup>15</sup>We say that  $z(t)$  converges to  $\mathcal{C}$  if  $\inf_{c \in \mathcal{C}} \|z(t) - c\| \rightarrow 0$  as  $t \rightarrow \infty$ .

<sup>16</sup>Recall that by Remark 1, if a player has positive maximal regret, then it remains positive forever. So we can consider histories from a distant enough period  $t_0$  where both players have positive regrets and (Q2) plays no role. If  $t_0$  does not exist, i.e., some player *always* has nonpositive maximal regret, then Proposition 1 and (Q2) imply that her play is constant, whereas her opponent's play must approach a best reply to it, leading to Nash equilibrium. By replacing (Q2) by (Q2') we avoid dealing with this issue.

**Proposition 3.** *For every curb set  $B$ , the set  $H_B$  is eventually attracting under every no-regret dynamics in  $\mathcal{R}$ .*

The proof is deferred to the Appendix. Note that Proposition 3 *does not* claim that if the play is locked on some curb set, all actions of that set will be played infinitely often. For example, a no-regret trajectory may lock on a Nash equilibrium which is not strict, hence not curb, thus being properly contained in some minimal curb set whose other actions are never played.

## 5 Continuous-time and expected no-regret dynamics

We now prove an analog of Theorem 1 for continuous-time dynamics (Hart and Mas-Colell, 2003) and the expected version of discrete-time dynamics. Consider a continuous-time dynamics

$$\dot{z}(t) = \frac{1}{t}(q(t) - z(t)) \quad (8)$$

where  $q(t) \in \prod_i \Delta(A_i)$  is the (independent) joint play at time  $t$  and  $z(t) \in \Delta(A)$  the average correlated play. Note that  $q_i(t)$  is now a *mixed* action. As in continuous fictitious play, start at time 1 with some initial condition  $z(1) \in \Delta(A)$ . Assume that whenever  $\mathbb{R}_{i,\max}(t) > 0$

$$q_{i,k}(t) = \frac{\nabla_k P_i(R_i(t))}{\sum_{s \in A_i} \nabla_s P_i(R_i(t))}, \quad k \in A_i \quad (9)$$

where  $P_i$  is a  $C^1$  potential function satisfying (R1), (R2) and the technical condition:

**(P4')** There exists  $0 < \rho_2 < \infty$  such that  $\nabla P_i(x) \cdot x \leq \rho_2 P_i(x)$  for all  $x \in \mathbb{R}_+^{A_i}$ .

This is a part of condition (P4) in Hart and Mas-Colell (2003). Note that we do not need the better-reply condition (R3).

**Proposition 4.** *Let  $z(t)$  be a solution of (8) and (9) with  $P_i$  satisfying conditions (R1), (R2) and (R4) for all  $i = 1, 2$ . Assume that the initial condition  $z(1)$  is such that both players have some positive regrets:  $R_{i,\max}(1) > 0$  for all  $i = 1, 2$ . Then the conclusion of Theorem 1 applies.*

**Proof.** (We advise the reader to first read Appendix A.1.) Let  $\varepsilon_i(t) := \max_{k \in A_i} R_{i,k}(t)$ . Hart and Mas-Colell (2003, Theorem 3.1 and Lemma 3.3<sup>17</sup>) show that if  $\varepsilon_i(1) > 0$ , then

---

<sup>17</sup>Note a typo in the proof of Lemma 3.3 in Hart and Mas-Colell (2003): (P3) should be replaced by (P4); beside, only our condition (P4') is used.

$\varepsilon_i(t) > 0$  for all  $t$ , and that  $\varepsilon_i(t) \rightarrow 0$  as  $t \rightarrow +\infty$ . Moreover, by (R2) applied to  $x = R_i(t)$  and definition of  $q_i$ , we have:  $u_i(q_i, z_{-i}) - u_i(z) = q_i \cdot R_i > 0$  (this is equation (3.3) in Hart and Mas-Colell, 2001). Thus by Lemma 1,  $q_i \in BR_i^{\varepsilon_i(t)}(z_{-i})$ . Together with Lemma 3 of Appendix A.1, this implies that  $(z_1(\cdot), z_2(\cdot))$  is a perturbed solution of CFP. The result then follows from Theorem 3.6 of Benaïm et al. (2005). ■

**Remark 5.** Assume that if all initial regrets of a player are nonpositive then the dynamics is defined as in Hart and Mas-Colell (2003), equation (4.9). Then it is easily seen that Theorem 1 applies for any initial condition  $z(1)$ .

**Expected discrete-time dynamics.** The expected motion in (1) is described by

$$z(t) - z(t-1) = \frac{1}{t} (q_i(t) - z(t-1)). \quad (10)$$

where  $q_i \in \Delta(A_i)$  is the expectation of  $a$ . Assume that  $q_i$  is derived by (Q1) from a potential function satisfying (R1)–(R2). Let  $\varepsilon_i(t) := R_{i,\max}(t)$ . It is easily seen that, as for continuous-time dynamics,  $\varepsilon_i(t) \rightarrow 0$  as  $t \rightarrow +\infty$ , and if  $\varepsilon_i(1) > 0$ , then  $\varepsilon_i(t) > 0$  for all  $t$  and  $q_i \in BR_i^{\varepsilon_i(t)}(z_{-i})$ . It follows that (for a good choice of behavior when all regrets are initially nonpositive) Theorem 1 holds for the expected process (10).

This shows that the reason why the proof of Theorem 1 is easier in continuous-time is not continuous-time per se. It is that (8), as (10), is an expected version of (1).

## Appendix

### A.1 Proof of Theorem 1

Denote by  $\widehat{BR}_i^\varepsilon(x)$  the correspondence whose graph is the  $\varepsilon$ -neighborhood of the graph of  $BR_i$ :

$$\widehat{BR}_i^\varepsilon(x_{-i}) = \left\{ x_i \in \Delta(A_i) \left| \begin{array}{l} \exists (x_i^*, x_{-i}^*) \in \Delta(A_1) \times \Delta(A_2) \text{ s.t.} \\ x_i^* \in BR_i(x_{-i}^*), \text{ and } \|(x_i^*, x_{-i}^*) - (x_i, x_{-i})\|_\infty \leq \varepsilon \end{array} \right. \right\} \quad (11)$$

Let  $\widehat{BR}^\varepsilon(x) = \widehat{BR}_1^\varepsilon(x_2) \times \widehat{BR}_2^\varepsilon(x_1)$ . In words, action  $y_i$  is an  $\varepsilon$ -graph perturbed best reply to  $x_{-i}$  if there is an action  $\varepsilon$ -close to  $y_i$  which is an exact best-reply to an action  $\varepsilon$ -close to  $x_{-i}$ . This is the notion of perturbation used in the theory of perturbed differential inclusions (Benaïm et al., 2005, 2006). As illustrated by the following example, it is different from the notion of perturbation of payoffs in the  $\varepsilon$ -best reply correspondence,

i.e.  $BR^\varepsilon(x) = BR_1^\varepsilon(x_2) \times BR_2^\varepsilon(x_1)$  with

$$BR_i^\varepsilon(x_{-i}) = \left\{ \sigma_i \in \Delta(A_i) \mid u_i(\sigma_i, x_{-i}) \geq \max_{k \in A_i} u_i(k, x_{-i}) - \varepsilon \right\}, \quad i = 1, 2. \quad (12)$$

	$L$	$R$
$T$	1	0
$C$	0	1
$B$	$\frac{1}{2} - \eta$	$\frac{1}{2} - \eta$

Fig. 5

Consider a game where the payoffs of player 1 are given by Fig 5. Let  $\varepsilon \in (0, 1/2)$  and let  $x_2^\varepsilon = (\frac{1}{2} + \varepsilon)L + (\frac{1}{2} - \varepsilon)R$ . The pure action  $C$  is a  $2\varepsilon$ -best reply to  $x_2^\varepsilon$ . Using the sup norm, it is at distance 1 from pure action  $T$ , the unique exact best reply to  $x_2^\varepsilon$ . Nevertheless,  $C$  is an  $\varepsilon$ -graph perturbed best reply, because it is an exact best reply to  $x_2^0$ , which is  $\varepsilon$ -close (in sup norm) to  $x_2^\varepsilon$ . By contrast, for all  $\eta > 0$ , action  $B$  is an  $(\varepsilon + \eta)$ -best reply, but only a  $1$ -graph perturbed best reply to  $x_2^\varepsilon$ .

A discrete-time trajectory  $(x_1(t), x_2(t))_{t=1}^\infty$  on  $\Delta(A_1) \times \Delta(A_2)$  is a *payoff perturbed* fictitious play trajectory if there exists a positive sequence  $(\varepsilon_t)$  converging to zero such that (4) holds with  $q_i(t) \in A_i \cap BR_i^{\varepsilon_t}(x_{-i}(t-1))$  for all  $i = 1, 2$  and all  $t \in \mathbb{N}^*$ . It is a *graph perturbed* fictitious play trajectory if the same holds but replacing  $BR_i^{\varepsilon_t}$  with  $\widehat{BR}_i^{\varepsilon_t}$ . A trajectory  $(z(t))_{t=1}^\infty$  on  $\Delta(A)$  generates a *sequence of beliefs*  $(z_1(t), z_2(t))$  in  $\Delta(A_1) \times \Delta(A_2)$

The proof goes as follows. Lemma 2 shows that the sequence of beliefs generated by a no-regret dynamics is a payoff perturbed FP trajectory. Together with Lemma 3, this implies that it is a graph-perturbed FP trajectory (Lemma 4). It follows that the interpolated process of a no-regret dynamics trajectory is a perturbed solution of CFP (Lemma 5). The result then follows from Benaïm et al. (2005).

**Lemma 2.** *The sequence of beliefs of a solution of a no-regret dynamics in class  $\mathcal{R}$  is almost surely a payoff perturbed DFP trajectory.*

**Proof.** If  $R_{i,\max}(t) \leq 0$  for all  $t$ , then by Remark 2, player  $i$  plays an  $\varepsilon(t)$ -best reply for some  $\varepsilon(t)$  converging to zero. Otherwise,  $R_{i,\max}(t_0) > 0$  for some  $t_0 \in \mathbb{N}^*$ . Then for all times  $t \geq t_0$ ,  $R_{i,\max}(t) > 0$  (by Remark 1) and player  $i$  plays an  $R_{i,\max}(t)$ -best reply by Lemma 1 and condition (R3). Since  $R_{i,\max}(t) \rightarrow 0$  almost surely, the result follows. ■

**Lemma 3.** *Let  $X$  be a compact subset of  $\mathbb{R}^m$  and  $F$  a correspondence from  $X$  to itself. For any  $\delta \geq 0$ , let  $\hat{F}_\delta : X \rightrightarrows X$  denote the correspondence whose graph is the  $\delta$ -neighborhood*

of the graph of  $F$ :

$$\hat{F}_\delta(x) = \left\{ y \in X \mid \exists (x^*, y^*) \in X^2 \text{ s.t. } y^* \in F(x^*) \text{ and } \|(x^*, y^*) - (x, y)\|_\infty \leq \delta \right\}.$$

For any  $\alpha > 0$ , let  $G_\alpha$  be an u.s.c. correspondence from  $X$  to itself. Assume that for each  $x$  in  $X$ :

- (i)  $\alpha < \alpha' \Rightarrow G_\alpha(x) \subset G_{\alpha'}(x)$  (that is,  $(G_\alpha)_{\alpha>0}$  is increasing w.r.t. inclusion);
- (ii)  $\bigcap_{\alpha>0} G_\alpha(x) \subset F(x)$ .

Then for every  $\delta > 0$  there exists  $\alpha > 0$  such that for each  $x$  in  $X$ ,  $G_\alpha(x) \subset \hat{F}_\delta(x)$ .

**Proof.** By contradiction, assume that there exists  $\delta > 0$ , a decreasing sequence  $(\alpha_n)$  converging to zero, and sequences  $(x_n)$  and  $(y_n)$  of points in  $X$  such that  $y_n \in G_{\alpha_n}(x_n) \setminus \hat{F}_\delta(x_n)$  for all  $n$ . By compactness of  $X$ , we can assume that  $(x_n)$  and  $(y_n)$  converge respectively to  $x^*$  and  $y^*$ . Fix  $k \in \mathbb{N}$ . For all  $n \geq k$ ,  $y_n \in G_{\alpha_n}(x_n) \subset G_{\alpha_k}(x_n)$  by (i). Since  $G_{\alpha_k}$  is u.s.c., it follows that  $y^* \in G_{\alpha_k}(x^*)$ . Therefore, by (i) and (ii)

$$y^* \in \bigcap_{k \in \mathbb{N}} G_{\alpha_k}(x^*) = \bigcap_{\alpha > 0} G_\alpha(x^*) \subset F(x^*)$$

But for  $n$  large enough,  $\|(x^*, y^*) - (x_n, y_n)\|_\infty < \delta$ , hence  $y_n \in \hat{F}_\delta(x_n)$ , a contradiction. ■

Applied to the best-reply correspondence, Lemma 3 means that for any  $\delta > 0$ , an  $\varepsilon$ -perturbed best-reply is a  $\delta$ -graph perturbed best-reply, provided  $\varepsilon$  is small enough. Thus we have the next result.

**Lemma 4.** *Any payoff perturbed DFP trajectory is a graph perturbed DFP trajectory.*

**Proof.** Let  $\varepsilon_t \rightarrow 0$ . Let

$$\delta_t = \min \left\{ \delta \geq 0 \mid \forall i = 1, 2, \forall x \in \Delta(A_1) \times \Delta(A_2), BR_i^{\varepsilon_t}(x_{-i}) \subset \widehat{BR}_i^\delta(x_{-i}) \right\}.$$

Applying Lemma 3 with  $X = \Delta(A_1) \times \Delta(A_2)$ ,  $G_\varepsilon = BR^\varepsilon$  and  $F = BR$ , we obtain that  $\delta_t \rightarrow 0$ . The result follows. ■

Given a discrete-time trajectory  $x(n) = (x_1(n), x_2(n))$  on  $\Delta(A_1) \times \Delta(A_2)$ , with  $n \in \mathbb{N}^*$ , define its interpolated process  $x : [1, +\infty[ \rightarrow \Delta(A_1) \times \Delta(A_2)$  as follows. For all  $t \in [n, n+1)$  let  $tx(t) = nx(n) + (t-n)q(n)$ , where  $q_i(n) = (n+1)x_i(n+1) - nx_i(n)$ ,  $i = 1, 2$ . This is equivalent to

$$x_i(t) - x_i(n) = \frac{t-n}{t}(q_i(n) - x_i(t)), \quad i = 1, 2.$$

Hence for all  $t \in (n, n + 1)$  we have  $\|x(t) - x(n)\|_\infty \leq \frac{1}{n+1}$  and

$$\dot{x}(t) = \frac{1}{t}(q(n) - x(t)) \quad (13)$$

An absolutely continuous function  $x : [1, +\infty[ \rightarrow \Delta(A_1) \times \Delta(A_2)$  is a perturbed solution of CFP if there exists a vanishing function  $\varepsilon : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  such that for almost all  $t$ ,

$$\dot{x} \in \frac{1}{t} \left( \widehat{BR}^{\varepsilon(t)}(x) - x \right) \quad \text{where } x = x(t). \quad (14)$$

**Lemma 5.** *The interpolated process of a graph perturbed DFP trajectory is a perturbed solution of CFP.*

**Proof.** Consider a discrete time trajectory  $(x_1(n), x_2(n))_{n \in \mathbb{N}}$  such that

$$x_i(n) - x_i(n - 1) = \frac{1}{n} (q_i(n) - x_i(n)), \quad i = 1, 2,$$

with  $q_i(n) \in \widehat{BR}_i^{\varepsilon_n}(x_{-i}(n - 1))$  and  $\varepsilon_n \rightarrow 0$ . For all  $n$  and all  $t \in [n, n + 1)$ , let  $\varepsilon(t) = \varepsilon_n + 2/n$ . Obviously,  $\varepsilon(t) \rightarrow 0$  as  $t \rightarrow \infty$ . Moreover, for all  $t \in (n, n + 1)$ , the interpolated process satisfies  $\|x_{-i}(t) - x_{-i}(n - 1)\|_\infty \leq \frac{1}{n} + \frac{1}{n+1} < 2/n$ , so  $q_i(n) \in \widehat{BR}_i^{\varepsilon(t)}(x_{-i}(t))$ . Therefore (13) implies (14). ■

We can now prove Theorem 1. By Lemmata 2 and 4, the sequence of beliefs of a solution of a no-regret dynamics in class  $\mathcal{R}$  is almost surely a graph perturbed DFP trajectory. Hence, by Lemma 5, its interpolated process  $x(t)$  is a perturbed solution of CFP. This implies that  $x(e^t)$  is almost surely a perturbed solution of the best-reply dynamics, in the sense of Benaïm et al. (2005, Definition II). Theorem 1 now follows from Theorem 3.6 of Benaïm et al. (2005).<sup>18</sup>

**Remark 6.** Assume that at stage  $t$ , for each  $i = 1, 2$ , player  $i$  chooses a pure action according to a mixed action  $q_i(t)$  that depends on the previous history  $h(t - 1)$ . Do not assume conditions (R1)–(R3) and (Q1), but assume that there exists a vanishing sequence  $\varepsilon_t$  such that for any previous history  $h(t - 1)$ ,  $q_i(t) \in BR_i^{\varepsilon_t}(z_{-i}(t - 1))$  for all  $t > 1$  and all  $i = 1, 2$ . Then it follows from Lemma 3, the above proof and Benaïm et al. (2005, Proposition 1.4 and a variant of Proposition 1.3) that Theorem 1 applies. As is well known, this is the case for the exponential weight algorithm (Freund and Schapire, 1999;

---

<sup>18</sup>The definition of perturbed solution in Benaïm et al. (2005) is different from ours but equivalent.

Littlestone and Warmuth, 1994) that corresponds to

$$q_{i,k}(t) := \frac{\exp \beta_t u_i(k, z_{-i})}{\sum_{s \in A_i} \exp(\beta_t u_i(s, z_{-i}))}$$

with  $z_{-i} = z_{-i}(t-1)$ ,  $\beta_t \rightarrow_{t \rightarrow +\infty} +\infty$ , and  $\beta_t < t^\alpha$  for some  $\alpha \in ]0, 1[$  to ensure that this is a no-regret dynamics (see, e.g., Benaïm and Faure, 2011). The above assumptions are not (or not trivially) satisfied by no-regret dynamics in class  $\mathcal{R}$ . Indeed, the rate at which the maximal regret vanishes, hence the value  $\varepsilon_t$  such that  $q_i(t) \in BR_i^{\varepsilon_t}(z_{-i}(t))$ , may depend on the trajectory.

## A.2 ICT sets when all solutions converge to Nash equilibria

The fact that all solutions of the best-reply dynamics converge to the set of Nash equilibria does not guarantee that ICT sets contain only Nash equilibria. We provide counterexamples below.

**Example 1** (single-population dynamics). Consider the following symmetric  $3 \times 3$  game:

	<i>A</i>	<i>B</i>	<i>C</i>
<i>A</i>	0	0	0
<i>B</i>	0	0	0
<i>C</i>	-1	0	0

Denote a mixed action by  $x = (x_A, x_B, x_C)$ . Then  $(x, x)$  is a Nash equilibrium if and only if  $x_A = 0$  or  $x_C = 0$ . It is easily seen that all solutions of the best-reply dynamics converge to the set of symmetric Nash equilibria. However, the whole state space is ICT. Indeed, any mixed action  $x$  can be connected to any other mixed action  $y$  as follows: starting from  $x$ , follow a solution pointing towards the edge  $x_C = 0$ , then jump on this edge and follow a solution pointing towards the pure strategy  $B$ ; once close to  $B$ , jump on the edge  $x_A = 0$ , and follow a solution pointing towards  $C$ ; once close to  $C$ , make a small jump to reach a point from which a solution points toward  $y$ ; follow this solution and if needed (i.e. if  $y_C = 0$ ), make one more jump to reach  $y$ .

This example is also valid for the replicator dynamics and any payoff monotone dynamics in the sense of, e.g., Hofbauer and Weibull (1996). The only difference is that traveling from  $A$  to  $B$  and from  $B$  to  $C$  cannot be done by following solutions of the dynamics but only through long sequences of jumps. Note also that in an inward cycling Rock-Paper-Scissors game (see e.g., Hofbauer and Sigmund, 1998, or Weibull, 1995), all



solutions of the replicator dynamics converge to one of the four rest points but the whole boundary of the state space is ICT (for the replicator dynamics).

**Example 2** ( $n$ -population dynamics). Similarly, in the bimatrix version of example 1, all solutions of the two-population best-reply dynamics converge to the set of Nash equilibria but the whole state space is ICT. Again, this is true for all payoff monotone dynamics. Similar examples can be given for  $n$ -population dynamics for any  $n \geq 1$ .

At least for the best-reply dynamics,  $2 \times 2$  examples can also be given. Consider, for instance, the  $2 \times 2$  game:

	$L$	$R$
$T$	$0, 0$	$0, 0$
$B$	$0, 0$	$-1, 0$

Denote mixed actions of players 1 and 2 by  $x = (x_T, x_B)$  and  $y = (y_L, y_R)$ , respectively. The set of Nash equilibria is the union of the edges  $x_B = 0$  and  $y_R = 0$  and all solutions of the two-population best-reply dynamics converge to this set. However, the whole triangle  $x_T + y_L \geq 1$  is ICT.

In these examples, a direct analysis shows that all solutions of no-regret dynamics in class  $\mathcal{R}$  converge to the set of Nash equilibria. Thus we do not know whether, in general, convergence of all solutions of CFP to the set of Nash equilibria entails convergence of no-regret dynamics. The point is that this is not guaranteed by Theorem 1.

### A.3 Proof of Proposition 3

We need some notation. For  $z \in \Delta(A)$  and  $a \in A$ , let  $z_a$  denote the probability of  $a$  under the correlated action  $z$ . Let  $\mathcal{U}_\gamma(H_B)$  be the neighborhood of  $H_B$  in which the total weight on action profiles outside of  $B$  and the potential of each player are below  $\gamma$ :

$$\mathcal{U}_\gamma(H_B) = \left\{ z \in \Delta(A) \left| \begin{array}{l} \sum_{a \notin B} z_a < \gamma, \text{ and} \\ P_i(R_i(z)) < \gamma, \quad i = 1, 2, \end{array} \right. \right\}$$

where  $R_i(z)$  is the regret vector of player  $i$ :  $R_i(z) = (u_i(s, z_{-i}) - u_i(z))_{s \in A_i}$ .

Let  $B = B_1 \times B_2$  be a curb set. Let

$$\delta_B = \min_{i=1,2} \min_{z_{-i} \in \Delta(B_{-i})} \left\{ \max_{s \in A_i} u_i(s, z_{-i}) - \max_{k \in A_i \setminus B_i} u_i(k, z_{-i}) \right\}$$

and note that  $\delta_B > 0$ , since  $B$  is curb. Now, consider a no-regret dynamics in  $\mathcal{R}$  defined

by potentials  $P_i$ ,  $i = 1, 2$ , with trajectory  $(z(t))_{t \geq 1}$ . Let  $\rho_i(\gamma)$  be the smallest number such that for all  $z \in \Delta(A)$

$$P_i(z) \leq \gamma \implies R_{i,\max}(z) \leq \rho_i(\gamma),$$

and let  $\rho(\gamma) = \max\{\rho_1(\gamma), \rho_2(\gamma)\}$ . Let  $\gamma_B$  be the solution of

$$(2\bar{U} + \delta_B)\gamma_B + \rho(\gamma_B) - \delta_B = 0 \tag{15}$$

where  $\bar{U} = \max_{i=1,2} \max_{a \in A} |u_i(a)|$  is a payoff bound. Since  $\rho(\gamma)$  is weakly increasing in  $\gamma$  and  $\rho(0) = 0$ , there exists a unique solution  $\gamma_B$  of (15) and  $\gamma_B > 0$ .

Consider the following event  $\mathcal{E}_t$ :

$$P_i(R_i(t+n)) < \gamma_B \text{ for each } i = 1, 2 \text{ and all } n \in \mathbb{N}^*. \tag{\mathcal{E}_t}$$

The statement of Proposition 3 is immediate by the following claims and Corollary 1.

**Claim 1.** *If  $z(t) \in \mathcal{U}_{\gamma_B}(H_B)$  and event  $\mathcal{E}_t$  holds, then  $a(t+n) \in B$  for all  $n \in \mathbb{N}^*$ .*

**Claim 2.** *For every  $\pi \in (0, 1]$  and every  $\gamma \in (0, \gamma_B)$  there exists  $t_0$  such that for every  $t \geq t_0$ , if  $z(t) \in \mathcal{U}_\gamma(B)$ , then event  $\mathcal{E}_t$  holds with probability at least  $1 - \pi$ .*

For the proof of Claim 1 we need the following lemma.

**Lemma 6.** *For any  $t$ , if  $z(t) \in \mathcal{U}_{\gamma_B}(H_B)$ , then  $a(t+1) \in B$ .*

**Proof.** Let  $z \in \Delta(A)$ . If  $z$  is close enough to  $B$ , then  $\max_{s \in A_i} u_i(s, z_{-i}) - \max_{k \in A_i \setminus B_i} u_i(k, z_{-i}) > \delta_B/2$  for all  $i = 1, 2$ . If  $z$  is close enough to  $H$ , then  $\max_{s \in A_i} u_i(s, z_{-i}) - u_i(z) < \delta_B/2$ . Thus in the neighborhood of  $H_B$ ,  $\max_{k \in A_i \setminus B_i} u_i(k, z_{-i}) < u_i(z)$  hence  $R_{i,k}(z) < 0$  for all  $k \in A_i \setminus B_i$ . In particular, this holds if  $z \in U_{\gamma_B}(H_B)$  (we omit the proof: easy but lengthy). It follows that if  $z(t) \in U_{\gamma_B}(H_B)$ , then by conditions (R3) and (Q2'),  $a_i(t+1) \in B_i$  for each  $i = 1, 2$ . ■

**Proof of Claim 1.** Suppose that  $\mathcal{E}_t$  holds and let  $z(t) \in \mathcal{U}_{\gamma_B}(B)$ . Then  $a(t+1) \in B$  by Lemma 6. We proceed by induction. Assume  $a(t+1), \dots, a(t+n) \in B$  for some  $n \in \mathbb{N}^*$ . Since  $z(t) \in \mathcal{U}_{\gamma_B}(B)$ ,

$$\sum_{a \in A \setminus B} z_a(t+n) < \sum_{a \in A \setminus B} z_a(t) < \gamma_B.$$

Together with  $\mathcal{E}_t$ , this implies that  $z(t+n) \in \mathcal{U}_{\gamma_B}(B)$ . Consequently, by Lemma 6,  $a(t+n+1) \in B$ . ■

The proof of Claim 2 builds up on the proof of Theorem 2.1 of Hart and Mas-Colell (2001). It is different though, since we need to find the convergence rate of the maximal regret *conditional* on a given initial history (in particular, on those where the past average play is close to a curb set), which Hart and Mas-Colell (2001) do not provide. So our result cannot be directly derived from their proof.

For the proof of Claim 2 we need the following lemmata.

**Lemma 7.** *Let  $x_1, x_2, \dots$  be a sequence of real random variables with  $E[x_n | x_{n-1}, \dots, x_1] = 0$  and  $Var[x_n] \leq \bar{\sigma}^2$  for all  $n$ . Then for every  $\pi > 0$  and every  $m = 1, 2, \dots$*

$$\Pr \left[ \max_{n > m} \left| \frac{1}{n} \sum_{k=m+1}^n x_k \right| \geq \frac{\bar{\sigma}}{\sqrt{m\pi}} \right] \leq \pi.$$

**Proof.** Hájek-Rényi inequality (e.g., Bullen, 1998) implies

$$\Pr \left[ \max_{m < k \leq n} c_k |x_{m+1} + \dots + x_k| \geq \varepsilon \right] \leq \frac{1}{\varepsilon^2} \sum_{k=m+1}^n c_k^2 Var[x_k].$$

Using  $c_k = 1/k$  and  $Var[x_k] \leq \bar{\sigma}^2$ , the right-hand side can be bounded as follows,

$$\sum_{k=m+1}^n c_k^2 Var[x_k] \leq \bar{\sigma}^2 \sum_{k=1}^{n-m} \frac{1}{(m+k)^2} \leq \bar{\sigma}^2 \frac{1}{m}.$$

Taking the limit  $n \rightarrow \infty$  yields

$$\Pr \left[ \max_{k > m} \frac{1}{k} |x_{m+1} + \dots + x_k| \geq \varepsilon \right] \leq \frac{\bar{\sigma}^2}{m\varepsilon^2}.$$

The result is immediate by substitution  $\pi = \bar{\sigma}^2/(m\varepsilon^2)$ . ■

Define  $\xi(1) = P_i(R_i(1))$  and for all  $t = 2, 3, \dots$

$$\xi(t) = tP_i(R_i(t)) - (t-1)P_i(R_i(t-1)). \quad (16)$$

**Lemma 8.**  *$\xi(t)$  is uniformly bounded and  $E[\xi(t) | h(t-1)] \leq C/t$  holds for some constant  $C$  uniformly for all  $t$ .*

**Proof.** Let  $x_0 = R_i(t-1)$  and  $x = R_i(t)$ . Note that  $R_i(t) = \frac{t-1}{t}R_i(t-1) + \frac{1}{t}r_i$ , where  $r_i = [u_i(k, a_{-i}(t)) - u_i(a(t))]_{k \in A_i}$ . Hence

$$x - x_0 = \frac{1}{t}(r_i - x_0). \quad (17)$$

The regret for an action is bounded by  $2\bar{U}$  and the difference between two regret terms by  $4\bar{U}$ . Thus, in sup norm,  $\|r_i - x_0\| \leq 4\bar{U}$  and  $\|x - x_0\| \leq 4\bar{U}/t$ .

Since  $P_i$  is  $C^2$ , there exist constants  $c$ ,  $c'$  and  $c''$  such that if  $\|y\| \leq 4\bar{U}$ ,

$$\|P_i(y)\| \leq c, \quad \|\nabla P_i(y) \cdot y\| \leq c'\|y\|, \quad \text{and} \quad \|y \cdot \nabla^2 P_i(y)y\| \leq c''\|y\|^2.$$

Moreover,  $\xi(t) = P_i(x_0) + t(P_i(x) - P_i(x_0))$  hence  $|\xi(t)| \leq c + tc'\|x - x_0\| \leq c + 4\bar{U}c'$ . Thus  $\xi(t)$  is uniformly bounded.

We now show that  $E[\xi(t)|h(t-1)] \leq C/t$  for  $C = 8\bar{U}^2c''$ . By definition of  $c''$  and Taylor-Lagrange theorem,

$$P_i(x) \leq P_i(x_0) + \nabla P_i(x_0) \cdot (x - x_0) + \frac{1}{2}c''\|x - x_0\|^2.$$

Using (17) we get:

$$P_i(x) \leq \frac{t-1}{t}P_i(x_0) + \frac{1}{t}(P_i(x_0) - \nabla P_i(x_0) \cdot x_0) + \frac{1}{t}\nabla P_i(x_0) \cdot r_i(t) + \frac{C}{t^2}.$$

Since  $P_i$  is convex and  $P_i(0) = 0$ , we have:

$$P_i(x_0) - \nabla P_i(x_0) \cdot x_0 = P_i(x_0) + \nabla P_i(x_0) \cdot (0 - x_0) \leq P_i(0) = 0.$$

Therefore

$$P_i(x) \leq \frac{t-1}{t}P_i(x_0) + \frac{1}{t}\nabla P_i(x_0) \cdot r_i + \frac{C}{t^2},$$

so that

$$\xi(t) = tP_i(x) - (t-1)P_i(x_0) \leq \nabla P_i(x_0) \cdot r_i + \frac{C}{t}.$$

To prove that  $E[\xi(t)|h(t-1)] \leq C/t$ , it suffices to show that  $E[\nabla P_i(x_0) \cdot r_i|h(t-1)] = 0$ .

To see this, note that  $E[r_{i,k}|h(t-1)] = u_i(k, a_{-i}(t)) - u_i(q_i(t), a_{-i}(t))$  hence

$$E[q_i(t) \cdot r_i|h(t-1)] = q_i(t) \cdot E[r_i|h(t-1)] = \sum_{k \in A_i} q_{i,k}[u_i(k, a_{-i}) - u_i(q_i, a_{-i})] = 0.$$

Since  $q_i(t)$  is proportional to  $\nabla P_i(x_0)$ , the result follows. ■

**Proof of Claim 2.** By (16) we have

$$\begin{aligned} P_i(R_i(t+n)) &= \frac{1}{t+n} \sum_{s=t+1}^{t+n} \xi(s) + \frac{t}{t+n} P_i(R_i(t)) \\ &= \frac{1}{t+n} \sum_{s=t+1}^{t+n} \zeta(s) + \frac{1}{t+n} \sum_{s=t+1}^{t+n} E[\xi(s)|h(s-1)] + \frac{t}{t+n} P_i(R_i(t)), \end{aligned}$$

where  $\zeta(s) = \xi(s) - E[\xi(s)|h(s-1)]$ . As we have assumed  $z(t) \in \mathcal{U}_\gamma(B)$ , we have

$$\frac{t}{t+n} P_i(R_i(t)) < \frac{t}{t+n} \gamma \leq \gamma.$$

Next, by Lemma 8,

$$\frac{1}{t+n} \sum_{s=t+1}^{t+n} E[\xi(s)|h(s-1)] \leq \frac{1}{t+n} \sum_{s=t+1}^{t+n} \frac{C}{s} \leq C \frac{\ln(t+n) - \ln t}{t+n}.$$

Maximizing  $\frac{\ln(t+n) - \ln t}{t+n}$  among all  $n > 0$  yields  $\frac{\ln(t+n) - \ln t}{t+n} \leq (te)^{-1}$ , hence

$$\frac{1}{t+n} \sum_{s=t+1}^{t+n} E[\xi(s)|h(s-1)] \leq \frac{C}{te}.$$

Let  $\bar{\sigma}^2$  be a bound on  $\text{Var}[\zeta(s)]$  for all  $s$  (this bound exists, since by Lemma 8 variables  $\xi(s)$  are uniformly bounded). Applying Lemma 7, we obtain that for every  $\pi > 0$  and every  $t$ ,

$$\Pr \left[ \max_{n \in \mathbb{N}} \left| \frac{1}{t+n} \sum_{s=t+1}^{t+n} \zeta(s) \right| \geq \frac{\bar{\sigma}}{\sqrt{t\pi/2}} \right] \leq \frac{\pi}{2}.$$

Hence with probability at least  $1 - \pi/2$  the following holds for all  $n \in \mathbb{N}^*$ ,

$$P_i(R_i(t+n)) < \frac{\sqrt{2}\bar{\sigma}}{\sqrt{t\pi}} + \frac{C}{te} + \gamma.$$

and it holds for both  $i = 1, 2$  simultaneously with probability at least  $(1 - \pi/2)^2 \geq 1 - \pi$ . Choosing  $t_0 = t_0(\pi, \gamma)$  so large that  $\frac{\sqrt{2}\bar{\sigma}}{\sqrt{t_0\pi}} + \frac{C}{t_0e} \leq \gamma_B - \gamma$ , we obtain that event  $\mathcal{E}_i$ :

$$P_i(R_i(t+n)) < \gamma_B \quad \text{for each } i = 1, 2 \text{ and all } n \in \mathbb{N}^*$$

occurs with probability at least  $(1 - \pi/2)^2 \geq 1 - \pi$ . ■

## References

- Aubin, J.-P. and A. Celina (1984). *Differential Inclusions*. Springer.
- Balkenborg, D., J. Hofbauer, and C. Kuzmics (2011). Refined best reply correspondence and dynamics. Mimeo.
- Basu, K. and J. W. Weibull (1991). Strategy subsets closed under rational behavior. *Economic Letters* 36, 141–146.
- Benaïm, M. and M. Faure (2011). Consistency of vanishing smooth fictitious play. Mimeo.
- Benaïm, M., J. Hofbauer, and S. Sorin (2005). Stochastic approximations and differential inclusions. *SIAM Journal on Control and Optimization* 44, 328–348.
- Benaïm, M., J. Hofbauer, and S. Sorin (2006). Stochastic approximations and differential inclusions. Part II: Applications. *Mathematics of Operations Research* 31, 673–695.
- Berger, U. (2005). Fictitious play in  $2 \times n$  games. *Journal of Economic Theory* 120, 139–154.
- Berger, U. (2007). Two more classes of games with the continuous-time fictitious play property. *Games and Economic Behavior* 60, 247–261.
- Blum, A., E. Even-Dar, and K. Ligett (2006). Routing without regret: on convergence to nash equilibria of regret-minimizing algorithms in routing games. In *Proceedings of the 25th Annual ACM Symposium on Principles of Distributed Computing*, pp. 45–52.
- Brown, G. (1951). Iterative solutions of games by fictitious play. In T. Koopmans (Ed.), *Activity Analysis of Production and Allocation*, Volume 13 of *Cowles Commission Monograph*, pp. 374–376. New York: Wiley.
- Bullen, P. (1998). *A Dictionary of Inequalities*. Addison Wesley Longman, Harlow.
- Cesa-Bianchi, N. and G. Lugosi (2003). Potential-based algorithms in on-line prediction and game theory. *Machine Learning* 51, 239–261.
- Cesa-Bianchi, N. and G. Lugosi (2006). *Prediction, Learning, and Games*. Cambridge University Press.
- Chen, Y. and J. W. Vaughan (2010). A new understanding of prediction markets via no-regret learning. In *Proceedings of the 11th ACM Conference on Electronic Commerce*, pp. 189–198. mimeo.
- Clemen, R. T. and R. L. Winkler (2007). Aggregating probability distributions. In W. Edwards, R. Miles, and D. von Winterfeldt (Eds.), *Advances in Decision Analysis*, pp. 154–176. Cambridge University Press.

- DeMarzo, P., I. Kremer, and Y. Mansour (2006). Online trading algorithms and robust option pricing. In *Proceedings of the 38th Annual ACM Symposium on Theory of Computing*, pp. 477–486.
- Foster, D. and R. Vohra (1993). A randomization rule for selecting forecasts. *Operations Research* 41, 704–709.
- Foster, D. and R. Vohra (1999). Regret in the online decision problem. *Games and Economic Behavior* 29, 7–35.
- Freund, Y. and R. Schapire (1999). Adaptive game playing using multiplicative weights. *Games and Economic Behavior* 29, 79–103.
- Fudenberg, D. and D. Levine (1995). Consistency and cautious fictitious play. *Journal of Economic Dynamics and Control* 19, 1065–1089.
- Gaunersdorfer, A. and J. Hofbauer (1995). Fictitious play, Shapley polygons, and the replicator equation. *Games and Economic Behavior* 11, 279–303.
- Gilboa, I. and A. Matsui (1991). Social stability and equilibrium. *Econometrica* 59, 859–867.
- Hannan, J. (1957). Approximation to Bayes risk in repeated play. In M. Dresher, A. W. Tucker, and P. Wolfe (Eds.), *Contributions to the Theory of Games, Vol. III*, Annals of Mathematics Studies 39, pp. 97–139. Princeton University Press.
- Hart, S. and A. Mas-Colell (2000). A simple adaptive procedure leading to correlated equilibrium. *Econometrica* 68, 1127–1150.
- Hart, S. and A. Mas-Colell (2001). A general class of adaptive procedures. *Journal of Economic Theory* 98, 26–54.
- Hart, S. and A. Mas-Colell (2003). Continuous-time regret-based dynamics. *Games and Economic Behavior* 45, 375–394.
- Hofbauer, J. and W. H. Sandholm (2009). Stable games and their dynamics. *Journal of Economic Theory* 144, 1665–1693.
- Hofbauer, J. and K. Sigmund (1998). *Evolutionary Games and Population Dynamics*. Cambridge University Press.
- Hofbauer, J. and S. Sorin (2006). Best response dynamics for continuous zero-sum games. *Discrete and Continuous Dynamical Systems B* 6, 215–224.
- Hofbauer, J., S. Sorin, and Y. Viossat (2009). Time average replicator and best reply dynamics. *Mathematics of Operations Research* 34, 263–269.
- Hofbauer, J. and J. W. Weibull (1996). Evolutionary selection against dominated strategies. *Journal of Economic Theory* 71, 558–573.

- Irani, S. and A. Karlin (1996). On-line computation. In D. Hochbaum (Ed.), *Approximation Algorithms for NP-Hard Problems*, pp. 521–564. Boston: PWS-Kent.
- Krishna, V. and T. Sjöström (1998). On the convergence of fictitious play. *Mathematics of Operations Research* 23, 479–511.
- Larrick, R. P. and J. B. Soll (2006). Intuitions about combining opinions: misappreciation of the averaging principle. *Management Science* 52, 111–127.
- Lehrer, E. (2003). A wide range no-regret theorem. *Games and Economic Behavior* 42, 101–115.
- Mansour, Y. (2010). Regret minimization and job scheduling. In *Proceedings of the 36th Conference on Current Trends in Theory and Practice of Computer Science*, pp. 71–76. Springer.
- Matsui, A. (1992). Best response dynamics and socially stable strategies. *Journal of Economic Theory* 57, 343–362.
- Monderer, D., D. Samet, and A. Sela (1997). Belief affirming in learning processes. *Journal of Economic Theory* 73, 438–452.
- Moulin, H. and J. P. Vial (1978). Strategically zero-sum games: the class of games whose completely mixed equilibria cannot be improved upon. *International Journal of Game Theory* 7, 201–221.
- Selten, R. (1995). An axiomatic theory of a risk dominance measure for bipolar games with linear incentives. *Games and Economic Behavior* 8, 213–263.
- Shapley, L. S. (1964). Some topics in two person games. In M. Dresher, L. S. Shapley, and A. W. Tucker (Eds.), *Advances in Game Theory*, pp. 1–28. Princeton University Press.
- Sparrow, C., S. van Strien, and C. Harris (2008). Fictitious play in  $3 \times 3$  games: The transition between periodic and chaotic behaviour. *Games and Economic Behavior* 63, 259–291.
- Timmerman, A. (2006). Forecast combinations. In G. Elliott, C. W. Granger, and A. Timmermann (Eds.), *Handbook of Economic Forecasting*. Elsevier.
- Tirole, J. (1988). *The Theory of Industrial Organization*. MIT Press.
- Weibull, J. W. (1995). *Evolutionary Game Theory*. Cambridge, MA: Cambridge University Press.
- Yanovskaya, E. (1968). Equilibrium points in polymatrix games (in Russian). *Litovskii Matematicheskii Sbornik* 8, 381–384.
- Young, H. P. (1993). The evolution of conventions. *Econometrica* 61, 57–84.
- Young, H. P. (2004). *Strategic Learning and Its Limits*. Oxford University Press.