



HAL
open science

Estimation of a cumulative distribution function under interval censoring "case 1" via warped wavelets

Christophe Chesneau, Thomas Willer

► **To cite this version:**

Christophe Chesneau, Thomas Willer. Estimation of a cumulative distribution function under interval censoring "case 1" via warped wavelets. 2012. hal-00715260v1

HAL Id: hal-00715260

<https://hal.science/hal-00715260v1>

Preprint submitted on 6 Jul 2012 (v1), last revised 1 Dec 2013 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Estimation of a cumulative distribution function under interval censoring “case 1” via warped wavelets

Christophe Chesneau¹ and Thomas Willer²

Abstract: The estimation of an unknown cumulative distribution function in the interval censoring “case 1” model from dependent sequences is considered. We construct a new adaptive estimator based on a warped wavelet basis and a hard thresholding rule. Under mild assumptions on the parameters of the model, considering the \mathbb{L}_2 risk and the weighted Besov balls, we prove that the estimator attains a sharp rate of convergence. We also investigate its practical performances thanks to simulation experiments.

Key words and phrases: Adaptive estimation, Strongly mixing, Interval censoring, Warped wavelets, Hard thresholding.

AMS 2000 Subject Classifications: 62G05, 62G20.

1 Introduction

The mathematical context of the interval censoring “case 1” model can be described as follows: let $(\delta_i, U_i)_{i \in \mathbb{Z}}$ be a stationary process where, for any $i \in \mathbb{Z}$,

$$\delta_i = \mathbf{1}_{\{X_i \leq U_i\}},$$

$\mathbf{1}_{\mathcal{A}}$ is the indicator function on any random event \mathcal{A} , $(X_i)_{i \in \mathbb{Z}}$ and $(U_i)_{i \in \mathbb{Z}}$ are independent, and $(X_i)_{i \in \mathbb{Z}}$ is a stationary process with common unknown cumulative distribution F . We assume that U_1 admits a density, denoted by g , and we denote by G its cumulative distribution function. Our goal is to estimate F under mild assumptions on g from n observations $(\delta_1, U_1), \dots, (\delta_n, U_n)$ of $(\delta_i, U_i)_{i \in \mathbb{Z}}$. This model has applications in Demography and Biology. See e.g. [9] and [13], and the references there in.

For recent statistical results, we refer to [19] and [1]. In particular, considering the independent case, [1] have constructed adaptive penalized minimum contrast estimators built on trigonometric, polynomial or wavelet spaces. Using the \mathbb{L}_2 risk over Besov balls, under some boundedness assumptions on g , [1, Corollary 3.1] proves that it attains the standard rate of convergence “ $n^{-2s/(2s+1)}$ ” where s characterizes the smoothness of F .

However, the independence assumption on $(\delta_i, U_i)_{i \in \mathbb{Z}}$ is often stringent in applications. In this study, we investigate the adaptive estimation of F in a dependent setting (including the independent one). The so-called *strong mixing case* is considered. Examples and applications of this kind of dependence can be found in [3] and [11].

Assuming that g is known but with no boundedness assumptions on it, we develop a new adaptive estimator based on a warped wavelet basis and a hard thresholding rule. The features of this basis consist of a standard wavelet basis and of the definition of G

¹Université de Caen, LMNO, Campus II, Science 3, 14032, Caen, France

²Université de Provence, LATP, CMI, Technopôle Château Gombert 39, rue F. Joliot Curie, 13453, Marseille, France

related to the model. This enables us to give a significant stability to our thresholding algorithm. Such a technique has been already used with success in the framework of nonparametric regression with random design by [14]. Recent works on warped wavelet basis in nonparametric statistics can be found in [4], [5], [2] and [18].

Considering the \mathbb{L}_2 risk over weighted Besov balls, we prove that our estimator attains the rate of convergence “ $(\ln n/n)^{2s/(2s+1)}$ ”, where s characterizes the smoothness of F . This rate of convergence corresponds to the one attained in the *i.i.d.* case (see [1]) up to an extra logarithmic term. Finally, we explore the numerical performances of the estimator.

The rest of the paper is organized as follows. Section 2 introduces notations and assumptions on the model. In Section 3, we describe warped wavelet bases on $[0, 1]$ and weighted Besov balls. Our adaptive wavelet estimator is defined in Section 4. Theoretical and practical results are presented Section 5. The proofs are postponed to Section 6.

2 Notations and assumptions

2.1 Assumptions on the dependence structure of the process

For any $m \in \mathbb{Z}$, we define the m -th strongly mixing coefficient of $(\delta_i, U_i)_{i \in \mathbb{Z}}$ by

$$a_m = \sup_{(A,B) \in \mathcal{F}_{-\infty,0}^{(\delta,U)} \times \mathcal{F}_{m,\infty}^{(\delta,U)}} |\mathbb{P}(A \cap B) - \mathbb{P}(A)\mathbb{P}(B)|, \quad (2.1)$$

where $\mathcal{F}_{-\infty,0}^{(\delta,U)}$ is the σ -algebra generated by the pairs of random variables $\dots, (\delta_{-1}, U_{-1}), (\delta_0, U_0)$ and $\mathcal{F}_{m,\infty}^{(\delta,U)}$ is the σ -algebra generated by the random variables $(\delta_m, U_m), (\delta_{m+1}, U_{m+1}), \dots$

We consider the exponentially strongly mixing case: there exist two known constants $\gamma > 0$ and $c > 0$ such that, for any integer $m \geq 1$,

$$a_m \leq \gamma \exp(-cm). \quad (2.2)$$

This assumption is not very restrictive; some examples of processes satisfying such conditions can be found in e.g. [20], [11], [17] and [3].

2.2 Assumptions on the densities

For the sake of simplicity, we suppose that all the considered random variables take their values in $[0, 1]$.

In the main part of the study, we assume that g is known. The unknown case will only be explored in the simulation study in subsection 6.2.

We suppose that, for any interval $[a, b] \subseteq [0, 1]$, there exists a constant $C > 0$ such that

$$\left(\frac{1}{b-a} \int_a^b g(x)^2 dx \right)^{1/2} \leq C \frac{1}{b-a} \int_a^b g(x) dx. \quad (2.3)$$

This “reverse Hölder inequality” is related to the Muckenhoupt weights theory. It includes a wide variety of densities, non-necessarily bounded from above and/or below. For instance, $g(x) = (u+1)x^u$, $x \in [0, 1]$ and $u \in (0, 1)$ satisfies (2.3). Further details can be found in [14, subsection 4.1].

For any $m \in \{1, \dots, n\}$, let $\mathbf{f}_{(\delta_0, U_0, \delta_m, U_m)}$ be the density of $(\delta_0, U_0, \delta_m, U_m)$, $\mathbf{f}_{(\delta_0, U_0)}$ the density of (δ_0, U_0) and, for any $(y, x, y_*, x_*) \in [0, 1]^4$,

$$h_m(y, x, y_*, x_*) = \mathbf{f}_{(\delta_0, U_0, \delta_m, U_m)}(y, x, y_*, x_*) - \mathbf{f}_{(\delta_0, U_0)}(y, x)\mathbf{f}_{(\delta_0, U_0)}(y_*, x_*). \quad (2.4)$$

We suppose that there exists a known constant $C > 0$ such that

$$\sup_{m \in \{1, \dots, n\}} \sup_{(y, x, y_*, x_*) \in [0, 1]^4} \frac{1}{g(x)g(x_*)} |h_m(y, x, y_*, x_*)| \leq C. \quad (2.5)$$

Note that, in the independent case, we have $h_m(y, x, y_*, x_*) = 0$ and (2.5) is satisfied. Moreover, functions g satisfying (2.3) and (2.5) are not necessarily bounded from below and above. Hence our conditions are less restrictive than [1, Assumption A1].

3 Warped wavelets and weighted Besov balls

Let N be a positive integer. We consider an orthonormal wavelet basis generated by dilations and translations of a "father" Daubechies-type wavelet ϕ and a "mother" Daubechies-type wavelet ψ of the family $db2N$. In particular, ϕ and ψ have compact supports, $\int \phi(x)dx = 1$ and, for any $r \in \{0, \dots, N-1\}$, $\int x^r \psi(x)dx = 0$.

Set

$$\phi_{j,k}(x) = 2^{j/2}\phi(2^j x - k), \quad \psi_{j,k}(x) = 2^{j/2}\psi(2^j x - k).$$

Suppose that (2.3) holds and recall that

$$G(x) = \mathbb{P}(U_1 \leq x) = \int_0^x g(u)du, \quad x \in \mathbb{R}.$$

Then, with an appropriate treatment at the boundaries, there exists an integer τ satisfying $2^\tau \geq 2N$ such that, for any integer $j_* \geq \tau$, any $h \in \mathbb{L}_2([0, 1])$ can be expanded into a warped wavelet series as

$$h(x) = \sum_{k=0}^{2^{j_*}-1} \alpha_{j_*,k} \phi_{j_*,k}(G(x)) + \sum_{j=j_*}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k} \psi_{j,k}(G(x)), \quad x \in [0, 1],$$

where

$$\alpha_{j,k} = \int_0^1 h(G^{-1}(x))\phi_{j,k}(x)dx, \quad \beta_{j,k} = \int_0^1 h(G^{-1}(x))\psi_{j,k}(x)dx. \quad (3.1)$$

See [14, subsection 3.3].

Let $M > 0$ and $s > 0$. We say that a function h in $\mathbb{L}_2([0, 1])$ belongs to the weighted Besov ball $B_{s,\infty}^w(M)$ if there exists a constant $M > 0$ such that the associated wavelet coefficients (3.1) satisfy, for any integer $j \geq \tau$,

$$\sum_{k=0}^{2^j-1} \beta_{j,k}^2 w_{j,k} \leq M2^{-j(2s+1)},$$

where

$$w_{j,k} = \int_{k/2^j}^{(k+1)/2^j} \frac{1}{g(G^{-1}(x))} dx. \quad (3.2)$$

In this expression, s is a smoothness parameter. Details concerning the warped wavelets and the analytic definition of weighted Besov balls can be found in [14, Section 7]. For the standard wavelet basis on $[0, 1]$, see e.g. [16] and [6].

4 Estimators

For any integer $j \geq \tau$ and any $k \in \{0, \dots, 2^j - 1\}$, we estimate the unknown warped wavelet coefficients of F i.e. $\alpha_{j,k} = \int_0^1 F(G^{-1}(x))\phi_{j,k}(x)dx$ and $\beta_{j,k} = \int_0^1 F(G^{-1}(x))\psi_{j,k}(x)dx$ by respectively

$$\hat{\alpha}_{j,k} = \frac{1}{n} \sum_{i=1}^n \delta_i \phi_{j,k}(G(U_i)), \quad \hat{\beta}_{j,k} = \frac{1}{n} \sum_{i=1}^n \delta_i \psi_{j,k}(G(U_i)). \quad (4.1)$$

Some of their statistical properties are investigated in Propositions 6.1, 6.2 and 6.3 below.

We estimate F by the following hard thresholding estimator \hat{F} :

$$\hat{F}(x) = \sum_{k=0}^{2^\tau-1} \hat{\alpha}_{\tau,k} \phi_{\tau,k}(G(x)) + \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^j-1} \hat{\beta}_{j,k} \mathbf{1}_{\{|\hat{\beta}_{j,k}| \geq \kappa \rho_n\}} \psi_{j,k}(G(x)), \quad (4.2)$$

where $x \in [0, 1]$, $\hat{\alpha}_{\tau,k}$ and $\hat{\beta}_{j,k}$ are defined by (4.1), j_1 is the integer satisfying

$$\frac{1}{2} \frac{n}{(\ln n)^3} < 2^{j_1} \leq \frac{n}{(\ln n)^3}, \quad (4.3)$$

κ is a large enough constant (the one in Proposition 6.3 below) and ρ_n denotes the “universal threshold”, i.e.

$$\rho_n = \sqrt{\frac{\ln n}{n}}. \quad (4.4)$$

Naturally, for $x < 0$, we put $\hat{F}(x) = 0$ and, for $x > 1$, $\hat{F}(x) = 1$.

Note that \hat{F} is adaptive: its construction does not depend on the smoothness of F .

The general idea in the construction of \hat{F} is to apply a term-by-term selection on the unknown wavelet coefficients of F : only the most significant are kept. The reason is that only these few coefficients contain the main characteristics of F . For the construction of hard thresholding wavelet estimators in the standard nonparametric models, see e.g. [10], [8] and [12].

5 Performances of \hat{F}

5.1 Theoretical results

Theorem 5.1 *Suppose that the assumptions of Section 2 hold. Let \hat{F} be (4.2). Suppose that $F \in B_{s,\infty}^w(M)$ with $s > 0$. Then there exists a constant $C > 0$ such that*

$$\mathbb{E} \left(\int_0^1 \left(\hat{F}(x) - F(x) \right)^2 dx \right) \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}.$$




Figure 1: Four target cumulative distribution functions

The proof of Theorem 5.1 uses a suitable decomposition of the \mathbb{L}_2 risk including some geometrical properties of the warped wavelet basis and the statistical properties of the wavelet coefficients estimators presented in Propositions 6.1, 6.2 and 6.3.

Theorem 5.1 shows that, under mild assumptions on the dependence of the observations and on g , \hat{F} attains a rate of convergence close to the one for the *i.i.d.* case i.e. $n^{-2s/(2s+1)}$. The difference is the “negligible” logarithmic term $(\ln n)^{2s/(2s+1)}$.

Let us recall that, if we restrict our study to the independent case, contrary to [1, Assumption A1], Theorem 5.1 holds without boundedness assumptions on g .

5.2 Practical results

This section is devoted to the numerical performances of \hat{F} .

5.2.1 Target functions and data

We consider four target functions F which correspond to the following formula with values $q = 0.2, 1, 5, 50$:

$$F_q(t) = \begin{cases} 2^{q-1}t^q, & t \in [0, 0.5], \\ 1 - 2^{q-1}(1-t)^q, & t \in [0.5, 1]. \end{cases} \quad (5.1)$$

These four functions are plotted in Figure 1, and their wavelets detail coefficients are plotted in Figure 2. One can remark that F_q has high detail coefficients near the edges of the interval $[0, 1]$ when q is small, and at the middle of the interval $[0, 1]$ when q is large (see Figure 2).

An example of the data on which the estimators are based is given in Figures 4 and 5 (for the cdf F_{50} and for a uniform distribution of the U_i s). Figure 4 represents the couples (U_i, δ_i) . Figure 5 gives the detail wavelet coefficients of the sequence $(\delta_{\sigma_U(i)})_{i \in \{1, \dots, n\}}$, where σ_U is the following permutation: $\sigma_U(i)$ is the index j such that U_j is the i^{st} largest element of the sequence U . These coefficients are approximations of the ones given in equation (4.1), as their construction consists in replacing G by the empirical pdf \hat{G}_n in their expressions. Then the estimation consists in trying to select the coefficients corresponding to the signal F , and to put to zero those corresponding to the “noise”. We consider first the case of a known uniform density of the U_i s, then the case of a known varying density, and lastly the case of an unknown density.




Figure 2: Detail wavelet coefficients of each target function: the first row relates to $F_{0.2}$, the second to F_1 , the third to F_5 and the fourth to F_{50} .




Figure 3: Data with (U_i) s (uniformly distributed) on axis 1, and their corresponding (δ_i) s on axis 2, for the F_{50} function.




Figure 4: Detail wavelet coefficients of the data (δ_i) generated thanks to the F_{50} cdf in the uniform case.

5.2.2 Uniform density of the design

First let us look at the performances of the main estimator presented in Section 4, in the case of a uniform distribution of the U_i s. The calibration of the threshold $\kappa\rho_n$ and of the cutoff level j_1 are important practical issues. The values given by the theory are not useful in practice. Indeed the thresholds are defined up to some intricate "large enough constant". Moreover the cutoff level j_1 defined in (4.3) is too small in practice, as it is even lower than the minimal coarsest level value. Thus we try $\kappa = \sqrt{2}$ (universal thresholding) and $j_1 = \log_2(n) - 1$ (the maximal possible level), hoping the high resolution "noise" is filtered thanks to thresholding.

We give an example of the performances for F_{50} . Figure 5 represents the true coefficients above, and the noisy ones below, along with the thresholds (horizontal bars). Figure 6 represents the target function (dashed line) and the estimator (solid line). One can see that the estimator fails to recover F properly in the middle of the interval, as it leaves a lot of noise unfiltered in the high resolution scales. On the other hand the estimator behaves properly outside of the middle of the interval.

The main problem with this estimator is that high resolution scales contain huge noise coefficients that one cannot filter with thresholding, except if we put huge thresholds as well, but then low scale information would be lost. Thus one may ask oneself if the results could be improved by using other calibrations of κ and j_1 . For this purpose, one can compute some "oracle" estimator in the sense that it uses the threshold and the cutoff values which minimize the mean square error among all estimators of the type of Section 4, except with κ and j_1 left free. Of course this estimator is completely inaccessible in practice. In the case of the F_{50} distribution, simulations show that this oracle estimator is obtained approximately with a threshold $t = 0.5$ and a cutoff level $j_1 = 3$. This means that only some of the coarsest resolution detail coefficients should be kept. As can be seen in Figure 7, this oracle estimator is better than the previous one.

We briefly look at the performances of the estimator for the three other distributions F_q . The high resolution noise problem is far more present than for F_{50} (first row of Figure 8). The oracle strategy consists in cutting all the details. The functions obtained this way are plotted in the second row of Figure 8.

5.2.3 Non-uniform density of the design

We now investigate the performances of the estimators for non-constant densities of the U_i s. We consider the four densities plotted in Figure 9, named Bump1, Bump2, Pit1 and Pit2. They correspond to the following formulas (up to normalization constants):

- Bump1 $g(x) = \exp(-(100 * (x - 0.5)^2)) + 1$,
- Pit1 $g(x) = -\exp(-(100 * (x - 0.5)^2)) + 1.05$,
- Bump2 $g(x) = \exp(-(100 * x^2)) + \exp(-(100 * (x - 1)^2)) + 1$,
- Pit2 $g(x) = -\exp(-(100 * x^2)) - \exp(-(100 * (x - 1)^2)) + 1.05$.

The effects of non uniformity seem to be multiple for the performances of the estimator. Let us focus on the two cdf F_5 and F_{50} , and the Bump1 and Pit1 density. The Bump1 density implies a surplus (resp. a lack) of observations in the middle of the interval, where most important wavelet coefficients are located. On the one hand, low scale




Figure 5: Theoretical thresholds (horizontal bars), wavelet detail coefficients of F_{50} (first row of graphics), wavelet detail coefficients of the (δ_i) s (second row of graphics).




Figure 6: Target function (dashed line) and the estimator of section 4 (solid line) for F_{50} in the uniform case for U_i .




Figure 7: Target function (dashed line) and the oracle estimator (solid line) for F_{50} in the uniform case.




Figure 8: Target functions (dashed line), estimator of section 4 (solid line top), oracle estimator (solid line bottom) for $F_{0.2}$, F_1 , F_5 (from left to right) in the uniform case.




Figure 9: Densities of the (U_i) s.




Figure 10: Realizations of the estimator of Section 4 (top) and oracle estimator (bottom) for respectively (from left to right): F_5 and Bump1 density, F_5 and Pit1 density, F_{50} and Bump1 density, F_{50} and Pit1 density.

coefficients are better estimated thanks to a bump than to a pit (see the second row of Figure 10). For the Pit1 density, the lack of observation in the middle of the interval causes the estimator to fail locating the jump of the target function. But on the other hand, the bump attributes numerous data near the median of F , which creates much variability in the observations. Thus many artifacts appear in the middle of the interval for the bump density, while no artifact remains for the pit density for the cdf F_{50} (see the first row of Figure 10).

Moreover, the two other functions F_1 or $F_{0.2}$ remain very hard to estimate by the techniques of Section 4. We consider the Bump2 (resp Pit2) density, which attributes more (resp less) observations to the beginning and to the end of the interval (where important detail coefficients lie for these two target functions). Then one really cannot distinguish the true function in the first row of Figure 11. The oracle estimators are plotted in the second row. Once again simulations show that the oracle strategy consists in putting all the detail coefficients to zero. The oracle estimator behaves similarly whether we use the Pit2 or the Bump2 density.

5.2.4 Unknown density of the design

When g is unknown, an intuitive adaptive estimator of F is (4.2) but, instead of G , we consider its empirical version:

$$\hat{G}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbf{1}_{\{U_i \leq x\}}.$$

This plug-in method yields the hard thresholding estimator

$$\hat{F}^*(x) = \sum_{k=0}^{2^\tau-1} \hat{\alpha}_{\tau,k}^* \phi_{\tau,k}(\hat{G}_n(x)) + \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^j-1} \hat{\beta}_{j,k}^* \mathbf{1}_{\{|\hat{\beta}_{j,k}| \geq \kappa \rho_n\}} \psi_{j,k}(\hat{G}_n(x)),$$




Figure 11: Realizations of the estimator of Section 4 (top) and oracle estimator (bottom) for respectively: $F_{0,2}$ and Bump2 density, $F_{0,2}$ and Pit2 density, F_1 and Bump2 density, F_1 and Pit2 density.

$x \in [0, 1]$, where

$$\hat{\alpha}_{j,k}^* = \frac{1}{n} \sum_{i=1}^n \delta_i \phi_{j,k}(\hat{G}_n(U_i)), \quad \hat{\beta}_{j,k}^* = \frac{1}{n} \sum_{i=1}^n \delta_i \psi_{j,k}(\hat{G}_n(U_i)).$$

Like previously, we look at the performances of the estimator for the F_{50} and F_5 target functions, with three different densities g (Bump1, Uniform, Pit1). The results are given in Figure 12. We obtain similar performances to the ones in the case of known g , the main general remarks remain true here.

6 Proofs

In this section, we suppose that the assumptions of Section 2 hold. Moreover, C denotes any constant that does not depend on j , k and n . Its value may change from one term to another and may depend on ϕ or ψ .

6.1 Auxiliary results on (4.1)

Proposition 6.1 *Suppose that the assumptions of Section 2 hold. For any integer $j \geq \tau$ such that $2^j \leq n$ and any $k \in \{0, \dots, 2^j - 1\}$, let $\alpha_{j,k} = \int_0^1 F(G^{-1}(x)) \phi_{j,k}(x) dx$ and $\hat{\alpha}_{j,k}$ be (4.1). Then $\hat{\alpha}_{j,k}$ is an unbiased estimator for $\alpha_{j,k}$ and there exists a constant $C > 0$ such that*

$$\text{Var}(\hat{\alpha}_{j,k}) \leq C \frac{1}{n},$$

Mention that Proposition 6.1 holds with $\hat{\beta}_{j,k}$ (4.1) instead of $\hat{\alpha}_{j,k}$ and $\beta_{j,k} = \int_0^1 F(G^{-1}(x)) \psi_{j,k}(x) dx$ instead of $\alpha_{j,k}$.




Figure 12: Target function (dashed line), estimator of section 4 (solid line) for the cdf F_{50} (first row) and F_5 (second row) with unknown g equal to Bump1, Uniform, Pit1 (from left to right).

Proof of Proposition 6.1. We have

$$\begin{aligned}
\mathbb{E}(\hat{\alpha}_{j,k}) &= \mathbb{E}(\delta_1 \phi_{j,k}(G(U_1))) = \mathbb{E}(\mathbf{1}_{\{X_i \leq U_i\}} \phi_{j,k}(G(U_1))) \\
&= \int_0^1 \int_0^u f(x) \phi_{j,k}(G(u)) g(u) dx du = \int_0^1 F(u) \phi_{j,k}(G(u)) g(u) du \\
&= \int_0^1 F(G^{-1}(u)) \phi_{j,k}(u) du = \alpha_{j,k}.
\end{aligned}$$

An elementary covariance inequality yields

$$\text{Var}(\hat{\alpha}_{j,k}) = \frac{1}{n^2} \sum_{v=1}^n \sum_{\ell=1}^n \text{Cov}(\delta_v \phi_{j,k}(G(U_v)), \delta_\ell \phi_{j,k}(G(U_\ell))) \leq \mathbf{S} + \mathbf{T}, \quad (6.1)$$

where

$$\mathbf{S} = \frac{1}{n} \text{Var}(\delta_1 \phi_{j,k}(G(U_1)))$$

and

$$\mathbf{T} = \frac{2}{n^2} \left| \sum_{v=2}^n \sum_{\ell=1}^{v-1} \text{Cov}(\delta_v \phi_{j,k}(G(U_v)), \delta_\ell \phi_{j,k}(G(U_\ell))) \right|.$$

Since $\delta_1 \leq 1$, we have

$$\begin{aligned}
\mathbf{S} &\leq \frac{1}{n} \mathbb{E}((\delta_1 \phi_{j,k}(G(U_1)))^2) \leq \frac{1}{n} \mathbb{E}((\phi_{j,k}(G(U_1)))^2) \\
&= \frac{1}{n} \int_0^1 (\phi_{j,k}(G(x)))^2 g(x) dx = \frac{1}{n} \int_0^1 (\phi_{j,k}(x))^2 dx = \frac{1}{n}.
\end{aligned} \quad (6.2)$$

It follows from the stationarity of $(\delta_i, U_i)_{i \in \mathbb{Z}}$ and $2^j \leq n$ that

$$\begin{aligned} \mathbf{T} &= \frac{2}{n^2} \left| \sum_{m=1}^n (n-m) \text{Cov}(\delta_0 \phi_{j,k}(G(U_0)), \delta_m \phi_{j,k}(G(U_m))) \right| \\ &\leq \frac{2}{n} \sum_{m=1}^n |\text{Cov}(\delta_0 \phi_{j,k}(G(U_0)), \delta_m \phi_{j,k}(G(U_m)))| = \mathbf{T}_1 + \mathbf{T}_2, \end{aligned} \quad (6.3)$$

where

$$\mathbf{T}_1 = \frac{2}{n} \sum_{m=1}^{2^j-1} |\text{Cov}(\delta_0 \phi_{j,k}(G(U_0)), \delta_m \phi_{j,k}(G(U_m)))|$$

and

$$\mathbf{T}_2 = \frac{2}{n} \sum_{m=2^j}^n |\text{Cov}(\delta_0 \phi_{j,k}(G(U_0)), \delta_m \phi_{j,k}(G(U_m)))|.$$

Upper bound for \mathbf{T}_1 . Using (2.4), (2.5) and the change of variables $y = 2^j x - k$, we obtain

$$\begin{aligned} &|\text{Cov}(\delta_0 \phi_{j,k}(G(U_0)), \delta_m \phi_{j,k}(G(U_m)))| \\ &= \left| \int_0^1 \int_0^1 \int_0^1 \int_0^1 h_m(y, x, y_*, x_*) (y \phi_{j,k}(G(x)) y_* \phi_{j,k}(G(x_*))) dy dx dy_* dx_* \right| \\ &\leq \int_0^1 \int_0^1 \int_0^1 \int_0^1 |h_m(y, x, y_*, x_*)| |y \phi_{j,k}(G(x)) y_* \phi_{j,k}(G(x_*))| dy dx dy_* dx_* \\ &\leq C \left(\int_0^1 y dy \right)^2 \left(\int_0^1 |\phi_{j,k}(G(x))| g(x) dx \right)^2 \leq C \left(\int_0^1 |\phi_{j,k}(x)| dx \right)^2 \leq C 2^{-j}. \end{aligned}$$

Therefore

$$\mathbf{T}_1 \leq C \frac{1}{n} 2^{-j} 2^j = C \frac{1}{n}. \quad (6.4)$$

Upper bound for \mathbf{T}_2 . Applying the Davydov inequality for strongly mixing processes (see [7]), for any $q \in (0, 1)$, we have

$$\begin{aligned} &|\text{Cov}(\delta_0 \phi_{j,k}(G(U_0)), \delta_m \phi_{j,k}(G(U_m)))| \\ &\leq C a_m^q \left(\mathbb{E} \left(|\delta_0 \phi_{j,k}(G(U_0))|^{2/(1-q)} \right) \right)^{1-q} \\ &\leq C a_m^q \left(\sup_{(x,y) \in [0,1]^2} |y \phi_{j,k}(G(x))| \right)^{2q} \left(\mathbb{E} \left((\delta_0 \phi_{j,k}(G(U_0)))^2 \right) \right)^{1-q}. \end{aligned}$$

Observe that

$$\sup_{(x,y) \in [0,1]^2} |y \phi_{j,k}(G(x))| \leq \sup_{x \in [0,1]} |\phi_{j,k}(x)| \leq C 2^{j/2}$$

and, by (6.2),

$$\mathbb{E} \left((\delta_0 \phi_{j,k}(G(U_0)))^2 \right) \leq 1.$$

Therefore

$$|\text{Cov}(\delta_0 \phi_{j,k}(G(U_0)), \delta_m \phi_{j,k}(G(U_m)))| \leq C 2^{jq} a_m^q.$$

Hence

$$\mathbf{T}_2 \leq C \frac{1}{n} 2^{qj} \sum_{m=2^j}^n a_m^q \leq C \frac{1}{n} \sum_{m=2^j}^n m^q a_m^q \leq C \frac{1}{n}. \quad (6.5)$$

It follows from (6.3), (6.4) (6.5) that

$$\mathbf{T} \leq C \frac{1}{n}. \quad (6.6)$$

Combining (6.1), (6.2) and (6.6), we obtain

$$\text{Var}(\hat{\alpha}_{j,k}) \leq C \frac{1}{n}.$$

The proof of Proposition 6.1 is complete. □

Proposition 6.2 *Suppose that the assumptions of Section 2 hold. For any integer $j \geq \tau$ such that $2^j \leq n$ and any $k \in \{0, \dots, 2^j - 1\}$, let $\beta_{j,k} = \int_0^1 F(G^{-1}(x))\psi_{j,k}(x)dx$ and $\hat{\beta}_{j,k}$ be (4.1). Then there exists a constant $C > 0$ such that*

$$\mathbb{E} \left(\left(\hat{\beta}_{j,k} - \beta_{j,k} \right)^4 \right) \leq C 2^j \frac{1}{n}.$$

Proof of Proposition 6.2. Observe that

$$\begin{aligned} |\hat{\alpha}_{j,k} - \alpha_{j,k}| &\leq |\hat{\alpha}_{j,k}| + |\alpha_{j,k}| \leq \sup_{(x,y) \in [0,1]^2} |y\phi_{j,k}(G(x))| + C \\ &\leq \sup_{x \in [0,1]} |\phi_{j,k}(x)| + C \leq C 2^{j/2}. \end{aligned} \quad (6.7)$$

Using (6.7) and Proposition 6.1, we obtain

$$\mathbb{E} \left((\hat{\alpha}_{j,k} - \alpha_{j,k})^4 \right) \leq C 2^j \mathbb{E} \left((\hat{\alpha}_{j,k} - \alpha_{j,k})^2 \right) \leq C 2^j \frac{1}{n}.$$

The proof of Proposition 6.2 is complete. □

Proposition 6.3 *Suppose that the assumptions of Section 2 hold. For any $j \in \{\tau, \dots, j_1\}$ and any $k \in \{0, \dots, 2^j - 1\}$, let $\beta_{j,k} = \int_0^1 F(G^{-1}(x))\psi_{j,k}(x)dx$, $\hat{\beta}_{j,k}$ be (4.1) and ρ_n be (4.4). Then there exist two constants, $\kappa > 0$ and $C > 0$, such that*

$$\mathbb{P} \left(|\hat{\beta}_{j,k} - \beta_{j,k}| \geq \kappa \rho_n / 2 \right) \leq C \frac{1}{n^4}.$$

Proof of Proposition 6.3. We shall use the Bernstein inequality for exponentially strongly mixing process presented in Lemma 6.1 below. The proof can be found [15].

Lemma 6.1 ([15]) *Let $\gamma > 0$, $c > 0$ and $(Z_i)_{i \in \mathbb{Z}}$ be a stationary process with the m -th strongly mixing coefficient (2.1). Let n be a positive integer, $h : \mathbb{R} \rightarrow \mathbb{R}$ be a measurable function and, for any $i \in \mathbb{Z}$, $V_i = h(Z_i)$. We assume that $\mathbb{E}(V_1) = 0$ and there exists a constant $M > 0$ satisfying $|V_1| \leq M$. Then, for any $m \in \{1, \dots, n\}$ and any $\lambda > 0$, we have*

$$\mathbb{P} \left(\left| \frac{1}{n} \sum_{i=1}^n V_i \right| \geq \lambda \right) \leq 4 \exp \left(-\frac{\lambda^2 n}{16(D_m/m + \lambda M m/3)} \right) + 32 \frac{M}{\lambda} n a_m,$$

where $D_m = \max_{l \in \{1, \dots, 2m\}} \text{Var} \left(\sum_{i=1}^l V_i \right)$.

For any $i \in \{1, \dots, n\}$, set

$$V_i = \delta_i \psi_{j,k}(G(U_i)) - \beta_{j,k}.$$

Then

$$\mathbb{P} \left(|\hat{\beta}_{j,k} - \beta_{j,k}| \geq \kappa \rho_n / 2 \right) = \mathbb{P} \left(\left| \frac{1}{n} \sum_{i=1}^n V_i \right| \geq \kappa \rho_n / 2 \right), \quad (6.8)$$

V_1, \dots, V_n are identically distributed, depend on the stationary strongly mixing process $(\delta_i, U_i)_{i \in \mathbb{Z}}$ satisfying (2.2), Propositions 6.1 and 6.2 give

$$\mathbb{E}(V_1) = 0, \quad \mathbb{E}(V_1^2) \leq \mathbb{E}((\delta_1 \phi_{j,k}(G(U_1)))^2) \leq 1,$$

using similar arguments to (6.7), $|V_1| \leq C 2^{j/2} \leq C 2^{j_1/2} \leq C(n/(\ln n)^3)^{1/2}$, and, proceeding as in Proposition 6.1 with l instead of n , we have

$$\max_{l \in \{1, \dots, 2m\}} \text{Var} \left(\sum_{i=1}^l V_i \right) \leq C \max_{l \in \{1, \dots, 2m\}} l \leq Cm.$$

It follows from Lemma 6.1 applied with these V_1, \dots, V_n , $\lambda = \kappa C \rho_n$, $\rho_n = (\ln n/n)^{1/2}$, $m = u \ln n$ with $u > 0$ (chosen later), $M = C(n/(\ln n)^3)^{1/2}$ and (2.2) that

$$\begin{aligned} & \mathbb{P} \left(|\hat{\beta}_{j,k} - \beta_{j,k}| \geq \kappa \rho_n / 2 \right) \\ & \leq C \left(\exp \left(-C \frac{\kappa^2 \rho_n^2 n}{1 + \kappa \rho_n m M} \right) + \frac{n^{1/2}}{\rho_n (\ln n)^{3/2}} n \exp(-cm) \right) \\ & \leq C \left(\exp \left(-C \frac{\kappa^2 \ln n}{1 + \kappa (\ln n/n)^{1/2} u \ln n (n/(\ln n)^3)^{1/2}} \right) + n \exp(-cu \ln n) \right) \\ & \leq C \left(n^{-C \kappa^2 / (1 + \kappa u)} + n^{1-cu} \right). \end{aligned}$$

Taking κ and u large enough, we obtain

$$\mathbb{P} \left(|\hat{\beta}_{j,k} - \beta_{j,k}| \geq \kappa \rho_n / 2 \right) \leq C \frac{1}{n^4}.$$

This ends the proof of Proposition 6.3.

□

6.2 Proof of Theorem 5.1

Proof of Theorem 5.1. Since $F \in \mathbb{L}_2([0, 1])$, we can write

$$F(x) = \sum_{k=0}^{2^\tau-1} \alpha_{\tau,k} \phi_{\tau,k}(G(x)) + \sum_{j=\tau}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k} \psi_{j,k}(G(x)), \quad x \in [0, 1],$$

where $\alpha_{\tau,k} = \int_0^1 F(G^{-1}(x)) \phi_{\tau,k}(x) dx$ and $\beta_{j,k} = \int_0^1 F(G^{-1}(x)) \psi_{j,k}(x) dx$.

We have, for any $x \in [0, 1]$,

$$\begin{aligned} & \hat{F}(x) - F(x) \\ &= \sum_{k=0}^{2^\tau-1} (\hat{\alpha}_{\tau,k} - \alpha_{\tau,k}) \phi_{\tau,k}(G(x)) + \sum_{j=\tau}^{j_1} \sum_{k=0}^{2^j-1} \left(\hat{\beta}_{j,k} \mathbf{1}_{\{|\hat{\beta}_{j,k}| \geq \kappa \rho_n\}} - \beta_{j,k} \right) \psi_{j,k}(G(x)) \\ & - \sum_{j=j_1+1}^{\infty} \sum_{k=0}^{2^j-1} \beta_{j,k} \psi_{j,k}(G(x)). \end{aligned}$$

We now need the following lemma which is an immediate consequence of [14, Lemma 2] and the Minkowski inequality.

Lemma 6.2 ([14]) *Suppose that (2.3) holds. Then, for any sequences $(u_{j,k}) \in \ell_2(\mathbb{N}^2)$ and any integers j_0 and j_1 such that $j_1 > j_0 \geq \tau$, there exists a constant $C > 0$ such that*

$$\int_0^1 \left(\sum_{j=j_0}^{j_1} \sum_{k=0}^{2^j-1} u_{j,k} \psi_{j,k}(G(x)) \right)^2 dx \leq C \left(\sum_{j=j_0}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} u_{j,k}^2 w_{j,k} \right)^{1/2} \right)^2,$$

where $w_{j,k}$ is defined by (3.2).

It follows from Lemma 6.2 and the Cauchy-Schwarz inequality that

$$\mathbb{E} \left(\int_0^1 \left(\hat{F}(x) - F(x) \right)^2 dx \right) \leq C(\mathbf{F} + \mathbf{G} + \mathbf{H}), \quad (6.9)$$

where

$$\begin{aligned} \mathbf{F} &= 2^\tau \sum_{k=0}^{2^\tau-1} \mathbb{E} \left((\hat{\alpha}_{\tau,k} - \alpha_{\tau,k})^2 \right) w_{\tau,k}, \\ \mathbf{G} &= \left(\sum_{j=\tau}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \mathbb{E} \left(\left(\hat{\beta}_{j,k} \mathbf{1}_{\{|\hat{\beta}_{j,k}| \geq \kappa \rho_n\}} - \beta_{j,k} \right)^2 \right) w_{j,k} \right)^{1/2} \right)^2 \end{aligned}$$

and

$$\mathbf{H} = \left(\sum_{j=j_1+1}^{\infty} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \beta_{j,k}^2 w_{j,k} \right)^{1/2} \right)^2.$$

Using Proposition 6.1 and $\sum_{k=0}^{2^\tau-1} w_{\tau,k} = 1$, we obtain

$$\mathbf{F} \leq C \frac{1}{n} \sum_{k=0}^{2^\tau-1} w_{\tau,k} \leq C \frac{1}{n} \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}. \quad (6.10)$$

Since $F \in B_{s,\infty}^w(M)$, we have

$$\begin{aligned} \mathbf{H} &\leq C \left(\sum_{j=j_1+1}^{\infty} 2^{-js} \right)^2 \leq C 2^{-2j_1 s} \leq C \left(\frac{(\ln n)^3}{n} \right)^{2s} \\ &\leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}. \end{aligned} \quad (6.11)$$

Let us now bound the term \mathbf{G} . Observe that

$$\mathbf{G} \leq C(\mathbf{G}_1 + \mathbf{G}_2 + \mathbf{G}_3 + \mathbf{G}_4), \quad (6.12)$$

where

$$\mathbf{G}_1 = \left(\sum_{j=\tau}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \mathbb{E} \left(\left(\hat{\beta}_{j,k} - \beta_{j,k} \right)^2 \mathbf{1}_{\{|\hat{\beta}_{j,k}| \geq \kappa \rho_n\}} \mathbf{1}_{\{|\beta_{j,k}| < \kappa \rho_n / 2\}} \right) w_{j,k} \right)^{1/2} \right)^2,$$

$$\mathbf{G}_2 = \left(\sum_{j=\tau}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \mathbb{E} \left(\left(\hat{\beta}_{j,k} - \beta_{j,k} \right)^2 \mathbf{1}_{\{|\hat{\beta}_{j,k}| \geq \kappa \rho_n\}} \mathbf{1}_{\{|\beta_{j,k}| \geq \kappa \rho_n / 2\}} \right) w_{j,k} \right)^{1/2} \right)^2,$$

$$\mathbf{G}_3 = \left(\sum_{j=\tau}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \mathbb{E} \left(\beta_{j,k}^2 \mathbf{1}_{\{|\hat{\beta}_{j,k}| < \kappa \rho_n\}} \mathbf{1}_{\{|\beta_{j,k}| \geq 2\kappa \rho_n\}} \right) w_{j,k} \right)^{1/2} \right)^2$$

and

$$\mathbf{G}_4 = \left(\sum_{j=\tau}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \mathbb{E} \left(\beta_{j,k}^2 \mathbf{1}_{\{|\hat{\beta}_{j,k}| < \kappa \rho_n\}} \mathbf{1}_{\{|\beta_{j,k}| < 2\kappa \rho_n\}} \right) w_{j,k} \right)^{1/2} \right)^2.$$

Upper bounds for $\mathbf{G}_1 + \mathbf{G}_3$. Note that

$$\left\{ |\hat{\beta}_{j,k}| < \kappa \rho_n, |\beta_{j,k}| \geq 2\kappa \rho_n \right\} \subseteq \left\{ |\hat{\beta}_{j,k} - \beta_{j,k}| > \kappa \rho_n / 2 \right\},$$

$$\left\{ |\hat{\beta}_{j,k}| \geq \kappa \rho_n, |\beta_{j,k}| < \kappa \rho_n / 2 \right\} \subseteq \left\{ |\hat{\beta}_{j,k} - \beta_{j,k}| > \kappa \rho_n / 2 \right\}$$

and

$$\left\{ |\hat{\beta}_{j,k}| < \kappa \rho_n, |\beta_{j,k}| \geq 2\kappa \rho_n \right\} \subseteq \left\{ |\beta_{j,k}| \leq 2|\hat{\beta}_{j,k} - \beta_{j,k}| \right\}.$$

So

$$\mathbf{G}_1 + \mathbf{G}_3 \leq C \left(\sum_{j=\tau}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \mathbb{E} \left(\left(\hat{\beta}_{j,k} - \beta_{j,k} \right)^2 \mathbf{1}_{\{|\hat{\beta}_{j,k} - \beta_{j,k}| > \kappa \rho_n / 2\}} \right) w_{j,k} \right)^{1/2} \right)^2.$$

It follows from the Cauchy-Schwarz inequality, Proposition 6.2, Proposition 6.3 and $2^j \leq 2^{j_1} \leq n$ that

$$\begin{aligned} & \mathbb{E} \left(\left(\hat{\beta}_{j,k} - \beta_{j,k} \right)^2 \mathbf{1}_{\{|\hat{\beta}_{j,k} - \beta_{j,k}| > \kappa \rho_n / 2\}} \right) \\ & \leq \left(\mathbb{E} \left(\left(\hat{\beta}_{j,k} - \beta_{j,k} \right)^4 \right) \right)^{1/2} \left(\mathbb{P} \left(|\hat{\beta}_{j,k} - \beta_{j,k}| > \kappa \rho_n / 2 \right) \right)^{1/2} \\ & \leq C \left(2^j \frac{1}{n} \right)^{1/2} \left(\frac{1}{n^4} \right)^{1/2} \leq C \frac{1}{n^2}. \end{aligned}$$

Since $\sum_{k=0}^{2^j-1} w_{j,k} = 1$, we have

$$\begin{aligned} \mathbf{G}_1 + \mathbf{G}_3 & \leq C \frac{1}{n^2} \left(\sum_{j=\tau}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} w_{j,k} \right)^{1/2} \right)^2 = C \frac{1}{n^2} \left(\sum_{j=\tau}^{j_1} 2^{j/2} \right)^2 \\ & \leq C \frac{1}{n^2} 2^{j_1} \leq C \frac{1}{n} \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}. \end{aligned} \quad (6.13)$$

Upper bound for \mathbf{G}_2 . Using again Proposition 6.2, we obtain

$$\mathbb{E} \left(\left(\hat{\beta}_{j,k} - \beta_{j,k} \right)^2 \right) \leq C \frac{1}{n} \leq C \frac{\ln n}{n}.$$

Hence

$$\mathbf{G}_2 \leq C \frac{\ln n}{n} \left(\sum_{j=\tau}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \mathbf{1}_{\{|\beta_{j,k}| > \kappa \rho_n / 2\}} w_{j,k} \right)^{1/2} \right)^2.$$

Let j_2 be the integer defined by

$$\frac{1}{2} \left(\frac{n}{\ln n} \right)^{1/(2s+1)} < 2^{j_2} \leq \left(\frac{n}{\ln n} \right)^{1/(2s+1)}. \quad (6.14)$$

We have

$$\mathbf{G}_2 \leq C(\mathbf{G}_{2,1} + \mathbf{G}_{2,2}),$$

where

$$\mathbf{G}_{2,1} = \frac{\ln n}{n} \left(\sum_{j=\tau}^{j_2} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \mathbf{1}_{\{|\beta_{j,k}| > \kappa \rho_n / 2\}} w_{j,k} \right)^{1/2} \right)^2$$

and

$$\mathbf{G}_{2,2} = \frac{\ln n}{n} \left(\sum_{j=j_2+1}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \mathbf{1}_{\{|\beta_{j,k}| > \kappa \rho_n / 2\}} w_{j,k} \right)^{1/2} \right)^2.$$

Using $\mathbf{1}_{\{|\beta_{j,k}| > \kappa \rho_n / 2\}} \leq 1$ and $\sum_{k=0}^{2^j-1} w_{j,k} = 1$,

$$\begin{aligned} \mathbf{G}_{2,1} &\leq C \frac{\ln n}{n} \left(\sum_{j=\tau}^{j_2} 2^{j/2} \left(\sum_{k=0}^{2^j-1} w_{j,k} \right)^{1/2} \right)^2 = C \frac{\ln n}{n} \left(\sum_{j=\tau}^{j_2} 2^{j/2} \right)^2 \\ &\leq C \frac{\ln n}{n} 2^{j_2} \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)} \end{aligned}$$

and, since $F \in B_{s,\infty}^w(M)$,

$$\begin{aligned} \mathbf{G}_{2,2} &\leq C \frac{\ln n}{n \rho_n^2} \left(\sum_{j=j_2+1}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \beta_{j,k}^2 w_{j,k} \right)^{1/2} \right)^2 \leq C \left(\sum_{j=j_2+1}^{j_1} 2^{-js} \right)^2 \\ &\leq C 2^{-2j_2 s} \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}. \end{aligned}$$

So

$$\mathbf{G}_2 \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}. \quad (6.15)$$

Upper bound for \mathbf{G}_4 . We have

$$\mathbf{G}_4 \leq \left(\sum_{j=\tau}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \beta_{j,k}^2 \mathbf{1}_{\{|\beta_{j,k}| < 2\kappa \rho_n\}} w_{j,k} \right)^{1/2} \right)^2.$$

Let j_2 be the integer (6.14). Then

$$\mathbf{G}_4 \leq C(\mathbf{G}_{4,1} + \mathbf{G}_{4,2}),$$

where

$$\mathbf{G}_{4,1} = \left(\sum_{j=\tau}^{j_2} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \beta_{j,k}^2 \mathbf{1}_{\{|\beta_{j,k}| < 2\kappa \rho_n\}} w_{j,k} \right)^{1/2} \right)^2$$

and

$$\mathbf{G}_{4,2} = \left(\sum_{j=j_2+1}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \beta_{j,k}^2 \mathbf{1}_{\{|\beta_{j,k}| < 2\kappa \rho_n\}} w_{j,k} \right)^{1/2} \right)^2.$$

Using $\beta_{j,k}^2 \mathbf{1}_{\{|\beta_{j,k}| < 2\kappa \rho_n\}} \leq C \rho_n^2$ and $\sum_{k=0}^{2^j-1} w_{j,k} = 1$, we have

$$\begin{aligned} \mathbf{G}_{4,1} &\leq C \rho_n^2 \left(\sum_{j=\tau}^{j_2} 2^{j/2} \left(\sum_{k=0}^{2^j-1} w_{j,k} \right)^{1/2} \right)^2 = C \frac{\ln n}{n} \left(\sum_{j=\tau}^{j_2} 2^{j/2} \right)^2 \\ &\leq C \frac{\ln n}{n} 2^{j_2} \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}. \end{aligned}$$

Since $F \in B_{s,\infty}^w(M)$, we have

$$\begin{aligned} \mathbf{G}_{4,2} &\leq \left(\sum_{j=j_2+1}^{j_1} 2^{j/2} \left(\sum_{k=0}^{2^j-1} \beta_{j,k}^2 w_{j,k} \right)^{1/2} \right)^2 \leq \left(\sum_{j=j_2+1}^{j_1} 2^{-js} \right)^2 \\ &\leq C 2^{-2j_2 s} \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}. \end{aligned}$$

So

$$\mathbf{G}_4 \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}. \quad (6.16)$$

It follows from (6.12), (6.13), (6.15) and (6.16) that

$$\mathbf{G} \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}. \quad (6.17)$$

Combining (6.9), (6.10), (6.11) and (6.17), we have

$$\mathbb{E} \left(\int_0^1 \left(\hat{F}(x) - F(x) \right)^2 dx \right) \leq C \left(\frac{\ln n}{n} \right)^{2s/(2s+1)}.$$

The proof of Theorem 5.1 is complete. □

References

- [1] Brunel, E. and Comte, F. (2009). Cumulative distribution function estimation under interval censoring case 1. *Electron. J. Stat.*, 3, 1-24.
- [2] Brutti, P. (2008). Warped wavelets and vertical thresholding. Preprint. arXiv:0801.3319v1.
- [3] Carrasco, M. and Chen, X. (2002). Mixing and moment properties of various GARCH and stochastic volatility models. *Econometric Theory*, 18, 17-39.
- [4] Chesneau, C. (2007). A maxiset approach of a Gaussian noise model. (2007). *TEST*, 16, 3, 523-546.
- [5] Chesneau, C. and Willer, T. (2007). Numerical performances of a warped wavelet estimation procedure for regression in random design, Preprint, HAL:hal-00133831.
- [6] Cohen, A., Daubechies, I., Jawerth, B. and Vial, P. (1993). Wavelets on the interval and fast wavelet transforms. *Applied and Computational Harmonic Analysis*, 24, 1, 54-81.
- [7] Davydov, Y. (1970). The invariance principle for stationary processes. *Theor. Probab. Appl.*, 15, 3, 498-509.

- [8] Delyon, B. and Juditsky, A. (1996). On minimax wavelet estimators. *Applied Computational Harmonic Analysis*, 3, 215-228.
- [9] Diamond, I.D. and McDonald, J.W. (1991). The analysis of current status data. *Demo-graphic applications of event history analysis*. (eds J. Trussell, R. Hankinson & J. Tilton). Oxford University Press, Oxford.
- [10] Donoho, D.L., Johnstone, I.M., Kerkyacharian, G. and Picard, D. (1996). Density estimation by wavelet thresholding. *Annals of Statistics*, 24, 508–539.
- [11] Doukhan, P. (1994). *Mixing. Properties and Examples*. Lecture Notes in Statistics 85. Springer Verlag, New York.
- [12] Härdle, W., Kerkyacharian, G., Picard, D. and Tsybakov, A. (1998). *Wavelet, Approximation and Statistical Applications*. Lectures Notes in Statistics New York 129, Springer Verlag.
- [13] Jewell, N. P. and van der Laan, M. (2004). *Current status data: review, recent developments and open problems*. Advances in survival analysis, 625-642, Handbook of Statist., 23, Elsevier, Amsterdam.
- [14] Kerkyacharian, G. and Picard, D. (2004). Regression in random design and warped wavelets. *Bernoulli*, 10(6) :1053-1105.
- [15] Liebscher, E. (2001). Estimation of the density and the regression function under mixing conditions. *Statist. Decisions*, 19, (1), 9-26.
- [16] Meyer, Y. (1992). *Wavelets and Operators*. Cambridge University Press, Cambridge.
- [17] Modha, D. and Masry, E. (1996). Minimum complexity regression estimation with weakly dependent observations. *IEEE Trans. Inform. Theory*, 42, 2133-2145.
- [18] Pham Ngoc, T. M. (2009). Regression in random design and Bayesian warped wavelets estimators. *Electronic Journal of Statistics*, 3, 1084-1112.
- [19] van der Vaart, A. and van der Laan, M. J. (2006). Estimating a survival distribution with current status data and high-dimensional covariates. *Int. J. Biostat.*, 2, Art 9, 42pp.
- [20] Withers, C. S. (1981). Conditions for linear processes to be strong-mixing. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 57, 477-480.