



Coupling geochemical and biological approaches to assess the availability of cadmium in freshwater sediment

A. Dabrin, C. Durand, J. Garric, Olivier Geffard, B.J.D. Ferrari, Marina Coquery

► To cite this version:

A. Dabrin, C. Durand, J. Garric, Olivier Geffard, B.J.D. Ferrari, et al.. Coupling geochemical and biological approaches to assess the availability of cadmium in freshwater sediment. Science of the Total Environment, 2012, 424, p. 308 - p. 315. 10.1016/j.scitotenv.2012.02.069 . hal-00715728

HAL Id: hal-00715728

<https://hal.science/hal-00715728>

Submitted on 9 Jul 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Coupling geochemical and biological approaches to assess the availability of cadmium in freshwater sediment

Aymeric Dabrin, Cyrielle L. Durand, Jeanne Garric, Olivier Geffard, Benoit J.D. Ferrari, Marina Coquery

Cemagref, UR MALY, 3bis quai Chauveau, CP 220, F-69336 Lyon, France

Corresponding author: phone: +33 4 72 20 89 33

aymeric.dabrin@cemagref.fr

Abstract

Sediments are considered as a sink for metals, and the assessment of metal bioavailability for benthic organisms represents a great challenge. Diffusive Gradient in Thin films (DGT), developed to measure labile metals in aquatic media, have more recently been applied to sediment. Nevertheless, few studies have determined the relation between measurements from DGT and bioaccumulation in different benthic organisms. The aim of our work was to determine if labile metal measured by DGT in sediment is representative of bioavailable metal for benthic organisms. We focused our work on Cd and chose to use the diversity of ecological traits from different organisms to better understand the measurement given by DGT. We exposed simultaneously DGT and 3 macroinvertebrates species (the chironomid, *C. riparius*; the amphipod, *G. fossarum*; the mudsnail, *P. antipodarum*) to a natural sediment Cd-spiked at environmental relevant concentrations. The nature of sediment-bound Cd was also determined by means of sequential extractions in order to better interpret DGT measurements. Cadmium concentrations were determined in DGT and in the 3 organisms after one week of exposure. Results provided by DGT indicated that Cd was poorly released from particulate phase to pore water, suggesting that Cd measured by DGT was representative of the pore waters labile fraction. Sequential extractions showed that the percentage of Cd bound to carbonate fraction increased simultaneously with Cd-spiking level; hence, this Cd fraction was poorly reactive to supply DGT demand. Cadmium accumulation rates were similar between DGT measurements and *P. antipodarum*, suggesting that labile Cd in pore waters was representative of bioavailable Cd for this species. Cadmium accumulation rates in *C. riparius* were higher than in DGT, demonstrating that *C. riparius* can mobilize Cd bound to carbonate phase. *G. fossarum* showed the lowest Cd accumulation rates, suggesting that they were mainly exposed to Cd from overlaying waters.

Keywords: spiked-sediment, cadmium, DGT, sequential extractions, bioaccumulation, macroinvertebrates, ecological traits

1. Introduction

Sediment is generally considered as an important compartment to be investigated for evaluating the ecological integrity of an aquatic ecosystem. It provides a habitat for a large diversity of communities, but also serves as a reservoir for many pollutants. Sediment contamination primarily occurs via hydrodynamic processes during which many pollutants present in water are readily adsorbed on colloids that can aggregate with each other and settle to the bottom of aquatic systems (Buffle et al., 1998; Vignati et al., 2005). Once in sediment, the evaluation of pollutants mobility and bioavailability remains difficult because of the complex interactions between their own intrinsic properties, the geochemical/climatic factors and the biodiversity on-place (Munkittrick and McCarty, 1995; Chapman et al., 2003). In particular, the lability and the bioavailability of trace metals in sediments are function of the targeted metal, its colloidal, dissolved and particulate partition, its speciation, sediment physico-chemical characteristics (i.e., porosity, grain size, organic matter content), redox and pH conditions, and the physiological and ecological characteristics of the exposed organisms. Traditionally, the assessment of metal availability in sediments using a geochemical approach includes measurements of metals in dissolved and particulate phases and chemical extractions of available fractions (Tessier et al., 1979; Buykx et al., 2000). However, the total metal concentration in sediments is generally not a good proxy to predict bioaccumulation or biological effects (Di Toro et al., 1992). It was demonstrated in several case studies that pore waters metal concentration was a better tool to assess metal bioavailability in sediment (Di Toro et al., 1992; Ankley et al., 1993a; Berry et al., 1996); nonetheless, pore waters metal concentration did not reflect the supply of metals from the particulate phase when particles are ingested by organisms, such as chironomids larvae (Bervoets et al., 1997; Warren et al., 1998). Another parameter which strongly controls metal mobility in sediments is the acid-volatile sulphides (AVS). Acid-volatile sulphides are known to bind trace metals (i.e., Cd, Cu, Ni, Pb and Zn) in marine or freshwater sediments, resulting in insoluble sulphides complexes which reduce metal mobility and toxicity (Di Toro et al., 1992; Ankley et al., 1993b; Ankley et al., 1996; Burton et al., 2005). However, the AVS model did not support the prediction of metals bioaccumulation in benthic organisms for heavily contaminated freshwater sediments (Ankley, 1996; Warren et al., 1998; Lee et al., 2000; De Jonge et al., 2009). More recently, the Diffusive Gradient in Thin films (DGT), initially developed in the 1990's to measure labile metals in surface waters (Zhang and Davison, 1995), has been applied to soil and sediment (Harper et al., 2000; Zhang and Davison, 2000; Zhang et al., 2004; Rachou et al.,

2007). When a DGT is introduced into sediments, trace metals of the pore waters are rapidly bound by the resin of the device; this results in a decrease of pore waters trace metals concentrations, inducing a variable resupply from the particulate phase (Zhang et al., 1995). Metal accumulation in DGT and in terrestrial plants has been successfully compared because it operates similarly to active transport across a cell membrane (Zhang et al., 2001; Almås et al., 2006). The comparison is more difficult with benthic organisms, since metal transfer pathways are more complex, involving dissolved, colloidal and also particulate forms when sediment is ingested.

The aim of our study was to determine if the measurement of the labile Cd fraction in freshwater sediment using the DGT device is representative of the metallic bioavailable fraction for benthic organisms. We chose to use the diversity of ecological traits and habitats from different organisms to better understand the measurement given by the DGT tool in term of bioavailability. *Chironomus riparius* (insect larvae) are a relevant indicator of metal bioaccumulation because they live within the sediment during their larvae development and a large part of the metals they accumulate comes from the particulate phase due to particles ingestion (Bervoets et al., 1997; Warren et al., 1998). *Gammarus fossarum* (amphipod) and *Potamopyrgus antipodarum* (mudsnail) have been used in ecotoxicological test, especially for the evaluation of metals exposure from the water column (Geffard et al., 2010; Gust et al., 2011), and they have also been recently used to assess sediment toxicity (Mazurová et al., 2008; Mazurová et al., 2010; Schmitt et al., 2010). *G. fossarum* are epibenthic organisms living at the sediment-water interface; hence, they are mostly influenced by metals from the overlying waters. In contrast, *P. antipodarum* are known to live at the sediment-water interface and as well within the first millimeters depth of the sediment (Michaut, 1968); so they are also exposed to metals from pore waters. The nature of sediment-bound Cd was determined by means of sequential extractions in order to better interpret DGT measurements. Therefore, we performed the following laboratory tests under controlled conditions: parallel experiments were conducted by coupling DGT measurements with bioaccumulation measurements on *C. riparius*, *G. fossarum* and *P. antipodarum*. Since several physico-chemical parameters (AVS concentration, organic matter concentration, carbonates...) could induce a variability of geochemical and ecotoxicological results, we decided to use a unique low contaminated natural sediment, Cd-spiked at environmentally relevant concentrations.

2. Material and methods

2.1 Sediment sampling and Cd spiking

The sediment used in this study (PG) was collected in October 2009 from the Ain River which is a tributary of the Rhône River in the Southeast of France (N 45°809; E 5°209). This sediment was chosen since concentrations of inorganic (Table 1) and organic contaminants were low and under Threshold Effect Concentrations (TEC; Mc Donald et al., 2000). Surface sediment (10 cm) was sampled using a Van Veen grab (Hydrobios, GmbH), then it was immediately sieved through a 2 mm mesh. It was kept at 4°C before experiments. Five environmentally relevant nominal spiking Cd concentrations (0.5, 1.3, 3.1, 7.8 and 19.5 mg kg⁻¹ dry weight) were defined in order to cover a concentration range evenly spread between the Threshold Effect Concentration (TEC; 0.99 mg kg⁻¹) and the Probable Effect Concentration (PEC; 4.98 mg kg⁻¹) defined by Mc Donald et al. (2000). A Control was prepared in the same way as the spiked concentrations. Cadmium spiking (CdCl₂ salts) was realised by mixing wet sediment with the spiked solution using a liquid-solid ratio of 3.6. The water (FOS) used to proceed to the spiking mixing step corresponded to a spring-water adjusted to 300 µS cm⁻¹ with purified water. The composition and characteristics of this water were regularly checked during experimental protocol. The pH was circa 6.9; Ca²⁺ and Mg²⁺ concentrations averaged 35 mg L⁻¹ and 2.3 mg L⁻¹, respectively. Dissolved cadmium concentrations were below the limit of quantification (LQ = 0.01 µg L⁻¹). The content was mixed on a rotating wheel for 6 hours. After particles settling (48 hours), the supernatant water was removed using a vacuum pump and the sediment was homogenized with a plastic spoon prior to its introduction in test beakers. The same procedure was used for the Control sediment, but with Cd-free FOS water.

2.2 Experimental protocol

Subsamples of 100 ml of homogenized spiked sediment were distributed in 500 ml polypropylene beakers. An amount of 400 ml of FOS water was slowly added over the sediment minimising the suspension of the sediment. All beakers were placed in temperature regulated water baths at 21 ± 0.5 C, except for *G. fossarum* (12 ± 1°C), and with a 16:8 h artificial light:dark photoperiod. In order to equilibrate, systems were maintained during 10 days before introduction of DGT and organisms. Each beaker was continuously supplied with FOS water using a flow through system to obtain 4 renewals of overlaying waters per day.

2.3 DGT preparation

The DGT probe consists in a plastic piston loaded with a diffusive gel layer backed by an ion-exchange resin gel (Chelex 100) and a plastic cap with a 2 cm diameter window. A protective 0.45 μm cellulose nitrate filter (0.13 mm thickness, Millipore) separates the diffusive gel from the solution. Diffuse gels (0.8 mm thickness) and resin gels were purchased from DGT Research Ltd (Lancaster, UK). In order to prevent introduction of oxygen during the deployment within the sediment, DGT probes were deoxygenated by immersion in a suspension of 5 g L⁻¹ of Chelex-100 resin (Sigma) in 0.01 M NaCl, bubbled with nitrogen during 24 hours. Before their deployment into the sediment, the probes were transferred in a glove box under nitrogen atmosphere, put in clean plastic jars which were immediately sealed. The probes were exposed to air and oxygenated water only for a few seconds before the insertion into the sediment.

2.4 Geochemical approach

Geochemical experiments were only realised for the Control and for the 3 highest concentrations. DGT probes were introduced in the sediment (one DGT per beaker) by pressing them carefully at the sediment-water boundary layer with a polypropylene stick. The window of measurement of the DGT was positioned at a sediment depth of about 1 cm. DGT devices were retrieved at 2, 4, 8, 12, 24, 48, 72 and 144 hours of deployment (except at 2 and 4 hours for the Control sediment). One replicate was used at each time for the Control sediment and two replicates were retrieved for the Cd-spiked sediments at 4, 12, 24 and 144 hours. After removal, the DGT probes were thoroughly rinsed with MilliQ water and stored at 4°C into clean plastic bags. DGT resin gels were eluted with 2.5 ml of 1 M HNO₃ (Merck, Suprapur) into acid-cleaned polypropylene tubes. At the start and at the end of the DGT deployment period, pore waters were extracted from beakers by centrifugation (10 000 g, 30 min) and filtration (0.45 μm) under nitrogen atmosphere. Additionally, surface waters were sampled and filtered (0.45 μm) in duplicate at the start of the experiment. At the end of the deployment time, an aliquot of each sediment was sampled under nitrogen conditions and immediately kept at 4°C for sequential extractions and at -19°C for AVS extraction.

2.5 Bioaccumulation approach

C. riparius were obtained from the laboratory culture. *G. fossarum* and *P. antipodarum* were collected from natural populations in the Bourbre River and the Rhône River (France), respectively. The 3 test organisms were exposed to the Control sediment and to the 4 concentrations of Cd-spiked sediments in separate beakers from DGT exposure.

Cadmium bioaccumulation was measured on 3 replicate organisms pools (from 3 independent beakers) at the beginning and after 7 days (*C. riparius*, *G. fossarum*) or 10 days (*P. antipodarum*) of exposure. Depending on the mortality rates, 35 unshelled *P. antipodarum* (average of 0.033 g d.w.) , 10 *C. riparius* (average of 0.005 g d.w.) and 10 *G. fossarum* (average of 0.050 g d.w.) were retrieved for each replicate pool sample. All the organisms were thoroughly rinsed successively with FOS water, EDTA (3 mM L⁻¹, diluted in FOS water) and FOS water. The replicates were then freeze-dried (-20°C) before analysis.

2.6 Physico-chemical analyses

Sediment granulometry was determined using a Coulter LS-100. Acid-volatile sulfides (AVS) were determined on wet sediments using the procedure developed by Allen et al. (1993) and the limit of quantification was 0.12 µmol g⁻¹. Carbonates content was determined by addition of cold HCl to sediment and by measurement of liberated CO₂ volume. Loss on ignition (LOI) was realised by weighting sediment before and after heating at 500°C in order to estimate total organic matter content. In order to better assess the partitioning of Cd among the different carrier phases of the sediment, sequential extractions were realised according to Roullet et al. (2010). In order to preserve the sub-anoxic conditions in the sediment, sequential extractions procedure was realised under nitrogen atmosphere in a box glove using extractants previously deoxygenated. Duplicate samples were run throughout the procedure. To assess the percentage of recovery of the sequential extractions procedure, the bulk sediment was also mineralized with *aqua regia* (HNO₃: HCl, 3:1) using a microwave oven (MARS-5, CEM).

Organisms were mineralized with 2 mL of HNO₃ (Suprapur, 65%) using a microwave oven (MARS-5, CEM). Metal concentrations in surface waters, pore waters, DGT eluates, sediments extracts from sequential extractions and in biological samples were analysed by inductively coupled plasma mass spectrometry (ICP-MS, Thermo X7 series^{II}). Blanks and certified reference materials (TM-28.3, lake water, Environment Canada; IAEA-158, marine sediment; NRCC, TORT-2, lobster hepatopancreas) were systematically used to control analytical accuracy and precision. For sediment, typical expanded uncertainties (i.e., 95%

confidence intervals) were 5%. Relative standard deviations of Cd concentration for triplicate biological samples ranged between 6 and 38% for *G. fossarum*, between 1 and 13% for *P. antipodarum* and between 26 and 52% for *C. riparius*.

2.7 Data analysis

The flux of metals from sediment to the DGT device (F_{DGT}) provides information on the ability of the solid phase to resupply the pore waters. F_{DGT} was calculated as follows:

$$F_{DGT} = \frac{M}{A \times t} \quad (1)$$

Where M is the accumulated mass of Cd for a resin surface A (cm²), over the deployment time t (s). The time-averaged concentration at the interface of the sediment and the diffusive gel, C_{DGT} (in µg l⁻¹) was calculated as follows (Eq. 2), with D corresponding to the diffusive coefficient of the free metal ion (cm² s⁻¹) in the diffusive layer of thickness Δg (cm):

$$C_{DGT} = \frac{M \times \Delta g}{D \times t \times A} \quad (2)$$

C_{DGT} generally decreases during the deployment time because the pore waters concentration near the DGT probe progressively decreases. The R ratio, which corresponds to the ratio between C_{DGT} and the initial metal concentration in pore waters (C_{pw}) before the introduction of the DGT probes (Eq. 3), gives an indication of the depletion at the DGT interface (Harper et al., 2000):

$$R = \frac{C_{DGT}}{C_{pw}} \quad (3)$$

R can be simulated with the 2D-DIFS model (DGT-induced flux in soils) proposed by Sochaczewski et al. (2007). This model is based upon solute exchange between solid phase and pore waters, coupled to diffusion supply of metal to the interface and across the diffusion layer to the resin gel. DIFS-2D was used by fitting experimental values of R over time allowing to derive the parameters conditioning the response of the metal release from the sediment: the size of the particulate labile pool defined by the distribution coefficient K_{dl} (L kg⁻¹) and the characteristic response time (T_c) of the assumed 1st order reversible exchange process between dissolved and solid phases.

3. Results

3.1 Geochemical approaches

Physico-chemical characteristics and particulate metal concentrations of the studied sediment are presented in Table 1. The sediment showed a relatively fine composition since 70% of particles are represented by silts and clays ($< 63 \mu\text{m}$). Acid volatile sulfides concentrations in the sediment were similar between the Control sediment and spiked sediments, with a mean concentration of $8.1 \pm 0.5 \mu\text{mol g}^{-1}$. Calcium carbonates were the main component of the sediment (54%). Particulate trace metal concentrations (Cd, Cr, Ni, Cu, Zn, As and Pb) were low and all below the TEC. Particulate Cd concentrations obtained for the 5 Cd-spiked concentrations were close to the nominal ones, respectively 0.71, 1.35, 2.97, 6.97 and 17.5 mg kg^{-1} . Dissolved Cd concentrations in overlaying waters were below the limit of quantification ($0.010 \mu\text{g L}^{-1}$) in beakers containing the Control sediment and the 2 lowest Cd-spiked concentrations. Cadmium concentrations in overlaying waters increased with Cd spiking concentrations, with 0.014, 0.020 and $0.035 \mu\text{g L}^{-1}$ for PG3.1, PG7.8 and PG19.5, respectively. Cadmium pore waters concentrations ranged between $0.02 \mu\text{g L}^{-1}$ (Control sediment) and $1.82 \mu\text{g L}^{-1}$ (PG19.5). Only for PG19.5, Cd pore waters concentrations increased at the end of the experiment ($4.19 \mu\text{g L}^{-1}$). For the other sediments, Cd pore waters concentrations were similar between the start (T0) and the end (T1: 144 hours) of the experiment. The mass of Cd accumulated by the DGT from the Control sediment was too low to be detected in the eluates ($< 0.010 \mu\text{g L}^{-1}$). In contrast, for PG3.1, PG7.8 and PG19.5 Cd-spiked sediments, the mass of Cd trapped by the DGT increased with time up to 48 hours, then remained constant (Figure 1a). Cadmium fluxes through DGT rapidly decreased to reach low and similar values for the 3 highest spiked sediments (Cd fluxes = $0.18 \cdot 10^{-5} \pm 0.10 \text{ ng cm}^{-2} \text{ s}^{-1}$) after 144 hours of deployment (Figure 1b).

Experimental R data, obtained from the ratio between C_{DGT} and the initial pore waters concentrations (Equation 3), were fitted from inverse 2D-DIFS modelling, which allowed to estimate the range of the partition coefficient of the particulate labile Cd pool (K_{dl}) and the response time of the sorption processes (T_c) for the 3 highest Cd-spiked sediments (Table 2). The response times of sorption processes were relatively high ($34 \text{ s} < T_c < 12500 \text{ s}$) and the labile partition coefficients (K_{dl}) very low (< 2.79). The distribution of particulate Cd concentrations obtained by sequential extractions showed that Cd was not present under exchangeable form (F1) or bound to the residual phase (F5) whatever the level of Cd spiking

(Figure 2a). Particulate Cd concentrations in F2, F3 and F4 fractions increased with the level of Cd-spiking (Figure 2a), reaching up to 9.34, 5.77 and 1.94 mg kg⁻¹ respectively, for the highest level of spiking (PG19.5). The relative contribution of each fraction (Figure 2b) showed a progressive increase of F2 fraction from 16% (Control) to 54% (PG19.5), at the expense of F3 and F4 fractions, which decreased from 54% to 34% and from 30% to 11%, respectively.

3.2 Cd bioaccumulation

Internalised Cd bioaccumulation for *C. riparius*, *G. fossarum* and *P. antipodarum* are reported in Table 3. Since exposure time was different between the tested organisms and in order to compare results, we calculated Cd bioaccumulation rates (Figure 3). Cadmium bioaccumulation rates were the highest in *C. riparius*, ranging from 0.051 ± 0.019 µg g⁻¹ day⁻¹ for the Control sediment to 3.68 ± 1.92 µg g⁻¹ day⁻¹ for PG19.5. Cadmium bioaccumulation rates ranged between 0.018 ± 0.007 µg g⁻¹ day⁻¹ and 0.195 ± 0.040 µg g⁻¹ day⁻¹ for *G. fossarum* and between 0.060 ± 0.039 µg g⁻¹ day⁻¹ and 0.393 ± 0.008 µg g⁻¹ day⁻¹ for *P. antipodarum*, depending on exposure concentration.

4. Discussion

4.1 Supply of Cd from the particulate phase

For the 3 highest Cd-spiked sediments, the mass of Cd trapped by the DGT after 144 hours of deployment was 1.3 times higher between PG3.1 and PG7.8 and 2.0 times higher between PG7.8 and PG19.5. Since total particulate Cd concentrations were about 2.5 times higher between each studied sediment, this suggests that DGT measurement did not reflect the particulate Cd concentrations. Cadmium fluxes towards DGT as a function of time (Figure 1b) showed a typical distribution, corresponding to a rapid decrease during the first hours of deployment followed by a plateau. This distribution reflects a consumption of Cd from pore waters and a poor resupply from particles. In contrast, results obtained by Roulier et al. (2010) on natural sediments, exposed in similar conditions, showed an increase of Cd fluxes between 48 and 72 h followed by a decrease. The authors proposed that the increase of Cd fluxes was linked to Mn and Fe oxides reduction at the DGT/sediment interface. They obtained Cd fluxes ranging from ~0.1 to 0.7 10⁻⁵ ng cm⁻² s⁻¹ between 24 and 144 h, corresponding to particulate

Cd concentrations of 1.4 to 1.8 mg kg^{-1} . In our experiment, despite Cd particulate concentrations reaching up to 17.5 mg kg^{-1} , Cd fluxes were similar, ranging from 0.1 to $1.2 \cdot 10^{-5} \text{ ng cm}^{-2} \text{ s}^{-1}$. Moreover, the total mass of Cd was mainly trapped by DGT during the first 24 hours of deployment (38%: PG3.1; 74%: PG7.8; 71%: PG19.5). This suggests that Cd trapped by DGT mainly originated from the pore waters.

Pore waters Cd concentrations include all the metal phases (colloidal forms, inert and labile complexes, free ions), while DGT measures only free ions and labile metal. Although few data are available on Cd partition between colloidal and truly dissolved phase in sediment, Cd is known to be poorly present in colloidal complexes in surface waters (Sañudo-Wilhelmy et al., 2002). In our study, we analysed pore waters Cd concentrations at the start and at the end of the DGT deployment. Relative percent deviation (RPD) was below 10% for PG3.1 and PG7.8, showing a good reproducibility between beakers and that equilibration steady state was reached at the onset of the experiment. However, a large difference was noted for PG19.5 (RPD = 79%). This indicates an evolution of the sediment at high Cd-spiked level, probably due to a pH decrease (Hutchins et al., 2007). Indeed, metal-spiking might result in a drop of pH in relation with (i) hydrolysis of metal-spiked (ii) oxidative precipitation of Fe in pore waters and (iii) competition between protons H^+ and metal-spiked for particulate sorption sites.

Nonetheless the plot of experimental R ratio calculated for PG19.5 using initial and final pore waters concentrations showed very similar distributions (Figure 4). Moreover, the R ratio distribution for PG19.5 fitted very well with those of PG3.1 and PG7.8, calculated using means of initial and final pore waters concentrations. The distribution of the R ratio showed a fast decrease during the first 8 hours of DGT deployment, reaching a very small R ratio (< 0.16). Then, the R ratio slowly decreased and stabilized below 0.04 after 144 hours of DGT deployment. Apparently, whatever the level of Cd-spiking, Cd in pore waters was rapidly bound by the DGT in the first hours of deployment, followed by a very low resupply from the particulate phase. Ernstberger et al. (2002) exposed DGT to alluvial soil from 4 hours to 19.5 days; the R ratio was about 0.6 at the start of the experiment, then reached about 0.35 after 144 hours of deployment. In a similar study, Ernstberger et al. (2005) exposed DGT to 5 different soils with similar particulate Cd concentrations ($3.24 \pm 0.07 \text{ mg kg}^{-1}$), but with variable pH, granulometry and organic carbon content. The R ratios were higher than our results, ranging between ~ 0.8 and ~ 0.4 at the start of the experiment, and between ~ 0.6 and ~ 0.2 after 144 hours. The highest R ratios were obtained for the clayey soils with neutral pH and organic matter content from 2.6 to 5.8%. We obtained significantly lower R ratio, in spite

of high clay ($<63 \mu\text{m} = 70\%$) and high organic carbon content ($\text{LOI} = 13\%$). This suggests that in our case, spiked Cd was probably bound to unreactive particulate phases.

The 2D-DIFS model was used to model the R ratio distribution in order to determine K_{dl} and T_c parameters. An accurate determination of these two parameters was difficult due to a large number of possible combinations (K_{dl} , T_c) which can fit the experimental data (Lehto et al., 2008). Thus, several response times (T_c) and partition coefficient (K_{dl}) were proposed (Table 2). We obtained very low values for K_{dl} (< 2.79) and high T_c values (ranging between 34 and 12500 s) for the 3 highest Cd-spiked sediments. In Roulier et al. (2008), DGT were deployed in 6 polluted sediments and only one couple of K_{dl} (1) and T_c (5000) was comparable to our results. For the other sediments, K_{dl} ranged between 5 and 250, corresponding to small response times (<10 s in most cases). The highest K_{dl} values were obtained for sediments with high organic contents. Similarly, Ernstberger et al. (2005) showed that soils with poor resupply from the particulate phase were sandy and characterised by low pH and low organic matter content. Our tested sediments were fine with relatively high organic matter content (Table 1), but K_{dl} and T_c inferred from the R ratio distributions denoted a poor resupply from the particulate phase.

In our study, only total Cd concentration was modified; whatever the level of Cd-spiking, particulate Cd was strongly bound to the particles in response to the equilibrium modification induced by DGT insertion. Acid volatile sulphides are known to strongly bind Cd, reducing its mobility and its toxicity; but AVS concentrations were low in all Cd-spiked sediments ($8.1 \pm 0.5 \mu\text{mol g}^{-1}$; Table 1) and could not explain the equivalent low resupply from particulate phase obtained via DGT.

In all tested sediments, Cd was not present in the more potentially mobile particulate phase nor bound to the residual fraction. Cadmium was distributed between the F2, F3 and F4 particulate fractions, corresponding to carbonates, Fe and Mn oxides, organic matter and sulfides, respectively (Figure 2). With higher Cd-spiked level, the F2 fraction increased at the expense of F3 and F4 fractions. The significant F2 fraction was due to the high carbonate contents of sediments (54%). Indeed, Cd is known to have a great affinity with carbonates. Thakur et al. (2006) showed that Cd is effectively retained on CaCO_3 by the mechanism of chemisorption at low Cd concentrations, while at higher concentrations, precipitation of CdCO_3 on CaCO_3 surface phase predominated. This phenomenon was also reported in several studies, which showed that Cd was not only bound to sediments by sorption but as well by precipitation of secondary minerals such as otavite (CdCO_3) (Papadopoulos and Rowell, 1988; Komárek and Zeman, 2004; Cubillas et al., 2005). In summary, Cd spiked to this highly

carbonated sediment was firstly chemisorbed by carbonates surface, followed by CdCO_3 precipitation, inducing an exponential increase of the F2 fraction between each spiked sediment. This F2 fraction, presenting a great affinity for Cd, seems to be poorly reactive to supply the decrease of Cd in pore waters induced by the DGT demand.

4.2 Cd bioavailability

The slope of the regression between Cd accumulation in organisms, function of total particulate Cd, dissolved Cd in pore waters and in overlaying waters were more pronounced for *C. riparius* as compared to *G. fossarum* and *P. antipodarum* (Figure 3a, b, c), suggesting that *C. riparius* was exposed to a pool of Cd which, in contrast, was not available for the 2 others species. By means of in situ experiments, Warren et al. (1998) showed that prediction of Cd concentrations in most benthic animals would be more accurate if they were based on water column rather than on sedimentary Cd concentrations, except for sediment-feeding species such as Chironomids and Tubificidae, which also get Cd from sediments. Indeed, *C. riparius* live in close contact with sediments (Young and Harvey, 1991; Charbonneau et al., 1998), burrowing into sediments and creating their own microenvironment by irrigating and ingesting particles. Thus, ligands present in the gut fluid of deposit feeders, provide a complexation capacity for mobilization of sediments-bound metals (Chen et al., 2000). Roulier et al. (2008) reported a significant positive relationship between Cd concentrations in *C. riparius* and particulate Cd concentrations in freshwater contaminated sediments. Our data perfectly fit with this previous study performed under similar exposure conditions (Figure 5). Clearly, these results show that total Cd concentrations in sediments are a good predictor of Cd accumulation in *C. riparius*. Fitting our data with those of Roulier et al. (2008) indicates that particles ingestion by chironomids mobilize more Cd from the particulate phase than DGT. Since sequential extractions showed that Cd-spiked was mainly bound to the carbonate phase, we believe that chironomids are able to assimilate part of the Cd bound to carbonates. To our knowledge, this is the first time that a relationship is established between Cd bioaccumulation in chironomids and a specific Cd particulate fraction. In a similar way, Baumann and Fisher (2011) showed that bioaccumulated Cd in *Nereis Succinea* was related to exchangeable Cd and to Cd bound to the carbonate phase. Cadmium bioaccumulated in *G. fossarum* in the Control sediment (Table 3: 0.12 mg kg^{-1}) was comparable to that obtained (0.28 mg kg^{-1}) in the study of Felten et al. (2008), for the same exposure time in a control solution ($\text{Cd} < 0.30 \mu\text{g L}^{-1}$). Cadmium concentrations in *G.*

fossarum from the Control sediment, PG0.5 (0.204 mg kg⁻¹) and PG1.3 (0.299 mg kg⁻¹) were in the range of concentrations obtained for organisms collected on the upstream Bourbre River reference site (0.10 to 0.25 mg kg⁻¹), (Geffard et al., 2007). *G. fossarum* showed higher Cd concentration only starting from PG3.1 (0.441 mg kg⁻¹). Since *G. fossarum* are epibenthic species living at the water-sediment interface, they should mainly be influenced by Cd concentrations in overlaying waters. For PG19.5, overlaying waters Cd concentration was 0.035 µg L⁻¹ and could explain Cd content in *G. fossarum* (1.37 mg kg⁻¹). In fact, on the Morcille River (Beaujolais region, France), dissolved Cd concentration was about 0.040 µg L⁻¹ and Cd concentration in *G. fossarum* averaged 0.80 mg kg⁻¹ (personal communication). If *G. fossarum* were exposed to Cd from pore waters (1.82 µg L⁻¹) of PG19.5 sediment, Cd concentrations in *G. fossarum* should have been higher. Indeed, for example, after one week of field exposure at 1 µg L⁻¹, Lacaze et al. (2011) obtained Cd concentrations in *G. fossarum* ranging between 4 and 12 mg kg⁻¹. Pellet et al. (2009) also reported that after one week of laboratory exposure at 1 µg L⁻¹, Cd concentration in *G. fossarum* was about 2.5 mg kg⁻¹. Clearly, all these data strongly support the conclusion that during our experiment, *G. fossarum* exposed to Cd-spiked sediment were mainly influenced by Cd present in overlying waters.

The slope of Cd bioaccumulation in *P. antipodarum* as a function of particulate Cd concentrations was higher than the slope of Cd bioaccumulation in *G. fossarum* (Figure 3d). It looks like *P. antipodarum* were affected by a pool of Cd that was not seen by *G. fossarum*. Since *P. antipodarum* live at the sediment-water interface and within the first millimeters of the sediment (Michaut, 1968), the exposed organisms were apparently influenced by Cd from pore waters (0.46 to 4.19 µg L⁻¹). However, few data on Cd bioaccumulation in *P. antipodarum* are available to reinforce this hypothesis. Gust et al. (2011) showed on a French river with dissolved Cd concentrations of 0.6 µg L⁻¹ that Cd concentrations in *P. antipodarum* were about 1.82 mg kg⁻¹. These data are comparable with Cd concentrations in *P. antipodarum* of PG3.1 (1.02 mg kg⁻¹) and PG7.8 (2.12 mg kg⁻¹), corresponding to pore waters Cd concentrations of 0.45 and 1.38 µg L⁻¹.

Cadmium accumulation rates in DGT after 144h hours as a function of total particulate concentration showed that the slope of the regression was 2 orders of magnitude lower than the slope obtained for *C. riparius* (Figure 3d). These results corroborates our hypothesis that in a highly carbonated sediment, Cd was not released from the particulate phase to the pore waters in response to the locally depleted dissolved Cd concentration at the DGT-sediment interface, whereas *C. riparius* could bioaccumulate Cd from the particulate phase. Thus, in

highly carbonated sediment, DGT is a poor predictor for Cd bioaccumulation in chironomids. This is in accordance with results from Roulier et al. (2008), which did not show a good relationship between Cd bioaccumulated in chironomids and Cd trapped by DGT. For *G. fossarum*, the slope of the relationship was lower than the slope for DGT, corroborating the idea that *G. fossarum* exposed to Cd-spiked sediment were mainly influenced by Cd present in overlying waters, while DGT reflected labile Cd from pore waters. Thus, DGT is a poor predictor for Cd bioaccumulation in *G. fossarum* in this highly carbonated sediment. In contrast, the slope of Cd accumulation rates as a function of particulate Cd concentrations was similar between DGT and *P. antipodarum* (Figure 3d). As previously seen, Cd trapped by DGT was represented by labile Cd from pore waters. In conclusion, among the 3 tested species, *P. antipodarum* reflected the best Cd bioavailability from sediment pore waters which was assessed by DGT in this highly carbonated sediment.

5. Conclusions

Results of our geochemical approach, coupling DGT measurements and sequential extractions, showed that in a highly carbonated sediment, Cd bound to carbonates increased simultaneously with Cd-spiking level and was poorly reactive to supply the decrease of Cd in pore waters induced by DGT demand. This suggests that, in this kind of sediment, DGT measurement correctly reflects labile Cd from pore waters.

Our coupled biological approach, which included the confrontation of Cd accumulation rates in DGT and bioaccumulation for 3 benthic organisms with different ecological life traits, showed that *C. riparius* were exposed to a larger pool of Cd because they ingest particles. Since Cd was mainly bound to the carbonate fraction, *C. riparius* were in fact able to desorb Cd from carbonates. In contrast, Cd accumulation rates in *P. antipodarum* and DGT showed the same slope, suggesting that *P. antipodarum* were exposed to pore waters only and that the Cd pore waters labile fraction measured by DGT represented the bioavailable Cd fraction for this species. Cd accumulation rates were the lowest for *G. fossarum*; hence, this species was mainly influenced by Cd from overlying waters. Our results are consistent with the known behaviour of the 3 organisms.

Therefore, the proposed DGT technique proved its usefulness for sediment quality monitoring and bioaccumulation prediction; nevertheless, its relevance for Cd and others metals would need to be further assessed by testing DGT with a panel of sediments with contrasted characteristics. Moreover, it might be more accurate, although more complex to apply, if

DGT measurement was realised on the whole sediment-water interface, in connection with the living area of the studied species.

Acknowledgements

We gratefully acknowledge the following colleagues for sampling assistance, contribution to field work and analytical support: Jean-Louis Roulier, Raphaël Mons, Lysiane Dherret, Josiane Gahou, Eloise Vray, Bernard Motte and Rebecca Flück from Cemagref Lyon and Bernard Loizeau from Forel Institute (Switzerland). This study was supported by a grant from the French National Research Agency (ANR) in the framework of the “DIESE” program.

References

- Allen HE, Fu G, Deng B. Analysis of acid-volatile sulfide (AVS) and simultaneously extracted metals (SEM) for the estimation of potential toxicity in aquatic sediments. *Environ Toxicol Chem* 1993; 12: 1441-1453.
- Almås ÅR, Lombnæs P, Sogn TA, Mulder J. Speciation of Cd and Zn in contaminated soils assessed by DGT-DIFS, and WHAM/Model VI in relation to uptake by spinach and ryegrass. *Chemosphere* 2006; 62: 1647-1655.
- Ankley GT. Evaluation of metal/acid-volatile sulfide relationships in the prediction of metal bioaccumulation by benthic macroinvertebrates. *Environ Toxicol Chem* 1996; 15: 2138-2146.
- Ankley GT, Mattson VR, Leonard EN, West CW, Bennett JL. Predicting the acute toxicity of copper in freshwater sediments: Evaluation of the role of acid-volatile sulfide. *Environ Toxicol Chem* 1993a; 12: 315-320.
- Ankley GT, Liber K, Call DJ, Markee TP, Canfield TJ, Ingersoll CG. A field investigation of the relationship between zinc and acid volatile sulfide concentrations in freshwater sediments. *Journal of Aquatic Ecosystem Stress and Recovery* 1996; 5: 255-264.
- Ankley GT, Benoit DA, Hoke RA, Leonard EN, West CW, Phipps GL, Mattson VR, Anderson LA. Development and evaluation of test methods for benthic invertebrates and sediments: Effects of flow rate and feeding on water quality and exposure conditions. *Arch Environ Contam Toxicol* 1993b; 25: 12-19.

- 512 Baumann Z, Fisher NS. Relating the sediment phase speciation of arsenic, cadmium, and
513 chromium with their bioavailability for the deposit-feeding polychaete *Nereis*
514 *succinea*. *Environ Toxicol Chem* 2011; 30: 747-756.
- 515 Berry WJ, Hansen DJ, Mahony JD, Robson DL, Di Toro DM, Shipley BP, Rogers B, Corbin
516 JM, Boothman WS. Predicting the toxicity of metal-spiked laboratory sediments using
517 acid- volatile sulfide and interstitial water normalizations. *Environ Toxicol Chem*
518 1996; 15: 2067-2079.
- 519 Bervoets L, Blust R, De Wit M, Verheyen R. Relationships between river sediment
520 characteristics and trace metal concentrations in tubificid worms and chironomid
521 larvae. *Environ Pollut* 1997; 95: 345-356.
- 522 Buffle J, Wilkinson KJ, Stoll S, Filella M, Zhang J. A generalized description of aquatic
523 colloidal interactions: The three- colloidal component approach. *Environ Sci Technol*
524 1998; 32: 2887-2899.
- 525 Burton ED, Phillips IR, Hawker DW. Reactive sulfide relationships with trace metal
526 extractability in sediments from southern Moreton Bay, Australia. *Mar Pollut Bull*
527 2005; 50: 589-595.
- 528 Buykx SEJ, Bleijenberg M, Van den Hoop MAGT, Loch JPG. The effect of oxidation and
529 acidification on the speciation of heavy metals in sulfide-rich freshwater sediments
530 using a sequential extraction procedure. *J Environ Monitor* 2000; 2: 23-27.
- 531 Chapman AS, Foster IDL, Lees JA, Hodgkinson RJ, Jackson RH. Sediment and phosphorus
532 delivery from field to river via land drains in England and Wales. A risk assessment
533 using field and national databases. *Soil Use and Management* 2003; 19: 347-355.
- 534 Charbonneau P, Hare L, Carignan R. Use of x-ray images and a contrasting agent to study the
535 behavior of animals in soft sediments. *Limnol Oceanogr* 1998; 42: 1823-1828.
- 536 Chen Z, Mayer LM, Quetel C, Donard OFX, Self RFL, Jumars PA, Weston DP. High
537 concentrations of complexed metals in the guts of deposit feeders. *Limnol Oceanogr*
538 2000; 45: 1358-1367.
- 539 Cubillas P, Köhler S, Prieto M, Causserand C, Oelkers EH. How do mineral coatings affect
540 dissolution rates? An experimental study of coupled CaCO_3 dissolution - CdCO_3
541 precipitation. *Geochim Cosmochim Acta* 2005; 69: 5459-5476.
- 542 De Jonge M, Dreesen F, De Paepe J, Blust R, Bervoets L. Do acid volatile sulfides (AVS)
543 influence the accumulation of sediment-bound metals to benthic invertebrates under
544 natural field conditions? *Environ Sci Technol* 2009; 43: 4510-4516.

- 545 Di Toro DM, Mahony JD, Hansen DJ, Scott KJ, Carlson AR, Ankley GT. Acid volatile
546 sulfide predicts the acute toxicity of cadmium and nickel in sediments. *Environ Sci*
547 *Technol* 1992; 26: 96-101.
- 548 Ernstberger H, Davison W, Zhang H, Andrew TYE, Young S. Measurement and dynamic
549 modeling of trace metal mobilization in soils using DGT and DIFS. *Environ Sci*
550 *Technol* 2002; 36: 349-354.
- 551 Ernstberger H, Zhang H, Tye A, Young S, Davison W. Desorption kinetics of Cd, Zn, and Ni
552 measured in soils by DGT. *Environ Sci Technol* 2005; 39: 1591-1597.
- 553 Felten V, Charmantier G, Mons R, Geffard A, Rousselle P, Coquery M, Garric J, Geffard O.
554 Physiological and behavioural responses of *Gammarus pulex* (Crustacea: Amphipoda)
555 exposed to cadmium. *Aquat Toxicol* 2008; 86: 413-425.
- 556 Geffard A, Quéau H, Dedourge O, Biagianti-Risboug S, Geffard O. Influence of biotic and
557 abiotic factors on metallothionein level in *Gammarus pulex*. *Comparative*
558 *Biochemistry and Physiology - C Toxicology and Pharmacology* 2007; 145: 632-640.
- 559 Geffard O, Xuereb B, Chaumot A, Geffard A, Biagianti S, Noël C, Abbaci K, Garric J,
560 Charmantier G, Charmantier-Daures M. Ovarian cycle and embryonic development in
561 *Gammarus fossarum*: Application for reproductive toxicity assessment. *Environ*
562 *Toxicol Chem* 2010; 29: 2249-2259.
- 563 Gust M, Buronfosse T, Geffard O, Coquery M, Mons R, Abbaci K, Giamberini L, Garric J.
564 Comprehensive biological effects of a complex field poly-metallic pollution gradient
565 on the New Zealand mudsnail *Potamopyrgus antipodarum* (Gray). *Aquat Toxicol*
566 2011; 101: 100-108.
- 567 Harper MP, Davison W, Tych W. DIFS - A modelling and simulation tool for DGT induced
568 trace metal remobilisation in sediments and soils. *Environmental Modelling and*
569 *Software* 2000; 15: 55-66.
- 570 Hutchins CM, Teasdale PR, Lee J, Simpson SL. The effect of manipulating sediment pH on
571 the porewater chemistry of copper- and zinc-spiked sediments. *Chemosphere* 2007;
572 69: 1089-1099.
- 573 Komárek M, Zeman J. Dynamics of Cu, Zn, Cd, and Hg release from sediments at surface
574 conditions. *Bulletin of Geosciences* 2004; 79: 99-106.
- 575 Lacaze E, Devaux A, Mons R, Bony S, Garric J, Geffard A, Geffard O. DNA damage in
576 caged *Gammarus fossarum* amphipods: A tool for freshwater genotoxicity assessment.
577 *Environ Pollut* 2011; 159: 1682-1691.

- Lee BG, Lee JS, Luoma SN, Choi HJ, Koh CH. Influence of acid volatile sulfide and metal concentrations on metal bioavailability to marine invertebrates in contaminated sediments. *Environ Sci Technol* 2000; 34: 4517-4523.
- Lehto NJ, Sochaczewski Ł, Davison W, Tych W, Zhang H. Quantitative assessment of soil parameter (KD and TC) estimation using DGT measurements and the 2D DIFS model. *Chemosphere* 2008; 71: 795-801.
- MacDonald DD, Ingersoll CG, Berger TA. Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. *Arch Environ Contam Toxicol* 2000; 39: 20-31.
- Mazurová E, Hilscherová K, Jállová V, Köhler HR, Triebskorn R, Giesy JP, Bláha L. Endocrine effects of contaminated sediments on the freshwater snail *Potamopyrgus antipodarum* in vivo and in the cell bioassays in vitro. *Aquat Toxicol* 2008; 89: 172-179.
- Mazurová E, Hilscherová K, Šídlová-Štěpánková T, Köhler H, Triebskorn R, Jungmann D, Giesy JP, Bláha L. Chronic toxicity of contaminated sediments on reproduction and histopathology of the crustacean *Gammarus fossarum* and relationship with the chemical contamination and in vitro effects. *Journal of Soils and Sediments* 2010; 10: 423-433.
- Michaut P. Données biologiques sur un Gastéropode Prosobranche récemment introduit en Côte-d'Or, *Potamopyrgus jenkinsi*. *Hydrobiol* 1968; 32: 513-527.
- Munkittrick KR, McCarty LS. An integrated approach to aquatic ecosystem health: top-down, bottom-up or middle-out? *J Aquat Ecosyst Health* 1995; 4: 77-90.
- Papadopoulos P, Rowell DL. The reactions of cadmium with calcium carbonate surfaces. *J SOIL SCI* 1988; 39: 23-36.
- Pellet B, Geffard O, Lacour C, Kermoal T, Gourlay-FrancÉ C, Tusseau-Vuillemin MH. A model predicting waterborne cadmium bioaccumulation in *gammarus pulex*: the effects of dissolved organic ligands, calcium, and temperature. *Environ Toxicol Chem* 2009; 28: 2434-2442.
- Rachou J, Hendershot W, Sauvé S. Soil organic matter impacts upon fluxes of cadmium in soils measured using diffusive gradients in thin films. *Communications in Soil Science and Plant Analysis* 2007; 38: 1619-1636.
- Roulier JL, Belaud S, Coquery M. Comparison of dynamic mobilization of Co, Cd and Pb in sediments using DGT and metal mobility assessed by sequential extraction. *Chemosphere* 2010; 79: 839-843.

- 612 Roullet JL, Tusseau-Vuillemin MH, Coquery M, Geffard O, Garric J. Measurement of
613 dynamic mobilization of trace metals in sediments using DGT and comparison with
614 bioaccumulation in *Chironomus riparius*: First results of an experimental study.
615 *Chemosphere* 2008; 70: 925-932.
- 616 Sañudo-Wilhelmy SA, Rossi FK, Bokuniewicz H, Paulsen RJ. Trace metal levels in
617 uncontaminated groundwater of a coastal watershed: Importance of colloidal forms.
618 *Environ Sci Technol* 2002; 36: 1435-1441.
- 619 Schmitt C, Balaam J, Leonards P, Brix R, Streck G, Tuikka A, Bervoets L, Brack W, van
620 Hattum B, Meire P, de Deckere E. Characterizing field sediments from three European
621 river basins with special emphasis on endocrine effects - A recommendation for
622 *Potamopyrgus antipodarum* as test organism. *Chemosphere* 2010; 80: 13-19.
- 623 Sochaczewski Ł, Tych W, Davison B, Zhang H. 2D DGT induced fluxes in sediments and
624 soils (2D DIFS). *Environmental Modelling and Software* 2007; 22: 14-23.
- 625 Tessier A, Campbell PGC, Blsson M. Sequential extraction procedure for the speciation of
626 particulate trace metals. *Anal Chem* 1979; 51: 844-851.
- 627 Thakur SK, Tomar NK, Pandeya SB. Influence of phosphate on cadmium sorption by calcium
628 carbonate. *Geoderma* 2006; 130: 240-249.
- 629 Vignati DAL, Dworak T, Ferrari B, Koukal B, Loizeau JL, Minouflet M, Camusso MI,
630 Polesello S, Dominik J. Assessment of the geochemical role of colloids and their
631 impact on contaminant toxicity in freshwaters: An example from the Lambro-Po
632 system (Italy). *Environ Sci Technol* 2005; 39: 489-497.
- 633 Warren LA, Tessier A, Hare L. Modelling cadmium accumulation by benthic invertebrates in
634 situ: The relative contributions of sediment and overlying water reservoirs to organism
635 cadmium concentrations. *Limnol Oceanogr* 1998; 43: 1442-1454.
- 636 Young LB, Harvey HH. Metal concentrations in chironomids in relation to the geochemical
637 characteristics of surficial sediments. *Arch Environ Contam Toxicol* 1991; 21: 202-
638 211.
- 639 Zhang H, Davison W. Performance characteristics of diffusion gradients in thin films for the
640 in situ measurement of trace metals in aqueous solution. *Anal Chem* 1995; 67: 3391-
641 3400.
- 642 Zhang H, Davison W. Direct in situ measurements of labile inorganic and organically bound
643 metal species in synthetic solutions and natural waters using diffusive gradients in thin
644 films. *Anal Chem* 2000; 72: 4447-4457.

- Zhang H, Davison W, Miller S, Tych W. In situ high resolution measurements of fluxes of Ni, Cu, Fe, and Mn and concentrations of Zn and Cd in porewaters by DGT. *Geochim Cosmochim Acta* 1995; 59: 4181-4192.
- Zhang H, Lombi E, Smolders E, McGrath S. Kinetics of Zn release in soils and prediction of Zn concentration in plants using diffusive gradients in thin films. *Environ Sci Technol* 2004; 38: 3608-3613.
- Zhang H, Zhao FJ, Sun B, Davison W, McGrath SP. A new method to measure effective soil solution concentration predicts copper availability to plants. *Environ Sci Technol* 2001; 35: 2602-2607.

Table captions

Table 1: Physico-chemical characteristics and particulate trace metal concentrations of the studied sediment. Cadmium concentrations in pore waters and in overlaying waters were determined during the experiment.

Table 2: Partition coefficient (K_d) and parameters (K_{dl} : labile partition coefficient; T_c : characteristic response time) deduced from 2D-DIFS modelling for the Control sediment and the 3 highest Cd-spiked sediment concentrations.

Table 3: Concentrations of Cd bioaccumulated in *C. riparius*, *G. fossarum* and *P. antipodarum*.

Figure captions

Fig. 1: (a) Mass of Cd accumulated in the DGT resin and (b) fluxes of Cd through the DGT as a function of deployment time for the 3 highest Cd-spiked sediment concentrations (PG3.1, PG7.8, PG19.5).

Fig. 2: (a) Cd concentrations in F2, F3 and F4 particulate fractions (mg kg^{-1}) and (b) relative contributions (%) of the 3 fractions to total particulate Cd concentrations (for the Control sediment and 3 highest Cd-spiked sediments PG3.1, PG7.8 and PG19.5).

Fig. 3: Bioaccumulation Cd rates ($\text{mean} \pm \text{sd}$, $n=3$) in *C. riparius*, *G. fossarum* and *P. antipodarum* as a function of (a) total particulate Cd concentrations, (b) dissolved pore waters Cd concentrations and (c) dissolved Cd concentration in overlaying waters. Accumulation Cd rates in DGT after 144 hours of deployment are reported for the 3 highest Cd-spiked sediment concentrations (d).

Fig 4: Distribution of the R ratio for sediments Cd-spiked at 3.1, 7.8 and 19.5 mg kg^{-1} .

Fig. 5: Cadmium concentrations in *C. riparius* as a function of particulate Cd concentrations for (●) this study and for (○) the study of Roulier et al. (2008).




Fig. 1.




Fig. 2.




Fig. 3:




Fig 4:




Fig. 5: