



HAL
open science

Parabolic geodesics as parallel curves in parabolic geometries

Marc Herzlich

► **To cite this version:**

Marc Herzlich. Parabolic geodesics as parallel curves in parabolic geometries. *International Journal of Mathematics*, 2013, 24 (9), pp.1350067. 10.1142/S0129167X13500675 . hal-00718588

HAL Id: hal-00718588

<https://hal.science/hal-00718588>

Submitted on 17 Jul 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PARABOLIC GEODESICS AS PARALLEL CURVES IN PARABOLIC GEOMETRIES

MARC HERZLICH

ABSTRACT. We give a simple characterization of the parabolic geodesics introduced by Čap, Slovák and Žádník for all parabolic geometries. This goes through the definition of a natural connection on the space of Weyl structures. We then show that parabolic geodesics can be characterized as the following data: a curve on the manifold and a Weyl structure along the curve, so that the curve is a geodesic for its companion Weyl structure and the Weyl structure is parallel along the curve and in the direction of the tangent vector of the curve.

INTRODUCTION

Parabolic geometries have attracted much interest in the recent years, providing an efficient framework for tackling problems in various geometries as, *e.g.*, those induced by conformal or Cauchy-Riemann (CR) structures. There exist curves in parabolic geometries (hereafter called parabolic geodesics) that play a similar role as geodesics in Riemannian geometry: for instance, they provide natural local charts adapted to the geometry at hand, see [16] for applications.

In conformal geometry, conformal geodesics may have been known to Cartan [8] and their study has been revived in the 80's by Ferrand [14]. For CR structures, canonical curves, or chains, were defined by Cartan [9, 10] in dimension 3, and by Chern and Moser [11] in any dimension. Another definition was then introduced by Fefferman [13] and more general curves were studied by Koch [21, 22]. A general definition of distinguished curves for all parabolic geometries, or *parabolic geodesics*, was then given by Čap, Slovák, and Žádník [6, 7].

However, and contrarily to geodesics, parabolic geodesics are not defined directly as solutions of a natural differential equation on the manifold itself, but rather as projections of special curves on a natural principal bundle (the so-called Cartan bundle) over the base manifold. To the knowledge of the author, natural differential equations for parabolic geodesics have been derived only in the case of conformal or CR geometry [1, 11, 13, 14, 21, 22]. The goal of this short paper is then to provide such a simple definition. To

Date: July 17th, 2012.

2010 Mathematics Subject Classification. 53B25, 53A55, 53A30.

Key words and phrases. Parabolic geometry, parabolic geodesics.

achieve this, we shall need to study in some detail a special class of connections in parabolic geometries: the Weyl structures, introduced by Čap and Slovák [5] and independently by Calderbank, Diemer and Souček [3]. The set of Weyl structures forms a bundle, and we shall show that this bundle admits a natural connection. In a second step, we shall prove the following

Main Theorem. *Distinguished curves (parabolic geodesics) on a manifold endowed with a parabolic geometry are exactly the curves that satisfy the following requirements:*

- (i) *they are parallel for a Weyl structure ;*
- (ii) *this Weyl structure is itself parallel along the curve and in the direction of the tangent vector of the curve for the natural connection acting on the bundle of Weyl structures.*

The paper is organised as follows. Section 1 recalls the basic facts on parabolic geometries and their Weyl structures. Sections 2 and 3 yield the definition of the natural connection on the bundle of Weyl structures. Our main result is proved in section 4. The paper ends in section 5 with a few examples of computations, where we shall re-derive from our setting known equations in the most simple cases.

Acknowledgements. – This note emerged from a series of discussions with Paul Gauduchon and the author is happy to acknowledge his influence on this work. He also thanks Olivier Biquard and Charles Francès for their interest and comments.

1. PARABOLIC GEOMETRIES

We recall here a few basic facts on parabolic geometries. Our main references in this section are [3, 5, 6]. In all what follows, we shall denote by M a connected manifold (of dimension n), by G a (real) semi-simple Lie group, and by P a closed parabolic subgroup of G , whose Lie algebras will be denoted by \mathfrak{g} and \mathfrak{p} . This implies the existence of a filtration of \mathfrak{g} :

$$\mathfrak{g} = \mathfrak{g}_{(-k)} \supset \mathfrak{g}_{(-k+1)} \supset \cdots \supset \mathfrak{g}_{(0)} = \mathfrak{p} \supset \mathfrak{g}_{(1)} \supset \cdots \supset \mathfrak{g}_{(k)}$$

for some positive integer k . Here one has $\mathfrak{g}_{(1)} = [\mathfrak{p}, \mathfrak{p}]$ (thus $\mathfrak{g}_{(1)}$ is the maximal nilpotent ideal in \mathfrak{p} and one has $\mathfrak{g}_{(1)} = \mathfrak{p}^\perp$ with respect to the Killing form of \mathfrak{g}), $\mathfrak{g}_{(2)} = [\mathfrak{g}_{(1)}, \mathfrak{g}_{(1)}]$, and the subsequent $\mathfrak{g}_{(j)}$ for $j > 0$ are the elements of the descending central series of $\mathfrak{g}_{(1)}$; for $j < 0$, the terms of the filtration are defined by $\mathfrak{g}_{(j)} = (\mathfrak{g}_{(-j-1)})^\perp$. The associated graded Lie algebra $\text{gr}(\mathfrak{g})$ factors as

$$\text{gr}(\mathfrak{g}) = \mathfrak{g}^{-k} \oplus \mathfrak{g}^{-k+1} \oplus \cdots \oplus \mathfrak{g}^0 \oplus \mathfrak{g}^1 \oplus \cdots \oplus \mathfrak{g}^k$$

where $\mathfrak{g}^j = \mathfrak{g}_{(j)}/\mathfrak{g}_{(j+1)}$. Thus, $[\mathfrak{g}^i, \mathfrak{g}^j] \subset \mathfrak{g}^{i+j}$ for any i and j in $\{-k, \dots, k\}$.

Any choice of a Cartan involution in \mathfrak{g} induces a Lie algebra isomorphism between \mathfrak{g} and $\text{gr}(\mathfrak{g})$. There is no canonical choice of a Cartan involution, but any two such choices are related by the adjoint action of an element of P .

For reasons that will be clear below, the choice of such an identification will be harmless for our purposes, hence we shall assume that such a choice has been made once and for all. It will also be useful to consider the subalgebras $\mathfrak{g}^+ = \mathfrak{g}^0 \oplus \mathfrak{g}^1 \oplus \cdots \oplus \mathfrak{g}^k$ and $\mathfrak{g}^- = \mathfrak{g}^{-k} \oplus \cdots \oplus \mathfrak{g}^{-1} \oplus \mathfrak{g}^0$ in \mathfrak{g} . The smaller subalgebras $\mathfrak{g}^{>0}$ and $\mathfrak{g}^{<0}$ are defined by removing the 0-th factor in the above formulas. We shall denote by P^0 the closed subgroup of P whose adjoint action preserves the grading of \mathfrak{g} into the \mathfrak{g}^0 -modules \mathfrak{g}^i ($i = -k, \dots, k$); its Lie algebra is of course \mathfrak{g}^0 . In the same vein, $P^{>0} = \exp(\mathfrak{g}^{>0})$ is the connected subgroup of P whose Lie algebra is $\mathfrak{g}^{>0}$. It is then clear that $P_0 = P/P^{>0}$ is isomorphic to P^0 ; this is the group known as the *Levi factor*.

Definition 1.1. A parabolic geometry of type (G, P) on M is given by a P -principal bundle $\mathcal{G} \rightarrow M$ and a section ω (a *Cartan connection*) of the bundle of 1-forms on \mathcal{G} with values in \mathfrak{g} such that

- (a) ω is equivariant with respect to the natural action of P , *i.e.*

$$(R_p)^*\omega = \text{Ad}(p^{-1}) \circ \omega \quad \text{for any } p \in P;$$

- (b) ω reproduces the Maurer-Cartan form of P on vertical vector fields ;
(c) $\omega_e : T_e\mathcal{G} \rightarrow \mathfrak{g}$ is an isomorphism at any e in \mathcal{G} .

Under a mild cohomological condition on the Lie algebras \mathfrak{g} and \mathfrak{p} , the geometric structure can be entirely described at the bottom level, *i.e.* on the manifold M itself, as a filtration of the tangent bundle TM of the form

$$TM = T_{(-k)}M \supset \cdots \supset T_{(-1)}M,$$

induced by the identification of TM with the associated bundle $\mathcal{G} \times_P \mathfrak{g}/\mathfrak{p}$, together with a reduction of the structure group of the associated graded bundle $\text{gr}(TM)$ to P_0 . A fundamental theorem due to Čap-Schichl [4] (see also [6] for a presentation in book form with a different proof) asserts that under another, similar, algebraic condition on the pair $(\mathfrak{g}, \mathfrak{p})$, any such filtered structure on TM which satisfies a simple compatibility condition between the Lie algebraic bracket of \mathfrak{g} and the bracket of vector fields (this condition is called *regularity*) induces a parabolic geometry in the sense given above. Moreover, the latter is unique up to isomorphism if the Cartan connection is required to be *normal*, another algebraic condition on the curvature of ω . All classical examples of parabolic geometries (*e.g.* conformal geometries of any signature, CR geometry, etc.) belong to this setting. Another example is projective geometry (*i.e.* the geometry induced by a projective class of torsion-free connections on the tangent bundle), with the special feature that the first cohomological condition alluded to above is not satisfied in this case, hence the underlying geometric structure does not reduce to a filtration of the tangent bundle and a reduction of the structure group of the induced graded bundle; apart from this, it however fits into the parabolic setting rather nicely, and we shall also discuss this example later.

The last tool for our construction will be a subgroup P^- of G whose Lie algebra is \mathfrak{g}^- . We shall moreover assume that there is such a subgroup so

that $P \cap P^- = P^0$. This is a mild algebraic condition which is satisfied in all classical cases; note that it is also satisfied when all three groups P , P^0 , and P^- are connected.

2. WEYL STRUCTURES

Similarly to the whole family of Levi-Civita connections of all metrics in a conformal class, very parabolic geometry admits a family of distinguished connections called *Weyl structures*. There are different ways to define them, but the following, due to Čap and Slovák [5], will be the most useful here.

Definition 2.1. A Weyl structure is a reduction \mathcal{G}^0 of the P -principal bundle \mathcal{G} to the structure group P^0 . This can be equivalently described as a P^0 -equivariant bundle map $\sigma : \mathcal{G}_0 \rightarrow \mathcal{G}$, where $\mathcal{G}_0 = \mathcal{G}/P^{>0}$.

Any Weyl structure induces a principal connection on \mathcal{G} , whose construction is done as follows : at any point $e = \sigma(e_0)$ of the image of σ in \mathcal{G} , the tangent bundle $T\mathcal{G}$ splits as

$$T_e\mathcal{G} = T_e^V\mathcal{G} \oplus T_e^H\mathcal{G}$$

where $T_e^V\mathcal{G}$ is the vertical bundle and $T_e^H\mathcal{G} = (\omega_e)^{-1}(\mathfrak{g}^{<0})$. Since ω is equivariant and P^0 stabilizes every graded subspace \mathfrak{g}^j in \mathfrak{g} , this splitting is invariant by the right action of P^0 on $\sigma(\mathcal{G}_0)$ and it can be extended as an equivariant connection on \mathcal{G} by the right action. This of course yields connections on bundles associated to \mathcal{G} through a representation of P , such as, e.g., the tangent or cotangent bundles of the base manifold M . We shall term it the *Weyl connection* induced by σ to make it clear that it depends on the choice of σ .

An equivalent definition of Weyl structures has been given in [3]. Weyl structures are defined there directly as special P -equivariant splittings of $T_e\mathcal{G}$ compatible with the natural filtration of \mathfrak{g} . It is an easy exercise to show that any such splitting is necessarily induced by a unique section of \mathcal{G} over \mathcal{G}_0 , thus yielding the equivalence between the two definitions.

From now on, we shall make extensive use of bundles associated with the Cartan bundle \mathcal{G} . We shall need the following notation: for any space X endowed with a P -action, the bundle $\mathcal{G} \times_P X$ is the quotient of the product space $\mathcal{G} \times X$ by the equivalence relation $(e, x) \sim (ep, p^{-1}x)$ for all $p \in P$. We denote by $[e, x]$ the class of (e, x) in $\mathcal{G} \times_P X$, and sections of $\mathcal{G} \times_P X$ are given by P -equivariant maps from \mathcal{G} to X .

It is well known that reductions to P^0 of the structure group of a P -principal bundle are classified by sections of the associated bundle

$$\mathcal{W} = \mathcal{G} \times_P P/P^0.$$

Indeed, given a reduction σ , one can build a map \mathfrak{p} from \mathcal{G} to P/P^0 which is P -equivariant as follows: one begins by fixing $\mathfrak{p}(\sigma(e_0)) = [1]_{P/P^0}$ at any point $\sigma(e_0)$ in the image of σ ; note that here and henceforth, an element

between brackets will always denote its class in the quotient space indicated as an index to the right bracket. One then extends \mathfrak{p} equivariantly by letting

$$\mathfrak{p}(\sigma(e_0)p) = p^{-1}[1]_{P/P^0} = [p^{-1}]_{P/P^0}.$$

In other words, σ induces a section $\tau = [\sigma(e_0), [1]_{P/P^0}] = [e, \mathfrak{p}(e)]$ of the bundle \mathcal{W} .

Definition 2.2. The bundle \mathcal{W} is called the *bundle of Weyl structures* of the parabolic geometry.

The space of sections of \mathcal{W} admits a natural right action of the space of sections of the bundle in nilpotent groups $\mathcal{G}_0 \times_{P^0} P^{>0} = \mathcal{G} \times_P P^{>0}$, where the action of p in P or in P^0 on an element n of $P^{>0}$ is by inner automorphisms: $n \mapsto \text{Int}_p(n) = pn p^{-1}$. Hence our claim here is that there is a right action of the space of sections of $\mathcal{G} \times_{\text{Int}} P^{>0}$ on $\mathcal{W} = \mathcal{G} \times_L P/P^0$ (where L stands for the left action). Indeed, a section of $\mathcal{G} \times_{\text{Int}} P^{>0}$, seen as an equivariant map $n : \mathcal{G} \rightarrow P^{>0}$, acts on the right on

$$\mathfrak{p} : \mathcal{G} \rightarrow P/P^0, \quad \mathfrak{p}(e) = [p(e)]_{P/P^0}$$

to yield a map

$$\mathfrak{p} \circ n : \mathcal{G} \rightarrow P/P^0, \quad (\mathfrak{p} \circ n)(e) = [n^{-1}(e)p(e)]_{P/P^0}$$

This last section of \mathcal{W} corresponds to the Weyl structure $\sigma \cdot n$ where the right factor n acts by right multiplication on the map $\sigma : \mathcal{G}_0 \rightarrow \mathcal{G}$. This description of the action of $\mathcal{G} \times_P P^{>0}$ on the bundle of Weyl structures is equivalent to that given in [3, §2.4–2.5].

The bundle \mathcal{W} is associated to the principal bundle \mathcal{G} through a P -action on P/P^0 which does not extend to a G -action. Since \mathcal{G} does not have a true connection but only a Cartan connection, it does not induce a natural connection on \mathcal{W} . Of course, any Weyl structure induces a connection on \mathcal{W} , but since there is no natural choice of a Weyl structure this does not help for our purposes. However, there is a special class of associated bundles that are naturally endowed with a canonical connection: these are the bundles induced from a G -action.

3. THE NATURAL CONNECTION ON THE SPACE OF WEYL STRUCTURES

Let X be a manifold endowed with an action of the group G . The bundle $\mathcal{G} \times_P X$ is then endowed with a natural connection, which can be defined as follows: $\mathcal{G} \times_P X$ is a quotient of the product space $\mathcal{G} \times X$ and the kernel of the differential of the projection map $\tilde{\pi} : \mathcal{G} \times X \rightarrow \mathcal{G} \times_P X$ at (e, x) is the range of the injective map

$$\mathfrak{p} \rightarrow T_{(e,x)}(\mathcal{G} \times X) = T_e \mathcal{G} \times T_x X, \quad \xi \mapsto (\tilde{\xi}_e, -\check{\xi}_x)$$

where $\tilde{\xi}_e$ and $\check{\xi}_x$ are the natural vectors induced by the actions:

$$\tilde{\xi}_e = \frac{d}{dt} (e \exp(t\xi))|_{t=0} \quad \text{and} \quad \check{\xi}_x = \frac{d}{dt} (\exp(t\xi) x)|_{t=0}.$$

Any element ξ of \mathfrak{p} yields a vector field $\check{\xi} = \omega^{-1}(\xi)$ on \mathcal{G} , which coincides with the previously defined $\check{\xi}$. Thus, the kernel of the projection map is also identified to the range of the map $\xi \mapsto (\omega_e^{-1}(\xi), -\check{\xi}_x)$ from \mathfrak{p} to $T_e\mathcal{G} \times T_xX$.

Definition 3.1. The horizontal space $T^H(\mathcal{G} \times_P X)$ is the projection through $\tilde{\pi}$ of $T^H(\mathcal{G} \times X) = \text{Range}(\xi \in \mathfrak{g} \mapsto (\omega_e^{-1}(\xi), -\check{\xi}_x))$.

Thus, the horizontal space $T^H(\mathcal{G} \times_P X)$ is naturally isomorphic to the quotient space $T^H(\mathcal{G} \times X) / \text{Ker } \tilde{\pi}_*$. Its dimension equals that of $\mathfrak{g}/\mathfrak{p}$ hence that of M , and it projects itself isomorphically to TM through the map

$$(\omega_e^{-1}(\xi), -\check{\xi}_x) \mapsto \pi_*\left(\omega_e^{-1}(\xi)\right)$$

tangent to the bundle projection $\pi : \mathcal{G} \rightarrow M$. The connection induced by this choice of horizontal space will be called *the canonical (tractor) connection of the space $\mathcal{G} \times_P X$* .

We can now apply this construction to the case where $X = G/P^-$. The tangent space at $[g]_{G/P^-}$ to X is the image of the map $\xi \mapsto \check{\xi}_{[g]}$ already defined above. By analogy with conformal geometry (where G/P^- is the Moebius sphere up to a covering), we shall make the following definition.

Definition 3.2. The bundle $\mathcal{S} = \mathcal{G} \times_P G/P^-$ is called the *Moebius bundle* of the parabolic geometry.

Proposition 3.3. *The bundle of Weyl structures \mathcal{W} is naturally embedded as an open set in the Moebius bundle \mathcal{S} .*

Proof. – This is clear since one may send any element $[e, [p]_{P/P^0}]$ of \mathcal{W} to $[e, [p]_{G/P^-}]$, which is a well defined map from a P -bundle to another. The embedding is nothing but the natural embedding of P/P^0 in G/P^- in each fiber, whose differential is obviously injective. \square

Corollary 3.4. *The bundle of Weyl structures admits a natural tractor connection inherited from its embedding as an open set of the Moebius bundle.*

We shall now denote by \mathcal{D} the tractor connection on Weyl structures. Choose now a (local) Weyl structure σ (thought as a P^0 -equivariant section $\sigma : \mathcal{G}_0 \rightarrow \mathcal{G}$), and a vector X at a point x in M . Then one may lift X to a vector \hat{X}_0 in \mathcal{G}_0 at a point e_0 in \mathcal{G}_0 , and the tractor derivative $\mathcal{D}_X\sigma$ of σ at x in the direction of X is given by the following element of the cotangent bundle $T^*M = \mathcal{G} \times_P \mathfrak{g}^{>0}$

$$\mathcal{D}_X\sigma = \left[\sigma(e_0), \text{proj}_{\mathfrak{g}^{>0}} \circ \omega_{\sigma(e_0)}(\sigma_*(\hat{X}_0)) \right].$$

With this in mind, we can now prove a helpful fact. We begin by recalling that any choice of Weyl structure $\sigma : \mathcal{G}_0 \rightarrow \mathcal{G}$ induces a natural connection on \mathcal{G} obtained by choosing to split the tangent bundle of \mathcal{G} at any point in the image of σ as the direct sum of the vertical subspace $\omega^{-1}(\mathfrak{p})$ with the horizontal subspace $\omega^{-1}(\mathfrak{g}^{<0})$. Thus, it is given by the connection 1-form $\omega^\sigma = \omega^+ = \text{proj}^{\mathfrak{p}} \circ \omega$ where $\text{proj}^{\mathfrak{p}}$ denotes the projection onto \mathfrak{p} in \mathfrak{g} .

The Weyl connection is a \mathcal{G}_0 -connection: indeed, by the very definition of P^0 , the right P^0 -action on \mathcal{G} preserves the decomposition of $T\mathcal{G} = \omega^{-1}(\mathfrak{g})$ into its graded components. Hence the connection preserves the P^0 -bundle given by the image of σ . Equivalently, the Weyl connection can be pulled back to \mathcal{G}_0 , and the connection 1-form associated to this connection on \mathcal{G}_0 is given by the projection $\omega^0 = \text{proj}^{\mathfrak{g}^0} \circ \omega$ onto \mathfrak{g}^0 of ω .

Lemma 3.5. *The tractor derivative of a Weyl structure $\sigma : \mathcal{G}_0 \rightarrow \mathcal{G}$ measures the failure of the Weyl structure to lift horizontal vectors in \mathcal{G}_0 to horizontal vectors in \mathcal{G} .*

More precisely, for any vector Y in TM and any horizontal lift \hat{Y}_0 in \mathcal{G}_0 for the \mathcal{G}_0 -Weyl connection induced by σ , the vertical part of $\sigma_(\hat{Y}_0)$ for the \mathcal{G} -Weyl connection equals $\mathcal{D}_Y\sigma$.*

Proof. – Obvious corollary of the remarks above. \square

Weyl structures whose tractor-derivative vanishes in every direction were called *normal* by Čap and Slovák [6, Definition 5.1.12].

4. PARABOLIC GEODESICS AS PARALLEL CURVES FOR THE NATURAL CONNECTION

We can now proceed to the main result of this note. Following Čap, Slovák, and Žádník [6, 7], we recall the definition of parabolic geodesics.

Definition 4.1. For any P^0 -invariant subset \mathfrak{a} of $\mathfrak{g}^{<0}$, let

$$\mathcal{U}^{\mathfrak{a}} = \{ \xi \in \mathfrak{g} \setminus \{0\} \mid \exists p \in P \text{ such that } \text{Ad}_{p^{-1}}(\xi) \in \mathfrak{a} \}.$$

If $\mathfrak{a} = \mathfrak{g}^{<0}$, then we shall simply denote $\mathcal{U} = \mathcal{U}^{\mathfrak{g}^{<0}}$.

We recall that a *constant* vector field on \mathcal{G} is a section of $T\mathcal{G}$ of the form $\tilde{\xi} = \omega^{-1}(\xi)$ where ξ is a fixed element of \mathfrak{g} .

Definition 4.2. Let \mathfrak{a} be a P^0 -invariant subset of $\mathfrak{g}^{<0}$. A distinguished curve in a parabolic geometry or *parabolic geodesic* of type \mathfrak{a} is the projection on M of an integral curve in \mathcal{G} of a constant vector field $\tilde{\xi} = \omega^{-1}(\xi)$, where ξ is an element of $\mathcal{U}^{\mathfrak{a}}$ (hence it is not zero).

The most natural choice is $\mathfrak{a} = \mathfrak{g}^{<0}$. For simplicity's sake, we shall only consider this case here and omit the type in the notation.

This definition provides a very general setting encompassing all examples of natural curves used in the most studied examples of parabolic geometries such as, *e.g.*, conformal geodesics of Ferrand [14] in conformal geometries (of any signature), or chains [9, 10, 13] or horizontal curves of Koch [21, 22] in CR geometry. For more details, we refer to Čap and Slovák's book [6]. We can now prove our main result.

Theorem 4.3. *Parabolic geodesics of general type are exactly the curves satisfying the following requirements:*

- (i) *they are parallel for a Weyl structure ;*

- (ii) *the Weyl structure itself is parallel, among the curve, in the direction of the tangent vector of the curve for the natural tractor connection of the bundle of Weyl structures.*

Proof. – Let γ be a parabolic geodesic in the sense of Čap and Slovák, and denote by $\hat{\gamma}$ a lift on \mathcal{G} that is an integral curve of a constant vector field $\tilde{\xi} = \omega^{-1}(\xi)$, for a fixed ξ in \mathfrak{g} . This immediately defines a Weyl structure along γ : indeed, $\hat{\gamma}$ projects itself onto a curve $\hat{\gamma}_0$ in \mathcal{G}_0 , and one may define a section $\sigma : \mathcal{G}_{0|\gamma} \rightarrow \mathcal{G}_\gamma$ as follows: for any e_0 in $\mathcal{G}_{0|\gamma(s)}$, there is p^0 in P^0 such that $e_0 = \hat{\gamma}_0(s)p^0$, and one sets $\sigma(e_0) = \hat{\gamma}(s)p^0$. As an example, we have $\sigma(\hat{\gamma}_0(s)) = \hat{\gamma}(s)$.

One now needs to slightly change the lift: as ξ is not zero and in \mathcal{U} , one may choose p in P such that $\text{Ad}_{p^{-1}}(\xi)$ belongs to $\mathfrak{g}^{<0} \setminus \{0\}$. Thus $\hat{\gamma}p$ is another lift of γ which is an integral curve of $\text{Ad}_{p^{-1}}(\xi)$. This shows that one may always suppose that ξ belongs to $\mathfrak{g}^{<0}$. We shall now make this choice and denote by $\hat{\gamma}$ a lift of γ satisfying this extra property, and use the Weyl structure induced by this choice of the lift.

At $\hat{\gamma}(s)$, the horizontal space in \mathcal{G} for the so defined Weyl structure is $\omega_{\hat{\gamma}(s)}^{-1}(\mathfrak{g}^{<0})$. Recall now that any vector on M is an element of $\mathcal{G} \times_P \mathfrak{g}/\mathfrak{p}$: more precisely, any $[e, \bar{\eta}]$ in $\mathcal{G} \times_P \mathfrak{g}/\mathfrak{p}$ corresponds to a vector

$$X = \pi_* \circ \omega_e^{-1}(\eta),$$

where η is any lift of $\bar{\eta}$ in \mathfrak{g} . Thus, the vector field $\gamma'(s)$ may be thought as $[\hat{\gamma}(s), [\xi]_{\mathfrak{g}/\mathfrak{p}}]$ and its covariant derivative in its own direction with respect to the previously defined Weyl structure is given by the usual computation

$$[\hat{\gamma}(s), \text{ad}(\theta(\hat{\gamma}'(s)))([\xi]_{\mathfrak{g}/\mathfrak{p}})],$$

where θ is the connection 1-form of the Weyl structure. But since $\omega(\hat{\gamma}')$ belongs to the space $\mathfrak{g}^{<0}$, the curve $\hat{\gamma}$ is an horizontal curve in \mathcal{G} and this implies that $\theta(\hat{\gamma}'(s)) = 0$. Thus the curve γ on M is a geodesic for this choice of Weyl structure.

We now prove that the Weyl structure itself, seen as a section of \mathcal{W} , is parallel in the direction of γ' with respect to the tractor connection. Along γ , the section τ associated to the Weyl structure is given by

$$\tau_\gamma(\gamma(s)) = [\hat{\gamma}(s), [1]_{P/P^0}],$$

or, seen as a section of the Moebius bundle $\mathcal{G} \times_P G/P^-$,

$$\tau_\gamma(\gamma(s)) = [\hat{\gamma}(s), [1]_{G/P^-}].$$

Lifting this curve to $\mathcal{G} \times G/P^-$, we can compute its tangent vector:

$$(\hat{\gamma}'(s), 0) = (\omega_e^{-1}(\xi), 0).$$

Since $\xi \in \mathfrak{g}^{<0}$,

$$\frac{d}{dt} [\exp(t\xi)1]_{G/P^-} = 0$$

and it follows that τ is a horizontal section along γ .

Conversely, let γ be a curve on M such that there exists a Weyl structure for which the curve γ is a geodesic and which is tractor-parallel along γ in the direction of γ' . Let $\hat{\gamma}_0(s)$ any local horizontal lift of $\gamma(s)$ to \mathcal{G}_0 . From the very definition of the Weyl connection on \mathcal{G}_0 (see above), the fact that $\hat{\gamma}'_0(s)$ is horizontal in \mathcal{G}_0 is equivalent to the fact that $\omega^0(\hat{\gamma}'_0(s)) = 0$, and since $\gamma'(s)$ is represented in $TM = \mathcal{G}_0 \times_{p^0} \mathfrak{g}^{<0} = \mathcal{G}_0 \times_{p^0} \mathfrak{g}/\mathfrak{p}$ by

$$\gamma'(s) = [e_0, \omega^{<0}(\hat{\gamma}'_0(s)) + \mathfrak{p}],$$

one deduces that $\omega^{<0}(\hat{\gamma}'_0(s))$ is constant.

We now define $\hat{\gamma}(s) = \sigma \circ \hat{\gamma}_0(s)$. This is a lift of $\gamma(s)$ on \mathcal{G} , and Lemma 3.5 shows that $\hat{\gamma}'(s)$ is horizontal for the Weyl connection. Thus, $\omega^+(\hat{\gamma}'(s))$ vanishes, and

$$\omega(\hat{\gamma}'(s)) = \omega^{<0}(\hat{\gamma}'(s))$$

is a constant element of \mathfrak{g} . Consequently, $\gamma(s)$ is a parabolic geodesic. \square

5. EXAMPLES

We shall consider here three classical examples which historically served as compelling motivations for the study of general parabolic geometries: (pseudo-)conformal geometry, projective classes of torsion free connections, and Cauchy-Riemann structures. Of course the results are already known in all these cases, but this is precisely the reason why they may be considered as interesting settings to test the result of this note. Moreover, one may choose the group G to be a linear group in those three cases, and this simplifies a lot the task of computing the Cartan connection as one may use the natural representation to shift from the principal bundle viewpoint into a vector bundle approach which is easier to cope with.

5.1. Conformal geodesics. These are the oldest known examples of distinguished curves in a parabolic geometry. They appear in the 1982 paper by Ferrand [14], but Bailey and Eastwood [1] mention that they might have been known earlier to Penrose [23, 24]. Some authors moreover claim that they are not-so-hidden in Cartan's works on conformal geometry. We shall here show that our approach gives the usual equations for conformal geodesics as given by Bailey and Eastwood [1], Ferrand [15], or Gauduchon [19].

If $(M, [g])$ is a conformal manifold of signature (p, q) ($n = p + q \geq 3$), one may take $G = O(p + 1, q + 1)$ and P is the isotropy subgroup of an isotropic ray (half-line) in \mathbb{R}^{p+q+2} . Thus $\mathfrak{g} = \mathfrak{g}^{-1} \oplus \mathfrak{g}^0 \oplus \mathfrak{g}^1$ where $\mathfrak{g}^{-1} \simeq \mathbb{R}^n$, $\mathfrak{g}^0 \simeq \mathfrak{co}(p, q)$, $\mathfrak{g}^1 \simeq (\mathbb{R}^n)^*$ and the subgroup P^0 of P preserving the grading is $P^0 = CO(p, q) = \mathbb{R}_{>0} \times O(p, q)$.

The Moebius bundle is more easily identified with the help of the so-called tractor or Cartan bundle [2, 5, 19], and the existence of such a bundle motivated our choice of groups, see Graham and Willse [18] for subtleties in this issue. The tractor bundle is the vector bundle $\mathcal{G} \times_P \mathbb{R}^{p+q+2}$. As it is

induced by the (faithful) standard representation of G , the tractor connection may rather be defined on this vector bundle rather than on \mathcal{G} . For this purpose, we notice that there is a natural filtration

$$0 \subset \Lambda \subset \Lambda^\perp \subset \mathbb{T},$$

where Λ is an oriented isotropic line bundle and Λ^\perp is its orthogonal subspace for the natural pseudo-Riemannian metric of \mathbb{T} . Any choice of metric g in $[g]$ induces a Weyl structure, which turns this filtration into a graduation

$$\mathbb{T} \simeq_g \mathbf{1} \oplus T^*M \oplus \mathbf{1},$$

where each $\mathbf{1}$ denotes a trivial line bundle, the first one being nothing but the already introduced Λ . Under this isomorphism, the tractor connection on a section $u = (\lambda, \alpha, \mu)$ of \mathbb{T} is explicitly computed as

$$\mathcal{D}_X u = \begin{pmatrix} d\lambda(X) + S^g(\alpha^\sharp, X) \\ D_X^g \alpha - \lambda g(X, \cdot) + \mu S^g(X, \cdot) \\ d\mu(X) - \alpha(X) \end{pmatrix},$$

where $S^g = \frac{1}{n-2} \left(\text{Ric}^g - \frac{\text{Scal}^g}{2(n-1)} g \right)$ is the Schouten tensor (sometimes called the Rho-tensor) of the metric g and X is any tangent vector. The bundle \mathbb{T} is endowed with a pseudo-Riemannian metric of signature $(p+1, q+1)$ given by $\mathbf{H}(u, u) = g^{-1}(\alpha, \alpha) - 2\lambda\mu$ and the P -bundle \mathcal{G} is the subbundle of the orthonormal frame bundle of \mathbb{T} whose elements are isometries from \mathbb{R}^{p+q+2} to \mathbb{T} sending a fixed isotropic ray $\mathbb{R}_{>0}e_0$ in \mathbb{R}^{p+q+2} to the ray $\Lambda_{>0}$.

One may now take P^- to be the subgroup of $G = O(p+1, q+1)$ fixing an isotropic ray $\mathbb{R}_{>0}e_\infty$, where e_∞ is any vector in \mathbb{R}^{p+q+2} such that $\langle e_0, e_\infty \rangle = 1$. One has then $P \cap P^- = P^0$ as requested and one concludes that the Moebius bundle $\mathcal{S} = \mathcal{G} \times_P G/P^-$ is the bundle of isotropic rays in \mathbb{T} .

When the signature is Riemannian ($q = 0$), each fiber of the Moebius bundle consists in two copies of the Moebius sphere \mathbb{S}^n , seen as the set of isotropic lines in \mathbb{R}^{n+2} with Lorentzian signature $(+\dots+)$. The bundle of Weyl structures \mathcal{W} is then identified to $\mathcal{S}^+ \setminus \{\Lambda_{>0}\}$, where \mathcal{S}^+ is the connected component of \mathcal{S} containing $\Lambda_{>0}$. For arbitrary p and q , each fiber of \mathcal{W} is the image of the embedding of T^*M into \mathcal{S} defined by $\beta \mapsto \mathbb{R} \left(\frac{1}{2}g^{-1}(\beta, \beta), \beta, 1 \right)$, which reproduces the usual conformal embedding of the flat pseudo-Riemannian space \mathbb{R}^{p+q} of signature (p, q) into the set of isotropic rays of the space \mathbb{R}^{p+q+2} with signature $(p+1, q+1)$. This embedding will also be used below to compute explicitly the tractor derivative of a Weyl structure, as it identifies the cotangent bundle T^*M to the vertical tangent space $\mathcal{G} \times_P (g/p^-) = \mathcal{G} \times_P g^{>0}$ of the Moebius bundle.

Weyl structures may be retrieved in the following way: a choice of metric g fixes a P^0 -subbundle (hence a background Weyl structure) given by the frames adapted to the splitting $\mathbb{T} \simeq_g (\mathbf{1} \oplus \mathbf{1}) \oplus^\perp T^*M$ where the trivial line factors are the same as the ones given above. Any other Weyl structure is induced in this trivialization by the choice of a field of 1-forms α . This

yields another splitting $\mathbb{T} \simeq (\mathbf{1} \oplus \mathbb{R}w) \oplus^\perp T^*M$ where the first $\mathbf{1}$ is again the trivial line bundle Λ but the second trivial line is now generated by

$$w = \left(\frac{1}{2}g^{-1}(\alpha, \alpha), -\alpha, 1\right)$$

where α belongs to T^*M (the sign in front of the second factor being chosen here to simplify further computations). The bundle of orthonormal frames adapted to this decomposition is again a P^0 -subbundle of \mathcal{G} which is the image in \mathcal{G} of a new Weyl structure $\sigma : \mathcal{G}_0 \rightarrow \mathcal{G}$.

To make the picture complete, we note that a field of 1-forms may be seen as a section of the bundle in nilpotent groups $\mathcal{G} \times_P P^{>0}$: the filtration of the Lie algebra \mathfrak{g} has depth 1 in the conformal case, which means that one has $\mathfrak{g} = \mathfrak{g}^{-1} \oplus \mathfrak{g}^0 \oplus \mathfrak{g}^1$ and \mathfrak{g}^1 is the abelian algebra \mathbb{R}^{p+q} . Hence, $\mathcal{G} \times_P P^{>0}$ is nothing but the bundle in abelian groups $T^*M = \mathcal{G} \times_P \mathfrak{g}^1$ which we already know as a vector bundle. As a result, we recover the simply transitive action of the bundle of nilpotent groups on the bundle of Weyl structures discussed in section 2. In the conformal case, this action turns \mathcal{W} into an bundle in affine spaces whose associated vector bundle is T^*M .

It now remains to compute explicitly the Weyl connection on the tangent bundle, but this is well-known. Starting with a metric g in the conformal class, the Weyl connection induced by the Weyl structure translated from g by a 1-form α is

$$D_X^\alpha Y = D_X^g Y + \alpha(X)Y + \alpha(Y)X - g(X, Y)\alpha^\sharp$$

(if α had been chosen above rather than $-\alpha$ in the formula yielding the embedding of T^*M into \mathcal{S} , then all occurrences of α here should be changed to $-\alpha$). Thus one may fix a background Riemannian metric g_0 (and its Levi-Civita connection D^0), and using either the fact that the projection into the $\mathfrak{g}^{>0}$ -part of the Cartan connection of conformal geometry is the Schouten tensor or the formulas given above for the tractor connection on \mathbb{T} , one easily deduces from the definition of the tractor connection given above that the equations for a conformal geodesic are

$$\begin{cases} 0 = D_{\gamma'}^\alpha \gamma' = D_{\gamma'}^0 \gamma' + 2\alpha(\gamma')\gamma' - g_0(\gamma', \gamma')\alpha^\sharp, \\ 0 = S^{D^\alpha}(\gamma', \cdot) = S^{g_0}(\gamma', \cdot) - D_{\gamma'}^0 \alpha - \frac{1}{2}g^{-1}(\alpha, \alpha)(\gamma')^b + \alpha(\gamma')\alpha. \end{cases}$$

It is easily checked that either $g_0(\gamma', \gamma') = 0$ at time $s = 0$ and it remains so along the solution curve, or $g_0(\gamma', \gamma')$ never vanishes. In the first case, the first equation is turned into $D_{\gamma'}^0 \gamma' + 2\alpha(\gamma')\gamma' = 0$, which is the (unparametrised) geodesic equation for any Weyl structure, and one recovers the set of null geodesic curves (which forms a conformally invariant set of curves). In the second case, one may always solve the first equation for α and get

$$\alpha = \frac{1}{g_0(\gamma', \gamma')} \left((D_{\gamma'}^0 \gamma')^b - 2 \frac{g_0(D_{\gamma'}^0 \gamma', \gamma')}{g_0(\gamma', \gamma')} (\gamma')^b \right).$$

Thus, the parabolic geodesics are exactly the solutions of the following well-known third-order equation [1, 14, 19]

$$\gamma''' + \frac{3}{2} \frac{|\gamma''|_{g_0}^2}{|\gamma'|_{g_0}^2} \gamma' - 3 \frac{g_0(\gamma'', \gamma')}{|\gamma'|_{g_0}^2} \gamma'' - |\gamma'|_{g_0}^2 S^{g_0}(\gamma', \cdot)^\# + S^{g_0}(\gamma', \gamma') \gamma' = 0,$$

where we have of course denoted $D_{\gamma'}^0 \gamma' = \gamma''$ and $D_{\gamma'}^0 (D_{\gamma'}^0 \gamma') = \gamma'''$.

5.2. Great circles in projective geometry. We now consider the geometric structure given by a projective class of linear connections, *i.e.* a class of torsion-free connections on TM sharing the same family of unparametrized geodesics. Of course, this example is in a sense vacuous for our purposes as we expect that the family of (unparametrized) parabolic geodesics will be the same as the common family of geodesics for all connections in the projective class. It is however interesting to see how they explicitly appear in the computations.

Following for instance [2] or [6, chapter 4], we may now take the group $G = SL(n+1, \mathbb{R})$ and the isotropy subgroup P of a fixed ray $(\mathbb{R}_{>0} e_0, \text{ say})$ in \mathbb{R}^{n+1} , where n is the dimension of the underlying manifold. The Lie algebra \mathfrak{g} splits as $\mathfrak{g} = \mathfrak{g}^{-1} \oplus \mathfrak{g}^0 \oplus \mathfrak{g}^1$ where one has $\mathfrak{g}^{-1} \simeq \mathbb{R}^n$, $\mathfrak{g}^0 \simeq \mathfrak{gl}(n, \mathbb{R})$ and $\mathfrak{g}^1 \simeq (\mathbb{R}^n)^*$. The group P^0 preserving the grading is isomorphic to $GL^+(n, \mathbb{R})$. Fixing a basis (e_0, \dots, e_n) of \mathbb{R}^{n+1} , one takes P^- to be the subgroup of G preserving the hyperplane generated by (e_1, \dots, e_n) together with the orientation induced by the n -form induced from the determinant of \mathbb{R}^{n+1} by the choice of e_0 .

There is again a tractor bundle $\mathbb{T} = \mathcal{G} \times_P \mathbb{R}^{n+1}$, which contains a natural oriented line bundle $\Lambda \subset \mathbb{T}$, see [2]. Choosing a Weyl structure amounts to choosing an hyperplane in \mathbb{T} transverse to the line Λ , and this induces a splitting $\mathbb{T} \simeq \Lambda \oplus (TM \otimes \Lambda)$ reproducing the reduction of the structure group from P to P^0 . Given such a splitting (*i.e.* given a background Weyl structure), choosing another Weyl structure then amounts to choosing a 1-form $u \in (\mathbb{R}^n)^*$ since the induced hyperplane is the kernel of the map:

$$(\lambda, \xi) \in \Lambda \oplus (TM \otimes \Lambda) \mapsto \ell(\lambda) - \ell \otimes u(\xi).$$

Equivalently, this defines in $\mathbb{T}^* \simeq (T^*M \otimes \Lambda^*) \oplus \Lambda^*$ the ray whose elements are $(-u \otimes \ell, \ell)$, where ℓ runs through all positive elements in Λ^* .

On the background manifold, the torsion-free connections on TM are the same as the set of connections induced by Weyl structures: two Weyl structures differ one from another by a 1-form, and the relation between the Weyl connection ∇^u induced by the choice of a 1-form u and a background Weyl connection ∇ is given by

$$\nabla_X^u Y = \nabla_X Y + u(X)Y + u(Y)X, \quad X \in TM, Y \in \Gamma(TM).$$

The tractor connection has been computed for instance in [2]. To state the result, one defines a background Weyl structure (*i.e.* a connection ∇ in the projective class) and one denotes by \mathbb{P} its modified Ricci tensor (or

Schouten tensor) [2, p. 1209]. As explained there, it is always possible to choose ∇ such that \mathbf{P} is symmetric, and we shall indeed do so. It is then a simple task to check that a curve γ is a parabolic geodesic if it satisfies the equations

$$\begin{cases} \nabla_{\gamma'}\gamma' + 2u(\gamma')\gamma' = 0, \\ \nabla_{\gamma'}u - \mathbf{P}(\gamma', \cdot) - u(\gamma')u = 0. \end{cases}$$

The first equation shows that any solution is an (*a priori* non affinely parametrized) geodesic for ∇ , and it is again easy to check that one may always choose the parametrization such that the second equation is satisfied. This substantiate our previous claim that the set of parabolic geodesics here is exactly the same as the set of *great circles*, or common geodesics to all connections in the projective class.

5.3. Parabolic geodesics in strictly pseudo-convex CR geometry. The situation in the complex case of positive definite signature is very similar to the conformal case: if $n = 2m + 1$ is the dimension of the underlying CR manifold, one may take the semi-simple group $G = SU(m + 1, 1)$ (formally, taking this group amounts to choosing a slightly refined description of CR geometry as a parabolic geometry, since one also assumes existence of a $(m + 2)$ -root of the CR-canonical bundle) and P and P^- are the isotropy groups of the complex lines $\mathbb{C}e_0$ and $\mathbb{C}e_\infty$ in \mathbb{C}^{m+2} , where we fixed a basis $(e_0, e_1, \dots, e_m, e_\infty)$ and the hermitian form given by $\langle e_0, e_0 \rangle = 0$, $\langle e_\infty, e_\infty \rangle = 0$, $\langle e_0, e_\infty \rangle = \langle e_\infty, e_0 \rangle = 1$, and $\langle e_i, e_j \rangle = \delta_{ij}$ (Kronecker symbol) whenever (i, j) belongs to $\{0, \dots, m, \infty\}^2 \setminus \{(0, \infty), (\infty, 0)\}$; as a result, $P \cap P^- = P^0$.

As in the real case, it is easier to work with the tractor vector bundle \mathbb{T} , see [17, 20]. If (M, H, J) is a strictly pseudo-convex CR manifold of dimension $n = 2m + 1$, the tractor bundle is a rank $(m + 2)$ complex vector bundle endowed with a hermitian form of signature $(m + 1, 1)$. There is again an exact sequence

$$0 \subset \Lambda \subset \Lambda^\perp \subset \mathbb{T},$$

where Λ is an isotropic complex line. Choosing a contact form on M whose kernel is the contact distribution H yields a splitting

$$\mathbb{T} \simeq_g \mathbf{1} \oplus (H^*M)^{1,0} \oplus \mathbf{1},$$

where the symbol $\mathbf{1}$ now denotes a trivial complex line bundle and $(H^*M)^{1,0}$ is the space of 1-forms of type $(1, 0)$ on the contact hyperplane H .

The description of Weyl structures is formally very similar to that given in the conformal case: the Moebius bundle \mathcal{S} is the bundle of isotropic complex lines in \mathbb{T} , the bundle of Weyl structures is $\mathcal{S} \setminus \{\Lambda\}$, and much of the considerations given above also apply here *mutatis mutandis*. For instance, once a reference contact form θ has been fixed, one may describe \mathcal{W} as the image of the embedding $\alpha = \alpha_{|H} + \alpha_0\theta \mapsto (\frac{1}{2}|\alpha_{|H}^{1,0}|^2 + i\alpha_0, \alpha_{|H}^{1,0}, 1)$ of T^*M in \mathcal{S} . This reproduces the standard embedding of the Heisenberg

algebra in G/P^- also gives an identification between the cotangent bundle and the vertical bundle of the Moebius bundle.

Explicit equations for a curve γ to be a parabolic geodesic are much longer to obtain in this case than in the previous ones, so we shall skip leave the detailed computations to the reader. Let us just indicate that explicit forms of the tractor connection can be found in [17, formula (3.3)] or in [20, formula (5.11)] (both with respect to the Tanaka-Webster connection), thus leading with some effort to the equations asserting that the Weyl structure associated to a choice of 1-form α is tractor parallel along the curve γ . To get the full system of equations for a parabolic geodesic, one must complement by the equation ensuring that γ is the geodesic for the Weyl structure defined by the 1-form α , and this can be calculated by using for instance [3, §4] which easily leads to explicit formulas for Weyl covariant derivative on the tangent bundle.

REFERENCES

- [1] T. N. Bailey and M. G. Eastwood, *Conformal circles and parametrizations of curves in conformal manifolds*, Proc. Amer. Math. Soc. **108** (1990), 215–221.
- [2] T. N. Bailey, M. G. Eastwood and A.R. Gover, *Thomas' structure bundle for conformal, projective, and related structures*, Rocky Mountain J. Math. **24** (1994), 61–103.
- [3] D. M. J. Calderbank, T. Diemer, and V. Souček, *Ricci-corrected derivatives and invariant differential operators*, Diff. Geom. Appl. **23** (2005), 149–175.
- [4] A. Čap and H. Schichl, *Parabolic geometries and canonical Cartan connection*, Hokkaido Math. J. **29** (2000), 453–505.
- [5] A. Čap and J. Slovák, *Weyl structures for parabolic geometries*, Math. Scand. **93** (2003), 53–90.
- [6] A. Čap and J. Slovák, *Parabolic geometries, I: background and general theory*, Math. Surv. Monogr., vol. 154, Amer. Math. Soc., Providence, RI, 2009.
- [7] A. Čap, J. Slovák, and V. Žádník, *On distinguished curves in parabolic geometries*, Transf. Groups **9** (2004), 143–166.
- [8] E. Cartan, *Les espaces à connexion conforme*, Ann. Soc. Pol. Math. **2** (1923), 171–221, *Œuvres complètes*, t.III.1, Gauthier-Villars, Paris, 1955, 747–797.
- [9] E. Cartan, *Sur la géométrie pseudo-conforme des hypersurfaces de deux variables complexes, I*, Ann. Mat. Pura Appl. **11** (1932), 17–90.
- [10] E. Cartan, *Sur la géométrie pseudo-conforme des hypersurfaces de deux variables complexes, II*, Ann. Sc. Norm. Sup. Pisa, Ser. II, **1** (1932), 333–354.
- [11] S.-S. Chern and J. K. Moser, *Real hypersurfaces in complex manifolds*, Acta Math. **133** (1974), 219–271.
- [12] B. Doubrov and I. Zelenko, *Geometry of curves in parabolic homogeneous spaces*, preprint, ArXiv:1110.0226 [math.DG], 2011.
- [13] C. L. Fefferman, *Monge-Ampère equations, the Bergman kernel, and the geometry of pseudoconvex domains*, Annals of Math. **103** (1976), 395–416, *Erratum*, Annals of Math. **104** (1976), 393–394.
- [14] J. Ferrand, *Les géodésiques des structures conformes*, C. R. Acad. Sci. Paris Sér. I Math. **294** (1982), 629–632.
- [15] J. Ferrand, *Un invariant conforme lié aux géodésiques conformes*, Complex analysis, Fifth Romanian-Finnish seminar, Part 1 (Bucharest, 1981), Lecture Notes in Math., vol. 1013, Springer, Berlin, 1983, 76–86.

- [16] C. Francès, *Sur le groupe d'automorphismes des géométries paraboliques de rang 1*, Ann. Sci. École Norm. Sup. **40** (2007), 741–764.
- [17] A. R. Gover and C. R. Graham, *CR invariant powers of the sub-Laplacian*, J. Reine Angew. Math. **583** (2005), 1–27.
- [18] C. R. Graham and T. Willse, *Subtleties concerning conformal tractor bundles*, preprint, ArXiv:1201.2670 [math.DG], 2012.
- [19] P. Gauduchon, *Connexion canonique et structures de Weyl en géométrie conforme*, unpublished lecture notes, 1990.
- [20] M. Herzlich, *The canonical Cartan bundle and connection in CR geometry*, Math. Proc. Camb. Phil. Soc. **146** (2009), 415–434.
- [21] L. K. Koch, *Chains in CR manifolds and Lorentz geometry*, Trans. Amer. Math. Soc. **308** (1988), 827–841.
- [22] L. K. Koch, *Chains, null-chains and CR geometry*, Trans. Amer. Math. Soc. **338** (1993), 245–261.
- [23] R. Penrose and W. Rindler, *Spinors and space-time, 1: two-spinor calculus and relativistic fields*, Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1987.
- [24] R. Penrose and W. Rindler, *Spinors and space-time, 2: spinor and twistor methods in space-time geometry*, Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1988.

INSTITUT DE MATHÉMATIQUES ET DE MODÉLISATION DE MONTPELLIER, CNRS & UNIVERSITÉ
MONTPELLIER 2

E-mail address: `herzlich@math.univ-montp2.fr`