

Heteromerization of human cytomegalovirus encoded chemokine receptors

Pia Tschische, Kenjiro Tadagaki, Maud Kamal, Ralf Jockers, & Maria

Waldhoer

► To cite this version:

Pia Tschische, Kenjiro Tadagaki, Maud Kamal, Ralf Jockers, & Maria Waldhoer. Heteromerization of human cytomegalovirus encoded chemokine receptors. Biochemical Pharmacology, 2011, 82 (6), pp.610. 10.1016/j.bcp.2011.06.009 . hal-00722841

HAL Id: hal-00722841 https://hal.science/hal-00722841

Submitted on 5 Aug 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Accepted date:

Title: Heteromerization of human cytomegalovirus encoded chemokine receptors

Authors: Pia Tschische, Kenjiro Tadagaki, Maud Kamal, Ralf Jockers, & Maria Waldhoer

| PII: | S0006-2952(11)00374-1 |
|----------------|-------------------------------|
| DOI: | doi:10.1016/j.bcp.2011.06.009 |
| Reference: | BCP 10933 |
| To appear in: | ВСР |
| Received date: | 13-4-2011 |
| Revised date: | 28-5-2011 |

2-6-2011



Please cite this article as: Tschische P, Tadagaki K, Kamal M, Jockers R, Waldhoer M, Heteromerization of human cytomegalovirus encoded chemokine receptors, *Biochemical Pharmacology* (2010), doi:10.1016/j.bcp.2011.06.009

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Heteromerization of human cytomegalovirus encoded chemokine receptors Pia Tschische^{a,e,*}, Kenjiro Tadagaki^{b,c,d}, Maud Kamal^{b,c,d}, Ralf Jockers^{b,c,d} & Maria Waldhoer^a ^a Institute for Experimental and Clinical Pharmacology, Medical University of Graz, Universitaetsplatz 4, 8010 Graz, Austria ^b Inserm, U1016, Institut Cochin, 22 Rue Méchain, 75014 Paris, France ^c CNRS UMR 8104, Paris, France ^d Université Paris Descartes, Paris, France * Corresponding author: Pia Tschische, PhD ^e Present address: Ernest Gallo Clinic and Research Center University of California San Francisco 5858 Horton Street, Suite 200 Emeryville, CA 94608, USA Phone: +1 510 985 3142 E-mail: ptschische@gallo.ucsf.edu E-mail addresses of authors: kenjiro.tadagaki@inserm.fr maud.kamal@inserm.fr ralf.jockers@inserm.fr maria.waldhoer@medunigraz.at

Keywords: Human cytomegalovirus, chemokine receptors, signaling, post-endocytic sorting, heteromerization, G-protein coupled receptor

Category: Inflammation and Immunopharmacology

coordination of the second sec

Abstract

Human cytomegalovirus (HCMV) is a widespread pathogen that infects up to 80% of the human population and causes severe complications in immunocompromised patients. HCMV expresses four seven transmembrane (7TM) spanning/G protein-coupled receptors (GPCRs) – US28, US27, UL33 and UL78 – that show close homology to human chemokine receptors. While US28 was shown to bind several chemokines and to constitutively activate multiple signaling cascades, the function(s) of US27, UL33 and UL78 in the viral life cycle have not yet been identified. Here we investigated the possible interaction/heteromerization of US27, UL33 and UL78 with US28 and the functional consequences thereof. We provide evidence that these receptors not only co-localize, but also heteromerize with US28 *in vitro*. While the constitutive activation of the US28-mediated Gaq/phospholipase C pathway was not affected by receptor heteromerization, UL33 and UL78 were able to silence US28mediated activation of the transcription factor NF- κ B. Summarized, we provide evidence that these orphan viral receptors have an important regulatory capacity on the function of US28 and as a consequence, may ultimately impact on the viral life cycle of HCMV.

A COX

1. Introduction

Human cytomegalovirus (HCMV), also known as human herpesvirus 5 (HHV-5), is a large and highly species-specific virus that belongs to the family of beta-herpesviruses. Infection with HCMV is asymptomatic and rarely causes complications in immunocompetent hosts. In immunosuppressed hosts, however, HCMV infection causes severe complications, including graft rejection or systemic infections [1-4]. HCMV expresses four viral seven transmembrane (7TM) spanning/G protein-coupled receptors (GPCRs), i.e. UL33 (<u>u</u>nique <u>long 33</u>), UL78, US27 (<u>u</u>nique <u>short 27</u>) and US28, which show high homology to human chemokine receptors [5, 6].

US28 is the best characterized HCMV encoded 7TM/GPCR and was suggested to i) be important for the viral life cycle by enhancing cell-cell fusion, thus promoting viral spread [7-10] and ii) to activate the immediate early HCMV promoter, which generally leads to transactivation of other viral genes [11]. US28 is constitutively active – i.e. transduces signals in a ligand-independent way. The receptor activates the Gaq/phospholipase C (PLC) pathway and induces several transcription factors, such as NF- κ B (nuclear factor kappa B), CREB (cyclic AMP responsive element binding) [12, 13], NFAT (nuclear factor of activated T cells) [14] or SRF (serum response factor) [15]. Moreover, US28 is constitutively endocytosed, which results in the localization of the receptor in the membranes of intracellular organelles, especially late endosomes/lysosomes and multi-vesicular bodies (MVBs) [13, 16, 17], where it has been suggested that the virions of HCMV may be assembled [7, 8].

US27, UL33 and UL78 are still "orphan" receptors, since no endogenous ligands have been identified to date. Of these, UL33 was reported to constitutively induce inositol phosphate (IP) accumulation via coupling to Gaq and Gai/o-proteins [13, 18] and to activate CREB via Gaq, Gas, G $\beta\gamma$ and p38 kinase [13, 18], as well as to co-localize with endosomes

in HCMV infected cells [7, 8]. US27 was shown to be expressed during late HCMV infection states and found to be present on HCMV virions [7, 19, 20]. UL78 shares only very limited homology with endogenous chemokine receptors [21] and was shown to be dispensable for viral replication [22].

Orphan 7TM/GPCRs have recently been shown to influence the signal activity of nonorphan receptors through heteromerization. One prominent example is the orphan receptor GPR50, which was reported to antagonize the functions of the melatonin MT_1 receptor. Heteromerization of GPR50 with the MT_1 receptor prevented the agonist binding and G protein coupling [23].

Here we set out to test whether the orphan viral receptors US27, UL33 and UL78 possibly interact/heteromerize with US28 and whether this had functional consequences on the signaling and/or endocytic properties of US28.

2. Materials and Methods

2.1 Reagents

Mouse anti-Flag M1 and M2 monoclonal antibody, gelatine from bovine skin Type B, Triton X-100, 3, 3', 5, 5'-Tetramethylbenzidine liquid substrate system were purchased from Sigma Aldrich (Austria). Cell culture media (DMEM), foetal bovine serum (FBS), Dulbecco's phosphate buffered saline (DPBS), Lipofectamine 2000, Alexa Fluor 594 nm conjugated IgG1a, 488 nm conjugated IgG2b and 594 nm conjugated donkey anti rabbit IgG were purchased from Invitrogen (Austria). HRP-conjugated anti-mouse antibody was obtained from Jackson Immuno Research (Dianova, Germany). Vectashield mounting medium and Vectastain ABC Kit were purchased from Vector Laboratories (Szabo-Scandic, Austria). The white 96-well culture plates and the Steadylite Plus Assay Kit, high sensitivity luminescence

reporter gene assay system, [³H]myo-inositol and EasyTag ³⁵S-methionine were purchased from PerkinElmer (Austria). Yttrium silicate (YSi) SPA beads were from GE Healthcare (Austria). NaOH, KCl and CaCl₂*2H₂O were obtained from Merck (Austria), NaCl, Tris and Formaldehyde were from Roth (Lactan, Austria). Coelenterazine-h powder (1 mg) was purchased from Interchim (France). Monoclonal rat anti-HA and monoclonal mouse anti-GFP antibodies were purchased from Roche Diagnostics (France). Monoclonal rabbit anti-HA antibody was purchased from Cell signaling (France). Polyclonal rabbit anti-Flag antibody, protein G sepharose beads and poly-L-lysine were obtained from Sigma Aldrich (France). Phusion High-Fidelity DNA Polymerase was purchased from Finnzymes (France). IRdye 800-conjugated anti-rat, IRdye 680-conjugated anti-rabbit, IRdye680-conjugated anti-mouse antibodies were purchased from Thermo Scientific (France). JetPEI transfection reagent was obtained from Polyplus Transfection (France).

2.2 DNA constructs

A Flag-tagged version of US28 (Flag-US28) was cloned by using the BamHI and XbaI restriction sites in the SS-Flag pcDNA3.1 zeo (+) vector (16). The cDNA encoding UL78 was kindly provided by Dr. Detlef Michel (University of Ulm, Germany) and was cloned in the SS-Flag pcDNA3.1 zeo (+) and SS-HA pcDNA3.1 zeo (+) vectors using BamHI and EcoRV restriction sites. UL33 (exon 1+2) was cloned in the SS-HA pcDNA3.1 zeo (+) vector using BamHI and XhoI restriction sites. A HA-tagged version of US27 (exon 2) was cloned by using the BamHI and EcoRV restriction sites in SS-HA pcDNA3.1 zeo (+) vector. Flag-US28-YFP, HA-US28-Rluc, HA-UL33, HA-UL33-Rluc, Flag-UL78 and HA-UL78-Rluc constructs were obtained using the latter cDNAs as templates and the Phusion High-Fidelity DNA Polymerase. The Flag or HA tags were inserted after the ATG in frame of the US28, UL33 or UL78 sequences. Flag-US28, HA-US28, HA-UL33, Flag-UL78 and HA-UL78 were amplified by PCR. The amplified products were subcloned directly into a pCR2.1 vector

(Invitrogen, CA). After sequencing, the fragments were digested with the restriction enzymes HindIII and BamHI and subcloned into pcDNA3.1 vector or pcDNA3.1 vector containing either YFP or Rluc sequences. The vasopressin V2-YFP fusion protein has been described previously [24]. All DNA constructs were verified by sequencing.

2.3 Antibody feeding experiments

HEK293 cells were grown on gelatin coated coverslips and transiently transfected with Flag-US28 (1.5 μg/coverslip) in combination with HA-UL33 (1.5 μg/coverslip), HA-UL78 (1.5 μg/coverslip) or HA-US27 (1.5 μg/coverslip) using Lipofectamine 2000. 24 hours posttransfection, antibody feeding experiments were conducted essentially as previously described [25]. In brief, living cells were incubated with anti-Flag M1 (1:500) and anti-HA HA11 (1:500) antibodies for 45 minutes, fixed with 3.7 % formaldehyde in PBS and permeabilized in blotto (50 mM Tris-HCl, pH 7.5, 1 mM CaCl₂, 0.2 % Triton X-100 and 3 % milk). Subsequently, the cells were stained with the subtype-selective antibodies Alexa Fluor 488conjugated IgG2b against the Flag-tag (1:1000, 20 minutes) and Alexa Fluor 594-conjugated IgG1a against the HA-tag (1:1000, 20 minutes). Finally, cells were mounted with Vecta shield mounting medium and analysed using a Olympus IX70 fluorescence microscope.

2.4 Immunoprecipitation

For co-immunoprecipitation assays, HEK293T cells were seeded in 10 cm dishes and cotransfected with 8 μ g of the indicated plasmids. 48 hours after transfection, cells were washed twice in PBS, and lysed in 1 ml cold lysis buffer (25 mM Tris (pH 7.4), 2 mM EDTA, 10 mM MgCl₂, protease inhibitor cocktail and 1 % Triton). After solubilization for 2 hours at 4 °C, lysates were centrifuged at 12 000 g during 1 hour at 4 °C and the supernatant was collected for the immunoprecipitation assay. Immunoprecipitations were performed using 2 μ g of the indicated antibodies pre-adsorbed on protein G sepharose beads for 2 hours at 4 °C.

Immunoprecipitated proteins were eluted with Laemmli buffer 2X (Tris-HCl pH 6.8, 125 mM, SDS 10 %, Glycerol 10 %, bromophenol blue 1 % and DTT 100 mM) and subjected to 10 % or 7.5 % SDS–PAGE and immunoblotting. Immunoblotting was performed using the indicated antibodies and immunoreactivity was revealed using secondary antibodies coupled to 680 or 800 nm fluorophores using the Odyssey LI-COR infrared fluorescent scanner (ScienceTec).

2.5 BRET (Bioluminescence resonance energy transfer) measurement

HEK293T cells were grown in complete medium (Dulbecco's modified Eagle's medium supplemented with 10 % (v/v) fetal bovine serum, 4.5 g/L glucose, 100 U/mL penicillin, 0.1 mg/mL streptomycin and 1 mM glutamine) (Invitrogen, CA). Transient transfections were performed using JetPEI transfection reagent according to manufacturer's instructions.

For BRET donor saturation curves, HEK293T cells were seeded in 12-well plates and transiently transfected with 200 ng of HA-UL33-Rluc or 100 ng of HA-UL78-Rluc and 100 to 1800 ng of YFP plasmids. For BRET competition assays, HEK293T cells were seeded in 6-well plates. Cells were transiently transfected with (i) 200 ng HA-UL33-Rluc and 1000 ng Flag-US28-YFP in presence or absence of 1800 ng Flag-UL78 or (ii) 100 ng HA-UL78-Rluc and 1000 ng Flag-US28-YFP in presence or absence of 1900 ng HA-UL33 plasmids. 24 hours after transfection, cells were transferred into a 96-well white Optiplate pre-coated with 10 mg/ml poly-L-lysine. Following 24 hours incubation, BRET measurements were conducted. Luminescence and fluorescence were measured simultaneously using the lumino/fluorometer MithrasTM (Berthold) that allows the sequential integration of luminescence signals detected with two filter settings (Rluc filter, 485±10 nm; YFP filter, 530±12.5 nm). Emission signals at 530 nm were divided by emission signals at 485 nm and the difference between this emission ratio (obtained with donor and acceptor fused or co-expressed and that obtained with the donor protein expressed alone) was defined as the BRET ratio. The results were expressed in

milliBRET units (mBU) corresponding to the BRET ratio values multiplied by 1000. Total fluorescence was measured with the fluorometer FusionTM (Packard Instrument Company).

2.6 Reporter gene assays

Transcription factor luciferase assays were performed essentially as described in [13]. In brief, HEK293 cells (30000 cells/well) were seeded on gelatin coated 96-well plates and transiently transfected with 75 ng/well Flag-US28, the cis-reporter plasmid for NF- κ B (50 ng/well) and either pcDNA3.1 (75 ng/well) or the respective receptors (HA-UL33, HA-UL78 or HA-US27; 75 ng/well). Controls were transfected with 150 ng/well of either pcDNA3.1 alone, or pcDNA3.1 (75 ng/well) in combination with UL33, UL78 or US27 (75 ng/well), but in the absence of US28. In parallel to each assay, an additional plate was prepared for ELISA (see below) to control for receptor expression levels. The luciferase reporter gene assays were conducted 24 hours post-transfection according to the manufacturer's guidelines. In brief, cells were washed twice with PBS and the cell number was determined by optical density in a FlexStation II Device (Molecular Probes, Endpoint Measurement). Then, 100 µl/well Steadylight luciferase assay reagent were added to 100 µl/ well PBS. Following a 10 min incubation period, luminescence was measured using a TopCounter Device (TopCount NXT, PerkinElmer).

2.7 Inositol Phosphate (IP) accumulation assay

HEK293 cells were seeded on gelatin coated 96-well plates (30000 cells/well) and transfected with 75 ng/well Flag-US28 and either pcDNA3.1 (75 ng/well) or the respective orphan receptor (HA-UL33, HA-UL78 or HA-US27; 75 ng/well). Controls were transfected with 150ng/well of either pcDNA3.1 alone, or pcDNA3.1 (75 ng/well) in combination with UL33, UL78 or US27 (75 ng/well), but in the absence of US28. In parallel to each assay, an additional plate was prepared for ELISA (see below) to control for receptor expression levels.

24 hours post-transfection the cells were loaded with 1 μ Ci/well [³H]myo-inositol in 100 μ l/well Optimem and incubated overnight at 37 °C and 5 % CO₂. The next day, the labeling medium was aspirated, 100 μ l/well HBSS buffer (including CaCl₂ and MgCl₂) containing 10 mM LiCl were added and the cells were incubated for 45 minutes at 37 °C and 5 % CO₂. The reaction was terminated by aspiration and the cell number was determined by optical density in a FlexStation II Device. Subsequently, cells were lysed with 50 μ l/well 10 mM formic acid for 90 minutes on ice. 40 μ l of the resulting cell extract were transferred to 160 μ l of YSi-SPA beads (12.5 mg/ml) and shaken for 90 minutes at 4°C. The plates were then stored at 4°C and the accumulation of inositol phosphate was counted in a TopCount microplate scintillation counter (TopCount NXT, PerkinElmer) 24 hours later.

2.8 ELISA (enzyme linked immunosorbent assay)

The ELISA assays were performed on the same day as the corresponding signaling experiments (see above). Cells were fixed with 3.7 % formaldehyde in PBS and permeabilized in blotto (50 mM Tris-HCl, pH7.5, 1 mM CaCl₂, 0.2 % Triton X-100 and 3 % milk) for one hour. Next, the cells were incubated with either anti-Flag M1 (1:500) or anti-HA HA11 (1:1000) antibody overnight at 4 °C. Then cells were washed with TBS (25 mM Tris base, 135 mM NaCl, 2.5 mM KCl, 1 mM CaCl₂*2H₂O, pH 7.4) and incubated with HRP-conjugated anti-mouse antibody (1:2500) in blotto (50 mM Tris-HCl, pH 7.5, 1 mM CaCl₂ and 1.5 % milk) for two hours at room temperature. After three washes with TBS, the cell number was determined by optical density in a FlexStation II Device. Then, 75 μ /well TMB (3, 3', 5, 5'-Tetramethylbenzidine) substrate were added and the coloring reaction was stopped by the addition of 50 μ /well 0.5 M sulfuric acid after 2 minutes at room temperature. Receptor levels were measured at the optical density of 450 nm in a BioRad xMark Microplate Spectrophotometer.

2.9 Statistical analysis

Statistical analyses were performed using ANOVA analysis of variance for comparisons between multiple groups, followed by a Bonferroni's post-hoc analysis using GraphPad Prism software.

3. Results

3.1 Co-localization of HCMV encoded chemokine receptors

US28, UL33 and US27 have previously been shown to constitutively internalize and localize in intracellular compartments [7, 13, 16]. Here we set out to investigate a possible colocalization of US28 with the three orphan HCMV encoded receptors US27, UL33 and UL78. Therefore, the respective receptors were transiently co-expressed with US28 in HEK293 cells at a 1:1 ratio and antibody feeding experiments were conducted (see methods). These experiments revealed that US28 co-internalizes and co-localizes with UL33 (Fig. 1A) and UL78 (Fig. 1B) in HEK293 cells. As described before [26], US28 was also found to co-localize with US27 (Fig. 1C).

3.2 Dimerization of US28 with US27, UL33 and UL78

We were able to demonstrate that after constitutive internalization US28 co-localizes with all three orphan receptors in endosomal compartments (Fig. 1). Next, we conducted coimmunoprecipitation (co-IP) experiments to test whether the receptors are not only colocalizing, but furthermore able to physically interact with each other. Therefore, HEK293T cells were transiently co-transfected with Flag-US28-YFP and HA-US28 (A), HA-UL33-Rluc (B), HA-UL78-Rluc (C) or HA-US27 (D). These experiments revealed that US28 not only forms homomers (Fig. 2A), but moreover interacts with all three orphan receptors (Figs. 2B, C & D).

Next, Bioluminescence Resonance Energy Transfer (BRET) assays were performed to confirm the co-IP results. Therefore, UL33 or UL78 were fused at their C-terminus to Renilla luciferase (HA-UL33-Rluc, HA-UL78-Rluc) and used as energy donors. US28 was fused at its C-terminus to YFP (Flag-US28-YFP) and used as the energy acceptor. A possible heteromerization between US28 and US27 could not be validated using the BRET approach, since the co-expression of US27-Rluc led to a significant decrease in US28-YFP expression levels (data not shown).

Donor saturation curves were performed by co-transfecting a fixed amount of HA-UL33-Rluc or HA-UL78-Rluc in the presence of increasing amounts of Flag-US28-YFP in HEK293T cells. The vasopressin V2 receptor YFP (V2-YFP) fusion protein was used as a negative control. A specific interaction between BRET donor and acceptor pairs is reflected by a hyperbolic donor saturation curve, which reaches an asymptote with increasing YFP/Rluc ratios. This is shown for the US28/UL33 couple (Fig. 3A) and the US28/UL78 couple (Fig. 3B), respectively. In contrast, a non-specific interaction, due to random collision, was observed with V2-YFP where a quasi-linear increase in BRET was observed with increasing YFP/Rluc ratios (Fig. 3B). BRET competition assays were then performed to assess, if UL78 and UL33 can disrupt US28/UL33 and US28/UL78 heteromerization, respectively. The YFP/Rluc ratio in the ascending portion of the saturation curve before the plateau was selected and the corresponding amounts of donor and acceptor plasmids were transfected in the presence of saturating amounts of the competitor receptor (Flag-UL78 in the case of HA-UL33-Rluc/Flag-US28-YFP and HA-UL33 in the case of HA-UL78-Rluc/Flag-US28-YFP). The BRET signal was significantly decreased in the presence of the competitor receptor indicating that UL78 (Fig. 3C) and UL33 (Fig. 3D) interfere with US28/UL33 and US28/UL78 heteromerization, respectively.

Summarized, the co-IP and BRET experiments show that US28 forms homomers and is furthermore able to heteromerize with the three viral HCMV encoded receptors US27, UL33 and UL78.

3.3 US28-mediated transcription factor activation, but not G protein-mediated inositol phosphate accumulation, is altered by receptor heteromerization

It has previously been demonstrated that receptor heteromerization alters the trafficking and signaling properties of 7TM/GPCRs [27-29]. Importantly, orphan receptors – such as GPR50 - have been demonstrated to modulate the activity of non-orphan receptors through heteromerization [30].

Hence, we next tested the functional consequences of the heteromerization of US28 with US27, UL33 and UL78. First, the ability of US28 to mediate the Ga_q /phospholipase C (PLC) dependent accumulation of inositol phosphate (IP) was assessed in cells co-expressing the orphan HCMV encoded receptors. US28 constitutively induced IP turnover in the absence (Figs. 4 A, D & G, black bars) and presence (Figs. 4 A, D & G, grey bars) of the other viral receptors. The expression levels of Flag-tagged US28 (Figs. 4 B, E & H) as well as the HA-tagged orphan receptors (Figs. 4 C, F & I) were monitored by ELISA. These results suggest that the orphan receptors had no direct influence on the G protein mediated signaling capacity of US28.

US28 has been reported to activate downstream transcription factors such as NF- κ B, CREB [12, 13], NFAT [14] or SRF [15]. We next tested whether the heteromerization of US28 with the orphan HCMV encoded receptors alters the downstream signal transduction profile of US28. Hence, US28 was co-expressed with the respective orphan receptors and reporter gene assays were performed. Interestingly, the co-expression of UL33 and UL78 were found to significantly decrease the activation of the transcription factor NF- κ B by US28 (Figs. 5A & 5D, compare black bars vs. grey bars). The reduction of US28 signaling capacity

in the presence of UL33 and UL78 was unlikely to be due to altered expression levels of the receptors, since they remained relatively stable under the different experimental conditions (see Figs. 5B, C, E & F). Co-expression of US27 on the other hand, did not alter the signaling capacity of US28 (Fig. 5G compare black bar vs. gray bar).

Summarized, these data suggest that the constitutive signaling activity of US28 is selectively modulated via heteromerization with the orphan HCMV-encoded UL33 and UL78. The activation of the Gaq/PLC/IP3 pathway via US28 was not altered, whereas the activation of the transcription factor NF- κ B by US28 was almost completely blocked by UL33 or UL78, respectively.

4. Discussion

Out of the four HCMV encoded 7TM/GPCRs, US28 is the only receptor that has been shown to bind endogenous chemokines [13, 31, 32]. The other three receptors - US27, UL33 and UL78 - are yet orphan receptors with mainly unknown function [5, 6]. Here we tested whether these orphan receptors – similar to recent reports on other orphan 7TM/GPCRs [[23] - may have ligand-independent functions by modulating the signaling and trafficking capacities of US28.

Antibody feeding experiments revealed that all four viral 7TM/GPCRs are predominantly found in intracellular compartments. This is in accordance with previous data on US27, US28 and UL33 [[7, 33], which have all been described to constitutively internalize. Here we show for the first time that US28 co-localizes with all three orphan receptors in HEK293 cells (Fig. 1).

Heteromerization and thus a direct protein-protein interaction between these HCMV encoded receptors were confirmed by co-IP and BRET experiments (Figs. 2 & 3). To date it is

widely accepted that most, if not all, 7TM/GPCRs are organized in dimeric/oligomeric complexes. The importance of 7TM/GPCR oligomerization is most convincingly illustrated by the functional alterations induced when two different 7TM/GPCRs associate to form heteromeric complexes. Modifications in pharmacological, signaling and trafficking properties have been observed for many 7TM/GPCRs when engaged into heteromers [27-29, 34]. For instance, the chemokine receptor CXCR4 was shown to heteromerize with CCR5 [35, 36], whereby the CCR5/CXCR4 heteromer was causally involved in the modulation of T lymphocyte responses [37]. Moreover, homomerization of CCR5 was shown to prevent infection with HIV-1 [38].

In addition, orphan receptors were shown to be capable of influencing the activity of non-orphan receptors through heteromerization. The orphan receptor GPR50, for example, was reported to antagonize the functions of the melatonin MT_1 receptor via heteromerization. Heteromerization of GPR50 with the MT_1 receptor prevented the agonist binding and G protein coupling [23]. Moreover, it was found that the carboxy-terminus of GPR50 is involved in these inhibitory effects. As opposed to the wild type receptor, heteromerization with a carboxy-terminally deleted GPR50 did not alter MT_1 related functions [23].

To evaluate the functional consequences of viral HCMV receptor heteromerization, the Gaq/PLC/IP pathway and the activation of the downstream transcription factor NF- κ B were monitored. Surprisingly, the IP accumulation was not altered upon co-expression of US28 with the orphan receptors (Fig. 4). On the other hand, the constitutive activation of NF- κ B was almost completely blocked by the co-expression of UL33 (Fig. 5A) and UL78 (Fig. 5D). US27, the HCMV encoded receptor that shares the highest sequence homology with US28, did neither alter the IP accumulation (Fig. 4G), nor the activation of the transcription factor (Fig. 5G).

The NF-κB pathway is known to be an important regulator of the innate and the adaptive immune response, inflammation and cell survival [39]. The major immediate early

promoter (MIEP) region of HCMV, which controls the expression of immediate early (IE) genes, contains four binding sites for the transcription factor NF-kB. The role of these binding sites is not completely established. In some studies, inhibition of NF-kB signaling yielded an impaired viral replication [40, 41], whereas other reports show that the deletion of the binding sites within the MIEP region had no effects [42, 43]. The mechanisms employed by HCMV for the establishment of viral latency and reactivation are still an important source of speculation. In general, HCMV infection and reactivation were shown to yield the enhancement of cellular responses, including the activation of transcription factors, accumulation of cAMP, inositol phosphate hydrolysis and the metabolism of arachidonic acid [44, 45]. For γ-herpesviruses, such as KSHV (Kaposi's sarcoma-associated herpesvirus), EBV (Epstein-Barr virus) or γ -herpesvirus 68 (γ -HV68), the NF- κ B pathway was reported to be involved in cell growth as well as in the establishment of viral latency [46-50]. Hence, it is tempting to speculate that the selective modulation of the NF-kB activation via heteromerization with US28 might be important in viral reactivation or latency. Indeed, heteromerization of US28 and UL78, which is co-expressed during the early phases of HCMV infection, could provide a way to downregulate the NF-kB activation of US28. In addition, during late stages of HCMV infection, the constitutive activity of US28 could be modulated via heteromerization with UL33. After the initial re-programming of newly infected cells, the downregulation of US28 via heteromerization with the orphan receptors UL33 and UL78 might thus represent a mechanism to negatively modulate the NF- κ B pathway.

Recent reports suggest that US28 might play a causative role in multiple diseases associated with HCMV infection. Human cytomegalovirus has been shown to be present in various malignancies, including breast cancer [51], colon cancer [52] or malignant glioblastoma [53]. Indeed, US28 seems to be involved in tumorigenesis via the up-regulation of COX-2 [54, 55]. Moreover, Bongers and co-workers showed that the receptor also promotes intestinal neoplasia and cancer in transgenic mice [56]. In addition, US28 might

also represent the causative link between HCMV infection and the accelerated progression of vascular diseases, since it is able to mediate the migration of vascular smooth muscle cells [57].

Hence, the finding that US28 is engaged in hetero-/oligomeric complexes provides important information for future drug design. Furthermore, the manipulation of the NF- κ B pathway via heteromerization with the other HCMV encoded receptors might contribute to understanding the mechanisms underlying viral latency and reactivation *in vivo*.

5. Acknowledgements

This work was supported by grants from the Austrian Science Fund (P18723 to MW), the Jubiläumsfonds of the Austrian National Bank (to MW), the Lanyar Stiftung Graz, Austria (to MW and PT), the 'Molecular Medicine Ph.D. program' of the Medical University of Graz, Austria (to PT), grants from the Fondation Recherche Médicale ("Equipe FRM"), the Association pour la Recherche sur le Cancer (ARC, n° 5051), the Institut National de la Santé et de la Recherche Médicale (INSERM) and the Centre National de la Recherche Scientifique (CNRS). PT and KT were supported by research fellowships of the Medical University of Graz (Austria) and the Université Paris Descartes (France), respectively.

6. References

- [1] Humar A and Michaels M. American Society of Transplantation recommendations for screening, monitoring and reporting of infectious complications in immunosuppression trials in recipients of organ transplantation. Am J Transplant 2006;6(2):262-74.
- [2] Gandhi MK and Khanna R. Human cytomegalovirus: clinical aspects, immune regulation, and emerging treatments. Lancet Infect Dis 2004;4(12):725-38.
- [3] Ross SA and Boppana SB. Congenital cytomegalovirus infection: outcome and diagnosis. Semin Pediatr Infect Dis 2005;16(1):44-9.
- [4] Griffiths PD. CMV as a cofactor enhancing progression of AIDS. J Clin Virol 2006;35(4):489-92.
- [5] Rosenkilde MM, Waldhoer M, Luttichau HR and Schwartz TW. Virally encoded 7TM receptors. Oncogene 2001;20(13):1582-93.
- [6] Rosenkilde MM, Smit MJ and Waldhoer M. Structure, function and physiological consequences of virally encoded chemokine seven transmembrane receptors. Br J Pharmacol 2008;153 Suppl 1:S154-66.
- [7] Fraile-Ramos A, Pelchen-Matthews A, Kledal TN, Browne H, Schwartz TW and Marsh M. Localization of HCMV UL33 and US27 in endocytic compartments and viral membranes. Traffic 2002;3(3):218-32.
- [8] Cepeda V, Esteban M and Fraile-Ramos A. Human cytomegalovirus final envelopment on membranes containing both trans-Golgi network and endosomal markers. Cell Microbiol 12(3):386-404.
- [9] Casarosa P, Waldhoer M, LiWang PJ, Vischer HF, Kledal T, Timmerman H, et al. CC and CX3C chemokines differentially interact with the N terminus of the human cytomegalovirus-encoded US28 receptor. J Biol Chem 2005;280(5):3275-85.

- [10] Pleskoff O, Treboute C and Alizon M. The cytomegalovirus-encoded chemokine receptor US28 can enhance cell-cell fusion mediated by different viral proteins. J Virol 1998;72(8):6389-97.
- [11] Boomker JM, The TH, de Leij LF and Harmsen MC. The human cytomegalovirusencoded receptor US28 increases the activity of the major immediate-early promoter/enhancer. Virus Res 2006;118(1-2):196-200.
- [12] Casarosa P, Bakker RA, Verzijl D, Navis M, Timmerman H, Leurs R, et al.
 Constitutive signaling of the human cytomegalovirus-encoded chemokine receptor
 US28. J Biol Chem 2001;276(2):1133-7.
- [13] Waldhoer M, Kledal TN, Farrell H and Schwartz TW. Murine cytomegalovirus
 (CMV) M33 and human CMV US28 receptors exhibit similar constitutive signaling activities. J Virol 2002;76(16):8161-8.
- [14] McLean KA, Holst PJ, Martini L, Schwartz TW and Rosenkilde MM. Similar activation of signal transduction pathways by the herpesvirus-encoded chemokine receptors US28 and ORF74. Virology 2004;325(2):241-51.
- [15] Moepps B, Tulone C, Kern C, Minisini R, Michels G, Vatter P, et al. Constitutive serum response factor activation by the viral chemokine receptor homologue pUS28 is differentially regulated by Galpha(q/11) and Galpha(16). Cell Signal 2008;20(8):1528-37.
- [16] Fraile-Ramos A, Kledal TN, Pelchen-Matthews A, Bowers K, Schwartz TW and Marsh M. The human cytomegalovirus US28 protein is located in endocytic vesicles and undergoes constitutive endocytosis and recycling. Mol Biol Cell 2001;12(6):1737-49.
- [17] Tschische P, Moser E, Thompson D, Vischer HF, Parzmair GP, Pommer V, et al. The G-protein coupled receptor associated sorting protein GASP-1 regulates the signalling and trafficking of the viral chemokine receptor US28. Traffic 11(5):660-74.

- [18] Casarosa P, Gruijthuijsen YK, Michel D, Beisser PS, Holl J, Fitzsimons CP, et al. Constitutive signaling of the human cytomegalovirus-encoded receptor UL33 differs from that of its rat cytomegalovirus homolog R33 by promiscuous activation of G proteins of the Gq, Gi, and Gs classes. J Biol Chem 2003;278(50):50010-23.
- [19] Margulies BJ and Gibson W. The chemokine receptor homologue encoded by US27 of human cytomegalovirus is heavily glycosylated and is present in infected human foreskin fibroblasts and enveloped virus particles. Virus Res 2007;123(1):57-71.
- [20] O'Connor CM and Shenk T. Human Cytomegalovirus pUS27 G Protein-coupled Receptor Homologue Is Required for Efficient Spread by the Extracellular Route but Not for Direct Cell-to-Cell Spread. J Virol.
- [21] Vischer HF, Leurs R and Smit MJ. HCMV-encoded G-protein-coupled receptors as constitutively active modulators of cellular signaling networks. Trends Pharmacol Sci 2006;27(1):56-63.
- [22] Michel D, Milotic I, Wagner M, Vaida B, Holl J, Ansorge R, et al. The human cytomegalovirus UL78 gene is highly conserved among clinical isolates, but is dispensable for replication in fibroblasts and a renal artery organ-culture system. J Gen Virol 2005;86(Pt 2):297-306.
- [23] Levoye A, Dam J, Ayoub MA, Guillaume JL, Couturier C, Delagrange P, et al. The orphan GPR50 receptor specifically inhibits MT1 melatonin receptor function through heterodimerization. Embo J 2006;25(13):3012-23.
- [24] Terrillon S, Durroux T, Mouillac B, Breit A, Ayoub MA, Taulan M, et al. Oxytocin and vasopressin V1a and V2 receptors form constitutive homo- and heterodimers during biosynthesis. Mol Endocrinol 2003;17(4):677-91.
- [25] Martini L, Waldhoer M, Pusch M, Kharazia V, Fong J, Lee JH, et al. Ligand-induced down-regulation of the cannabinoid 1 receptor is mediated by the G-protein-coupled receptor-associated sorting protein GASP1. Faseb J 2007;21(3):802-11.

- [26] Fraile-Ramos A, Kohout TA, Waldhoer M and Marsh M. Endocytosis of the viral chemokine receptor US28 does not require beta-arrestins but is dependent on the clathrin-mediated pathway. Traffic 2003;4(4):243-53.
- [27] Neote K, DiGregorio D, Mak JY, Horuk R and Schall TJ. Molecular cloning, functional expression, and signaling characteristics of a C-C chemokine receptor. Cell 1993;72(3):415-25.
- [28] Ferre S, Baler R, Bouvier M, Caron MG, Devi LA, Durroux T, et al. Building a new conceptual framework for receptor heteromers. Nat Chem Biol 2009;5(3):131-4.
- [29] Milligan G. G protein-coupled receptor dimerization: function and ligand pharmacology. Mol Pharmacol 2004;66(1):1-7.
- [30] Levoye A, Dam J, Ayoub MA, Guillaume JL and Jockers R. Do orphan G-proteincoupled receptors have ligand-independent functions? New insights from receptor heterodimers. EMBO Rep 2006;7(11):1094-8.
- [31] Kledal TN, Rosenkilde MM and Schwartz TW. Selective recognition of the membrane-bound CX3C chemokine, fractalkine, by the human cytomegalovirusencoded broad-spectrum receptor US28. FEBS Lett 1998;441(2):209-14.
- [32] Kuhn DE, Beall CJ and Kolattukudy PE. The cytomegalovirus US28 protein binds multiple CC chemokines with high affinity. Biochem Biophys Res Commun 1995;211(1):325-30.
- [33] Waldhoer M, Casarosa P, Rosenkilde MM, Smit MJ, Leurs R, Whistler JL, et al. The carboxyl terminus of human cytomegalovirus-encoded 7 transmembrane receptor US28 camouflages agonism by mediating constitutive endocytosis. J Biol Chem 2003;278(21):19473-82.
- [34] Prinster SC, Hague C and Hall RA. Heterodimerization of g protein-coupled receptors: specificity and functional significance. Pharmacol Rev 2005;57(3):289-98.

- [35] Guo W, Shi L, Filizola M, Weinstein H and Javitch JA. Crosstalk in G proteincoupled receptors: changes at the transmembrane homodimer interface determine activation. Proc Natl Acad Sci U S A 2005;102(48):17495-500.
- [36] Sohy D, Yano H, de Nadai P, Urizar E, Guillabert A, Javitch JA, et al. Heterooligomerization of CCR2, CCR5, and CXCR4 and the protean effects of "selective" antagonists. J Biol Chem 2009;284(45):31270-9.
- [37] Contento RL, Molon B, Boularan C, Pozzan T, Manes S, Marullo S, et al. CXCR4 CCR5: a couple modulating T cell functions. Proc Natl Acad Sci U S A
 2008;105(29):10101-6.
- [38] Vila-Coro AJ, Mellado M, Martin de Ana A, Lucas P, del Real G, Martinez AC, et al. HIV-1 infection through the CCR5 receptor is blocked by receptor dimerization. Proc Natl Acad Sci U S A 2000;97(7):3388-93.
- [39] Ashall L, Horton CA, Nelson DE, Paszek P, Harper CV, Sillitoe K, et al. Pulsatile stimulation determines timing and specificity of NF-kappaB-dependent transcription. Science 2009;324(5924):242-6.
- [40] DeMeritt IB, Milford LE and Yurochko AD. Activation of the NF-kappaB pathway in human cytomegalovirus-infected cells is necessary for efficient transactivation of the major immediate-early promoter. J Virol 2004;78(9):4498-507.
- [41] Caposio P, Musso T, Luganini A, Inoue H, Gariglio M, Landolfo S, et al. Targeting the NF-kappaB pathway through pharmacological inhibition of IKK2 prevents human cytomegalovirus replication and virus-induced inflammatory response in infected endothelial cells. Antiviral Res 2007;73(3):175-84.
- [42] Benedict CA, Angulo A, Patterson G, Ha S, Huang H, Messerle M, et al. Neutrality of the canonical NF-kappaB-dependent pathway for human and murine cytomegalovirus transcription and replication in vitro. J Virol 2004;78(2):741-50.

- [43] Gustems M, Borst E, Benedict CA, Perez C, Messerle M, Ghazal P, et al. Regulation of the transcription and replication cycle of human cytomegalovirus is insensitive to genetic elimination of the cognate NF-kappaB binding sites in the enhancer. J Virol 2006;80(19):9899-904.
- [44] Evers DL, Wang X and Huang ES. Cellular stress and signal transduction responses to human cytomegalovirus infection. Microbes Infect 2004;6(12):1084-93.
- [45] Zhu H, Cong JP, Mamtora G, Gingeras T and Shenk T. Cellular gene expression altered by human cytomegalovirus: global monitoring with oligonucleotide arrays.
 Proc Natl Acad Sci U S A 1998;95(24):14470-5.
- [46] Chaudhary PM, Jasmin A, Eby MT and Hood L. Modulation of the NF-kappa B pathway by virally encoded death effector domains-containing proteins. Oncogene 1999;18(42):5738-46.
- [47] Matta H and Chaudhary PM. Activation of alternative NF-kappa B pathway by human herpes virus 8-encoded Fas-associated death domain-like IL-1 beta-converting enzyme inhibitory protein (vFLIP). Proc Natl Acad Sci U S A 2004;101(25):9399-404.
- [48] Mitchell T and Sugden B. Stimulation of NF-kappa B-mediated transcription by mutant derivatives of the latent membrane protein of Epstein-Barr virus. J Virol 1995;69(5):2968-76.
- [49] Lei X, Bai Z, Ye F, Xie J, Kim CG, Huang Y, et al. Regulation of NF-kappaB inhibitor IkappaBalpha and viral replication by a KSHV microRNA. Nat Cell Biol 12(2):193-9.
- [50] Krug LT, Moser JM, Dickerson SM and Speck SH. Inhibition of NF-kappaB activation in vivo impairs establishment of gammaherpesvirus latency. PLoS Pathog 2007;3(1):e11.
- [51] Soderberg-Naucler C. HCMV microinfections in inflammatory diseases and cancer. J Clin Virol 2008;41(3):218-23.

- [52] Harkins L, Volk AL, Samanta M, Mikolaenko I, Britt WJ, Bland KI, et al. Specific localisation of human cytomegalovirus nucleic acids and proteins in human colorectal cancer. Lancet 2002;360(9345):1557-63.
- [53] Sabatier J, Uro-Coste E, Pommepuy I, Labrousse F, Allart S, Tremoulet M, et al. Detection of human cytomegalovirus genome and gene products in central nervous system tumours. Br J Cancer 2005;92(4):747-50.
- [54] Maussang D, Langemeijer E, Fitzsimons CP, Stigter-van Walsum M, Dijkman R, Borg MK, et al. The human cytomegalovirus-encoded chemokine receptor US28 promotes angiogenesis and tumor formation via cyclooxygenase-2. Cancer Res 2009;69(7):2861-9.
- [55] Maussang D, Verzijl D, van Walsum M, Leurs R, Holl J, Pleskoff O, et al. Human cytomegalovirus-encoded chemokine receptor US28 promotes tumorigenesis. Proc Natl Acad Sci U S A 2006;103(35):13068-73.
- [56] Bongers G, Maussang D, Muniz LR, Noriega VM, Fraile-Ramos A, Barker N, et al. The cytomegalovirus-encoded chemokine receptor US28 promotes intestinal neoplasia in transgenic mice. J Clin Invest 120(11):3969-78.
- [57] Streblow DN, Vomaske J, Smith P, Melnychuk R, Hall L, Pancheva D, et al. Human cytomegalovirus chemokine receptor US28-induced smooth muscle cell migration is mediated by focal adhesion kinase and Src. J Biol Chem 2003;278(50):50456-65.

Figure captions

Figure 1. US28 co-localizes with UL33, UL78 and US27. HEK293 cells transiently transfected with Flag-US28 and either HA-UL33 (A), HA-UL78 (B) or HA-US27 (C) were 'fed' antibody to the extracellular Flag-tag and HA-tag for 60 minutes. The cells were then fixed and stained with the secondary antibodies under permeabilizing conditions. Finally, cells were mounted with vectashield (+DAPI, blue) and analyzed by fluorescence microscopy. Merge: yellow. Scale bars = $10 \mu m$.

Figure 2 A. Detection of US28/US28 homomers using co-immunoprecipitation. HEK293T cells were transfected with HA-US28 in the presence or absence of Flag-US28-YFP. Lysates were immunoblotted with monoclonal rabbit anti-HA (bottom blot) and polyclonal rabbit anti-Flag antibodies (middle blot). Lysates were immunoprecipitated with monoclonal rat anti-HA, resolved by SDS-PAGE and immunoblotted with polyclonal rabbit anti-Flag antibodies (top blot). B. Detection of US28/UL33 heteromers using co-immunoprecipitation. HEK293T cells were transfected with HA-UL33-Rluc in the presence or absence of Flag-US28-YFP. The expression of HA-UL33-Rluc was monitored by measuring Renilla luciferase activity following coelenterazine H addition (bottom graph). Lysates were immunoblotted with monoclonal mouse anti-GFP antibodies (middle blot). Lysates were immunoprecipitated with monoclonal rat anti-HA, resolved by SDS-PAGE and immunoblotted with monoclonal mouse anti-GFP antibodies (top blot). C. Detection of US28/UL78 heteromers using coimmunoprecipitation. HEK293T cells were transfected with HA-UL78-Rluc in the presence or absence of Flag-US28-YFP. The expression of HA-UL78-Rluc was monitored by measuring Renilla luciferase activity following coelenterazine H addition (bottom graph). Lysates were immunoblotted with polyclonal rabbit anti-Flag antibodies (middle blot).

Lysates were immunoprecipitated with monoclonal rat anti-HA, resolved by SDS–PAGE and immunoblotted with polyclonal rabbit anti-Flag antibodies (top blot). All immunoblots shown are representative of three independent experiments. **D. Detection of US28/US27 heteromers using co-immunoprecipitation.** HEK293T cells were transfected with HA-US27 in the presence or absence of Flag-US28-YFP to assess US27/US28 heteromer formation. Lysates were immunoblotted with monoclonal rat anti-HA (bottom blot) and polyclonal rabbit anti-Flag antibodies (middle blot). Lysates were immunoprecipitated with monoclonal rat anti-HA, resolved by SDS–PAGE and immunoblotted with polyclonal rabbit anti-Flag antibodies (top blot).

Figure 3. Heteromerization of US28 with UL33 and UL78. A. Donor saturation curves were performed by co-transfecting a fixed amount of HA-UL33-Rluc in presence of increasing amounts of Flag-US28-YFP in HEK 293T cells. The saturation curves are obtained from three independent experiments. B. Donor saturation curves were performed by cotransfecting a fixed amount of HA-UL78-Rluc in presence of increasing amounts of Flag-US28-YFP (●) in HEK 293T cells. V2-YFP (■) was used as a negative control. The saturation curves are obtained from three independent experiments. C. Bret competition assays were performed by co-transfecting a fixed amount of HA-UL33-Rluc and Flag-US28-YFP corresponding to a YFP/Rluc ratio in the ascending portion of the saturation curve before the plateau in presence of saturating amounts of the competitor receptor (Flag-UL78). Lysates were immunoblotted with polyclonal rabbit anti-Flag antibody (bottom blot) to assess the expression levels of the competitor receptor. **D.** Bret competition assays were performed by co-transfecting a fixed amount of HA-UL78-Rluc and Flag-US28-YFP corresponding to a YFP/Rluc ratio in the ascending portion of the saturation curve before the plateau in presence of saturating amounts of the competitor receptor (HA-UL33). Lysates were immunoblotted with monoclonal rabbit anti-HA antibody (bottom blot) to assess the expression levels of the

competitor receptor. Statistical differences were assessed using the student t-test (***p <0.001).

Figure 4. US28-mediated inositol phosphate (IP) accumulation is not altered by heteromerization with UL33, UL78 or US27. A, D & G. IP accumulation in HEK293 cells co-expressing receptor heteromers. HEK293 cells were transiently transfected with Flag-US28 (75 ng/well) and either pcDNA3.1 (black bars, 75 ng/well) or 75ng/well of HA-UL33 (A), HA-UL78 (D) or HA-US27 (G), respectively (grey bars). Controls were transfected with either pcDNA3.1 alone (150 ng/well, white bars) or pcDNA3.1 (75 ng/well) in combination with 75 ng/well of HA-UL33 (A), HA-UL78 (D) or HA-US27 (G) (striped bars) in the absence of Flag-US28. IP production mediated by US28 was allowed to accumulate for 45 minutes. Values were normalized to the cell number, and 100% corresponds to the basal activity of US28 (black bars). Data are means of four experiments \pm SEM carried out in quadruplicates. B, C, E, F, H & I. Receptor expression levels evaluated by ELISA. HEK293 cells were transfected as described for the respective IP accumulation assay. The expression levels of US28 were assayed by ELISA against the Flag epitope tag of US28 (B, E & H). The levels of UL33 (C), UL78 (F) and US27 (I) were assessed by ELISA against the HA epitope tag of the respective receptors. Values were normalized to the cell number. Data are means ± SEM of four independent experiments carried out in quadruplicates.

Figure 5. UL33 and UL78 alter US28 mediated NF-κB activation. A, D & G. NF-κB activation in HEK293 cells co-expressing receptor heteromers. HEK293 cells were transiently transfected with Flag-US28 (75 ng/well), the cis-reporter luciferase plasmid for NF-κB (50 ng/well) and either pcDNA3.1 (black bars, 75 ng/well) or 75ng/well of HA-UL33 (A), HA-UL78 (D) or HA-US27 (G), respectively (grey bars). Controls were transfected with

either pcDNA3.1 alone (150 ng/well, white bars) or pcDNA3.1 (75 ng/well) in combination with 75 ng/well of HA-UL33 (**A**), HA-UL78 (**D**) or HA-US27 (**G**) (striped bars) in the absence of Flag-US28. A luciferase reporter assay was conducted 24 hours post-transfection. Values were normalized to the cell number, whereby 100% corresponds to the basal activity of US28 (black bars). Data are means of three independent experiments \pm SEM carried out in quadruplicates. *** P<0.001. **B**, **C**, **E**, **F**, **H** & **I**. **Receptor expression levels evaluated by ELISA**. HEK293 cells were transfected as described for the respective NF-κB assay. The expression levels of US28 were assayed by ELISA against the Flag epitope tag of US28 (**B**, **E** & **H**). The levels of UL33 (**C**), UL78 (**F**) and US27 (**I**) were assessed by ELISA against the HA epitope tag of the respective receptors. Values were normalized to the cell number. Data are means \pm SEM of four independent experiments carried out in quadruplicates.

Figure 1





Figure Click here to download high resolution image















