



HAL
open science

Cellular accumulation of fluoroquinolones is not predictive of their intracellular activity: studies with gemifloxacin, moxifloxacin and ciprofloxacin in a pharmacokinetic/pharmacodynamic model of uninfected and infected macrophages

Coralie M. Vallet, Béatrice Marquez, Eva Ngabirano, Sandrine Lemaire, Marie-Paule Mingeot-Leclercq, Paul M. Tulkens, Françoise van Bambeke

► To cite this version:

Coralie M. Vallet, Béatrice Marquez, Eva Ngabirano, Sandrine Lemaire, Marie-Paule Mingeot-Leclercq, et al.. Cellular accumulation of fluoroquinolones is not predictive of their intracellular activity: studies with gemifloxacin, moxifloxacin and ciprofloxacin in a pharmacokinetic/pharmacodynamic model of uninfected and infected macrophages. *International Journal of Antimicrobial Agents*, 2011, <10.1016/j.ijantimicag.2011.05.011>. <hal-00722865>

HAL Id: hal-00722865

<https://hal.science/hal-00722865v1>

Submitted on 6 Aug 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Accepted Manuscript

Title: Cellular accumulation of fluoroquinolones is not predictive of their intracellular activity: studies with gemifloxacin, moxifloxacin and ciprofloxacin in a pharmacokinetic/pharmacodynamic model of uninfected and infected macrophages



Authors: Coralie M. Vallet, Béatrice Marquez, Eva Ngabirano, Sandrine Lemaire, Marie-Paule Mingeot-Leclercq, Paul M. Tulkens, Françoise Van Bambeke

PII: S0924-8579(11)00252-4
DOI: doi:10.1016/j.ijantimicag.2011.05.011
Reference: ANTAGE 3636

To appear in: *International Journal of Antimicrobial Agents*

Received date: 22-3-2011
Revised date: 14-5-2011
Accepted date: 17-5-2011

Please cite this article as: Vallet CM, Marquez B, Ngabirano E, Lemaire S, Mingeot-Leclercq M-P, Tulkens PM, Van Bambeke F, Cellular accumulation of fluoroquinolones is not predictive of their intracellular activity: studies with gemifloxacin, moxifloxacin and ciprofloxacin in a pharmacokinetic/pharmacodynamic model of uninfected and infected macrophages, *International Journal of Antimicrobial Agents* (2010), doi:10.1016/j.ijantimicag.2011.05.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Cellular accumulation of fluoroquinolones is not predictive of their**
2 **intracellular activity: studies with gemifloxacin, moxifloxacin, and**
3 **ciprofloxacin in a pharmacokinetic/pharmacodynamic model of**
4 **uninfected and infected macrophages.**

5
6 Coralie M. Vallet, Béatrice Marquez,[§] Eva Ngabirano, Sandrine Lemaire, Marie-Paule
7 Mingeot-Leclercq, Paul M. Tulkens*, and Françoise Van Bambeke

8
9 Pharmacologie cellulaire et moléculaire, Louvain Drug Research Institute, Université
10 catholique de Louvain, Brussels, Belgium

11
12 **Running title:** Cellular pharmacokinetics and activity of fluoroquinolones

13
14
15 [§] Present address: Grupo Tumorigenesis endocrina y regulación hormonal del cáncer,
16 Instituto de Biomedicina de Sevilla - Hospitales Universitarios Virgen del Rocío, Avenida
17 Manuel Siurot s/n, 41013 Sevilla, Spain.

18
19
20
21 *** Corresponding author**

22 Pharmacologie cellulaire et moléculaire,
23 Université catholique de Louvain,
24 Avenue E. Mounier 73 bte B1.73.05,
25 B-1200 Bruxelles, Belgium.

26 Phone: +32-2-764-73-71.

27 Fax: +32-2-764-73-73.

28 E-mail: tulkens@facm.ucl.ac.be

29

30 **ABSTRACT**

31 Fluoroquinolones enter eukaryotic cells but the correlation between cellular accumulation
32 and activity remains poorly established. Gemifloxacin is known to accumulate to a larger
33 extent than most other fluoroquinolones in tissues. Using murine J774 macrophages and
34 human THP-1 monocytes, we show that gemifloxacin accumulates more than ciprofloxacin
35 and even moxifloxacin. While showing indistinguishable kinetics of accumulation in J774
36 macrophages, gemifloxacin was released at an approximately two-fold slower rate than
37 ciprofloxacin and its release was only partial. Gemifloxacin was also a weaker substrate
38 than ciprofloxacin for the efflux transporter Mrp4 active in J774 macrophages. In cells
39 infected with *L. monocytogenes* or *S. aureus* (typical cytoplasmic and phagolysosomal
40 organisms, respectively), gemifloxacin was equipotent to moxifloxacin and ciprofloxacin in
41 concentration-dependent experiments if data are normalized based on MIC in broth. Thus,
42 larger cellular concentrations of gemifloxacin than of moxifloxacin or ciprofloxacin were
43 needed to obtain a similar target effect. Fractionation studies showed a similar subcellular
44 distribution for all 3 fluoroquinolones, with about 2/3 of the cell-associated drug recovered in
45 the soluble fraction (cytosol). The data suggest that the cellular accumulation of
46 fluoroquinolones is largely a self-defeating process as far as activity is concerned, with the
47 intracellular drug made inactive in proportion to its accumulation level. While these
48 observations do not decrease the intrinsic value of fluoroquinolones for treatment of
49 intracellular infections, they indicate that ranking fluoronolones based on cell accumulation
50 data without measuring the corresponding intracellular activity may lead to incorrect
51 conclusions concerning their real potential.

52

53 **Keywords:** ciprofloxacin, moxifloxacin, gemifloxacin, Mrp4, Staphylococcus aureus, Listeria
54 monocytogenes, macrophages

55

56

57 **1. Introduction**

58 Fluoroquinolone antibiotics are important in our current therapeutic arsenal, because of their
59 broad spectrum, highly bactericidal activity, and favourable pharmacokinetic properties [1].
60 Their wide tissue distribution allows them reaching therapeutic concentrations in deep body
61 compartments as well as in the intracellular milieu, which may be an advantage in the
62 treatment of intracellular infections. Accumulation and activity in cells are usually linked
63 when considering a given fluoroquinolone in a specific cell type, as demonstrated for
64 ciprofloxacin *vis-à-vis* the intracellular forms of *L. monocytogenes* in J774 macrophages in
65 experiments where its cellular concentration was modulated by inhibition or overexpression
66 of the constitutive ciprofloxacin efflux transporter Mrp4 [2;3]. There is, however, a lack of
67 quantitative data comparing distinct fluoroquinolones in this context.

68
69 Gemifloxacin [4] accumulates to high levels in human PMN and is active against intracellular
70 bacteria [5;6]. This prompted us to compare it to other fluoroquinolones for cellular
71 pharmacokinetics and activity in an established model of murine J774 macrophages [7].
72 Ciprofloxacin and moxifloxacin, **when needed**, were used comparators, as these show a low
73 and large accumulation, respectively, in relation to differential susceptibility to efflux [8-11].
74 We also examined THP-1 cells, where no active fluoroquinolone efflux has been evidenced
75 so far. We found that gemifloxacin accumulates to higher levels than ciprofloxacin and
76 moxifloxacin in both cell types and that all three drugs have a similar subcellular distribution.
77 Yet, gemifloxacin showed no improved activity against two types of intracellular bacteria,
78 *L. monocytogenes* and *S. aureus*, localized in the cytosol and in phagolysosomes,
79 respectively.

80

81 2. Materials and methods

82 2.1. Antibiotics and main reagents

83 Gemifloxacin mesylate (LG Life Sciences, Seoul, Korea), and ciprofloxacin-HCl and
84 moxifloxacin-HCl (Bayer HealthCare AG, Leverkusen, Germany) were obtained as
85 microbiological standards (potencies: 79 %, 85 %, and 91 %). Gemfibrozil was from Sigma-
86 Aldrich (St-Louis, MO, USA), human serum from Lonza Ltd (Basel, Switzerland), and cell
87 culture media and sera from Invitrogen Corp. (Carlsbad, CA).

88

89 2.2. Cell lines

90 Murine J774 macrophages (wild-type cells [9]) and their ciprofloxacin-resistant derivatives
91 overexpressing Mrp4 efflux transporter [8;11] were used for most experiments. Human THP-
92 1 cells (ATCC TIB-202, American Tissue Culture Collection, Manassas, VA) [12;13] were
93 used for comparison purposes. ATP depletion was obtained as previously described [9].

94

95 2.3. Determination of the cellular accumulation of fluoroquinolones.

96 We used a previously described protocol [9;14]. Cell -associated fluoroquinolones were
97 assayed by fluorimetry (see [10] for ciprofloxacin and moxifloxacin; for gemifloxacin, the
98 conditions were: $\lambda_{\text{ex.}} = 270\text{nm}$; $\lambda_{\text{em.}} = 402\text{ nm}$ [lowest limit of detection: 50 $\mu\text{g/L}$; linearity: 0-
99 1.5 mg/L]). The cell drug content was expressed by reference to the total cell protein content
100 [15]. The apparent total cellular concentration was then calculated using a conversion factor
101 of 3.08 μL of cell volume per mg of cell protein [9].

102

103 2.4. Cell fractionation studies in J774 cells

104 The main subcellular organelles were separated by differential centrifugation as previously
105 described [2]. The protein and antibiotic content of each fraction was determined in parallel
106 with the activity of marker enzymes of the main organelles (cytochrome c-oxidase for
107 mitochondria; N-acetyl- β -hexosaminidase for lysosomes, and lactate dehydrogenase for
108 cytosol [7]).

109 *2.5. Bacterial strains and susceptibility testings.*

110 We used *L. monocytogenes* strain EGD and *S. aureus* strain ATCC 25923. MIC
111 determinations were made according to CLSI guidelines [16], using Tryptic Soy broth for *L.*
112 *monocytogenes* [13] and Mueller Hinton broth for *S. aureus* [14].

113

114 *2.6. Cell infection and assessment of antibiotic intracellular activities.*

115 Cell infection was performed as described previously [2], with pharmacological comparison
116 between drugs and bacteria based on concentration-dependent effects analyses [14], to
117 determine (i) the relative minimal and maximal efficacies (E_{\min} / E_{\max} , in \log_{10} units), and (ii)
118 the relative potencies (EC_{50}) and static concentrations). This type of analysis and its
119 usefulness for comparing antibiotics and the response of different bacteria has been
120 described in details in previous publications [14;17-19]). As discussed previously [20], the
121 large dilution of samples before spreading on agar plates for cfu counting ensures an
122 absence of carry-over effect.

123

124 *2.7. Curve fitting and statistical analyses*

125 Curve-fitting analyses were made using GraphPad Prism® version 4.03, GraphPad
126 Software, San Diego, CA, USA. Statistical analyses were made with the same software for
127 comparing concentration-response functions and with GraphPad InStat® version 3.06
128 (GraphPad Software) for other studies.

129

130 3. Results

131 3.1. Cellular pharmacokinetics

132 We first compared the cellular accumulation of gemifloxacin with that of ciprofloxacin and
133 moxifloxacin, and examined the influence of gemfibrozil, a broad spectrum inhibitor of anion
134 transporters including the Mrp transporters, on this accumulation. Figure 1A (upper panels)
135 shows that (i) gemifloxacin accumulated to a larger extent than the other two
136 fluoroquinolones in both J774 and THP-1 cells; (ii) the accumulation of gemifloxacin and
137 moxifloxacin was not influenced by gemfibrozil; (iii) in contrast, ciprofloxacin, which
138 accumulated to the lowest extent in J774 macrophages, reached a cellular concentration
139 similar to that of moxifloxacin in these cells in the presence of gemfibrozil, as already
140 observed in the same model [10]; (iv) the level of accumulation of ciprofloxacin was similar to
141 that of moxifloxacin in THP-1 cells and not influenced by the addition of gemfibrozil.

142
143 We then compared the kinetics of accumulation and efflux of gemifloxacin with that of
144 ciprofloxacin using J774 macrophages only as this is where the largest difference of
145 accumulation was observed. Figure 1B shows that the two fluoroquinolones could not be
146 distinguished with respect to accumulation kinetics but displayed marked differences for
147 efflux. Thus, gemifloxacin release (i) occurred at the same rate as its uptake (compare k_{in}
148 and k_{out} parameters); (ii) was about 2-fold slower than that of ciprofloxacin, including at the
149 very initial period phase (see inset); (iii) was only partial, with about 25 % of the accumulated
150 drug remaining cell-associated in apparent stable fashion after 30 min incubation in drug-free
151 medium vs. negligible amounts for ciprofloxacin.

152
153 We next measured the level of accumulation of gemifloxacin compared to that of
154 ciprofloxacin in J774 macrophages overexpressing the ciprofloxacin efflux transporter Mrp4
155 (ciprofloxacin-resistant cells), using normal conditions and conditions of ATP depletion
156 (which inhibits all ATP-dependent active transporters including Mrp4). Figure 2A (upper

157 panels) shows that (i) gemifloxacin accumulation was reduced (but in a non-statistically
158 significant manner) in ciprofloxacin-resistant cells compared to wild-type cells; (ii) ATP
159 depletion increased its accumulation in both wild-type and ciprofloxacin-resistant cells, but
160 with a significant difference in the latter cells only; (iii) ciprofloxacin accumulation was
161 significantly reduced in ciprofloxacin-resistant cells, but was markedly increased by ATP
162 depletion, reaching a value similar to that observed in wild-type cells after ATP depletion; (iv)
163 in line with our previous observations [11], ATP depletion markedly increased the
164 accumulation of ciprofloxacin in wild-type cells.

165
166 Because the ciprofloxacin efflux transporter is saturable in a 10-200 mg/L range [9], we
167 measured the accumulation of gemifloxacin in both wild-type J774 macrophages and
168 ciprofloxacin-resistant cells over increasing concentrations of gemifloxacin in that range.
169 Figure 2 (lower panel) shows that while gemifloxacin accumulation was not significantly
170 influenced by its extracellular concentration in wild-type cells, there was a significant increase
171 over the range of concentrations investigated for ciprofloxacin-resistant cells. In contrast,
172 and as described earlier [9], ciprofloxacin showed a marked increase in its accumulation over
173 the same concentration range in wild-type cells. For ciprofloxacin-resistant cells, the
174 increase in cell accumulation of ciprofloxacin was much less marked in the range of drug
175 concentrations investigated due to overexpression of the Mrp4 transporter (see [11]).

176
177 These results suggest that gemifloxacin could be a poor, albeit still recognized substrate for
178 efflux transport in J774 macrophages if Mrp4 is overexpressed. We, therefore, compared the
179 kinetics of gemifloxacin efflux in ciprofloxacin-resistant vs. wild-type cells. While the *plateau*
180 values observed at 30 min remained close from each other, denoting an incomplete release
181 of gemifloxacin in both cases, its rate of efflux was significantly accelerated in ciprofloxacin-
182 resistant cells compared to wild-type cells ($k_{out} = 2.393 \pm 0.907$ vs. 0.403 ± 0.122 min⁻¹;
183 $p < 0.001$; see graphical representation in Figure SP1 in the Supplementary Material).

184 3.2. Intracellular activity

185 To examine the correlation between cellular accumulation and intracellular activity, we
186 compared all 3 fluoroquinolones in our pharmacological model of intracellular infection
187 [14;17], using J774 macrophages since this is where the largest differences in accumulation
188 levels had been observed. *L. monocytogenes* and *S. aureus* were selected as bacterial
189 targets as they represent a typical cytoplasmic and phagolysosomal organism, respectively.
190 Data presented in Figure 3A (with analysis of the key pharmacological descriptors in Table 1)
191 show that all 3 antibiotics induced essentially a similar response when expressed as a
192 function of equipotent extracellular concentrations (multiples of MIC). Thus, in all cases, a
193 single sigmoid function could be fitted to the individual responses of each antibiotic (see
194 Figure SP2 in the Supplementary Material and the pertinent regression parameters and
195 pharmacological descriptors in Table 1). As no statistically significant difference was
196 observed between the 3 antibiotics, all data were pooled to fit a single function shown in
197 Figure 3A. Thus, for each bacterium, the relative minimal efficacies (E_{\min} [growth in the
198 absence of antibiotic]), maximal relative efficacies (E_{\max} [maximal antibiotic-related killing]),
199 relative potencies (E_{50}), and static concentrations (C_s ; [in multiples of MIC) were not
200 statistically significantly different. We then calculated for each fluoroquinolone which cellular
201 drug concentration would be needed to reach two predefined pharmacodynamic targets
202 (static effect and a 1 or 2 \log_{10} cfu decrease). The results (with the mode of calculation) are
203 presented in Figure 3B and show that the potencies of the drugs with respect to their
204 intracellular targets is in inverse proportion to their respective cellular accumulations.

205

206 3.3. Subcellular distribution

207 Lastly, we compared the subcellular distributions of ciprofloxacin, moxifloxacin, and
208 gemifloxacin. Figure 4 shows that all 3 fluoroquinolones shared essentially the same
209 distribution, with about 70 % recovered in the soluble fraction, about 10 % of ciprofloxacin
210 and gemifloxacin and 18 % of moxifloxacin in the nuclei/unbroken cells fraction, and the

211 remaining in the organelles/membranes fraction. As previously described [7], lactate
212 dehydrogenase was mostly recovered in the soluble fraction, and cytochrome oxydase and
213 N-acetyl- β -hexosaminidase, in the granules/membranes fraction, indicating that the
214 fractionation method effectively separated the corresponding subcellular entities with only a
215 very low proportion of unbroken cells left after homogenization.

216

Accepted Manuscript

217 **4. Discussion**

218 Gemifloxacin, approved for clinical use in over 27 countries [21], is characterized by very low
219 MICs against Gram-positive bacteria [22;23] related to the presence of an oximinomethyl
220 group [4] in its C7 side-chain and by a high tissular accumulation [24]. Human PK/PD
221 studies show that gemifloxacin achieves higher AUC/MIC ratios in epithelial lining fluid and
222 alveolar macrophages than other currently used fluoroquinolones, suggesting an advantage
223 in terms of availability and efficacy at the site of infection [25;26]. The present study,
224 however, shows that the higher accumulation of gemifloxacin in J774 macrophages (i) is not
225 associated with differences in influx rates compared to a fluoroquinolone with lower
226 accumulation (ciprofloxacin); (ii) does not preclude and cannot be explained by differences in
227 active efflux transport (in comparison with moxifloxacin); (iii) does not lead to a higher
228 intracellular activity. This goes against commonly accepted pharmacokinetic and
229 pharmacodynamic concepts that tend to link accumulation and lack of efflux on the one
230 hand, and accumulation and activity on the other hand. Our model may be questioned, but it
231 is important to note that it reproduces (i) with respect to pharmacokinetics, what is observed
232 in human alveolar macrophages where the concentrations of ciprofloxacin, moxifloxacin, and
233 gemifloxacin are respectively 2-5 x, 20-40 x, and 90 x higher than serum levels [26-28], and
234 (ii) with respect to intracellular activity what has been observed in human polymorphonuclear
235 leucocytes infected by *S. aureus* [5].

236
237 Mechanistically, differences in accumulation of drugs in cells and tissues usually result from
238 commensurate differences in influx or efflux rates, or from differential trapping by intracellular
239 organelles or constituents.

240
241 Considering influx first, faster drug accumulation is usually related to a higher lipophilicity
242 (which is supposed to facilitate transmembrane diffusion) or from the activity of transporters.
243 This does not seem to apply to gemifloxacin, as this fluoroquinolone (i) is not globally more
244 lipophilic than ciprofloxacin (see Table SP1 for experimental and calculated log P and log D

245 values), (ii) is probably not the substrate of a specific influx transporter when compared to
246 ciprofloxacin (same rate accumulation constants). Non-specific influx transporter(s)
247 observed in PMN and human monocytes [29-31] can probably be dismissed here as these
248 belong to the Solute Carrier Organic Anion (SLCO) family [32] that is inhibited by gemfibrozil,
249 which was not the case here.

250
251 Considering efflux, Mrp4 has been proposed as the main transporter responsible for the
252 lower accumulation of ciprofloxacin in J774 macrophages compared to levofloxacin,
253 garenoxacin, and moxifloxacin. These fluoroquinolones, indeed, reach a similar level of
254 accumulation when Mrp4 is made inactive by ATP depletion or addition of gemfibrozil [10].
255 Moreover, ciprofloxacin accumulation is significantly increased by silencing the gene coding
256 for Mrp4 [8]. The present data show that this conclusion cannot be generalized to all
257 fluoroquinolones and all situations. Thus gemifloxacin not only accumulates more than
258 moxifloxacin in J774 macrophages under conditions of ATP depletion or in the presence of
259 gemfibrozil, but also in THP-1 macrophages in which no gemfibrozil-inhibited efflux can be
260 demonstrated. Another compelling reason to disregard efflux as being the main cause for
261 the differential accumulation of gemifloxacin vs. ciprofloxacin and moxifloxacin is that
262 gemifloxacin actually seems a weak but nevertheless effective substrate of Mrp4 in J774
263 macrophages, whereas we know that moxifloxacin is not. Thus, globally and in contrast to
264 what we proposed for moxifloxacin, the higher cellular concentration of gemifloxacin
265 compared to other fluoroquinolones must find an explanation beyond considerations of influx
266 and efflux rates only.

267
268 Considering intracellular trapping, a model has been presented [33] that relates
269 fluoroquinolone accumulation in eukaryotic cells to their trapping under a protonated form in
270 lysosomes due to the acid pH (~ 5.4) prevailing therein. This, however, is unlikely because
271 fluoroquinolones are not weak bases but zwitterionic compounds. Moreover, differences in
272 accumulation of drugs in acidic membrane-bounded compartments should result from

273 commensurate differences in the number and/or the pK_a of their basic functions (see [34]),
274 which is not the case for the 3 fluoroquinolones studied here (see individual basic pK_a values
275 in Table SP1). More factually, cell fractionation studies show a predominant association of
276 the cell-associated fluoroquinolones with the cytosol rather than with lysosomes, in line with
277 the results of previous studies with ciprofloxacin [2;35] (studies using the same technique
278 have shown that macrolides are predominantly associated with lysosomes in J774
279 macrophages [2;36;37]). Lastly, experimental studies have shown a lack of effect of
280 monensin (a H^+ ionophore that collapses the cytosolic-lysosomal ΔpH) on ciprofloxacin
281 accumulation under conditions in which it drastically reduces the accumulation of
282 azithromycin in J774 macrophages [9].

283
284 Actually, a more likely explanation for the larger cellular accumulation of gemifloxacin
285 compared to moxifloxacin and ciprofloxacin could be its tighter binding to still undefined
286 cellular constituents such as soluble proteins. This hypothesis would account for the
287 pharmacokinetic and subcellular distribution data presented here, including (i) the lower
288 efflux rate of gemifloxacin compared to ciprofloxacin (which, however, may also result from
289 the less efficient recognition of gemifloxacin by the Mrp4 efflux transporter, both mechanisms
290 being not mutually exclusive) and, (ii) its incomplete release upon transfer of the cells to
291 drug-free medium. It is also consistent with the larger serum protein binding of gemifloxacin
292 (55-73 %) compared to moxifloxacin (39-52 %) and ciprofloxacin (30 % only) [38;39].

293
294 Determining the molecular nature of the intracellular binding sites for fluoroquinolones still
295 require further investigations, but the mechanism proposed provides a rational explanation
296 for the main critical observation made here, namely that all 3 fluoroquinolones are equipotent
297 against intracellular bacteria in spite of their differences in cellular accumulation. We show,
298 indeed, that it is the MIC of each drug that drives its intracellular potency (as defined by the
299 C_s and EC_{50} pharmacological descriptors) since all 3 fluoroquinolones show superimposable
300 concentration-effects relationships once the data are normalized on basis of multiples of the

301 MIC. MICs are measured in broth where little protein binding takes place, which means that
302 their values must essentially be interpreted as corresponding to free drug levels [40].
303 Intracellularly, a static effect (C_s) for gemifloxacin was obtained for an extracellular
304 concentration corresponding to its MIC, although its intracellular concentration is much
305 larger. It is, therefore, tempting to speculate that only a fraction of the total intracellular
306 gemifloxacin is available for activity, corresponding essentially to its free form. Moxifloxacin
307 should show an intermediate behaviour with intracellular activity also driven by its MIC
308 (measured in broth), which is what we observe. Thus, the larger cellular accumulation of
309 some fluoroquinolones, taking gemifloxacin an example, would essentially be a self-defeating
310 process as far as activity is concerned (assuming all comparisons are made on basis of the
311 MIC), leading to a larger concentration of bound drugs with, however, no or little difference in
312 their free forms. This confirms and extends previous work that showed that the intracellular
313 activity of fluoroquinolones was weaker and not in proportion to what could be anticipated
314 from the level of their cellular accumulation [13;41-43].

315

316 In conclusion, the present work documents that (i) recording the cellular accumulation of
317 fluoroquinolones does not allow to predict their intracellular activity; (ii) a higher cellular
318 accumulation may depend from other parameters than influx and efflux rates and/or the
319 activity of specific transporters. This calls for both more mechanistic studies and more
320 comprehensive structure-activity analyses where these two important elements of the
321 pharmacological properties of fluoroquinolones will be examined in a systematic fashion.

322 **Acknowledgments**

323 We are grateful to M.C. Cambier, C. Misson, and M. Vergauwen for dedicated technical
324 assistance. We thank the Oscient Pharmaceuticals Corporation and Bayer HealthCare for
325 the kind gift of gemifloxacin, and of moxifloxacin and ciprofloxacin, respectively.

326

327 **Declarations**

328 Funding: C.M.V. is Boursier of the Belgian *Fonds pour la Recherche dans l'Industrie et*
329 *l'Agriculture* (F.R.I.A.), B.M. was post-doctoral fellow of the First post-doc programme of the
330 Belgian *Région wallonne*, S.L. and F.V.B. are *Chargé de Recherches* and *Maître de*
331 *Recherches* of the Belgian *Fonds de la Recherche Scientifique (F.R.S.-FNRS)*, respectively.

332 This work was supported by the Belgian *Fonds de la Recherche Scientifique Médicale*
333 (grants no. 3.4.597.06 and 3.4.583.08) and the Belgian *Fonds de la Recherche Scientifique*
334 (grant no. 1.5.195.07).

335 **Competing interests:** None

336 **Ethical approval:** Not applicable

337

338 **References**

- 339 [1] Van Bambeke F, Michot JM, Van Eldere J, Tulkens PM. Quinolones in 2005: an
340 update. *Clin Microbiol Infect* 2005; 11:256-280.
- 341 [2] Seral C, Carryn S, Tulkens PM, Van Bambeke F. Influence of P-glycoprotein and MRP
342 efflux pump inhibitors on the intracellular activity of azithromycin and ciprofloxacin in
343 macrophages infected by *Listeria monocytogenes* or *Staphylococcus aureus*. *J*
344 *Antimicrob Chemother* 2003; 51:1167-1173.
- 345 [3] Lismond A, Tulkens PM, Mingeot-Leclercq MP, Courvalin P, Van Bambeke F.
346 Cooperation between prokaryotic (Lde) and eukaryotic (MRP) efflux transporters in
347 J774 macrophages infected with *Listeria monocytogenes*: studies with ciprofloxacin and
348 moxifloxacin. *Antimicrob Agents Chemother* 2008; 52:3040-3046.
- 349 [4] Hong CY, Kim YK, Chang JH, Kim SH, Choi H, Nam DH et al. Novel fluoroquinolone
350 antibacterial agents containing oxime-substituted (aminomethyl)pyrrolidines: synthesis
351 and antibacterial activity of 7-(4-(aminomethyl)-3-(methoxyimino)pyrrolidin-1-yl)-1-
352 cyclopropyl-6- fluoro-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxylic acid (LB20304).
353 *J Med Chem* 1997; 40:3584-3593.
- 354 [5] Garcia I, Pascual A, Ballesta S, Joyanes P, Perea EJ. Intracellular penetration and
355 activity of gemifloxacin in human polymorphonuclear leukocytes. *Antimicrob Agents*
356 *Chemother* 2000; 44:3193-3195.
- 357 [6] Baltch AL, Bopp LH, Smith RP, Michelsen PB, Ritz WJ. Antibacterial activities of
358 gemifloxacin, levofloxacin, gatifloxacin, moxifloxacin and erythromycin against
359 intracellular *Legionella pneumophila* and *Legionella micdadei* in human monocytes. *J*
360 *Antimicrob Chemother* 2005; 56:104-109.

- 361 [7] Renard C, Vanderhaeghe HJ, Claes PJ, Zenebergh A, Tulkens PM. Influence of
362 conversion of penicillin G into a basic derivative on its accumulation and subcellular
363 localization in cultured macrophages. *Antimicrob Agents Chemother* 1987; 31:410-416.
- 364 [8] Marquez B, Caceres NE, Mingeot-Leclercq MP, Tulkens PM, Van Bambeke F.
365 Identification of the efflux transporter of the fluoroquinolone antibiotic ciprofloxacin in
366 murine macrophages: studies with ciprofloxacin-resistant cells. *Antimicrob Agents*
367 *Chemother* 2009; 53:2410-2416.
- 368 [9] Michot JM, Van Bambeke F, Mingeot-Leclercq MP, Tulkens PM. Active efflux of
369 ciprofloxacin from J774 macrophages through an MRP-like transporter. *Antimicrob*
370 *Agents Chemother* 2004; 48:2673-2682.
- 371 [10] Michot JM, Seral C, Van Bambeke F, Mingeot-Leclercq MP, Tulkens PM. Influence of
372 efflux transporters on the accumulation and efflux of four quinolones (ciprofloxacin,
373 levofloxacin, garenoxacin, and moxifloxacin) in J774 macrophages. *Antimicrob Agents*
374 *Chemother* 2005; 49:2429-2437.
- 375 [11] Michot JM, Heremans MF, Caceres NE, Mingeot-Leclercq MP, Tulkens PM, Van
376 Bambeke F. Cellular accumulation and activity of quinolones in ciprofloxacin-resistant
377 J774 macrophages. *Antimicrob Agents Chemother* 2006; 50:1689-1695.
- 378 [12] Tsuchiya S, Yamabe M, Yamaguchi Y, Kobayashi Y, Konno T, Tada K. Establishment
379 and characterization of a human acute monocytic leukemia cell line (THP-1). *Int J*
380 *Cancer* 1980; 26:171-176.
- 381 [13] Carryn S, Van Bambeke F, Mingeot-Leclercq MP, Tulkens PM. Comparative
382 intracellular (THP-1 macrophage) and extracellular activities of beta-lactams,
383 azithromycin, gentamicin, and fluoroquinolones against *Listeria monocytogenes* at
384 clinically relevant concentrations. *Antimicrob Agents Chemother* 2002; 46:2095-2103.

- 385 [14] Barcia-Macay M, Seral C, Mingeot-Leclercq MP, Tulkens PM, Van Bambeke F.
386 Pharmacodynamic evaluation of the intracellular activities of antibiotics against
387 *Staphylococcus aureus* in a model of THP-1 macrophages. *Antimicrob Agents*
388 *Chemother* 2006; 50:841-851.
- 389 [15] Lowry OH, N.J.Rosebrough, A.L.Farr, R.J.Randall. Protein measurement with the Folin
390 phenol reagent. *J Biol Chem* 1951; 193:265-275.
- 391 [16] Anonymous. *Methods for dilution antimicrobial susceptibility tests for bacteria that grow*
392 *aerobically - Approved standard*. Clinical and Laboratory Standard Institute, Wayne,
393 PA, 2010.
- 394 [17] Lemaire S, Van Bambeke F, Mingeot-Leclercq MP, Tulkens PM. Activity of three {beta}-
395 lactams (ertapenem, meropenem and ampicillin) against intraphagocytic *Listeria*
396 *monocytogenes* and *Staphylococcus aureus*. *J Antimicrob Chemother* 2005; 55:897-
397 904.
- 398 [18] Lemaire S, Van Bambeke F, Mingeot-Leclercq MP, Glupczynski Y, Tulkens PM. Role of
399 acidic pH in the susceptibility of intraphagocytic methicillin-resistant *Staphylococcus*
400 *aureus* strains to meropenem and cloxacillin. *Antimicrob Agents Chemother* 2007;
401 51:1627-1632.
- 402 [19] Lemaire S, Kosowska-Shick K, Appelbaum PC, Glupczynski Y, Van Bambeke F,
403 Tulkens PM. Activity of moxifloxacin against intracellular community-acquired
404 methicillin-resistant *Staphylococcus aureus*: comparison with clindamycin, linezolid and
405 co-trimoxazole and attempt at defining an intracellular susceptibility breakpoint. *J*
406 *Antimicrob Chemother* 2011; 66:596-607.
- 407 [20] Nguyen HA, Denis O, Vergison A, Theunis A, Tulkens PM, Struelens MJ et al.
408 Intracellular activity of antibiotics in a model of human THP-1 macrophages infected by
409 a *Staphylococcus aureus* small-colony variant strain isolated from a cystic fibrosis

- 410 patient: pharmacodynamic evaluation and comparison with isogenic normal-phenotype
411 and revertant strains. *Antimicrob Agents Chemother* 2009; 53:1434-1442.
- 412 [21] LG Life Sciences. LG Life Sciences 2009 Annual Report. 2010. Seoul, Korea, p. 18.
- 413 [22] Cormican MG, Jones RN. Antimicrobial activity and spectrum of LB20304, a novel
414 fluoronaphthyridone. *Antimicrob Agents Chemother* 1997; 41:204-211.
- 415 [23] Oh JI, Paek KS, Ahn MJ, Kim MY, Hong CY, Kim IC et al. In vitro and in vivo
416 evaluations of LB20304, a new fluoronaphthyridone. *Antimicrob Agents Chemother*
417 1996; 40:1564-1568.
- 418 [24] Roy B, Das A, Bhaumik U, Sarkar AK, Bose A, Mukharjee J et al. Determination of
419 gemifloxacin in different tissues of rat after oral dosing of gemifloxacin mesylate by LC-
420 MS/MS and its application in drug tissue distribution study. *J Pharm Biomed Anal* 2010;
421 52:216-226.
- 422 [25] Saravolatz LD, Leggett J. Gatifloxacin, gemifloxacin, and moxifloxacin: the role of 3
423 newer fluoroquinolones. *Clin Infect Dis* 2003; 37:1210-1215.
- 424 [26] Mandell LA, Iannini PB, Tillotson GS. Respiratory fluoroquinolones: differences in the
425 details. *Clin Infect Dis* 2004; 38:1331-1332.
- 426 [27] Capitano B, Mattoes HM, Shore E, O'Brien A, Braman S, Sutherland C et al. Steady-
427 state intrapulmonary concentrations of moxifloxacin, levofloxacin, and azithromycin in
428 older adults. *Chest* 2004; 125:965-973.
- 429 [28] Schuler P, Zemper K, Borner K, Koeppe P, Schaberg T, Lode H. Penetration of
430 sparfloxacin and ciprofloxacin into alveolar macrophages, epithelial lining fluid, and
431 polymorphonuclear leucocytes. *Eur Respir J* 1997; 10:1130-1136.

- 432 [29] Hara T, Takemura H, Kanemitsu K, Yamamoto H, Shimada J. Comparative uptake of
433 grepafloxacin and ciprofloxacin by a human monocytic cell line, THP-1. *J Infect*
434 *Chemother* 2000; 6:162-167.
- 435 [30] Bounds SJ, Nakkula R, Walters JD. Fluoroquinolone transport by human monocytes:
436 characterization and comparison to other cells of myeloid lineage. *Antimicrob Agents*
437 *Chemother* 2000; 44:2609-2614.
- 438 [31] Memin E, Panteix G, Revol A. Carrier-mediated system for pefloxacin uptake in human
439 monocytes. *J Antimicrob Chemother* 1997; 40:263-268.
- 440 [32] Skazik C, Heise R, Bostanci O, Paul N, Denecke B, Jousseen S et al. Differential
441 expression of influx and efflux transport proteins in human antigen presenting cells. *Exp*
442 *Dermatol* 2008; 17:739-747.
- 443 [33] Fresta M, Guccione S, Beccari AR, Furneri PM, Puglisi G. Combining molecular
444 modeling with experimental methodologies: mechanism of membrane permeation and
445 accumulation of ofloxacin. *Bioorg Med Chem* 2002; 10:3871-3889.
- 446 [34] de Duve C, de Barse T, Poole B, Trouet A, Tulkens P, Van Hoof F. Commentary.
447 Lysosomotropic agents. *Biochem Pharmacol* 1974; 23:2495-2531.
- 448 [35] Carlier MB, Scorneaux B, Zenebergh A, Desnottes JF, Tulkens PM. Cellular uptake,
449 localization and activity of fluoroquinolones in uninfected and infected macrophages. *J*
450 *Antimicrob Chemother* 1990; 26 Suppl B:27-39.
- 451 [36] Carlier MB, Zenebergh A, Tulkens PM. Cellular uptake and subcellular distribution of
452 roxithromycin and erythromycin in phagocytic cells. *J Antimicrob Chemother* 1987; 20
453 Suppl B:47-56.

- 454 [37] Carlier MB, Garcia-Luque I, Montenez JP, Tulkens PM, Piret J. Accumulation, release
455 and subcellular localization of azithromycin in phagocytic and non-phagocytic cells in
456 culture. *Int J Tissue React* 1994; 16:211-220.
- 457 [38] Lode H, Hoffken G, Boeck M, Deppermann N, Borner K, Koeppe P. Quinolone
458 pharmacokinetics and metabolism. *J Antimicrob Chemother* 1990; 26 Suppl B:41-49.
- 459 [39] Dudley MN. Pharmacokinetics of fluoroquinolones. In: Hooper DC, Wolfson JS, editors.
460 Quinolone Antimicrobial Agents. Washington, DC: ASM Press, 2003: 115-132.
- 461 [40] Schmidt S, Rock K, Sahre M, Burkhardt O, Brunner M, Lobmeyer MT et al. Effect of
462 protein binding on the pharmacological activity of highly bound antibiotics. *Antimicrob
463 Agents Chemother* 2008; 52:3994-4000.
- 464 [41] Paillard D, Grellet J, Dubois V, Saux MC, Quentin C. Discrepancy between uptake and
465 intracellular activity of moxifloxacin in a *Staphylococcus aureus*-human THP-1
466 monocytic cell model. *Antimicrob Agents Chemother* 2002; 46:288-293.
- 467 [42] Seral C, Barcia-Macay M, Mingeot-Leclercq MP, Tulkens PM, Van Bambeke F.
468 Comparative activity of quinolones (ciprofloxacin, levofloxacin, moxifloxacin and
469 garenoxacin) against extracellular and intracellular infection by *Listeria monocytogenes*
470 and *Staphylococcus aureus* in J774 macrophages. *J Antimicrob Chemother* 2005;
471 55:511-517.
- 472 [43] Van de Velde S, Nguyen HA, Van Bambeke F, Tulkens PM, Grellet J, Dubois V et al.
473 Contrasting effects of human THP-1 cell differentiation on levofloxacin and moxifloxacin
474 intracellular accumulation and activity against *Staphylococcus aureus* and *Listeria
475 monocytogenes*. *J Antimicrob Chemother* 2008; 62:518-521.

476

477

478 **Table 1:** Pertinent regression parameters ^a (with confidence intervals [CI]) and statistical analyses of data from experiments examining the
 479 concentration-response activities of ciprofloxacin, moxifloxacin and gemifloxacin (shown in Figure 3A). The 3 first rows show the
 480 analysis for each individual antibiotic and the last row the analysis made for all pooled data.

antibiotic	<i>L. monocytogenes</i>					<i>S. aureus</i>				
	R ²	E _{min} ^b (CI)	E _{max} ^c (CI)	EC ₅₀ ^d (CI)	C _s ^e	R ²	E _{min} ^b (CI)	E _{max} ^c (CI)	EC ₅₀ ^d (CI)	C _s ^e
ciprofloxacin	0.95	2.82 (1.83 to 3.81)	-3.96 (-5.44 to -2.48)	2.94 (1.00 to 8.66)	0.49	0.96	3.80 (2.92 to 4.68)	-1.60 (-2.26 to -0.94)	1.36 (0.64 to 2.89)	3.1
moxifloxacin	0.95	2.84 (2.05 to 3.63)	-4.48 (-5.21 to -3.56)	1.30 (0.62 to 2.74)	0.47	0.95	2.98 (2.11 to 3.86)	-1.85 (-2.28 to -1.43)	1.62 (0.78 to 3.38)	2.6
gemifloxacin	0.97	3.03 (1.79 to 4.27)	-3.55 (-4.28 to -2.82)	0.65 (0.27 to 1.59)	0.74	0.98	3.07 (2.32 to 3.83)	-1.30 (-1.64 to -0.97)	2.02 (1.01 to 4.03)	4.8
all 3	0.91	3.44 (2.95 to 3.94)	-3.92 (-4.53 to -3.32)	1.44 (0.95 to 2.19)	0.93	0.94	3.44 (2.95 to 3.94)	-1.55 (-1.81 to -1.29)	1.44 (0.95 to 2.19)	3.2

481

482 ^a by use of all data for antibiotic concentrations ranging from approx. 0.01 to approx. 1,000 x MIC (ciprofloxacin: 0.01 to 100 mg/L

483 [*L. monocytogenes*] and 0.001 to 100 mg/L [*S. aureus*]; moxifloxacin: 0.005 to 30 mg/L [*L. monocytogenes*] and 0.001 to 100 mg/L

484 [*S. aureus*]; gemifloxacin: 0.005 to 150 mg/L. [*L. monocytogenes*] and 0.0008 to 20 mg/L [*S. aureus*]).

485 ^b relative minimal efficacy: change in cfu (in log₁₀ units) at time 24 h from the initial, post-phagocytosis inoculum, as extrapolated for an infinitely
 486 low antibiotic concentration

487 ^c relative maximal efficacy: change in cfu (in log₁₀ units) at time 24 h from the initial, post-phagocytosis inoculum, as extrapolated for an
 488 infinitely large antibiotic concentration

489 ^d relative potency: extracellular concentration (in multiple of MIC) yielding a change in cfu at time 24 h half way between E_{\min} and E_{\max}
490 ^e static concentration: concentration (in multiples of MIC) resulting no apparent bacterial growth (no change in cfu) from the initial, post-
491 phagocytosis inoculum), as determined by graphical intrapolation (MIC values (mg/L) are : 1 and 0.125 (ciprofloxacin), 0.5 and 0.03
492 (moxifloxacin), and 0.5 and 0.008 (gemifloxacin) for *L. monocytogenes* and *S. aureus* respectively.


493

494 Statistical analysis:

495 The raw data obtained for each individual antibiotic, and the corresponding Hill functions were compared using one way ANOVA (parametric)
496 and Kuskal-Wallis (non-parametric) tests found to be not significantly different (*L. monocytogenes*: $p = 0.420$ and 0.152 , respectively: *S.*
497 *aureus*: $p = 0.351$ and 0.249 , respectively). The analysis was then repeated for comparison of antibiotic pairs (ciprofloxacin vs. gemifloxacin;
498 ciprofloxacin vs. moxifloxacin; moxifloxacin vs. gemifloxacin) using unpaired t-test and showed no significant difference for any comparison
499 ($p > 0.18$).

500

501 Figure 1
502



503


504

505 Caption to Figure 1: Accumulation and efflux of fluoroquinolones.

506 **A** (upper panel): accumulation of gemifloxacin (GMF), moxifloxacin (MXF), and
507 ciprofloxacin (CIP) in wild-type J774 mouse macrophages (left) and THP-1 human
508 monocytes (right) incubated during 2 h with 20 mg/L in control conditions (open bars)
509 or in the presence of the Mrp inhibitor gemfibrozil (500 µM; hatched bars). All values
510 are the means of 3 independent determinations ± SD. Statistical analysis (ANOVA):
511 control vs. gemfibrozil, *** $p < 0.001$; comparison of fluoroquinolones: bars with
512 different letters are different from one another ($p < 0.001$; caps letters, control
513 conditions; small letters, + gemfibrozil).

514 **B** (lower panel): kinetics of accumulation (left) and efflux (right) of gemifloxacin
 515 **compared** to ciprofloxacin in J774 macrophages (see [10] for efflux of moxifloxacin) .
 516 For accumulation studies, cells were transferred to medium containing a fixed
 517 amount of drug (20 mg/L) and collected at the times indicated in the abscissa. For
 518 efflux, cells were first exposed to the drug for 2 h at a concentration of 20 mg/L,
 519 gently washed, transferred to drug free medium and collected at the times indicated
 520 in the abscissa. Data were used to fit a one phase exponential association function
 521 for influx [$y = y_{max} x (1 - e^{-k_{in} x t})$] and a one phase exponential decay function
 522 for efflux [$y = y_{max} x e^{-k_{out} x t} + plateau$] by nonlinear regression. Regression
 523 parameters for influx: (a) gemifloxacin, $R^2 = 0.780$, $k_{in} = 0.386 \pm 0.123 \text{ min}^{-1}$; (b)
 524 ciprofloxacin, $R^2 = 0.922$, $k_{in} = 0.348 \pm 0.066 \text{ min}^{-1}$. Regression parameters for efflux:
 525 (1) main graph, (a) gemifloxacin, $R^2 = 0.897$, $k_{out} = 0.403 \pm 0.122 \text{ min}^{-1}$, $plateau =$
 526 25.71 ± 4.63 ; (b) ciprofloxacin, $R^2 = 0.949$, $k_{out} = 0.949 \pm 0.204 \text{ min}^{-1}$, $plateau =$
 527 3.56 ± 3.24); (2) Inset: data for the initial stage of efflux (0 – 5 min) and corrected for
 528 differences in plateau reached after 10 min, (a) gemifloxacin, $R^2 = 0.658$,
 529 $k_{out} = 0.571 \pm 0.138 \text{ min}^{-1}$; (b) ciprofloxacin, $R^2 = 0.909$, $k_{out} = 1.216 \pm 0.209 \text{ min}^{-1}$.
 530 Statistical analysis (paired *t*- test two-tailed): Influx, no significant difference in rate
 531 constants; absolute values of plateaus of accumulation were different and in line with
 532 data of Figure 1). Efflux: Main graph, comparison of all values: $p < 0.001$), plateaus
 533 values only: $p < 0.001$, k values only: $p < 0.001$; Inset: comparison for all values:
 534 $p = 0.016$, k_{out} values only: $p < 0.001$.
 535

536 Figure 2




537

538 Caption to Figure 2: Cellular accumulation of gemifloxacin as compared to
 539 ciprofloxacin in wild-type or ciprofloxacin-resistant J774 mouse macrophages (see
 540 [11] for the corresponding data with moxifloxacin) **A** (upper panel): cells were
 541 incubated for 2 h in control conditions (open bars) or in ATP-depleted cells (gray
 542 hatched bars) with a fixed concentration (20 mg/L) of gemifloxacin (left) or
 543 ciprofloxacin (right) with wild-type cells (WT) or ciprofloxacin-resistant cells (CIP-R).
 544 Data are expressed in percentage of the value measured in wild-type cells in control
 545 conditions for each fluoroquinolone. All values are the means of 3 independent
 546 determinations \pm SD. Statistical analysis (ANOVA): control vs. ATP-depletion, *** p
 547 < 0.001 ; WT vs. CIP-R cells: bars with different letters are different from one another
 548 ($p < 0.05$; upper case letters, control conditions; lower case letters, ATP-depletion).
 549 **B** (lower panel): Influence of the extracellular concentration of gemifloxacin (left) and
 550 ciprofloxacin (right) on their cellular to extracellular concentration ratio in wild-type

551 (closed symbols) or ciprofloxacin-resistant (open symbols) J774 mouse
552 macrophages, measured after 2 h of incubation. The cellular concentration was
553 expressed as μg per mg protein. Data are expressed in percentage of the highest
554 value observed in wild-type cells for each fluoroquinolone. All values are the means
555 of 3 independent determinations \pm SD.
556
557

Accepted Manuscript

558 Figure 3
559



560

561 Caption to Figure 3: Concentration-response of the activities of gemifloxacin,
562 moxifloxacin, and ciprofloxacin against phagocytized *L. monocytogenes* EGD (left)
563 and *S. aureus* ATCC25923 (right) in wild-type J774 macrophages.


564 **A** (top): after phagocytosis and elimination of the extracellular bacteria, cells were
565 incubated for 24 h with increasing concentrations of antibiotic (total drug) covering a
566 ~ 0.01 to ~ 1,000 x MIC range (MIC [mg/L] were 1 and 0.125 [ciprofloxacin], 0.5 and
567 0.03 [moxifloxacin], and 0.5 and 0.008 [gemifloxacin] for *L. monocytogenes* and
568 *S. aureus* respectively). The graphs show the change in the number of cfu (log
569 scale) per mg of cell protein compared to the initial post-phagocytosis inoculum

570 (ordinate) as a function of the extracellular concentration of each drug expressed in
571 multiple of its MIC (abscissa). In each graph, the horizontal dotted line corresponds
572 to an apparent static effect and the vertical line to the MIC of the drug. A single
573 sigmoidal regression has been fit to all data sets (see Figure SP2 for individual
574 regression curves. The pertinent regression parameters and numerical values of the
575 4 key pharmacological descriptors (E_{\min} , E_{\max} ; EC_{50} , C_s) are shown Table 1 for each
576 drug - bacteria combination.

577 **B** (bottom): the ordinates show the calculated cellular concentrations (total drug; in
578 multiples of MIC) needed to achieve two predefined activity levels (targets) shown on
579 the abscissa (static effect [no apparent change in cfu]; 2 (*L. monocytogenes*) or 1
580 (*S. aureus*) \log_{10} cfu decrease compared to the initial, post-phagocytosis inoculum).
581 The cellular concentrations were calculated by (i) using the concentration-response
582 curves shown in **A** to determine the extracellular concentrations needed to achieve
583 the target effects (graphical intrapolation), and (ii) using the data of Figure 2B (wild
584 type cells) to calculate the corresponding apparent total cellular concentrations of
585 gemifloxacin and ciprofloxacin (for moxifloxacin, we used the accumulation data
586 published in [10]) based on a conversion factor of 3.08 μ L of total cell volume per mg
587 protein as determined experimentally for wild type J774 macrophages in previous
588 studies [9].

589

590 **Figure 4**
591



592
593 Caption for Figure 4: Subcellular distribution of gemifloxacin (GMF), moxifloxacin
594 (MXF), and ciprofloxacin (CIP) in J774 mouse macrophages incubated for 2 h with
595 50 mg/L of each drug. The upper panel shows the antibiotic content in the
596 nuclear/unbroken cells, organelles, and soluble fractions expressed in percentage of
597 the total recovered amount (each bar corresponds to a separate experiment). The
598 lower panel shows the distribution of lactate dehydrogenase (LDH; marker of the
599 cytosol), cytochrome-c-oxidase (CYTOX; marker of mitochondria), and N-acetyl- β -
600 hexosaminidase (NAB; marker of lysosomes) as the mean values (\pm SD) of the 3
601 experiments (corresponding to each of the individual experiment shown in the upper
602 panel).
603


604 Vallet *et al.* Intracellular activity of fluoroquinolones - Supplementary Material

605

606

607

Figure SP1



608


609 Caption to Figure SP1: Kinetics of efflux of gemifloxacin from wild-type and

610 ciprofloxacin-resistant J774 cells. Cells were exposed to gemifloxacin (20 mg/L) for

611 2 h and then transferred to antibiotic-free medium for up to 30 min. Regression

612 parameters: (a) wild-type cells (same data as in Figure 2), $R^2 = 0.896$,613 $k_{out} = 0.403 \pm 0.122 \text{ min}^{-1}$, $plateau = 25.71 \pm 4.63$; (b) ciprofloxacin-resistant cells,614 $R^2 = 0.830$, $k_{out} = 2.39 \pm 0.907 \text{ min}^{-1}$, $plateau = 31.6 \pm 4.0$).

615

616 **Figure SP2**
617

618
619 Caption to Figure SP2: Concentration-response of the activities of gemifloxacin,
620 moxifloxacin, and ciprofloxacin (CIP) against *S. aureus* ATCC25923 (top) and
621 *L. monocytogenes* EGD (bottom) in wild-type J774 macrophages. Cells were
622 incubated with increasing concentrations of antibiotic (total drug) for 24 h. Each
623 graph shows the change in the number of cfu (log scale) per mg of cell protein
624 compared to the initial post-phagocytosis inoculum (ordinate) as a function of the
625 extracellular concentration of each drug expressed in multiples of its MIC (abscissa).
626 In each graph, the horizontal dotted line corresponds to an apparent static effect and
627 the vertical line to the MIC of the drug. A sigmoidal regression has been fitted to
628 each set of data (see Table 1 for the pertinent regression parameters and numerical
629 values of the four key pharmacological descriptors (E_{\min} , E_{\max} , EC_{50} , C_s)).

630
631

632 **Table SP1: Physico-chemical properties of fluoroquinolones and azithromycin at physiologically-relevant pHs**

633 The data indicate that (i) ciprofloxacin, moxifloxacin, and gemifloxacin display quite similar biophysical properties although showing distinct cellular
 634 accumulation levels (gemifloxacin > moxifloxacin > ciprofloxacin; see Results) that are not correlated to the minor differences seen; (ii) these
 635 properties are very different from those of azithromycin, a drug known to accumulate extensively in lysosomes by proton-trapping (see
 636 Discussion). The pHs considered are those of the extracellular (7-7.4) and of the lysosomal (5-5.4) milieus, respectively.

637

Drug	pKa ₁ ^a (acidic)	pKa ₂ ^a (basic)	species in solution (calculated %) ^a						logP ^b		calculated logD ^{a,c}	
			pH 7.4			pH 5.4			calculated ^{a,d}	experimental ^e	pH 7	pH 5
			cationic	zwitterionic	anionic	cationic	zwitterionic	anionic				
ciprofloxacin	5.8	8.7	2	93	5	69	30	0	1.63 ^c	2.30	-1.38	-1.62
moxifloxacin	5.6	9.4	2	97	1	66	34	0	1.90	2.90	-1.72	-1.33
gemifloxacin	5.5	9.5	1	98	1	47	52	0	1.04	2.30	-2.54	-1.78
azithromycin	-	8.9 ^f 9.6	96.98^g	0.02	0	99.97^g	0	0	2.44	4.02	-1.99	-4.41

638

639 ^a calculated using Reaxys (<http://www.reaxys.com>) with the ChemAxon's Marvin plug-in calculators (<http://www.chemaxon.com/marvin>). The
 640 actual values of the pK_a of the acidic function may be about 0.5 units higher due to the influence of the vicinal carbonyl function [1].

641 ^b logP: partition coefficient (log of the ratio of the concentrations of the unionized compound between a non polar [octanol] and a polar [water]
 642 phases);

643 ^c logD: distribution coefficient (log of the ratio of the sum of the concentrations of all forms of the compound [ionized plus un-ionized] in each of
644 the two phases at a given pH)

645 ^d These values are the arithmetic average of three methods of calculations (Viswanadhan's fragmentation; Klopman's fragmentation; and
646 PHYSPROP© database [see https://www.reaxys.com/static/marvin/marvin_5_3_7/help/calculations/partitioning.html for details]).

647 ^e value as reported in Drugbank (see <http://www.drugbank.ca> and [2])

648 ^f azithromycin is a dicationic drug

649 ^g dicationic form (monocationic form: 3 % at pH 7.4 and 0.03 % at pH 5.4; a zwitterionic form is virtually inexistent (< 0.001 %) at these pH
650 values).

651

652 References

653

654 [1] Nikaido H, Thanassi DG. Penetration of lipophilic agents with multiple protonation sites into bacterial cells: tetracyclines and
655 fluoroquinolones as examples. *Antimicrob Agents Chemother.* 1993 Jul;37(7):1393-9. [PMID: 8363364]

656

657 [2] Wishart DS, Knox C, Guo AC, Shrivastava S, Hassanali M, Stothard P, Chang Z, Woolsey J: DrugBank, a comprehensive resource for *in*
658 *silico* drug discovery and exploration, *Nucleic Acids Res.* 2006 Jan 1;34 (Database issue): D668-72 [PMID: 16381955]

659

660