



HAL
open science

Description of the minimizers of least squares regularized with ℓ_0 -norm. Uniqueness of the global minimizer

Mila Nikolova

► To cite this version:

Mila Nikolova. Description of the minimizers of least squares regularized with ℓ_0 -norm. Uniqueness of the global minimizer. 2013. hal-00723812v4

HAL Id: hal-00723812

<https://hal.science/hal-00723812v4>

Preprint submitted on 15 Jan 2013 (v4), last revised 10 Nov 2014 (v7)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Description of the minimizers of least squares regularized with ℓ_0 -norm. Uniqueness of the global minimizer

Mila NIKOLOVA

CMLA, ENS Cachan, CNRS, 61 Avenue du President Wilson, F-94230 Cachan, France
nikolova@cmla.ens-cachan.fr (Submitted: November 10, 2011. Revised: January 11, 2013)

Abstract. We have an $M \times N$ real-valued arbitrary matrix A (e. g. a dictionary) with $M < N$ and data d describing the sought-after object with the help of A . This work provides an in-depth analysis of the (local and global) minimizers of an objective function \mathcal{F}_d combining a quadratic data-fidelity term and an ℓ_0 penalty applied to each entry of the sought after solution, weighted by a regularization parameter $\beta > 0$. For several decades, this objective focuses a ceaseless effort to conceive algorithms approaching a good minimizer. Our theoretical contributions, summarized below, shed new light on the existing algorithms and can help the conception of innovative numerical schemes.

To solve the normal equation associated with any M -row submatrix of A is equivalent to compute a local minimizer \hat{u} of \mathcal{F}_d . (Local) minimizers \hat{u} of \mathcal{F}_d are strict if and only if the submatrix, composed of those columns of A whose indexes form the support of \hat{u} , has full column rank. An outcome is that strict local minimizers of \mathcal{F}_d are easily computed without knowing the value of β . Each strict local minimizer is linear in data. It is proved that \mathcal{F}_d has global minimizers and that they are always strict. They are studied in more details under the (standard) assumption that $\text{rank}(A) = M < N$. The global minimizers with M -length support are seen to be impractical. Given d , critical values β_K for any $K \leq M - 1$ are exhibited such that if $\beta > \beta_K$, all global minimizers of \mathcal{F}_d are K -sparse.

An assumption on A is adopted and proved to fail only on a closed negligible subset. Then for all data d beyond a closed negligible subset, the objective \mathcal{F}_d for $\beta > \beta_K$, $K \leq M - 1$, has a unique global minimizer and the latter is K -sparse.

Instructive small-size (5×10) numerical illustrations confirm the main theoretical results.

Key words: asymptotically level stable functions, global minimizers, local minimizers, ℓ_0 regularization, nonconvex nonsmooth minimization, perturbation analysis, quadratic programming, solution analysis, sparse recovery, strict minimizers, under-determined linear systems, uniqueness of the solution, variational methods.

1 Introduction

Let A be an arbitrary matrix (e. g. a dictionary) such that

$$A \in \mathbb{R}^{M \times N} \quad \text{for} \quad M < N,$$

where the positive integers M and N are fixed. Given a data vector $d \in \mathbb{R}^M$, we consider an objective function $\mathcal{F}_d : \mathbb{R}^N \rightarrow \mathbb{R}$ of the form

$$\begin{aligned} \mathcal{F}_d(u) &= \|Au - d\|_2^2 + \beta \|u\|_0, \quad \beta > 0, \\ \|u\|_0 &= \#\sigma(u), \end{aligned} \tag{1}$$

where $u \in \mathbb{R}^N$ contains the coefficients describing the sought-after object, β is a regularization parameter, $\#$ stands for cardinality and $\sigma(u)$ is the support of u (i. e. the set of all $i \in \{1, \dots, N\}$ for which the i th entry of u meets $u[i] \neq 0$). By an abuse of language, the penalty in (1) is called the ℓ_0 -norm. Define $\phi : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\phi(t) \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } t = 0, \\ 1 & \text{if } t \neq 0. \end{cases} \quad (2)$$

Then $\|u\|_0 = \sum_{i=1}^N \phi(u[i]) = \sum_{i \in \sigma(u)} \phi(u[i])$, so \mathcal{F}_d in (1) equivalently reads

$$\mathcal{F}_d(u) = \|Au - d\|_2^2 + \beta \sum_{i=1}^N \phi(u[i]) \quad (3)$$

$$= \|Au - d\|_2^2 + \beta \sum_{i \in \sigma(u)} \phi(u[i]). \quad (4)$$

We focus on all (local and global) minimizers \hat{u} of an objective \mathcal{F}_d of the form (1):

$$\hat{u} \in \mathbb{R}^N \text{ such that } \mathcal{F}_d(\hat{u}) = \min_{u \in \mathcal{O}} \mathcal{F}_d(u), \quad (5)$$

where \mathcal{O} is an open neighborhood of \hat{u} . We note that finding the global minimizer of \mathcal{F}_d must be an *NP-hard* computational problem [10, 39].

The function ϕ in (2) served as a regularizer for a long time. In the context of Markov random fields it was used by Geman and Geman in 1984 [19] and Besag in 1986 [5] as a prior in MAP energies to restore labeled images. The MAP objective reads

$$\mathcal{F}_d(u) = \|Au - d\|_2^2 + \beta \sum_k \phi(D_k u), \quad (6)$$

where D_k is a finite differences operator and ϕ is given by (2). This label-designed form is known as the Potts prior model, or as the multi-level logistic model [6, 23]. Various stochastic and deterministic algorithms were considered to minimize (6). Leclerc [22] proposed in 1989 a deterministic continuation method to restore piecewise constant images. Robini and Magnin [32] introduced the stochastic continuation approach and successfully used it to reconstruct 3D tomographic images. They refined the method and the theory in [33]. Very recently, Robini and Reissman [34] gave theoretical results relating the probability for global convergence and the computation speed.

The problem stated in (1) and (5)—to (locally) minimize \mathcal{F}_d —arises when *sparse* solutions are desired. Typical application fields are signal and image processing, morphologic component analysis, compression, dictionary building, inverse problems, compressive sensing, machine learning, model selection, classification, subset selection, among others. The original hard-thresholding method proposed by Donoho and Johnstone [14] amounts to¹ minimize \mathcal{F}_d where d contains the coefficients of a signal or an image expanded in a wavelet basis ($M = N$). When $M < N$ various (usually strong) restrictions on $\|u\|_0$ (often $\|u\|_0$ is replaced by a less irregular function) and on A (e.g. RIP-like criteria, conditions on $\|A\|$, etc.) are needed to conceive

¹For reminder: if d are some noisy coefficients, the restored coefficients \hat{u} minimize $\|u - d\|^2 + \beta\|u\|_0$ and read $\hat{u}[i] = 0$ if $|d[i]| \leq \sqrt{\beta}$ and $\hat{u}[i] = d[i]$ otherwise.

numerical schemes approximating a minimizer of \mathcal{F}_d , to establish local convergence and derive the asymptotic of the obtained solution. In statistics the problem was widely considered for subset selection and numerous algorithms were designed, with limited theoretical production, as explained in the book of Miller [29]. More recently, Haupt and Nowak [21] investigate the statistical performances of the global minimizer of \mathcal{F}_d and propose an iterative bound-optimization procedure. Fan and Li [16] discuss a variable splitting and penalty decomposition minimization technique for (1), along with other approximations of the ℓ_0 -norm. Liu and Wu [24] mix the ℓ_0 and the ℓ_1 penalties, establish some asymptotic properties of the new estimator and use mixed integer programming aimed at global minimization. For model selection, Lv and Fan [26] approximate the ℓ_0 penalty using functions that are concave on \mathbb{R}_+ and prove a non asymptotic nearly oracle property of the resultant estimator. Thiao *et al.* [38] reformulate the problem so that an approximate solution can be found using difference of convex functions programming. Blumensath and Davies [7] propose an iterative thresholding scheme to approximate a solution and prove convergence to a local minimizer of \mathcal{F}_d . Lu and Zhang [25] suggest a penalty decomposition method to minimize \mathcal{F}_d . Fornasier and Ward [17] propose an iterative thresholding algorithm for finding local minimizers of \mathcal{F}_d where ℓ_0 is replaced by a reasonable sparsity-promoting relaxation given by $\phi(t) = \min\{t, 1\}$; convergence to a local minimizer is established. In a recent report, Chouzenoux *et al.* [9], a mixed $\ell_2 - \ell_0$ regularization is considered: a slightly smoothed version of the objective is analyzed and optimization using a majorize-minimize subspace approach is shown to converge to a local minimizer in a finite number of iterations. Since the submission of our paper, image reconstruction methods were designed where ℓ_0 regularization is applied to the coefficients of the expansion of the sought-after image in a wavelet frame [42, 13]: the provided numerical results outperform ℓ_1 regularization for a reasonable computational cost achieved using penalty decomposition techniques. In a general study on the convergence of descent methods for nonconvex objectives, Attouch *et al.* [1] apply an inexact forward-backward splitting scheme to find a critical point of \mathcal{F}_d . Several other references can be evoked, e.g. [30, 18].

Even though overlooked for several decades, the objective \mathcal{F}_d was essentially considered from a numerical standpoint. The motivation naturally comes from the promising applications and the intrinsic difficulty to minimize \mathcal{F}_d .

The goal of this work is to analyze the (local and global) minimizers \hat{u} of objectives \mathcal{F}_d of the form (1).

We provide detailed results on the minimization problem.

The uniqueness of the global minimizer of \mathcal{F}_d is examined as well.

We do not propose an algorithm. However, our theoretical results raise salient cross-questions to the existing algorithms and can help the conception of innovative numerical schemes.

The minimization of \mathcal{F}_d in (1) might seem close to its constraint variants:

$$\begin{aligned} \text{given } \varepsilon \geq 0, \quad & \text{minimize } \|u\|_0 \quad \text{subject to } \|Au - d\|^2 \leq \varepsilon, \\ \text{given } K \in \mathbb{I}_M, \quad & \text{minimize } \|Au - d\|^2 \quad \text{subject to } \|u\|_0 \leq K. \end{aligned} \tag{7}$$

The latter problems are abundantly studied in the context of sparse recovery in different fields. An excellent account is given in [8], see also the book [27]. For recent achievements, we refer to [40]. It is worth emphasizing that in general, *there is no equivalence between the problems stated in (7) and the minimization of \mathcal{F}_d in (1)* because all these problems are nonconvex.

1.1 Main notations and definitions

We recall that if \hat{u} is a (local) *minimizer* of \mathcal{F}_d , the value $\mathcal{F}_d(\hat{u})$ is a (local) minimum² of \mathcal{F}_d reached at (possibly numerous) points \hat{u} . Saying that a (local) minimizer \hat{u} is *strict* means that there is a neighborhood $\mathcal{O} \subset \mathbb{R}^N$, containing \hat{u} , such that $\mathcal{F}_d(\hat{u}) < \mathcal{F}_d(v)$ for any $v \in \mathcal{O} \setminus \{\hat{u}\}$. So \hat{u} is an isolated minimizer.

Let K be any positive integer. The expression $\{u \in \mathbb{R}^K : u \text{ satisfying property } \mathfrak{P}\}$ designates the subset of \mathbb{R}^K formed from all elements u that meet \mathfrak{P} . The identity operator on \mathbb{R}^K is denoted by I_K . The entries of a vector $u \in \mathbb{R}^K$ read $u[i]$, for any i . The i th vector of the canonical basis³ of \mathbb{R}^K is denoted by $e_i \in \mathbb{R}^K$. Given $u \in \mathbb{R}^K$ and $\rho > 0$, the *open ball* at u of radius ρ with respect to the ℓ_p -norm for $1 \leq p \leq \infty$ reads

$$B_p(u, \rho) \stackrel{\text{def}}{=} \{v \in \mathbb{R}^K : \|v - u\|_p < \rho\} .$$

To simplify the notations, the ℓ_2 -norm is *systematically* denoted by

$$\|\cdot\| \stackrel{\text{def}}{=} \|\cdot\|_2 .$$

We denote by \mathbb{I}_K the *totally and strictly ordered* index set

$$\mathbb{I}_K \stackrel{\text{def}}{=} (\{1, \dots, K\}, <) . \quad (8)$$

where the symbol “ $<$ ” stands for the natural order of the positive integers. Accordingly,

any subset $\omega \subseteq \mathbb{I}_N$ inherits the property of being totally and strictly ordered.

We shall often consider the index set \mathbb{I}_N . The complement of $\omega \subseteq \mathbb{I}_N$ in \mathbb{I}_N is denoted by

$$\omega^c = \mathbb{I}_N \setminus \omega \subseteq \mathbb{I}_N .$$

Definition 1 For any $u \in \mathbb{R}^N$, the *support* $\sigma(u)$ of u is defined by

$$\sigma(u) = \{i \in \mathbb{I}_N : u[i] \neq 0\} \subseteq \mathbb{I}_N .$$

If $u = 0$, clearly $\sigma(u) = \emptyset$.

The i th column in a matrix $A \in \mathbb{R}^{M \times N}$ is denoted by a_i . It is *systematically* assumed that

$$a_i \neq 0, \quad \forall i \in \mathbb{I}_N . \quad (9)$$

For a matrix $A \in \mathbb{R}^{M \times N}$ and a vector $u \in \mathbb{R}^N$, with any $\omega \subseteq \mathbb{I}_N$, we associate the *submatrix* A_ω and the *subvector* u_ω given by

$$A_\omega \stackrel{\text{def}}{=} (a_{\omega[1]}, \dots, a_{\omega[\#\omega]}) \in \mathbb{R}^{M \times \#\omega} , \quad (10)$$

$$u_\omega \stackrel{\text{def}}{=} (u[\omega[1]], \dots, u[\omega[\#\omega]]) \in \mathbb{R}^{\#\omega} , \quad (11)$$

respectively, as well as the zero padding operator $Z_\omega : \mathbb{R}^{\#\omega} \rightarrow \mathbb{R}^N$ that inverts (11):

$$u = Z_\omega(u_\omega) \quad u[i] = \begin{cases} 0 & \text{if } i \notin \omega , \\ u_\omega[k] & \text{for the unique } k \text{ such that } \omega[k] = i . \end{cases} \quad (12)$$

²These two terms are often assimilated in the literature.

³More precisely, for any $i \in \mathbb{I}_K$, the vector $e_i \in \mathbb{R}^K$ is defined by $e_i[i] = 1$ and $e_i[j] = 0, \forall j \in \mathbb{I}_K \setminus \{i\}$.

Thus for $\omega = \emptyset$ one finds $u_\emptyset = \emptyset$ and $u = Z_\emptyset(u_\emptyset) = 0 \in \mathbb{R}^N$.

Using Definition 1 and the notations in (10)-(11), for any $u \in \mathbb{R}^N \setminus \{0\}$ we have

$$\omega \in \mathbb{I}_N \text{ and } \omega \supseteq \sigma(u) \Rightarrow Au = A_\omega u_\omega. \quad (13)$$

So as to simplify the presentation, we adopt the following *definitions*⁴:

$$\begin{aligned} (a) \quad & A_\emptyset = [] \in \mathbb{R}^{M \times 0}, \\ (b) \quad & \text{rank}(A_\emptyset) = 0. \end{aligned} \quad (14)$$

In order to avoid possible ambiguities⁵, we set

$$A_\omega^T \stackrel{\text{def}}{=} (A_\omega)^T,$$

where the superscript “ T ” stands for transposed. If A_ω is invertible, similarly $A_\omega^{-1} \stackrel{\text{def}}{=} (A_\omega)^{-1}$.

In the course of this work, we shall frequently refer to the constrained quadratic optimization problem stated next.

Given $d \in \mathbb{R}^M$ and $\omega \subseteq \mathbb{I}_N$, problem (\mathcal{P}_ω) reads:

$$(\mathcal{P}_\omega) \quad \boxed{\begin{cases} \min_{u \in \mathbb{R}^N} \|Au - d\|^2, \\ \text{subject to } u[i] = 0, \forall i \in \omega^c. \end{cases}} \quad (15)$$

The definition below will be used to evaluate the extent of some subsets and assumptions.

Definition 2 *A property (or an assumption) is called generic on \mathbb{R}^K if it holds true on a dense and open subset of \mathbb{R}^K .*

As usual, a subset $\mathcal{S} \subset \mathbb{R}^K$ is said to be *negligible* in \mathbb{R}^K if there exists $\mathcal{Z} \subset \mathbb{R}^K$ whose Lebesgue measure in \mathbb{R}^K is $\mathbb{L}^K(\mathcal{Z}) = 0$ and $\mathcal{S} \subseteq \mathcal{Z}$. If a property fails only on a negligible set, then it is said to hold *almost everywhere*, meaning “with probability one”.

Definition 2 requires much more than *almost everywhere*:

If a property holds true for all $v \in \mathbb{R}^K \setminus \mathcal{S}$, where $\mathcal{S} \subseteq \mathcal{Z} \subset \mathbb{R}^K$, \mathcal{Z} is closed in \mathbb{R}^K and $\mathbb{L}^K(\mathcal{Z}) = 0$, then this property is *generic* on \mathbb{R}^K .

Indeed, $\mathbb{R}^K \setminus \mathcal{Z}$ contains a *dense open* subset of \mathbb{R}^K . So if a property is generic on \mathbb{R}^K , then it holds true almost everywhere on \mathbb{R}^K . But the converse is false: an almost everywhere true property fails on a negligible subset whose closure may have positive measure⁶ in which case $\mathbb{R}^K \setminus \mathcal{Z}$ does not contain an open subset of \mathbb{R}^K .

In this sense, a generic property is stable with respect to the objects it concerns.

The elements of a set $\mathcal{S} \subset \mathbb{R}^K$ where a generic property fails are highly exceptional in \mathbb{R}^K .

The chance that a truly random $v \in \mathbb{R}^K$ —i. e. a v following a non singular probability distribution on \mathbb{R}^K —comes across such an \mathcal{S} can be ignored in practice.

⁴Note that (a) corresponds to the zero mapping on \mathbb{R}^0 and that (b) is the usual definition for the rank of an empty matrix.

⁵In the light of (10), A_ω^T could also mean $(A^T)_\omega$.

⁶There are many examples—e.g. $\mathcal{Z} = \{x \in [0, 1] : x \text{ is rational}\}$, then $\mathbb{L}^1(\mathcal{Z}) = 0$ and $\mathbb{L}^1(\text{closure}(\mathcal{Z})) = 1$.

1.2 Content of the paper

The main result in section 2 tells us that finding a solution of (\mathcal{P}_ω) for $\omega \subset \mathbb{I}_N$ is *equivalent* to computing a (local) minimizer of \mathcal{F}_d . In section 3 we prove that the (local) minimizers \hat{u} of \mathcal{F}_d are *strict* if and only if the submatrix $A_{\sigma(\hat{u})}$ has full column rank. The strict minimizers of \mathcal{F}_d are shown to be linear in data d . The importance of the $(M-1)$ -sparse strict minimizers is emphasized. The global minimizers of \mathcal{F}_d are studied in section 4. Their existence is proved. They are shown to be strict for any d and for any $\beta > 0$. More details are provided under the standard assumption that $\text{rank}(A) = M < N$. Given $d \in \mathbb{R}^M$, critical values β_K for $K \in \mathbb{I}_{M-1}$ are exhibited such that all global minimizers of \mathcal{F}_d are K -sparse⁷ if $\beta > \beta_K$.

In section 5, a gentle assumption on A is shown to be *generic* for all $M \times N$ real matrices. Under this assumption, for all data $d \in \mathbb{R}^M$ beyond a closed negligible subset, the objective \mathcal{F}_d for $\beta > \beta_K$, $K \in \mathbb{I}_{M-1}$, has a unique global minimizer and the latter is K -sparse.

Small size (A is 5×10) numerical tests in section 6 illustrate the main theoretical results.

2 All minimizers of \mathcal{F}_d

2.1 Preliminary results

First, we give some basic facts on problem (\mathcal{P}_ω) as defined in (15) needed for later use. If $\omega = \emptyset$, then $\omega^c = \mathbb{I}_N$, so the unique solution of (\mathcal{P}_ω) is $\hat{u} = 0$. For an arbitrary $\omega \subset \mathbb{I}_N$ meeting $\#\omega \geq 1$, (\mathcal{P}_ω) amounts to minimize a quadratic term with respect to only $\#\omega$ components of u , the remaining entries being null. The latter *quadratic* problem (\mathcal{Q}_ω) reads

$$(\mathcal{Q}_\omega) \quad \boxed{\min_{v \in \mathbb{R}^{\#\omega}} \|A_\omega v - d\|^2, \quad \#\omega \geq 1} \quad (16)$$

and it always admits a solution. Using the zero-padding operator Z_ω in (12), we have

$$\left[\hat{u}_\omega \in \mathbb{R}^{\#\omega} \text{ solves } (\mathcal{Q}_\omega) \text{ and } \hat{u} = Z_\omega(\hat{u}_\omega) \right] \Leftrightarrow \left[\hat{u} \in \mathbb{R}^N \text{ solves } (\mathcal{P}_\omega), \quad \#\omega \geq 1 \right].$$

The optimality conditions for (\mathcal{Q}_ω) , combined with the definition in (14)(a), give rise to the following equivalence which holds true for any $\omega \subseteq \mathbb{I}_N$:

$$\left[\hat{u} \in \mathbb{R}^N \text{ solves } (\mathcal{P}_\omega) \right] \Leftrightarrow \left[\hat{u}_\omega \in \mathbb{R}^{\#\omega} \text{ solves } A_\omega^T A_\omega v = A_\omega^T d \text{ and } \hat{u} = Z_\omega(\hat{u}_\omega) \right]. \quad (17)$$

Clearly, problem (\mathcal{P}_ω) in (15) always admits a solution. Note that $A_\omega^T A_\omega v = A_\omega^T d$ in (17) is the normal equation associated to $A_\omega v = d$. The remark below shows that the optimal value of (\mathcal{P}_ω) in (15) can also be seen as an orthogonal projection problem.

Remark 1 Let $r \stackrel{\text{def}}{=} \text{rank}(A_\omega)$ and $B_\omega \in \mathbb{R}^{M \times r}$ be an orthonormal basis for $\text{range}(A_\omega)$. Then $A_\omega = B_\omega H_\omega$ for a uniquely defined matrix $H_\omega \in \mathbb{R}^{r \times \#\omega}$ with $\text{rank}(H_\omega) = r$. Using (17), we have

$$A_\omega^T A_\omega \hat{u}_\omega = A_\omega^T d \Leftrightarrow H_\omega^T H_\omega \hat{u}_\omega = H_\omega^T B_\omega^T d \Leftrightarrow H_\omega \hat{u}_\omega = B_\omega^T d \Leftrightarrow A_\omega \hat{u}_\omega = B_\omega B_\omega^T d.$$

Let $\Pi_{\text{range}(A_\omega)} = B_\omega B_\omega^T$ be the orthogonal projector onto the subspace spanned by the columns of A_ω , see e.g. [28]. The expression above combined with (17) shows that

$$\left[\hat{u} \in \mathbb{R}^N \text{ solves } (\mathcal{P}_\omega) \right] \Leftrightarrow \left[\varpi \subseteq \omega, \text{ range}(A_\varpi) = \text{range}(A_\omega), A_\varpi \hat{u}_\varpi = \Pi_{\text{range}(A_\omega)} d, \hat{u} = Z_\varpi(\hat{u}_\varpi) \right].$$

⁷As usual, a vector u is said to be K -sparse if $\|u\|_0 \leq K$.

In words⁸, $A\hat{u} = A_\omega\hat{u}_\omega$ is the orthogonal projection of d onto the basis B_ω .

For $\omega \subseteq \mathbb{I}_N$, let K_ω denote the vector subspace

$$K_\omega \stackrel{\text{def}}{=} \{v \in \mathbb{R}^N : v[i] = 0, \forall i \in \omega^c\}. \quad (18)$$

This notation enables problem (\mathcal{P}_ω) in (15) to be rewritten as

$$\min_{u \in K_\omega} \|Au - d\|^2. \quad (19)$$

The technical lemma below will be used in the sequel. We emphasize that its statement is independent of the vector $\hat{u} \in \mathbb{R}^N \setminus \{0\}$.

Lemma 1 *Let $d \in \mathbb{R}^M$, $\beta > 0$ and $\hat{u} \in \mathbb{R}^N \setminus \{0\}$ be arbitrary. For $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u})$, set*

$$\rho \stackrel{\text{def}}{=} \min \left\{ \min_{i \in \hat{\sigma}} |\hat{u}[i]|, \frac{\beta}{2(\|A^T(A\hat{u} - d)\|_1 + 1)} \right\}. \quad (20)$$

Then $\rho > 0$.

(i) For ϕ as defined in (2), we have

$$v \in B_\infty(0, \rho) \Rightarrow \sum_{i \in \mathbb{I}_N} \phi(\hat{u}[i] + v[i]) = \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i]) + \sum_{i \in \hat{\sigma}^c} \phi(v[i]);$$

(ii) For $K_{\hat{\sigma}}$ defined according to (18), \mathcal{F}_d satisfies

$$v \in B_\infty(0, \rho) \cap (\mathbb{R}^N \setminus K_{\hat{\sigma}}) \Rightarrow \mathcal{F}_d(\hat{u} + v) \geq \mathcal{F}_d(\hat{u}),$$

where the inequality is strict whenever $\hat{\sigma}^c \neq \emptyset$.

The proof is outlined in Appendix 8.1, p. 29.

2.2 The (local) minimizers of \mathcal{F}_d solve quadratic problems

It is worth emphasizing that no special assumptions on the matrix A are adopted.

We begin with an easy but warning result.

Lemma 2 *For any $d \in \mathbb{R}^M$ and $\forall \beta > 0$, \mathcal{F}_d has a strict (local) minimum at $\hat{u} = 0 \in \mathbb{R}^N$.*

Proof. Using that $\mathcal{F}_d(0) = \|d\|^2 \geq 0$, we have

$$\mathcal{F}_d(v) = \|Av - d\|^2 + \beta\|v\|_0 = \mathcal{F}_d(0) + \mathcal{R}_d(v), \quad (21)$$

$$\text{where } \mathcal{R}_d(v) = \|Av\|^2 - 2\langle v, A^T d \rangle + \beta\|v\|_0. \quad (22)$$

Noticing that $\beta\|v\|_0 \geq \beta > 0$ for $v \neq 0$ leads to

$$v \in B_2\left(0, \frac{\beta}{2\|A^T d\| + 1}\right) \setminus \{0\} \Rightarrow \mathcal{R}_d(v) \geq -2\|v\| \|A^T d\| + \beta > 0.$$

Inserting the latter implication into (21) entails the statement. \square

⁸We note that $B_\omega B_\omega^T$ is the orthogonal projector onto $\text{range}(B_\omega)$.

For any $\beta > 0$ and $d \in \mathbb{R}^M$, the sparsest strict local minimizer of \mathcal{F}_d reads $\hat{u} = 0$.

Initialization with zero of a suboptimal algorithm should generally be a bad choice.

Indeed, experiments have shown that such an initialization can be harmful, see e. g. [29, 7].

The next proposition states a (might be intuitive) result that is often evoked in this work.

Proposition 1 *Let $d \in \mathbb{R}^M$. Given an $\omega \subseteq \mathbb{I}_N$, let \hat{u} solve problem (\mathcal{P}_ω) as formulated in (15). Then for any $\beta > 0$, the objective \mathcal{F}_d in (1) reaches a (local) minimum at \hat{u} and*

$$\sigma(\hat{u}) \subseteq \omega, \quad (23)$$

where $\sigma(\hat{u})$ reads according to Definition 1.

Proof. Let \hat{u} solve problem (\mathcal{P}_ω) and $\beta > 0$. The constraint in (\mathcal{P}_ω) entails (23).

Consider that $\hat{u} \neq 0$ in which case for $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u})$ we have $1 \leq \#\hat{\sigma} \leq \#\omega$. Using the equivalent formulation of (\mathcal{P}_ω) given in (18)-(19), p. 7, yields

$$v \in K_\omega \quad \Rightarrow \quad \|A(\hat{u} + v) - d\|^2 \geq \|A\hat{u} - d\|^2. \quad (24)$$

The inclusion in (23) is equivalent to $\omega^c \subseteq \hat{\sigma}^c$. Let $K_{\hat{\sigma}}$ read according to (18) as well. Then

$$\hat{u} \in K_{\hat{\sigma}} \subseteq K_\omega.$$

Combining the latter relation with (24) entails that

$$v \in K_{\hat{\sigma}} \quad \Rightarrow \quad \|A(\hat{u} + v) - d\|^2 \geq \|A\hat{u} - d\|^2. \quad (25)$$

Let ρ read as in (20), Lemma 1. Noticing that by (2) and (18)

$$v \in K_{\hat{\sigma}} \quad \Rightarrow \quad \phi(v[i]) = 0, \quad \forall i \in \hat{\sigma}^c, \quad (26)$$

the following inequality chain is derived:

$$\begin{aligned} v \in B_\infty(0, \rho) \cap K_{\hat{\sigma}} \quad \Rightarrow \quad \mathcal{F}_d(\hat{u} + v) &= \|A(\hat{u} + v) - d\|^2 + \beta \sum_{i \in \mathbb{I}_N} \phi(\hat{u}[i] + v[i]) \\ &\stackrel{\text{[by Lemma 1(i)]}}{=} \|A(\hat{u} + v) - d\|^2 + \beta \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i]) + \beta \sum_{i \in \hat{\sigma}^c} \phi(v[i]) \\ &\stackrel{\text{[by (26)]}}{=} \|A(\hat{u} + v) - d\|^2 + \beta \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i]) \\ &\stackrel{\text{[by (25)]}}{\geq} \|A\hat{u} - d\|^2 + \beta \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i]) \\ &\stackrel{\text{[by (4), p. 2]}}{=} \mathcal{F}_d(\hat{u}). \end{aligned}$$

Combining the latter implication with Lemma 1(ii), p. 7, shows that

$$\mathcal{F}_d(\hat{u} + v) \geq \mathcal{F}_d(\hat{u}), \quad \forall v \in B_\infty(0, \rho).$$

If $\hat{u} = 0$, this is a (local) minimizer of \mathcal{F}_d by Lemma 2. □

Many authors mention that initialization is paramount for the success of approximate algorithms minimizing \mathcal{F}_d . In view of Proposition 1, if one already has a well elaborated initialization, it could be enough to solve the relevant problem (\mathcal{P}_ω) .

The statement reciprocal to Proposition 1 is obvious but it helps the presentation.

Lemma 3 For $d \in \mathbb{R}^M$ and $\beta > 0$, let \mathcal{F}_d have a (local) minimum at \hat{u} . Then \hat{u} solves $(\mathcal{P}_{\hat{\sigma}})$ for $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u})$.

Proof. Let \hat{u} be a (local) minimizer of \mathcal{F}_d . Denote $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u})$. Then \hat{u} solves the problem:

$$\min_{u \in \mathbb{R}^N} \left\{ \|Au - d\|^2 + \beta \# \hat{\sigma} \right\} \quad \text{subject to } u[i] = 0, \forall i \in \hat{\sigma}^c.$$

Since $\# \hat{\sigma}$ is a constant, \hat{u} solves $(\mathcal{P}_{\hat{\sigma}})$. □

By Proposition 1 and Lemma 3, to solve (\mathcal{P}_ω) for some $\omega \subseteq \mathbb{I}_N$ and then to apply the zero padding operator Z_ω in (12) is equivalent to find a (local) minimizer of \mathcal{F}_d .

This equivalence underlies most of the theory developed in this work.

Corollary 1 For $d \in \mathbb{R}^M$ and $\beta > 0$, let \hat{u} be a (local) minimizer of \mathcal{F}_d . Set $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u})$. Then

$$\hat{u} = Z_{\hat{\sigma}}(\hat{u}_{\hat{\sigma}}) \quad \text{where } \hat{u}_{\hat{\sigma}} \text{ satisfies } A_{\hat{\sigma}}^T A_{\hat{\sigma}} \hat{u}_{\hat{\sigma}} = A_{\hat{\sigma}}^T d. \quad (27)$$

Conversely, if $\hat{u} \in \mathbb{R}^N$ meets (27) for $\hat{\sigma} = \sigma(\hat{u})$, then \hat{u} is a (local) minimizer of \mathcal{F}_d .

Proof. By Lemma 3, \hat{u} solves $(\mathcal{P}_{\hat{\sigma}})$. The equation in (27) follows directly from (17) on p. 6. The last claim is a straightforward consequence of (17) and Proposition 1. □

Equation (27) shows that a (local) minimizer \hat{u} of \mathcal{F}_d follows a *pseudo-hard* thresholding scheme⁹: the nonzero part $\hat{u}_{\hat{\sigma}}$ of \hat{u} is the least squares solution with respect to the submatrix $A_{\hat{\sigma}}$ and the whole data vector d is involved in its computation. Unlike the hard thresholding scheme in [14], insignificant or purely noisy data entries can hardly be discarded from \hat{u} and they risk to pollute its nonzero part $\hat{u}_{\hat{\sigma}}$. Noisy data d should degrade $\hat{u}_{\hat{\sigma}}$ and this effect is stronger if $A_{\hat{\sigma}}^T A_{\hat{\sigma}}$ is ill-conditioned [12]. The quality of the outcome critically depends on the selected (local) minimizer and on the pertinence of A . See also Remark 2 on p. 12.

It may be interesting to evoke another consequence of Proposition 1:

Given $d \in \mathbb{R}^M$, for any $\omega \subseteq \mathbb{I}_N$, \mathcal{F}_d has a (local) minimizer \hat{u} such that $\sigma(\hat{u}) \subseteq \omega$.

3 The strict minimizers of \mathcal{F}_d

We remind, yet again, that no special assumptions on $A \in \mathbb{R}^{M \times N}$ are adopted.

Strict minimizers of an objective function enable unambiguous solutions of inverse problems. We focus on the strict minimizers of \mathcal{F}_d . The definition below is practical to characterize them.

Definition 3 Given a matrix $A \in \mathbb{R}^{M \times N}$, for any $r \in \mathbb{I}_M$ we define Ω_r as the subset of all r -length supports that correspond to full column rank $M \times r$ submatrices of A , i. e.

$$\Omega_r = \left\{ \omega \subset \mathbb{I}_N : \# \omega = r = \text{rank}(A_\omega) \right\}.$$

⁹In a Bayesian setting, the quadratic data fidelity term in \mathcal{F}_d models data corrupted with Gaussian i.i.d. noise.

Set $\Omega_0 = \emptyset$ and define as well

$$\boxed{\Omega \stackrel{\text{def}}{=} \bigcup_{r=0}^{M-1} \Omega_r \quad \text{and} \quad \Omega_{\max} \stackrel{\text{def}}{=} \Omega \cup \Omega_M .}$$

Definition 3 entails that for any $r \in \mathbb{I}_M$,

$$\text{rank}(A) = r \geq 1 \quad \Leftrightarrow \quad \Omega_r \neq \emptyset \quad \text{and} \quad \Omega_t = \emptyset, \quad \forall t \geq r + 1 .$$

3.1 How to recognize a strict minimizer of \mathcal{F}_d ?

The theorem below gives an exhaustive answer to this question.

Theorem 1 *Given $d \in \mathbb{R}^M$ and $\beta > 0$, let \hat{u} be a (local) minimizer of \mathcal{F}_d . Define*

$$\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u}) .$$

The following statements are equivalent:

- (i) The (local) minimum that \mathcal{F}_d has at \hat{u} is strict;
- (ii) $\boxed{\text{rank}(A_{\hat{\sigma}}) = \#\hat{\sigma}}$;
- (iii) $\hat{\sigma} \in \Omega_{\max}$.

If \hat{u} is a strict (local) minimizer of \mathcal{F}_d , then it reads

$$\hat{u} = Z_{\hat{\sigma}}(\hat{u}_{\hat{\sigma}}) \quad \text{for} \quad \hat{u}_{\hat{\sigma}} = (A_{\hat{\sigma}}^T A_{\hat{\sigma}})^{-1} A_{\hat{\sigma}}^T d \quad (28)$$

and satisfies $\#\hat{\sigma} = \|\hat{u}\|_0 \leq M$.

Proof. We recall that by the rank-nullity theorem [20, 28]

$$\dim \ker(A_{\hat{\sigma}}) = \#\hat{\sigma} - \text{rank}(A_{\hat{\sigma}}) . \quad (29)$$

Proof of (i) \Rightarrow (ii). Let¹⁰ $\hat{u} \neq 0$ be a strict (local) minimizer of \mathcal{F}_d . Assume that (ii) fails. Using (29) the latter means that

$$\dim \ker(A_{\hat{\sigma}}) \geq 1 . \quad (30)$$

By Lemma 3, \hat{u} solves $(\mathcal{P}_{\hat{\sigma}})$. Let ρ read as in (20) and $K_{\hat{\sigma}}$ be defined according to (18), p. 7. Noticing that

$$v \in K_{\hat{\sigma}}, \quad \hat{\sigma} \neq \emptyset \quad \Rightarrow \quad Av = A_{\hat{\sigma}} v_{\hat{\sigma}} , \quad (31)$$

Lemma 1(i) shows that

$$\begin{aligned} \left\{ \begin{array}{l} v \in B_{\infty}(0, \rho) \cap K_{\hat{\sigma}} , \\ v_{\hat{\sigma}} \in \ker(A_{\hat{\sigma}}) \end{array} \right. & \Rightarrow \quad \mathcal{F}_d(\hat{u} + v) = \|A_{\hat{\sigma}}(\hat{u}_{\hat{\sigma}} + v_{\hat{\sigma}}) - d\|^2 + \beta \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i] + v[i]) \\ & \quad \left[\text{by Lemma 1(i)} \right] = \|A_{\hat{\sigma}}\hat{u}_{\hat{\sigma}} - d\|^2 + \beta \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i]) + \beta \sum_{i \in \hat{\sigma}^c} \phi(v[i]) \\ & \quad \left[\text{by (26), p. 8} \right] = \|A_{\hat{\sigma}}\hat{u}_{\hat{\sigma}} - d\|^2 + \beta \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i]) \\ & \quad \left[\text{by (4), p. 2} \right] = \mathcal{F}_d(\hat{u}) , \end{aligned}$$

¹⁰This part can alternatively be proven using Remark 1, p. 6.

i. e. that \hat{u} is not a strict minimizer, which contradicts (i). Hence the assumption in (30) is false. Using (29), (ii) holds true.

If $\hat{u} = 0$, then $\hat{\sigma} = \emptyset$, hence $A_{\hat{\sigma}} \in \mathbb{R}^{M \times 0}$ and $\text{rank}(A_{\hat{\sigma}}) = 0 = \#\hat{\sigma}$ after (14).

Proof of (ii) \Rightarrow (i). Let \hat{u} be a minimizer of \mathcal{F}_d that meets (ii). To have $\#\hat{\sigma} = 0$ is equivalent to $\hat{u} = 0$. By Lemma 2, \hat{u} is a strict minimizer. Focus on $\#\hat{\sigma} \geq 1$. Since $\text{rank}(A_{\hat{\sigma}}) = \#\hat{\sigma} \leq M$ and problem (\mathcal{Q}_ω) in (16), p. 6, is strictly convex for $\omega = \hat{\sigma}$, its unique solution $\hat{u}_{\hat{\sigma}}$ satisfies

$$v \in \mathbb{R}^{\#\hat{\sigma}} \setminus \{0\} \quad \Rightarrow \quad \|A_{\hat{\sigma}}(\hat{u}_{\hat{\sigma}} + v) - d\|^2 > \|A_{\hat{\sigma}}\hat{u}_{\hat{\sigma}} - d\|^2 .$$

Using (31), p. 10, the latter is equivalent to

$$v \in K_{\hat{\sigma}} \setminus \{0\} \quad \Rightarrow \quad \|A(\hat{u} + v) - d\|^2 = \|A_{\hat{\sigma}}(\hat{u}_{\hat{\sigma}} + v_{\hat{\sigma}}) - d\|^2 > \|A_{\hat{\sigma}}\hat{u}_{\hat{\sigma}} - d\|^2 = \|A\hat{u} - d\|^2 . \quad (32)$$

Lemma 1(i), along with (26), p. 8, yields

$$\begin{aligned} v \in B_\infty(0, \rho) \cap K_{\hat{\sigma}} \setminus \{0\} \quad \Rightarrow \quad \mathcal{F}_d(\hat{u} + v) &= \|A(u + v) - d\|^2 + \beta \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i]) \\ &\quad \left[\text{by (32)} \right] > \|A\hat{u} - d\|^2 + \beta \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i]) \\ &= \mathcal{F}_d(\hat{u}) . \end{aligned}$$

Since $\#\hat{\sigma} \leq M \leq N - 1$, we have $\hat{\sigma}^c \neq \emptyset$. So Lemma 1(ii) tells us that

$$v \in B_\infty(0, \rho) \setminus K_{\hat{\sigma}} \quad \Rightarrow \quad \mathcal{F}_d(\hat{u} + v) > \mathcal{F}_d(\hat{u}) .$$

Combining the last two implications proves (i).

Proof of (ii) \Leftrightarrow (iii). Comparing (iii) with Definitions 1 (p. 4) and 3 shows the statement.

Equation (28). It follows from (27) in Corollary 1 (p. 9) where¹¹ $A_{\hat{\sigma}}^T A_{\hat{\sigma}}$ is invertible. □

Theorem 1 provides a simple rule enabling to verify if a numerical scheme has reached or not a strict (local) minimizer of \mathcal{F}_d .

The notations Ω_r , Ω and Ω_{\max} are frequently used in this paper. Their interpretation is obvious in the light of Theorem 1.

For any $d \in \mathbb{R}^M$ and $\forall \beta > 0$, the set Ω_{\max} is composed of the supports of all possible strict (local) minimizers of \mathcal{F}_d , while Ω is the subset of those ones that are $(M - 1)$ -sparse.

An easy and useful corollary is presented next.

Corollary 2 *Let $d \in \mathbb{R}^M$. Given an arbitrary $\omega \in \Omega_{\max}$, let \hat{u} solve (\mathcal{P}_ω) . Then*

¹¹For $\hat{\sigma} = \emptyset$, (12) and (14)(a) show that (28) yields $\hat{u} = 0$.

(i) \hat{u} reads

$$\hat{u} = Z_\omega(\hat{u}_\omega) \quad \text{where} \quad \hat{u}_\omega = (A_\omega^T A_\omega)^{-1} A_\omega^T d, \quad (33)$$

and meets $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u}) \subseteq \omega$ and $\hat{\sigma} \in \Omega_{\max}$;

(ii) for any $\beta > 0$, \hat{u} is a strict (local) minimizer of \mathcal{F}_d ;

(iii) \hat{u} solves $(\mathcal{P}_{\hat{\sigma}})$.

Proof. Using (17) (see p. 6), \hat{u} meets (i) since $A_\omega^T A_\omega$ is invertible and $\sigma(\hat{u}) \subseteq \omega$ by the constraint in (\mathcal{P}_ω) . If $\hat{\sigma} = \emptyset$, (ii) follows from Lemma 2. For $\#\hat{\sigma} \geq 1$, $A_{\hat{\sigma}}$ is an $M \times \#\hat{\sigma}$ submatrix of A_ω . Since $\text{rank}(A_\omega) = \#\omega$, we have $\text{rank}(A_{\hat{\sigma}}) = \#\hat{\sigma}$ and so $\hat{\sigma} \in \Omega_{\max}$. By Proposition 1 \hat{u} is a (local) minimizer of \mathcal{F}_d and Theorem 1 leads to (ii). Lemma 3 and Corollary 2(ii) yield (iii). \square

The following consequence of Corollary 2 might be striking:

One can easily compute a strict (local) minimizer \hat{u} of \mathcal{F}_d without knowing the value of the regularization parameter β . Just data d and an $\omega \in \Omega_{\max}$ are needed.

Clearly, the support $\sigma(\hat{u})$ of a nonstrict local minimizer \bar{u} of \mathcal{F}_d contains some sub-supports yielding strict (local) minimizers of \mathcal{F}_d . It is easy to see that among them, there is $\hat{\sigma} \subsetneq \sigma(\bar{u})$ such that the corresponding \hat{u} given by (28) *strictly* decreases the value of \mathcal{F}_d , i.e. $\mathcal{F}_d(\hat{u}) < \mathcal{F}_d(\bar{u})$.

3.2 Every strict (local) minimizer of \mathcal{F}_d is linear in d

Here we explore the behavior of the strict (local) minimizers of \mathcal{F}_d with respect to variations of d . An interesting sequel of Theorem 1 is presented below.

Corollary 3 For $d \in \mathbb{R}^M$ and $\beta > 0$, let \hat{u} be a (local) minimizer of \mathcal{F}_d satisfying $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u}) \in \Omega$. Define

$$N_{\hat{\sigma}} \stackrel{\text{def}}{=} \ker(A_{\hat{\sigma}}^T) \subset \mathbb{R}^M .$$

We have $\dim N_{\hat{\sigma}} = M - \#\hat{\sigma} \geq 1$ and

$$d' \in N_{\hat{\sigma}} \quad \Rightarrow \quad \mathcal{F}_{d+d'} \quad \text{has a strict (local) minimum at} \quad \hat{u} .$$

Proof. Since $\hat{\sigma} \in \Omega$, the minimizer \hat{u} is strict by Theorem 1. By $d' \in \ker(A_{\hat{\sigma}}^T)$ we find $A_{\hat{\sigma}}^T(d + d') = A_{\hat{\sigma}}^T d$ for any $d' \in N_{\hat{\sigma}}$. Inserting this into (28) in Theorem 1 yields the result. \square

All data living in the vector subspace $N_{\hat{\sigma}} \supsetneq \{0\}$ yield the same strict (local) minimizer \hat{u} .

Remark 2 If data contain some noise n , it can be decomposed in a unique way as $n = n_{N_{\hat{\sigma}}} + n_{N_{\hat{\sigma}}^\perp}$ where $n_{N_{\hat{\sigma}}} \in N_{\hat{\sigma}}$ and $n_{N_{\hat{\sigma}}^\perp} \in N_{\hat{\sigma}}^\perp$. By Corollary 3, the component $n_{N_{\hat{\sigma}}}$ will be removed.

We shall use the definition given below.

Definition 4 Let $\mathcal{O} \subseteq \mathbb{R}^M$ be an open domain. We say that $\mathcal{U} : \mathcal{O} \rightarrow \mathbb{R}^N$ is a local minimizer function for the family of objectives $\mathfrak{F}_{\mathcal{O}} \stackrel{\text{def}}{=} \{\mathcal{F}_d : d \in \mathcal{O}\}$ if for any $d \in \mathcal{O}$, the function \mathcal{F}_d reaches a strict (local) minimum at $\mathcal{U}(d)$.

Corollary 2 shows that for any $d \in \mathbb{R}^M$, each strict (local) minimizer of \mathcal{F}_d is entirely described by an $\omega \in \Omega_{\max}$ via equation (33) in the same corollary.

Consequently, a local minimizer function \mathcal{U} is associated with every $\omega \in \Omega_{\max}$.

Lemma 4 For some arbitrarily fixed $\omega \in \Omega_{\max}$ and $\beta > 0$, the family of functions $\mathfrak{F}_{\mathbb{R}^M}$ has a linear (local) minimizer function $\mathcal{U} : \mathbb{R}^M \rightarrow \mathbb{R}^N$ that reads

$$\forall d \in \mathbb{R}^M, \quad \mathcal{U}(d) = Z_\omega(U_\omega d) \quad \text{where} \quad U_\omega = (A_\omega^T A_\omega)^{-1} A_\omega^T \in \mathbb{R}^{\#\omega \times M}. \quad (34)$$

Proof. Clearly, the function \mathcal{U} in (34) is linear in d . From Corollary 2 (p. 11), for any $\beta > 0$ and for any $d \in \mathbb{R}^M$, \mathcal{F}_d has a strict (local) minimum at $\hat{u} = \mathcal{U}(d)$. Hence \mathcal{U} fits Definition 4. \square

Even if \mathcal{F}_d has many strict (local) minimizers, each one of them is linear in d .

Next we exhibit a closed negligible subset of \mathbb{R}^M , associated with a nonempty $\omega \in \Omega_{\max}$, whose elements are data d leading to $\|\mathcal{U}(d)\|_0 < \#\omega$.

Lemma 5 For any $\omega \in \Omega_{\max}$, define the subset $D_\omega \subset \mathbb{R}^M$ by

$$D_\omega \stackrel{\text{def}}{=} \bigcup_{i=1}^{\#\omega} \left\{ g \in \mathbb{R}^M : \left\langle e_i, (A_\omega^T A_\omega)^{-1} A_\omega^T g \right\rangle = 0 \right\}. \quad (35)$$

Then D_ω is closed in \mathbb{R}^M and $\mathbb{L}^M(D_\omega) = 0$.

Proof. If $\omega = \emptyset$ then $D_\omega = \emptyset$, hence the claim. Let $\#\omega \geq 1$. For some $i \in \mathbb{I}_{\#\omega}$, set

$$\begin{aligned} D &\stackrel{\text{def}}{=} \left\{ g \in \mathbb{R}^M : \left\langle e_i, (A_\omega^T A_\omega)^{-1} A_\omega^T g \right\rangle = 0 \right\} \\ &= \left\{ g \in \mathbb{R}^M : \left\langle A_\omega (A_\omega^T A_\omega)^{-1} e_i, g \right\rangle = 0 \right\}. \end{aligned}$$

Since $\text{rank} \left(A_\omega (A_\omega^T A_\omega)^{-1} \right) = \#\omega$, then $\ker \left(A_\omega (A_\omega^T A_\omega)^{-1} \right) = \{0\}$. Therefore $A_\omega (A_\omega^T A_\omega)^{-1} e_i \neq 0$. Hence D is a vector subspace of \mathbb{R}^M of dimension $M - 1$ and so $\mathbb{L}^M(D) = 0$. The conclusion follows from the fact that D_ω in (35) is the union of $\#\omega$ subsets like D (see e. g. [35, 15]). \square

Proposition 2 For some arbitrarily fixed $\omega \in \Omega_{\max}$ and $\beta > 0$, let $\mathcal{U} : \mathbb{R}^M \rightarrow \mathbb{R}^N$ be the relevant (local) minimizer function for $\mathfrak{F}_{\mathbb{R}^M}$ as given in (34) (Lemma 4). Let D_ω read as in (35). Then the function $d \mapsto \mathcal{F}_d(\mathcal{U}(d))$ is \mathcal{C}^∞ on $\mathbb{R}^M \setminus D_\omega$ and

$$d \in \mathbb{R}^M \setminus D_\omega \quad \Rightarrow \quad \sigma(\mathcal{U}(d)) = \omega,$$

where the set $\mathbb{R}^M \setminus D_\omega$ contains an open and dense subset of \mathbb{R}^M .

Proof. The statement about $\mathbb{R}^M \setminus D_\omega$ is a direct consequence of Lemma 5.

If $\omega = \emptyset$, then $\mathcal{U}(d) = 0$, $\forall d \in \mathbb{R}^M$, so the implication is trivial. Consider that $\#\omega \geq 1$. For any $i \in \mathbb{I}_{\#\omega}$, the $\omega[i]$ th component of $\mathcal{U}(d)$ reads (see Lemma 4)

$$\mathcal{U}_{\omega[i]}(d) = \left\langle e_i, (A_\omega^T A_\omega)^{-1} A_\omega^T d \right\rangle.$$

The definition of D_ω shows that

$$d \in \mathbb{R}^M \setminus D_\omega \quad \text{and} \quad i \in \mathbb{I}_{\# \omega} \quad \Rightarrow \quad \mathcal{U}_{\omega[i]}(d) \neq 0 ,$$

whereas $\mathcal{U}_i(d) = 0$ for all $i \in \omega^c$. Consequently,

$$\omega \in \Omega_{\max} \quad \text{and} \quad d \in \mathbb{R}^M \setminus D_\omega \quad \Rightarrow \quad \sigma(\mathcal{U}(d)) = \omega .$$

Then $\|\mathcal{U}(d)\|_0$ is constant on $\mathbb{R}^M \setminus D_\omega$ and

$$\omega \in \Omega_{\max} \quad \text{and} \quad d \in \mathbb{R}^M \setminus D_\omega \quad \Rightarrow \quad \mathcal{F}_d(\mathcal{U}(d)) = \|A\mathcal{U}(d) - d\|^2 + \beta \# \omega .$$

We infer from (34) that $d \mapsto \|A\mathcal{U}(d) - d\|^2$ is \mathcal{C}^∞ on \mathbb{R}^M , so $d \mapsto \mathcal{F}_d(\mathcal{U}(d))$ is \mathcal{C}^∞ on $\mathbb{R}^M \setminus D_\omega$. \square

A generic property is that a local minimizer function produces solutions sharing the same support.

The application $d \mapsto \mathcal{F}_d(\mathcal{U}(d))$ is discontinuous on the *closed negligible* subset D_ω where the support of $\mathcal{U}(d)$ is shrunk.

3.3 Strict minimizers with an M-length support

Proposition 3 *Let $\text{rank}(A) = M$, $\beta > 0$ and for $d \in \mathbb{R}^M$ set*

$$U_M \stackrel{\text{def}}{=} \{ \hat{u} \in \mathbb{R}^N : \hat{u} \text{ is a strict (local) minimizer of } \mathcal{F}_d \text{ meeting } \|\hat{u}\|_0 = M \} .$$

Put

$$Q_M \stackrel{\text{def}}{=} \bigcup_{\omega \in \Omega_M} \bigcup_{i \in \mathbb{I}_M} \{ g \in \mathbb{R}^M : \langle e_i, A_\omega^{-1} g \rangle = 0 \} . \quad (36)$$

Then $\mathbb{R}^M \setminus Q_M$ contains a dense open subset of \mathbb{R}^M and

$$d \in \mathbb{R}^M \setminus Q_M \quad \Rightarrow \quad \# U_M = \# \Omega_M \quad \text{and} \quad \mathcal{F}_d(\hat{u}) = \beta M, \quad \forall \hat{u} \in U_M .$$

Proof. Using the notation in (35), Q_M reads

$$Q_M = \bigcup_{\omega \in \Omega_M} D_\omega .$$

The claim on $\mathbb{R}^M \setminus Q_M$ follows from Lemma 5. Since $\text{rank}(A) = M$, we have $\# \Omega_M \geq 1$.

Consider that $d \in \mathbb{R}^M \setminus Q_M$. By Proposition 2

$$d \in \mathbb{R}^M \setminus Q_M \quad \text{and} \quad \omega \in \Omega_M \quad \Rightarrow \quad \mathcal{F}_d \text{ has a strict (local) minimizer } \hat{u} \text{ obeying } \sigma(\hat{u}) = \omega .$$

Hence $\hat{u} \in U_M$. Therefore, we have a mapping $b : \Omega_M \rightarrow U_M$ such that $\hat{u} = b(\omega) \in U_M$. Using Lemma 3 and Corollary 2, it reads

$$b(\omega) = Z_\omega(A_\omega^{-1} d) .$$

For $(\omega, \varpi) \in \Omega_M \times \Omega_M$ with $\varpi \neq \omega$ one obtains $\hat{u} = b(\omega) \in U_M$, $\bar{u} = b(\varpi) \in U_M$ and $\hat{u} \neq \bar{u}$, hence b is one-to-one. Conversely, for any $\hat{u} \in U_M$ there is $\omega \in \Omega_M$ such that $\hat{u} = b(\omega)$ and $\sigma(\hat{u}) = \omega$ (because $d \notin Q_M$). It follows that b maps Ω_M onto U_M . Therefore, Ω_M and U_M are in one-to-one correspondence, i.e. $\# \Omega_M = \# U_M$.

Last, it is clear that $\omega \in \Omega_M$ and $d \notin Q_M$ lead to $\|A\hat{u} - d\|^2 = 0$ and $\mathcal{F}_d(\hat{u}) = \beta M$. \square

For any $\beta > 0$, a generic property of \mathcal{F}_d is that it has $\sharp\Omega_M$ strict minimizers \hat{u} obeying $\|\hat{u}\|_0 = M$ and $\mathcal{F}_d(\hat{u}) = \beta M$. It is hard to discriminate between all these minimizers.

Hence the interest in minimizers with supports living in Ω , i. e. strict $(M - 1)$ -sparse minimizers of \mathcal{F}_d .

4 On the global minimizers of \mathcal{F}_d

The statement below gives a *necessary condition* for a global minimizer of \mathcal{F}_d . It follows directly from [31, Proposition 3.4] where¹² the regularization term is $\|Du\|_0$ for a full row rank matrix D . For \mathcal{F}_d in (1) with $\|a_i\|_2 = 1, \forall i \in \mathbb{I}_N$, a simpler condition was derived later on in [39, Theorem 12], using different tools. For completeness, the proof for a general A is outlined in Appendix 8.2 on p. 30.

Proposition 4 For $d \in \mathbb{R}^M$ and $\beta > 0$, let \mathcal{F}_d have a global minimum at \hat{u} . Then¹³

$$i \in \sigma(\hat{u}) \quad \Rightarrow \quad |\hat{u}[i]| \geq \frac{\sqrt{\beta}}{\|a_i\|}. \quad (37)$$

Observe that the lower bound on $\{|\hat{u}[i]| : i \in \sigma(\hat{u})\}$ given in (37) is independent of d . This means that in general, (37) provides a pessimistic bound.

The proof of the proposition shows that (37) is met also by all (local) minimizers of \mathcal{F}_d satisfying

$$\mathcal{F}_d(\hat{u}) \leq \mathcal{F}_d(\hat{u} + \rho e_i), \quad \forall \rho \in \mathbb{R}, \quad \forall i \in \mathbb{I}_N.$$

4.1 The global minimizers of \mathcal{F}_d are strict

Remark 3 Let $d \in \mathbb{R}^M$ and $\beta > \|d\|^2$. Then \mathcal{F}_d has a strict global minimum at $\hat{u} = 0$. Indeed,

$$v \neq 0 \quad \Rightarrow \quad \|v\|_0 \geq 1 \quad \Rightarrow \quad \mathcal{F}_d(0) = \|d\|^2 < \beta \leq \|Av - d\|^2 + \beta\|v\|_0.$$

For least-squares regularized with a more regular ϕ , one usually gets $\hat{u} = 0$ asymptotically as $\beta \rightarrow +\infty$ but $\hat{u} \neq 0$ for finite values of β . The latter does not hold for \mathcal{F}_d (Remark 3).

Some theoretical results on the global minimizers of \mathcal{F}_d were obtained by the past [31, 21, 39, 7]. Surprisingly, the question about the *existence of global minimizers of \mathcal{F}_d* has never been raised. We answer this question using the notion of *asymptotically level stable functions* introduced by Auslender [2] in 2000. Since then, it was revealed that many problems related to the existence of optimal solutions and their stability under perturbations are easy to solve for this wide class of functions. As usual,

$$\text{lev}(\mathcal{F}_d, \lambda) \stackrel{\text{def}}{=} \{v \in \mathbb{R}^N : \mathcal{F}_d(v) \leq \lambda\} \quad \text{for } \lambda > \inf \mathcal{F}_d.$$

The definition below is taken from [3, p. 94].

¹²Just set $g_i = e_i, P = I_M$ and $H = I_N$ in [31, Proposition 3.4].

¹³Recall that $a_i \neq 0$ for all $i \in \mathbb{I}_N$ by (9) and that $\|\cdot\| = \|\cdot\|_2$.

Definition 5 Let $\mathcal{F}_d : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ be lower semicontinuous and proper. Then \mathcal{F}_d is said asymptotically level stable if for each $\rho > 0$, each bounded sequence $\{\lambda_k\} \in \mathbb{R}$ and each sequence $\{v_k\} \in \mathbb{R}^N$ satisfying

$$v_k \in \text{lev}(\mathcal{F}_d, \lambda_k), \quad \|v_k\| \rightarrow +\infty, \quad v_k \|v_k\|^{-1} \rightarrow \bar{v} \in \ker((\mathcal{F}_d)_\infty), \quad (38)$$

where $(\mathcal{F}_d)_\infty$ denotes the asymptotic (or recession) function of \mathcal{F}_d , there exists k_0 such that

$$v_k - \rho \bar{v} \in \text{lev}(\mathcal{F}_d, \lambda_k) \quad \forall k \geq k_0. \quad (39)$$

One can note that a coercive function is asymptotically level stable, since (38) is empty. We prove that our discontinuous noncoercive objective \mathcal{F}_d is asymptotically level stable, as well.

Proposition 5 Let $\mathcal{F}_d : \mathbb{R}^N \rightarrow \mathbb{R}$ be of the form (1). Then $\ker((\mathcal{F}_d)_\infty) = \ker(A)$ and \mathcal{F}_d is asymptotically level stable.

The proof is outlined in Appendix 8.3, p. 31.

Theorem 2 Let $d \in \mathbb{R}^M$ and $\beta > 0$. Then

(i) the set of the global minimizers of \mathcal{F}_d

$$\hat{U} \stackrel{\text{def}}{=} \left\{ \hat{u} \in \mathbb{R}^N : \hat{u} = \min_{u \in \mathbb{R}^N} \mathcal{F}_d(u) \right\} \quad (40)$$

is nonempty;

(ii) every $\hat{u} \in \hat{U}$ is a strict minimizer of \mathcal{F}_d , i. e.

$$\sigma(\hat{u}) \in \Omega_{\max},$$

hence $\|\hat{u}\|_0 \leq M$.

Proof. For claim (i), we use the following statement¹⁴, whose proof can be found in the monograph of Auslender and Teboulle [3].

[3, Corollary 3.4.2] Let $\mathcal{F}_d : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$ be asymptotically level stable with $\inf \mathcal{F}_d > -\infty$. Then the optimal set \hat{U} —as given in (40)—is nonempty.

From Proposition 5, \mathcal{F}_d is asymptotically level stable and $\inf \mathcal{F}_d \geq 0$ from (1). Hence $\hat{U} \neq \emptyset$.

(ii). Let \hat{u} be a global minimizer of \mathcal{F}_d . Set $\hat{\sigma} = \sigma(\hat{u})$.

If $\hat{u} = 0$, (ii) follows from Lemma 2 (p. 7). Let $\hat{u} \neq 0$ be a *nonstrict* minimizer of \mathcal{F}_d . Then Theorem 1(ii) (p. 10), fails to hold, hence

$$\dim \ker(A_{\hat{\sigma}}) \geq 1. \quad (41)$$

Choose $v_{\hat{\sigma}} \in \ker(A_{\hat{\sigma}}) \setminus \{0\}$ and set $v = Z_{\hat{\sigma}}(v_{\hat{\sigma}})$. Select an $i \in \hat{\sigma}$ satisfying $v[i] \neq 0$. Define \tilde{u} by

$$\tilde{u} \stackrel{\text{def}}{=} \hat{u} - \hat{u}[i] \frac{v}{v[i]}. \quad (42)$$

¹⁴This result was originally exhibited in [4] (without the notion of asymptotically level stable functions).

We have $\tilde{u}[i] = 0$ and $\hat{u}[i] \neq 0$. Set $\tilde{\sigma} \stackrel{\text{def}}{=} \sigma(\tilde{u})$. Then

$$\tilde{\sigma} \subsetneq \hat{\sigma} \quad \text{hence} \quad \#\tilde{\sigma} \leq \#\hat{\sigma} - 1. \quad (43)$$

From $v_{\hat{\sigma}} \frac{\hat{u}[i]}{v[i]} \in \ker(A_{\hat{\sigma}})$, using (13), p. 5 and Remark 1 (p. 6) show that¹⁵ $A\hat{u} = A_{\hat{\sigma}}\hat{u}_{\hat{\sigma}} = A_{\tilde{\sigma}}\tilde{u}_{\tilde{\sigma}} = A\tilde{u}$. Then

$$\begin{aligned} \mathcal{F}_d(\tilde{u}) &= \|A\tilde{u} - d\|^2 + \beta \#\tilde{\sigma} \\ &\leq \mathcal{F}_d(\hat{u}) - \beta = \|A\hat{u} - d\|^2 + \beta(\#\hat{\sigma} - 1). \end{aligned}$$

It follows that \hat{u} is not a global minimizer, hence (41) is false. Therefore $\text{rank}(A_{\hat{\sigma}}) = \#\hat{\sigma}$ and \hat{u} is a strict minimizer of \mathcal{F}_d (Theorem 1). \square

One can note that if $\text{rank}(A) = M$, any global minimizer \hat{u} of \mathcal{F}_d obeys

$$\mathcal{F}_d(\hat{u}) \leq \beta M.$$

According to Theorem 2, the global minimizers of \mathcal{F}_d are strict and their number is finite: this is a nice property that fails for many convex nonsmooth optimization problems.

4.2 Can we get K-sparse global minimizers for $K \leq M - 1$?

In order to simplify the presentation, in what follows we consider that

$$\text{rank}(A) = M < N.$$

Since \mathcal{F}_d has a large number (typically equal to $\#\Omega_M$) of strict minimizers with $\|\hat{u}\|_0 = M$ yielding the same value $\mathcal{F}_d(\hat{u}) = \beta M$ (see Proposition 3 on p. 14 and the comments next to it), it is important to get sure that the global minimizers of \mathcal{F}_d are $(M - 1)$ -sparse.

We introduce a notation which is used in the rest of this article. For any $K \in \mathbb{I}_{M-1}$, put

$$\overline{\Omega}_K \stackrel{\text{def}}{=} \bigcup_{r=0}^K \Omega_r, \quad (44)$$

where Ω_r read according to Definition 3 (p. 9). Theorem 1 gives a clear meaning of the sets¹⁶ $\overline{\Omega}_K$.

For any $d \in \mathbb{R}^M$ and any $\beta > 0$, for any fixed $K \in \mathbb{I}_{M-1}$, the set $\overline{\Omega}_K$ is composed of the supports of all possible K-sparse strict (local) minimizers of \mathcal{F}_d .

The next propositions checks the existence of $\beta > 0$ ensuring that all the global minimizers of \mathcal{F}_d are K-sparse, for some $K \in \mathbb{I}_{M-1}$.

¹⁵In detail we have $A\hat{u} = A_{\hat{\sigma}}\hat{u}_{\hat{\sigma}} = A_{\tilde{\sigma}}\left(\hat{u}_{\hat{\sigma}} - v_{\tilde{\sigma}} \frac{\hat{u}[i]}{v[i]}\right) = A_{\tilde{\sigma}}\tilde{u}_{\tilde{\sigma}} = A_{\tilde{\sigma}}\tilde{u}_{\tilde{\sigma}} = A\tilde{u}$.

¹⁶Clearly, $\overline{\Omega}_{M-1} = \Omega$.

Proposition 6 Let $d \in \mathbb{R}^M$. For any $K \in \mathbb{I}_{M-1}$, there exists $\beta_K \geq 0$ such that if $\beta > \beta_K$, then each global minimizer \hat{u} of \mathcal{F}_d meets

$$\|\hat{u}\|_0 \leq K \quad \text{and} \quad \sigma(\hat{u}) \in \bar{\Omega}_K. \quad (45)$$

One can choose $\beta_K = \|A\tilde{u} - d\|^2$ where \tilde{u} solves (\mathcal{P}_ω) for some $\omega \in \Omega_K$.

The proof is given in Appendix 8.4 (p. 32). The value of β_K in the statement is easy to compute but in general it is not sharp¹⁷.

5 Uniqueness of the global minimizer of \mathcal{F}_d

The presentation is simplified using the notation introduced next.

Given a matrix $A \in \mathbb{R}^{M \times N}$, with any $\omega \in \Omega$ (see Definition 3), we associate the $M \times M$ matrix Π_ω that yields the orthogonal projection¹⁸ onto the subspace spanned by the columns of A_ω :

$$\Pi_\omega \stackrel{\text{def}}{=} A_\omega (A_\omega^T A_\omega)^{-1} A_\omega^T. \quad (46)$$

For $\omega \in \Omega$, the projector in Remark 1 (p. 6) reads $\Pi_{\text{range}(A_\omega)} = \Pi_\omega$.

Checking if a global minimizer \hat{u} of \mathcal{F}_d is unique requires to compare its value $\mathcal{F}_d(\hat{u})$ with the values $\mathcal{F}_d(\bar{u})$ of the concurrent strict minimizers \bar{u} . Let \hat{u} be an $(M-1)$ -sparse strict (local) minimizer of \mathcal{F}_d . Then $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u}) \in \Omega$. Using Remark 1 (p. 6) shows that

$$\begin{aligned} \mathcal{F}_d(\hat{u}) &= \|A_{\hat{\sigma}}\hat{u} - d\|^2 + \beta \sharp \hat{\sigma} = \|\Pi_{\hat{\sigma}}d - d\|^2 + \beta \sharp \hat{\sigma} \\ &= d^T (I_M - \Pi_{\hat{\sigma}}) d + \beta \sharp \hat{\sigma}. \end{aligned} \quad (47)$$

Let \bar{u} be another $(M-1)$ -sparse strict minimizer of \mathcal{F}_d ; set $\bar{\sigma} \stackrel{\text{def}}{=} \sigma(\bar{u})$. Then

$$\mathcal{F}_d(\hat{u}) - \mathcal{F}_d(\bar{u}) = d^T (\Pi_{\bar{\sigma}} - \Pi_{\hat{\sigma}}) d + \beta (\sharp \hat{\sigma} - \sharp \bar{\sigma}).$$

If both \hat{u} and $\bar{u} \neq \hat{u}$ are global minimizers of \mathcal{F}_d , the equality given above yields

$$\mathcal{F}_d(\hat{u}) = \mathcal{F}_d(\bar{u}) \quad \Leftrightarrow \quad d^T (\Pi_{\bar{\sigma}} - \Pi_{\hat{\sigma}}) d = -\beta (\sharp \hat{\sigma} - \sharp \bar{\sigma}). \quad (48)$$

Equation (48) reveals that the uniqueness of the global minimizer of \mathcal{F}_d cannot be guaranteed without suitable assumptions on A and on d .

5.1 A generic assumption on A

We adopt an assumption on the matrix A in \mathcal{F}_d in order to restrict the cases when (48) takes place for some supports $\hat{\sigma} \neq \bar{\sigma}$ meeting $\sharp \hat{\sigma} = \sharp \bar{\sigma}$.

H1 The matrix $A \in \mathbb{R}^{M \times N}$ is such that for some given $K \in \mathbb{I}_{M-1}$,

$$r \in \mathbb{I}_K \quad \text{and} \quad (\omega, \varpi) \in (\Omega_r \times \Omega_r), \quad \omega \neq \varpi \quad \Rightarrow \quad \Pi_\omega \neq \Pi_\varpi. \quad (49)$$

¹⁷For $\beta \gtrsim \beta_K$ the global minimizers of \mathcal{F}_d might be k -sparse for $k \ll K$. A sharp β_K can be obtained as

$$\beta_K = \min_{\omega \in \Omega_K} \{ \|A\tilde{u} - d\|^2 : \tilde{u} \text{ solve } (\mathcal{P}_\omega) \text{ for } \omega \in \Omega_K \}.$$

¹⁸If $\omega = \emptyset$, we have $A_\omega \in \mathbb{R}^{M \times 0}$ and so $\Pi_{\hat{\sigma}}$ is an $M \times M$ null matrix.

Assumption H1 means that the angle (or the gap) between the equidimensional subspaces $\text{range}(A_\omega)$ and $\text{range}(A_\varpi)$ must be nonzero [28]. For instance, if $(i, j) \in \mathbb{I}_\mathbb{N} \times \mathbb{I}_\mathbb{N}$ meet $i \neq j$, H1 implies that $a_i \neq \kappa a_j$ for any $\kappa \in \mathbb{R} \setminus \{0\}$ since $\Pi_{\{i\}} = a_i a_i^T / \|a_i\|^2$.

Checking if H1 holds for a given matrix A requires a combinatorial search over all possible couples $(\varpi, \omega) \in (\Omega_r \times \Omega_r)$ meeting $\varpi \neq \omega, \forall r \in \mathbb{I}_\mathbb{K}$. This is hard to do. Instead, we wish to know whether or not H1 is a *practical* limitation. Using some auxiliary claims, we shall show that H1 *fails only for a closed negligible subset of matrices* in the space of all $\mathbb{M} \times \mathbb{N}$ real matrices.

Lemma 6 *Given $r \in \mathbb{I}_{\mathbb{M}-1}$ and $\varpi \in \Omega_r$, define the following set of submatrices of A :*

$$\mathcal{H}_\varpi = \left\{ A_\omega : \omega \in \Omega_r \text{ and } \Pi_\omega = \Pi_\varpi \right\} .$$

Then \mathcal{H}_ϖ belongs to an $(r \times r)$ -dimensional subspace of the space of all $\mathbb{M} \times r$ matrices.

Proof. Using that $\varpi \in \Omega_r$ and $\omega \in \Omega_r$, we have¹⁹

$$\Pi_\omega = \Pi_\varpi \quad \Leftrightarrow \quad A_\omega = \Pi_\varpi A_\omega . \quad (50)$$

Therefore \mathcal{H}_ϖ equivalently reads

$$\mathcal{H}_\varpi = \left\{ A_\omega : \omega \in \Omega_r \text{ and } A_\omega = \Pi_\varpi A_\omega \right\} . \quad (51)$$

Let $A_\omega \in \mathcal{H}_\varpi$. Denote the columns of A_ω by \tilde{a}_i for $i \in \mathbb{I}_r$. Then (51) yields

$$\Pi_\varpi \tilde{a}_i = \tilde{a}_i, \quad \forall i \in \mathbb{I}_r \quad \Rightarrow \quad \tilde{a}_i \in \text{range}(A_\varpi), \quad \forall i \in \mathbb{I}_r .$$

Hence all $\tilde{a}_i, i \in \mathbb{I}_r$, live in the r -dimensional vector subspace $\text{range}(A_\varpi)$. All the columns of each matrix $A_\omega \in \mathcal{H}_\varpi$ belong to $\text{range}(A_\varpi)$ as well. It follows that \mathcal{H}_ϖ belongs to a (closed) subspace of dimension $r \times r$ in the space of all $\mathbb{M} \times r$ matrices, where $r \leq \mathbb{M} - 1$. \square

More precisions on the submatrices of A living in \mathcal{H}_ϖ are given next.

Remark 4 *The closed negligible subset \mathcal{H}_ϖ in Lemma 6 is formed from all the submatrices of A that are column equivalent to A_ϖ (see [28, p. 171]), that is*

$$A_\omega \in \mathcal{H}_\varpi \quad \Leftrightarrow \quad \exists P \in \mathbb{R}^{r \times r} \text{ meeting } \text{rank}(P) = r \text{ and } A_\omega = A_\varpi P . \quad (52)$$

Observe that P has r^2 unknowns that must satisfy $\mathbb{M}r$ equations and that P must be invertible. It should be quite exceptional that such a matrix P does exist.

This remark can help to discern structured dictionaries that meet or not H1.

Next we inspect the set of all matrices A failing assumption H1.

¹⁹Using (46) (p. 18), as well as the fact that $A_\omega = \Pi_\omega A_\omega, \forall \omega \in \Omega_r$, one easily derives that

$$\Pi_\omega = \Pi_\varpi \quad \Leftrightarrow \quad \left\{ \begin{array}{l} A_\omega (A_\omega^T A_\omega)^{-1} A_\omega^T = \Pi_\varpi \\ A_\varpi (A_\varpi^T A_\varpi)^{-1} A_\varpi^T = \Pi_\omega \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} A_\omega = \Pi_\varpi A_\omega \\ A_\varpi = \Pi_\omega A_\varpi \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \Pi_\omega = \Pi_\varpi \Pi_\omega \\ \Pi_\varpi = \Pi_\omega \Pi_\varpi \end{array} \right\} \Rightarrow \Pi_\omega = \Pi_\varpi .$$

Lemma 7 Consider the set \mathcal{H} formed from $M \times N$ real matrices described next:

$$\mathcal{H} \stackrel{\text{def}}{=} \left\{ A \in \mathbb{R}^{M \times N} : \exists r \in \mathbb{I}_{M-1}, \exists (\varpi, \omega) \in \Omega_r \times \Omega_r, \varpi \neq \omega \text{ and } \Pi_{\varpi} = \Pi_{\omega} \right\} .$$

Then \mathcal{H} belongs to a finite union of vector subspaces in $\mathbb{R}^{M \times N}$ whose Lebesgue measure in the latter space is null.

Proof. Let $A \in \mathcal{H}$. Then there exists at least one integer $r \in \mathbb{I}_{M-1}$ and at least one pair $(\varpi, \omega) \in \Omega_r \times \Omega_r$ such that $\varpi \neq \omega$ and $\Pi_{\varpi} = \Pi_{\omega}$. Using Lemma 6, A contains (at least) one $M \times r$ submatrix A_{ϖ} belonging to an $r \times r$ vector subspace in the space of all $M \times r$ real matrices. Identifying A with an MN -length vector, its entries are included in a vector subspace of \mathbb{R}^{MN} of dimension no larger than $MN - 1$. The claim of the lemma is straightforward. \square

We can now pleasantly clarify assumption H1.

Theorem 3 Given an arbitrary $K \in \mathbb{I}_{M-1}$, consider the set of $M \times N$ real matrices below

$$\mathcal{A}_K \stackrel{\text{def}}{=} \left\{ A \in \mathbb{R}^{M \times N} : A \text{ satisfies H1 for } K \right\} .$$

Then \mathcal{A}_K contains an open and dense subset in the space of all $M \times N$ real-valued matrices.

Proof. The complement of \mathcal{A}_K in the space of all $M \times N$ real matrices reads

$$\mathcal{A}_K^c = \left\{ A \in \mathbb{R}^{M \times N} : \text{H1 fails for } A \text{ and } K \right\} .$$

It is clear that $\mathcal{A}_K^c \subset \mathcal{H}$, where \mathcal{H} is described in Lemma 7. By the same lemma, \mathcal{A}_K^c is included in a closed subset of vector subspaces in $\mathbb{R}^{M \times N}$ whose Lebesgue measure in the latter space is null. Consequently, \mathcal{A}_K fits the statement. \square

For any $K \in \mathbb{I}_{M-1}$, H1 is a *generic* property of all $M \times N$ real matrices meeting $M < N$.

We can note that

$$\mathcal{A}_{K+1} \subseteq \mathcal{A}_K, \quad \forall K \in \mathbb{I}_{M-2} .$$

One can hence presume that H1 is weakened as K decreases. This issue is illustrated in section 6.

5.2 A generic assumption on d

A preliminary result is stated next.

Lemma 8 Let $(\omega, \varpi) \in \overline{\Omega}_K \times \overline{\Omega}_K$ for $\omega \neq \varpi$ and H1 hold for $K \in \mathbb{I}_{M-1}$. Given $\kappa \in \mathbb{R}$, define

$$\mathbb{T}_{\kappa} \stackrel{\text{def}}{=} \{g \in \mathbb{R}^M : g^T (\Pi_{\omega} - \Pi_{\varpi}) g = \kappa\} .$$

Then \mathbb{T}_{κ} is a closed subset of \mathbb{R}^M and $\mathbb{L}^M(\mathbb{T}_{\kappa}) = 0$.

Proof. Define $f : \mathbb{R}^M \rightarrow \mathbb{R}$ by $f(g) = g^T (\Pi_{\omega} - \Pi_{\varpi}) g$. Then

$$\mathbb{T}_{\kappa} = \{g \in \mathbb{R}^M : f(g) = \kappa\} . \tag{53}$$

Using H1, T_κ is closed in \mathbb{R}^M . Set

$$Q = \{g \in \mathbb{R}^M : \nabla f(g) \neq 0\} \quad \text{and} \quad Q^c = \mathbb{R}^M \setminus Q.$$

Consider an arbitrary $g \in T_\kappa \cap Q$. From H1, $\text{rank}(\nabla f(g)) = 1$. For simplicity, assume that

$$\nabla f(g)[M] = \frac{df(g)}{dg[M]} \neq 0.$$

By the implicit functions theorem, there are open neighborhoods $\mathcal{O}_g \subset Q \subset \mathbb{R}^M$ and $\mathcal{V} \subset \mathbb{R}^{M-1}$ of g and $g_{\mathbb{I}_{M-1}}$, respectively, and a unique \mathcal{C}^1 -function $h_g : \mathcal{V} \rightarrow \mathbb{R}$ with ∇h_g bounded, such that

$$\gamma = (\gamma_{\mathbb{I}_{M-1}}, \gamma[M]) \in \mathcal{O}_g \quad \text{and} \quad f(\gamma) = \kappa \quad \iff \quad \gamma_{\mathbb{I}_{M-1}} \in \mathcal{V} \quad \text{and} \quad \gamma[M] = h_g(\gamma_{\mathbb{I}_{M-1}}). \quad (54)$$

From (53) and (54) it follows that²⁰

$$\mathcal{O}_g \cap T_\kappa = \psi^g \left(\mathcal{O}_g \cap (\mathbb{R}^{M-1} \times \{0\}) \right),$$

where ψ^g is a diffeomorphism on \mathcal{O}_g given by

$$\psi_i^g(\gamma) = \gamma[i], \quad 1 \leq i \leq M-1 \quad \text{and} \quad \psi_M^g(\gamma) = h_g(\gamma_{\mathbb{I}_{M-1}}) + \gamma[M]. \quad (55)$$

Since $\mathbb{L}^M(\mathcal{O}_g \cap (\mathbb{R}^{M-1} \times \{0\})) = 0$ and $\nabla \psi^g$ is bounded on \mathcal{O}_g , it follows from [36, Lemma 7.25] that²¹ $\mathbb{L}^M(\mathcal{V}_g \cap T_\kappa) = 0$. We have thus obtained that

$$S \subset Q, \quad S \text{ bounded} \quad \Rightarrow \quad \mathbb{L}^M(S \cap T_\kappa) = 0. \quad (56)$$

Using that every open subset of \mathbb{R}^M can be written as a countable union²² of cubes in \mathbb{R}^M [35, 15, 37], (56) entails that $\mathbb{L}^M(T_\kappa \cap Q) = 0$.

Next, $Q^c = \ker(\Pi_\omega - \Pi_\varpi)$ where $\dim \ker(\Pi_\omega - \Pi_\varpi) \leq M-1$ by H1. Then $\mathbb{L}^M(T_\kappa \cap Q^c) = 0$. The statement follows from the equality $\mathbb{L}^M(T_\kappa) = \mathbb{L}^M(T_\kappa \cap Q) + \mathbb{L}^M(T_\kappa \cap Q^c)$. \square

We exhibit a *closed negligible* subset of data in \mathbb{R}^M that can still meet the equality in (48), p. 18.

Proposition 7 For $\beta > 0$ and $K \in \mathbb{I}_{M-1}$, put

$$\Sigma_K \stackrel{\text{def}}{=} \bigcup_{n=-K}^K \bigcup_{\omega \in \bar{\Omega}_K} \bigcup_{\varpi \in \bar{\Omega}_K} \left\{ g \in \mathbb{R}^M : \omega \neq \varpi \quad \text{and} \quad g^T (\Pi_\omega - \Pi_\varpi) g = n\beta \right\}, \quad (57)$$

where $\bar{\Omega}_K$ is given in (44), p. 17. Let H1 hold for K . Then Σ_K is closed in \mathbb{R}^M and $\mathbb{L}^M(\Sigma_K) = 0$.

Proof. For some $n \in \{-K, \dots, K\}$ and $(\omega, \varpi) \in (\bar{\Omega}_K \times \bar{\Omega}_K)$ meeting $\omega \neq \varpi$, put

$$\Sigma \stackrel{\text{def}}{=} \left\{ g \in \mathbb{R}^M : g^T (\Pi_\omega - \Pi_\varpi) g = n\beta \right\}.$$

If $\sharp \omega \neq \sharp \varpi$, then $\text{rank}(\Pi_\omega - \Pi_\varpi) \geq 1$. If $\sharp \omega = \sharp \varpi$, H1 guarantees that $\text{rank}(\Pi_\omega - \Pi_\varpi) \geq 1$, yet again. The number $n\beta \in \mathbb{R}$ is given. According to Lemma 8, Σ is a closed subset of \mathbb{R}^M and $\mathbb{L}^M(\Sigma) = 0$. The conclusion follows from the fact that Σ_K is a finite union of subsets like Σ . \square

We assume thereafter that if H1 holds for some $K \in \mathbb{I}_{M-1}$, data d satisfy

$$d \in \{g \in \mathbb{R}^M : g \notin \Sigma_K\} = \mathbb{R}^M \setminus \Sigma_K.$$

²⁰From (54), \mathcal{V} is the restriction of \mathcal{O}_g to \mathbb{R}^{M-1} .

²¹The same result follows from the change-of-variables theorem for the Lebesgue integral, see e.g. [36].

²²From (56), adjacent cubes can also intersect in our case.

5.3 The unique global minimizer of \mathcal{F}_d is K -sparse for $K \leq (M - 1)$

We are looking for guarantees that \mathcal{F}_d has a *unique* global minimizer \hat{u} obeying

$$\|\hat{u}\|_0 \leq K \text{ for some fixed } K \in \mathbb{I}_{M-1} .$$

This is the aim of the next theorem.

Theorem 4 *Given $K \in \mathbb{I}_{M-1}$, let H1 hold for K , $\beta > \beta_K$ where β_K meets Proposition 6 (p. 18) and $\Sigma_K \subset \mathbb{R}^M$ read as in (57). Consider that*

$$d \in \mathbb{R}^M \setminus \Sigma_K .$$

Then

- (i) the set $\mathbb{R}^M \setminus \Sigma_K$ is open and dense in \mathbb{R}^M ;
- (ii) \mathcal{F}_d has a unique global minimizer \hat{u} satisfying $\|\hat{u}\|_0 \leq K$.

Proof. Statement (i) follows from Proposition 7.

Since $\beta > \beta_K$, all global minimizers of \mathcal{F}_d have their support in $\bar{\Omega}_K$ (Proposition 6). Using that data meet $d \in \mathbb{R}^M \setminus \Sigma_K$, the definition of Σ_K in (57) shows that

$$-K \leq n \leq K \text{ and } (\omega, \varpi) \in (\bar{\Omega}_K \times \bar{\Omega}_K), \omega \neq \varpi \Rightarrow d^T (\Pi_\omega - \Pi_\varpi) d \neq n\beta . \quad (58)$$

The proof is conducted by contradiction. Let \hat{u} and $\bar{u} \neq \hat{u}$ be two global minimizers of \mathcal{F}_d . Then

$$\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u}) \in \bar{\Omega}_K \text{ and } \bar{\sigma} \stackrel{\text{def}}{=} \sigma(\bar{u}) \in \bar{\Omega}_K ,$$

and $\hat{\sigma} \neq \bar{\sigma}$. By $\mathcal{F}_d(\hat{u}) = \mathcal{F}_d(\bar{u})$, equation (48) (p. 18) reads

$$d^T (\Pi_{\hat{\sigma}} - \Pi_{\bar{\sigma}}) d = -\beta (\#\hat{\sigma} - \#\bar{\sigma}) . \quad (59)$$

An enumeration of all possible values of $\#\hat{\sigma} - \#\bar{\sigma}$ shows that

$$\beta (\#\hat{\sigma} - \#\bar{\sigma}) = n\beta \text{ for some } n \in \{-K, \dots, K\} .$$

Inserting this equation into (59) shows that

$$d^T (\Pi_{\hat{\sigma}} - \Pi_{\bar{\sigma}}) d = n\beta \text{ for some } n \in \{-K, \dots, K\} .$$

The last result contradicts (58), hence it violates the assumptions H1 and $d \in \mathbb{R}^M \setminus \Sigma_K$. Consequently, \mathcal{F}_d cannot have two global minimizers. Since \mathcal{F}_d has always global minimizers (Theorem 2(i), p. 16), it follows that \mathcal{F}_d has a unique global minimizer. \square

For $\beta > \beta_K$, the objective \mathcal{F}_d in (1) has a unique K -sparse global minimizer where $K \leq M - 1$. For all $K \in \mathbb{I}_{M-1}$, the claim holds true in a generic sense.

The proof of Theorem 4 shows a wider result: if H1 holds for some $K \in \mathbb{I}_{M-1}$, $\beta > \beta_K$ and $d \notin \Sigma_K$, any two *distinct* K -sparse strict (local) minimizers \hat{u} and \bar{u} of \mathcal{F}_d satisfy $\mathcal{F}_d(\hat{u}) \neq \mathcal{F}_d(\bar{u})$.

6 Numerical illustrations

6.1 On assumption H1

Assumption H1 (p. 18) requires that $\Pi_\omega \neq \Pi_\varpi$ when $(\omega, \varpi) \in \Omega_r \times \Omega_r$, $\omega \neq \varpi$ for all $r \leq K \in \mathbb{I}_{M-1}$. From a practical viewpoint, the magnitude of $(\Pi_\omega - \Pi_\varpi)$ should be discernible. One way to assess the viability of H1 for a matrix A and $K \in \mathbb{I}_{M-1}$ is to calculate

$$\xi_K(A) \stackrel{\text{def}}{=} \min_{r \in \mathbb{I}_K} \mu_r(A), \quad (60)$$

where

$$\mu_r(A) = \min_{\substack{(\omega, \varpi) \in \Omega_r \times \Omega_r \\ \omega \neq \varpi}} \|\Pi_\omega - \Pi_\varpi\|_2, \quad \forall r \in \mathbb{I}_K.$$

In fact, $\|\Pi_\omega - \Pi_\varpi\|_2 = \sin(\theta)$ where $\theta \in [0, \pi/2]$ is the maximum angle between $\text{range}(A_\omega)$ and $\text{range}(A_\varpi)$, see [28, p. 456]. The latter subspaces have the same dimension and $\Pi_\omega \neq \Pi_\varpi$ when $(\omega, \varpi) \in \Omega_r \times \Omega_r$, $\omega \neq \varpi$ and $r \in \mathbb{I}_K$, hence $\theta \in (0, \pi/2]$. Consequently,

$$\text{H1 holds for } K \in \mathbb{I}_{M-1} \Rightarrow \mu_r(A) \in (0, 1], \quad \forall r \in \mathbb{I}_K \Rightarrow \xi_K(A) \in (0, 1].$$

According to (60), we have $\xi_K \geq \xi_{K+1}$, $\forall K \in \mathbb{I}_{M-2}$. Our guess that *assumption H1 is lightened when K decreases* (see the comments following Theorem 3 on p. 20) means that

$$\xi_1(A) > \dots > \xi_{M-1}(A). \quad (61)$$

We provide numerical tests on two subsets of real-valued random matrices for $M = 5$ and $N = 10$, denoted by \mathcal{A}_{20}^N and \mathcal{A}_{1000}^U . The values of $\xi_K(\cdot)$, $K \in \mathbb{I}_{M-1} = \mathbb{I}_4$, for every matrix in \mathcal{A}_{20}^N and in \mathcal{A}_{1000}^U , were calculated using *exhaustive combinatorial search*. All tested matrices satisfy *assumption H1*, which confirms Theorem 3 (p. 20) and its consequences. In order to evaluate the extent of H1, we computed the *worst* and the *best* values of $\xi_K(\cdot)$ over these sets:

$$\begin{cases} \xi_K^{\text{worst}} = \min_{A \in \mathcal{A}} \xi_K(A), \\ \xi_K^{\text{best}} = \max_{A \in \mathcal{A}} \xi_K(A), \end{cases} \quad \forall K \in \mathbb{I}_{M-1}, \quad \mathcal{A} \in \{\mathcal{A}_{20}^N, \mathcal{A}_{1000}^U\}. \quad (62)$$

Set \mathcal{A}_{20}^N . This set was formed from 20 matrices A^n , $n \in \mathbb{I}_{20}$ of size 5×10 . The components of each matrix A^n were independent and uniformly drawn from the standard normal distribution with mean zero and variance one. The values of $\xi_K(\cdot)$ are depicted in Fig. 1. We have²³ $\xi_1(A^{10}) = \xi_2(A^{10})$ and $\xi_1(A^{17}) = \xi_2(A^{17})$. In all other cases (61) is satisfied. Fig. 1 clearly shows that in general $\xi_K(\cdot)$ increases substantially as K decreases (from $M - 1$ to 1).

The *worst* and the *best* values of $\xi_K(\cdot)$ over the whole set \mathcal{A}_{20}^N are displayed in Table 1.

	$K = 1$	$K = 2$	$K = 3$	$K = 4$
ξ_K^{worst}	0.3519	0.1467	0.0676	0.0072
ξ_K^{best}	0.8666	0.5881	0.3966	0.0785

Table 1: The worst and the best values of $\xi_K(A)$, for $K \in \mathbb{I}_{M-1}$, over the set \mathcal{A}_{20}^N , see (62).

²³This is why on the figure, in columns 10 and 17, the green “o” and the red “◇” overlap.




Figure 1: x -axis: the list of the 20 random matrices in \mathcal{A}_{20}^N . (a) y -axis: the value $\xi_K(A^n)$ according to (60), $\forall K \in \mathbb{I}_{M-1}$ and $\forall n \in \mathbb{I}_{20}$. The plot in (b) is a zoom of (a) along the y -axis.

Set \mathcal{A}_{1000}^U . The set \mathcal{A}_{1000}^U was composed of one thousand 5×10 matrices A^n , $n \in \mathbb{I}_{1000}$. The entries of each matrix A^n were independent and uniformly sampled on $[-1, 1]$. The obtained values for ξ_K^{worst} and ξ_K^{best} , calculated according to (62), are shown in Table 2. For $K \in \mathbb{I}_3$, the

	K = 1	K = 2	K = 3	K = 4
ξ_K^{worst}	0.1085	0.0235	0.0045	0.0001
ξ_K^{best}	0.9526	0.8625	0.5379	0.1152

Table 2: The worst and the best values of $\xi_K(A)$, for $K \in \mathbb{I}_{M-1}$, over the set \mathcal{A}_{1000}^U .

best values of $\xi_K(\cdot)$ were obtained for the same matrix, A^{964} . Note that $\xi_4(A^{964}) = 0.0425 \gg \xi_4^{worst}$. The *worst* values in Table 2 are smaller than those in Table 1 while the *best* values in Table 2 are larger than those in Table 1; one credible reason is that \mathcal{A}_{1000}^U is much larger than \mathcal{A}_{20}^N .

Overall, equation (61) is satisfied on \mathcal{A}_{1000}^U —the percentages in Table 3 are pretty small.

	$\xi_1(A^n) = \xi_2(A^n)$	$\xi_2(A^n) = \xi_3(A^n)$	$\xi_3(A^n) = \xi_4(A^n)$
occurrences $\{n\}$	5 %	1.6 %	0.1 %

Table 3: Percentage of the cases in \mathcal{A}_{1000}^U when (61) fails to hold.

Table 3 and Figure 1 agree with our guess that H1 is more viable for smaller values of K .

Based on the magnitudes for ξ_K^{best} in Tables 1 and 2, one can expect that there are some classes of matrices (random or not) that fit H1 for larger values of $\xi_K(\cdot)$.

6.2 On the global minimizers of \mathcal{F}_d

Here we summarize the outcome of a series of experiments corresponding to several matrices $A \in \mathbb{R}^{M \times N}$ where $M = 5$ and $N = 10$, satisfying H1 for $K = M - 1$, different original vectors $\ddot{u} \in \mathbb{R}^N$ and data samples $d = A\ddot{u} + \text{noise}$, for various values of $\beta > 0$. In each experiment, we computed the complete list of all different strict (local) minimizers of \mathcal{F}_d , say $(\hat{u}^i)_{i=1}^n$. Then the sequence of values $(\mathcal{F}_d(\hat{u}^i))_{i=1}^n$ was sorted in increasing order, $\mathcal{F}_d(\hat{u}^{i_1}) \leq \mathcal{F}_d(\hat{u}^{i_2}) \leq \dots \leq \mathcal{F}_d(\hat{u}^{i_n})$. A global minimizer \hat{u}^{i_1} is unique provided that $\mathcal{F}_d(\hat{u}^{i_1}) < \mathcal{F}_d(\hat{u}^{i_2})$. In order to discard numerical errors, we also checked if $|\mathcal{F}_d(\hat{u}^{i_1}) - \mathcal{F}_d(\hat{u}^{i_2})|$ is well detectable.

In all experiments we carried out, the following facts were observed:

- The global minimizer of \mathcal{F}_d was unique—manifestly data d did never belong to the *closed negligible* subset Σ_K in Proposition 7 (p. 21). This confirms Theorem 4 (p. 22).
- The global minimizers of \mathcal{F}_d remained unchanged under large variations of β .
- The necessary condition for a global minimizer in Proposition 4 was met.

Next we present in detail two of these experiments where \mathcal{F}_d is defined using

$$A = \begin{bmatrix} 7 & 2 & 4 & 9 & 0 & 3 & 3 & 6 & 6 & 7 \\ 3 & 4 & 9 & 3 & 3 & 9 & 1 & 3 & 1 & 5 \\ 5 & 4 & 2 & 4 & 0 & 7 & 1 & 9 & 2 & 9 \\ 8 & 4 & 0 & 9 & 6 & 0 & 4 & 2 & 3 & 7 \\ 6 & 3 & 6 & 5 & 0 & 9 & 0 & 0 & 3 & 8 \end{bmatrix} \quad \begin{aligned} d &= A\ddot{u} + n, \\ \text{where } n &\text{ is noise and} \\ \ddot{u} &= (0, 1, 8, 0, 3, 0, 0, 0, 0, 9)^T. \end{aligned} \quad (63)$$

Only integers appear in (63) for better readability. We have $\text{rank}(A) = M = 5$. An exhaustive combinatorial test shows that the arbitrary matrix A in (63) satisfies H1 for $K = M - 1$. The values of $\xi_K(A)$ are seen in Table 4. One notes that $\mu_2(A) > \mu_1(A)$, hence $\xi_1(A) = \xi_2(A)$.

	$K = 1$	$K = 2$	$K = 3$	$K = 4$
$\xi_K(A)$	0.2737	0.2737	0.2008	0.0564
$\mu_K(A)$	0.2737	0.2799	0.2008	0.0564

Table 4: The values of $\xi_K(A)$ and $\mu_K(A)$, $\forall K \in \mathbb{I}_{M-1}$, for the matrix A in (63).

A way to rate the quality of the solution provided by a global minimizer \hat{u} of \mathcal{F}_d is to compute the mean square error (MSE) of the retained non-zero values $\hat{u}_{\sigma(\hat{u})}$ with respect to $\ddot{u}_{\sigma(\hat{u})}$. So

$$\text{MSE}(\ddot{u}_{\hat{\sigma}}, \hat{u}_{\hat{\sigma}}) = \frac{1}{\#\hat{\sigma}} \sum_{i \in \hat{\sigma}} (\hat{u}[i] - \ddot{u}[i])^2 \quad \text{where } \hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u}). \quad (64)$$

The rationale is that one expects that the global minimizer \hat{u} of \mathcal{F}_d obeys $\hat{\sigma} \subseteq \sigma(\ddot{u})$ where \ddot{u} is the original in (63) and that the vanished entries of \hat{u} correspond to the leas entries of \ddot{u} .

The experiments described hereafter correspond to two data samples relevant to (63)—without and with noise—and to several values of $\beta > 0$.

β	The global minimizer \hat{u} of \mathcal{F}_d (<i>row vector</i>)									$\ \hat{u}\ _0$	$\mathcal{F}_d(\hat{u})$	MSE($\hat{u}_{\hat{\sigma}}, \hat{u}_{\hat{\sigma}}$)	
1	0	1	8	0	3	0	0	0	0	9	4	4	0
10^2	0	0	8.12	0	3.31	0	0	0	0	9.33	3	301.52	0.073
10^3	0	0	0	0	0	12.58	20.28	0	0	0	2	2179.3	284.7
10^4	0	29.95	0	0	0	0	0	0	0	0	1	14144	838.2
$7 \cdot 10^4$	0	0	0	0	0	0	0	0	0	0	0	58864	—
\ddot{u}	= 0	1	8	0	3	0	0	0	0	9			

Table 5: The global minimizer \hat{u} of \mathcal{F}_d and its value $\mathcal{F}_d(\hat{u})$ for the noise-free data d in (65) for different values of β . Last column: the error defined in (64). Last row: the original \ddot{u} in (63).

Noise-free data. The noise-free data in (63) read:

$$d = A\ddot{u} = (97, 130, 101, 85, 123)^T. \quad (65)$$

For different values of β , the global minimizer \hat{u} is given in Table 5. Since $\sigma(\ddot{u}) \in \Omega$ and data are noise-free, \mathcal{F}_d does not have global minimizers with $\|\hat{u}\|_0 = 5$. Actually, applying Proposition 6(i) for $\tilde{u} = \ddot{u}$ yields $\beta_{M-1} = 0$, hence for any $\beta > 0$ all global minimizers of \mathcal{F}_d have a support in $\Omega = \overline{\Omega}_{M-1}$ (see Definition 3 on p. 9 and (44) on p. 17). The global minimizer \hat{u} for $\beta = 1$ meets $\hat{u} = \ddot{u}$. For $\beta = 100$, the global minimizer \hat{u} obeys $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u}) = \{3, 5, 10\} \subsetneq \sigma(\ddot{u})$ and $\|\hat{u}\|_0 = 3$ —the least nonzero entry of the original \ddot{u} is canceled, which is reasonable. The global minimizers corresponding to $\beta \geq 300$ are meaningless. We could not find any value of β giving better 2-sparse global minimizers. Reminding that data are noise-free, this confirms the discussion next to Corollary 1 (p. 9): the global minimizers of \mathcal{F}_d realize an only *pseudo-hard* thresholding. For $\beta \geq 7 \cdot 10^4 > \|d\|^2$, the global minimizer of \mathcal{F}_d is $\hat{u} = 0$ which confirms Remark 3 (p. 15).

Noisy data. Now we consider *noisy data* in (63) for

$$n = (4, -1, 2, -3, 5)^T. \quad (66)$$

This arbitrary noise yields a signal-to-noise ratio²⁴ (SNR) equal to 14.07 dB. If $\beta \leq 0.04$, \mathcal{F}_d has 252 different strict global minimizers \hat{u} obeying $\|\hat{u}\|_0 = M$ and $\mathcal{F}_d(\hat{u}) = \beta M$ (remind Proposition 3 on p. 14). For $\beta \geq 0.05$, the global minimizer \hat{u} of \mathcal{F}_d is unique and meets $\sigma(\hat{u}) \in \Omega$. For several values of $\beta \geq 0.05$, it is given in Table 6. For $\beta = 1$, the global minimizer is meaningless. We

β	The global minimizer \hat{u} of \mathcal{F}_d (<i>row vector</i>)									$\ \hat{u}\ _0$	$\mathcal{F}_d(\hat{u})$	MSE($\hat{u}_{\hat{\sigma}}, \hat{u}_{\hat{\sigma}}$)	
1	0	6.02	2.66	6.43	0	6.85	0	0	0	0	4	4.0436	35.5
10^2	0	0	8.23	0	2.3	0	0	0	0	9.71	3	301.94	0.35
10^3	0	0	8.14	0	0	0	0	0	0	10.25	2	2174.8	0.79
10^4	0	0	0	0	0	0	0	0	0	14.47	1	14473	29.9
$7 \cdot 10^4$	0	0	0	0	0	0	0	0	0	0	0	60559	—
\ddot{u}	= 0	1	8	0	3	0	0	0	0	9			

Table 6: The global minimizer \hat{u} of \mathcal{F}_d and its value $\mathcal{F}_d(\hat{u})$ for noisy data given by (63) and (66), for different values of β . Last column: the error defined in (64). Last row: the original \ddot{u} .

could not find any value of β yielding a better global minimizer with a 4-length support. For

²⁴Let us denote $\ddot{d} = A\ddot{u}$ and $d = \ddot{d} + n$. The SNR reads [41] $\text{SNR}(\ddot{d}, d) = 10 \log_{10} \frac{\sum_{i=1}^M (\ddot{d}[i] - \frac{1}{M} \sum_{i=1}^M \ddot{d}[i])^2}{\sum_{i=1}^M (\ddot{d}[i] - \ddot{d}[i])^2}$.

the other values of β , the global minimizer \hat{u} meets $\hat{\sigma} \stackrel{\text{def}}{=} \sigma(\hat{u}) \subsetneq \sigma(\ddot{u})$ and its vanished entries correspond to the least entries in the original \ddot{u} . For $\beta = 100$, the global minimizer seems to furnish a good approximation to \ddot{u} . Observe that the last entry of the global minimizer $\hat{u}[10]$, corresponding to the largest magnitude in \ddot{u} , freely increases when β increases from 10^2 to 10^4 . We tested a tight sequence of intermediate values of β without finding better results. Yet again, $\beta \geq 7 \cdot 10^4 > \|d\|^2$ leads to a unique null global minimizer (see Remark 3).




Figure 2: All **638** strict (local) minima of \mathcal{F}_d in (63) for $\beta = 100$ and data d corrupted with the arbitrary noise in (66). The x -axis lists all strict (local) minimizers $\{\hat{u}\}$ of \mathcal{F}_d sorted according to their ℓ_0 -norm $\|\hat{u}\|_0$ in increasing order. (a) The y -axis shows the value $\mathcal{F}_d(\hat{u})$ of these minimizers marked with a star. The value of \mathcal{F}_d for $\hat{u} = 0$ is not shown because it is too large ($\mathcal{F}_d(0) = 60559 = \|d\|^2$). (b) A zoom of (a) along the y -axis. It clearly shows that \mathcal{F}_d has a well recognizable unique global minimizer.

Figure 2 shows the value $\mathcal{F}_d(\hat{u})$ of all the strict local minimizers of \mathcal{F}_d for $\beta = 100$. On the zoom in Figure 2(b) it is well seen that the global minimizer is unique (remember Theorem 4). It meets $\|\hat{u}\|_0 = 3$ and $\mathcal{F}_d(\hat{u}) = 301.94$. One observes that \mathcal{F}_d has $252 = \sharp\Omega_M$ different strict local minimizers \hat{u} meeting $\|\hat{u}\|_0 = 5 = M$ and $\mathcal{F}_d(\hat{u}) = 500 = \beta M$. This confirms Proposition 3 (p. 14)—obviously d does not belong to the closed negligible subset \mathcal{Q}_M described in (36).

7 Conclusions and perspectives

We provided a detailed analysis of the (local and global) minimizers of a regularized objective \mathcal{F}_d composed of an ℓ_2 data fidelity term and an ℓ_0 penalty weighted by a parameter $\beta > 0$, as given in (1), p. 1. We exhibited easy necessary and sufficient conditions ensuring that a (local) minimizer \hat{u} of \mathcal{F}_d is strict (Theorem 1). The global minimizers of \mathcal{F}_d (whose existence was proved) were shown to be strict as well (Theorem 2). Under very mild conditions, \mathcal{F}_d was shown to have a unique global minimizer (Theorem 4). Other interesting results were listed in the Abstract. Below we evoke some perspectives and open questions raised by this work.

- The relationship between the value of the regularization parameter β and the sparsity of the global minimizers of \mathcal{F}_d (Proposition 6, p. 18) can be improved.

- The *generic* linearity in data d of each strict (local) minimizer of \mathcal{F}_d (subsection 3.2, p. 12) should be exploited to better characterize the global minimizers of \mathcal{F}_d .
- Is there a simple way to check if assumption H1 (p. 18) is satisfied by a given matrix $A \in \mathbb{R}^{M \times N}$ when $M < N$ are large? Remark 4 and in particular equation (52), p. 19, could help to discard some non-random matrices. Conversely, one can ask if there is a systematic way to construct matrices A that satisfy H1.

An alternative would be to exhibit families of matrices that satisfy H1 for large values of $\xi_{\mathcal{K}}(\cdot)$, where the latter quantifiers are defined in equation (60) (p. 23).

- A proper adaptation of the results to matrices A and data d with complex entries should not present inherent difficulties.
- The theory developed here can be extended to MAP energies of the form evoked in (6), p. 2. This is important for the imaging applications mentioned there.
- Based on the comments next to Corollary 1 and the numerical tests in subsection 6.2, one has right of asking for conditions ensuring that the global minimizers of \mathcal{F}_d perform a honorable work. Given the high quality of the numerical results provided in [32, 33], the question deserves attention.

There exist numerous algorithms aimed at approximating a (local) minimizer of \mathcal{F}_d . As a by-product of our research, we got simple rules to verify whether or not an algorithm could find

- a (local) minimizer \hat{u} of \mathcal{F}_d —by checking if \hat{u} satisfies (27) in Corollary 1 (p. 9);
- and if this local minimizer is strict—by testing if the submatrix whose columns are indexed by the support of \hat{u} (i. e. $A_{\hat{\sigma}}$) has full column rank (Theorem 1, p. 10).

Some properties of the minimizers of \mathcal{F}_d given in this work can be inserted in numerical schemes in order to quickly escape from shallow local minimizers.

Many existing numerical methods involve a studious choice of the regularization parameter β and some of them are proved to converge to a local minimizer of \mathcal{F}_d . *We have seen that finding a (strict or nonstrict) local minimizer of \mathcal{F}_d is easy and that it is independent of the value of β* (Corollary 1, p. 9 and Corollary 2, p. 11). It is therefore obscure what meaning to attach to “choosing a good β and proving convergence”. There is a *theoretical gap* that needs clarification. Other successful algorithms are not guaranteed to converge to a local minimizer of \mathcal{F}_d . Whenever algorithms do a good job, the choice of β , the assumptions on A and on $\|\hat{u}\|_0$, the iterative scheme and its initialization *obviously* provide a tool for selecting a meaningful solution by minimizing \mathcal{F}_d .

The connection between the existing algorithms and the description of the minimizers exposed in this article deserves a deep exploration.

What conditions ensure that an algorithm minimizing \mathcal{F}_d yields meaningful solutions?

One can expect that such a research gives rise to innovative and more efficient algorithms enabling to compute relevant solutions by minimizing the tricky objective \mathcal{F}_d .

8 Appendix

8.1 Proof of Lemma 1, p. 7

Since $\hat{u} \neq 0$, the definition of $\hat{\sigma}$ shows that $\min_{i \in \hat{\sigma}} |\hat{u}[i]| > 0$. Then ρ in (20) meets $\rho > 0$.

Proof of (i). Since $\#\hat{\sigma} \geq 1$, we have

$$\begin{aligned}
i \in \hat{\sigma}, v \in B_\infty(0, \rho) &\Rightarrow \max_{j \in \hat{\sigma}} |v[j]| < \rho \\
&\Rightarrow \max_{j \in \hat{\sigma}} |v[j]| < \min_{j \in \hat{\sigma}} |\hat{u}[j]| \\
&\Rightarrow |\hat{u}[i] + v[i]| \geq |\hat{u}[i]| - |v[i]| \\
&\quad \geq \min_{j \in \hat{\sigma}} |\hat{u}[j]| - \max_{j \in \hat{\sigma}} |v[j]| \geq \rho - \max_{j \in \hat{\sigma}} |v[j]| > 0 \\
&\Rightarrow \hat{u}[i] + v[i] \neq 0 \\
\left[\text{by (2), p. 2} \right] &\Rightarrow \phi(\hat{u}[i] + v[i]) = \phi(\hat{u}[i]) = 1. \tag{67}
\end{aligned}$$

If $\hat{\sigma}^c = \emptyset$ the result is proved. Let $\hat{\sigma}^c \neq \emptyset$. Then $\hat{u}[i] = 0 = \phi(\hat{u}[i])$, $\forall i \in \hat{\sigma}^c$. Statement (i) is obtained by inserting the latter and (67) into

$$\sum_{i \in \mathbb{I}_N} \phi(\hat{u}[i] + v[i]) = \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i] + v[i]) + \sum_{i \in \hat{\sigma}^c} \phi(\hat{u}[i] + v[i]).$$

Proof of (ii). Using that $\|A(\hat{u} + v) - d\|^2 = \|A\hat{u} - d\|^2 + \|Av\|^2 + 2\langle Av, A\hat{u} - d \rangle$, one obtains

$$\begin{aligned}
v \in B_\infty(0, \rho) \setminus K_{\hat{\sigma}} &\Rightarrow \mathcal{F}_d(\hat{u} + v) = \|A\hat{u} - d\|^2 + \|Av\|^2 + 2\langle Av, A\hat{u} - d \rangle \\
\left[\text{by Lemma 1(i)} \right] &\quad + \beta \sum_{i \in \hat{\sigma}} \phi(\hat{u}[i]) + \beta \sum_{i \in \hat{\sigma}^c} \phi(v[i]) \\
\left[\text{using (4), p. 2} \right] &= \mathcal{F}_d(\hat{u}) + \|Av\|^2 + 2\langle Av, A\hat{u} - d \rangle + \beta \sum_{i \in \hat{\sigma}^c} \phi(v[i]) \\
&\geq \mathcal{F}_d(\hat{u}) - |2\langle v, A^T(A\hat{u} - d) \rangle| + \beta \|v_{\hat{\sigma}^c}\|_0 \\
\left[\text{by Hölder's inequality} \right] &\geq \mathcal{F}_d(\hat{u}) - 2\|v\|_\infty \|A^T(A\hat{u} - d)\|_1 + \beta \|v_{\hat{\sigma}^c}\|_0. \tag{68}
\end{aligned}$$

If $\#\hat{\sigma}^c = 0$ then $K_{\hat{\sigma}} = \mathbb{R}^N$, so $v \in \mathbb{R}^0$ and $\|v\|_\infty = 0$, hence the inequality.

Let $\#\hat{\sigma}^c \geq 1$. For $v \notin K_{\hat{\sigma}}$, there at least one index $i \in \hat{\sigma}^c$ such that $v[i] \neq 0$, hence $\|v_{\hat{\sigma}^c}\|_0 \geq 1$. The definition of ρ in (20) shows that

$$\begin{aligned}
v \in B_\infty(0, \rho) \setminus K_{\hat{\sigma}} &\Rightarrow -\|v\|_\infty > -\rho \geq -\frac{\beta}{2(\|A^T(A\hat{u} - d)\|_1 + 1)} \\
&\Rightarrow -2\|v\|_\infty \|A^T(A\hat{u} - d)\|_1 + \beta \|v_{\hat{\sigma}^c}\|_0 > -\frac{2\beta \|A^T(A\hat{u} - d)\|_1}{2(\|A^T(A\hat{u} - d)\|_1 + 1)} + \beta > 0.
\end{aligned}$$

Introducing the last inequality into (68) shows that for $\#\hat{\sigma}^c \geq 1$, the inequality in (ii) is strict.

8.2 Proof of Proposition 4, p. 15

If $\hat{u} = 0$, the statement is obvious. We focus on $\hat{u} \neq 0$. For an arbitrary $i \in \mathbb{I}_N$, define

$$\hat{u}^{(i)} \stackrel{\text{def}}{=} (\hat{u}[1], \dots, \hat{u}[i-1], 0, \hat{u}[i+1], \dots, \hat{u}[N]) \in \mathbb{R}^N .$$

We shall use the equivalent formulation of \mathcal{F}_d given in (3). Clearly²⁵,

$$\mathcal{F}_d(\hat{u}) = \mathcal{F}_d(\hat{u}^{(i)} + e_i \hat{u}[i]) = \|A\hat{u}^{(i)} + a_i \hat{u}[i] - d\|^2 + \beta \sum_{j \in \mathbb{I}_N} \phi(\hat{u}^{(i)}[j]) + \phi(\hat{u}[i]) .$$

Consider $f_i : \mathbb{R} \rightarrow \mathbb{R}$ as given below

$$f_i(t) \stackrel{\text{def}}{=} \mathcal{F}_d(\hat{u}^{(i)} + e_i t) . \quad (69)$$

Since \hat{u} is a global minimizer of \mathcal{F}_d , for any $i \in \mathbb{I}_N$, we have

$$\begin{aligned} f_i(\hat{u}[i]) &= \mathcal{F}_d(\hat{u}^{(i)} + e_i \hat{u}[i]) \\ &\leq \mathcal{F}_d(\hat{u}^{(i)} + e_i t) = f_i(t), \quad \forall t \in \mathbb{R} . \end{aligned}$$

Equivalently, for any $i \in \mathbb{I}_N$, $f_i(\hat{u}[i])$ is the global minimum of $f_i(t)$ on \mathbb{R} . Below we will determine the global minimizer(s) $\hat{t} = \hat{u}[i]$ of f_i as given in (69), i. e.

$$\hat{t} = \hat{u}[i] = \arg \min_{t \in \mathbb{R}} f_i(t) .$$

Let us focus on an arbitrary $i \in \mathbb{I}_N$. In detail, the function f_i reads

$$\begin{aligned} f_i(t) &= \|A\hat{u}^{(i)} + a_i t - d\|^2 + \beta \sum_{j \in \mathbb{I}_N} \phi(\hat{u}^{(i)}[j]) + \beta \phi(t) \\ &= \|A\hat{u}^{(i)} - d\|^2 + \|a_i\|^2 t^2 + 2t \langle a_i, A\hat{u}^{(i)} - d \rangle + \beta \sum_{j \in \mathbb{I}_N} \phi(\hat{u}^{(i)}[j]) + \beta \phi(t) \\ &= \|a_i\|^2 t^2 + 2t \langle a_i, A\hat{u}^{(i)} - d \rangle + \beta \phi(t) + C_i , \end{aligned} \quad (70)$$

where

$$C_i = \|A\hat{u}^{(i)} - d\|^2 + \beta \sum_{j \in \mathbb{I}_N} \phi(\hat{u}^{(i)}[j]) .$$

Note that C_i does not depend on t . The function f_i has two local minimizers denoted by \hat{t}_0 and \hat{t}_1 . The first one is

$$\hat{t}_0 = 0 \quad \Rightarrow \quad f_i(\hat{t}_0) = C_i . \quad (71)$$

The other one, $\hat{t}_1 \neq 0$, corresponds to $\phi(t) = 1$. From (70), \hat{t}_1 solves

$$2\|a_i\|^2 t + 2\langle a_i, A\hat{u}^{(i)} - d \rangle = 0 .$$

Recalling that $a_i \neq 0, \forall i \in \mathbb{I}_N$ —see (9)—it follows that

$$\hat{t}_1 = -\frac{\langle a_i, A\hat{u}^{(i)} - d \rangle}{\|a_i\|^2} \quad \Rightarrow \quad f_i(\hat{t}_1) = -\frac{\langle a_i, A\hat{u}^{(i)} - d \rangle^2}{\|a_i\|^2} + \beta + C_i . \quad (72)$$

²⁵Using the definition of $\hat{u}^{(i)}$, we have $\hat{u}^{(i)} = A_{(\mathbb{I}_N \setminus \{i\})} \hat{u}_{(\mathbb{I}_N \setminus \{i\})}$, hence $A\hat{u}^{(i)}$ is independent of $\hat{u}[i]$.

Next we check if \hat{t}_0 or \hat{t}_1 is a global minimizer of f_i . From (71) and (72) we get

$$f_i(\hat{t}_0) - f_i(\hat{t}_1) = \frac{\langle a_i, A\hat{u}^{(i)} - d \rangle^2}{\|a_i\|^2} - \beta .$$

Furthermore,

$$\begin{aligned} f_i(\hat{t}_0) < f_i(\hat{t}_1) &\Rightarrow \hat{u}[i] = \hat{t}_0 = 0 , \\ f_i(\hat{t}_1) < f_i(\hat{t}_0) &\Rightarrow \hat{u}[i] = \hat{t}_1 = -\frac{\langle a_i, A\hat{u}^{(i)} - d \rangle}{\|a_i\|^2} , \\ f_i(\hat{t}_0) = f_i(\hat{t}_1) &\Rightarrow \hat{t}_0 \text{ and } \hat{t}_1 \text{ are global minimizers of } f_i . \end{aligned} \tag{73}$$

In particular, we have

$$\begin{aligned} f_i(\hat{t}_1) \leq f_i(\hat{t}_0) &\Leftrightarrow \langle a_i, A\hat{u}^{(i)} - d \rangle^2 \geq \beta \|a_i\|^2 \\ \text{[by (73)]} &\Rightarrow |\hat{u}[i]| = \frac{|\langle a_i, A\hat{u}^{(i)} - d \rangle|}{\|a_i\|^2} \\ \text{[by (74)]} &\geq \frac{\sqrt{\beta} \|a_i\|}{\|a_i\|^2} = \frac{\sqrt{\beta}}{\|a_i\|} . \end{aligned} \tag{74}$$

It is clear that the conclusion holds true for any $i \in \mathbb{I}_N$.

8.3 Proof of Proposition 5, p. 16

The asymptotic function $(\mathcal{F}_d)_\infty(v)$ of \mathcal{F}_d can be calculated according to²⁶ [3, Theorem 2.5.1]

$$(\mathcal{F}_d)_\infty(v) = \liminf_{\substack{v' \rightarrow v \\ t \rightarrow \infty}} \frac{\mathcal{F}_d(tv')}{t} .$$

Then

$$\begin{aligned} (\mathcal{F}_d)_\infty(v) &= \liminf_{\substack{v' \rightarrow v \\ t \rightarrow \infty}} \frac{\|Av' - d\|^2 + \beta \|v'\|_0}{t} \\ &= \liminf_{\substack{v' \rightarrow v \\ t \rightarrow \infty}} \left(t \|Av'\|^2 - 2\langle d, Av' \rangle + \frac{\|d\|^2 + \beta \|v'\|_0}{t} \right) \\ &= \begin{cases} 0 & \text{if } v \in \ker(A) , \\ +\infty & \text{if } v \notin \ker(A) . \end{cases} \end{aligned}$$

Hence

$$\ker((\mathcal{F}_d)_\infty) = \ker(A) , \tag{75}$$

where $\ker((\mathcal{F}_d)_\infty) = \{v \in \mathbb{R}^N : (\mathcal{F}_d)_\infty(v) = 0\}$.

Let $\{v_k\}$ satisfy (38) with $v_k \|v_k\|^{-1} \rightarrow \bar{v} \in \ker(A)$. Below we compare the numbers $\|v_k\|_0$ and $\|v_k - \rho \bar{v}\|_0$ where $\rho > 0$. There are two options.

²⁶In the nonconvex case, the notion of asymptotic function and the representation formula were first given by J.P. Dedieu [11].

1. Consider that $i \in \sigma(\bar{v})$, that is, $\bar{v}[i] = \lim_{k \rightarrow \infty} v_k[i] \|v_k\|^{-1} \neq 0$. Then $|v_k[i]| > 0$ for all but finitely many k as otherwise, $v_k[i] \|v_k\|^{-1}$ would converge to 0. Therefore, there exists k_i such that

$$|v_k[i] - \rho \bar{v}[i]| \geq 0 \quad \text{and} \quad |v_k[i]| > 0, \quad \forall k \geq k_i. \quad (76)$$

2. If $i \in (\sigma(\bar{v}))^c$, i.e. $\bar{v}[i] = 0$, then clearly

$$v_k[i] - \rho \bar{v}[i] = v_k[i]. \quad (77)$$

Combining (76) and (77), the definition of $\|\cdot\|_0$ using ϕ in (2) (p. 2) shows that

$$\|v_k - \rho \bar{v}\|_0 \leq \|v_k\|_0, \quad \forall k \geq k_0 \stackrel{\text{def}}{=} \max_{i \in \sigma(\bar{v})} k_i. \quad (78)$$

By (75), $A\bar{v} = 0$, which jointly with (78) entails that

$$\begin{aligned} \mathcal{F}_d(v_k - \rho \bar{v}) &= \|A(v_k + \rho \bar{v}) - d\|^2 + \beta \|v_k - \rho \bar{v}\|_0 \\ &= \|Av_k - d\|^2 + \beta \|v_k - \rho \bar{v}\|_0 \\ &\leq \|Av_k - d\|^2 + \beta \|v_k\|_0 = \mathcal{F}_d(v_k) \quad \forall k \geq k_0. \end{aligned}$$

It follows that for any $k \geq k_0$ we have

$$v_k \in \text{lev}(\mathcal{F}_d, \lambda_k) \quad \Rightarrow \quad v_k - \rho \bar{v} \in \text{lev}(\mathcal{F}_d, \lambda_k)$$

and thus \mathcal{F}_d satisfies Definition 5.

8.4 Proof of Proposition 6, p. 18

Given $K \in \mathbb{I}_{M-1}$, set

$$U_{K+1} \stackrel{\text{def}}{=} \bigcup_{\omega \subset \mathbb{I}_N} \{\bar{u} : \bar{u} \text{ solves } (\mathcal{P}_\omega) \text{ and } \|\bar{u}\|_0 \geq K+1\}. \quad (79)$$

- Let $U_{K+1} \neq \emptyset$. By Proposition 1 (p. 8), for any $\beta > 0$, \mathcal{F}_d has a (local) minimum at each $\bar{u} \in U_{K+1}$. Thus

$$\bar{u} \text{ is a (local) minimizer of } \mathcal{F}_d \text{ and } \|\bar{u}\|_0 \geq K+1 \quad \Leftrightarrow \quad \bar{u} \in U_{K+1}. \quad (80)$$

Then for any $\beta > 0$

$$\mathcal{F}_d(\bar{u}) \geq \beta(K+1), \quad \forall \bar{u} \in U_{K+1}. \quad (81)$$

Let \tilde{u} be defined by²⁷:

$$\tilde{u} \text{ solves } (\mathcal{P}_\omega) \text{ for some } \omega \in \Omega_K.$$

Then

$$\|\tilde{u}\|_0 \leq K. \quad (82)$$

Set β and β_K according to

$$\beta > \beta_K \stackrel{\text{def}}{=} \|A\tilde{u} - d\|^2. \quad (83)$$

²⁷Such a \tilde{u} always exists as explained in subsection 2.1. By Proposition 1 and Theorem 1, it is uniquely defined.

For such a β we have

$$\begin{aligned} \mathcal{F}_d(\tilde{u}) &= \|A\tilde{u} - d\|^2 + \beta\|\tilde{u}\|_0 \\ \left[\text{by (82) and (83)} \right] &< \beta + \beta\mathbf{K} = \beta(\mathbf{K} + 1) \\ \left[\text{by (81)} \right] &\leq \mathcal{F}_d(\bar{u}), \quad \forall \bar{u} \in \mathbf{U}_{\mathbf{K}+1}. \end{aligned}$$

Let \hat{u} be a global minimizer of \mathcal{F}_d . Then

$$\mathcal{F}_d(\hat{u}) \leq \mathcal{F}_d(\tilde{u}) < \mathcal{F}_d(\bar{u}), \quad \forall \bar{u} \in \mathbf{U}_{\mathbf{K}+1}.$$

Using (79)-(80), we find

$$\|\hat{u}\|_0 \leq \mathbf{K}.$$

- $\mathbf{U}_{\mathbf{K}+1} = \emptyset$ entails that²⁸

$$\bar{u} \text{ solves } (\mathcal{P}_\omega) \text{ for } \omega \subset \mathbb{I}_N, \quad \#\omega \geq \mathbf{K} + 1 \quad \Rightarrow \quad \|\bar{u}\|_0 \leq \mathbf{K}. \quad (84)$$

Let \hat{u} be a global minimizer of \mathcal{F}_d . By (84) we have

$$\|\hat{u}\|_0 \leq \mathbf{K}.$$

According to Theorem 2(ii) (p. 16), any global minimizer of \mathcal{F}_d is strict, hence $\sigma(\hat{u}) \in \bar{\Omega}_{\mathbf{K}}$.

References

- [1] H. ATTOUCH, J. BOLTE, AND F. SVAITER, *Convergence of descent methods for semi-algebraic and tame problems: proximal algorithms, forwardbackward splitting, and regularized gaussseidel methods*, Mathematical Programming, (2011).
- [2] A. AUSLENDER, *Existence of optimal solutions and duality results under weak conditions*, Mathematical Programming, 88, Serie A (2000), pp 45–59.
- [3] A. AUSLENDER AND M. TEBOULLE, *Asymptotic Cones and Functions in Optimization and Variational Inequalities*, Springer, New York, 2003.
- [4] C. BAIOCCHI, G. BUTTAZO, F. GASTALDI, AND F. TOMARELLI, *General existence theorems for unilateral problems in continuum mechanics*, Arch. Rational Mech. Anal., (1998), pp. 149–189.
- [5] J. E. BESAG, *On the statistical analysis of dirty pictures (with discussion)*, Journal of the Royal Statistical Society B, 48 (1986), pp. 259–302.
- [6] ———, *Digital image processing : Towards Bayesian image analysis*, Journal of Applied Statistics, 16 (1989), pp. 395–407.
- [7] T. BLUMENSATH AND M. DAVIES, *Iterative thresholding for sparse approximations*, Journal of Fourier Analysis and Applications, 14 (2008), pp. 629–654.

²⁸Let $A \in \mathbb{R}^{4 \times 5}$ read $A = (e_1, e_2, e_3, e_4, e_1)$ and $d = e_1 \in \mathbb{R}^4$. For $\mathbf{K} = \mathbf{M} - 1 = 3$ one can check that $\mathbf{U}_{\mathbf{K}+1} = \emptyset$.

- [8] A. M. BRUCKSTEIN, D. L. DONOHO, AND M. ELAD, *From sparse solutions of systems of equations to sparse modeling of signals and images*, SIAM Review, 51 (2009), pp. 34–81.
- [9] E. CHOZENOUX, A. JEZIERKA, J.-C. PESQUET, AND H. TALBOT, *A majorize-minimize subspace approach for $\ell_2 - \ell_0$ image regularization*, tech. report, arXiv:1112.6272, 2011.
- [10] G. DAVIS, S. MALLAT, AND M. AVELLANEDA, *Adaptive greedy approximations*, Constructive approximation, 13 (1997), pp. 57–98.
- [11] J. DEDIEU, *Cône asymptotique d'un ensemble non convexe. Application à l'optimisation.*, Comptes-rendus de l'académie des sciences, 287 (1977), pp. 91–103.
- [12] G. DEMOMENT, *Image reconstruction and restoration : Overview of common estimation structure and problems*, IEEE Transactions on Acoustics Speech and Signal Processing, ASSP-37 (1989), pp. 2024–2036.
- [13] B. DONG AND Y. ZHANG, *An efficient algorithm for ℓ_0 minimization in wavelet frame based image restoration*, Journal of Scientific Computing, (online) (2012).
- [14] D. L. DONOHO AND I. M. JOHNSTONE, *Ideal spatial adaptation by wavelet shrinkage*, Biometrika, 81 (1994), pp. 425–455.
- [15] L. C. EVANS AND R. F. GARIEPY, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics, CRC Press, Boca Baton, FL, 1992.
- [16] J. FAN AND R. LI, *Statistical challenges with high dimensionality: feature selection in knowledge discovery*, in Proceedings of the International Congress of Mathematicians, Madrid, Spain, vol. 3, European Mathematical Society, 2006, pp. 595–622.
- [17] M. FORNASIER AND R. WARD, *Iterative thresholding meets free-discontinuity problems*, Foundations of Computational Mathematics, 10 (2010), pp. 527–567.
- [18] G. GASSO, A. RAKOTOMAMONJY, AND S. CANU, *Recovering sparse signals with a certain family of non-convex penalties and DC programming*, IEEE Transactions on Signal Processing, 57 (2009), pp. 4686–4698.
- [19] S. GEMAN AND D. GEMAN, *Stochastic relaxation, Gibbs distributions, and the Bayesian restoration of images*, IEEE Transactions on Pattern Analysis and Machine Intelligence, PAMI-6 (1984), pp. 721–741.
- [20] G. GOLUB AND C. VAN LOAN, *Matrix Computations*, The Johns Hopkins University Press, Baltimore and London, 3 ed., 1996.
- [21] J. HAUPT AND R. NOWAK, *Signal reconstruction from noisy random projections*, IEEE Transactions on Information Theory, 52 (2006), pp. 4036–4048.
- [22] Y. LECLERC, *Constructing simple stable descriptions for image partitioning*, International Journal of Computer Vision, 3 (1989), pp. 73–102.
- [23] S. LI, *Markov Random Field Modeling in Computer Vision*, Springer-Verlag, London, UK, 1 ed., 1995.

- [24] Y. LIU AND Y. WU, *Variable selection via a combination of the l_0 and l_1 penalties*, Journal of Computational and Graphical Statistics, 16 (2007), pp. 782–798.
- [25] Z. LU AND Y. ZHANG, *Sparse approximation via penalty decomposition methods* (revised), arXiv:1205.2334 (2012).
- [26] J. LV AND Y. FAN, *A unified approach to model selection and sparse recovery using regularized least squares*, The Annals of Statistics, 37 (2009), pp. 3498–3528.
- [27] S. MALLAT, *A Wavelet Tour of Signal Processing (The sparse way)*, Academic Press, London, 3 ed., 2008.
- [28] C. D. MEYER, *Matrix Analysis and Applied Linear Algebra*, SIAM, 2000.
- [29] A. J. MILLER, *Subset Selection in Regression*, Chapman and Hall, London, U.K., 2 ed., 2002.
- [30] J. NEUMANN, C. SCHÖRR, AND G. STEIDL, *Combined SVM-based Feature Selection and classification*, Machine Learning 61, 2005, pp. 129–150.
- [31] M. NIKOLOVA, *Analysis of the recovery of edges in images and signals by minimizing non-convex regularized least-squares*, SIAM Journal on Multiscale Modeling and Simulation, 4 (2005), pp. 960–991.
- [32] M. ROBINI, A. LACHAL, AND I. MAGNIN, *A stochastic continuation approach to piecewise constant reconstruction*, IEEE Transactions on Image Processing, 16 (2007), pp. 2576–2589.
- [33] M. C. ROBINI AND I. E. MAGNIN, *Optimization by stochastic continuation*, SIAM Journal on Imaging Sciences, 3 (2010), pp. 1096–1121.
- [34] M. C. ROBINI AND P.-J. REISSMAN, *From simulated annealing to stochastic continuation: a new trend in combinatorial optimization*, J. of Global Optimization (to appear), (2012).
- [35] W. RUDIN, *Principles of Mathematical analysis*, Mathematics Series, McGraw-Hill, Berlin, 1976.
- [36] ———, *Real and Complex Analysis*, Mathematics Series, McGraw-Hill, Berlin, 1987.
- [37] E. M. STEIN AND R. SHAKARCHI, *Real Analysis: Measure Theory, Integration, and Hilbert Spaces*, Princeton University Press, 2005.
- [38] M. THIAO, T. P. DINH, AND A. L. THI, *DC Programming Approach for a Class of Nonconvex Programs Involving l_0 Norm*, Communications in Computer and Information Science, vol. 14, Springer, 2008, pp. 348–357.
- [39] J. TROPP, *Just relax: convex programming methods for identifying sparse signals in noise*, IEEE Transactions on Information Theory, 52 (2006), pp. 1030–1051.
- [40] URL, *Compressive Sensing Resources - Most Recent Postings*, http://dsp.rice.edu/cs/ref_chron, 2012.
- [41] M. VETTERLI AND KOVAČEVIĆ, *Wavelets and subband coding*, Prentice Hall PTR, 1995.
- [42] Y. ZHANG, B. DONG, AND Z. LU, *l_0 minimization of wavelet frame based image restoration*, Mathematics of Computation (to appear), (2012).