



**HAL**  
open science

# Perfect forms, K-theory and the cohomology of modular groups

Philippe Elbaz-Vincent, Herbert Gangl, Christophe Soulé

► **To cite this version:**

Philippe Elbaz-Vincent, Herbert Gangl, Christophe Soulé. Perfect forms, K-theory and the cohomology of modular groups. 2012. hal-00724128

**HAL Id: hal-00724128**

**<https://hal.science/hal-00724128v1>**

Preprint submitted on 17 Aug 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# PERFECT FORMS, K-THEORY AND THE COHOMOLOGY OF MODULAR GROUPS

PHILIPPE ELBAZ-VINCENT, HERBERT GANGL, AND CHRISTOPHE SOULÉ

## CONTENTS

1. Introduction	1
2. Voronoï's reduction theory	4
3. The Voronoï complex	5
4. The Voronoï complex in dimensions 5, 6 and 7	7
5. Explicit homology classes	12
6. Splitting off the Voronoï complex $\text{Vor}_N$ from $\text{Vor}_{N+1}$ for small $N$	14
7. The Cohomology of modular groups	16
8. Homology of modular groups with coefficients in the Steinberg module	18
9. Application to K-theory	32
References	35

ABSTRACT. For  $N = 5, 6$  and  $7$ , using the classification of perfect quadratic forms, we compute the homology of the Voronoï cell complexes attached to the modular groups  $SL_N(\mathbb{Z})$  and  $GL_N(\mathbb{Z})$ . From this we deduce the rational cohomology of those groups and some information about  $K_m(\mathbb{Z})$ , when  $m = 5, 6$  and  $7$ .

## 1. INTRODUCTION

Let  $N \geq 1$  be an integer and let  $SL_N(\mathbb{Z})$  be the modular group of integral matrices with determinant one. Our goal is to compute its cohomology groups with trivial coefficients, i.e.  $H^q(SL_N(\mathbb{Z}), \mathbb{Z})$ . The case  $N = 2$  is well-known and follows from the fact that  $SL_2(\mathbb{Z})$  is the amalgamated product of two finite cyclic groups ([29], [7], II.7, Ex.3, p.52). The case  $N = 3$  was done in [31]: for any  $q > 0$  the group  $H^q(SL_3(\mathbb{Z}), \mathbb{Z})$  is killed by 12. The case  $N = 4$  has been studied by Lee and Szczarba in [19]: modulo 2, 3 and 5-torsion, the cohomology group  $H^q(SL_4(\mathbb{Z}), \mathbb{Z})$  is trivial whenever  $q > 0$ , except that  $H^3(SL_4(\mathbb{Z}), \mathbb{Z}) = \mathbb{Z}$ . In Theorem 7.3 below, we solve the cases  $N = 5, 6$  and  $7$ .

For these calculations we follow the method of [19], i.e. we use the perfect forms of Voronoï. Recall from [34] and [20] that a perfect form in  $N$  variables is a positive definite real quadratic form  $h$  on  $\mathbb{R}^N$  which is uniquely determined (up to a scalar) by its set of integral minimal vectors. Voronoï proved in [34] that there are finitely many perfect forms of rank  $N$ , modulo the action of  $SL_N(\mathbb{Z})$ . These are known for  $N \leq 8$  (see §2 below).

---

1991 *Mathematics Subject Classification.* 11H55, 11F75, 11F06, 11Y99, 55N91, 19D50, 20J06, 57-04.

*Key words and phrases.* Perfect forms, Voronoï complex, group cohomology, modular groups, Steinberg modules, K-theory of integers, machine calculations.

Voronoi used perfect forms to define a cell decomposition of the space  $X_N^*$  of positive real quadratic forms, the kernel of which is defined over  $\mathbb{Q}$ . This cell decomposition (cf. §3) is invariant under  $SL_N(\mathbb{Z})$ , hence it can be used to compute the equivariant homology of  $X_N^*$  modulo its boundary. On the other hand, this equivariant homology turns out to be isomorphic to the groups  $H_q(SL_N(\mathbb{Z}), St_N)$ , where  $St_N$  is the Steinberg module (see [6] and §3.4 below). Finally, Borel–Serre duality [6] asserts that the homology  $H_*(SL_N(\mathbb{Z}), St_N)$  is dual to the cohomology  $H^*(SL_N(\mathbb{Z}), \mathbb{Z})$  (modulo torsion).

To perform these computations for  $N \leq 7$ , we needed the help of a computer. The reason is that the Voronoi cell decomposition of  $X_N^*$  gets soon very complicated when  $N$  increases. For instance, when  $N = 7$ , there are more than two million orbits of cells of dimension 18, modulo the action of  $SL_N(\mathbb{Z})$  (see Figure 2 below). For this purpose, we have developed a C library [23], which uses PARI [22] for some functionalities. The algorithms are based on exact methods. As a result we get the full Voronoi cell decomposition of the spaces  $X_N^*$  for  $N \leq 7$  (with either  $GL_N(\mathbb{Z})$  or  $SL_N(\mathbb{Z})$  action). Those decompositions are summarized in the figures and tables below. The computations were done on several computers using different processor architectures (which is useful for checking the results) and for  $N = 7$  the overall computational time was more than a year.

The paper is organized as follows. In §2, we recall the Voronoi theory of perfect forms. In §3, we introduce a complex of abelian groups that we call the “Voronoi complex” which computes the homology groups  $H_q(SL_N(\mathbb{Z}), St_N)$ . In §4, we explain how to get an explicit description of the Voronoi complex in rank  $N = 5, 6$  or  $7$ , starting from the description of perfect forms available in the literature (especially in the work of Jaquet [15]). In Figures 1 and 2 we display the rank of the groups in the Voronoi complex and in Tables 1–5 we give the elementary divisors of its differentials. The homology of the Voronoi complex (hence the groups  $H_q(SL_N(\mathbb{Z}), St_N)$ ) follows from this. It is given in Theorem 4.3.

We found two methods to test whether our computations are correct. First, checking that the virtual Euler characteristic of  $SL_N(\mathbb{Z})$  vanishes leads to a mass formula for the orders of the stabilizers of the cells of  $X_N^*$  (cf. §4.5). Second, the identity  $d_{n-1} \circ d_n = 0$  for the differentials in the Voronoi complex is a non-trivial equality when these differentials are written as explicit (large) matrices.

In §5 we give an explicit formula for the top homology group of the Voronoi complex (Theorem 5.1). In §6 we prove that the Voronoi complex of  $GL_5(\mathbb{Z})$  is a direct factor of the Voronoi complex of  $GL_6(\mathbb{Z})$  shifted by one. In §7 we explain how to compute the cohomology of  $SL_N(\mathbb{Z})$  and  $GL_N(\mathbb{Z})$  (modulo torsion) from our results on the homology of the Voronoi complex in §4. Our main result is stated in Theorem 7.3. In §8 we compute some homology groups of  $GL_N(\mathbb{Z})$  with coefficients the Steinberg module. In §9, we use these results to get some information on  $K_m(\mathbb{Z})$ , when  $m = 5, 6$  and  $7$ . Some of these results had already been announced in [10].

**Acknowledgments:** The first two authors are particularly indebted to the IHES for its hospitality. The second author thanks the Institute for Experimental Mathematics (Essen), acknowledging financial support by the DFG and the European Commission as well as hospitality of the Newton Institute in Cambridge and the

MPI for Mathematics in Bonn. The authors are grateful to B. Allombert, J.-G. Dumas, M. Dutour, D.-O. Jaquet, J.-C. König, J. Martinet, S. Morita, A. Rahm, J.-P. Serre and B. Souvignier for helpful discussions and comments. The computations of the Voronoï cell decomposition were performed on the computers of the MPI, the Institut Fourier in Grenoble and the Centre de Calcul Médecis; we are grateful to those institutions.

*Notation:* For any positive integer  $n$  we let  $\mathcal{S}_n$  be the class of finite abelian groups the order of which has only prime factors less than or equal to  $n$ .

## 2. VORONOÏ'S REDUCTION THEORY

**2.1. Perfect forms.** Let  $N \geq 2$  be an integer. We let  $C_N$  be the set of positive definite real quadratic forms in  $N$  variables. Given  $h \in C_N$ , let  $m(h)$  be the finite set of minimal vectors of  $h$ , i.e. vectors  $v \in \mathbb{Z}^N$ ,  $v \neq 0$ , such that  $h(v)$  is minimal. A form  $h$  is called *perfect* when  $m(h)$  determines  $h$  up to scalar: if  $h' \in C_N$  is such that  $m(h') = m(h)$ , then  $h'$  is proportional to  $h$ .

**Example 2.1.** The form  $h(x, y) = x^2 + y^2$  has minimum 1 and precisely 4 minimal vectors  $\pm(1, 0)$  and  $\pm(0, 1)$ . This form is not perfect, because there is an infinite number of positive definite quadratic forms having these minimal vectors, namely the forms  $h(x, y) = x^2 + axy + y^2$  where  $a$  is a non-negative real number less than 1. By contrast, the form  $h(x, y) = x^2 + xy + y^2$  has also minimum 1 and has exactly 6 minimal vectors, viz. the ones above and  $\pm(1, -1)$ . This form is perfect, the associated lattice is the ‘‘honeycomb lattice’’.

Denote by  $C_N^*$  the set of non-negative real quadratic forms on  $\mathbb{R}^N$  the kernel of which is spanned by a proper linear subspace of  $\mathbb{Q}^N$ , by  $X_N^*$  the quotient of  $C_N^*$  by positive real homotheties, and by  $\pi : C_N^* \rightarrow X_N^*$  the projection. Let  $X_N = \pi(C_N)$  and  $\partial X_N^* = X_N^* - X_N$ . Let  $\Gamma$  be either  $GL_N(\mathbb{Z})$  or  $SL_N(\mathbb{Z})$ . The group  $\Gamma$  acts on  $C_N^*$  and  $X_N^*$  on the right by the formula

$$h \cdot \gamma = \gamma^t h \gamma, \quad \gamma \in \Gamma, h \in C_N^*,$$

where  $h$  is viewed as a symmetric matrix and  $\gamma^t$  is the transpose of the matrix  $\gamma$ . Voronoï proved that there are only finitely many perfect forms modulo the action of  $\Gamma$  and multiplication by positive real numbers ([34], Thm. p.110).

The following table gives the current state of the art on the enumeration of perfect forms.

rank	1	2	3	4	5	6	7	8	9
#classes	1	1	1	2	3	7	33	10916	$\geq 500000$

The classification of perfect forms of rank 8 was achieved by Dutour, Schürmann and Vallentin in 2005 [9], [28]. They have also shown that in rank 9 there are at least 500000 classes of perfect forms. The corresponding classification for rank 7 was completed by Jaquet in 1991 [15], for rank 6 by Barnes [2], for rank 5 and 4 by Korkine and Zolotarev [16], [17], for dimension 3 by Gauss [13] and for dimension 2 by Lagrange [18]. We refer to the book of Martinet [20] for more details on the results up to rank 7.

**2.2. A cell complex.** Given  $v \in \mathbb{Z}^N - \{0\}$  we let  $\hat{v} \in C_N^*$  be the form defined by

$$\hat{v}(x) = (v | x)^2, \quad x \in \mathbb{R}^N,$$

where  $(v | x)$  is the scalar product of  $v$  and  $x$ . The *convex hull in  $X_N^*$*  of a finite subset  $B \subset \mathbb{Z}^N - \{0\}$  is the subset of  $X_N^*$  which is the image under  $\pi$  of the quadratic forms  $\sum_j \lambda_j \hat{v}_j \in C_N^*$ , where  $v_j \in B$  and  $\lambda_j \geq 0$ . For any perfect form  $h$ , we let  $\sigma(h) \subset X_N^*$  be the convex hull of the set  $m(h)$  of its minimal vectors. Voronoï proved in [34], §§8-15, that the cells  $\sigma(h)$  and their intersections, as  $h$  runs over all perfect forms, define a cell decomposition of  $X_N^*$ , which is invariant under the action of  $\Gamma$ . We endow  $X_N^*$  with the corresponding CW-topology. If  $\tau$  is a closed cell in  $X_N^*$  and  $h$  a perfect form with  $\tau \subset \sigma(h)$ , we let  $m(\tau)$  be the set of vectors  $v$  in

$m(h)$  such that  $\hat{v}$  lies in  $\tau$ . Any closed cell  $\tau$  is the convex hull of  $m(\tau)$ , and for any two closed cells  $\tau, \tau'$  in  $X_N^*$  we have  $m(\tau) \cap m(\tau') = m(\tau \cap \tau')$ .

### 3. THE VORONOÏ COMPLEX

**3.1. An explicit differential for the Voronoi complex.** Let  $d(N) = N(N+1)/2 - 1$  be the dimension of  $X_N^*$  and  $n \leq d(N)$  a natural integer. We denote by  $\Sigma_n^* = \Sigma_n^*(\Gamma)$  a set of representatives, modulo the action of  $\Gamma$ , of those cells of dimension  $n$  in  $X_N^*$  which meet  $X_N$ , and by  $\Sigma_n = \Sigma_n(\Gamma) \subset \Sigma_n^*(\Gamma)$  the cells  $\sigma$  for which any element of the stabilizer  $\Gamma_\sigma$  of  $\sigma$  in  $\Gamma$  preserves the orientation. Let  $V_n$  be the free abelian group generated by  $\Sigma_n$ . We define as follows a map

$$d_n : V_n \rightarrow V_{n-1}.$$

For each closed cell  $\sigma$  in  $X_N^*$  we fix an orientation of  $\sigma$ , i.e. an orientation of the real vector space  $\mathbb{R}(\sigma)$  of symmetric matrices spanned by the forms  $\hat{v}$  with  $v \in m(\sigma)$ . Let  $\tau \in \Sigma_n$  and let  $\tau'$  be a face of  $\sigma$  which is equivalent under  $\Gamma$  to an element in  $\Sigma_{n-1}$  (i.e.  $\tau'$  neither lies on the boundary nor has elements in its stabilizer reversing the orientation). Given a positive basis  $B'$  of  $\mathbb{R}(\tau')$  we get a basis  $B$  of  $\mathbb{R}(\sigma) \supset \mathbb{R}(\tau')$  by appending to  $B'$  a vector  $\hat{v}$ , where  $v \in m(\sigma) - m(\tau')$ . We let  $\varepsilon(\tau', \sigma) = \pm 1$  be the sign of the orientation of  $B$  in the oriented vector space  $\mathbb{R}(\sigma)$  (this sign does not depend on the choice of  $v$ ).

Next, let  $\tau \in \Sigma_{n-1}$  be the (unique) cell equivalent to  $\tau'$  and let  $\gamma \in \Gamma$  be such that  $\tau' = \tau \cdot \gamma$ . We define  $\eta(\tau, \tau') = 1$  (resp.  $\eta(\tau, \tau') = -1$ ) when  $\gamma$  is compatible (resp. incompatible) with the chosen orientations of  $\mathbb{R}(\tau)$  and  $\mathbb{R}(\tau')$ .

Finally we define

$$(1) \quad d_n(\sigma) = \sum_{\tau \in \Sigma_{n-1}} \sum_{\tau'} \eta(\tau, \tau') \varepsilon(\tau', \sigma) \tau,$$

where  $\tau'$  runs through the set of faces of  $\sigma$  which are equivalent to  $\tau$ .

**3.2. A spectral sequence.** According to [7], VII.7, there is a spectral sequence  $E_{pq}^r$  converging to the equivariant homology groups  $H_{p+q}^\Gamma(X_N^*, \partial X_N^*; \mathbb{Z})$  of the homology pair  $(X_N^*, \partial X_N^*)$ , and such that

$$E_{pq}^1 = \bigoplus_{\sigma \in \Sigma_p^*} H_q(\Gamma_\sigma, \mathbb{Z}_\sigma),$$

where  $\mathbb{Z}_\sigma$  is the orientation module of the cell  $\sigma$  and, as above,  $\Sigma_p^*$  is a set of representatives, modulo  $\Gamma$ , of the  $p$ -cells  $\sigma$  in  $X_N^*$  which meet  $X_N$ . Notice that the action of  $\Gamma_\sigma$  on  $\mathbb{Z}_\sigma$  is given by  $\eta$  described above. Since  $\sigma$  meets  $X_N$ , its stabilizer  $\Gamma_\sigma$  is finite and, by Lemma 7.1 in §7 below, the order of  $\Gamma_\sigma$  is divisible only by primes  $p \leq N+1$ . Therefore, when  $q$  is positive, the group  $H_q(\Gamma_\sigma, \mathbb{Z}_\sigma)$  lies in  $\mathcal{S}_{N+1}$ .

When  $\Gamma_\sigma$  happens to contain an element which changes the orientation of  $\sigma$ , the group  $H_0(\Gamma_\sigma, \mathbb{Z}_\sigma)$  is killed by 2, otherwise  $H_0(\Gamma_\sigma, \mathbb{Z}_\sigma) \cong \mathbb{Z}_\sigma$ . Therefore, modulo  $\mathcal{S}_2$ , we have

$$E_{n0}^1 = \bigoplus_{\sigma \in \Sigma_n} \mathbb{Z}_\sigma,$$

and the choice of an orientation for each cell  $\sigma$  gives an isomorphism between  $E_{n0}^1$  and  $V_n$ .

**3.3. Comparison.** We claim that the differential

$$d_n^1 : E_{n0}^1 \rightarrow E_{n-1,0}^1$$

coincides, up to sign, with the map  $d_n$  defined in 3.1. According to [7], VII, Prop. (8.1), the differential  $d_n^1$  can be described as follows.

Let  $\sigma \in \Sigma_n^*$  and let  $\tau'$  be a face of  $\sigma$ . Consider the group  $\Gamma_{\sigma\tau'} = \Gamma_\sigma \cap \Gamma_{\tau'}$  and denote by

$$t_{\sigma\tau'} : H_*(\Gamma_\sigma, \mathbb{Z}_\sigma) \rightarrow H_*(\Gamma_{\sigma\tau'}, \mathbb{Z}_\sigma)$$

the transfer map. Next, let

$$u_{\sigma\tau'} : H_*(\Gamma_{\sigma\tau'}, \mathbb{Z}_\sigma) \rightarrow H_*(\Gamma_{\tau'}, \mathbb{Z}_{\tau'})$$

be the map induced by the natural map  $\mathbb{Z}_\sigma \rightarrow \mathbb{Z}_{\tau'}$ , together with the inclusion  $\Gamma_{\sigma\tau'} \subset \Gamma_{\tau'}$ . Finally, let  $\tau \in \Sigma_{n-1}^*$  be the representative of the  $\Gamma$ -orbit of  $\tau'$ , let  $\gamma \in \Gamma$  be such that  $\tau' = \tau \cdot \gamma$ , and let

$$v_{\tau'\tau} : H_*(\Gamma_{\tau'}, \mathbb{Z}_{\tau'}) \rightarrow H_*(\Gamma_\tau, \mathbb{Z}_\tau)$$

be the isomorphism induced by  $\gamma$ . Then the restriction of  $d_n^1$  to  $H_*(\Gamma_\sigma, \mathbb{Z}_\sigma)$  is equal, up to sign, to the sum

$$(2) \quad \sum_{\tau'} v_{\tau'\tau} u_{\sigma\tau'} t_{\sigma\tau'},$$

where  $\tau'$  runs over a set of representatives of faces of  $\sigma$  modulo  $\Gamma_\sigma$ .

To compare  $d_n^1$  with  $d_n$  we first note that, when  $\tau \in \Sigma_{n-1}$ ,

$$v_{\tau'\tau} : H_0(\Gamma_{\tau'}, \mathbb{Z}_{\tau'}) = \mathbb{Z} \rightarrow H_0(\Gamma_\tau, \mathbb{Z}_\tau) = \mathbb{Z}$$

is the multiplication by  $\eta(\tau, \tau')$ , as defined in §3.1. Next, when  $\sigma \in \Sigma_n$ , the map

$$u_{\sigma\tau'} : H_0(\Gamma_{\sigma\tau'}, \mathbb{Z}_\sigma) = \mathbb{Z}_\sigma = \mathbb{Z} \rightarrow H_0(\Gamma_{\tau'}, \mathbb{Z}_{\tau'}) = \mathbb{Z}$$

is the multiplication by  $\varepsilon(\tau', \sigma)$ , up to a sign depending on  $n$  only. Finally, the transfer map

$$t_{\sigma\tau'} : H_0(\Gamma_\sigma, \mathbb{Z}_\sigma) = \mathbb{Z} \rightarrow H_0(\Gamma_{\sigma\tau'}, \mathbb{Z}_\sigma) = \mathbb{Z}$$

is the multiplication by  $[\Gamma_\sigma : \Gamma_{\sigma\tau'}]$ . Multiplying the sum (2) by this number amounts to the same as taking the sum over all faces of  $\sigma$  as in (1). This proves that  $d_n$  coincides, up to sign, with  $d_n^1$  on  $E_{n0}^1 = V_n$ .  $\square$

In particular, we get that  $d_{n-1} \circ d_n = 0$ . Note that this identity will give us a non-trivial test of our explicit computations of the complex.

*Notation:* The resulting complex  $(V_\bullet, d_\bullet)$  will be denoted by  $\text{Vor}_\Gamma$ , and we call it the *Voronoi complex*.

**3.4. The Steinberg module.** Let  $T_N$  be the spherical Tits building of  $SL_N$  over  $\mathbb{Q}$ , i.e. the simplicial set defined by the ordered set of non-zero proper linear subspaces of  $\mathbb{Q}^N$ . The reduced homology  $\tilde{H}_q(T_N, \mathbb{Z})$  of  $T_N$  with integral coefficients is zero except when  $q = N - 2$ , in which case

$$\tilde{H}_{N-2}(T_N, \mathbb{Z}) = \text{St}_N$$

is by definition the Steinberg module [6]. According to [30], Prop. 1, the relative homology groups  $H_q(X_N^*, \partial X_N^*; \mathbb{Z})$  are zero except when  $q = N - 1$ , and

$$H_{N-1}(X_N^*, \partial X_N^*; \mathbb{Z}) = \text{St}_N.$$

From this it follows that, for all  $m \in \mathbb{N}$ ,

$$H_m^\Gamma(X_N^*, \partial X_N^*; \mathbb{Z}) = H_{m-N+1}(\Gamma, \text{St}_N)$$

(see e.g. [30], §3.1). Combining this equality with the previous sections we conclude that, modulo  $\mathcal{S}_{N+1}$ ,

$$(3) \quad H_{m-N+1}(\Gamma, \text{St}_N) = H_m(\text{Vor}_\Gamma).$$

#### 4. THE VORONOÏ COMPLEX IN DIMENSIONS 5, 6 AND 7

In this section, we explain how to compute the Voronoï complexes of rank  $N \leq 7$ .

**4.1. Checking the equivalence of cells.** As a preliminary step, we develop an effective method to check whether two cells  $\sigma$  and  $\sigma'$  of the same dimension are equivalent under the action of  $\Gamma$ . The cell  $\sigma$  (resp.  $\sigma'$ ) is described by its set of minimal vectors  $m(\sigma)$  (resp.  $m(\sigma')$ ). We let  $b$  (resp.  $b'$ ) be the sum of the forms  $\hat{v}$  with  $v \in m(\sigma)$  (resp.  $m(\sigma')$ ). If  $\sigma$  and  $\sigma'$  are equivalent under the action of  $\Gamma$  the same is true for  $b$  and  $b'$ , and the converse holds true since two cells of the same dimension are equal when they have an interior point in common.

To compare  $b$  and  $b'$  we first check whether or not they have the same determinant. In case they do, we let  $M$  (resp.  $M'$ ) be the set of numbers  $b(x)$  with  $x \in m(\sigma)$  (resp.  $b'(x)$  with  $x \in m(\sigma')$ ). If  $b$  and  $b'$  are equivalent, then the sets  $M$  and  $M'$  must be equal.

Finally, if  $M = M'$  we check if  $b$  and  $b'$  are equivalent by applying an algorithm of Plesken and Souvignier [24] (based on an implementation of Souvignier).

**4.2. Finding generators of the Voronoï complex.** In order to compute  $\Sigma_n$  (and  $\Sigma_n^*$ ), we proceed as follows. Fix  $N \leq 7$ . Let  $\mathcal{P}$  be a set of representatives of the perfect forms of rank  $N$ . A choice of  $\mathcal{P}$  is provided by Jaquet [15]. Furthermore, for each  $h \in \mathcal{P}$ , Jaquet gives the list  $m(h)$  of its minimal vectors, and the list of all perfect forms  $h'\gamma$  (one for each orbit under  $\Gamma_{\sigma(h)}$ ), where  $h' \in \mathcal{P}$  and  $\gamma \in \Gamma$ , such that  $\sigma(h)$  and  $\sigma(h')\gamma$  share a face of codimension one. This provides a complete list  $C_h^1$  of representatives of codimension one faces in  $\sigma(h)$ .

From this, one deduces the full list  $\mathcal{F}_h^1$  of faces of codimension one in  $\sigma(h)$  as follows: first list all the elements in the automorphism group  $\Gamma_{\sigma(h)}$ ; this can be obtained by using a second procedure implemented by Souvignier [24] which gives generators for  $\Gamma_{\sigma(h)}$ . We represent the latter generators as elements in the symmetric group  $\mathfrak{S}_M$ , where  $M$  is the cardinality of  $m(h)$ , acting on the set  $m(h)$  of minimal vectors. Using those generators, we let GAP [12] list all the elements of  $\Gamma_{\sigma(h)}$ , viewed as elements of the symmetric group above.

The next step is to create a shortlist  $\mathcal{F}_h^2$  of codimension 2 facets of  $\sigma(h)$  by intersecting all the translates under  $\mathfrak{S}_M$  of codimension 1 facets with each member of  $C_h^1$  and only keeping those intersections with the correct rank ( $=d(N) - 2$ ). The resulting shortlist is reasonably small and we apply the procedure of 4.1 to reduce the shortlist to a set of representatives  $C_h^2$  of codimension 2 facets.

We then proceed by induction on the codimension to define a list  $\mathcal{F}_h^p$  of cells of codimension  $p > 2$  in  $\sigma(h)$ . Given  $\mathcal{F}_h^p$ , we let  $C_h^p \subset \mathcal{F}_h^p$  be a set of representatives for the action of  $\Gamma$ . We then let  $\mathcal{F}_h^{p+1}$  be the set of cells  $\varphi \cap \tau$ , with  $\varphi \in \mathcal{F}_h^p$ , and  $\tau \in C_h^p$ . As a result, we get directly the cellular structure of the quotient space  $(X_N^*, \partial X_N^*)/\Gamma$  without computing the full cellular structure of  $X_N^*$  which is not required (and of greater computational complexity).



n	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
$\Sigma_n^*(GL_5(\mathbb{Z}))$	2	5	10	16	23	25	23	16	9	4	3						
$\Sigma_n(GL_5(\mathbb{Z}))$					1	7	6	1	0	2	3						
$\Sigma_n^*(GL_6(\mathbb{Z}))$		3	10	28	71	162	329	589	874	1066	1039	775	425	181	57	18	7
$\Sigma_n(GL_6(\mathbb{Z}))$						3	46	163	340	544	636	469	200	49	5		
$\Sigma_n^*(SL_6(\mathbb{Z}))$		3	10	28	71	163	347	691	1152	1532	1551	1134	585	222	62	18	7
$\Sigma_n(SL_6(\mathbb{Z}))$			3	10	18	43	169	460	815	1132	1270	970	434	114	27	14	7

FIGURE 1. Cardinality of  $\Sigma_n$  and  $\Sigma_n^*$  for  $N = 5, 6$  (empty slots denote zero).

n	6	7	8	9	10	11	12	13	14	15	16
$\Sigma_n^*$	6	28	115	467	1882	7375	26885	87400	244029	569568	1089356
$\Sigma_n$				1	60	1019	8899	47271	171375	460261	955128

n	17	18	19	20	21	22	23	24	25	26	27
$\Sigma_n^*$	1683368	2075982	2017914	1523376	876385	374826	115411	24623	3518	352	33
$\Sigma_n$	1548650	1955309	1911130	1437547	822922	349443	105054	21074	2798	305	33

FIGURE 2. Cardinality of  $\Sigma_n$  and  $\Sigma_n^*$  for  $GL_7(\mathbb{Z})$ .

Next, we let  $\Sigma_n^*$  be a system of representatives modulo  $\Gamma$  in the union of the sets  $C_h^{d(N)-n}$ ,  $h \in \mathcal{P}$ . We then compute generators of the stabilizer of each cell in  $\Sigma_n^*$  with the help of another algorithm developed by Plesken and Souvignier in [24], and we check whether all generators preserve the orientation of the cell. This gives us the set  $\Sigma_n$  as the set of those cells which pass that check.

**Proposition 4.1.** *The cardinality of  $\Sigma_n$  and  $\Sigma_n^*$  is displayed in Figure 1 for rank  $N = 5, 6$  and in Figure 2 for rank  $N = 7$ .*

**Remark 4.2.** The first line in Figure 1 has already been computed by Batut (cf. [3], p.409, second column of Table 2). The running time for the computation of the cell structure (with the differentials and the checking) for  $N = 7$  using [23] was 18 months on several servers including quadri-processors computers, while for  $N = 6$  this can be done in a few seconds.

**4.3. The differential.** The next step is to compute the differentials of the Voronoï complex by using formula (1) above. In Table 3, we give information on the differentials in the Voronoï complex of rank 6. For instance the second line, denoted  $d_{11}$ , is about the differential from  $V_{11}$  to  $V_{10}$ . In the bases  $\Sigma_{11}$  and  $\Sigma_{10}$ , this differential is given by a matrix  $A$  with  $\Omega = 513$  non-zero entries, with  $m = 46 = \text{card}(\Sigma_{10})$  rows and  $n = 163 = \text{card}(\Sigma_{11})$  columns. The rank of  $A$  is 42, and the rank of its kernel is 121. The elementary divisors of  $A$  are 1 (multiplicity 40) and 2 (multiplicity 2).

$A$	$\Omega$	$n$	$m$	rank	ker	elementary divisors
$d_4$	0	1	0	0	1	
$d_5$	1	1	1	1	0	1(1)
$d_6$	0	1	1	0	1	
$d_7$	0	0	1	0	0	
$d_8$	0	1	0	0	1	
$d_9$	2	2	1	1	1	2(1)

TABLE 1. Results on the rank and elementary divisors of the differentials for  $SL_4(\mathbb{Z})$ .

$A$	$\Omega$	$n$	$m$	rank	ker	elementary divisors
$d_8$	0	1	0	0	1	
$d_9$	2	7	1	1	6	1(1)
$d_{10}$	18	6	7	5	1	1(4), 2(1)
$d_{11}$	5	1	6	1	0	1(1)
$d_{12}$	0	0	1	0	0	
$d_{13}$	0	2	0	0	2	
$d_{14}$	4	3	2	2	1	5(1), 15(1)

TABLE 2. Results on the rank and elementary divisors of the differentials for  $GL_5(\mathbb{Z})$ .

The cases of  $SL_4(\mathbb{Z})$ ,  $GL_5(\mathbb{Z})$  and  $SL_6(\mathbb{Z})$  are treated in Table 1, Table 2 and Table 4, respectively.

Our results on the differentials in rank 7 are shown in Table 5. While the matrices are sparse, they are not sparse enough for efficient computation. They have a poor conditioning with some dense columns or rows (this is a consequence of the fact that the complex is not simplicial and non-simplicial cells can have a large number of non-trivial intersections with the faces). We have obtained full information on the rank of the differentials. For the computation of the elementary divisors complete results have been obtained in the case of matrices of  $d_n$  except for  $n = 19$ . For this case, the computational cost is currently too high. The computations have required a full year on a parallel computer (including checking). For  $n = 19$  alone, the computational cost is equivalent to 3 CPU-years on a current processor. See [8, 33] for a detailed description of the computations.

**4.4. The homology of the Voronoï complexes.** From the computation of the differentials, we can determine the homology of Voronoï complex. Recall that if we have a complex of free abelian groups

$$\dots \rightarrow \mathbb{Z}^\alpha \xrightarrow{f} \mathbb{Z}^\beta \xrightarrow{g} \mathbb{Z}^\gamma \rightarrow \dots$$

with  $f$  and  $g$  represented by matrices, then the homology is

$$\ker(g)/\text{Im}(f) \cong \mathbb{Z}/d_1\mathbb{Z} \oplus \dots \oplus \mathbb{Z}/d_\ell\mathbb{Z} \oplus \mathbb{Z}^{\beta - \text{rank}(f) - \text{rank}(g)},$$

where  $d_1, \dots, d_\ell$  are the elementary divisors of the matrix of  $f$ .

We deduce from Tables 1–5 the following result on the homology of the Voronoï complex.

A	$\Omega$	$n$	$m$	rank	ker	elementary divisors
$d_{10}$	17	46	3	3	43	1(3)
$d_{11}$	513	163	46	42	121	1(40), 2(2)
$d_{12}$	2053	340	163	120	220	1(120)
$d_{13}$	4349	544	340	220	324	1(217), 2(3)
$d_{14}$	6153	636	544	324	312	1(320), 2(1), 6(2), 12(1)
$d_{15}$	5378	469	636	312	157	1(307), 2(3), 60(2)
$d_{16}$	2526	200	469	156	44	1(156)
$d_{17}$	597	49	200	44	5	1(41), 3(1), 6(1), 36(1)
$d_{18}$	43	5	49	5	0	1(5)

TABLE 3. Results on the rank and elementary divisors of the differentials for  $GL_6(\mathbb{Z})$ .

A	$\Omega$	$n$	$m$	rank	ker	elementary divisors
$d_7$	12	10	3	3	7	1(3)
$d_8$	48	18	10	7	11	1(7)
$d_9$	140	43	18	11	32	1(11)
$d_{10}$	613	169	43	32	137	1(32)
$d_{11}$	2952	460	169	136	324	1(129), 2(6), 6(1)
$d_{12}$	7614	815	460	323	492	1(318), 2(3), 4(2)
$d_{13}$	12395	1132	815	491	641	1(491)
$d_{14}$	14966	1270	1132	641	629	1(637), 3(3), 12(1)
$d_{15}$	12714	970	1270	629	341	1(621), 2(5), 6(1), 60(2)
$d_{16}$	6491	434	970	339	95	1(338), 2(1)
$d_{17}$	1832	114	434	95	19	1(92), 3(2), 18(1)
$d_{18}$	257	27	114	19	8	1(17), 2(2)
$d_{19}$	62	14	27	8	6	1(7), 10(1)
$d_{20}$	28	7	14	6	1	1(1), 3(4), 504(1)

TABLE 4. Results on the rank and elementary divisors of the differentials for  $SL_6(\mathbb{Z})$ .

**Theorem 4.3.** *The non-trivial homology of the Voronoi complexes associated to  $GL_N(\mathbb{Z})$  with  $N = 5, 6$  modulo  $\mathcal{S}_5$  is given by:*

$$H_n(\text{Vor}_{GL_5(\mathbb{Z})}) \cong \mathbb{Z}, \quad \text{if } n = 9, 14,$$

$$H_n(\text{Vor}_{GL_6(\mathbb{Z})}) \cong \mathbb{Z}, \quad \text{if } n = 10, 11, 15,$$

while in the case  $SL_6(\mathbb{Z})$  we get, modulo  $\mathcal{S}_7$ , that

$$H_n(\text{Vor}_{SL_6(\mathbb{Z})}) \cong \begin{cases} \mathbb{Z}, & \text{if } n = 10, 11, 12, 20, \\ \mathbb{Z}^2, & \text{if } n = 15. \end{cases}$$

Furthermore, for  $N = 7$  we get, modulo  $\mathcal{S}_7$ , that

$$H_n(\text{Vor}_{GL_7(\mathbb{Z})}) \cong \begin{cases} \mathbb{Z} & \text{if } n = 12, 13, 18, 22, 27, \\ 0 & \text{otherwise.} \end{cases}$$

Notice that, if  $N$  is odd,  $SL_N(\mathbb{Z})$  and  $GL_N(\mathbb{Z})$  have the same homology modulo  $\mathcal{S}_2$ . Notice also that, for simplicity, in the statement of the theorem we did not use the full information given by the list of elementary divisors in Tables 1–5.

A	$\Omega$	$n$	$m$	rank	ker	elementary divisors
$d_{10}$	8	60	1	1	59	1
$d_{11}$	1513	1019	60	59	960	1 (59)
$d_{12}$	37519	8899	1019	960	7939	1 (958), 2 (2)
$d_{13}$	356232	47271	8899	7938	39333	1 (7937), 2 (1)
$d_{14}$	1831183	171375	47271	39332	132043	1 (39300), 2 (29), 4 (3)
$d_{15}$	6080381	460261	171375	132043	328218	1(131993), 2(46), 12 (4)
$d_{16}$	14488881	955128	460261	328218	626910	1 (328183), 2 (33), 4(1), 12(1)
$d_{17}$	25978098	1548650	955128	626910	921740	1 (626857), 2(52), 4 (1)
$d_{18}$	35590540	1955309	1548650	921740	1033569	1 (921637), 2 (100), 42 (2), 252 (1)
$d_{19}$	37322725	1911130	1955309	1033568	877562	1 (1033458), 2 (110)
$d_{20}$	29893084	1437547	1911130	877562	559985	1 (877526), 2 (33), 6 (3)
$d_{21}$	18174775	822922	1437547	559985	262937	1 (559895), 2 (88), 6 (2)
$d_{22}$	8251000	349443	822922	262937	86506	1 (262835), 2 (98), 4 (3), 12 (1)
$d_{23}$	2695430	105054	349443	86505	18549	1 (86488), 2 (12), 6 (3), 42 (1), 84 (1)
$d_{24}$	593892	21074	105054	18549	2525	1 (18544), 2 (4), 4 (1)
$d_{25}$	81671	2798	21074	2525	273	1 (2507), 2 (18)
$d_{26}$	7412	305	2798	273	32	1 (258), 2 (7), 6 (7), 36 (1)
$d_{27}$	600	33	305	32	1	1 (23), 2 (4), 28 (3), 168 (1), 2016 (1)

TABLE 5. Results on the rank and elementary divisors of the differentials for  $GL_7(\mathbb{Z})$ , middle entries are cited from the thesis of A. Urbanska [33]. The elementary divisors for  $d_{19}$  were computed by B. Boyer and J.-G. Dumas using refinements of the techniques described in [8].

4.5. **Mass formulae for the Voronoï complex.** Let  $\chi(SL_N(\mathbb{Z}))$  be the virtual Euler characteristic of the group  $SL_N(\mathbb{Z})$ . It can be computed in two ways. First, the mass formula in [7] gives

$$\chi(SL_N(\mathbb{Z})) = \sum_{\sigma \in E} (-1)^{\dim(\sigma)} \frac{1}{|\Gamma_\sigma|} = \sum_{n=N}^{d(N)} (-1)^n \sum_{\sigma \in \Sigma_n^*} \frac{1}{|\Gamma_\sigma|},$$

where  $E$  is a family of representatives of the cells of the Voronoï complex of rank  $N$  modulo the action of  $SL_N(\mathbb{Z})$ , and  $\Gamma_\sigma$  is the stabilizer of  $\sigma$  in  $SL_N(\mathbb{Z})$ . Second, by a result of Harder [14], we know that

$$\chi(SL_N(\mathbb{Z})) = \prod_{k=2}^N \zeta(1-k),$$

hence  $\chi(SL_N(\mathbb{Z})) = 0$  if  $N \geq 3$ .

A non-trivial check of our computations is to test the compatibility of these two formulas, and the corresponding check for rank  $N = 5$  had been performed by Batut (cf. [3], where a proof of an analogous statement, for any  $N$ , but instead pertaining to *well-rounded* forms, which in our case are precisely the ones in  $\Sigma_\bullet^*$ , is attributed to Bavard [4]).

If we add together the terms  $\frac{1}{|\Gamma_\sigma|}$  for cells  $\sigma$  of the same dimension to a single term, then we get for  $N = 6$ , starting with the top dimension,

$$\begin{aligned}
& \frac{45047}{1451520} - \frac{10633}{11520} + \frac{6425}{576} - \frac{12541}{192} \\
& + \frac{7438673}{34560} - \frac{3841271}{8640} + \frac{9238}{15} - \frac{266865}{448} + \frac{14205227}{34560} - \frac{14081573}{69120} \\
& + \frac{830183}{11520} - \frac{205189}{11520} + \frac{61213}{20736} - \frac{1169}{3840} + \frac{17}{1008} - \frac{1}{2880} \\
& = \chi(SL_6(\mathbb{Z})) = 0.
\end{aligned}$$

For  $N = 7$  we obtain similarly

$$\begin{aligned}
& -\frac{290879}{107520} + \frac{13994381}{103680} - \frac{31815503}{13824} + \frac{1362329683}{69120} - \frac{6986939119}{69120} \\
& + \frac{7902421301}{23040} - \frac{340039739981}{414720} + \frac{174175928729}{120960} - \frac{132108094091}{69120} \\
& + \frac{27016703389}{13824} - \frac{13463035571}{8640} + \frac{14977461287}{15360} - \frac{22103821919}{46080} \\
& + \frac{8522164169}{46080} - \frac{17886026827}{322560} + \frac{1764066533}{138240} - \frac{101908213}{46080} + \frac{12961451}{46080} \\
& - \frac{10538393}{414720} + \frac{162617}{103680} - \frac{721}{11520} + \frac{43}{32256} \\
& = \chi(SL_7(\mathbb{Z})) = 0.
\end{aligned}$$

## 5. EXPLICIT HOMOLOGY CLASSES

### 5.1. Equivariant fundamental classes.

**Theorem 5.1.** *The top homology group  $H_{d(N)}(\text{Vor}_{SL_N(\mathbb{Z})} \otimes \mathbb{Q})$  has dimension 1. When  $N = 4, 5, 6$  or  $7$ , it is represented by the cycle*

$$\sum_{\sigma} \frac{1}{|\Gamma_{\sigma}|} [\sigma],$$

where  $\sigma$  runs through the perfect forms of rank  $N$  and the orientation of each cell is inherited from the one of  $X_N/\Gamma$ .

*Proof.* The first assertion is clear since, by (3) above and (6) below we have

$$H_{d(N)}(\text{Vor}_{SL_N(\mathbb{Z})} \otimes \mathbb{Q}) \cong H_{d(N)-N+1}(SL_N(\mathbb{Z}), St_N \otimes \mathbb{Q}) \cong H^0(SL_N(\mathbb{Z}), \mathbb{Q}) \cong \mathbb{Q}.$$

In order to prove the second claim, write the differential between codimension 0 and codimension 1 cells as a matrix  $A$  of size  $n_1 \times n_0$ , with  $n_i = |\Sigma_{d(N)-i}(\Gamma)|$  denoting the number of codimension  $i$  cells in the Voronoi cell complex. It can be checked that in each of the  $n_1$  rows of  $A$  there are precisely two non-zero entries. Moreover, the absolute value of the  $(i, j)$ -th entry of  $A$  is equal to the quotient  $|\Gamma_{\sigma_j}|/|\Gamma_{\tau_i}|$  (an integer), where  $\sigma_j \in \Sigma_{d(N)}(\Gamma)$  and  $\tau_i \in \Sigma_{d(N)-1}(\Gamma)$ . Finally, one can multiply some columns by  $-1$  (which amounts to changing the orientation of the corresponding codimension 0 cell) in such a way that each row has exactly one positive and one negative entry.  $\square$

**Example 5.2.** For  $N = 5$  the differential matrix  $d_{14}$  (cf. Table 2) between codimension 0 and codimension 1 is given by

$$\begin{pmatrix} 40 & 0 & -15 \\ 40 & -15 & 0 \end{pmatrix},$$

so the kernel is generated by  $(3, 8, 8) = 11520 \left( \frac{1}{3840}, \frac{1}{1440}, \frac{1}{1440} \right)$ , while the orders of the three automorphism groups are 3840, 1440 and 1440, respectively.

**Example 5.3.** Similarly, the differential  $d_{20} : V_{20} \rightarrow V_{19}$  for rank  $N = 6$  (cf. Table 3) is represented by the matrix

$$\begin{pmatrix} 0 & 0 & 96 & 0 & 0 & 0 & -21 \\ 3240 & 0 & 0 & 0 & -21 & 0 & 0 \\ 0 & 0 & 1440 & 0 & 0 & -3 & 0 \\ 0 & 0 & 0 & 18 & 0 & -6 & 0 \\ -12960 & 0 & 0 & 0 & 0 & 12 & 0 \\ -3240 & 0 & 0 & 9 & 0 & 0 & 0 \\ 0 & -360 & 0 & 1 & 0 & 0 & 0 \\ -4320 & 0 & 0 & 12 & 0 & 0 & 0 \\ 0 & 0 & 960 & -6 & 0 & 0 & 0 \\ 0 & -216 & 96 & 0 & 0 & 0 & 0 \\ -45 & 45 & 0 & 0 & 0 & 0 & 0 \\ -2592 & 0 & 1152 & 0 & 0 & 0 & 0 \\ -3240 & 0 & 1440 & 0 & 0 & 0 & 0 \\ -432 & 0 & 192 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Its kernel is generated by

$$(28, 28, 63, 10080, 4320, 30240, 288)$$

while the orders of the corresponding automorphism groups are, respectively,

$$103680, 103680, 46080, 288, 672, 96, 10080,$$

and we note that  $28 \cdot 103680 = 63 \cdot 46080 = 10080 \cdot 288 = 4320 \cdot 672 = 30240 \cdot 96$ .

**5.2. An explicit non-trivial homology class for rank  $N = 5$ .** The integer kernel of the  $7 \times 1$ -matrix of  $d_9$  for  $GL_5(\mathbb{Z})$ , given by  $(0, 0, 0, 0, -1, 0, 1)$ , is spanned by the image of  $d_{10}$  (the latter being given, up to permutation of rows and columns, by the transpose of the matrix (4) below), together with  $(2, 1, -1, -1, -1, 1, 1)$ . The latter vector therefore provides the coefficients of a non-trivial homology class in  $H_9(\text{Vor}_{GL_5(\mathbb{Z})}) \cong H^5(GL_5(\mathbb{Z}), \mathbb{Z})$  (modulo  $\mathcal{S}_5$ ), given as a linear combination of cells (in terms of minimal vectors) as follows:

$$\begin{aligned} & 2\varphi([e_1, e_2, \bar{e}_{23}, \bar{e}_{13}, e_3, \bar{e}_{34}, \bar{e}_{14}, \bar{e}_{45}, \bar{e}_{35}, \bar{e}_{25}]) \\ & + \varphi([e_1, e_2, e_3, e_4, e_{24}, e_{34}, e_5, e_{15}, e_{35}, e_{1245}]) \\ & - \varphi([e_1, \bar{e}_{12}, e_2, \bar{e}_{23}, e_3, \bar{e}_{34}, \bar{e}_{14}, \bar{e}_{45}, \bar{e}_{35}, \bar{e}_{25}]) \\ & - \varphi([e_1, e_2, e_3, e_4, e_{14}, e_{24}, e_{34}, e_5, e_{35}, e_{1245}]) \\ & - \varphi([e_1, \bar{e}_{12}, e_2, \bar{e}_{13}, e_3, \bar{e}_{14}, e_4, u, \bar{e}_{45}, v]) \\ & + \varphi([e_1, e_2, e_3, e_{14}, e_{24}, e_{34}, e_5, e_{15}, e_{35}, e_{1245}]) \\ & + \varphi([e_1, e_2, e_3, e_4, e_{24}, e_{34}, e_{25}, e_{35}, e_{1245}, e_{1345}]). \end{aligned}$$

where we denote the standard basis vectors in  $\mathbb{R}^5$  by  $e_i$ , and we put  $e_{ij} = e_i + e_j$ ,  $\bar{e}_{ij} = -e_i + e_j$  and  $e_{ijkl} = e_i + e_j + e_k + e_l$ , as well as  $u = e_5 - e_1 - e_4$  and  $v = e_5 - e_2 - e_3$ .

## 6. SPLITTING OFF THE VORONOÏ COMPLEX $\text{Vor}_N$ FROM $\text{Vor}_{N+1}$ FOR SMALL $N$

In this section, we will be concerned with  $\Gamma = GL_N(\mathbb{Z})$  only and we adopt the notation  $\Sigma_n(N) = \Sigma_n(GL_N(\mathbb{Z}))$  for the sets of representatives.

**6.1. Inflating well-rounded forms.** Let  $A$  be the symmetric matrix attached to a form  $h$  in  $C_N^*$ . Suppose the cell associated to  $A$  is *well-rounded*, i.e., its set of minimal vectors  $S = S(A)$  spans the underlying vector space  $\mathbb{R}^N$ . Then we can associate to it a form  $\tilde{h}$  with matrix  $\tilde{A} = \begin{pmatrix} A & 0 \\ 0 & m(A) \end{pmatrix}$  in  $C_{N+1}^*$ , where  $m(A)$  denotes the minimum positive value of  $A$  on  $\mathbb{Z}^N$ . The set  $\tilde{S}$  of minimal vectors of  $\tilde{A}$  contains the ones from  $S$ , each vector being extended by an  $(N+1)$ -th coordinate 0. Furthermore,  $\tilde{S}$  contains the additional minimal vectors  $\pm e_{N+1} = \pm(0, \dots, 0, 1)$ , and hence it spans  $\mathbb{R}^{N+1}$ , i.e.,  $\tilde{A}$  is well-rounded as well. In the following, we will call forms like  $\tilde{A}$  as well as their associated cells *inflated*.


The stabilizer of  $h$  in  $GL_N(\mathbb{Z})$  thereby embeds into the one of  $\tilde{h}$  inside  $GL_{N+1}(\mathbb{Z})$  (at least modulo  $\pm \text{Id}$ ) under the usual stabilization map.


Note that, by iterating the same argument  $r$  times,  $A$  induces a well-rounded form also in  $\Sigma_\bullet^*(N+r)$  which, for  $r \geq 2$ , does not belong to  $\Sigma_\bullet(N+r)$  since there is an obvious orientation-reversing automorphism of the inflated form, given by the permutation which swaps the last two coordinates.

## 6.2. The case $N = 5$ .

**Theorem 6.1.** *The complex  $\text{Vor}_{GL_5(\mathbb{Z})}$  is isomorphic to a direct factor of  $\text{Vor}_{GL_6(\mathbb{Z})}$ , with degrees shifted by 1.*

*Proof.* The Voronoï complex of  $GL_5(\mathbb{Z})$  can be represented by the following weighted graph with levels





Here the nodes in line  $j$  (marked on the left) represent the elements in  $\Sigma_{d(N)-j}(5)$ , i.e. we have 3, 2, 0, 1, 6, 7 and 1 cells in codimensions 0, 1, 2, 3, 4, 5 and 6, respectively, and arrows show incidences of those cells, while numbers attached to arrows give the corresponding incidence multiplicities. Since entering the multiplicities relating codimensions 4 and 5 would make the graph rather unwieldy, we give them instead in terms of the matrix corresponding to the differential  $d_{10}$  connecting dimension 10 to 9 (columns refer, in this order, to  $\sigma_5^1, \dots, \sigma_5^7$ , while rows refer to  $\sigma_4^1, \dots, \sigma_4^6$ )

$$(4) \quad \begin{pmatrix} -5 & 0 & -5 & 0 & -1 & 0 & 0 \\ 0 & -2 & 0 & 2 & -2 & 0 & 0 \\ 2 & -2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 1 & 0 & 0 & 0 \\ -1 & -2 & 1 & 0 & 1 & 0 & 0 \\ 0 & 4 & 0 & 0 & 0 & -1 & -1 \end{pmatrix}.$$

As is apparent from the picture, there are two connected components in that graph. The corresponding graph for  $GL_6(\mathbb{Z})$  has three connected components, two of which are "isomorphic" (as weighted graphs with levels) to the one above for  $GL_5(\mathbb{Z})$ , except for a shift in codimension by 5 (e.g. codimension 0 cells in  $\Sigma_{\bullet}(5)$  correspond to codimension 5 cells in  $\Sigma_{\bullet}(6)$ ), i.e. a shift in dimension by 1.

In fact, it is possible, after appropriate coordinate transformations, to identify the minimal vectors (viewed up to sign) of any given cell in the two inflated components of  $\Sigma_{\bullet}(6)$  alluded to above with the minimal vectors of another cell which is inflated from one in  $\Sigma_{\bullet}(5)$ , except precisely *one* minimal vector (up to sign) which is *fixed* under the stabilizer of the cell.

Let us illustrate this correspondence for the top-dimensional cell  $\sigma$  of the perfect form  $P_5^1 \in \Sigma_{14}(5)$ , also denoted  $P(5, 1)$  in [15] and  $D_5$  in [19], with the list  $m(P_5^1)$  of minimal vectors given already at the end of §5.2.

Using the algorithm described in §4.1, the corresponding inflated cell  $\tilde{\sigma}$  in  $\Sigma_{15}(6)$  can be found to be, in terms of its 21 minimal vectors of the perfect form  $P_6^1$  in Jaquet's notation (see [15] and §5.2 for the full list  $m(P_6^1)$ ),



$v_1$	$v_2$	$v_4$	$v_5$	$v_{10}$	$v_{12}$	$v_{13}$	$v_{14}$	$v_{16}$	$v_{17}$	$v_{18}$	$v_{22}$	$v_{24}$	$v_{25}$	$v_{26}$	$v_{27}$	$v_{29}$	$v_{33}$	$v_{34}$	$v_{35}$	$v_{36}$
1	-1	0	-1	0	0	0	-1	1	1	0	1	1	0	1	0	0	1	0	0	1
0	1	-1	0	0	0	-1	0	1	0	1	1	0	1	0	1	0	0	1	0	1
0	0	1	1	0	-1	0	0	-1	0	0	-1	0	0	-1	-1	0	-1	-1	0	-1
0	0	0	0	1	0	0	0	-1	-1	-1	0	0	0	-1	-1	-1	0	0	0	-1
0	0	0	0	0	1	1	1	-1	-1	-1	-1	-1	-1	0	0	0	0	0	0	-1
0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1	2

The transformation

$$\gamma = \begin{pmatrix} 0 & -1 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & -1 & -1 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 \\ -1 & -1 & -1 & 0 & -1 & 0 \end{pmatrix}$$

sends  $v_1$  to  $(0, 0, 0, 0, 0, 1)$  and sends each of the other vectors to the corresponding one of the form  $(v, 0)$  where  $v$  is the corresponding minimal vector for  $P_5^1$  (in the order given above).

One can verify that the other two perfect forms  $P_5^2$  and  $P_5^3$  (denoted by Voronoï  $A_5$  and  $\varphi_2$ , respectively) give rise to a corresponding inflated cell in  $\Sigma_{15}(6)$  in a similar way.

Concerning the cells of *positive* codimension in  $\Sigma_\bullet(5)$ , it turns out that these all have a representative which is a facet in  $\sigma$ . Furthermore, the matrix  $\gamma$  induces an isomorphism from the subcomplex of  $\Sigma_\bullet(6)$  spanned by  $\tilde{\sigma}$  and all its facets to the complex obtained by inflation, as in §6.1 above, from the complex spanned by  $\sigma_5$  and all its facets. Finally, one can verify that the cells attached to  $P_5^2$  and  $P_5^3$  are conjugate, after inflation, to cells in  $\Sigma_{15}(6)$ , and that the differentials for  $\text{Vor}_{GL_5}$  and  $\text{Vor}_{GL_6}$  agree on these. This ends the proof of the theorem.  $\square$

**6.3. Other cases.** A similar situation holds for  $\Sigma_\bullet(3)$  and  $\Sigma_\bullet(4)$ , but as  $\Sigma_\bullet(3)$  consists of a single cell only, the picture is far less significant.

For  $N = 4$ , there is only one cell leftover in  $\Sigma_\bullet(4)$ , in fact in  $\Sigma_6(4)$ , and it is already inflated from  $\Sigma_5(3)$ . Hence its image in  $\Sigma_7^*(5)$  will allow an orientation reversing automorphism and hence will not show up in  $\Sigma_7(5)$ . This illustrates the remark at the end of 6.1.

Finally, for  $N = 6$ , the cells in the third component of the incidence graph for  $GL_6(\mathbb{Z})$  mentioned in the proof of Theorem 6.1 above appear, in inflated form, in the Voronoï complex for  $GL_7(\mathbb{Z})$  which inherits the homology of that component, since in the weighted graph of  $GL_7(\mathbb{Z})$ , which is connected, there is only one incidence of an inflated cell with a non-inflated one. Therefore we do not have a splitting in this case.

## 7. THE COHOMOLOGY OF MODULAR GROUPS

**7.1. Preliminaries.** Recall the following simple fact:

**Lemma 7.1.** *Assume that  $p$  is a prime and  $g \in GL_N(\mathbb{R})$  has order  $p$ . Then  $p \leq N + 1$ .*

*Proof.* The minimal polynomial of  $g$  is the cyclotomic polynomial  $x^{p-1} + x^{p-2} + \dots + 1$ . By the Cayley-Hamilton theorem, this polynomial divides the characteristic polynomial of  $g$ . Therefore  $p - 1 \leq N$ .  $\square$

We shall also need the following result:

**Lemma 7.2.** *The action of  $\mathrm{GL}_N(\mathbb{R})$  on the symmetric space  $X_N$  preserves its orientation if and only if  $N$  is odd.*

*Proof.* The subgroup  $\mathrm{GL}_N(\mathbb{R})^+ \subset \mathrm{GL}_N(\mathbb{R})$  of elements with positive determinant is the connected component of the identity, therefore it preserves the orientation of  $X_N$ . Any  $g \in \mathrm{GL}_N(\mathbb{R})$  which is not in  $\mathrm{GL}_N(\mathbb{R})^+$  is the product of an element of  $\mathrm{GL}_N(\mathbb{R})^+$  with the diagonal matrix  $\varepsilon = \mathrm{diag}(-1, 1, \dots, 1)$ , so we just need to check when  $\varepsilon$  preserves the orientation of  $X_N$ . The tangent space  $TX_N$  of  $X_N$  at the origin consists of real symmetric matrices  $m = (m_{ij})$  of trace zero. The action of  $\varepsilon$  is given by  $m \cdot \varepsilon = \varepsilon^t m \varepsilon$  (cf. §2.1) and we get

$$(m \cdot \varepsilon)_{ij} = m_{ij}$$

unless  $i = 1$  or  $j = 1$  and  $i \neq j$ , in which case  $(m \cdot \varepsilon)_{ij} = -m_{ij}$ . Let  $\delta_{ij}$  be the matrix with entry 1 in row  $i$  and column  $j$ , and zero elsewhere. A basis of  $TX_N$  consists of the matrices  $\delta_{ij} + \delta_{ji}$ ,  $i \neq j$ , together with  $N - 1$  diagonal matrices. For this basis, the action of  $\varepsilon$  maps  $N - 1$  vectors  $v$  to their opposite  $-v$  and fixes the other ones. The lemma follows.  $\square$

**7.2. Borel/Serre duality.** According to Borel and Serre ([6], Thm. 11.4.4 and Thm. 11.5.1), the group  $\Gamma = \mathrm{SL}_N(\mathbb{Z})$  or  $\mathrm{GL}_N(\mathbb{Z})$  is a virtual duality group with dualizing module

$$H^{v(N)}(\Gamma, \mathbb{Z}[\Gamma]) = \mathrm{St}_N \otimes \tilde{\mathbb{Z}},$$

where  $v(N) = N(N - 1)/2$  is the virtual cohomological dimension of  $\Gamma$  and  $\tilde{\mathbb{Z}}$  is the orientation module of  $X_N$ . It follows that there is a long exact sequence

$$(5) \quad \dots \rightarrow H_n(\Gamma, \mathrm{St}_N) \rightarrow H^{v(N)-n}(\Gamma, \tilde{\mathbb{Z}}) \rightarrow \hat{H}^{v(N)-n}(\Gamma, \tilde{\mathbb{Z}}) \rightarrow H_{n-1}(\Gamma, \mathrm{St}_N) \rightarrow \dots$$

where  $\hat{H}^*$  is the Farrell cohomology of  $\Gamma$  [11]. From Lemma 7.1 and the Brown spectral sequence ([7], X (4.1)) we deduce that  $\hat{H}^*(\Gamma, \tilde{\mathbb{Z}})$  lies in  $\mathcal{S}_{N+1}$ . Therefore

$$(6) \quad H_n(\Gamma, \mathrm{St}_N) \equiv H^{v(N)-n}(\Gamma, \tilde{\mathbb{Z}}), \text{ modulo } \mathcal{S}_{N+1}.$$

When  $N$  is odd, then  $\mathrm{GL}_N(\mathbb{Z})$  is the product of  $\mathrm{SL}_N(\mathbb{Z})$  by  $\mathbb{Z}/2$ , therefore

$$H^m(\mathrm{GL}_N(\mathbb{Z}), \mathbb{Z}) \equiv H^m(\mathrm{SL}_N(\mathbb{Z}), \mathbb{Z}), \text{ modulo } \mathcal{S}_2.$$

When  $N$  is even, then the action of  $\mathrm{GL}_N(\mathbb{Z})$  on  $\tilde{\mathbb{Z}}$  is given by the sign of the determinant (see Lemma 7.2) and Shapiro's lemma gives

$$(7) \quad H^m(\mathrm{SL}_N(\mathbb{Z}), \mathbb{Z}) = H^m(\mathrm{GL}_N(\mathbb{Z}), M),$$

with

$$M = \mathrm{Ind}_{\mathrm{SL}_N(\mathbb{Z})}^{\mathrm{GL}_N(\mathbb{Z})} \mathbb{Z} \equiv \mathbb{Z} \oplus \tilde{\mathbb{Z}}, \text{ modulo } \mathcal{S}_2.$$

**7.3. The cohomology of modular groups.** When  $\Gamma = SL_N(\mathbb{Z})$  or  $GL_N(\mathbb{Z})$ , where  $N \leq 7$ , we know  $H^m(\Gamma, \mathbb{Z})$  by combining (3) (end of §3.4), Theorem 4.3 and (6). As shown above, this allows us to compute the cohomology of  $\Gamma$  with trivial coefficients. The results are given in Theorem 7.3 below.

**Theorem 7.3.** (i) *Modulo  $\mathcal{S}_5$  we have*

$$H^m(SL_5(\mathbb{Z}), \mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } m = 0, 5, \\ 0 & \text{otherwise.} \end{cases}$$

(ii) *Modulo  $\mathcal{S}_7$  we have*

$$H^m(GL_6(\mathbb{Z}), \mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } m = 0, 5, 8, \\ 0 & \text{otherwise,} \end{cases}$$

and

$$H^m(SL_6(\mathbb{Z}), \mathbb{Z}) = \begin{cases} \mathbb{Z}^2 & \text{if } m = 5, \\ \mathbb{Z} & \text{if } m = 0, 8, 9, 10, \\ 0 & \text{otherwise.} \end{cases}$$

(iii) *Modulo  $\mathcal{S}_7$  we get that*

$$H^m(SL_7(\mathbb{Z}), \mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } m = 0, 5, 9, 14, 15, \\ 0 & \text{if } 1 \leq m \leq 20. \end{cases}$$

For the proof of the final statement on integral cohomology (modulo  $\mathcal{S}_7$ ) we use the fact that there are no primes  $p > 7$  that divide the elementary divisors of the corresponding differentials or the order of the stabilizer of a cell in  $\Sigma_{27-m}$ .

**Remark 7.4.** Morita asks in [21] whether the class of infinite order in  $H^5(GL_5(\mathbb{Z}), \mathbb{Z})$  survives in the cohomology of the group of outer automorphisms of the free group of rank six.

**Remark 7.5.** It was shown by A. Borel [5] that, for  $N$  large enough,  $H^5(SL_N(\mathbb{Z}), \mathbb{Q})$  has dimension one. In view of Theorem 7.3 it is tempting to believe that the restriction map from  $H^5(SL_N(\mathbb{Z}), \mathbb{Q})$  to  $H^5(SL_5(\mathbb{Z}), \mathbb{Q})$  is an isomorphism. We have been unable to show that. An analogous statement holds for  $H^9(SL_N(\mathbb{Z}), \mathbb{Q})$ . Theorem 7.3 suggests that the non-trivial cohomology class already occurs when  $N = 6$  and 7, i.e., in the “non-stable range”.

## 8. HOMOLOGY OF MODULAR GROUPS WITH COEFFICIENTS IN THE STEINBERG MODULE

In this section we compute some homology groups of  $GL_N(\mathbb{Z})$  with coefficients the Steinberg module. Note that, when  $N \geq 1$ , the group  $H_0(GL_N(\mathbb{Z}), St_N)$  vanishes [19].

**Theorem 8.1.** (i) *Modulo  $\mathcal{S}_2$  we have*

$$(8) \quad H_3(GL_3(\mathbb{Z}), St_3) \cong \mathbb{Z}$$

and

$$(9) \quad H_3(GL_4(\mathbb{Z}), St_4) \cong \mathbb{Z}.$$

(ii) *The following groups lie in  $\mathcal{S}_2$ :*

$$\begin{aligned} &H_4(GL_2(\mathbb{Z}), St_2), & H_5(GL_2(\mathbb{Z}), St_2), \\ &H_4(GL_3(\mathbb{Z}), St_3), \\ &H_2(GL_4(\mathbb{Z}), St_4), \\ &H_1(GL_5(\mathbb{Z}), St_5), & H_2(GL_5(\mathbb{Z}), St_5), \\ &H_1(GL_6(\mathbb{Z}), St_6). \end{aligned}$$

(iii) *The groups  $H_2(GL_6(\mathbb{Z}), St_6)$  and  $H_1(GL_7(\mathbb{Z}), St_7)$  lie in  $\mathcal{S}_5$ .*

In order to prepare for the proof, we first compute several terms in the spectral sequence  $E_{pq}^1$  of §3.2. This is done in five lemmas, dealing with  $GL_4(\mathbb{Z})$ ,  $GL_5(\mathbb{Z})$ ,  $GL_6(\mathbb{Z})$  (separating the cases  $p + q = 6$  and  $p + q = 7$ ) and  $GL_7(\mathbb{Z})$ , respectively. We will show that the  $E^1$  terms of the respective equivariant spectral sequences, in the desired ranges, are all zero modulo some torsion classes (mostly  $\mathcal{S}_2$ ) which will allow us to deduce the claims. The general strategy is as follows: if  $G$  is the stabilizer of a cell, we will construct the maximal normal subgroup  $H$  of  $G$  which acts trivially on the cell. The quotient  $G/H$  will be in  $\mathcal{S}_2$ . Hence, using the Lyndon/Hochschild/Serre spectral sequence (denoted LHS in the remaining of the paper), the computation of the homology of  $G$  with coefficients in  $\tilde{\mathbb{Z}}$  (i.e.,  $\mathbb{Z}$  endowed with the  $G$ -action on the cell) will be reduced to the computation of the homology of  $H$  with trivial coefficients. It will result that, in general, the corresponding homology groups lie in  $\mathcal{S}_2$ . We start by giving two general lemma, with straightforward proofs, that will be systematically used in our arguments.

**Lemma 8.2.** *Let  $\Gamma$  be a subgroup of  $GL_N(\mathbb{Z})$  and let  $\sigma$  be a cell of  $\Sigma_n^*(\Gamma)$ , for some  $n$ . Let  $\Gamma_\sigma$  be the stabilizer of the cell  $\sigma$  in  $\Gamma$ . Then there exists a normal subgroup  $H$  of  $\Gamma_\sigma$ , acting trivially on the cell  $\sigma$  and with quotient  $\Gamma_\sigma/H$  isomorphic to  $\mathbb{Z}/2$ .*

*Proof.* The action on the cell is given by  $\eta$  (see §3.1). It defines a morphism  $\Gamma_\sigma \rightarrow \mathbb{Z}/2$  mapping  $\gamma$  to  $\eta(\gamma \cdot \sigma, \sigma)$ . We define  $H$  to be the kernel of this map.  $\square$

**Lemma 8.3.** *Consider a short exact sequence of finite groups*

$$1 \rightarrow H \rightarrow G \rightarrow Q \rightarrow 1.$$

*Assume  $Q \in \mathcal{S}_p$  for some prime  $p$ . Let  $M$  be a  $G$ -module and  $k$  a positive integer. If  $H_i(H, M) \in \mathcal{S}_p$  for all positive  $i \leq k$ , then  $H_k(G, M) \in \mathcal{S}_p$ .*

Now, we can compute the relevant parts of the equivariant spectral sequences of §3.2.

**Lemma 8.4.** *The terms  $E_{5,1}^1$ ,  $E_{4,2}^1$ ,  $E_{3,3}^2$ ,  $E_{4,1}^1$  and  $E_{3,2}^1$  of the equivariant spectral sequence associated to  $\Gamma = GL_4(\mathbb{Z})$  lie in  $\mathcal{S}_2$ .*

*Proof.* • *Computation of  $E_{5,1}^1$ .* According to [19], Lemma 3.2, the set  $\Sigma_5^*(SL_4(\mathbb{Z}))$  consists of four cells, denoted  $\sigma_i^5$  ( $i = 2, 3, 4, 5$ ) in op.cit.

The stabilizer of  $\sigma_5^5$  in  $PGL_4(\mathbb{Z})$  is isomorphic to  $\mathfrak{S}_2 \times \mathfrak{S}_3$  (op.cit., p.121), each factor acting non-trivially on the orientation module of  $\sigma_5^5$ . It follows that  $Stab(\sigma_5^5)$  contains a subgroup isomorphic to  $\mathfrak{S}_3$  and preserving the orientation of  $\sigma_5^5$ . Therefore, modulo  $\mathcal{S}_2$ , we get

$$H_1(Stab(\sigma_5^5), \tilde{\mathbb{Z}}) = H_1(\mathfrak{S}_3, \mathbb{Z}).$$

From the exact sequence

$$1 \longrightarrow \mathbb{Z}/3 \longrightarrow \mathfrak{S}_3 \longrightarrow \mathbb{Z}/2 \longrightarrow 1$$

we deduce that, modulo  $\mathcal{S}_2$ ,

$$H_1(\mathfrak{S}_3, \mathbb{Z}) = H_0(\mathbb{Z}/2, H_1(\mathbb{Z}/3, \mathbb{Z})) = 0.$$

The stabilizer  $Stab(\sigma_4^5)$  in  $GL_4(\mathbb{Z})$  has order 32. Therefore its first homology group lies in  $\mathcal{S}_2$ .

The cell  $\sigma_3^5$  of [19] (p.110) has its stabilizer (in  $GL_4(\mathbb{Z})$ ) generated by the matrices

$$g_{2,1} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ -1 & -1 & 0 & -1 \\ 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad g_{2,2} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & -1 & -2 & -2 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad g_{2,3} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad g_{2,4} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & -1 & -2 & -2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Denote by  $G_2$  this group. It is of order  $288 = 2^5 \cdot 3^2$ . All the generators have a non-trivial action on the cell, except  $g_{2,2}$ . Let  $H_2$  be the subgroup of  $G_2$  generated by  $g_{2,1}g_{2,3}$ ,  $g_{2,2}$  and  $g_{2,1}g_{2,4}$ . By construction  $H_2$  acts trivially on the cell. Using GAP, we can check that  $H_2$  is normal in  $G_2$  and the quotient  $G_2/H_2$  is isomorphic to  $\mathbb{Z}/2$ . Furthermore, the derived subgroup of  $H_2$  is isomorphic to  $\mathbb{Z}/6 \times \mathbb{Z}/3$  and its abelianization is isomorphic to  $\mathbb{Z}/4 \times \mathbb{Z}/2$ . As a result  $H_1(G_2, \mathbb{Z}) = H_0(\mathbb{Z}/2, H_1(H_2, \mathbb{Z})) = H_0(\mathbb{Z}/2, H_1(H_2; \mathbb{Z})) = 0 \pmod{\mathcal{S}_2}$ .

The last cell to consider is  $\sigma_2^5$ . Let  $G_4$  be the stabilizer of this cell. A set of generators of  $G_4$  is given by the matrices

$$g_{4,1} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad g_{4,2} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

$$g_{4,3} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & -1 & 0 & -1 \\ 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad g_{4,4} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & -1 & -1 & -1 \end{pmatrix}, \quad g_{4,5} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 0 & -1 & -1 \\ -1 & -1 & 0 & -1 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Its order is  $96 = 2^5 \cdot 3$ . The group  $G_4$  is isomorphic to  $\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathfrak{S}_4$ . Among the generators, only  $g_{4,3}$  and  $g_{4,5}$  have a non-trivial action. The subgroup generated by  $g_{4,1}, g_{4,2}, g_{4,4}, g_{4,3}g_{4,5}$  is normal and isomorphic to  $\mathbb{Z}/2 \times \mathfrak{S}_4$ . Its abelianization is isomorphic to  $\mathbb{Z}/2 \times \mathbb{Z}/2$ . We deduce that  $H_1(G_4, \mathbb{Z}) = H_0(\mathbb{Z}/2, H_1(\mathbb{Z}/2 \times \mathfrak{S}_4, \mathbb{Z})) = 0 \pmod{\mathcal{S}_2}$ , and this ends the computations of  $E_{5,1}^1$ .

- *Computation of  $E_{4,2}^1$  and  $E_{4,1}^1$ .* According to [19], Lemma 3.2, the set  $\Sigma_4^*(SL_4(\mathbb{Z}))$  consists of the three cells  $\sigma_2^4$ ,  $\sigma_3^4$  and  $\sigma_4^4$ . The stabilizer of  $\sigma_4^4$  in  $PGL_4(\mathbb{Z})$  is isomorphic to  $\mathfrak{S}_5$  (op.cit., p.121). Modulo  $\mathcal{S}_2$ , the group  $H_2(Stab(\sigma_4^4), \mathbb{Z})$  is thus a quotient of  $H_2(\mathbb{Z}/5, \mathbb{Z}) \oplus H_2(\mathbb{Z}/3, \mathbb{Z}) = 0$ .

Furthermore, the alternating subgroup  $\mathfrak{A}_5 \subset \mathfrak{S}_5$  preserves the orientation of  $\sigma_4^4$  ([19], Lemma 3.4), and it is equal to its commutator subgroup.

Therefore, modulo  $\mathcal{S}_2$ ,

$$H_1(\text{Stab}(\sigma_4^4), \tilde{\mathbb{Z}}) = H_0(\mathbb{Z}/2, H_1(\mathfrak{A}_5, \mathbb{Z})) = 0.$$

- *Computation of  $E_{3,3}^2$ .* The only cell in  $\Sigma_3^*(GL_4(\mathbb{Z}))$  is the cell  $\sigma_3^3$  of [19], Lemma 3.2. The action of  $\text{Stab}(\sigma_3^3)$  on the orientation module is not trivial ([19], Lemma 3.3). According to [30], §3.2, we have

$$H_3(\text{Stab}(\sigma_3^3), \tilde{\mathbb{Z}}) \cong H_3(\mathfrak{A}_4, \mathbb{Z}) = \mathbb{Z}/3$$

modulo  $\mathcal{S}_2$ , and the differential

$$d^1 : H_3(\text{Stab}(\sigma_3^4), \tilde{\mathbb{Z}}) \rightarrow H_3(\text{Stab}(\sigma_3^3), \tilde{\mathbb{Z}})$$

is surjective. Therefore  $E_{3,3}^2$  lies in  $\mathcal{S}_2$ .

- *Computation of  $E_{3,2}^1$ .* Modulo  $\mathcal{S}_2$ , we get that  $E_{3,2}^1 = H_2(\text{Stab}(\sigma_3^3), \tilde{\mathbb{Z}})$  is a quotient of  $H_2(\mathbb{Z}/3, \mathbb{Z}) = 0$ .

□

**Lemma 8.5.** *The terms  $E_{4,1}^1$ ,  $E_{4,2}^1$ ,  $E_{5,1}^1$  and  $E_{6,0}^1$  of the equivariant spectral sequence associated to  $\Gamma = GL_5(\mathbb{Z})$  are zero modulo  $\mathcal{S}_2$ .*

*Proof.* • As none of the cells of  $\Sigma_6^*$  has its orientation preserved by the action of its stabilizer (see Fig.1), we have  $E_{6,0}^1 = 0 \pmod{\mathcal{S}_2}$ .

- *Computation of  $E_{5,1}^1$ .* We need to know the group  $H_1(\text{Stab}_\Gamma(\sigma), \tilde{\mathbb{Z}})$  for all five cells  $\sigma \in \Sigma_5^*$  (cf. Table 2). Up to equivalence under  $GL_5(\mathbb{Z})$ , these cells are contained in  $\sigma(P_5^2)$ , where  $P_5^2$  is the perfect form of rank 5 described in [20], §6.4 (and mentioned above at the end of §6.2). We will denote these five cells by  $\sigma'_i$  with  $i = 1, \dots, 5$ .

*Analyzing the cell  $\sigma'_1$ .* First, let us describe  $\sigma'_1$ . The 15 minimal vectors of  $P_5^2$  are given below, together with their label:

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	0	0	0	1	0	0	0	1	0	0	1	1	0	1
0	1	0	0	0	1	0	0	0	1	0	1	0	1	1
0	0	1	0	0	0	1	0	0	0	1	0	1	1	1
0	0	0	1	1	1	1	0	0	0	0	1	1	1	1
0	0	0	0	0	0	0	1	1	1	1	1	1	1	1

The vertices of the cell  $\sigma'_1$  are the forms  $\hat{v}$  where  $v$  is one of the vectors labelled by 1, 2, 3, 4, 5 and 8. Set  $G'_1 = \text{Stab}_\Gamma(\sigma'_1)$ . A set of generators of  $G'_1$  is given by the following six matrices of  $GL_5(\mathbb{Z})$ , of respective order 6,

2, 2, 2, 2, 6 :

$$g'_{1,1} = \begin{pmatrix} -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix}, g'_{1,2} = \begin{pmatrix} -1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}, g'_{1,3} = \begin{pmatrix} -1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$g'_{1,4} = \begin{pmatrix} 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, g'_{1,5} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, g'_{1,6} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

The order of  $G'_1$  is  $576 = 2^6 \cdot 3^2$ . Thus, *a priori*, we could expect some 3-torsion in the homology of this group. Only  $g'_{1,2}$  and  $g'_{1,6}$  have a trivial action on the cells. Using GAP [12], we get that the group  $G'_1$  is isomorphic to  $\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathfrak{S}_3 \times \mathfrak{S}_4$ . Let  $H'_1$  be the subgroup of  $G'_1$  generated by  $g'_{1,2}, g'_{1,6}, g'_{1,1}g'_{1,3}, g'_{1,1}g'_{1,4}, g'_{1,1}g'_{1,5}$ . By construction this subgroup has trivial action on the cell. It is normal and has order 288. We then have  $G'_1/H'_1 = \mathbb{Z}/2$ . Furthermore, the derived subgroup of  $H'_1$  is isomorphic to  $\mathbb{Z}/3 \times \mathfrak{A}_4$  and the quotient  $H'_1/[H'_1, H'_1]$  is isomorphic to the product of three copies of  $\mathbb{Z}/2$ . Thus the first homology group of  $H'_1$  with trivial coefficients is zero modulo  $\mathcal{S}_2$ . By Lemma 8.3, we get

$$H_1(G'_1; \tilde{\mathbb{Z}}) = H_0(\mathbb{Z}/2, H_1(H'_1; \mathbb{Z})) = 0 \pmod{\mathcal{S}_2}.$$

*Analyzing the cell  $\sigma'_2$ .* The cell  $\sigma'_2$  is given by the vectors labelled by 1, 2, 3, 5, 6 and 8. Denote its stabilizer by  $G'_2$ . A set of generators of  $G'_2$  consists of the following six matrices of  $GL_5(\mathbb{Z})$ , of respective order 2, 4, 4, 4, 2, 2:

$$g'_{2,1} = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 \end{pmatrix}, g'_{2,2} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}, g'_{2,3} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 \end{pmatrix},$$

$$g'_{2,4} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, g'_{2,5} = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, g'_{2,6} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

The order of the stabilizer is  $384 = 2^7 \cdot 3$ .

Using GAP we get that  $G'_2$  is isomorphic to  $\mathbb{Z}/2 \times \mathfrak{S}_4 \times D_8$ . The generators  $g'_{2,1}, g'_{2,2}$  and  $g'_{2,6}$  act trivially on the cell. Consider the subgroup  $H'_2$  of  $G'_2$  generated by  $g'_{2,1}, g'_{2,2}, g'_{2,6}, g'_{2,3}g'_{2,4}$  and  $g'_{2,3}g'_{2,5}$ . This subgroup is normal and acts trivially on the cell. Its order is 192, thus the quotient  $G'_2/H'_2$  is of order 2. We can check with GAP that the abelianization of  $H'_2$  is isomorphic to  $(\mathbb{Z}/2)^3$ . We deduce, by Lemma 8.3, that

$$H_1(G'_2; \tilde{\mathbb{Z}}) = H_0(\mathbb{Z}/2; H_1(H'_2; \mathbb{Z})) = 0 \pmod{\mathcal{S}_2}.$$

*Analyzing the cell  $\sigma'_3$ .* The cell  $\sigma'_3$  is given by the vectors labelled by 2, 3, 5, 6, 8, 9. Denote its stabilizer by  $G'_3$ . A set of generators of  $G'_3$  consists of

the following three matrices of  $GL_5(\mathbb{Z})$ , of respective order 2, 10, 4:

$$g'_{3,1} = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 \end{pmatrix}, \quad g'_{3,2} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 1 & 1 \\ 0 & -1 & 0 & 1 & 0 \end{pmatrix}, \quad g'_{3,3} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

The order of the stabilizer is  $480 = 2^5 \cdot 3 \cdot 5$ . Using GAP, we see that the group  $G'_3$  is isomorphic to  $\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathfrak{S}_5$ . Among the generators only  $g'_{3,3}$  has a non-trivial action on the cell. Let us consider the subgroup of  $G'_3$ , denoted  $H'_3$ , generated by  $g'_{3,1}$ ,  $g'_{3,2}$  and  $g'_{3,3}{}^2$ . The subgroup  $H'_3$  acts trivially on the cell, it is normal and isomorphic to  $\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathfrak{A}_5$ . Thus  $G'_3/H'_3 = \mathbb{Z}/2$  and, as  $\mathfrak{A}_n$  is perfect for  $n \geq 5$ , the abelianization of  $H'_3$  is isomorphic to  $\mathbb{Z}/2 \times \mathbb{Z}/2$ . By Lemma 8.3, we get

$$H_1(G'_3; \tilde{\mathbb{Z}}) = H_0(\mathbb{Z}/2; H_1(H'_3; \mathbb{Z})) = 0 \pmod{S_2}.$$

*Analyzing the cell  $\sigma'_4$ .* The cell  $\sigma'_4$  is given by the vectors labelled by 1, 2, 3, 7, 11, 12. A set of generators of its stabilizer is given by the following two matrices of  $GL_5(\mathbb{Z})$ , of respective order 6, 2:

$$g'_{4,1} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 1 \\ -1 & -1 & -1 & 1 & 1 \\ 0 & -1 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad g'_{4,2} = \begin{pmatrix} -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & -1 & -1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 & 0 \end{pmatrix}.$$

The order of the stabilizer is  $240 = 2^4 \cdot 3 \cdot 5$ . Denote  $G'_4$  this group, which is isomorphic to  $\mathbb{Z}/2 \times \mathfrak{S}_5$ . Only the generator  $g'_{4,1}$  acts non-trivially on the cell. Let  $H'_4$  be the subgroup generated by  $g'_{4,1}{}^2$  and  $g'_{4,2}$ . This subgroup is normal and isomorphic to  $\mathbb{Z}/2 \times \mathfrak{A}_5$ . So, as above, we get  $H_1(G'_4; \tilde{\mathbb{Z}}) = 0 \pmod{S_2}$ .

*Analyzing the cell  $\sigma'_5$ .* The cell  $\sigma'_5$  is given by the vectors labelled by 2, 3, 5, 7, 9, 10. A set of generators of its stabilizer, denoted  $G'_5$ , is given by the following three matrices of  $GL_5(\mathbb{Z})$ , of respective order 6, 6, 2:

$$g'_{5,1} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & -1 & 0 \\ -1 & -1 & 0 & 1 & 1 \\ 0 & -1 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad g'_{5,2} = \begin{pmatrix} -1 & -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 & 1 \end{pmatrix}, \quad g'_{5,3} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 & 1 \end{pmatrix}.$$

The stabilizer is of order  $1440 = 2^5 \cdot 3^2 \cdot 5$ . The group is isomorphic to  $\mathbb{Z}/2 \times \mathfrak{S}_6$ . Among the generators only  $g'_{5,3}$  has a non-trivial action. The subgroup generated by  $g'_{5,1}$  and  $g'_{5,2}$  is normal, it acts trivially on  $\sigma'_5$  and is isomorphic to  $\mathbb{Z}/2 \times \mathfrak{A}_6$ . So we get  $H_1(G'_5; \tilde{\mathbb{Z}}) = 0 \pmod{S_2}$ .

- Computation of  $E_{4,2}^1$ . The set  $\Sigma_4^\star$  consists of two cells contained (up to equivalence) in  $\sigma(P_5^2)$ . We will denote those cells by  $\tau'_i$  with  $i = 1, 2$ .

*Analyzing the cell  $\tau'_1$  and the cell  $\tau'_2$ .* The cell  $\tau'_1$  is given by the vectors labelled by 1, 2, 3, 4 and 8 in  $m(P_5^2)$ . A set of generators of its stabilizer is



given by the following three matrices of  $GL_5(\mathbb{Z})$ , of respective order 2, 6, 2:

$$t_{1,1} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad t_{1,2} = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad t_{1,3} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

The order of the stabilizer is  $3840 = 2^8 \cdot 3 \cdot 5$ . Furthermore, only  $t_{1,2}$  has a non-trivial action on the cell.

The cell  $\tau'_2$  is given by the vectors 1, 2, 7, 11 and 12. A set of generators of its stabilizer is given by the following three matrices of  $GL_5(\mathbb{Z})$ , of respective order 6, 4, 2:

$$t_{2,1} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 1 \\ 1 & -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 \\ 1 & 0 & 1 & -1 & 0 \end{pmatrix}, \quad t_{2,2} = \begin{pmatrix} -1 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & -1 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 1 \\ 0 & 1 & 0 & -1 & 0 \end{pmatrix}, \quad t_{2,3} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 1 & 0 \end{pmatrix}.$$

The order of the stabilizer is  $3840 = 2^8 \cdot 3 \cdot 5$ .

We need to analyze the first and second homology groups of the stabilizers of  $\tau'_1$  and  $\tau'_2$ . Using GAP, it is possible to show that these two stabilizers are isomorphic. Set

$$h_{1,1} = t_{1,2}^{-2} t_{1,1} t_{1,2} t_{1,1} t_{1,2}^{-1} t_{1,1} t_{1,2}^2 t_{1,1} t_{1,2} t_{1,1} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$h_{1,2} = t_{1,1} t_{1,3} t_{1,2}^{-1} t_{1,1} t_{1,2}^{-1} t_{1,1} t_{1,2} t_{1,1} = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix},$$

$$h_{2,1} = t_{2,3} t_{2,1} t_{2,2}^{-1} t_{2,1}^2 t_{2,2}^2 t_{2,1} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 \\ 0 & 1 & 1 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix},$$

$$h_{2,2} = t_{2,3} t_{2,1}^{-1} t_{2,3} t_{2,2}^{-1} t_{2,1}^2 = \begin{pmatrix} 0 & -1 & -1 & 1 & 0 \\ 1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 & 0 \end{pmatrix}.$$

Then the group generated by  $h_{1,1}$  and  $h_{1,2}$  (resp.  $h_{2,1}$  and  $h_{2,2}$ ) is isomorphic to  $Stab(\tau'_1)$  (resp.  $Stab(\tau'_2)$ ). We can check that the mapping sending  $h_{1,1}$  (resp.  $h_{1,2}$ ) to  $h_{2,1}$  (resp.  $h_{2,2}$ ) defines a group isomorphism. Hence it suffices to consider  $\tau'_1$ . Let  $H$  be the subgroup generated by  $t_{1,1}$ ,  $t_{1,3}$ ,  $(t_{1,1} t_{1,2})^2$  and  $(t_{1,3} t_{1,2})^2$ . Then  $H$  is normal, of order 1920 and it acts trivially on  $\tau'_1$ . Using

GAP, we can check that its abelianization is isomorphic to  $\mathbb{Z}/2$ . Using a composition series for  $H$ , we get a short exact sequence

$$0 \rightarrow (\mathbb{Z}/2)^5 \rightarrow H \rightarrow \mathfrak{A}_5 \rightarrow 1.$$

The homology of  $(\mathbb{Z}/2)^5$  is trivial modulo  $\mathcal{S}_2$  except

$$H_0((\mathbb{Z}/2)^5, \tilde{\mathbb{Z}}) = H_0((\mathbb{Z}/2)^5, \mathbb{Z}) = \mathbb{Z}.$$

But as  $\mathfrak{A}_5$  is simple, we deduce that it acts trivially on  $H_0((\mathbb{Z}/2)^5, \tilde{\mathbb{Z}})$ . Since  $H_i(\mathfrak{A}_5, \mathbb{Z})$  lies in  $\mathcal{S}_2$  for  $i = 1, 2$  [32], we deduce that

$$H_i(\mathfrak{A}_5, H_j((\mathbb{Z}/2)^5, \tilde{\mathbb{Z}})) = 0 \pmod{\mathcal{S}_2}, \text{ with } i + j = 1, 2.$$

Using the LHS spectral sequence associated to the above exact sequence, we get  $H_i(H, \tilde{\mathbb{Z}}) = 0$  modulo  $\mathcal{S}_2$  and by Lemma 8.3,  $H_i(\text{Stab}(\tau'_1), \tilde{\mathbb{Z}}) = 0$  mod  $\mathcal{S}_2$  for  $i = 1, 2$ .

□

**Lemma 8.6.** *The terms  $E_{5,1}^1$  and  $E_{6,0}^1$  of the equivariant spectral sequence associated to  $\Gamma = GL_6(\mathbb{Z})$  are zero modulo  $\mathcal{S}_2$ .*

*Proof.* The claim that  $E_{6,0}^1$  is zero modulo  $\mathcal{S}_2$  is again a consequence of the fact that none of the cells of  $\Sigma_6^*(GL_6(\mathbb{Z}))$  has its orientation preserved by the action of its stabilizer. It remains to show that  $E_{5,1}^1$  is zero modulo  $\mathcal{S}_2$ .

From our computations (cf. Fig.1), we know that  $\Sigma_5^*(GL_6(\mathbb{Z}))$  has three cell representatives which can be chosen inside  $\sigma(P_6^1)$ . We will denote these three cells by  $\tau_i$  ( $i = 1, 2, 3$ ). Here is the ordered list of minimal vectors of  $P_6^1$  that we shall use:

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1	-1	0	0	-1	0	0	0	-1	0	0	0	0	-1	0	1	1	0
0	1	1	-1	0	0	0	-1	0	0	0	0	-1	0	0	1	0	1
0	0	0	1	1	1	-1	0	0	0	0	-1	0	0	0	-1	0	0
0	0	0	0	0	0	1	1	1	1	-1	0	0	0	0	-1	-1	-1
0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	-1	-1	-1
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1

19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
1	1	0	1	0	1	0	1	0	1	0	1	0	0	1	0	0	1
1	0	1	1	0	0	1	0	1	1	0	0	1	0	0	1	0	1
0	-1	-1	-1	0	0	0	-1	-1	-1	0	0	0	-1	-1	-1	0	-1
-1	0	0	0	0	0	0	-1	-1	-1	-1	-1	-1	0	0	0	0	-1
-1	-1	-1	-1	-1	-1	-1	0	0	0	0	0	0	0	0	0	0	-1
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2

Analyzing the cell  $\tau_1$ . The cell  $\tau_1$  can be represented by the vectors 1, 15, 24, 25, 31, 34. Set  $G_1 = \text{Stab}_\Gamma(\tau_1)$ . A set of generators of  $G_1$  consists of the following

four matrices of  $GL_6(\mathbb{Z})$ , of respective order 4, 6, 4, 2:

$$g_{1,1} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -2 & -2 & -1 & -2 \\ 0 & -1 & 0 & 0 & 1 & 2 \end{pmatrix}, g_{1,2} = \begin{pmatrix} 1 & 1 & -1 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & -1 & -1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & -1 & -1 & -2 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$g_{1,3} = \begin{pmatrix} 0 & -1 & 1 & 0 & 0 & 1 \\ 1 & 0 & -1 & -2 & 0 & -1 \\ -1 & -1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 2 & 0 & 0 & -1 & -2 \\ 1 & -1 & -1 & -2 & 0 & 0 \end{pmatrix}, g_{1,4} = \begin{pmatrix} -1 & -2 & 0 & 0 & 0 & 2 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

The order of  $G_1$  is  $46080 = 2^{10} \cdot 3^2 \cdot 5$ . Only  $g_{1,3}$  has a non-trivial action on the cell. Consider the subgroup  $H_1$  of  $G_1$  generated by  $g_{1,1}, g_{1,2}, g_{1,4}$  and  $(g_{1,1}g_{1,3})^2$ . Then by construction, this subgroup acts trivially on the cell. Using GAP, we can check that  $G_1$  is isomorphic to  $GM_6(\mathbb{Z})$ , the subgroup of monomial matrices of  $GL_6(\mathbb{Z})$  (semi-direct product of  $\mathfrak{S}_6$  and  $\{\pm 1\}^6$ ), and  $H_1$  is normal, isomorphic to the semi-direct product of  $\mathfrak{A}_6$  and  $\{\pm 1\}^6$ . Thus the quotient  $G_1/H_1$  is isomorphic to  $\mathbb{Z}/2$ . Then, by the computation of the abelianization of semi-direct products, we get that  $H_1/[H_1, H_1] \cong \mathbb{Z}/2$ . We deduce that  $H_1(H_1, \tilde{\mathbb{Z}})$  lies in  $\mathcal{S}_2$  and by Lemma 8.3, we conclude that  $H_1(G_1, \tilde{\mathbb{Z}})$  lies in  $\mathcal{S}_2$ .

*Analyzing the cell  $\tau_2$ .* The cell  $\tau_2$  is given by the vectors 15, 24, 25, 28, 29, 34. Set  $G_2 = \text{Stab}_\Gamma(\tau_2)$ . A set of generators of  $G_2$  consists of the following four matrices of  $GL_6(\mathbb{Z})$ , of respective order 4, 6, 2, 2:

$$g_{2,1} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & -1 & -1 & 0 & -1 \\ -1 & 0 & -1 & 1 & -1 & 0 \end{pmatrix}, g_{2,2} = \begin{pmatrix} -1 & -1 & -1 & -1 & -1 & 0 \\ 0 & 1 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & -2 & -1 & -1 \end{pmatrix},$$

$$g_{2,3} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & -2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 2 & 2 & 0 & 1 & 0 \\ 0 & -2 & -2 & 0 & 0 & 1 \end{pmatrix}, g_{2,4} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ -1 & -1 & -2 & -1 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

The order of  $G_2$  is  $7680 = 2^9 \cdot 3 \cdot 5$ . Only the generators  $g_{2,3}$  and  $g_{2,4}$  have a non-trivial action on the cell. Denote by  $H_2$  the subgroup of  $G_2$  generated by  $g_{2,1}^2, g_{2,2}^2, g_{2,3}$  and  $g_{2,4}$ . Then  $H_2$  acts trivially on the cell and can be checked, using GAP, to be normal and of order 3840. Furthermore, its abelianization is isomorphic to  $\mathbb{Z}/2 \times \mathbb{Z}/2$ . Using Lemma 8.3 as above, we deduce that  $H_1(G_2, \tilde{\mathbb{Z}})$  lies in  $\mathcal{S}_2$ .

*Analyzing the cell  $\tau_3$ .* The cell  $\tau_3$  is given by the vectors 2, 15, 25, 28, 29, 34. Set  $G_3 = \text{Stab}_\Gamma(\tau_3)$ . A set of generators of  $G_3$  consists of the following three matrices

of  $GL_6(\mathbb{Z})$ , of respective order 6, 6, 2:

$$g_{3,1} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & -1 & -1 & -2 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & -1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & -1 & 0 & -2 \end{pmatrix}, g_{3,2} = \begin{pmatrix} -1 & -1 & -2 & 0 & -1 & 0 \\ 0 & 0 & 1 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & -1 & 0 \end{pmatrix},$$

$$g_{3,3} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ -1 & -1 & -2 & -1 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

The order of  $G_3$  is  $46080 = 2^{10} \cdot 3^2 \cdot 5$  and it is isomorphic to  $G_1$ . Only  $g_{3,3}$  has a trivial action on the cell. The subgroup of  $G_3$  generated by  $g_{3,1}^2, g_{3,2}^2, g_{3,1}g_{3,2}$  and  $g_{3,3}$  acts trivially on the cell, and using GAP, we can check that it is isomorphic to  $H_1$ . As a result, we conclude that  $H_1(G_3, \tilde{\mathbb{Z}})$  lies in  $\mathcal{S}_2$ . As all the terms of  $E_{5,1}^1$  lies in  $\mathcal{S}_2$ , the lemma is proved.  $\square$

**Lemma 8.7.** *The terms  $E_{5,2}^1, E_{6,1}^1$  and  $E_{7,0}^1$  of the equivariant spectral sequence associated to  $\Gamma = GL_6(\mathbb{Z})$  are zero modulo  $\mathcal{S}_5$ .*

*Proof.* Looking at the table of representatives for  $\Sigma_p^*(\Gamma)$  (cf. Figure 1), we see that there are three 5-cells, ten 6-cells and twenty-eight 7-cells. None of the 7-cells has its orientation preserved by its stabilizer. Thus  $H_0(Stab_\Gamma(\sigma), \tilde{\mathbb{Z}})$  lies in  $\mathcal{S}_2$  for all  $\sigma \in \Sigma_7^*(\Gamma)$ . Among the 6-cells, only one has a stabilizer with 7-torsion. It is the cell given by the minimal vectors

$$\begin{array}{ccccccc} 1 & 3 & 5 & 8 & 12 & 16 & 17 \\ \hline 1 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 \end{array}$$

from  $P_6^7$ . We will denote by  $G_0$  its stabilizer. It is generated by the following matrices:

$$g_{0,1} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & -1 & -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & -1 & -1 & -1 \\ 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}, g_{0,2} = \begin{pmatrix} -1 & 0 & -1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix},$$

$$g_{0,3} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

all of which have a non-trivial action on the cell. The order of  $G_0$  is  $10080 = 2^5 \times 3^2 \times 5 \times 7$ . From a composition series of  $G_0$ , we can deduce the exact sequence

$$1 \rightarrow H \rightarrow G_0 \rightarrow \mathbb{Z}/2 \rightarrow 1$$

where  $H \cong \mathfrak{A}_7 \times \mathbb{Z}/2$  and  $H$  is generated by  $g_{0,1}^2$ ,  $g_{0,2}^2$  and  $-\text{Id}$ . Hence the action of  $H$  on the cell is trivial. Furthermore, the quotient  $H/[H, H]$  is isomorphic to  $\mathbb{Z}/2$ . From the previous data, we deduce by a spectral sequence argument that  $H_1(G_0, \tilde{\mathbb{Z}}) = 0 \pmod{\mathcal{S}_2}$ . Finally, among the 5-cells, none of them has a stabilizer with 7-torsion. Lemma 8.7 follows.  $\square$

**Lemma 8.8.** *The terms  $E_{6,1}^1$  and  $E_{7,0}^1$  of the equivariant spectral sequence associated to  $\Gamma = GL_7(\mathbb{Z})$  are zero modulo  $\mathcal{S}_5$ .*

*Proof.* Looking at the table of representative for  $\Sigma_p^*(\Gamma)$  (cf. Figure 2), we see that there are twenty-eight 7-cells, none of them having its orientation preserved by the action of its stabilizer. As a result, we can deduce that  $E_{0,7}^1 = 0 \pmod{\mathcal{S}_2}$ . Among the six 6-cells, only three have a stabilizer of order divisible by 7. They are the ones to investigate.

1. The first cell is given by the following seven minimal vectors

1	2	3	4	5	6	7
1	1	1	1	1	2	1
1	1	1	1	2	1	1
-1	-1	-1	0	-1	-1	-1
-1	-1	0	-1	-1	-1	-1
-1	0	-1	-1	-1	-1	-1
0	-1	-1	-1	-1	-1	-1
2	2	2	2	2	2	2

of  $P_7^2$ . A set of generators for its stabilizer, that we will denote by  $G_1$ , consists of the following matrices

$$\begin{aligned}
g_{1,1} &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, & g_{1,2} &= \begin{pmatrix} -1 & -1 & -1 & -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 & 1 & 1 & 0 \end{pmatrix}, \\
g_{1,3} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, & g_{1,4} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \\
g_{1,5} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, & g_{1,6} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.
\end{aligned}$$

The group  $G_1$  is of order  $10080 = 2^5 \times 3^2 \times 5 \times 7$ . The generators  $g_{1,1}$ ,  $g_{1,2}$ ,  $g_{1,5}$  and  $g_{1,6}$  have a non-trivial action on the cell. A composition series of  $G_1$  is given by

$$1 \triangleleft \mathfrak{A}_7 \triangleleft \mathfrak{S}_7 \triangleleft G_1,$$

with  $G_1/\mathfrak{S}_7 \cong \mathbb{Z}/2$ . The group  $\mathfrak{A}_7$  is generated by  $(g_{1,1}g_{1,2})^2$  and  $g_{1,3}$ , and  $\mathfrak{S}_7$  is generated by  $(g_{1,1}g_{1,2})^2$ ,  $g_{1,3}$  and  $g_{1,1}$ . Using these generators, we deduce that the

action of  $\mathfrak{A}_7$  on the cell is trivial, while the one of  $\mathfrak{S}_7$  is not. There are two spectral sequences:

$$\begin{aligned} H_i(\mathbb{Z}/2; H_j(\mathfrak{S}_7; \tilde{\mathbb{Z}})) &\Longrightarrow H_{i+j}(G_1; \tilde{\mathbb{Z}}), \\ H_i(\mathbb{Z}/2; H_j(\mathfrak{A}_7; \tilde{\mathbb{Z}})) &\Longrightarrow H_{i+j}(\mathfrak{S}_7; \tilde{\mathbb{Z}}). \end{aligned}$$

The action of  $\mathfrak{A}_7$  is trivial and this group is perfect, so we get  $H_1(\mathfrak{S}_7; \tilde{\mathbb{Z}}) \cong \mathbb{Z}/2$ . We deduce that  $H_1(G_1; \tilde{\mathbb{Z}}) = 0 \pmod{\mathcal{S}_2}$ .

2. The second cell is given by the minimal vectors

$$\begin{array}{ccccccc} 1 & 2 & 3 & 5 & 8 & 12 & 23 \\ \hline -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & -1 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 3 & 1 & 1 & 1 & 1 & 1 & 0 \end{array}$$

from  $P_7^{12}$ . We will denote its stabilizer by  $G_2$ , which is of order  $645120 = 2^{11} \times 3^2 \times 5 \times 7$ . A set of generators for  $G_2$  is given by

$$\begin{aligned} g_{2,1} &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 & 0 & 1 & 0 \\ 0 & -1 & -1 & -1 & -1 & 1 & -1 \\ -1 & 0 & 0 & 0 & 0 & -3 & -1 \end{pmatrix}, & g_{2,2} &= \begin{pmatrix} -1 & -1 & -1 & 0 & -1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & -1 & -1 & -1 & 0 & -1 \\ 0 & 1 & 1 & -1 & 1 & 1 & 0 \end{pmatrix}, \\ g_{2,3} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

Only  $g_{2,2}$  has a non-trivial action on the cell. Using a composition series for  $G_2$ , we get the exact sequence

$$1 \rightarrow H \rightarrow G_2 \rightarrow \mathbb{Z}/2 \rightarrow 1$$

Using GAP, we can show that  $H$  is generated by  $g_{2,1}$ ,  $g_{2,3}$  and  $g_{2,2}^2$ . It follows that the action of  $H$  on the cell is trivial. We have the following spectral sequence

$${}'E_{i,j}^2 = H_i(\mathbb{Z}/2; H_j(H; \tilde{\mathbb{Z}})) \Longrightarrow H_{i+j}(G_2; \tilde{\mathbb{Z}}).$$

Furthermore, the group  $H/[H, H]$  is isomorphic to  $\mathbb{Z}/2$ . As a result, we get that  $H_1(G_2; \tilde{\mathbb{Z}}) = 0 \pmod{\mathcal{S}_2}$ .

3. The last cell is given by the minimal vectors

$$\begin{array}{ccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ \hline 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{array}$$

from  $P_7^{33}$ . We will denote its stabilizer by  $G_3$ . Its order is  $645120 = 2^{11} \times 3^2 \times 5 \times 7$ . The group  $G_3$  is spanned by the following six matrices:

$$\begin{aligned} g_{3,1} &= \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}, & g_{3,2} &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \\ g_{3,3} &= \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -2 & -2 & -2 & -2 & -2 & -1 \end{pmatrix}, & g_{3,4} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \\ g_{3,5} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, & g_{3,6} &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ -2 & -2 & -2 & -2 & -2 & -2 & -1 \end{pmatrix}. \end{aligned}$$

Only the generators  $g_{3,2}$  and  $g_{3,3}$  have a non-trivial action on the cell. The groups  $G_2$  and  $G_3$  turn out to be isomorphic. Hence the previous arguments will apply to  $G_3$ .

We have an exact sequence

$$1 \rightarrow H' \rightarrow G_3 \rightarrow \mathbb{Z}/2 \rightarrow 1,$$

where  $H'$  is generated by  $g_{3,1}$ ,  $g_{3,4}$ ,  $g_{3,5}$  and  $g_{3,6}$ , together with the product  $g_{3,2}g_{3,3}$ . As a result, the action of  $H'$  on the cell is trivial. Moreover, the quotient  $H'/[H', H']$  is isomorphic to  $\mathbb{Z}/2$  and we deduce that  $H_1(G_3; \tilde{\mathbb{Z}})$  lies in  $\mathcal{S}_2$ .  $\square$

Now we are ready to complete the proof of Theorem 8.1:

*Proof.* (of Theorem 8.1).

- In rank 2, we shall prove that  $H_4(GL_2(\mathbb{Z}), St_2)$  and  $H_5(GL_2(\mathbb{Z}), St_2)$  lie in  $\mathcal{S}_2$ . Let  $\tilde{\mathbb{Z}}$  be the orientation module of the symmetric space  $X_2$  and let  $\hat{H}^*(GL_2(\mathbb{Z}), \tilde{\mathbb{Z}})$  be the Farrell cohomology of  $GL_2(\mathbb{Z})$  [11]. From (5) in §7.2 it follows that

$$H_4(GL_2(\mathbb{Z}), St_2) \cong \hat{H}^{-3}(GL_2(\mathbb{Z}), \tilde{\mathbb{Z}}),$$

and

$$H_5(GL_2(\mathbb{Z}), St_2) \cong \hat{H}^{-4}(GL_2(\mathbb{Z}), \tilde{\mathbb{Z}}).$$

As to the first claim, since the only 3-group contained in  $GL_2(\mathbb{Z})$  is, up to conjugation,  $\mathbb{Z}/3$ , we get

$$\hat{H}^{-5}(GL_2(\mathbb{Z}), \tilde{\mathbb{Z}}) \subset \hat{H}^{-5}(\mathbb{Z}/3, \tilde{\mathbb{Z}}) = 0.$$

As to the second claim,  $GL_2(\mathbb{Z})$  is an extension

$$1 \longrightarrow SL_2(\mathbb{Z}) \longrightarrow GL_2(\mathbb{Z}) \longrightarrow \Delta \longrightarrow 1$$

with  $\Delta = \mathbb{Z}/2$ , and  $SL_2(\mathbb{Z})$  is the amalgamated product of  $\mathbb{Z}/4$  and  $\mathbb{Z}/6$  along  $\mathbb{Z}/2$  (see [29]). Therefore

$$\hat{H}^{-4}(SL_2(\mathbb{Z}), \mathbb{Z}) = \hat{H}^{-4}(GL_2(\mathbb{Z}), \mathbb{Z}) \oplus \hat{H}^{-4}(GL_2(\mathbb{Z}), \tilde{\mathbb{Z}})$$

(see §7.2) and, modulo  $\mathcal{S}_2$ ,

$$\hat{H}^{-4}(SL_2(\mathbb{Z}), \mathbb{Z}) = \mathbb{Z}/3.$$

Let  $\beta$  be a generator of  $\hat{H}^{-2}(SL_2(\mathbb{Z}), \mathbb{Z}) = \mathbb{Z}/3$ . Since  $\hat{H}^{-4}(SL_2(\mathbb{Z}), \mathbb{Z})$  is spanned by  $\beta^2$ , the action of  $\Delta$  on this group is trivial. Therefore

$$\hat{H}^{-4}(GL_2(\mathbb{Z}), \mathbb{Z}) = \mathbb{Z}/3$$

modulo  $\mathcal{S}_2$  and  $\hat{H}^{-4}(SL_2(\mathbb{Z}), \tilde{\mathbb{Z}})$  lies in  $\mathcal{S}_2$ .

- In rank 3, we know from [31], Thm. 5(iii), that  $H_3(GL_3(\mathbb{Z}), St_3) \cong \mathbb{Z}$  modulo  $\mathcal{S}_2$ . Moreover, from op.cit., Thm. 5(ii), we have

$$H_4(GL_3(\mathbb{Z}), St_3) \cong \hat{H}^{-2}(GL_3(\mathbb{Z}), \mathbb{Z}).$$

where  $\hat{H}^*$  denotes the Farrell cohomology, and from op.cit., Corollary (i) on p.9, we know that

$$\hat{H}^{-2}(GL_3(\mathbb{Z}), \mathbb{Z}) \text{ lies in } \mathcal{S}_2.$$

- In rank 4, we know from [19], Lemma 3.3, that  $\Sigma_p^*$  is empty when  $p < 3$ , hence  $E_{p,q}^1 = 0$  when  $p < 3$ . We proved in Lemma 8.4 that  $E_{p,q}^2$  lies in  $\mathcal{S}_2$  when  $q > 0$  and  $p + q = 5$  or  $p + q = 6$ . According to [19], Proposition 3.1,  $E_{0,6}^2 \cong \mathbb{Z}$  and  $E_{0,5}^2 = 0$  modulo  $\mathcal{S}_2$ . Therefore, modulo  $\mathcal{S}_2$ ,  $H_3(GL_4(\mathbb{Z}), St_4) \cong \mathbb{Z}$  and  $H_2(GL_4(\mathbb{Z}), St_4) = 0$ .
- In rank 5, we see in Fig.1 that  $\Sigma_p^*$  is empty when  $p < 4$ . Therefore

$$E_{p,q}^1 = \bigoplus_{\sigma \in \Sigma_p^*} H_q(\Gamma_\sigma, \tilde{\mathbb{Z}})$$

vanishes if  $p < 4$ . On the other hand, since  $\Sigma_5(GL_5(\mathbb{Z}))$  is empty, the group  $E_{5,0}^1$  lies in  $\mathcal{S}_2$ . We proved in Lemma 8.5 that  $E_{4,1}^1$  is in  $\mathcal{S}_2$ . Therefore  $E_{p,q}^1$  lies in  $\mathcal{S}_2$  when  $N = 5$  and  $p + q = 5$ , hence  $H_1(GL_5(\mathbb{Z}), St_5)$  lies in  $\mathcal{S}_2$ .

Similarly, we know from Lemma 8.5 that  $E_{p,q}^1$  is in  $\mathcal{S}_2$  when  $p \geq 4$  and  $p + q = 6$ . Therefore  $H_2(GL_5(\mathbb{Z}), St_5)$  lies in  $\mathcal{S}_2$ .



- In rank 6,  $\Sigma_p^\star$  is empty and  $E_{p,q}^1 = 0$  when  $p < 5$ .  
 Lemma 8.6 shows that also  $E_{p,q}^1 = 0$  when  $p \geq 5$  and  $p + q = 6$ .  
 Therefore  $H_1(GL_6(\mathbb{Z}), St_6)$  lies in  $\mathcal{S}_2$ .  
 When  $p + q = 7$  and  $p \geq 5$ , Lemma 8.7 shows that  $E_{p,q}^1$  lies in  $\mathcal{S}_5$ . Therefore  $H_2(GL_6(\mathbb{Z}), St_6)$  is in  $\mathcal{S}_5$ .
- In rank 7,  $\Sigma_p^\star$  is empty and  $E_{p,q}^1 = 0$  when  $p < 6$  (see Fig.1). On the other hand, we know from Lemma 8.8 that  $E_{6,1}^1$  and  $E_{7,0}^1$  lie in  $\mathcal{S}_5$ . Therefore  $H_1(GL_7(\mathbb{Z}), St_7)$  lies in  $\mathcal{S}_5$ .

□

## 9. APPLICATION TO K-THEORY

The homology of the general linear group with coefficients in the Steinberg module can also be used to compute the  $K$ -theory of  $\mathbb{Z}$ . Let  $P(\mathbb{Z})$  (resp.  $P_N(\mathbb{Z})$ ) be the exact category of free  $\mathbb{Z}$ -modules of finite rank (resp. of rank at most  $N$ ), let  $Q$  (resp.  $Q_N$ ) be the category obtained from  $P(\mathbb{Z})$  (resp.  $P_N(\mathbb{Z})$ ) by the  $Q$ -construction [25], and let  $BQ$  (resp.  $BQ_N$ ) be its classifying space. A definition of higher  $K$ -theory [25] is

$$K_m(\mathbb{Z}) = \pi_{m+1}(BQ), \quad m \geq 0.$$

On the other hand, Quillen proved in [26] that there are long exact sequences

$$(10) \quad \cdots \rightarrow H_m(BQ_{N-1}, \mathbb{Z}) \rightarrow H_m(BQ_N, \mathbb{Z}) \rightarrow H_{m-N}(GL_N(\mathbb{Z}), St_N) \\ \rightarrow H_{m-1}(BQ_{N-1}, \mathbb{Z}) \rightarrow \cdots,$$

and, according to Lee and Szczarba [19],  $H_0(GL_N(\mathbb{Z}), St_N) = 0$  when  $N \geq 1$ . Therefore we can compute  $K_m(\mathbb{Z})$  if we understand the Hurewicz map

$$h_m : K_m(\mathbb{Z}) \rightarrow H_{m+1}(BQ, \mathbb{Z})$$

and if we compute the groups  $H_{m+1-N}(GL_N(\mathbb{Z}), St_N)$  for all  $N \leq m$ .

**9.1. On the Hurewicz morphism.** Let  $BQ = BQP(\mathbb{Z})$  be the classifying space of Quillen's  $Q$ -construction on the exact category  $P(\mathbb{Z})$  of finitely generated free  $\mathbb{Z}$ -modules. By definition, for every integer  $m \geq 1$ ,

$$K_{m-1}(\mathbb{Z}) = \pi_m(BQ).$$

In this section we shall be interested in the kernel  $C_m$  of the Hurewicz map

$$h_m : \pi_m(BQ) \rightarrow H_m(BQ),$$

where  $H_m(X)$  stands for  $H_m(X; \mathbb{Z})$ .

**Proposition 9.1.** *The groups  $C_6$  and  $C_7$  lie in  $\mathcal{S}_2$ , and  $C_8$  lies in  $\mathcal{S}_5$ .*

9.2. **Proof.** To prove this proposition, we use a morphism of spectra

$$K(\mathcal{E}) \rightarrow K(\mathbb{Z})$$

introduced by Rognes in [27], § 4, where  $\mathcal{E}$  is the category of finite sets. At level zero this morphism is the map

$$\mathbb{Z} \times B\Sigma_\infty^+ \rightarrow \mathbb{Z} \times BGL(\mathbb{Z})^+,$$

where  $\Sigma_\infty$  is the infinite symmetric group,  $GL(\mathbb{Z})$  is the infinite general linear group over  $\mathbb{Z}$ , and  $( )^+$  is the +-construction of Quillen. Let  $\phi$  be the fiber of that map and consider the fibration

$$(11) \quad \begin{array}{ccc} B & \longrightarrow & BQ \\ & & \downarrow \\ & & \phi \end{array}$$

where  $B$  is the first level of  $K(\mathcal{E})$ . When  $m \geq 1$ , the group

$$\pi_{m+1}(B) = \pi_m(\mathbb{Z} \times B\Sigma_\infty^+) = \pi_m^s$$

is the  $m$ -th homotopy group of spheres by the Barratt/Priddy/Quillen theorem. The map

$$\pi_m^s \rightarrow K_m(\mathbb{Z})$$

is an isomorphism modulo  $\mathcal{S}_2$  when  $m \leq 4$ . Therefore the long exact sequence deduced from (11)

$$(12) \quad \cdots \rightarrow \pi_m^s \rightarrow K_m(\mathbb{Z}) \rightarrow \pi_{m+1}(\phi) \rightarrow \pi_{m-1}^s \rightarrow \cdots$$

implies that  $\pi_m(\phi)$  lies in  $\mathcal{S}_2$  when  $m \leq 5$ .

From [1] Theorem 1.5, which remains valid modulo a Serre class, it follows that the kernel of the Hurewicz map

$$\pi_m(\phi) \rightarrow H_m(\phi)$$

lies in  $\mathcal{S}_2$  when  $m = 6, 7$  or  $8$ . On the other hand,  $\pi_5^s$  and  $\pi_6^s$  lie in  $\mathcal{S}_2$ , while  $\pi_7^s$  lies in  $\mathcal{S}_5$ . Using (12), this implies that the kernel of the map

$$K_{m-1}(\mathbb{Z}) \rightarrow \pi_m(\phi)$$

lies in  $\mathcal{S}_2$  (resp  $\mathcal{S}_5$ ) when  $m = 6$  or  $7$  (resp.  $8$ ). The commutative diagram

$$\begin{array}{ccc} K_{m-1}(\mathbb{Z}) & \longrightarrow & \pi_m(\phi) \\ h_m \downarrow & & \downarrow \\ H_m(BQ) & \longrightarrow & H_m(\phi) \end{array}$$

concludes the proof.

**Theorem 9.2.** *We have  $K_5(\mathbb{Z}) \cong \mathbb{Z} \bmod \mathcal{S}_2$ ,  $K_6(\mathbb{Z})$  lies in  $\mathcal{S}_2$ , and  $K_7(\mathbb{Z})$  lies in  $\mathcal{S}_5$ .*

*Proof.* • First we compute  $K_5(\mathbb{Z})$ . From Theorem 8.1, (i) and (ii), we know that, modulo  $\mathcal{S}_2$ , the group  $H_{6-N}(GL_N(\mathbb{Z}), St_N)$  vanishes when  $N \leq 5$  and  $N \neq 3$ , and that  $H_3(GL_3(\mathbb{Z}), St_3) \cong \mathbb{Z}$ .

The exact sequence (10) for  $N = 2$  reads

$$\begin{aligned} H_6(BQ_1, \mathbb{Z}) &\rightarrow H_6(BQ_2, \mathbb{Z}) \rightarrow H_4(GL_2(\mathbb{Z}), St_2) \rightarrow \\ &\rightarrow H_5(BQ_1, \mathbb{Z}) \rightarrow H_5(BQ_2, \mathbb{Z}) \rightarrow H_3(GL_2(\mathbb{Z}), St_2) \end{aligned}$$

Since  $H_m(BQ_1, \mathbb{Z}) = 0$  when  $m > 0$ ,  $H_3(GL_2(\mathbb{Z}), St_2) \cong \hat{H}^{-3}(GL_2(\mathbb{Z}), St_2)$  is finite, and  $H_4(GL_2(\mathbb{Z}), St_2)$  lies in  $\mathcal{S}_2$ , we conclude that  $H_6(BQ_2, \mathbb{Z})$  lies in  $\mathcal{S}_2$  and  $H_5(BQ_2, \mathbb{Z})$  is finite.

The exact sequence (10) for  $N = 3$  gives

$$H_6(BQ_2, \mathbb{Z}) \rightarrow H_6(BQ_3, \mathbb{Z}) \rightarrow H_3(GL_3(\mathbb{Z}), St_3) \rightarrow H_5(BQ_2, \mathbb{Z}),$$

therefore  $H_6(BQ_3, \mathbb{Z}) \cong \mathbb{Z}$  modulo  $\mathcal{S}_2$ .

On the other hand, we deduce from (10) with  $N = 5, 6, 7$  and from Theorem 8.1 that, modulo  $\mathcal{S}_2$ ,

$$H_6(BQ, \mathbb{Z}) \cong H_6(BQ_5, \mathbb{Z}) \cong H_6(BQ_4, \mathbb{Z}),$$

as  $H_1(GL_5(\mathbb{Z}), St_5)$  and  $H_1(GL_6(\mathbb{Z}), St_6)$  are finite.

Now consider the exact sequence (10) for  $N = 4$ :

$$H_3(GL_4(\mathbb{Z}), St_4) \xrightarrow{\alpha} H_6(BQ_3, \mathbb{Z}) \rightarrow H_6(BQ_4, \mathbb{Z}) \rightarrow H_2(GL_4(\mathbb{Z}), St_4),$$

where the last group is in  $\mathcal{S}_2$  by Theorem 8.1(ii).

If  $\alpha$  were not zero modulo  $\mathcal{S}_2$  then we would conclude that  $H_6(BQ, \mathbb{Z}) \cong H_6(BQ_4, \mathbb{Z})$  is finite. But this is impossible since  $K_5(\mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Q} = \mathbb{Q}$  (Borel) and the Hurewicz map

$$h_6 : K_5(\mathbb{Z}) \rightarrow H_6(BQ, \mathbb{Z})$$

has finite kernel.

Therefore  $\alpha = 0$  modulo  $\mathcal{S}_2$ , and

$$H_6(BQ, \mathbb{Z}) \cong H_6(BQ_3, \mathbb{Z}) \cong \mathbb{Z}$$

modulo  $\mathcal{S}_2$ . The Hurewicz map  $h_6$  is an isomorphism modulo torsion, and its kernel  $C_6$  lies in  $\mathcal{S}_2$  (Proposition 9.1). Therefore  $K_5(\mathbb{Z})$  is the direct sum of  $\mathbb{Z}$  and a finite 2-group.

- Next, we compute  $K_6(\mathbb{Z})$ .

From Theorem 8.1(ii), we know that  $H_{7-N}(GL_N(\mathbb{Z}), St_N)$  lies in  $\mathcal{S}_2$  when  $N \neq 4$ , and, according to Theorem 8.1(i), we have  $H_3(GL_4(\mathbb{Z}), St_4) \cong \mathbb{Z}$  modulo  $\mathcal{S}_2$ .

From the exact sequence (10) for  $N = 2, 3$ , we conclude from (9) that  $H_7(BQ_3, \mathbb{Z})$  lies in  $\mathcal{S}_2$ . The exact sequence for  $N = 5$  gives

$$\begin{aligned} H_7(BQ_3, \mathbb{Z}) &\rightarrow H_7(BQ_4, \mathbb{Z}) \rightarrow H_3(GL_4(\mathbb{Z}), St_4) \xrightarrow{\alpha} \\ &\xrightarrow{\alpha} H_6(BQ_3, \mathbb{Z}) \rightarrow H_6(BQ_4, \mathbb{Z}) \end{aligned}$$

Since  $\alpha = 0$  modulo  $\mathcal{S}_2$ , we get  $H_7(BQ_4, \mathbb{Z}) \cong H_3(GL_4(\mathbb{Z}), St_4) \cong \mathbb{Z}$  modulo  $\mathcal{S}_2$ . Using the exact sequence (10) for  $N = 5$  and 6, we conclude that  $H_7(BQ, \mathbb{Z}) \cong \mathbb{Z}$  modulo  $\mathcal{S}_2$ .

Since  $K_6(\mathbb{Z})$  is finite (Borel) and the kernel  $C_7$  of the Hurewicz map

$$h_7 : K_6(\mathbb{Z}) \rightarrow H_7(BQ, \mathbb{Z})$$

lies in  $\mathcal{S}_2$ , we get that  $K_6(\mathbb{Z})$  is a finite 2-group.

- Finally, we show that  $K_7(\mathbb{Z})$  lies in  $\mathcal{S}_5$ . From Lemma 8.5 we deduce that the groups  $E_{pq}^r$  for  $GL_N(\mathbb{Z})$ ,  $N \leq 5$ ,  $r \geq 1$ , lie in  $\mathcal{S}_5$  when  $q > 0$ .

Using Theorem 8.1 and Lemma 8.8, we conclude that  $H_{8-N}(GL_N(\mathbb{Z}), St_N)$  lies in  $\mathcal{S}_5$  when  $N \leq 7$ . This implies that  $H_8(BQ, \mathbb{Z})$  is in  $\mathcal{S}_5$  and, since the kernel  $C_8$  of  $h_8$  lies in  $\mathcal{S}_5$  (Proposition 9.1), we conclude that  $K_7(\mathbb{Z})$  has no  $p$ -torsion with  $p > 5$ .

□

**Remark 9.3.** These three are already known:  $K_5(\mathbb{Z}) \cong \mathbb{Z}$ ,  $K_6(\mathbb{Z}) = 0$  and  $K_7(\mathbb{Z}) \cong \mathbb{Z}/240$  (see [35]). The group  $K_8(\mathbb{Z})$  still remains unknown.

## REFERENCES

- [1] **Arlettaz, D.**; *The Hurewicz homomorphism in algebraic K-theory*, *J. Pure Appl. Algebra* **71** (1991), p. 1–12.
- [2] **Barnes, E.S.**; *The complete enumeration of extreme senary forms*, *Phil. Trans. Roy. Soc. London* **249-A** (1957), 461–506.
- [3] **Batut, C.**; *Classification of quintic eutactic forms*, *Math. Comput.* **70**, no.233 (2001), 395–417.
- [4] **Bavard, C.**; *Classes minimales de réseaux et rétractions géométriques équivariantes dans les espaces symétriques*, *J. London Math. Soc. (2)* **64**, no. 2 (2001), 275–286.
- [5] **Borel, A.**; *Stable real cohomology of arithmetic groups*, *Ann. Sci. École Norm. Sup. (4)* **7** (1974), 235–272.
- [6] **Borel, A.; Serre, J-P.**; *Corners and arithmetic groups*, *Comment. Math. Helv.* **48** (1973), 436–491.
- [7] **Brown, K.**; *Cohomology of Groups*, *Graduate Texts in Mathematics* **87**, Springer-Verlag, New York, 1994.
- [8] **Dumas, J.-G.; Elbaz-Vincent, Ph.; Giorgi, P.; Urbanska, A.**; *Parallel Computation of the Rank of Large Sparse Matrices from Algebraic K-theory.*, PASCO 2007: Parallel Symbolic Computation '07, 26–27 July, Waterloo, Canada. Proceedings of the ACM (2007), 43–52.
- [9] **Dutour Sikiric, M.; Schürmann, A.; Vallentin, F.**; *Classification of eight dimensional perfect forms*, *Electron. Res. Announc. Amer. Math. Soc.* **13** (2007), 21–32.
- [10] **Elbaz-Vincent, Ph.; Gangl, H.; Soulé, C.**; *Quelques calculs de la cohomologie de  $GL_N(\mathbb{Z})$  et de la K-théorie de  $\mathbb{Z}$* , *C. R. Math. Acad. Sci. Paris* **335** (2002), no. 4, 321–324.
- [11] **Farrell, F. T.**; *An extension of Tate cohomology to a class of infinite groups*, *J. Pure Appl. Algebra* **10**, no. 2 (1977/78), 153–161.
- [12] **The GAP Group**; GAP – Groups, Algorithms, and Programming, Version 4.4.12; 2008. (<http://www.gap-system.org>).
- [13] **Gauss, C.G.**; *Untersuchungen über die Eigenschaften der positiven ternären quadratischen Formen von Ludwig August Seeber*, *J. Reine Angew. Math.* **20** (1840) 312–320.
- [14] **Harder, G.**; *Die Kohomologie S-arithmetischer Gruppen über Funktionenkörpern*, *Invent. Math.* **42** (1977), 135–175.
- [15] **Jaquet, D.-O.**; *Énumération complète des classes de formes parfaites en dimension 7*, Thèse de doctorat, Université de Neuchâtel (1991).
- [16] **Korkine, A.N. and E.I. Zolotarev**; *Sur les formes quadratiques positives*, *Math. Ann.* **11** (1877) 242–292.
- [17] **Korkine, A.N. and E.I. Zolotarev**; *Sur les formes quadratiques positives quaternaires*, *Math. Ann.* **5** (1872) 581–583.
- [18] **Lagrange, J.-L.**; *Démonstration d'un théorème d'arithmétique*, *Nouv. Mém. Acad. Berlin* (1770), in *Oeuvres de Lagrange III*, 189–201.
- [19] **Lee, R.; Szczarba, R. H.**; *On the torsion in  $K_4(\mathbb{Z})$  and  $K_5(\mathbb{Z})$* , *Duke Math. J.* **45** (1978), 101–129.
- [20] **Martinet, J.**; *Perfect Lattices in Euclidean Spaces*, Springer-Verlag, Grundlehren der Mathematischen Wissenschaften **327**, Heidelberg, 2003.
- [21] **Morita, S.**; *Cohomological structure of the mapping class group and beyond*. in: Problems on mapping class groups and related topics, *Proc. Sympos. Pure Math.*, **74**, Amer. Math. Soc., Providence, RI, 2006, 329–354.
- [22] **The PARI-Group**; PARI/GP, versions 2.1–2.4, Bordeaux, <http://pari.math.u-bordeaux.fr/>.
- [23] **PFPK**: A C library for computing Voronoi complexes, version 1.0.0, 2009.
- [24] **Plesken, W.; Souvignier, B.**; *Computing isometries of lattices*, *J. Symb. Comput.* **24** (1997), 327–334.
- [25] **Quillen, D.**; *Higher algebraic K-theory. I. Algebraic K-theory, I: Higher K-theories* (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), pp. 85–147. *Lecture Notes in Math.*, **341**, Springer, Berlin 1973.

- [26] **Quillen, D.;** *On the cohomology and K-theory of the general linear groups over a finite field.* *Ann. of Math. (2)* **96** (1972), 552–586.
- [27] **Rognes, J.;** *A spectrum level rank filtration in algebraic K-theory.* *Topology* **31** (1992), no. 4, 813–845.
- [28] **Schürmann, A.;** *Enumerating perfect forms.* arXiv:0901.1587 (math.NT).
- [29] **Serre, J-P.;** *Trees*, Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003.
- [30] **Soulé, C.;** *On the 3-torsion in  $K_4(\mathbb{Z})$ .* *Topology* **39**, no.2, (2000), 259–265.
- [31] **Soulé, C.;** *The cohomology of  $SL_3(\mathbb{Z})$ .* *Topology* **17**, no. 1 (1978), 1–22.
- [32] **Suzuki, M.;** *Group Theory I*, Springer-Verlag 1982.
- [33] **Urbanska-Marszalek, A.;** *Hybrid and adaptive algorithms in exact linear algebra*, Thesis, Université Joseph Fourier Grenoble, 2010.
- [34] **Voronoi, G.;** *Nouvelles applications des paramètres continus à la théorie des formes quadratiques I*, *Crelle* **133** (1907), 97–178.
- [35] **Weibel, C.;** *The algebraic K-theory of rings of integers in local and global fields.*

INSTITUT FOURIER, UMR 5582 (CNRS-UNIVERSITÉ DE GRENOBLE), 100 RUE DES MATHÉMATIQUES,  
DOMAINE UNIVERSITAIRE, BP 74, 38402 SAINT MARTIN D’HÈRES (FRANCE)  
*E-mail address:* Philippe.Elbaz-Vincent@ujf-grenoble.fr

DEPARTMENT OF MATHEMATICAL SCIENCES, SOUTH ROAD, UNIVERSITY OF DURHAM (UNITED KINGDOM)  
*E-mail address:* herbert.gangl@durham.ac.uk

IHES, LE BOIS-MARIE 35, ROUTE DE CHARTRES, 91440 BURES-SUR-YVETTE (FRANCE)  
*E-mail address:* soule@ihes.fr