



# Scattering theory for graphs isomorphic to a homogeneous tree at infinity

Yves Colin de Verdière, Francoise Truc

## ► To cite this version:

Yves Colin de Verdière, Francoise Truc. Scattering theory for graphs isomorphic to a homogeneous tree at infinity. 2012. hal-00728357v1

**HAL Id: hal-00728357**

**<https://hal.science/hal-00728357v1>**

Preprint submitted on 5 Sep 2012 (v1), last revised 17 May 2013 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Scattering theory for graphs isomorphic to a homogeneous tree at infinity <sup>\*†</sup>

Yves Colin de Verdière & Françoise Truc<sup>‡</sup>

September 5, 2012

## Abstract

We describe the spectral theory of the adjacency operator of a graph which is isomorphic to homogeneous trees at infinity. Using some combinatorics, we reduce the problem to a scattering problem for a finite rank perturbation of the adjacency operator on an homogeneous tree. We develop this scattering theory using the classical recipes for Schrödinger operators in Euclidian spaces.

## Introduction

The aim of this paper is to describe in an explicit way the spectral theory of the adjacency operator on an infinite graph  $\Gamma$  which, outside of a finite sub-graph  $\Gamma_0$ , looks like an homogeneous tree  $\mathbb{T}_q$  of degree  $q + 1$ . We mainly adapt the case of the Schrödinger operators as presented in [RS, Ike]. The proofs are often simpler here and the main results are similar. This paper can be read as an introduction to the scattering theory for differential operators on smooth manifolds. Even if we do not found our results in the literature, there is probably nothing really new for experts in the scattering theory of Schrödinger operators, except the combinatorial part in Section 4.

The main result is an explicit spectral decomposition: the Hilbert space  $l^2(\Gamma)$  splits into a sum of two invariant subspaces  $l^2(\Gamma) = \mathcal{H}_{ac} \oplus \mathcal{H}_{pp}$ . The first one is an absolutely continuous part isomorphic to that of the homogeneous tree of degree

---

**\*Keywords:** scattering on graphs, spectral measure, homogeneous tree, eigenfunction expansion.

**† Math Subject Classification (2000):** 05C12, 05C50, 05C63, 35J10, 39A12, 47B25, 81U20.

**‡**Grenoble University, Institut Fourier, Unité mixte de recherche CNRS-UJF 5582, BP 74, 38402-Saint Martin d'Hères Cedex (France); [francoise.truc@ujf-grenoble.fr](mailto:francoise.truc@ujf-grenoble.fr); <http://www-fourier.ujf-grenoble.fr/~trucfr/>

$q + 1$ , while the second one is finite dimensional and we have an upper bound on its dimension. The absolutely continuous part of the spectral decomposition is given in terms of explicit generalised eigenfunctions whose behaviour at infinity is described in terms of a scattering matrix.

We first introduce the setup, then we recall the spectral decomposition of the adjacency operator of a homogeneous tree using the Fourier-Helgason transform. In Section 3, we build the generalised eigenfunctions for a Schrödinger operator with a compactly supported potential on a homogeneous tree, we define a deformed Fourier-Helgason transform and get a spectral decomposition of this operator. As a consequence we write the point-to-point correlations of scattered waves in terms of the Green's function. Then we define the transmission coefficients, connect them to the scattering matrix, and get an explicit expression of them in terms of a Dirichlet-to-Neumann operator. In section 4, we deduce a similar spectral decomposition of any graph  $\Gamma$  asymptotic to an homogeneous tree  $\mathbb{T}_q$  by proving the following combinatorial result: any such graph  $\Gamma$  is isomorphic to a connected component of a graph  $\hat{\Gamma}$  so that the adjacency operator related to  $\hat{\Gamma}$  is a finite rank perturbation of that related to  $\mathbb{T}_q$ .

## 1 The setup: graphs asymptotic to a homogeneous tree

Let us consider a connected graph  $\Gamma = (V_\Gamma, E_\Gamma)$  with  $V_\Gamma$  the set of *vertices* and  $E_\Gamma$  the set of *edges*. We write  $x \sim y$  for  $\{x, y\} \in E_\Gamma$ .

**Definition 1.1** *Let  $q \geq 2$  be a fixed integer. We say that the infinite graph  $\Gamma$  is asymptotic to a homogeneous tree of degree  $q + 1$  if there exists a finite sub-graph  $\Gamma_0$  of  $\Gamma$  such that  $\Gamma' := \Gamma \setminus \Gamma_0$  is a disjoint union of a finite number of trees  $T_l$ ,  $l = 1, \dots, L$ , rooted at a vertex  $x_l$  linked to  $\Gamma_0$  and so that all vertices of  $T_l$  different from  $x_l$  are of degree  $q + 1$ . The trees  $T_l$ ,  $l = 1, \dots, L$ , are called the ends of  $\Gamma$ .*

*Equivalently,  $\Gamma$  is infinite, has a finite number of cycles and a maximal sub-tree of  $\Gamma$  has all vertices of degree  $q + 1$  except a finite number of them.*

**Definition 1.2** *We define the boundary  $\partial\Gamma_0$  of  $\Gamma_0$  as the set of edges of  $\Gamma$  connecting a vertex of  $\Gamma_0$  to a vertex of  $\Gamma'$ , namely one of the  $x_l$ 's. We denote by  $|x|_{\Gamma_0}$  the distance of  $x \in V_{\Gamma'}$  to  $\Gamma_0$ .*

The space of complex-valued functions on  $V_\Gamma$  is denoted

$$C(\Gamma) = \{f : V_\Gamma \longrightarrow \mathbb{C}\}$$

and  $C_0(\Gamma)$  is the subspace of functions with finite support. We define also

$$l^2(\Gamma) = \{f \in C(\Gamma); \sum_{x \in V_\Gamma} |f|^2(x) < \infty\}.$$

It is a Hilbert space when equipped with the inner product:

$$\langle f|g\rangle = \sum_{x \in V_\Gamma} \overline{f(x)} \cdot g(x) \quad .$$

Let us emphasize that we take the physicist's notation, as in [RS] for example: our inner product is conjugate-linear in the first vector and linear in the second. On  $C_0(\Gamma)$ , we define the adjacency operator  $A_\Gamma$ <sup>1</sup> by the formula:

$$(A_\Gamma f)(x) = \sum_{y \sim x} f(y) \tag{1}$$

The operator  $A_\Gamma$  is bounded on  $\ell^2(\Gamma)$  if and only if the degree of the vertices of  $\Gamma$  is bounded, which is the case here. In that case, the operator  $A_\Gamma$  is self-adjoint; otherwise, the operator  $A_\Gamma$  defined on  $C_0(\Gamma)$  could have several self-adjoint extensions.

For any  $\lambda$  outside the spectrum of  $A_\Gamma$ , we denote by  $R_\Gamma(\lambda)$  the resolvent  $(\lambda - A_\Gamma)^{-1}$  and by  $G_\Gamma(\lambda, x, y)$  the matrix of  $R_\Gamma(\lambda)$ , also called the Green's function.

## 2 The spectral decomposition of the adjacency matrix of the tree $\mathbb{T}_q$ and the Fourier-Helgason transform

### 2.1 Points at infinity

Let  $\mathbb{T}_q = (V_q, E_q)$  be the homogeneous tree of degree  $q + 1$  and let us choose an origin, also called a root,  $O$ . We denote by  $|x|$  the combinatorial distance of the vertex  $x$  to the root. The set of points at infinity denoted  $\Omega_O$  is the set of infinite simple paths starting from  $O$ . We will say that a sequence  $y_n \in V_q$  tends to  $\omega \in \Omega_O$  if, for  $n$  large enough,  $y_n$  belongs to the path  $\omega$  and is going to infinity along that path. If  $x$  is another vertex of  $V_q$ , the sets  $\Omega_O$  and  $\Omega_x$  are canonically identified by considering paths which coincide far from  $O$  and  $x$ . There is a canonical probability measure  $d\sigma_O$  on  $\Omega_O$ :  $d\sigma_O$  is the unique probability measure on  $\Omega_O$  which is invariant by the automorphisms of  $\mathbb{T}_q$  leaving  $O$  fixed. Later on we will always denote  $\Omega$  the set of points at infinity, because the root is fixed.

The *Busemann function*  $x \rightarrow b_\omega(x)$  associated to the point  $\omega \in \Omega_O$  is defined as follows: let us denote by  $x_\omega$  the last point lying on  $\omega$  in the geodesic path joining  $O$  to  $x$ , we put  $b_\omega(x) = |x_\omega| - d(x, x_\omega)$ . The level sets of  $b_\omega$  are the *horocycles* associated to  $\omega$ . The function  $b_\omega(x)$  goes to  $+\infty$  as  $x$  tends to  $\omega$ . As  $x$  tends to  $\omega' \neq \omega$ ,  $b_\omega(x)$  tends to  $-\infty$  and  $b_\omega(x) + |x|$  remains bounded.

---

<sup>1</sup>If  $B$  is a linear operator from  $C_0(\Gamma)$  into  $C(\Gamma)$ , we denote by  $[B](x, y)$ ,  $x, y \in V_\Gamma$  its matrix

## 2.2 The spectral Riemann surface

Let us define the Riemann surface  $S = \mathbb{R}/\tau\mathbb{Z} \times i\mathbb{R}$  with  $\tau = 2\pi/\log q$  and the holomorphic function  $s \rightarrow \lambda_s$  on  $S$  defined by  $\lambda_s = q^{\frac{1}{2}+is} + q^{\frac{1}{2}-is}$ . The physical sheet  $S^+ = \{s \in S \mid \Im s > 0\}$  is mapped bijectively onto  $\mathbb{C} \setminus I_q$  where, as we will see,  $I_q$  is the spectrum of  $A_0 := A_{\mathbb{T}_q}$ , namely the real segment  $[-2\sqrt{q}, +2\sqrt{q}]$ . We denote by  $S^0 = \mathbb{R}/\tau\mathbb{Z}$  the circle  $\Im s = 0$ . By the map  $s \rightarrow \lambda_s$ , the circle  $S^0$  is a double covering of  $I_q$ . If  $J$  is a subset of  $\mathbb{C}$ , we define by  $\hat{J}$  the subset of  $S$  defined by  $\hat{J} := \{s \in S \mid \lambda_s \in J\}$ .

## 2.3 Calculation of the Green's function

We denote by  $A_0$  (resp.  $G_0$ ) the adjacency operator (resp. the Green's function) on  $\mathbb{T}_q$ . We will compute explicitly  $G_0(\lambda, x, y)$ . Using the homogeneity of the tree  $\mathbb{T}_q$ , it is enough to compute  $G_0(\lambda, O, x)$  for an  $x \in V_q$ , that is the value  $f(x)$  of the  $l^2$  solution of

$$(\lambda - A_0)f = \delta_O. \quad (2)$$

The function  $f(x)$  depends only on the distance  $d(x, O) = |x|$  to the origin  $O$ , so we set  $f(x) = u_k$  if  $|x| = k$ ,  $k \in \mathbb{N}$ , and rewrite equation (2) as follows:

- i)  $\lambda u_k - q u_{k+1} - u_{k-1} = 0 \quad \text{for } k \geq 1$
- ii)  $\lambda u_0 - (q+1)u_1 = 1$
- iii)  $\sum_{n=0}^{\infty} (q+1)q^{n-1}u_n^2 < +\infty$

The last condition stands for  $f$  to be in  $l^2(\mathbb{T}_q)$ .

- If  $\lambda \notin I_q$ , the equation

$$q\alpha^2 - \lambda\alpha + 1 = 0$$

admits an unique solution  $\alpha$  such that  $|\alpha| < 1/\sqrt{q}$ . From i) and iii), we get that  $u_k = C\alpha^k$  and the constant  $C$  is determined by ii) :

$$C = C_\lambda = \frac{1}{\lambda - (q+1)\alpha}.$$

Therefore we have

$$G_0(\lambda, O, x) = \frac{2q\alpha^{|x|}}{\lambda(q-1) + (q+1)F(\lambda)}$$

where  $F(\lambda)$  denotes the determination of  $\sqrt{\lambda^2 - 4q}$  in  $\mathbb{C} \setminus I_q$  equivalent to  $\lambda$  as  $\lambda$  tends to infinity. Thus using the invariance of the Green's function by the group of automorphisms of the tree, we see that the Green's function

$G_0(\lambda, x, y)$  is a function of the distance  $d(x, y)$  and we have, for any  $x, y \in V(\mathbb{T}_q)$ ,

$$G_0(\lambda, x, y) = C_\lambda \alpha^{d(x, y)} . \quad (3)$$

The operator of matrix  $G_0(\lambda, ., .)$  is clearly bounded in  $l^2(\mathbb{T}_q)$  and  $\lambda$  is not in the spectrum of  $A_0$ .

- If  $\lambda \in I_q$ , there is no  $l^2$  solution of Equation (2) and  $\lambda$  is in the spectrum of  $A_0$ .

Using the parameter  $s \in S^+$ , we have

$$\alpha = q^{-\frac{1}{2}+is}, \quad C_{\lambda_s} := C(s) = \frac{1}{q^{\frac{1}{2}-is} - q^{-\frac{1}{2}+is}} \quad \text{and} \quad F(\lambda_s) = q^{\frac{1}{2}-is} - q^{\frac{1}{2}+is} .$$

**Theorem 2.1** *The spectrum of  $A_0$  is the interval  $I_q = [-2\sqrt{q}, +2\sqrt{q}]$ .*

*The Green's function of the tree  $\mathbb{T}_q$  is given, for  $s \in S^+$  by*

$$G_0(\lambda_s, x, y) = C(s) q^{(-\frac{1}{2}+is)d(x, y)} = \frac{q^{(-\frac{1}{2}+is)d(x, y)}}{q^{\frac{1}{2}-is} - q^{-\frac{1}{2}+is}} . \quad (4)$$

*As a function of  $s$ , the Green's function extends meromorphically to  $S$  with two poles  $-i/2$  and  $-i/2 + \tau/2$ .*

*Moreover we have, for any  $x \in V_q$  and any  $y$  belonging to the path  $\omega$ ,*

$$G_0(\lambda_s, x, y) = G_{rad}(\lambda_s, y) q^{(\frac{1}{2}-is)b_\omega(x)} \quad (5)$$

*with*

$$G_{rad}(\lambda_s, y) = C(s) q^{(-\frac{1}{2}+is)|y|} \quad (6)$$

*Proof.*–

The last result comes from the definition  $b_\omega(x) = |x_\omega| - d(x, x_\omega)$ .

□

## 2.4 The density of states

Let us recall how to introduce a notion of spectral measure (also called density of states) on the graph  $\Gamma$ . To a given continuous function  $\phi : \mathbb{R} \rightarrow \mathbb{R}$ , we associate by the functional calculus an operator  $\phi(A_\Gamma)$  on  $l^2(\Gamma)$ , which has a matrix  $[\phi(A_\Gamma)](x, x')$ . We consider then, for any  $x \in V_\Gamma$ , the linear form on  $C(\mathbb{R}, \mathbb{R})$

$$L_x(\phi) = [\phi(A_\Gamma)](x, x) .$$

$L_x$  is positive and verifies  $L_x(1) = 1$ , so we have  $L_x(\phi) = \int_{\mathbb{R}} \phi d e_x$  where  $d e_x$  is a probability measure on  $\mathbb{R}$ , supported by the spectrum of  $A_\Gamma$  which is called the spectral measure of  $\Gamma$  at the vertex  $x$ .

The density of states of  $\mathbb{T}_q$  is given by the

**Theorem 2.2** (See for example [CdV2]) *The spectral measure  $de_x$  of  $\mathbb{T}_q$  is independent of the vertex  $x$  and is given by*

$$de_x(\lambda) := de(\lambda) = \frac{(q+1)\sqrt{4q-\lambda^2}}{2\pi((q+1)^2-\lambda^2)}d\lambda \quad (7)$$

*Proof.*–

For the sake of clarity, we recall the main ingredients:

1) an explicit computation of the diagonal entries of the Green's function

$$G_0(\lambda, x, x) = \frac{2q}{\lambda(q-1) - (q+1)F(\lambda)}$$

where  $F(\lambda)$  denotes as previously the determination of  $\sqrt{\lambda^2 - 4q}$  in  $\mathbb{C}/I_q$  (with  $I_q = [-2\sqrt{q}, 2\sqrt{q}]$ ) equivalent to  $\lambda$  for great values of  $\lambda$ .

2) The expression of the spectral measure via Stone formula

$$de(\lambda) = \frac{-1}{2i\pi} (G(\lambda + i0, x, x) - G(\lambda - i0, x, x)) dt . \quad (8)$$

□

## 2.5 The Fourier-Helgason transform

Let us recall the definition of the Fourier-Helgason transform on the tree  $\mathbb{T}_q$  with the root  $O$ .

**Definition 2.1** *For any  $f \in C_0(\mathbb{T}_q)$ , the Fourier-Helgason transform  $\mathcal{FH}(f)$  is the function defined by*

$$\mathcal{FH}(f)(\omega, s) := \hat{f}(\omega, s) = \sum_{x \in V_q} f(x) q^{(1/2+is)b_\omega(x)} . \quad (9)$$

for any  $\omega \in \Omega_O$  and any  $s \in S$ .

Let us set

$$e_0(x, \omega, s) := q^{(1/2-is)b_\omega(x)} ,$$

and denote by  $e_0(\omega, s)$  the function  $x \rightarrow e_0(x; \omega, s)$ . If we restrict ourselves to  $s \in S^0$ , the previous definition writes

$$\hat{f}(\omega, s) = \langle e_0(\omega, s) | f \rangle = \sum_{x \in V_\Gamma} f(x) \overline{e_0(x, \omega, s)} . \quad (10)$$

We have

$$\forall s \in S_0, \quad A_0 e_0(\omega, s) = \lambda_s e_0(\omega, s) ,$$

and the completeness of the set  $\{e_0(\omega, s), s \in S^0, \omega \in \Omega\}$  is expressed by the following inversion formula (see [CMS]):

**Theorem 2.3** For any  $f \in C_0(\mathbb{T}_q)$ , the following inverse transform holds

$$f(x) = \int_{S^0} \int_{\Omega} e_0(x, \omega, s) \hat{f}(\omega, s) d\sigma_O(\omega) d\mu(s) \quad (11)$$

where

$$d\mu(s) = \frac{(q+1) \log q}{\pi} \frac{\sin^2(s \log q)}{q + q^{-1} - 2 \cos(2s \log q)} |ds| . \quad (12)$$

Moreover the Fourier-Helgason transform extends to a unitary map from  $l^2(\mathbb{T}_q)$  into  $L^2(\Omega \times S^0, d\sigma_O \otimes d\mu)$ .

The Fourier-Helgason transform is not surjective: its range is the subspace  $L^2_{\text{even}}(\Omega \times S^0, d\sigma_O \otimes d\mu)$  of the functions  $F$  of  $L^2(\Omega \times S^0, d\sigma_O \otimes d\mu)$  which satisfy the symmetry condition (see, for example, [CS] or [FN])

$$\int_{\Omega} e_0(x, \omega, s) F(\omega, s) d\sigma_O(\omega) = \int_{\Omega} e_0(x, \omega, -s) F(\omega, -s) d\sigma_O(\omega) .$$

The Fourier-Helgason transform provides a spectral resolution of  $A_0$ : if  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  is continuous,

$$\phi(A_0) = (\mathcal{FH})^{-1} \phi(\lambda_s) \mathcal{FH} ,$$

where  $\phi(\lambda_s)$  denotes the operator of multiplication by that function on  $L^2_{\text{even}}(\Omega \times S^0, d\sigma_0 \otimes d\mu)$ .

**Corollary 2.1** From the inverse Fourier-Helgason transform formula (11) we find back the expression of the spectral measure of  $\mathbb{T}_q$  (see Theorem 2.2).

*Proof.* –

By homogeneity of the tree  $\mathbb{T}_q$ , for any  $\phi \in C_0(\mathbb{T}_q)$ ,  $[\phi(A_0)](x, x)$  is independent of  $x$ . Using (11), we get

$$[\phi(A_0)](O, O) = \int_{\Omega} \int_{S^0} \Phi(\lambda_s) \overline{e_0(O, \omega, s)} e_0(O, \omega, s) d\sigma_O(\omega) d\mu(s) = \int_{S^0} \Phi(\lambda_s) d\mu(s).$$

Let us perform the change of variables

$$s = f_q(\lambda) := \frac{1}{\log q} \arccos \frac{\lambda}{2\sqrt{q}} .$$

Using (12) and the fact that, by the map  $s \rightarrow \lambda_s$ , the circle  $S^0$  is a double covering of  $I_q$ , we write

$$\begin{aligned} [\phi(A_0)](x, x) &= 2 \frac{(q+1) \log q}{\pi} \int_{I_q} \Phi(\lambda) \frac{1 - \lambda^2/4q}{q + q^{-1} + 2 - \lambda^2/q} f'_q(\lambda) d\lambda \\ &= 2 \frac{(q+1)}{4\pi} \int_{I_q} \frac{\sqrt{4q - \lambda^2}}{(q+1)^2 - \lambda^2} \Phi(\lambda) d\lambda , \end{aligned}$$

which actually implies formula (7).

□

### 3 A scattering problem for a Schrödinger operator with a compactly supported non local potential

We are concerned here with the scattering on  $\mathbb{T}_q$  between the adjacency operator  $A_0$  and the Schrödinger operator  $A = A_0 + W$ , where  $W$  is a compactly supported non local potential. More precisely the Hermitian matrix (also denoted  $W$ ) associated to this potential is supported by  $K \times K$  where  $K$  is a finite part of  $V_q$ . We assume in what follows that  $K$  is chosen minimal, so that:

$$K = \{x \in V_q \mid \exists y \in V_q \text{ with } V_{x,y} \neq 0\} .$$

Let us first describe the spectral theory of  $A$ : it follows from [RS], Sec. XI 3, and from the fact that  $A$  is a finite rank perturbation of  $A_0$  (see also Section 3.3) that the Hilbert space  $l^2(\mathbb{T}_q)$  admits an orthogonal decomposition into two subspaces invariant by  $A$ :  $l^2(\mathbb{T}_q) = \mathcal{H}_{ac} \oplus \mathcal{H}_{pp}$  where

- $\mathcal{H}_{ac}$  is the isometric image of  $l^2(\mathbb{T}_q)$  by the *wave operator*

$$\Omega^+ = s - \lim_{t \rightarrow -\infty} e^{itA} e^{-itA_0} .$$

We have  $A|_{\mathcal{H}_{ac}} = \Omega^+ A_0 (\Omega^+)^*$ , so that the corresponding part of the spectral decomposition is isomorphic to that of  $A_0$  which is an absolutely continuous spectrum on the interval  $I_q$ .

- The space  $\mathcal{H}_{pp}$  is finite dimensional, admits an orthonormal basis of  $l^2$  eigenfunctions associated to a finite set of eigenvalues, some of them can be embedded in the continuous spectrum  $I_q$ .

We will denote by  $P_{ac}$  and  $P_{pp}$  the orthogonal projections on both subspaces.

In order to make the spectral decomposition more explicit, we will introduce suitable generalised eigenfunctions of  $A$ . These generalised eigenfunctions are particular solutions of

$$(\lambda_s - A)e(., \omega, s) = 0 , \tag{13}$$

meaning not  $l^2$  solutions, but only point-wise solutions. For the adjacency operator  $A_0$ , these generalised eigenfunctions are called the “the plane waves” and given by the  $e_0(\omega, s)$ ’s with  $s \in S^0$  and  $\omega \in \Omega_O$ . They give the Fourier-Helgason transform which is the spectral decomposition of  $A_0$ .

We are going to prove a similar eigenfunction expansion theorem for  $A$ , using generalised eigenfunctions of  $A$ . We will mainly adapt the presentation of [RS], Sec. XI.6, for Schrödinger operators in  $\mathbb{R}^3$  (see also [Ike]). Our first goal is to build the generalised eigenfunctions  $x \rightarrow e(x, \omega, s)$  also denoted  $e(\omega, s)$ . We will derive and solve the so-called *Lippmann-Schwinger equation*. This is an integral equation that  $e(\omega, s)$  will satisfy.

### 3.1 Formal derivation of the Lippmann-Schwinger equation

Let us proceed first in a *formal way* by transferring the functions  $e_0(\omega, s)$  by the wave operator: if  $e(\omega, s)$  is the image of  $e_0(\omega, s)$  by the wave operator  $\Omega^+$  in some sense (they are not in  $l^2$ !), then we should have  $e_0(\omega, s) = \lim_{t \rightarrow -\infty} e^{itA_0} e^{-itA} e(\omega, s)$

$$= \lim_{t \rightarrow -\infty} [e(\omega, s) - i \int_0^t e^{iuA_0} W e^{-iuA} e(\omega, s) du]$$

$$= e(\omega, s) - i \lim_{\varepsilon \rightarrow 0} \int_0^{-\infty} e^{iuA_0} W e^{-iu\lambda_s} e^{\varepsilon u} e(\omega, s) du$$

$$= e(\omega, s) + \lim_{\varepsilon \rightarrow 0} [(A_0 - (\lambda_s + i\varepsilon))^{-1} W e](\omega, s) .$$

So  $e(\omega, s)$  should obey the following "Lippmann-Schwinger-type" equation

$$e(\omega, s) = e_0(\omega, s) + G_0(\lambda_s) W e(\omega, s) . \quad (14)$$

### 3.2 Existence and unicity of the solution for the modified "Lippmann-Schwinger-type" equation

Let  $\chi \in C_0(\mathbb{T}_q)$  be a compactly supported real-valued function so that  $W\chi = \chi W = W$ . For example  $\chi$  can be the characteristic function of  $K$ . We first introduce a modified "Lippmann-Schwinger-type" equation. If  $e(\omega, s)$  obeys (14) and  $a(\omega, s) = \chi e(\omega, s)$ , then  $a$  obeys

$$a(\omega, s) = \chi e_0(\omega, s) + \chi G_0(\lambda_s) W a(\omega, s) . \quad (15)$$

Let  $K_s$  be the finite rank operator on  $l^2(\mathbb{T}_q)$  defined by  $K_s = \chi G_0(\lambda_s) W$ . The map  $s \rightarrow K_s$  extends holomorphically to  $\Im s > -\frac{1}{2}$ . Equation (15) takes the form

$$a(., \omega, s) = \eta(., \omega, s) + K_s a(., \omega, s), \quad (16)$$

where  $\eta(., \omega, s) \in C_0(\mathbb{T}_q)$ . By the analytic Fredholm theorem ([RS], p 101), there exists a finite subset  $\hat{\mathcal{E}}$  of  $S^0$ , defined by  $\hat{\mathcal{E}} =: \{s \in S^0; \ker(\text{Id} - K_s) \neq 0\}$ , so that equation (15) has a unique solution  $a(\omega, s) \in C_0(\mathbb{T}_q)$  whenever  $s \notin \hat{\mathcal{E}}$ . Since  $K_s = K_{-s}$ , the subset  $\hat{\mathcal{E}}$  is invariant by  $s \rightarrow -s$  and consequently is the inverse image by  $s \rightarrow \lambda_s$  of a subset of  $I_q$  which we denote by  $\mathcal{E}$ .

For  $s \notin \hat{\mathcal{E}}$ , the function  $e(\omega, s) = e_0(\omega, s) + G_0(\lambda_s) W a(\omega, s)$  is the unique solution of the Lippmann-Schwinger equation (14).

### 3.3 The set $\mathcal{E}$ and the pure point spectrum

**Proposition 3.1** *The set  $\mathcal{E}$  as defined in Section 3.2 is independent of the choice of  $\chi$  with  $W\chi = \chi W = W$ .*

The “minimal”  $\chi$  is  $\chi_W = \mathbf{1}_K$ . If  $a$  is a non trivial solution of  $a - \chi_W G_0(\lambda_s) W a = 0$ , and  $\chi \chi_W = \chi_W$ ,  $a$  is also solution of  $a - \chi G_0(\lambda_s) W a = 0$ .

Conversely, if  $a - \chi G_0(\lambda_s) W a = 0$ , we have  $\chi_W a - \chi_W G_0(\lambda_s) W a = 0$ . We have to prove that  $\chi_W a \neq 0$ . If  $\chi_W a = 0$ , we would have  $W a = 0$  and  $a = 0$ .

**Proposition 3.2** *If  $(A - \lambda)f = 0$  with  $\lambda \in I_q$  and  $f \in l^2(\mathbb{T}_q)$ , then  $\text{Supp}(f) \subset \hat{K}$  where  $\hat{K}$  is the smallest subset of  $V_q$  so that  $\text{Supp}(W) \subset \hat{K} \times \hat{K}$  and all connected components of  $\mathbb{T}_q \setminus \hat{K}$  are infinite.*

*Proof.* –

We will proceed by contradiction. Let  $x \in V_q \setminus \hat{K}$  be so that  $f(x) \neq 0$ . Let us define an infinite sub-tree  $\mathbb{T}_x$  of  $\mathbb{T}_q$  as follows: let  $y_\alpha, \alpha = 1, \dots, a$  be the vertices of  $\mathbb{T}_q$  which satisfy  $y_\alpha \sim x$  and  $y_\alpha$  is closer to  $K$  than  $x$ . Then  $\mathbb{T}_x$  is the connected component of  $x$  in the graph obtained from  $\mathbb{T}_q$  by removing the edges  $\{x, y_\alpha\}$  for  $\alpha = 1, \dots, a$ . Let us consider the “averaged” function

$$n \in \mathbb{N} \rightarrow \bar{f}_x(n) := \frac{1}{q^n} \sum_{z \in \mathbb{T}_x, d(x,z)=n} f(z) .$$

Then  $\bar{f}_x$  satisfies the ordinary difference equation  $\lambda g(n) - qg(n+1) - g(n-1) = 0$ .

We thus get a contradiction, since this equation has no non-zero  $l^2$  solution when  $\lambda$  is in  $I_q$ .

□

**Corollary 3.1**  $\#\{\sigma_{\text{pp}}(A) \cap I_q\} \leq \#\hat{K}$ .

This holds because any eigenfunction associated to an eigenvalue in  $\{\sigma_{\text{pp}}(A) \cap I_q\}$  is supported in  $\hat{K}$  and the dimension of the vector space of functions supported in  $\hat{K}$  is  $\#\hat{K}$ .

**Theorem 3.1** *If  $s \in S^0$ ,  $(A - \lambda_s)f = 0$  and  $f \in l^2(\mathbb{T}_q) \setminus 0$ , then  $s \in \hat{\mathcal{E}}$ .*

*Conversely, if  $s \in \hat{\mathcal{E}} \subset S^0$ , there exists  $f \neq 0$  so that  $(A - \lambda_s)f = 0$  and  $f(x) = O(q^{-|x|/2})$ .*

*Proof.* –

Due to Proposition 3.2, the support of such an  $f$  is included in  $\hat{K}$  and  $(\lambda_s - A_0)f = Wf$ . We apply  $G_0(\lambda_s)$  to both sides of the equation, because every sum is finite. We have  $G_0(\lambda_s)(\lambda_s - A_0)f = f$ : this is true for  $\lambda_s \notin I_q$  because  $I_q$  is the spectrum of  $A_0$  and hence by continuity  $(G_0(\lambda_s)$  extends holomorphically near  $S^0$ ) for every  $\lambda_s$  since

$f$  is compactly supported. Hence  $f - G_0(\lambda_s)Wf = 0$ . If we choose for  $\chi$  the characteristic function of  $\hat{K}$ , we get  $f - \chi G_0(\lambda_s)Wf = 0$ . We have a non trivial solution of  $a - K_s a = 0$ , namely  $a = f$ .

Conversely, let us start from  $a$ , a non trivial solution of  $a - K_s a = 0$  and define  $f = G_0(\lambda_s)W a$ . Then  $(\lambda_s - A)f = 0$  by the same reasoning as before. Moreover  $f$  is a finite linear combination of the functions  $G_0(\lambda_s, \cdot, y)$ ,  $y \in \text{Supp}(W)$  and we use Equation (6) to get the bound in  $x$ .

□

**Theorem 3.2** *The pure point spectrum  $\sigma_{\text{pp}}(A)$  of  $A$  splits into 3 parts*

$$\sigma_{\text{pp}}(A) = \sigma_{\text{pp}}^-(A) \cup \sigma_{\text{pp}}^+(A) \cup \sigma_{\text{pp}}^0(A)$$

where  $\sigma_{\text{pp}}^-(A) = \sigma_{\text{pp}}(A) \cap ]-\infty, -2\sqrt{q}[$ ,  $\sigma_{\text{pp}}^+(A) = \sigma_{\text{pp}}(A) \cap ]2\sqrt{q}, +\infty[$ , and  $\sigma_{\text{pp}}^0(A) = \sigma_{\text{pp}}(A) \cap I_q$ . We have  $\#\sigma_{\text{pp}}^\pm(A) \leq \#\text{Supp}(W)$  and  $\#\sigma_{\text{pp}}^0(A) \leq \#\hat{K}$ .

The first estimate comes from the mini-max principle and the fact that  $W$  is a rank  $N$  perturbation of  $A_0$  with  $N = \#\text{Supp}(W)$ . The second one is already proved.

The reader could ask if there can really be some compactly supported eigenfunctions. They can exist as shown by the following 2 examples.

**Example 3.1**  $\Gamma$  is a tree with root 0 and  $W_{x,0} = W_{0,x} = -1$  for any  $x \sim 0$ . All other entries of  $W$  vanish. Then if  $H = A_\Gamma + W$ ,  $f = \delta(0)$ , we have  $Hf = 0$ .

**Example 3.2** The graph  $\Gamma$  is the union of a cycle with 4 vertices  $\{1, 2, 3, 4\}$  and a tree whose root is attached to 2 neighbouring vertices of the cycle. If  $f(p) = (-1)^p$  on the cycle and 0 on all other vertices,  $A_\Gamma f = 0$ .

However the proof of the following result is left to the reader:

**Proposition 3.3** *If  $\Gamma$  is an infinite tree, then  $A_\Gamma$  has no compactly supported eigenfunction.*

### 3.4 The deformed Fourier-Helgason transform

**Definition 3.1** *We define the deformed Fourier-Helgason transform  $\mathcal{FH}_{\text{sc}}$  of  $f \in C_0(\mathbb{T}_q)$  as the function  $\hat{f}_{\text{sc}}$  on  $\Omega \times (S^0 \setminus \hat{\mathcal{E}})$  defined by*

$$\hat{f}_{\text{sc}}(\omega, s) = \langle e(\omega, s) | f \rangle = \sum_{x \in V_\Gamma} f(x) \overline{e(x, \omega, s)}. \quad (17)$$

We want to prove the following

**Theorem 3.3** *For any  $f \in C_0(\mathbb{T}_q)$  and any closed interval  $J \subset I_q \setminus \mathcal{E}$ , if we denote by  $\hat{J}$  the inverse image of  $J$  by  $s \rightarrow \lambda_s$ , the following inverse transform holds*

$$P_J f(x) = \int_{\hat{J}} \int_{\Omega} e(x, \omega, s) \hat{f}_{\text{sc}}(\omega, s) d\sigma_O(\omega) d\mu(s) . \quad (18)$$

Moreover,  $f \rightarrow \hat{f}_{\text{sc}}$  extends to an isometry from  $\mathcal{H}_{\text{ac}}$  onto  $L^2_{\text{even}}(\Omega \times S^0, d\sigma_O \otimes d\mu)$ .

### 3.4.1 The relation of the deformed Fourier-Helgason transform with the resolvent

Denoting, with a slight abuse of notation, for  $s \in S^+$ , by  $G(s)$  the operator  $(\lambda_s - A)^{-1}$  and similarly by  $G_0(s)$  the operator  $(\lambda_s - A_0)^{-1}$ , we have the resolvent equation

$$G(s) = G_0(s) + G_0(s)WG(s) \quad (19)$$

For  $\sigma \in S^0$  and  $s$  in  $S^+$ , we set

$$h(s; \omega, \sigma) = (\lambda_s - \lambda_\sigma)G(s)e_0(\omega, \sigma) ,$$

where the right-handside is a convergent series which identifies to  $(\lambda_s - \lambda_\sigma)$ -times the inverse Fourier-Helgason transform of  $y \rightarrow G(s; x, y)$ .

Then equation (19) for  $G$  gives an integral equation for  $h$

$$h(s; \omega, \sigma) = e_0(\omega, \sigma) + G_0(s)Wh(s; \omega, \sigma)$$

and, if  $p(s; \omega, \sigma) = \chi h(s; \omega, \sigma)$ ,

$$p(s; \omega, \sigma) = \chi e_0(\omega, \sigma) + \chi G_0(s)Wp(s; \omega, \sigma) . \quad (20)$$

The key fact is the relation between (20) and the modified "Lippmann-Schwinger-type" equation (15). If  $s \in S^+$  is fixed and  $\sigma = s$ , then the equation for  $p(s; \omega, s)$  is identical to equation (15) for  $a(\omega, s)$ .

This can be used to prove

**Lemma 3.1** *Let  $f \in C_0(\mathbb{T}_q)$  be fixed and  $s \in S^+$ . Then the function*

$$\Phi(s; \omega, \sigma) = \sum_{x \in V_q} \overline{h(x; s; \omega, \sigma)} f(x)$$

*has a holomorphic extension in  $\sigma$  to  $S^+$  and*

$$\Phi(s; \omega, s) = \sum_{x \in V_q} \overline{e(x, \omega, s)} f(x) = \hat{f}_{\text{sc}}(\omega, s) .$$

We thus have related  $\hat{f}_{\text{sc}}$  to the resolvent.

### 3.4.2 End of the proof of Theorem 3.3

Let  $\lambda_s = \Lambda + i\varepsilon$  with  $\Lambda \in I_q \setminus \mathcal{E}$  and  $\varepsilon > 0$ , and  $s \in S^+$  (this implies  $0 < \Re s < \tau/2$ ). Up to a factor of  $(\Lambda + i\varepsilon - \lambda_\sigma)$ ,  $(\omega, \sigma) \rightarrow h(x; s; \omega, \sigma)$  is the inverse Fourier-Helgason transform of  $y \rightarrow G(\lambda_s, x, y)$ ; so the Plancherel theorem implies (after multiplying by  $\overline{f(x)}f(y)$ ) that

$$\begin{aligned} & (\lambda_s - \bar{\lambda}_s) \sum_{z \in V_q} \overline{G(\bar{\lambda}_s, x, z)} G(\bar{\lambda}_s, z, y) \overline{f(x)} f(y) = \dots \\ & \dots 2i\varepsilon \int_{S^0} \int_{\Omega} \frac{h(x; s; \omega, \sigma) \overline{h(y; s; \omega, \sigma)} \overline{f(x)} f(y)}{|\lambda_\sigma - \Lambda|^2 + \varepsilon^2} d\sigma_O(\omega) d\mu(\sigma) \end{aligned}$$

If we sum over all  $x$ 's and  $y$ 's, we obtain for the left-hand side

$$(\lambda_s - \bar{\lambda}_s) \langle G(\bar{\lambda}_s) f, G(\bar{\lambda}_s) f \rangle = (\lambda_s - \bar{\lambda}_s) \langle f | G(\lambda_s) G(\bar{\lambda}_s) f \rangle = \langle f | [G(\lambda_s) - G(\bar{\lambda}_s)] f \rangle$$

whereas the right-hand side becomes

$$\int_{S^0} \int_{\Omega} \frac{2i\varepsilon}{|\lambda_\sigma - \Lambda|^2 + \varepsilon^2} |\Phi(s; \omega, \sigma)|^2 d\sigma_O(\omega) d\mu(\sigma) .$$

We thus conclude that, for any closed sub-interval  $J$  of  $I_q$  disjoint from  $\mathcal{E}$ ,

$$\frac{1}{2\pi i} \int_J \langle f | [G(\Lambda + i\varepsilon) - G(\Lambda - i\varepsilon)] f \rangle d\Lambda = \frac{1}{\pi} \int_J d\Lambda \int_{S^0} \int_{\Omega} \frac{\varepsilon}{|\lambda_\sigma - \Lambda|^2 + \varepsilon^2} |\Phi(s, \omega, \sigma)|^2 d\sigma_O(\omega) d\mu(\sigma)$$

As  $\varepsilon \rightarrow 0$ , Stone's formula implies that the left-hand side approaches  $\|P_J f\|^2$ . Moreover the measures

$$dl_\varepsilon = \frac{\varepsilon d\Lambda}{\pi[|\lambda_\sigma - \Lambda|^2 + \varepsilon^2]}$$

converge weakly to  $\delta(\Lambda - \lambda_\sigma)$  as  $\varepsilon \rightarrow 0^+$ . So one has to put  $\Lambda = \lambda_\sigma$ , which implies  $\sigma = \pm s$ .

Thus one gets that the right-hand side tends to  $\int_J \int_{\Omega} |\hat{f}_{sc}(\omega, \sigma)|^2 d\sigma_O(\omega) d\mu(\sigma)$ , where  $\hat{J}$  is the inverse image of  $J$  by  $s \rightarrow \lambda_s$ .

## 3.5 Correlation of scattered plane waves

In the paper [CdV3], the first author computed the point-to-point correlations of the plane waves for a scattering problem in  $\mathbb{R}^d$  in terms of the Green's function: for a fixed spectral parameter, plane waves are viewed as random waves parametrised by the direction of their incoming part. The motivation comes from passive imaging in seismology, a method developped by Michel Campillo's seismology group in Grenoble, as described for example in the papers [CdV4, CdV5]. Following a similar method, we will compute the correlation of plane waves for our graphs viewed as random waves parametrised by points at infinity.

From (11), we get, for any  $\phi \in C_0(\mathbb{T}_q)$ , the following formula for the kernel of  $\phi(A)$ :

$$[\phi(A)](x, y) = \int_{\Omega} \int_{S^0} \Phi(\lambda_s) \overline{e(x, \omega, s)} e(y, \omega, s) d\sigma_O(\omega) d\mu(s)$$

Taking  $\phi = 1_I$ , the characteristic function of some interval  $I = [a, \lambda] \subset I_q$ , we get:

$$[\Pi_I](x, y) = 2 \int_{\Omega} \int_{f_q(\lambda)}^{f_q(a)} \overline{e(x, \omega, s)} e(y, \omega, s) d\sigma_O(\omega) d\mu(s)$$

where  $f_q(t) = \frac{1}{\log q} \text{Arccos} \frac{t}{2\sqrt{q}}$  (see section 2.5). In particular  $f_q(\lambda_s) = s$ .

If we consider the plane wave  $e(x, \omega, s(\lambda))$  as a random wave, we can define the point-to-point correlation  $C_{\lambda}^{sc}(x, y)$  of such a random wave in the usual way:

**Definition 3.2** *The point-to-point correlation  $C_{\lambda}^{sc}(x, y)$  of the random wave  $e(x, \omega, s(\lambda))$  is given by*

$$C_{\lambda}^{sc}(x, y) = \int_{\Omega} \overline{e(x, \omega, s(\lambda))} e(y, \omega, s(\lambda)) d\sigma(\omega) .$$

Denoting as usual by  $G$  the resolvent kernel of  $A$ :  $[(\lambda - A)^{-1}](x, y) := G(\lambda, x, y)$  for  $\text{Im} \lambda > 0$  we prove the

**Theorem 3.4** *For any  $\lambda \in I_q$  and any vertices  $x, y$  the point-to-point correlation can be expressed in terms of the Green's function as*

$$C_{\lambda}^{sc}(x, y) = -\frac{2(q^2 + 2q + 1 - \lambda^2)}{(q + 1)\sqrt{4q - \lambda^2}} \Im G(\lambda + i0, x, y) .$$

*Proof.* –

Taking the derivative with respect to  $\lambda$  in equation (3.5) yields:

$$\begin{aligned} \frac{d}{d\lambda} [\Pi_I](x, y) &= \dots \\ &= -2f'_q(\lambda) \frac{(q + 1) \log q}{\pi} \frac{\sin^2(s(\lambda) \log q)}{q + q^{-1} - 2 \cos(2s(\lambda) \log q)} \int_{\Omega} \overline{e(x, \omega, s(\lambda))} e(y, \omega, s(\lambda)) d\sigma_O(\omega) \\ &= \frac{q + 1}{2\pi} \frac{\sqrt{4q - \lambda^2}}{(q^2 + 2q + 1 - \lambda^2)} \int_{\Omega} \overline{e(x, \omega, s(\lambda))} e(y, \omega, s(\lambda)) d\sigma_O(\omega). \end{aligned}$$

(We have set  $f_q(\lambda) = s(\lambda)$  for simplicity.)

Thus we have

$$\frac{d}{d\lambda} [\Pi_I](x, y) = \frac{q + 1}{2\pi} \frac{\sqrt{4q - \lambda^2}}{(q^2 + 2q + 1 - \lambda^2)} C_{\lambda}^{sc}(x, y) .$$

Now we use the resolvent kernel of  $A : [\lambda - A]^{-1}(x, y) := G(\lambda, x, y)$  for  $\text{Im}\lambda > 0$  and Stone formula (8) to write

$$[\Pi_I](x, y) = -\frac{1}{\pi} \int_a^\lambda \Im G(t + i0, x, y) dt ,$$

and get the result. □

## 3.6 The T- matrix

### 3.6.1 Another feature of the deformed Fourier-Helgason transform

We will need in this section the following result

**Theorem 3.5** *For any  $f \in C_0(\mathbb{T}_q)$*

$$\mathcal{FH}_{\text{sc}}(\Omega^+ f)(\omega, s) = \hat{f}(\omega, s) \quad (21)$$

*Proof.*–

Suppose that we can prove

$$\mathcal{FH}((\Omega^+)^* f) = \mathcal{FH}_{\text{sc}}(f) ,$$

then (21) follows from

$$\mathcal{FH}_{\text{sc}}(\Omega^+ f) = \mathcal{FH}((\Omega^+)^* \Omega^+ f) = \hat{f} .$$

So, by Plancherel formula it is enough to prove that

$$\langle f | \Omega^+ g \rangle = \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}} f(\omega, s)} \hat{g}(\omega, s) d\sigma_O(\omega) d\mu(s) . \quad (22)$$

We have

$$\langle f | \Omega^+ g \rangle - \langle f | g \rangle = i \lim_{\varepsilon \rightarrow 0} \int_{-\infty}^0 e^{\varepsilon t} \langle f | e^{itA} W e^{-itA_0} g \rangle dt$$

But

$$\langle f | e^{itA} h \rangle = \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}} f(\omega, s)} e^{i\lambda_s t} \mathcal{FH}_{\text{sc}} h(\omega, s) d\sigma_O(\omega) d\mu(s)$$

if either  $f$  or  $h$  is in  $\mathcal{H}_{ac}$ .

As a result

$$\langle f | e^{itA} W e^{-itA_0} g \rangle = \sum_{x \in V_q} \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}} f(\omega, s)} e^{i\lambda_s t} (W e^{-itA_0} g)(x) \overline{e(x, \omega, s)} d\sigma_O(\omega) d\mu(s) .$$

Thus

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0} \int_{-\infty}^0 e^{\varepsilon t} \langle f | e^{itA} W e^{-itA_0} g \rangle dt \\
&= \lim_{\varepsilon \rightarrow 0} \sum_{x \in V_q} \int_{-\infty}^0 \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}} f(\omega, s)} W(x) e^{-it(A_0 - \lambda_s + i\varepsilon)} g(x) \overline{e(x, \omega, s)} d\sigma_O(\omega) d\mu(s) \\
&= -i \lim_{\varepsilon \rightarrow 0} \sum_{x \in V_q} \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}} f(\omega, s)} W(x) [(A_0 - \lambda_s + i\varepsilon)^{-1} g](x) \overline{e(x, \omega, s)} d\sigma_O(\omega) d\mu(s) \\
&= i \lim_{\varepsilon \rightarrow 0} \sum_{x, y \in V_q} \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}} f(\omega, s)} W(x) \overline{G_0(\lambda_s + i\varepsilon)}(x, y) g(y) \overline{e(x, \omega, s)} d\sigma_O(\omega) d\mu(s) \\
&= i \sum_{y \in V_q} \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}} f(\omega, s)} \left[ \sum_{x \in V_q} \overline{G_0(\lambda_s)}(x, y) W(x) \overline{e(x, \omega, s)} \right] g(y) d\sigma_O(\omega) d\mu(s) \\
&= i \sum_{y \in V_q} \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}} f(\omega, s)} [\overline{e(y, \omega, s)} - \overline{e_0(y, \omega, s)}] g(y) d\sigma_O(\omega) d\mu(s) \\
&= i \langle f | g \rangle - i \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}} f(\omega, s)} \hat{g}(\omega, s) d\sigma_O(\omega) d\mu(s) .
\end{aligned}$$

In the second line above we used the Lippmann-Schwinger equation and at the last step we used the isometric property of the deformed Fourier Helgason transform (Theorem 3.3).

□

The Lippmann-Schwinger eigenfunctions  $e(x, \omega, s)$  are especially useful to describe the so-called  $S$ -matrix ( $S = (\Omega^-)^* \Omega^+$ ). First we introduce the following object:

**Definition 3.3** Let  $(\omega, s)$  and  $(\omega', s')$  be in  $\Omega \times (S^0 \setminus \hat{\mathcal{E}})$ . Define

$$T(\omega, s; \omega', s') = \langle W e_0(\omega, s) | e(\omega', s') \rangle = \sum_{(x, y) \in V_q \times V_q} e(x, \omega', s') \overline{W(x, y) e_0(y, \omega, s)} .$$

$T(., .)$  is called the  $T$ -matrix.

**Theorem 3.6** For any  $f$  and  $g \in C_0(\mathbb{T}_q)$

$$(f | (S - I)g) = -2\pi i \int_{S^0 \times S^0} \int_{\Omega \times \Omega} T(\omega, s; \omega', s') \overline{\hat{f}(\omega, s)} \delta(\lambda_s - \lambda_{s'}) \hat{g}(\omega', s') d\sigma_O(\omega) d\mu(s) d\sigma_O(\omega') d\mu(s')$$

(See Appendix A for the precise definition of the measure  $\delta(\lambda_s - \lambda_{s'}) d\mu(s) d\mu(s')$ ).

This can be written symbolically by

$$S(\omega, s; \omega', s') = \delta(s - s') - 2\pi i T(\omega, s; \omega', s') \delta(\lambda_s - \lambda_{s'}) . \quad (23)$$

*Proof.*–

From the definition of  $S$  we get that

$$\begin{aligned}\langle f|(S-I)g\rangle &= \langle (\Omega^- - \Omega^+)f|\Omega^+g\rangle = \lim_{T \rightarrow \infty} \int_T^{-T} \langle e^{itA}(iW)e^{-itA_0}f|\Omega^+g\rangle dt \\ &= (-i)\lim_{\varepsilon \rightarrow 0} \int_{-\infty}^{+\infty} e^{-\varepsilon|t|} \langle e^{itA}We^{-itA_0}f|\Omega^+g\rangle dt \\ &= (-i)\lim_{\varepsilon \rightarrow 0} \int_{-\infty}^{+\infty} e^{-\varepsilon|t|} \mathcal{L}(t) dt ,\end{aligned}$$

with

$$\mathcal{L}(t) = \int_{S^0} \int_{\Omega} \overline{\mathcal{FH}_{\text{sc}}(e^{itA}We^{-itA_0}f)}(\omega', s') \mathcal{FH}_{\text{sc}}(\Omega^+g)(\omega', s') d\sigma_O(\omega') d\mu(s') \quad (24)$$

In the last step we used the isometric property of  $\mathcal{FH}_{\text{sc}}$  and the fact that  $\Omega^+g \in \mathcal{H}_{ac}$ . Moreover we have (Theorem 3.5)

$$\mathcal{FH}_{\text{sc}}(\Omega^+g)(\omega', s') = \hat{g}(\omega', s')$$

and

$$\begin{aligned}\mathcal{FH}_{\text{sc}}(e^{itA}We^{-itA_0}f)(\omega', s') &= e^{i\lambda_{s'}t} \mathcal{FH}_{\text{sc}}(We^{-itA_0}f)(\omega', s') \\ &= e^{i\lambda_{s'}t} \sum_{x \in V_q} (We^{-itA_0}f)(x) \overline{e(x, \omega', s')} \\ &= \sum_{x, y \in V_q} \int_{S^0} \int_{\Omega} e^{i(\lambda_{s'} - \lambda_s)t} W(x, y) e_0(y, \omega, s) \hat{f}(\omega, s) \overline{e(x, \omega', s')} d\sigma_O(\omega) d\mu(s) .\end{aligned}$$

Thus the expression in (24) (we set  $d\Sigma = d\sigma_O(\omega) d\mu(s) d\sigma_O(\omega') d\mu(s')$ ) is

$$\mathcal{L}(t) = \sum_{x, y \in V_q} \int_{S^0 \times S^0} \int_{\Omega \times \Omega} e^{i(\lambda_s - \lambda_{s'})t - \varepsilon|t|} \overline{V(x, y) e_0(y, \omega, s) \hat{f}(\omega, s)} e(x, \omega', s') \hat{g}(\omega', s') d\Sigma$$

and after doing the  $t$ -integration we get,

$$\langle f|(S-I)g\rangle = (-i)\lim_{\varepsilon \rightarrow 0} \int_{S^0 \times S^0} \int_{\Omega \times \Omega} T(\omega, s; \omega', s') \frac{2\varepsilon}{(\lambda_s - \lambda_{s'})^2 + \varepsilon^2} \overline{\hat{f}(\omega, s)} \hat{g}(\omega', s') d\Sigma .$$

We conclude by noticing as previously that the measures

$$dl_{\varepsilon} = \frac{2\varepsilon d\mu(s) d\mu(s')}{(\lambda_s - \lambda_{s'})^2 + \varepsilon^2}$$

converge weakly to  $2\pi\delta(\lambda_s - \lambda_{s'})$  as  $\varepsilon \rightarrow 0^+$ .

□

A consequence of the relation between  $T$  and  $S$  is the unitarity relation for  $T$ :

**Theorem 3.7** *Suppose  $\alpha \notin \mathcal{E}$ . Then for any  $s$  and  $s' \in S^+$  with  $\lambda_s = \lambda_{s'} = \alpha$ , and for any  $(\omega, \omega') \in \Omega \times \Omega$ ,*

$$\Im T(\omega, s; \omega', s') = \pi \int_{S^0} \int_{\Omega} \overline{T(\omega'', s''; \omega, s)} T(\omega'', s''; \omega', s') \delta(\lambda_{s''} - \alpha) d\mu(s'') d\sigma_O(\omega'') \quad (25)$$

*Proof.*–

By Theorem 3.6 we have

$$\overline{(\hat{S}f)}(\omega, s) = \hat{f}(\omega, s) - 2\pi i \int_{S^0} \int_{\Omega} T(\omega, s; \omega', s') \hat{f}(\omega', s') \delta(\lambda_s - \lambda_{s'}) d\sigma_O(\omega') d\mu(s') .$$

The adjoint of the map  $M : \hat{f} \rightarrow (\hat{S}f)$  is clearly given by

$$(M^*(g))(\omega, s) = g(\omega, s) + 2\pi i \int_{S^0} \int_{\Omega} \overline{T(\omega', s'; \omega, s)} g(\omega', s') \delta(\lambda_s - \lambda_{s'}) d\sigma_O(\omega') d\mu(s') .$$

The relation  $M^*M = I$ , which follows from  $S^*S = I$ , implies that (25) holds.

□

### 3.7 The S-matrix and the asymptotics of the deformed plane waves

**Theorem 3.8** *There exist “transmission coefficients”  $\tau(s, \omega, \omega')$  so that the solution of the Lippmann-Schwinger equation (14) writes*

$$e(x, \omega, s) = e_0(x, \omega, s) + \tau(s, \omega, \omega') q^{(-\frac{1}{2} + is)|x|}$$

for any  $x$  close enough to  $\omega'$ , and these coefficients are related to the scattering matrix by the following formula

$$S(\omega', -s; \omega, s) = -\frac{2i\pi}{C(s)} \tau(s, \omega, \omega') \quad (26)$$

with  $C(s) = \frac{1}{q^{\frac{1}{2}-is} - q^{-\frac{1}{2}+is}} .$

*Proof.*–

From the study of the Lippmann-Schwinger equation (14), we write the decomposition

$$e(x, \omega, s) = e_0(x, \omega, s) + e_{\text{scatt}}(x, \omega, s) ,$$

where

$$e_{\text{scatt}}(x, \omega, s) = \sum_{y \in K} G_0(\lambda_s, x, y) g(y, \omega, s)$$

where  $g(y, \omega, s) = \sum_{z \in K} W(y, z) e(z, \omega, s)$ .

Let us look at the asymptotic behaviour of  $e_{\text{scatt}}(x, \omega, s)$  as  $x \rightarrow \omega'$ . The Green's function  $G_0(\lambda_s; x, y)$  satisfies equation (4)

$$G_0(\lambda_s; x, y) = C(s) q^{(-\frac{1}{2} + is)d(x, y)};$$

then (5) and (6) imply that, if  $x \rightarrow \omega'$ ,

$$e_{\text{scatt}}(x, \omega, s) = \tau(s, \omega, \omega') q^{(-\frac{1}{2} + is)|x|} ,$$

with

$$\tau(s, \omega, \omega') = C(s) \sum_{y \in K} g(y, \omega, s) q^{(\frac{1}{2} - is)b_{\omega'}(y)} = C(s) \sum_{(y, z) \in K \times K} e(z, \omega, s) \overline{W(z, y)} e_0(y, \omega', s) .$$

Noticing that  $e_0(y, \omega', s) = \overline{e_0(y, \omega', -s)}$  we get that

$$\tau(s, \omega, \omega') = C(s) T(\omega', -s; \omega, s) .$$

and from (23) we derive formula (26).

□

**Remark 3.1** *The functions  $b_\omega(y)$  and  $b_{\omega'}(y)$  are equal if  $\omega$  and  $\omega'$  belong to the same end of  $\mathbb{T}_q \setminus K$ . This implies that the function  $\omega' \rightarrow \tau(s, \omega, \omega')$  is in fact constant in each end of  $\mathbb{T}_q \setminus K$ , so that the transmission coefficient  $\tau(s, \omega, \omega')$  can be written as a function  $\tau(s, \omega, l)$ . Moreover the reduced Lippmann-Schwinger equation depends only on the restriction of  $e_0$  to  $K$ , this implies that the function  $\omega \rightarrow \tau(s, \omega, l)$  is also constant in each end of  $\mathbb{T}_q \setminus K$ . Finally, we get an  $L \times L$  matrix depending on  $s$ , denoted by*

$$\tilde{S}(s) = (S(l', -s, l, s))_{l, l'} = -\frac{2i\pi}{C(s)} (\tau(s, l, l'))_{l, l'}$$

### 3.8 Computation of the transmission coefficients in terms of the Dirichlet-to Neumann operator

In this section, we compute the transmission coefficients following the method of [S].

We denote the distance between  $x$  and the root  $O$  of  $\mathbb{T}_q$  by  $|x|$ , and by  $B_{n-1}$  the ball of radius  $n-1$  where  $n-2$  is the supremum of  $|x|$  for  $x$  in  $K$ . We denote by  $T_l$  the ends of  $\mathbb{T}_q \setminus B_{n-1}$  ( $1 \leq l \leq L$ ), by  $x_l$  the root of  $T_l$  and by  $\Omega_l$  the boundary of  $T_l$ , which consists in the set of all geodesic rays starting from  $x_l$  and staying into  $T_l$ . The set of the roots  $\{x_l | l = 1, \dots, L\}$  is the circle of radius  $n$  and  $L = (q+1)q^{n-1}$ . From now on, we consider a fixed  $l$  ( $1 \leq l \leq L$ ), a fixed geodesic ray  $\omega$  in  $\Omega_l$ , and the associated "incoming plane wave"

$$\forall x \in V_q, \quad e_0(x, \omega, s) = q^{(1/2-is)b_\omega(x)},$$

where  $s \in S^0$ . Such a plane wave is a generalised eigenfunction for the adjacency operator  $A_0$  on  $\mathbb{T}_q$  in the sense that it satisfies

$$(\lambda_s - A_0)e_0(x, \omega, s) = 0 \quad (\lambda_s = 2\sqrt{q} \cos(s \log q)) ,$$

but is not in  $l^2$ . We are looking for solutions

$$e(., \omega, s) = e_0(., \omega, s) + e_{scat}(., \omega, s), \quad x \in V_q$$

of the equation

$$(\lambda_s - A)e(., \omega, s) = 0, \tag{27}$$

where the scattered wave  $e_{scat}(., \omega, s)$  satisfies:

$$e_{scat}(x, \omega, s) = \tau(s, \omega, l')\Phi_s(x) \quad \text{if} \quad x \in V(T_{l'}), \tag{28}$$

where

$$\Phi_s(x) = q^{(-1/2+is)|x|}$$

(the so-called radiation condition) and the coefficients  $\tau(s, \omega, l')$  are the transmission coefficients. These radial waves are generalised eigenfunctions of  $A_0$  in the sense defined previously. We want to get an explicit expression of the transmission vector

$$\overrightarrow{\tau(s, \omega)} := (\tau(s, \omega, 1), \dots, \tau(s, \omega, l'), \dots, \tau(s, \omega, L)). \tag{29}$$

As we shall see, the transmission vector does not depend on the choice of the geodesic ray  $\omega$ , it is uniquely determined by the choice of  $l$ ; we define

$$\overrightarrow{\tau(s, l)} := \overrightarrow{\tau(s, \omega)}, \quad \forall \omega \in \Omega_l. \tag{30}$$

We thus recover the result of the previous section, with the following relation

$$\forall l, l' \quad \tau(s, l, l') = -\frac{C(s)}{2i\pi} S(l', -s, l, s) .$$

We begin with noticing that  $b_\omega(x_{l'})$  does not depend of  $\omega \in \Omega_l$  for any  $l' \in \{1, \dots, L\}$ . We set

$$\vec{A}_l := (\alpha^{-b_\omega(x_1)}, \dots, \alpha^{-b_\omega(x_L)}) = (\alpha^{-b_l(x_1)}, \dots, \alpha^{-b_l(x_L)}) \quad (\alpha = q^{-1/2+is}),$$

and denote by  $\vec{E}_l$  the vector in  $\mathbb{R}^L$  having all null coordinates excepted the  $l$ -th coordinate, which is equal to 1.

We have the following

**Theorem 3.9** *Consider the integer  $n$  so that  $B_{n-2}$  is the smallest ball containing the finite graph  $K$ .*

*Set  $\Gamma = B_n$ ,  $\partial\Gamma = \{x_{l'}, 1 \leq l' \leq L\}$ , denote by  $\widehat{A}_n$  the restriction of  $A$  to  $B_n$  in the sense that  $\widehat{A}_n = (A_{x,y})_{(x,y) \in B_n}$ , define  $I_n$  in the same way, set  $B = \widehat{A}_n - \lambda_s I_n$  and denote by  $\mathcal{DN}_s$  the corresponding Dirichlet-to Neumann operator (see Definition 4.3, Appendix B).*

*Then  $\mathcal{DN}_s$  and the transmission vector  $\overrightarrow{\tau(s,l)}$  defined by (28), (29) and (30) exist for any*

$$s \notin \mathcal{E} = \{s \in S^0, \lambda_s \in \sigma(\widehat{A}_{n-1})\}$$

and

$$(\tau(s, l, l')) = -\alpha^{-2n} \left[ \frac{1}{C(s)} (\mathcal{DN}_s + q^{1/2+is} I)^{-1} + \mathcal{A} \right],$$

with  $\widehat{A}_{n-1} = (A_{x,y})_{(x,y) \in B_{n-1}}$ ,  $\mathcal{A} = (\mathcal{A}_{l,l'}) = (\alpha^{d(x_l, x_{l'})})$ ,  $\alpha = q^{-1/2+is}$ .

*Proof.*—

Let us recall that we have fixed  $l$  ( $1 \leq l \leq L$ ) and a geodesic ray  $\omega$  in  $\Omega_l$ . From now on we write  $e(x)$  instead of  $e(x, \omega, s)$  for any  $x \in V_q$  for simplicity.

Equation (27) splits into 3 expressions, depending on where  $x$  is taken.

- If  $x \notin B_n$  the equation is already verified, since  $A$  coincide with  $A_0$  on each end  $T_{l'}$ .
- if  $x \in B_{n-1}$ , the equation becomes the Dirichlet problem  $Ae(x) = \lambda_s e(x)$ ,  $x \in B_{n-1}$ ,  $e|_{\partial\Gamma} = f$ , and is uniquely solvable for any  $s$  outside  $\mathcal{E}$ , the prescribed values of the function  $f$  at the boundary being given by the  $L$ -vector

$$\vec{e}_l =: (e(x_1), \dots, e(x_L))$$

to be determined in the sequel.

- if  $x \in B_n \setminus B_{n-1}$  (which means that  $x$  is one of the roots  $x_{l'}$ ) equation (27) writes:

$$\forall l' \in \{1, \dots, L\}, \quad e(x_{l'}^-) + \sum_{x \sim x_{l'}, x \in T_{l'}} e(x) = \lambda_s e(x_{l'})$$

where  $x_{l'}^-$  is the unique interior neighbor of  $x_{l'}$ , and where we used that the potential  $W$  vanishes outside  $B_{n-2}$ .

According to definition 4.3, the corresponding Dirichlet-to Neumann operator  $\mathcal{DN}_s$  writes

$$\mathcal{DN}_s(\vec{e}_l)(x_{l'}) = e(x_{l'}^-) - \lambda_s e(x_{l'}) \quad \forall l' \in \{1, \dots, L\},$$

Therefore, if  $s \notin \mathcal{E}$ , Equation (3.8) can be rewritten as,

$$\forall l' \in \{1, \dots, L\}, \quad \mathcal{DN}_s(\vec{e}_l)(x_{l'}) + \sum_{x \sim x_{l'}, x \in T_{l'}} e(x) = 0 \quad (31)$$

Now it remains to compute  $\sum_{x \sim x_{l'}, x \in T_{l'}} e(x)$ . We have, for  $x \in T_{l'}$ ,  $e(x) = q^{(1/2-is)b_\omega(x)} + \tau(s, \omega, l')\Phi_s(x)$ . According to the expression of the radial function  $\Phi_s$ , we get, for any  $s \notin \mathcal{E}$  and  $l' \in \{1, \dots, L\}$ , that

$$\begin{aligned} \Phi_s(x_{l'}) &= q^{(-1/2+is)|x_{l'}|} = q^{n(-1/2+is)} \\ \Phi_s(x) &= q^{(-1/2+is)(n+1)} \quad \forall x \in T_{l'} \quad x \sim x_{l'}. \end{aligned}$$

Let us write the set  $N_l = \{x \in T_l, \quad x \sim x_l\}$  as  $N_l = \{y_l\} \cup \tilde{N}_l$ , where  $y_l$  belongs to the infinite path  $\omega$  whereas the  $q-1$  vertices of  $\tilde{N}_l$  do not. Then, using the properties of the Busemann function we have

$$\begin{aligned} b_\omega(x_l) &= b_l(x_l) = |x_l| = n \\ b_\omega(y_l) &= n+1 \\ b_\omega(x) &= b_l(x_l) - 1 = n-1, \quad \forall x \in \tilde{N}_l \\ b_\omega(x) &= b_l(x_{l'}) - 1 \quad \forall x \in N_{l'}, \quad l' \neq l. \end{aligned}$$

So we get for any  $l' \in \{1, \dots, L\}$  and after setting  $\alpha = q^{-1/2+is}$ ,

$$\begin{aligned} e(x_{l'}) &= \alpha^{-b_l(x_{l'})} + \tau(s, \omega, l')\alpha^n \\ e(x) &= \alpha^{-b_l(x_{l'})+\varepsilon} + \tau(s, \omega, l')\alpha^{n+1} \end{aligned}$$

with

$$\begin{aligned} \varepsilon &= 1 \quad \forall x \in \tilde{N}_l \\ \varepsilon &= 1, \quad \forall x \in N_{l'} \quad l' \neq l \\ \varepsilon &= -1 \quad \text{if } x = y_l. \end{aligned}$$

Hence we have

$$\sum_{x \sim x_{l'}, x \in T_{l'}} e(x) = q\alpha^{-b_l(x_{l'})+1} + q\tau(s, \omega, l')\alpha^{n+1} \quad \text{if } l' \neq l$$

$$\sum_{x \sim x_l, x \in T_l} e(x) = (q-1)\alpha^{-b_l(x_l)+1} + \alpha^{-b_l(x_l)-1} + q\tau(s, \omega, l')\alpha^{n+1}.$$

These equations can be summarised, for any  $l' \in \{1, \dots, L\}$ , as

$$\sum_{x \sim x_{l'}, x \in T_{l'}} e(x) = \alpha^{-b_l(x_{l'})+1}[q + \delta_{ll'}(\alpha^{-2} - 1)] + q\tau(s, \omega, l')\alpha^{n+1}$$

so that equation (31) gives, for any  $l' \in \{1, \dots, L\}$

$$\mathcal{DN}_s(\vec{e}_l)(x_{l'}) + \alpha^{-b_l(x_{l'})+1}[q + \delta_{ll'}(\alpha^{-2} - 1)] + q\tau(s, \omega, l')\alpha^{n+1} = 0. \quad (32)$$

Let us set  $\mathcal{DN}_s(\vec{e}_l) = (\mathcal{DN}_s(\vec{e}_l)(x_1), \dots, \mathcal{DN}_s(\vec{e}_l)(x_L))$ , and write  $\vec{e}_l = (e(x_1), \dots, e(x_L)) = \vec{A}_l + \alpha^n \overrightarrow{\tau(s, \omega)}$  (recall that  $\vec{A}_l$  and  $\overrightarrow{\tau(s, \omega)}$  are  $L$ -vectors having respectively  $\alpha^{-b_l(x_{l'})}$  and  $\tau(s, \omega, l')$  as their  $l'$ -coordinate).

Substituting in (32) and denoting by  $\vec{E}_l$  the vector in  $\mathbb{R}^L$  having all null coordinates except  $x_l = 1$ , we get

$$\mathcal{DN}_s[\vec{A}_l + \alpha^n \overrightarrow{\tau(s, \omega)}] + q\alpha \vec{A}_l + \alpha^{-n}(\alpha^{-1} - \alpha)\vec{E}_l + q\overrightarrow{\tau(s, \omega)}\alpha^{n+1} = 0$$

which yields

$$\alpha^n (\mathcal{DN}_s + q\alpha I) \overrightarrow{\tau(s, \omega)} = \alpha^{-n}(\alpha - \alpha^{-1})\vec{E}_l - (q\alpha I + \mathcal{DN}_s) \vec{A}_l.$$

Using the expression of  $\alpha$  and  $C(s)$ , we have then

$$\alpha^n (\mathcal{DN}_s + q^{1/2+is} I) \overrightarrow{\tau(s, \omega)} = -\frac{\alpha^{-n}}{C(s)} \vec{E}_l - (q^{1/2+is} I + \mathcal{DN}_s) \vec{A}_l.$$

Since the matrix  $\mathcal{DN}_s$  is real symmetric,  $\mathcal{DN}_s + q^{1/2+is} I$  is an invertible matrix for any  $s \in S^0$  so that  $\lambda_s \notin \sigma(\hat{A}_{n-1})$ , and

$$\overrightarrow{\tau(s, \omega)} = -\frac{\alpha^{-2n}}{C(s)} (\mathcal{DN}_s + q^{1/2+is} I)^{-1} \vec{E}_l - \alpha^{-n} \vec{A}_l.$$

We conclude the proof by noticing that, for any  $l' \in \{1, \dots, L\}$ ,  $b_l(x_{l'}) = n - d(x_l, x_{l'})$ .

□

### 3.9 Density of states

**Theorem 3.10** *When the vertex  $x$  tends to infinity, the densities of states  $\mathbf{d}\mu_x^A$  of  $\Gamma$  converge weakly to the density of states  $\mathbf{de}$  of  $\mathbb{T}_q$  computed in Section 2.4.*

*Proof.*—

It is enough to compute the limits of  $\int \lambda^n d\mu_x^A$  for  $n$  fixed and  $x \rightarrow \infty$ . By definition, we have  $\int t^n d\mu_x^A = [A^n](x, x)$ , and

$$[A^n](x, x) = \sum a_{x, x_1} a_{x_1, x_2} \cdots a_{x_{n-1}, x}$$

where the sum is on loops  $\gamma = (x, x_1, x_2, \dots, x_{n-1}, x)$  of length  $n$  based at  $x$ . If we assume that  $d(x, K) > n/2$ , the loops do not meet  $K$  and therefore  $[A^n](x, x) = [A_0^n](x, x)$ .

□

## 4 The spectral theory for a graph asymptotic to an homogeneous tree

We are concerned here with the spectral theory of the adjacency matrix of a graph  $\Gamma$  asymptotic to a homogeneous tree of degree  $q + 1$ , in the sense of Definition 1.1. We want to reduce the spectral theory of  $A_\Gamma$  to the situation studied in Section 3. For that, we need a preliminary combinatorial study which could be of independent interest.

### 4.1 Some combinatorics

We need the following combinatorial result:

**Theorem 4.1** *If  $\Gamma$  is asymptotic to a homogeneous tree of degree  $q + 1$ , then  $\Gamma$  is isomorphic to a connected component of a graph  $\hat{\Gamma}$  which can be obtained from  $\mathbb{T}_q$  by adding and removing a finite number of edges.*

In order to prove Theorem 4.1, we first introduce an integer  $\nu(\Gamma)$  associated to the graph  $\Gamma$ ; the integer  $\nu$  is a combinatorial analogue of the regularised total curvature of a Riemannian surface  $S$  which is of constant curvature  $\equiv K_0$  near infinity, namely  $\int_S (K - K_0) |d\sigma|$ .

**Definition 4.1** *If  $\Gamma$  is asymptotic to a homogeneous tree of degree  $q + 1$ , we define  $\nu(\Gamma)$  by*

$$\nu(\Gamma) = \sum_{x \in V_\Gamma} (q + 1 - d(x)) + 2b_1 ,$$

where  $d(x)$  is the degree of the vertex  $x$  and  $b_1$  is the first Betti number of  $\Gamma$  or equivalently the number of edges to be removed from  $\Gamma$  in order to get a tree.

Note that, if  $T$  is a maximal sub-tree of  $\Gamma$ ,  $\nu(T) = \nu(\Gamma)$ .

We will need the

**Lemma 4.1** *If, for  $r \geq 2$ ,  $B_r = \{x \in V_\Gamma \mid |x|_{\Gamma_0} \leq r\}$ , then we have*

$$\nu(\Gamma) = (q-1)m - M + 2 ,$$

where  $m$  is the number of inner vertices of  $B_r$  and  $M$  the number of boundary vertices (i.e. connected to a vertex of  $\Gamma \setminus B_r$ ) of  $B_r$ .

*Proof.*–

The  $M$  boundary vertices have  $q$  neighbours in  $\Gamma \setminus B_r$  and one in  $B_r$ . From Euler formula applied to  $\Gamma \cap B_r$ , we get

$$1 - b_1 = (m + M) - \frac{1}{2} \left( \sum_{|x|_{\Gamma_0} \leq r-1} d(x) + M \right) .$$

Thus

$$\nu(\Gamma) = \sum_{|x|_{\Gamma_0} \leq r-1} (q+1 - d(x)) + 2b_1$$

is equal to

$$\nu(\Gamma) = (q+1)m - (2m + M - 2 + 2b_1) + 2b_1 .$$

□

We will also need the:

**Lemma 4.2** *Let  $F$  be a finite tree whose all vertices are of degree  $q+1$  except the ends which are of degree 1. Let  $M$  be the number of ends and  $m$  the number of inner vertices of  $F$ . We have the relation*

$$M = 2 + (q-1)m . \tag{33}$$

*Conversely, for each choice of  $(m, M)$  satisfying Equation (33), there exists such a tree  $F$ .*

*Proof.*–

From Euler formula applied to  $F$ , we get  $1 = |V_F| - |E_F|$ . Moreover  $|V_F| = m + M$ . Let us choose a root inside  $F$  and orient the edges from that root. Then we count the edges by partitioning them with their  $m$  possible origins; this gives  $|E_F| = (q+1) + (m-1)q$ .

Conversely, the statement is true for  $m = 1$ ,  $M = q+1$  and we proceed by induction on  $m$  by adding  $q$  edges to a boundary vertex and the corresponding  $q$  boundary vertices, we have  $m \rightarrow m+1$ ,  $M \rightarrow M + (q-1)$ .

□

We will now make some modifications of  $\Gamma$  in order to get a new graph  $\hat{\Gamma}$  with  $\nu(\hat{\Gamma}) = 0$ .

**Lemma 4.3** *If  $\Gamma' = M_1(\Gamma)$  is defined by adding to  $\Gamma$  a vertex and an edge connecting that vertex to a vertex of  $\Gamma_0$ , then  $\nu(\Gamma') = \nu(\Gamma) + q - 1$ .*

*If  $\Gamma' = M_2(\Gamma)$  is defined by adding to  $\Gamma$  a tree whose root  $x$  is of degree  $q$  and all other vertices of degree  $q + 1$  and connecting  $x$  by an edge to a vertex of  $\Gamma_0$ ,  $\Gamma'$  is asymptotic to an homogeneous tree of degree  $q + 1$  and  $\nu(\Gamma') = \nu(\Gamma) - 1$ .*

This Lemma is quite easy to check.

*The proof of Theorem 4.1:* let us now write  $\nu(\Gamma) = N'' - (q - 1)N'$  with  $N' \geq 0$  and  $N'' \geq 0$ . By performing  $N'$  times the move  $M_1$  and  $N''$  times the move  $M_2$ , we arrive to a graph  $\hat{\Gamma}'$  with  $\nu(\hat{\Gamma}') = 0$ . Let  $\hat{\Gamma}$  be the graph obtained by removing from  $\hat{\Gamma}'$  the  $(N' + N'')$  edges not in  $E_\Gamma$ , one of whose vertices is in  $\Gamma_0$ . Then  $\Gamma$  is a connected component of  $\hat{\Gamma}$ .

We apply Lemma 4.2 and change  $\hat{\Gamma}'$  inside  $B_r$  by gluing to the boundary vertices a tree  $F$ . We get this way a tree  $\mathbb{T}_q$  which, by Lemma 4.2 has the same number of vertices inside  $B_r$  than  $\hat{\Gamma}'$ , so that one can recover  $\hat{\Gamma}'$  by removing and adding a finite number of edges to  $\mathbb{T}_q$ . Removing more edges, we get  $\hat{\Gamma}$  and  $\Gamma$  is a connected component of  $\hat{\Gamma}$ .

In particular  $\Gamma$  is isomorphic to a graph obtained from  $\mathbb{T}_q$  by removing or adding a finite number of edges if and only if  $\nu(\Gamma) = 0$ .

## 4.2 The spectral theory of $\Gamma$

From Theorem 4.1, we deduce the existence of a Hilbert space  $\mathcal{H}$  so that  $l^2(\hat{\Gamma}) = l^2(\Gamma) \oplus \mathcal{H}$  and this decomposition is invariant by  $A_{\hat{\Gamma}}$ . Moreover  $A_{\hat{\Gamma}}$  is a finite rank perturbation of  $A_0 = A_{\mathbb{T}_q}$ . This will allow us to describe the spectral theory of  $A_\Gamma$  by using the results of Section 3.

In order to get the spectral decomposition of  $A_\Gamma$  in terms of the spectral decomposition of  $A_{\hat{\Gamma}}$  given in Section 3.4, we will need the

**Lemma 4.4** *Let  $A_{\hat{\Gamma}} = A_{\mathbb{T}_q} + W$  with  $\text{Support}(W) \subset K \times K$  and  $K$  finite. Let  $\Gamma$  be an unbounded connected component of  $\hat{\Gamma}$  and  $\omega$  a point at infinity of  $\Gamma$ . Then, for any  $s \notin \hat{\mathcal{E}}$ , we have*

$$\text{support}(e(., s, \omega)) \subset V_\Gamma .$$

*Conversely, if  $\omega'$  is a point at infinity of  $\hat{\Gamma}$  which is not a point at infinity of  $\Gamma$  then*

$$\text{support}(e(., s, \omega')) \cap V_\Gamma = \emptyset .$$

Let us prove the first assertion, the proof of the second is similar. It is enough to prove it for  $s \in S^+$  close to  $S_0$ , because  $s \rightarrow e(x, s, \omega)$  is meromorphic on  $S$ . We have then

$$e(s, \omega) = e_0(s, \omega) + G_0(\lambda_s) W e(s, \omega) .$$

From the explicit expression of  $e_0$ , we get that the first term belongs to  $l^2(\hat{\Gamma} \setminus \Gamma)$ , and so does the second one, as the image of a compactly supported function by the Green's function for  $\lambda_s \notin I_q = \text{spectrum}(A_0)$ . This proves that the restriction of  $e(\cdot, s, \omega)$  to  $V_{\hat{\Gamma}} \setminus V_{\Gamma}$  is an  $l^2$  eigenfunction, with eigenvalue  $\lambda_s$ , of  $A_{\hat{\Gamma}}$ . Since  $A_{\hat{\Gamma}}$  has no eigenvalue  $\lambda_s$  for  $s \in S^+$  close to  $S_0$ , it follows that  $e(x, s, \omega)$  vanishes for  $s \in S^+$  close to  $S_0$  and  $x \notin V_{\Gamma}$ .

Theorem 4.1 allows to consider the set  $\Omega$  of points at infinity of  $\Gamma$  as a subset of the set  $\hat{\Omega}$  of the points at infinity of  $\hat{\Gamma}$ . The space  $l^2(\hat{\Gamma})$  splits as a direct sum  $l^2(\Gamma) \oplus l^2(\hat{\Gamma} \setminus \Gamma)$  which is preserved by the adjacency matrix. Lemma 4.4 shows that the support of the generalised eigenfunctions  $e(\cdot, s, \omega)$  for  $\omega \in \Omega$  is included in  $V_{\Gamma}$ . Using this, we can state the spectral decomposition of  $A_{\Gamma}$  as an immediate corollary of Theorem 3.3:

**Theorem 4.2** *The Hilbert space  $l^2(\Gamma)$  splits into a finite dimensional part  $\mathcal{H}_{\text{pp}}$  and an absolutely continuous part  $\mathcal{H}_{\text{ac}}$ . This decomposition is preserved by  $A_{\Gamma}$ . If  $f \in C_0(\Gamma)$  and, for  $\omega \in \Omega$ ,  $\hat{f}(s, \omega) = \langle e(\cdot, s, \omega), f \rangle$ , then the map  $f \rightarrow \hat{f}$  extends to an isometry from  $\mathcal{H}_{\text{ac}}$  onto  $L^2_{\text{even}}(S_0 \times \Omega, d\sigma_0 \otimes d\mu)$  which intertwines the action of  $A_{\gamma}$  with the multiplication by  $\lambda_s$ .*

## Appendix A: delta measures

The goal of this Appendix is to define in a precise way the meaning of the measures  $d\mu = \delta(S = 0)d\nu$  where  $d\nu = a(x)dx$  is absolutely continuous w.r. to the Lebesgue measure in  $\mathbb{R}^d$  and  $S$  is a  $C^1$  real valued function so that  $dS$  does not vanish on the hypersurface  $S = 0$ . The measure  $d\mu = \delta(S = 0)d\nu$  is supported by the hypersurface  $\Sigma := \{S = 0\}$ .

We can assume that  $\mathbb{R}^d$  and the hypersurface  $S = 0$  are oriented, so that we can play with differential forms instead of measures.

The proof of the following Lemma is left to the reader:

**Lemma 4.5** *There exists a differential form  $\beta$  defined in some neighborhood of  $\Sigma$  so that  $adx_1 \wedge \cdots \wedge dx_d = dS \wedge \beta$ . Moreover the restriction of  $\beta$  to  $\Sigma$  is uniquely defined.*

**Definition 4.2** *If  $\nu$  is the measure on  $\Sigma$  associated to the restriction of  $\beta$  to  $\Sigma$ . we can view  $\nu$  as a measure on  $\mathbb{R}^d$  denoted  $d\nu = \delta(S = 0)d\mu$ .*

We can view  $d\nu$  as weak limits: if  $f : \mathbb{R} \rightarrow \mathbb{R}^+$  is a positive  $L^1$  function of integral 1 and  $f_{\varepsilon}(t) = \varepsilon^{-1}f(\varepsilon^{-1}t)$  the measure  $\delta(S = 0)d\mu$  is the weak limit as

$\varepsilon \rightarrow 0$  of the measures  $d\nu_\varepsilon = f_\varepsilon(t)d\mu$  (For the proof, take local coordinates so that  $S = x_1$ ).

Usual choices are  $f_1$  the characteristic function of the interval  $[-\frac{1}{2}, \frac{1}{2}]$  and  $f_2(t) = \frac{1}{\pi} \frac{1}{1+t^2}$ .

## Appendix B: the Dirichlet-to Neumann operator $\mathcal{DN}$ on a finite graph

Let  $\Gamma = (V, E)$  be a connected finite graph and let  $\partial\Gamma$  be a subset of  $V$  called the "boundary of  $\Gamma$ ". Let  $B = (b_{i,j}) : \mathbb{R}^V \rightarrow \mathbb{R}^V$  be a symmetric matrix associated to  $\Gamma$ , namely

$$b_{i,j} = 0 \quad \text{if } i \neq j \quad \text{and} \quad \{i, j\} \notin E.$$

After setting  $V_0 = V \setminus \partial\Gamma$ , we define  $B_0 : \mathbb{R}^{V_0} \rightarrow \mathbb{R}^{V_0}$  as the restriction of  $B$  to the functions which vanish on  $\partial\Gamma$ .

We have the following

**Lemma 4.6** *Assume that  $B_0$  is invertible. Then, for any given  $f \in C(\partial\Gamma)$ , there exists a unique solution  $F \in C(\Gamma)$  of the Dirichlet problem*

$$(D_f) : F|_{\partial\Gamma} = f \text{ and } BF(l) = 0 \text{ if } l \in V_0, F|_{\partial\Gamma} = f.$$

The *Dirichlet-to Neumann operator*  $\mathcal{DN}$  associated to  $B$  is the linear operator from  $C(\partial\Gamma)$  to  $C(\partial\Gamma)$  defined as follows:

**Definition 4.3** *Assume that  $B_0$  is invertible. Let  $f \in C(\partial\Gamma)$ , and  $F$  be the unique solution of the Dirichlet problem  $(D_f)$ . Then, the Dirichlet-to-Neumann operator form  $\mathcal{DN} : \mathbb{R}^{\partial\Gamma} \rightarrow \mathbb{R}^{\partial\Gamma}$  is defined as follows: if  $l \in \partial\Gamma$ ,*

$$\mathcal{DN}(f)(l) = \sum_{i=1}^m b_{l,i} F(i) (= BF(l)).$$

## References

- [Bre] J. Breuer. *Singular continuous spectrum for the Laplacian on certain sparse trees*. Commun. Math. Phys. **269** (3):851–857 (2007).
- [CdV1] Y. Colin de Verdière. *Spectre de graphes*. Cours spécialisés **4**, Société mathématique de France (1998).
- [CdV2] Y. Colin de Verdière. *Distribution de points sur une sphère*. Séminaire N. Bourbaki, exposé **703**:83–93 (1988-89).
- [CdV3] Y. Colin de Verdière. *Mathematical models for passive imaging I: general background*. ArXiv 0610043.

- [CdV4] Y. Colin de Verdière. *Semiclassical analysis and passive imaging*. Nonlinearity **22**:45–75 (2009).
- [CdV5] Y. Colin de Verdière. *A Semi-classical calculus of correlations*. Thematic issue “Imaging and Monitoring with Seismic Noise” of the series “Comptes Rendus Géosciences”, from the Académie des sciences **343**:496–501 (2011).
- [CMS] M. Cowling, S. Meda & A. Setti. *An overview of harmonic analysis on the group of isometries of a homogeneous tree*. Exposition.Math., **16(5)**:385–423 (1998).
- [CS] M. Cowling & A. Setti. *The range of the Helgason-Fourier transformation on homogeneous trees*. Bull.Austral.Math.Soc., **59**:237–246 (1998).
- [FN] A. Figà-Talamanca & C. Nebbia, *Harmonic Analysis and representation theory for groups acting on homogeneous trees*. London Math. Soc. Lecture Notes Series, **162** Cambridge Univ. Press, 1991.
- [Ike] T. Ikebe. *Eigenfunction expansion associated with the Schrödinger operators and their applications to scattering theory*. Arch. Rational Mech. Anal., **5**:1–34 (1960).
- [RS] M. Reed & B. Simon. *Methods of Modern mathematical Physics III-Scattering theory*, (1980), New York, Academic Press.
- [S] U. Smilansky *Exterior-Interior Duality for Discrete Graphs* J. Phys. A: Math. Theor., **42**:035101 (2009).