



# Determinacy and Stability under Learning of Rational Expectations Equilibria

Stéphane Gauthier

## ► To cite this version:

Stéphane Gauthier. Determinacy and Stability under Learning of Rational Expectations Equilibria. Journal of Economic Theory, 2002, 102 (2), pp.354-374. 10.1006/jeth.2000.2768 . hal-00731065

**HAL Id: hal-00731065**

**<https://hal.science/hal-00731065>**

Submitted on 11 Sep 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Determinacy and Stability under Learning of Rational Expectations Equilibria<sup>□</sup>

Stéphane Gauthier<sup>γ</sup>

DELTA (Joint Research Unit CNRS-EHESS-ENS)

June 30, 2000

## Abstract

This paper studies relationships between the local determinacy of a stationary equilibrium in the perfect foresight dynamics, and its local stability in dynamics arising from econometric learning procedures. Attention is focused on linear scalar economies where agents forecast only one period ahead, and with an arbitrary, but fixed, number of predetermined variables. In such a framework, it is well known that there are no clear links between the determinacy of the stationary state in the perfect foresight dynamics on the levels of the state variable, and its stability under learning. The paper emphasizes, however, that this is not the right perfect foresight dynamics to look at whenever agents try to learn the coefficients of the perfect foresight dynamics restricted to an eigenspace of lower dimension. Indeed the paper introduces a growth rate perfect foresight dynamics on these coefficients and proves equivalence between determinacy in that dynamics and stability under learning provided that a simple sign condition is satisfied.

Journal of Economic Literature Classification Numbers: E32, D83

Key words: macroeconomic dynamics, rational expectations, determinacy, learning, expectational stability.

---

<sup>□</sup> I would like to thank my thesis supervisor Roger Guesnerie for his advices and Jean-Michel Grandmont for the kind interest he paid to my work. I am also grateful to Antoine d'Autume and Gabriel Desgranges for helpful discussions, and to an anonymous referee of this journal for his guidance in writing the paper. Of course the usual disclaimers apply.

<sup>γ</sup> DELTA, 48, bd Jourdan, 75014 Paris, France; phone number: 33 1 43 13 63 11; fax number: 33 1 43 13 63 10; e-mail address: gauthier@delta.ens.fr

# 1 Introduction

It is a commonplace knowledge that dynamic models with rational expectations exhibit a multiplicity of equilibrium paths. Several alternative devices have consequently been proposed for the selection of the solutions on which attention should be focused in practice. This paper is an attempt to confront two among these, the determinacy of an equilibrium and its stability in a learning dynamics. This purpose is related to Guesnerie's [9] dynamic equivalence principle which claims that local determinacy should be equivalent to local stability under learning provided that agents form "reasonably" their forecasts (see also Lucas [13]), and temporary equilibrium literature provides indeed many examples of reasonable learning rules (Grandmont and Laroque [6], Grandmont and Laroque [7] or Guesnerie and Woodford [10]). Nevertheless, the equivalence principle fails when agents employ recursive (econometric) learning rules (Duay [1], Evans and Honkapohja [3], Grandmont [5] or Grandmont and Laroque [8]) such as the ordinary least squares algorithm (Marcet and Sargent [14]), the Robbins and Monro [17] scheme (Woodford [20]), or the gradient one (Evans and Honkapohja [4]). Most of these are studied in Ljung and Soderström [12]. Our aim is to show that the determinacy criterion is not applied in a suitable way in these cases.

The intuition is easy to grasp in a linear one step forward looking economy where the current (univariate) state depends on  $L = 1$  predetermined variable (Reichlin [16], Woodford [18]). In such a model, the dynamics with perfect foresight is governed by two (local perfect foresight eigenvalues) growth rates  $\lambda_1$  and  $\lambda_2$  (with  $j_{\lambda_1} < j_{\lambda_2}$ ) in the immediate vicinity of a stationary state  $\bar{x}$  (where the state variable is equal to  $\bar{x}$  at all times). Usually the determinacy criterion is applied to  $\bar{x}$  in the perfect foresight dynamics on the levels of the state variable, whereas agents try to estimate  $\lambda_1$  and  $\lambda_2$  through a standard econometric procedure, the asymptotic behavior of which can be approximated by suitable continuous differential equations involving the expectational stability criterion (Evans [2]). It turns out that agents discover the growth rate of least modulus  $\lambda_1$  in such a specification, so that the learning dynamics may be stable ( $j_{\lambda_1} < 1$ ) even if  $\bar{x}$  is locally indeterminate ( $j_{\lambda_2} < 1$ ). One may wonder, however, whether this is the right perfect foresight dynamics to look at. The main innovation of the paper is indeed to apply the determinacy criterion to the fixed points in the learning dynamics ( $\lambda_1$  and  $\lambda_2$ ) by defining a perfect foresight dynamics of growth rates whose stationary equilibria are the perfect foresight growth rates. The outcome is particularly appealing within this simple economy (with only one predetermined variable) since a perfect foresight growth rate is stable under learning if and only if it is determinate in the perfect foresight dynamics of growth rates so defined.

In the more general framework where the current state depends on expectations of the next state and on an arbitrary, but fixed, number  $L \geq 0$  of predetermined variables,

the dynamics with perfect foresight involves  $(L + 1)$  local perfect foresight eigenvalues  $\lambda_{j_1}, \dots, \lambda_{j_{L+1}}$  (with  $j_{j_1} < \dots < j_{j_{L+1}}$ ) in the neighborhood of  $\bar{x}$ . In that case, agents are supposed to try to learn the  $L$  coefficients of the linear perfect foresight dynamics restricted to a  $L$ -dimensional eigenspace spanned by  $\prod_{i=1}^L$  eigenvectors among those associated with the  $(L + 1)$  local eigenvalues. When  $\lambda_{j_{L+1}}$  and  $\prod_{i=1}^{L+1} \lambda_{j_i}$  have the same sign (which must be the case if  $L = 1$ ), the eigenspace that is locally stable in the learning dynamics corresponds to the eigenvalues  $\lambda_{j_1}, \dots, \lambda_{j_L}$  of lowest modulus. Hence  $\bar{x}$  is locally stable in the learning dynamics if and only if  $j_{j_L} < 1$ , which encompasses both the saddle point determinate configuration ( $1 < j_{j_{L+1}}$ ) and the indeterminate configuration ( $j_{j_{L+1}} < 1$ ), for the perfect foresight dynamics. When  $\lambda_{j_{L+1}}$  and  $\prod_{i=1}^{L+1} \lambda_{j_i}$  have opposite signs, the eigenspace that is locally stable under learning always includes the eigenvector associated with  $\lambda_{j_{L+1}}$ , and the learning dynamics will be stable if and only if  $j_{j_{L+1}} < 1$ , i.e., in the locally indeterminate configuration for the perfect foresight dynamics. These results imply again that the learning dynamics may be stable even if  $\bar{x}$  is locally indeterminate in the perfect foresight dynamics on the levels of the state variable. Nevertheless one should instead look at a new perfect foresight dynamics, the extended growth rate perfect foresight dynamics, defined on the  $L$ -dimensional vectors of the coefficients that agents try to estimate. The issue is whether there is a neater relation between local stability under the considered class of learning algorithms, and local determinacy in this extended growth rate perfect foresight dynamics, of a particular eigenspace or of the  $L$ -dimensional vector of coefficients associated to it. The outcome here is still very simple since the  $L$ -dimensional eigenspace corresponding to the  $L$  perfect foresight growth rates  $\lambda_{j_1}, \dots, \lambda_{j_L}$  of lowest modulus, is the only one to be locally determinate. Therefore, if  $\lambda_{j_{L+1}}$  and  $\prod_{i=1}^{L+1} \lambda_{j_i}$  have the same sign (a condition that may be related to the one sided sign condition for stability of differential equations, and that is always satisfied when  $L = 1$ ), one gets indeed equivalence between the local determinacy of a particular eigenspace, or of the associated vector, and its local stability under learning. However this equivalence fails if  $\lambda_{j_{L+1}}$  and  $\prod_{i=1}^{L+1} \lambda_{j_i}$  have opposite signs.

In the paper, we first present the simple case where  $L = 1$  (Section 2), and then we turn to the more general one where  $L$  is arbitrary (Section 3). The conclusion (Section 4) will open a few leads about possible extensions.

## 2 A Preliminary Example

We shall suppose that the current state is a real number  $x_t$  linked with the forecast of the next state  $x_{t+1}^e$  and with the predetermined state  $x_{t-1}$  through the following map:

$$\phi x_{t+1}^e + x_t + \pm x_{t-1} = 0 \quad (1)$$

where  $\phi$  (with  $\phi = 0$ , i.e., expectations matter) and  $\pm$  represent the relative weights of future and past respectively. This equation stands for a first order approximation of a

temporary equilibrium dynamics in a small neighborhood of a locally unique stationary state ( $\bar{x} \neq 0$ ). We first focus interest on the relationships between the usual concept of the local determinacy of  $\bar{x}$ , in the perfect foresight dynamics generated by (1) with  $x_{t+1}^e = x_{t+1}$ , namely:

$$\phi x_{t+1} + x_t + \pm x_{t-1} = 0 \quad (2)$$

and its stability under learning. This concept of local determinacy is entirely governed by the perfect foresight roots  $\lambda_1$  and  $\lambda_2$  of the characteristic polynomial  $P(z) = \phi z^2 + z + \pm$  corresponding to (2). We let  $\lambda_1$  and  $\lambda_2$  be real, with  $|\lambda_1| < |\lambda_2|$ . If  $|\lambda_1| > 1$ , then  $\bar{x}$  is source determinate. If  $|\lambda_1| < 1 < |\lambda_2|$ , then it is saddle determinate and for every arbitrarily small neighborhood  $V(\bar{x})$  of  $\bar{x}$ , there is a unique equilibrium ( $x_t$ ) satisfying (2) and staying in  $V(\bar{x})$  at all times, for any initial condition  $x_{t-1}$  close to  $\bar{x}$ . If  $|\lambda_2| < 1$ , then  $\bar{x}$  is locally indeterminate. In that case, for any arbitrarily small neighborhood  $V(\bar{x})$  of  $\bar{x}$ , and any initial condition  $x_{t-1}$  close to  $\bar{x}$ , there are infinitely many perfect foresight equilibria staying in  $V(\bar{x})$  at all times.

One can interpret perfect foresight equilibria as a situation where traders believe that the state variable behavior is governed by:

$$x_t = \bar{\lambda} x_{t-1} \quad (3)$$

for every  $x_{t-1} \in V(\bar{x})$  and every  $t \geq 0$ , and set the growth rate  $\bar{\lambda}$  equal to  $\lambda_1$  or  $\lambda_2$ . Indeed agents are supposed to form their expectations by iterating twice (3) at time  $t$ , i.e.,  $x_{t+1}^e = \bar{\lambda}^2 x_{t-1}$  (note that they do not condition on  $x_t$ ), so that the actual dynamics comes by inserting  $x_{t+1}^e$  into (1):

$$x_t = \lambda_i (\phi \bar{\lambda}^2 + \pm) x_{t-1}. \quad (4)$$

Their belief is self-fulfilling for  $\bar{\lambda} = \lambda_i$  ( $i = 1; 2$ ) since then  $\lambda_i (\phi \bar{\lambda}^2 + \pm)$  equals  $\lambda_i$  by definition. One needs the additional condition  $|\lambda_i| < 1$  to get a locally feasible equilibrium that stays near the stationary state  $\bar{x}$  at all times. Thus the belief  $\bar{\lambda} = \lambda_1$  is the only one to be self-fulfilling and locally feasible when  $\bar{x}$  is saddle determinate. Both beliefs  $\bar{\lambda} = \lambda_1$  and  $\bar{\lambda} = \lambda_2$  are self-fulfilling and locally feasible in the indeterminate case. None of these self-fulfilling beliefs is locally feasible when  $\bar{x}$  is source determinate.

According to the above interpretation, perfect foresight requires that agents coordinate their expectations on some self-fulfilling belief  $\bar{\lambda} = \lambda_i$  ( $i = 1; 2$ ). This is a demanding hypothesis in a decentralized framework where agents may even not know the dynamic laws of their environment, summarized by  $\lambda_1$  and  $\lambda_2$ . It is thus natural to analyze how agents may in fact discover asymptotically  $\lambda_1$  and  $\lambda_2$  through some learning process, where they would formulate their expectations at each date  $t$  from beliefs (3) with  $\bar{\lambda} = \bar{\lambda}(t)$  and would revise them at the beginning of period  $(t+1)$  as a function of the actually observed

forecasting error  $(x_t - \bar{y}(t)x_{t-1})$  in period  $t$ . Here we shall consider the class of econometric learning algorithms whose recursive form is:

$$\bar{y}(t+1) = \bar{y}(t) + \theta(t)h(t)x_{t-1}[x_t - \bar{y}(t)x_{t-1}], \quad (5)$$

$$x_t = \alpha(\bar{y}(t)^2 + \beta)x_{t-1} \quad (6)$$

where  $\theta(t) > 0$  tends toward 0 as  $t$  becomes large, and where  $h(t)$  is some function of past history of the state variable. One must also impose that  $h(t) > 0$  for if agents overestimate the actual growth rate, i.e.,  $x_t < \bar{y}(t)x_{t-1}$ , then they set  $\bar{y}(t+1) < \bar{y}(t)$  in (5). A particular case of this formulation is for instance the ordinary least-squares learning scheme (Marcet and Sargent [14]) where  $h(t) = 1/(x_{t-1}^2 + \epsilon)$ .

If agents set  $\bar{y}(t) = \bar{y}_i$  in (5), then their belief is self-fulfilling, i.e.,  $x_t = \bar{y}_i x_{t-1}$  in (6), and they cease to revise their estimates  $\bar{y}(t)$  in (5), i.e.,  $\bar{y}(t+1) = \bar{y}(t) = \bar{y}_i$ , thus learning the whole trajectory  $x_t = \bar{y}_i x_{t-1}$ . Although the dynamics (5)-(6) seems, at first sight, complex to analyze because of the coupling between growth rates and levels of the state variable, it can be shown (Ljung's [11]) that, provided that  $|\bar{y}_i| < 1$  and  $\theta(t)$  goes to 0 but not too fast, local asymptotic stability of  $\bar{y}(t) = \bar{y}_i$  and of  $x_t = \bar{y}_i x_{t-1}$  in (5)-(6) are equivalent to local stability of  $\bar{y}(t) = \bar{y}_i$  in the simpler associated differential equation:

$$\frac{d\bar{y}(\zeta)}{d\zeta} = \lim_{t \rightarrow \infty} \theta(t) (\alpha(\bar{y}(\zeta)^2 + \beta) - \bar{y}(\zeta)) \quad (7)$$

where  $\zeta$  is a fictitious scale of time related to  $t$ , and  $\lim_{t \rightarrow \infty}$  is assumed to be the limit of some statistics of the process involving  $\theta(t)$ ,  $h(t)$  and  $x_t$ .<sup>1</sup> Provided that  $\lim_{t \rightarrow \infty} \theta(t) > 0$  (a condition that should be thought as a consistency requirement in the learning process), the root  $\bar{y}_i$  will be locally stable for (7) if and only if it is locally stable for the following expectational stability differential equation (Evans [2]):

$$\frac{d\bar{y}(\zeta)}{d\zeta} = \alpha(\bar{y}(\zeta)^2 + \beta) - \bar{y}(\zeta) \quad (8)$$

which makes the revision of growth rate estimates to depend directly on the discrepancy between the actual growth rate  $(\alpha(\bar{y}(\zeta)^2 + \beta))$  and the initial guess  $\bar{y}(\zeta)$ . The next lemma states the properties of (8) in the immediate vicinity of its rest points  $\bar{y}_1$  and  $\bar{y}_2$ .

**Lemma 1** Let  $\alpha = 0$ . The low perfect foresight root  $\bar{y}_1$  is locally stable while the high one  $\bar{y}_2$  is locally unstable in the dynamics with learning (8).

---

<sup>1</sup> The sufficient conditions for this result are  $|\bar{y}_i| < 1$ ,  $\theta(t)$  tends to 0 but the series  $\sum_{t=1}^{\infty} \theta(t)$  diverges to  $+\infty$ , and, for  $\theta(t) = 1/t$ ,

$$\eta = \lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T h_t x_{t-1}^2 \quad \text{as } T \rightarrow \infty.$$

For more general sequences  $(\theta(t))$ , see Ljung [11] (conditions C3, C5, C6)). For the case of weighted least-squares learning schemes where  $\eta = 1$ , see Marcet and Sargent [14].

Proof. Local stability of  $\bar{s}_i$  ( $i = 1; 2$ ) in (8) is obtained if and only if the first derivative of  $(i^{\circ-2} + \pm) i^{-}$  with respect to  $\bar{s}$  is negative for  $\bar{s} = \bar{s}_i$  (see, e.g., Theorem 1.1.1 in Wiggins [19]), namely:

$$\frac{d}{d\bar{s}} (i^{\circ-2} + \pm) i^{-} \Big|_{\bar{s}=\bar{s}_i} = \frac{2\bar{s}_i}{(\bar{s}_1 + \bar{s}_2)} i^{-1} < 0 \quad (9)$$

since  $(\bar{s}_1 + \bar{s}_2)^{-1} i^{-1} = 1$ . If  $\bar{s}_i = \bar{s}_1$ , then (9),  $\bar{s}_1 = \bar{s}_2 < 1$ , which is always satisfied. Otherwise, if  $\bar{s}_i = \bar{s}_2 > 0$ , then (9),  $\bar{s}_2 < \bar{s}_1$ , which never holds true. If  $\bar{s}_i = \bar{s}_2 < 0$ , then (9),  $\bar{s}_2 > \bar{s}_1$ , which is impossible too. ■

Thus the equivalence principle does not hold under the class of learning processes considered here, since agents then learn always the root  $\bar{s}_1$  of lowest modulus, and never discover  $\bar{s}_2$ . Indeed convergence of the trajectories  $(x_t)$  to the stationary state value  $\bar{x}$  imposes  $|j_{s,1}| < 1$ , but this is compatible with both the saddle indeterminate configuration ( $1 < j_{s,2}$ ) and with the indeterminate case ( $|j_{s,2}| < 1$ ). We argue in this paper that this failure is in large part due to the fact that the usual notion of determinacy, as recalled above, is stated in terms of the perfect foresight dynamics of the levels of the state variable  $x_t$ , while in fact traders try to discover some growth rate  $\bar{s} = \bar{s}_i$  ( $i = 1; 2$ ). We introduce now a new perfect foresight dynamics of growth rates, which is obtained by assuming that the traders' belief about the law of motion of the state variable fits (3) with  $\bar{s} = \bar{s}(t)$ :

$$x_t = \bar{s}(t) x_{t-1} \quad (10)$$

for every  $x_{t-1} \in V(\bar{x})$  and every  $t \geq 0$ . Perfect foresight of the state variable  $x$  induces a dynamics of the growth rate  $\bar{s}(t)$  whose fixed points are the roots  $\bar{s}_1$  and  $\bar{s}_2$ . The issue is to study whether there exist links between stability of these roots under learning and their determinacy properties in such a growth rate perfect foresight dynamics. It turns out that, in fact, a perfect foresight growth rate is locally stable under learning if and only if it is locally determinate in the growth rate perfect foresight dynamics.

Let us recall that (10) holds for every  $t \geq 0$ , so that traders expectations are:

$$x_t^e = \bar{s}(t) x_{t-1} \quad \text{and} \quad x_{t+1}^e = \bar{s}(t+1) x_t^e = \bar{s}(t+1) \bar{s}(t) x_{t-1}.$$

In the perfect foresight dynamics (2), these forecasts are equal to  $x_t$  and  $x_{t+1}$ , respectively. One can consequently rewrite (2) as a recursive equation of growth rates only.

**Definition 2** Let  $\phi = 0$ . Assume also that (10) holds for every  $x_{t-1} \in V(\bar{x})$  and every  $t \geq 0$ . The growth rate perfect foresight dynamics is a sequence of growth rates  $(\bar{s}(t))$  such that:

$$\phi \bar{s}(t+1) \bar{s}(t) + \bar{s}(t) + \pm = 0. \quad (11)$$

Taking  $\bar{r}(t) = \bar{r}(t+1)$  in (11) shows that the fixed points of (11) are  $r_1$  and  $r_2$ . This dynamics is well defined if and only if  $\bar{r}(t) = 0$  in each period. Therefore it must be the case that  $\bar{r}(t) = i \pm$  if  $\pm = 0$  (otherwise  $\bar{r}(t+1) = 0$ ), in which case (11) does not define a global dynamics but is yet well defined around  $r_1$  and  $r_2$  (both differ from  $i \pm$ ). If  $\pm = 0$ , then  $r_1 = 0$  (since then  $r_1, r_2 \neq i \pm = 0$  and  $j_{1j} < j_{2j}$ ) so that the dynamics is not well defined locally, but one may say that  $r_1$  is unstable while  $r_2$  is stable because either  $\bar{r}(t) = r_1$  at all times, or  $\bar{r}(t) = i 1 = r_2 = 0$  if  $\bar{r}(t) = 0$  at some date.

The dynamics (11) has the classical one-step forward looking structure without pre-determined variables (the current rate  $\bar{r}(t)$  is not given at outset of  $t$ ). So its fixed points are locally determinate if and only if they are locally unstable in (11), which allows us to state our equivalence result.

**Proposition 3** Let  $\phi = 0$ . The low perfect foresight root  $r_1$  is locally determinate in the growth rate perfect foresight dynamics (11) while the high one  $r_2$  is locally indeterminate in the same dynamics. Therefore, in view of Lemma 1, the root  $r_i$  ( $i = 1; 2$ ) is locally stable in the learning dynamics (8) if and only if it is locally determinate in the growth rate perfect foresight dynamics (11).

**Proof.** The growth rate perfect foresight dynamics in the neighborhood of  $r_i$  ( $i = 1; 2$ ) is obtained by linearizing (11) at this point:

$$\phi_{r,i}(\bar{r}(t+1) - r_i) + (\phi_{r,i} + 1)(\bar{r}(t) - r_i) = 0$$

The condition for  $r_i$  to be locally unstable in (11) is:

$$\frac{\phi_{r,i} + 1}{\phi_{r,i}} = \frac{\phi_{r,i} (r_1 + r_2)}{r_i} > 1 \quad (12)$$

If  $r_i = r_1$ , then (12),  $j_{2j} > j_{1j} > 1$ , which always holds. If  $r_i = r_2$ , then (12),  $j_{1j} > j_{2j} > 1$ , which never holds. ■

### 3 On the Dynamic Equivalence Principle

We now deal with economies where the current state depends on the forecast of the next state but also on an arbitrary number  $L$  of predetermined variables through the following map:

$$\phi x_{t+1}^e + x_t + \sum_{l=1}^L \pm_l x_{t-l} = 0 \quad (13)$$

where the parameter  $\pm_l$  ( $1 \leq l \leq L$ ) represents the relative contribution to  $x_t$  of the predetermined state  $x_{t-l}$  at  $t$ . We shall proceed here as in the previous section, namely we shall first state the lack of link between determinacy of the stationary state ( $\dot{x} = 0$ )



in the perfect foresight dynamics on the levels of the state variable  $x$ , and its stability under learning when agents try to estimate how the state of period  $t$  is related to the  $L$  past states  $x_{t-1}; \dots; x_{t-L}$ , i.e., try to discover the  $L$  coefficients of the linear perfect foresight dynamics restricted to a  $L$ -dimensional eigenspace. Then we shall define the perfect foresight dynamics on the  $L$ -dimensional vectors of these coefficients. Thus the issue will be to study whether there is a neater relationship between determinacy and stability under learning with this new perfect foresight dynamics.

### 3.1 State Variable Perfect Foresight Dynamics

The usual concept of local determinacy of the stationary state  $\bar{x}$  is defined from the perfect foresight dynamics on the levels of the state variable  $x$  obtained by setting  $x_{t+1}^e = x_{t+1}$  in (13), namely:

$$\phi x_{t+1} + x_t + \sum_{i=1}^P \alpha_i x_{t-i} = 0 \quad (14)$$

Indeed this concept only relies on the  $(L+1)$  perfect foresight roots  $\lambda_1; \dots; \lambda_{L+1}$  of the characteristic polynomial  $P(z) = \phi z^{L+1} + z^L + \sum_{i=1}^P \alpha_i z^{L-i}$  corresponding to (14). Let  $\lambda_i$  ( $i = 1; \dots; L+1$ ) be real, with  $\lambda_1 < \lambda_2 < \dots < \lambda_{L+1}$ . If  $\lambda_L > 1$ , then  $\bar{x}$  is source determinate. If  $\lambda_L < 1 < \lambda_{L+1}$ , then  $\bar{x}$  is saddle determinate and for every arbitrarily small neighborhood  $V(\bar{x})$  of  $\bar{x}$ , and for every initial condition  $(x_{t-1}; \dots; x_{t-L})$  close to  $(\bar{x}; \dots; \bar{x})$ , there is a unique perfect foresight equilibrium, i.e., a unique sequence  $(x_t)$  that satisfies (14) and stays in  $V(\bar{x})$  at all times  $t \geq 0$ . If  $\lambda_{L+1} < 1$ , then  $\bar{x}$  is indeterminate, i.e., for every arbitrarily small neighborhood  $V(\bar{x})$  of  $\bar{x}$ , and for every initial condition  $(x_{t-1}; \dots; x_{t-L})$  close to  $(\bar{x}; \dots; \bar{x})$ , there are infinitely many perfect foresight equilibria  $(x_t)$  that stay in  $V(\bar{x})$  at all dates  $t \geq 0$ .

Here again one can interpret perfect foresight as a situation where traders believe that the law of motion of the system is governed by:

$$x_t = \sum_{i=1}^P \alpha_i x_{t-i} \quad (15)$$

for every  $x_{t-i} \in V(\bar{x})$  and every  $t \geq 0$ , and where this belief is self-fulfilling. In that case, the expectation are formed at date  $t$  by iterating twice (15):

$$x_{t+1}^e = \sum_{i=1}^P \alpha_i x_{t+1-i} = \sum_{i=1}^{L+1} \left( \alpha_{i-1} + \alpha_{i+1} \right) x_{t-i} + \alpha_L x_{t-L}.$$

This forecast, once reintroduced into (13), generates the actual dynamics:

$$x_t = \sum_{i=1}^{L+1} \alpha_i \left( \alpha_{i-1} + \alpha_{i+1} \right) x_{t-i} + \alpha_L x_{t-L}. \quad (16)$$

Hence the initial belief (15) is self-fulfilling if and only if it coincides with (16), namely:

$$\alpha_i = \sum_{j=1}^{L+1} \alpha_j \left( \alpha_{j-1} + \alpha_{j+1} \right) + \alpha_L \quad \text{for } i = 1; \dots; L \quad (17)$$

with  $\bar{r}_{L+1} = 0$ . Vectors solutions to (17) will be called stationary extended growth rates (henceforth EGR(L)), and denoted  $\bar{r} = (\bar{r}_1; \dots; \bar{r}_L)$ . Intuitively the self-fulfilling belief (15) with  $\bar{r} = (\bar{r}_1; \dots; \bar{r}_L) = \bar{r}^k$  should correspond to the perfect foresight dynamics (14) restricted to an invariant subspace  $W$  of dimension  $L$ , i.e., to an eigenspace spanned by  $L$  eigenvectors among the  $(L + 1)$  eigenvectors  $u_i$  associated with the eigenvalues  $\lambda_i$  ( $i = 1; \dots; L + 1$ ). There are clearly  $(L + 1)$  such invariant  $L$ -dimensional eigenspaces  $W_k$  (each  $W_k$  is spanned by all the eigenvectors but  $u_k$  ( $k = 1; \dots; L + 1$ )). Thus one should expect (17) to have  $(L + 1)$  distinct vector solutions  $\bar{r}^k$ , where  $\bar{r}^k$  corresponds to the perfect foresight dynamics restricted to  $W_k$ . The next lemma makes precise this intuition and gives the expression of  $\bar{r}^k$  in terms of the perfect foresight roots  $\lambda_i$  ( $i = 1; \dots; L + 1$ ).

**Lemma 4** Assume that the characteristic polynomial  $P$  corresponding to the  $(L + 1)$ -dimensional difference equation (14) admits  $(L + 1)$  real and distinct roots  $\lambda_k$ , with  $\lambda_{j-1} < \lambda_j < \lambda_{j+1}$ . Let the  $(L + 1) \times 1$  eigenvector  $u_k$ ,  $1 \leq k \leq L + 1$ , be associated with  $\lambda_k$ . Finally let  $W_k \subset \mathbb{R}^L$ ,  $1 \leq k \leq L + 1$ , be the eigensubspace spanned by all the eigenvectors except  $u_k$ . The perfect foresight dynamics of the state variable restricted to  $W_k$  writes:

$$x_t = \sum_{i=1}^L \bar{r}_i^k x_{t-1}^i$$

where the  $i$ th entry  $\bar{r}_i^k$  of the stationary EGR(L)  $\bar{r}^k$  is:

$$\bar{r}_i^k = (\lambda_i - 1)^{L+1} \prod_{1 \leq i_1 < \dots < i_L \leq L+1} (\lambda_{i_1} \dots \lambda_{i_L} \lambda_i) \quad \text{for all } i_z = k, z = 1; \dots; L$$

The  $(L + 1)$  stationary EGR(L)  $\bar{r}^k$  are the solutions of the equations (17).

**Proof.** See in Appendix 5.1. ■

Of course, for the self-fulfilling belief (15) with  $\bar{r}^k$  to be locally feasible, one must also require that the perfect foresight dynamics of the state variable  $x$  in  $W_k$  be stable (all the perfect foresight roots different from  $\lambda_k$  must be stable, i.e.,  $\lambda_{i-1} < 1$  for  $i = k$ ). Therefore the self-fulfilling belief  $\bar{r} = \bar{r}^k$  with  $k = L + 1$  is the only one to be locally feasible in the saddle point configuration for the dynamics (14). Any self-fulfilling belief  $\bar{r}^k$  with  $k = 1; \dots; L + 1$  is locally feasible if  $\lambda$  is indeterminate in this dynamics, and none of these self-fulfilling beliefs is locally feasible when  $\lambda$  is source determinate.

## 3.2 Dynamics with Learning

Even if agents are initially aware of the  $(L + 1)$  stationary EGR(L) to the model (13), they have to coordinate their behavior on one among them, which may be quite demanding in a decentralized framework. In this section we shall instead assume that agents need learning how the state variable behaves in  $V(\lambda)$  through a process where they form at

the outset of period  $t$  their forecasts from the law (15) with some  $L$ -dimensional vector of estimates  $\bar{w}(t) = (\bar{w}_1(t); \dots; \bar{w}_L(t))^T$  and revise these estimates at the outset of period  $(t + 1)$  once the forecasting error  $(x_{t+1} - \bar{w}(t)^T x_{t+1})$  is actually observed (here  $x_{t+1}$  stands for the vector of lagged variables  $(x_{t+1-1}; \dots; x_{t+1-L})^T$ ), according to the class of recursive learning schemes:

$$\bar{w}(t + 1) = \bar{w}(t) + \alpha(t) H_t x_{t+1} (x_{t+1} - \bar{w}(t)^T x_{t+1}), \quad (18)$$

$$x_{t+1} = \sum_{i=1}^p h_i (\bar{w}_1(t) \bar{w}_{i+1}(t) + \bar{w}_{i+1}(t)) + \sum_{i=1}^L x_{t+1-i} - (\bar{w}(t))^T x_{t+1} \quad (19)$$

As in (5), the sequence of scalars  $\alpha(t) > 0$  in (18) is still assumed to tend toward 0 as time passes, and  $H_t$  is now a  $L \times L$  matrix related to past history of the economic system. The rule (18) is general enough to encompass, e.g., the weighted least-squares schemes where  $H_t = (x_{t+1-1} x_{t+1-1}^T + \dots + x_{t+1-L} x_{t+1-L}^T)^{-1}$ . Equation (19) is (16) with  $\bar{w}_i$  is replaced by  $\bar{w}_i(t)$  (for  $i = 1; \dots; L$ ), and with the convention that  $\bar{w}_{L+1}(t) = 0$  (hence the  $i$ th component of the  $L \times 1$  vector  $-(\bar{w}(t))$  is the actual weight of  $x_{t+1-i}$  in (19)).

If agents set  $\bar{w}(t) = w^k$  in (18) for some  $k$  ( $k = 1; \dots; L + 1$ ) at date  $t$ , then  $x_t = (w^k)^T x_{t+1}$  in (19). Therefore they do not revise their estimates, i.e., they set  $\bar{w}(t + 1) = \bar{w}(t) = w^k$  in (18), thus learning the law of motion of the state variable restricted to  $W_k$ . The dynamics (18)-(19) is, however, complex to analyze because of the coupling between components of  $\bar{w}(t)$  and levels of the state variable. Here again, as in Section 2, we may appeal to the existing theory on the convergence of learning algorithms (Ljung's [11]) to assert that local asymptotic stability in (18)-(19) of some particular  $w^k$  corresponding to a stable (feasible) dynamics in  $W_k$  is equivalent, provided that  $\alpha(t)$  goes to 0 but not too fast, to its local stability in the associated ordinary differential equation (see footnote 1 for details):

$$\dot{\bar{w}}(t) = \hat{\Phi}(-(\bar{w}(t))) \bar{w}(t) \quad (20)$$

where  $t$  is a fictitious continuous scale of time, and where the  $L \times L$  matrix  $\hat{\Phi}$  is assumed to be the limit of some statistics involving  $\alpha(t)$ ,  $H_t$  and the state variable  $x$ . We need here additional assumptions to be able to reduce the study of the stability in (20) to the differential expectational stability criterion (Evans [2]):

$$\dot{\bar{w}}(t) = -(\bar{w}(t)) \bar{w}(t). \quad (21)$$

An intuition for local equivalence between (20) and (21) is easy to grasp by considering Jacobian matrices  $\hat{\Phi}(D - (w^k)) \bar{w}(t)$  and  $(D - (w^k)) \bar{w}(t)$  that govern the dynamics near  $w^k$  in (20) and (21), respectively. One can show that  $(D - (w^k)) \bar{w}(t)$  has distinct real eigenvalues under the assumption of distinct real eigenvalues  $\lambda_i$  ( $i = 1; \dots; L + 1$ ) (see Appendix 5.2). It is therefore diagonalizable. In the corresponding basis, one should expect the asymptotic learning dynamics (20) to be consistent with that structure and to

involve correcting each “diagonal error” along the direction of the corresponding eigenvector of  $(D - (\lambda^k)_i I_L)$  with a positive weight. In other words, both  $\hat{C}$  and  $(D - (\lambda^k)_i I_L)$  should be diagonalizable in the same basis, and the diagonalized matrix  $\hat{C}$  should have only positive entries on its diagonal. Local stability of (20) and (21) are clearly equivalent in such circumstances, which are in principle more general than usual least-squares schemes where  $\hat{C}$  is in fact the identity matrix (see Marcet and Sargent, Proposition 3). Of course the dynamics (21) will be locally stable around  $\lambda^k$  if and only if the real part of all the eigenvalues of  $(D - (\lambda^k)_i I_L)$  is negative (see, e.g., Theorem 1.1.1 in Wiggins [19]).

**Theorem 5** Let  $\sigma = 0$ . Assume that beliefs  $\lambda_t$  (15) for every  $x_{t+1}$  in  $V(\lambda)$  and every  $t \geq 0$ . Then there exists a unique stationary  $EGR(L)^{\lambda^k}$  which is locally stable in the learning dynamics (21). It governs the behavior of the state variable restricted to the  $L$ -dimensional eigensubspace  $W_k$  of the state variable perfect foresight dynamics (14) spanned by all the eigenvectors except  $u_k$  where  $u_k$  is associated with the perfect foresight root  $\lambda_k$  which satisfies:

$$\lambda_k = \max_{i=1, \dots, L} \left( \frac{\lambda_i^{L+1}}{\lambda_i} \right)$$

Therefore the stationary  $EGR(L)^{\lambda^{L+1}}$  that governs the perfect foresight dynamics (14) restricted to the  $L$ -dimensional eigenspace associated with the  $L$  eigenvalues  $\lambda_1, \dots, \lambda_L$  of lowest modulus, is the unique stationary  $EGR(L)$  to be locally stable in the learning dynamics (21) if  $\frac{\lambda_{L+1}^{L+1}}{\lambda_{L+1}}$  is positive. Otherwise, i.e., if  $\frac{\lambda_{L+1}^{L+1}}{\lambda_{L+1}}$  is negative, then the stationary  $EGR(L)$  that is locally stable in the learning dynamics (21) governs the perfect foresight dynamics (14) restricted to an  $L$ -dimensional eigenspace that contains in particular the eigenvector  $u_{L+1}$  associated to the eigenvalue of largest modulus  $\lambda_{L+1}$ .

**Proof.** See in Appendix 5.2. ■

This result implies again that there are no simple links between the determinacy of  $\lambda$  in the perfect foresight dynamics (14) and its stability under learning. In the case where  $\lambda_{L+1}$  and  $\frac{\lambda_{L+1}^{L+1}}{\lambda_{L+1}}$  have the same sign (the reader will note that this condition is always met in the case of a single predetermined variable  $L = 1$  considered in the previous section), convergence of the state variable  $x_t$  to  $\lambda$  in the learning dynamics requires also that  $j_{L+1} < 1$ , but this condition is compatible with both the saddle determinate configuration  $1 < j_{L+1}$  and the indeterminate case  $j_{L+1} < 1$ . On the contrary, when  $\lambda_{L+1}$  and  $\frac{\lambda_{L+1}^{L+1}}{\lambda_{L+1}}$  have not the same sign (which may occur only if  $L \geq 2$ ), convergence of  $x_t$  to  $\lambda$  in the learning dynamics requires  $j_{L+1} < 1$ , which corresponds to the indeterminate case.

### 3.3 Extended Growth Rate Perfect Foresight Dynamics

We argue here again that the fact that there are no simple links between the determinacy of the stationary state  $\lambda$  and its stability under learning, may be due to the fact that

determinacy was applied to the perfect foresight dynamics (14) on the levels of the state variable  $x_t$ , while agents try to learn extended growth rates. We may thus expect that simpler relationships might arise if one considers instead determinacy in a perfect foresight dynamics on extended growth rates. Such a dynamics is constructed in assuming that the traders' beliefs .t:

$$x_t = \prod_{l=1}^L \bar{\omega}_l(t) x_{t_{l-1}} \quad (22)$$

for every  $x_{t_{l-1}}$  ( $l = 1; \dots; L$ ) in  $V(\bar{x})$  and every  $t \geq 0$ . The dynamics with perfect foresight (14) of the state variable induces a  $L$ -dimensional extended growth rate perfect foresight dynamics on the extended growth rates  $\bar{\omega}(t) = (\bar{\omega}_1(t); \dots; \bar{\omega}_L(t))$  whose fixed points are the stationary EGR( $L$ ). Applying the determinacy criterion to these vectors, in this extended growth rates dynamics, allows us to show by that the stationary EGR( $L$ ) that governs the perfect foresight dynamics restricted to the  $L$ -dimensional eigenspace associated with the  $L$  eigenvalues  $\lambda_1; \dots; \lambda_L$  of lowest modulus, is actually the only one that is locally determinate. In view of theorem 5, this fact implies that the equivalence between determinacy and stability under learning will be restored, at least when  $\lambda_{L+1}$  and  $\prod_{i=1}^{L+1} \lambda_i$  have the same sign.

Since (22) holds for each  $t$ , the corresponding expectations are as follows:

$$x_t^e = \prod_{l=1}^L \bar{\omega}_l(t) x_{t_{l-1}} \quad \text{and} \quad x_{t+1}^e = \bar{\omega}_1(t+1) x_t^e + \prod_{l=2}^L \bar{\omega}_l(t+1) x_{t+1_{l-1}}.$$

Under the perfect foresight hypothesis  $x_t^e$  and  $x_{t+1}^e$  are equal to  $x_t$  and  $x_{t+1}$  respectively so that (14) becomes:

$$(1 + \bar{\omega}_1(t+1))x_t = \prod_{l=1}^{L-1} (\bar{\omega}_{l+1}(t+1) + \pm_l) x_{t_{l-1}} \pm_L x_{t_L}. \quad (23)$$

Since (22) and (23) holds for every  $x_{t_{l-1}}$ , the EGR( $L$ ) perfect foresight dynamics is obtained whenever the coefficients of (22) and of (23) coincide.

**Definition 6** Let  $\omega = 0$ . Assume also that (22) holds for every  $x_{t_{l-1}}$  near  $\bar{x}$  ( $l = 1; \dots; L$ ) and every  $t \geq 0$ . The EGR( $L$ ) perfect foresight dynamics is a sequence of  $L$ -dimensional vectors  $\bar{\omega}(t)$  such that:

$$\bar{\omega}_l(t) = \prod_{i=1}^L \frac{\bar{\omega}_{l+1}(t+1) + \pm_i}{1 + \bar{\omega}_1(t+1)} \quad (24)$$

for each  $l = 1; \dots; L$ , and with the convention that  $\bar{\omega}_{L+1}(t+1) = 0$ .

This dynamics is well defined if and only if  $\bar{\omega}_L(t) = 0$  in each period. For in that case (24) for  $l = L$  determines  $\bar{\omega}_1(t+1)$  while the other equations determine  $\bar{\omega}_l(t+1)$  for  $l = 1; \dots; L-1$ . In the case  $L \geq 2$  (the case  $L = 1$  was dealt with Section 2),  $\bar{\omega}_L(t) = 0$  implies  $\bar{\omega}_L(t+1) = 0$  if and only if  $\bar{\omega}_L(t+1) = (\bar{\omega}_{L-1}(t) + \pm_L) \pm_{L-1} \pm_{L-1} = 0$ . So, when  $\pm_L = 0$ , global dynamics are not well defined but local dynamics near every stationary

EGR(L) are (see appendix 5.3). If  $\lambda_L = 0$ , then  $\lambda_1 = 0$  (since  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_{L+1} = \lambda_1 + \lambda_L = 0$  and  $j_{\lambda_1} < \lambda_2 < \dots < j_{\lambda_{L+1}}$ ), which implies (by using Lemma 4) that  $\lambda_1^+ = 0$  but  $\lambda_L^+ = 0$  for  $k = 1$  (i.e.,  $k = 2; \dots; L + 1$ ), so that local dynamics are not well defined around  $\lambda^k$  (for  $k = 1$ ) but one may say that  $\lambda^k$  is unstable for  $k = 1$  while  $\lambda^1$  is stable in the sense that either  $\lambda^-(t) = \lambda^k$  ( $k = 1$ ) at all times, or  $\lambda^-(t) = \lambda^1$  if  $\lambda^-(t) = 0$  at some date.

Since (24) displays a (multidimensional) one-step forward looking structure without predetermined variables, a stationary EGR(L) is locally determinate if and only if all the eigenvalues which govern (24) close to it have modulus greater than 1.

**Proposition 7** The stationary EGR(L)  $\lambda^{L+1}$  that governs the perfect foresight dynamics (14) restricted to the L-dimensional eigenspace associated to the L eigenvalues  $\lambda_1; \dots; \lambda_L$  of lowest modulus, is the unique stationary EGR(L) to be is locally determinate in the EGR(L) perfect foresight dynamics (24). Therefore, in view of Theorem 5, the dynamic equivalence principle holds true if  $\lambda_L$  has the same sign as  $\prod_{i=1}^{L+1} \lambda_i$ , and not otherwise.

Proof. See in Appendix 5.4. ■

## 4 Conclusion

The main innovation of the paper is to describe an extended growth rate perfect foresight dynamics whose fixed points are the L-dimensional vectors of coefficients which govern the local perfect foresight dynamics restricted to some L-dimensional eigensubspace. Such a dynamics allows us to save the equivalence between determinacy and stability under recursive learning under a simple one sided (sign) condition that may be related to the condition for stability of differential equations. It is clear, however, that this result depends on specific features of the model.

(i) Agents were assumed to condition forecasts at  $t$  only on past data up to date  $(t - 1)$ . An interesting topic for further research would be to investigate what would happen if agents may condition also on the current equilibrium state  $x_t$ .

(ii) Finally, agents were assumed to care about deviations from the stationary state value  $\bar{x}$  so that, in fact, they were a priori aware of this value. One may go beyond such an assumption by introducing an estimate  $\hat{A}$  of the value  $\bar{x}$  into the perceived laws of motion (15). Namely (15) rewrites:

$$x_t = \prod_{i=1}^P x_{t_i} + \hat{A}$$

One can prove that, in this case, saddle determinacy of the stationary state ( $\bar{x}$ ) is equivalent to its stability under learning if  $\lambda_{L+1} = \prod_{i=1}^{L+1} \lambda_i > 0$  and  $\prod_{i=1}^P \lambda_i = \lambda_1 = 0 > 0$ , i.e., if

$\lambda_{L+1} > 0$ . Actually learning a stationary extended growth rate does not rely on the fact that agents already know  $\bar{x}$ . Therefore the former condition ensures that agents discover the law of motion of the state variable restricted to the subspace corresponding to the  $L$  roots of lowest modulus  $\lambda_1; \dots; \lambda_L$ . The remaining one ( $\lambda < 0$ ) ensures that, in the model where  $L = 0$ , agents discover the value  $\bar{x}$  ( $\lambda > 1$ ) if and only if the stationary state ( $\bar{x}$ ) is locally determinate ( $|\lambda| < 1$ ).

## 5 Proofs

### 5.1 Proof of Lemma 4

We first transform the dynamics (14) into a vector first order difference equation:

$$x_{t+1} = T x_t$$

where:

$$x_t = \begin{pmatrix} x_t^0 \\ \vdots \\ x_t^L \end{pmatrix} \quad \text{and} \quad T = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & \lambda_1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_L \end{pmatrix}$$

The proof proceeds from the fact that  $x_t$  belongs to a  $L$ -dimensional eigensubspace  $W_k$  ( $k = 1; \dots; L + 1$ ) of (14) if and only if it is a linear combination of the  $k$  eigenvectors that span  $W_k$ . Of course the  $(L + 1)$  eigenvalues of the  $(L + 1) \times (L + 1)$  matrix  $T$  are the perfect foresight roots  $\lambda_i$  ( $i = 1; \dots; L + 1$ ) of the characteristic polynomial  $P$ . A convenient form for the  $(L + 1) \times 1$  eigenvector  $u_i$  ( $i = 1; \dots; L + 1$ ) associated with  $\lambda_i$  is obtained by using the relations between the coefficients and the roots of  $P$ . Namely:

$$P(\lambda) = 0, \quad \sum_{i=1}^{L+1} \lambda_i^L = 0, \quad \sum_{i=1}^{L+1} \lambda_i^{L-1} = 0, \quad \dots, \quad \sum_{i=1}^{L+1} \lambda_i = 0$$

Let us identify the coefficients in the expression above and those of  $P$ . One gets:

$$(\lambda_i - 1)^{L+1} \sum_{l=1}^{L+1} \lambda_i^l = 0 \quad \text{for } i = 0; \dots; L + 1, \text{ with } \lambda_0 = 1 \quad (25)$$

where  $\lambda$  is the set of all the perfect foresight roots, and  $\sum_{l=1}^{L+1} \lambda^l$  is the  $l$ th ( $l = 1; \dots; L + 1$ ) elementary symmetric polynomial of  $\lambda$ , i.e., the sum on all the possible products over  $l$  different elements of  $\lambda$ :

$$\sum_{l=1}^{L+1} \lambda^l \stackrel{\text{def}}{=} \sum_{1 \leq i_1 < \dots < i_l \leq L+1} (\lambda_{i_1} \dots \lambda_{i_l})$$

By definition  $u_i$  is such that  $T u_i = \lambda_i u_i$  ( $i = 1; \dots; L + 1$ ). It can be shown that:

$$u_i = (\lambda_i^L; \lambda_i^{L-1}; \dots; 1)^0$$

where the symbol  $^0$  denotes the (vector) transpose. The perfect foresight trajectory that is restricted to  $W_k$  is such that  $x_t$  is a linear combination of all the  $u_i$  but  $u_k$ . This condition writes  $\det(x_t; U_k) = 0$  where  $U_k$  is the  $(L + 1) \times L$  matrix whose columns are all the  $u_i$  but  $u_k$ . Developing  $\det(x_t; U_k)$  with respect to its first column  $x_t$  leads to:

$$x_t = \sum_{i=1}^L a_i x_{t_i} = \sum_{i=1}^L ((i-1)^{L+1} \phi_{i \neq k}) x_{t_i}$$

where  $\phi_0$  is the determinant of the  $(L+1) \times (L+1)$  Vandermonde matrix and  $\phi_i = \frac{1}{i} (\$)_k \phi_0$  (see respectively Ch. 10 and Ex. 10.12 in Ramis, Deschamps and Odoux [15]) and  $\$ _k$  is the set of all the perfect foresight roots but  $_{s,k}$ . Thus  $a_i$  is as stated in Lemma 1, namely:

$$a_i^k = (i-1)^{L+1} \frac{1}{i} (\$)_k \quad (26)$$

It remains consequently to prove that vectors  $(a_1^k; \dots; a_L^k)$  are the solutions to (17), i.e.,  $a_i^k = \frac{1}{i} (\$)_k$  ( $i = 1; \dots; L$  and  $k = 1; \dots; L+1$ ). We show this directly by replacing  $\bar{1}$  in (17) by the coefficient  $a_i^k$  given (26) for any  $k$  given. Observe first that, for  $\phi = 0$ , the  $L$ th equation of (17) rewrites:

$$(\phi + \bar{1}) \bar{1} = i \bar{1}_{i+1} i \pm \phi \quad (27)$$

Since  $a_i^k = \frac{1}{i} (\$)_k$  and  $(i-1) = \frac{1}{i} (\$)$ , we have  $(\phi + \bar{1}) = \frac{1}{i} (\$)_k$ . Hence the  $L$ th equation of (17) becomes (recall that  $\bar{1}_{L+1} = 0$ ):

$$\bar{1}_L = \frac{\pm \phi}{\$ _k} = \frac{1}{\$ _k} (i-1)^{L+1} \frac{1}{i} (\$) \quad (28)$$

where we used (25) with  $i = L$ . By definition,  $\frac{1}{i} (\$)$  is the product over all the perfect foresight roots. Therefore (28) rewrites:

$$\bar{1}_L = (i-1)^{L+1} \prod_{i=k}^{L+1} \frac{1}{i} \stackrel{\text{def}}{=} (i-1)^{L+1} \frac{1}{i} (\$)_k$$

As (26) shows,  $\bar{1}_L$  is equal  $a_L^k$ , and so  $a_L^k = \frac{1}{L} (\$)_k$ . Assume now that  $\bar{1}_{i+1} = a_{i+1}^k$  ( $i < L$ ). We are going to prove that  $\bar{1}_i = a_i^k$  ( $i < L$ ). Indeed, by using (25) and (26) in (27) for some  $i$ , one obtains:

$$\bar{1}_i = \frac{1}{\$ _k} ((i-1)^{L+2} \frac{1}{i} (\$)_k + (i-1)^{L+1} \frac{1}{i} (\$)) = \frac{1}{\$ _k} (i-1)^{L+1} (i \frac{1}{i} (\$)_k + \frac{1}{i} (\$))$$

But, by definition,  $\frac{1}{i} (\$)_k$  is the sum on all the products over  $(i+1)$  different elements of  $\$ _k$  while  $\frac{1}{i} (\$)$  is the sum on all the products over  $(i+1)$  different elements of  $\$$ . The difference between these sums is just the sum on all the products over  $(i+1)$  different perfect foresight roots, provided that each of the remaining products includes the root  $_{s,k}$ . Hence:

$$i \frac{1}{i} (\$)_k + \frac{1}{i} (\$) = \frac{1}{\$ _k} \frac{1}{i} (\$)_k \quad \bar{1}_i = (i-1)^{L+1} \frac{1}{i} (\$)_k = a_i^k$$

Since  $\bar{1}_i = a_i^k$ , we have  $\frac{1}{i} (\$)_k = a_i^k$  for any  $i$  ( $i = 1; \dots; L$ ) and for any given  $k$ , which completes the proof of Lemma 1.



## 5.2 Proof of Theorem 5

The dynamics (21) in the neighborhood of  $^{+k}$  ( $k = 1; \dots; L + 1$ ) is governed by the  $L$  eigenvalues  $^{1j}_k$  ( $j = 1; \dots; L$ ) of the  $L \times L$  Jacobian matrix  $^a_k$  of (21) calculated at  $^{+k}$ :

$$^a_k = \begin{pmatrix} 0 & i^{2^{+k}}_1 & 1 & i^\circ & 0 & \phi\phi\phi & 0 \\ i^{\circ+2k}_2 & i^{\circ+2k}_1 & 1 & i^\circ & \phi\phi\phi & 0 \\ i^{\circ+2k}_3 & 0 & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & i^\circ \\ i^{\circ+2k}_L & 0 & \phi\phi\phi & 0 & i^{\circ+2k}_1 & 1 \end{pmatrix}$$

These eigenvalues are the  $L$  solutions of the equation  $\det(^a_k i^{-1} I_L) = 0$ , where  $I_L$  denotes the  $L \times L$  identity matrix. Let develop  $\det(^a_k i^{-1} I_L)$  with respect to its first column:

$$\begin{aligned} & (i-1)^2 (i^{2^{+k}}_1 - 1 i^{-1}) \begin{vmatrix} i^{\circ+2k}_1 & 1 i^{-1} & i^\circ & \phi\phi\phi & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & & \ddots & i^\circ \\ 0 & \phi\phi\phi & \phi\phi\phi & i^{\circ+2k}_1 & 1 i^{-1} \end{vmatrix} \\ & + (i-1)^3 (i^{\circ+2k}_2) \begin{vmatrix} i^\circ & 0 & \phi\phi\phi & 0 \\ 0 & i^{\circ+2k}_1 & 1 i^{-1} & i^\circ \phi\phi\phi & 0 \\ \vdots & & \ddots & i^\circ \\ 0 & \phi\phi\phi & \phi\phi\phi & i^{\circ+2k}_1 & 1 i^{-1} \end{vmatrix} + \phi\phi\phi \\ & \phi\phi\phi + (i-1)^{L+1} (i^{\circ+2k}_L) \begin{vmatrix} i^{\circ+2k}_1 & 1 i^{-1} & i^\circ & 0 & \phi\phi\phi & 0 \\ \vdots & & \ddots & 0 \\ 0 & \phi\phi\phi & \phi\phi\phi & i^\circ \end{vmatrix} \end{aligned}$$

Therefore:

$$\det(^a_k i^{-1} I_L) = 0$$

$$, \quad Q_k(z) = (1 + i^{\circ+2k}_1 + z)^L i^{\frac{P}{L+1}} (i-1)^{L+1} i^{\frac{1}{L+1}} (1 + i^{\circ+2k}_1 + z)^{L+1} = 0 \quad (29)$$

The expression of  $^{+k}_i$  given in (26) lead to:

$$(i-1)^{L+1} i^{\frac{1}{L+1}} = (i^\circ)^L (i-1)^{L+1} i^{\frac{1}{L+1}} (\$_k) = (i-1)^{L+1} i^{\frac{1}{L+1}} (f_i^\circ \$_k g)$$

where the  $L$  elements of the set  $f_i^\circ \$_k g$  are  $(i^\circ, _j)$  for  $j = 1; \dots; L + 1$  and  $j \neq k$ . We reintroduce now the expression of  $(i-1)^\circ$  given in (25):

$$(i-1)^{L+1} i^{\frac{1}{L+1}} = (i-1)^{L+1} i^{\frac{1}{L+1}} \frac{\tilde{A}(\frac{1}{L+1})}{P_{L+1, i}} \$_k \quad (30)$$

Let now  $y = 1 + \alpha_1^{+k} + z$  so that (29) rewrites (with (30)):

$$Q_k(y) = y^L \prod_{i=1}^L (i-1)^{L+1} \frac{1}{\alpha_i^{+k}} \prod_{i=1}^{L+1} \frac{1}{\alpha_i^{+k}} \alpha_k = 0$$

This shows that the  $L$  roots of  $Q_k$  are  $\alpha_j = \frac{1}{\alpha_i^{+k}}$  for  $j = 1; \dots; L+1$  and  $j = k$ . Namely:

$$1 + \alpha_1^{+k} + \alpha_j^{+k} = \frac{1}{\alpha_i^{+k}}$$

Since  $\alpha_1^{+k} = \frac{1}{\alpha_i^{+k}} (\alpha_k) = \frac{1}{\alpha_i^{+k}} (\alpha)_{i=k}$ , one finally gets (by using again the expression of  $(i-1)^{+k}$  given in (25)):

$$\alpha_j^{+k} = \frac{1}{\alpha_i^{+k}} \prod_{i=1}^{L+1} (i-1)^{L+1} \frac{1}{\alpha_i^{+k}} = \frac{1}{\alpha_i^{+k}} \prod_{i=1}^{L+1} \frac{1}{\alpha_i^{+k}} + \frac{1}{\alpha_i^{+k}} \prod_{i=1}^{L+1} \frac{1}{\alpha_i^{+k}}$$

$$, \quad \alpha_j^{+k} = \frac{1}{\alpha_i^{+k}} \prod_{j=1}^{L+1} \frac{1}{\alpha_j^{+k}} \quad \text{for } j = k$$

A stationary  $EGR(L)^{+k}$  is locally stable under learning if and only if  $\alpha_j^{+k} < 0$  for each  $j = 1; \dots; L+1$  but  $j = k$  ( $k = 1; \dots; L+1$ ). The result follows.

### 5.3 Local Extended Growth Rate Dynamics

Recall that, from (25), we have:

$$\alpha_L^{+k} = (i-1)^{L+1} \frac{1}{\alpha_{L+1}^{+k}} (\alpha) \quad \text{and} \quad \alpha_{L+1}^{+k} = (i-1)^L \frac{1}{\alpha_L^{+k}} (\alpha) \quad (31)$$

If  $\alpha_L = 0$ , then  $\frac{1}{\alpha_{L+1}^{+k}} (\alpha) = \frac{1}{\alpha_{L+1}^{+k}} (\alpha) = 0$ , so that  $\alpha_i = 0$  whatever  $i$  is ( $i = 1; \dots; L+1$ ). Now, it follows from (26) that:

$$\alpha_L^{+k} = (i-1)^{L+1} \frac{1}{\alpha_{L+1}^{+k}} (\alpha_k) \quad \text{and} \quad \alpha_{L+1}^{+k} = (i-1)^L \frac{1}{\alpha_L^{+k}} (\alpha_k) \quad (32)$$

Since  $\frac{1}{\alpha_{L+1}^{+k}} (\alpha_k) = \frac{1}{\alpha_{L+1}^{+k}} (\alpha_k) = 0$  and since  $\alpha_i = 0$  when  $\alpha_L = 0$ , it follows that  $\alpha_L^{+k} = 0$  whatever  $k$  is ( $k = 1; \dots; L+1$ ). It remains consequently to show that, when  $\alpha_L = 0$ , we have:

$$\frac{\alpha_{L+1}^{+k}}{\alpha_L^{+k}} = \frac{\alpha_{L+1}^{+k}}{\alpha_L^{+k}} = \frac{\alpha_{L+1}^{+k}}{\alpha_L^{+k}} \quad (33)$$

Using definitions (31) and (32), this condition is equivalent to:

$$\frac{\frac{1}{\alpha_{L+1}^{+k}} (\alpha)}{\frac{1}{\alpha_L^{+k}} (\alpha_k)} = \frac{\frac{1}{\alpha_{L+1}^{+k}} (\alpha)}{\frac{1}{\alpha_L^{+k}} (\alpha_k)}$$

Observe now that  $\frac{1}{\alpha_{L+1}^{+k}} (\alpha) = \frac{1}{\alpha_{L+1}^{+k}} (\alpha_k) = \alpha_k$ , so that (33) rewrites:

$$\alpha_k \frac{1}{\alpha_{L+1}^{+k}} (\alpha_k) \prod_{i=1}^{L+1} \frac{1}{\alpha_i^{+k}} = 0, \quad \frac{1}{\alpha_L^{+k}} (\alpha_k) = 0$$

which is true when  $\alpha_L = 0$ . This proves the claim.

## 5.4 Proof of Proposition 7

The dynamics (24) in the neighborhood of  $^{+k}$  is obtained by linearizing (24) at  $^{+k}$ :

$${}^{\circ}B_k(\bar{^k}(t+1) | ^{+k}) = {}_1(1 + {}^{\circ}{}^{+k}_1)I_L(\bar{^k}(t) | ^{+k})$$

where:

$$\bar{^k}(t) = \begin{pmatrix} 0 & \bar{^k}_1(t) & 1 \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \\ 0 & \bar{^k}_{L+1}(t) & 1 \end{pmatrix}, \quad B_k = \begin{pmatrix} 0 & {}^{+k}_1 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & 0 & 1 & & \vdots \\ \vdots & \vdots & \vdots & & \ddots & 0 \\ 0 & {}^{+k}_{L+1} & 0 & \cdots & 0 & 1 \\ 0 & {}^{+k}_L & 0 & \cdots & \cdots & 0 \end{pmatrix}$$

Assume that  $\det B_k = {}^{+k}_L = 0$ , i.e.,  $\pm_L = 0$  (see appendix 4). One can express  $(\bar{^k}(t+1) | ^{+k})$  as a function of  $(\bar{^k}(t) | ^{+k})$  for  $\bar{^k}(t)$  near  $^{+k}$ . Namely:

$$(\bar{^k}(t+1) | ^{+k}) = F_k(\bar{^k}(t) | ^{+k}) \quad \text{where } F_k = {}_1 \frac{1 + {}^{\circ}{}^{+k}_1}{{}^{\circ}{}^{+k}_1} B_k {}_1$$

Let  $\mu_j^k$  and  $b_j^k$  ( $j = 1; \dots; L$ ) denote the  $L$  eigenvalues of  $F_k$  and  $B_k$  ( $k = 1; \dots; L+1$ ) respectively. Then we have:  $\mu_j^k = {}_1(1 + {}^{\circ}{}^{+k}_1)(b_j^k)$ . Remark now that  ${}^a_k = {}_1 {}^{\circ}B_k {}_1(1 + {}^{\circ}{}^{+k}_1)I_L$ , so that:  ${}^1_j = {}_1 {}^{\circ}b_j^k {}_1 {}^{\circ}{}^{+k}_1 {}_1$ . Hence:

$$\mu_j^k = \frac{1 + {}^{\circ}{}^{+k}_1}{(1 + {}^{\circ}{}^{+k}_1) + {}^1_j} = \frac{{}_j^k}{{}_j^k}$$

where the last equality comes by replacing  ${}^{+k}_1$  by  ${}_{L+1}({}_k)$  (see (26)). A given stationary EGR(L)  $^{+k}$  ( $k = 1; \dots; L+1$ ) is locally determinate in (24) if and only if  ${}^{\circ}\mu_j^k > 1$  ( $j = 1; \dots; L+1$  and  $j = k$ ), which holds true if and only if  $k = L+1$ . The result follows.

## References

- [1] J. Duay, On learning and the non-uniqueness of equilibrium in an overlapping generations model with fiat money, *J. Econ. Theory* 64 (1994), 541-553.
- [2] G. Evans, The fragility of sunspots and bubbles, *J. Monet. Econ.* 23 (1989), 297-317.
- [3] G. Evans and S. Honkapohja, On the robustness of bubbles in linear rational expectations models, *Int. Econ. Rev.* 33 (1992), 1-14.
- [4] G. Evans and S. Honkapohja, Stochastic gradient learning in the cobweb model, *Econ. Lett.* 61 (1998), 333-337.
- [5] J. M. Grandmont, Expectations formation and stability of large socioeconomic systems, *Econometrica* 66 (1998), 741-783.
- [6] J. M. Grandmont and G. Laroque, Stability of cycles and expectations, *J. Econ. Theory* 40 (1986), 138-151.
- [7] J. M. Grandmont and G. Laroque, Stability, expectations and predetermined variables, in "Essays in Honor of Edmond Malinvaud", vol. 1, (P. Champsaur, et al., Eds.), Cambridge MA, MIT press, 1990.
- [8] J. M. Grandmont and G. Laroque, Economic dynamics with learning: some instability examples, in "Equilibrium Theory and Applications: Proceedings of the Sixth International Symposium in Economic Theory and Econometrics" (W.A. Barnett et al., Eds.), Cambridge University Press, Cambridge U.K., 1991.
- [9] R. Guesnerie, Successes and failures in coordinating expectations, *Europ. Econ. Rev.* 37 (1992) 243-268.
- [10] R. Guesnerie and M. Woodford, Stability of cycles with adaptive learning rules, in "Equilibrium Theory and Applications: Proceedings of the Sixth International Symposium in Economic Theory and Econometrics" (W.A. Barnett et al., Eds.), Cambridge University Press, Cambridge U.K., 1991.
- [11] L. Ljung, Analysis of recursive stochastic algorithms, *IEEE Trans. Auto. Control* 22 (1977), 551-575.
- [12] L. Ljung and T. Soderström, "Theory and Practice of Recursive Identification", MIT press, Cambridge MA, 1983.
- [13] R. Lucas, Adaptive behavior and economic theory, *Journal of Business* 59 (1986), 401-426.

- [14] A. Marcet and T. Sargent, Convergence of least-squares learning mechanisms in self-referential stochastic models, *J. Econ. Theory* 48 (1989), 337-368.
- [15] E. Ramis, C. Deschamps and J. Odoux, “Cours de Mathématiques Spéciales”, Tome 1 Algèbre, Paris, Masson, 1974.
- [16] P. Reichlin, Equilibrium cycles in an overlapping generations economy with production, *J. Econ. Theory* 40 (1986), 89-102.
- [17] H. Robbins and S. Monro, A stochastic approximation method, *Annals of Math. Statistics* 22 (1951), 400-407.
- [18] M. Woodford, Stationary sunspot equilibria in a finance constrained economy, *J. Econ. Theory* 40 (1986), 128-137.
- [19] S. Wiggins, “Introduction to Applied Nonlinear Dynamical Systems and Chaos”, Springer Verlag (Texts in Applied Mathematics), 1990.
- [20] M. Woodford, Learning to believe in sunspots, *Econometrica* 58 (1990), 277-307.