



HAL
open science

A fractal approach to identifying urban boundaries

Cécile Tannier, Isabelle Thomas, Gilles Vuidel, Pierre Frankhauser

► **To cite this version:**

Cécile Tannier, Isabelle Thomas, Gilles Vuidel, Pierre Frankhauser. A fractal approach to identifying urban boundaries. *Geographical Analysis*, 2011, 43 (2), pp.211-227. 10.1111/j.1538-4632.2011.00814.x . hal-00731648

HAL Id: hal-00731648

<https://hal.science/hal-00731648>

Submitted on 14 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TANNIER C., THOMAS I., VUIDEL G., FRANKHAUSER P. (2011), A fractal approach to identifying urban boundaries, *Geographical Analysis*, vol. 43, n°2, pp. 211-227.

A Fractal Approach to Identifying Urban Boundaries

Cécile Tannier¹, Isabelle Thomas², Gilles Vuidel¹, Pierre Frankhauser¹

¹ThéMA, CNRS - University of Franche-Comté, Besançon, France, ²FRS-FNRS, CORE and Department of Geography, UCL, Louvain-la-Neuve, Belgium

Fractal geometry can be used for determining the morphological boundaries of metropolitan areas. A two-step method is proposed here: (1) Minkowski's dilation is applied to detect any multiscale spatial discontinuity; and, (2) a distance threshold is located on the dilation curve corresponding to a major change in its behavior. We therefore measure the maximum curvature of the dilation curve. The method is tested on theoretical urban patterns and on several European cities to identify their morphological boundaries and to track boundary changes over space and time. Results obtained show that cities characterized by comparable global densities may exhibit different distance thresholds. The less the distances separating buildings differ between an urban agglomeration and its surrounding built landscape, the greater the distance threshold. The fewer the buildings that are connected across scales, the greater the distance threshold.

Correspondence: Cécile Tannier, ThéMA CNRS-University of Franche-Comté, 32 rue Mégevand, F-25 030 Besançon Cedex, France
e-mail: cecile.tannier@univ-fcomte.fr

Submitted: May 29, 2009. Revised version accepted: April 1, 2010.

Introduction

Defining city boundaries is invariably an enthralling challenge for urban geographers and planners. Where do cities start and end? Is there a continuum of rural and urban land uses or a sharp divide between them? While some urban analyses (Parr 2007; Dujardin, Thomas, and Tulkens 2007) also include functional (e.g., socioeconomic, attractiveness) and operational (e.g., administrative, political) considerations, the focus here is on morphological (physical) criteria alone.

A general consensus does not appear to exist about the definition of the urban morphological agglomeration, or what Parr (2007) calls the “built city”. Working definitions often result from classifications or typologies of built patterns, and are based on applications of satellite and GIS technologies (e.g., Treitz, Howarth, and Gong 1992; Donnay, Barnsley, and Longley 2001; Abed and Kaysi 2003). Besides classifications, pre-defined morphological thresholds are commonly used (e.g., Donnay, Barnsley, and Longley 2001; Weber 2001). After classifying elementary spatial units so that similar ones can be grouped together, the criterion of contiguity—generally involving a distance threshold—is added. However, there is no consensus about the choice of contiguity criteria (Le Gléau, Pumain, and Saint-Julien 1997; Dujardin, Thomas, and Tulkens 2007), and the relevance of a distance threshold becomes questionable when analyzing urban fringes. European cities in particular exhibit irregular patterns where recent detached housing estates and traditional

rural buildings are mixed. Consequently, the spacing of neighboring buildings within sprawling patterns varies considerably, making application of a single distance threshold very difficult (Chaudhry and Mackaness 2008).

We suggest a method for identifying the morphological boundary of urban agglomerations that avoids the need for any pre-defined maximum distance threshold between buildings. Given that crucial spatial discontinuities can be identified by analyzing phenomena over a range of scales (Lam and Quattrochi 1992), we propose to use the conceptual and methodological tools of fractal geometry for defining the morphological boundaries of urban agglomerations.

Fractal geometry has been widely applied in geography for some thirty years now. Pioneering works argue that local and global spatial structures are interrelated at several scales (Goodchild 1980; Batty and Longley 1986; Arlinghaus 1985, 1987; Goodchild and Mark 1987; Frankhauser 1988). A string of publications specifically about urban geography shows that urban growth engenders a multiscale spatial organization (Fotheringham, Batty, and Longley 1989; Batty and Xie 1996, 1999; Benguigui et al. 2000; Shen 2002; Thomas, Frankhauser, and Biernaki 2008). Other contributions explore the fractal aspects of area–perimeter relations and the fractal boundary of built forms (Longley and Batty 1989; Arlinghaus and Nystuen 1990; Frankhauser 1994). Yet others discuss the scaling relationship between the number of built clusters and their size (White and Engelen 1993;

De Keersmaecker, Frankhauser, and Thomas 2003; Benguigui and Czamanski 2004; Benguigui, Blumenfeld-Lieberthal, and Czamanski 2006).

The main characteristic shared by fractals is their intrinsic multiscale spatial organization. They may, however, exhibit different kinds of multiscale behavior. For example, the elements of the familiar Sierpinski carpet are interconnected and actually form a single cluster (Mandelbrot 1982). By contrast, elements in a Fournier dust are unconnected and form clusters separated by empty lanes of different sizes: a Fournier dust contains just a few wide lanes and an increasing number of ever narrower ones. The lane widths (i.e., the distances separating the clusters) comply with a strict hierarchy (Mandelbrot 1982).

The hierarchy of empty lanes characterizing Fournier dusts can be explored by the dilation method for determining fractal dimensions proposed by Minkowski (1903) and further developed by Bouligand (1929). In dilation, each built element of a study area is surrounded by a buffer zone that widens at each step of the process. As dilation progresses, what were initially separate buildings become merged and clusters form. As these clusters expand, their number decreases. Dilation stops when just one big cluster remains. In fractals like Fournier dusts, the number of clusters N is related to the size of the dilation buffer ε according to a power-law function (equation 1).

$$N = a\varepsilon^D \tag{1}$$

where a is a constant of proportionality. The scaling exponent of this law is the fractal dimension D . If we take the logarithm of both sides, this expression has the form of a linear relationship with slope D (equation 2).

$$\log(N) = D \log(\epsilon) + \log(a) \quad (2)$$

In geography, dilation is used for analyzing and segmenting various types of images (e.g., remotely sensed images, aerial photographs). Some researchers (Pesaresi and Benediktsson 2000; Benediktsson, Pesaresi, and Arnason 2003) apply mathematical morphology (that uses dilation and erosion processes) to detect the boundaries of urban objects in remotely sensed images. With their method, local discontinuities emerge from an iterative transformation of the pattern itself and pre-determined thresholds are eschewed. However, few applications concern dilation for delimiting built agglomerations, and those applications that exist introduce pre-defined distance thresholds to identify morphological urban boundaries (e.g., Donnay 1994; Bailly 1996; Weber 2001; Guérois 2003).

Here we use dilation for identifying the morphological boundaries of urban agglomerations without introducing pre-defined distance thresholds. We begin with the log-log plot that counts the number of built clusters at each dilation step. If the built area of a city is fractal, then the number of clusters is related to their size by a power-law function corresponding to a linear relation visible on a log-log plot. We posit that the point at which the dilation curve

deviates most from a straight line is a crucial distance threshold, corresponding to the morphological multiscale boundary of an urban agglomeration. This boundary, referred to here as the *urban envelope*, is a spatial discontinuity across scales (Frankhauser and Tannier 2005). It separates two morphological spatial subsets that are distinct in fractal terms, and corresponds to a break point in the multiscale organization of a built pattern. Beyond this dilation threshold, the pattern is no longer self-similar, or its self-similarity changes dramatically.

Below a dilation threshold, the self-similarity of a pattern may be strictly fractal. However, the fractal dimension D is not expected to be constant in reality (Goodchild 1980); most often, it is constant over a limited range of scales, but varies somewhat over successive ranges of scales (Lam 1990; Lam and Quattrochi 1992; White and Engelen 1994; Tannier and Pumain 2005). Hence the fractality of a pattern below its dilation threshold is most probably multifractal.¹

Our aim is to identify an urban boundary, yet the possible existence of a rural–urban continuum cannot be ignored (Champion and Hugo 2004). Consequently, we do not assume that an urban envelope always exists—it exists only if two spatial organizations can be distinguished from a multiscale perspective. If this is the case, then a distance threshold can be identified; otherwise, in the case of a rural–urban continuum, detecting any discontinuity between the urban agglomeration and the outlying area is impossible.

Identifying an urban envelope

The data used are vector maps representing buildings in 2D (polygons) (Fig. 1). The spatial extent of a study area is necessarily large because it comprises an (either monopolar or multipolar) urban agglomeration and its hinterland (i.e., a suburban or rural area that is under the influence of its urban core).

A two-step method is adopted for identifying an urban envelope: (1) Minkowski's (1903) dilation is applied to detect the existence of any multiscale spatial discontinuity; and, (2) a distance threshold is located on the dilation curve corresponding to a major shift in its behavior. We therefore measure the maximum curvature of the dilation curve (Lowe 1989).




Figure 1. Sample of data used (excerpt from the 1957 map of Basel, see Fig. 7).

Minkowski dilation

A vector version of the Minkowski dilation is applied to each built polygon of a map. This consists in surrounding each built polygon with a buffer of increasing width. The buffer width increases according to a geometric logic, which corresponds to the fractal logic. The number of clusters is counted after each dilation step. The results are portrayed as a log-log plot, where the X-axis represents the width of the dilation buffer and the Y-axis represents the corresponding number of built clusters (Fig. 2a).

Identification of a significant threshold on a dilation curve

A significant threshold on a dilation curve is defined as the point characterized by the main value of curvature. The curvature function measures how a curve deviates to a greater or lesser extent from being straight at a given point (Lowe 1989). For each point on a curve, this value may be calculated as:

$$\kappa = \left(\frac{y''}{(1 + y'^2)^{3/2}} \right) \quad (3)$$

where y' (first derivative) measures the speed at which the number of built clusters decreases in the course of dilation, and y'' (second derivative) measures the acceleration of this decrease. The formula defines the curvature as a ratio of speed to acceleration. For a straight line, the speed is constant and the acceleration is zero. Hence, the main curvature

point corresponds to the maximum deviation of a curve with regard to a linear decrease in the number of built clusters in the course of dilation.

To identify the point of main curvature, a dilation curve is approximated by a series of polynomials of increasing degrees. We then select the polynomial of the lowest degree that best represents the original curve (Fig. 2b). This choice is based on Schwartz's (1978) Bayesian Information Criterion (BIC) for evaluating the trade-off between goodness-of-fit and complexity for statistical models. The BIC is stricter than Akaike's (1974) Information Criterion (AIC) in penalizing the loss of degrees of freedom by having more parameters in a fitted model. The polynomial degree chosen here corresponds to the point at which the BIC bottoms out or begins to increase. In any case, the adjusted correlation coefficient (R^2) between the real and estimated curves must be greater than 0.9.

Once the estimated curve is obtained, points of maximum curvature are identified (Fig. 2c): they are characterized by a zero derivation value of the curvature function. In order to avoid points of maximum curvature arising from estimation artifacts, the two extremities of an estimated curve are removed before calculating the curvature function (Fig. 2d). This adjustment implies that the distance threshold determining an urban envelope is neither a few meters (too small) nor several kilometers (too large). The point of main curvature has the highest absolute value of curvature among the points of maximum curvature. Finally, the point of main curvature is located on the estimated curve, which gives the distance

threshold for drawing an urban envelope (Fig. 2d).




Figure 2. Identification of a significant threshold on the dilation curve (the metropolitan area of Besançon, France example).

The step-by-step vector dilation and cluster count were performed for a computer application developed with the *Java Topology Suite* and *Geotools* library. The polynomial estimations of the dilation curve, the BIC, and the curvature function were computed using

Octave, an open source version of *Matlab*. The two applications are integrated in the software MorphoLim (<https://sourcesup.renater.fr/morpholim/>).



Application to theoretical urban patterns

Three theoretical patterns are studied here. Fig. 3a portrays a regular fractal pattern constructed in accordance with a strict iterative logic and characterized by strict self-similarity. This hybrid Sierpinski carpet combines a Sierpinski carpet consisting of a single cluster and a Fournier dust of unconnected elements at each step of iteration (Frankhauser and Tannier 2005). As with Fournier dusts, the hierarchy of distances between built clusters in the hybrid Sierpinski carpet obeys a power-law function. The hybrid carpet of Fig. 3a comprises 28,500 identical square buildings.


Fig. 3b portrays a random fractal pattern characterized by quasi-self-similarity. As with the hybrid carpet, its elements are partly connected and partly unconnected. A single scaling law defines the fractal organization of the pattern, but the respective locations of the elements vary randomly. As with a real city, elements are not all of the same size.

Fig. 3c portrays a regular fractal pattern in a nonfractal environment. It is the same hybrid carpet as in Fig. 3a, but this time it is surrounded by randomly and fairly uniformly distributed elements. These elements are like noise around the fractal pattern. Fig. 3c can be seen as a metaphor of a city surrounded by rural settlements.



(a) Regular fractal city



(b) Random fractal city



(c) Regular fractal city in a non fractal environment



In grey: spatial extent of the urban envelope

Figure 3. The envelopes of the three theoretical cities.

The multiscale envelope of each pattern has been identified by the method previously proposed. For the regular fractal city, the dilation curve exhibits some plateaus but is essentially linear: any shift along the X-axis has a corresponding similar shift along the Y-axis (Fig. 3a). This synchronized shifting is because a fractal relationship associates a geometric (hence multiplicative) change in the number of elements with a geometric change in their size (or in the distance between them). A log-log plot changes a geometric progression into a linear progression. The slope of the curve is the fractal dimension D , which is constant for a simple regular fractal pattern regardless of the scale. Consequently, the dilation curve is suitably estimated by a first degree polynomial (i.e., a linear function), as shown in Fig. 4. In this case, the second derivative is always zero; consequently, the curvature function is zero, too. No point of maximum curvature can be identified, and thus no significant threshold exists in the dilation curve. Therefore, an envelope is impossible to identify for a subset of elements of the image: the envelope here bounds the entire image.

In the case of the random fractal city, the dilation curve is smoother. The randomness of the pattern means that no plateau appears (Fig. 3b). However, the curve is linear (see Fig. 4—high adjusted R^2 for a first degree polynomial). No main curvature can be identified, and the envelope bounds the entire pattern. By contrast, for the hybrid carpet in a nonfractal environment, the dilation curve is no longer linear (Figs. 3c and 4). A clear threshold appears. This is an important feature arising from the spatial organization of the pattern under study—it reveals a shift in the behavior of the pattern in the course of dilation. For

simple theoretical fractal patterns, spatial relations between elements remain the same regardless of the scale of analysis, whereas for more complex patterns like the hybrid carpet in a nonfractal environment, the spatial organization between two scales may vary. Such a variation can be interpreted as a fundamental morphological discontinuity indicative of a sharp divide between two subsets of elements. Such a discontinuity corresponds to a change in the fractal dimension between two scales.




Figure 4. Adjusted R^2 for the series of polynomial estimations of the dilation curve.

Application to European cities

Six French and Belgian urban agglomerations

The usefulness and operability of the preceding method can be illustrated with some small- and medium-sized European urban agglomerations. Here we study three French cities (Besançon, Belfort, and Montbéliard) near Switzerland, and three Belgian cities (Namur, Liège, and Charleroi) in the former industrial belt in Wallonia. Their populations

range from about 80,000 inhabitants for Belfort to 500,000 inhabitants for Liège.

The study areas considered for each city are the French and Belgian *aires urbaines* (metropolitan areas) (INSEE 2009; Luyten and Van Hecke 2008). In France, an *aire urbaine* encompasses a densely-built *unité urbaine*—which is akin to a United States urban area, and its commuter belt. The Belgian definition of *aire urbaine* is similar to the French one. The spatial extent of the six agglomerations studied varies from 3.7 sq km (Belfort) to 28.5 sq km (Liège).

Data used for the French agglomerations come from the ‘BD Topo’ vectorial data base of the *Institut Géographique National* (IGN 2009). All buildings with footprints of more than about 1 sq m are mapped regardless of their function (e.g., residential, commercial, industrial). For the Belgian agglomerations, data are from the PLI (*Plan de Localisation Informatique*) developed by MRW-DGATLP (2004). The PLI is based on digitized topographic maps at a scale of 1:10,000 (IGN) and the land register (*cadastre*). Polygons representing buildings are stored in a vector data layer.

The shapes of the dilation curves obtained for each metropolitan area are very similar (Fig. 5), but the curves for the Belgian metropolitan areas are a little straighter, and with a lower main curvature value. This is understandable because Belgian metropolitan areas are very prone to urban sprawl (Caruso 2001; Champion 2001). Belgian suburban (sprawling) patterns, however, are mainly fractal and exhibit multiscale spatial contrasts (Thomas,

Frankhauser, and Biernaki 2008). By comparison, French urban agglomerations typically are individual cities that stand clearly apart from their rural surroundings. Belgium is a very small and highly urbanized country (approximately 30,000 sq km for 10 million inhabitants) compared to France (657,000 sq km for 65 million inhabitants); the structures of the two countries' urban systems are also different.




Figure 5. Dilation curves and distance thresholds for six Belgian and French agglomerations.

Besides the shape of the dilation curves, distance thresholds are clearly different, ranging from 465 m, for Namur, to 170 m, for Belfort. Study areas characterized by comparable global densities may exhibit different dilation thresholds. This tendency is consistent with the contention that a same density corresponds to various fractal dimensions. The information captured by fractal dimensions measured for areas is fundamentally different from the information given by density (for mathematical explorations of the differences and

relationship between density and fractal dimension, see Batty and Kim 1992; Batty and Xie 1996; Thomas, Frankhauser, and De Keersmaecker 2007). The less an urban agglomeration differs from its surrounding built landscape (in terms of variations of distances separating buildings), the greater the distance threshold. Moreover, the fewer the buildings in a study area that are connected across scales, the greater the distance threshold. The example of two fractal forms, the Fournier dust and the Sierpinski carpet (Mandelbrot 1982), is useful for illustrating this idea. The Fournier dust is made up of *unconnected* elements across scales, whereas the Sierpinski carpet is made up of *connected* elements across scales. European urban areas usually combine features of both Fournier dusts and Sierpinski carpets with regard to connectivity between buildings (Thomas, Frankhauser, and De Keersmaecker 2007). If an urban agglomeration is more like a Sierpinski carpet than a Fournier dust, and at the same time clearly different from its surrounding built landscape, then it exhibits a low dilation threshold (e.g., Belfort and Liège). In contrast, an urban agglomeration more like a Fournier dust than a Sierpinski carpet, and whose spatial organization is similar to that of its surrounding built pattern, exhibits a high dilation threshold (e.g., Montbéliard).

Diachronic analysis of the Basel urban agglomeration

We worked here with three vector maps of the built pattern of the Basel agglomeration and its hinterland for 1882, 1957 and 1994. The spatial extent of the study area is approximately 530 sq km. It extends over three European countries: Switzerland, France, and Germany. The urban agglomeration population was about 170,000 inhabitants in 1999. The original

data used for making the three maps are topographic maps from the Bundesamt für Landestopographie, 3084 Wabern, Germany. They are at different scales (from 1:25,000 to 1:50,000) and were drawn using different cartographic generalization methods (varying by date and country). Because of these differences, cartographic adjustments were required to develop a consistent set of vector built maps. Each polygon displayed with one of the finalized vector maps provides a fairly accurate representation of a building or part of a building. Small buildings, such as huts or sheds, are not mapped.

Results for the urban envelope of Basel at three points in time are presented in Fig. 6. The dilation curve for Basel in 1882 has two different sections. The first is characterized by a rather slow decline in the number of built clusters from one dilation step to the next, whereas the second exhibits a faster rate of decline. This trend is reminiscent of the spatial organization of the regular fractal city in a nonfractal environment (Fig. 3c). The two portions of the curve correspond to two spatial organizations of the elements on this map: a clearly delimited urban center, and small, well separated, built clusters. The distance threshold identified corresponds to the spatial boundary between these two spatial organizations.




Figure 6. Identification of a significant distance threshold in the dilation curves for Basel.

The spatial organization of Basel in 1957 and 1994 is markedly different from that in 1882.

The number of built clusters declines much faster from one dilation step to the next.

Distance thresholds are more difficult to identify because the spatial organization is more hierarchic. The metropolitan areas for both dates are close to a strict fractal hierarchy of the built clusters, as in the random fractal city in Fig. 3b.

Distance thresholds differ for each of the three dates: 300 m for 1882, 279 m for 1957, and 173 m for 1994. Hence, the morphological boundary of an urban agglomeration based on a single pre-defined distance threshold is not consistent with the morphological history of the urban area of Basel. Its distance threshold has decreased over time. Two phenomena explain this decrease: (1) distances between buildings were greater in 1882 than in 1957 or 1994, and (2) the hierarchy of distances between buildings remained the same both within

the urban agglomeration and outside it over a wide range of distances in 1882, but not in 1957 or 1994.

The shape of the urban envelope changed over time (Fig. 7). The urban agglomeration expanded significantly between 1882 and 1957, especially in Switzerland where population growth was particularly high in the 1950s (Bundesamt für Statistik 1997). The urban agglomeration was located entirely within Switzerland in 1882, but became transnational in the first half of the twentieth century. This change may be explained by the development of the tram network since 1895 serving both the French and German zones of the metropolitan area. A ring of hamlets around the urban agglomeration persisted in all three countries.

Between 1957 and 1994, the urban agglomeration underwent fewer changes, although the morphological evolutions are still noteworthy. The dominant trend is the expansion of the urban agglomeration in France and its contraction in Germany and Switzerland. Earlier gaps in the built pattern were infilled in the French part of the agglomeration (near Hegenheim and Huningue). Conversely, gaps widened or formed in the German and Swiss part of the agglomeration (e.g., in the south of Basel and in Weil am Rhein); built clusters that belong to the urban agglomeration in 1957 become separated by 1994 (e.g., Pratteln in the east of the Swiss part of the agglomeration). Differences in the evolution of the urban agglomeration between France, on the one hand, and Germany and Switzerland, on the other, can be attributed to differences in planning and economic policies in the three

countries. However, the ring of hamlets around the urban agglomeration in the three countries remained.

Conclusions drawn on the basis of the identification of the envelope of Basel at these three dates are consistent with existing knowledge about the evolution of this metropolitan area. In particular, Reitel (2006) shows the late but rapid growth of the metropolitan area after 1880 to be a consequence of the development of its chemical industry: at that time, the metropolitan area spread in both France (this part of France being German at that time) and Germany because of the establishment of industrial outlets for goods for the German market. The same study shows that more recently, in spite of the creation of a transnational framework, the spatial organization of the urban agglomeration remains clearly different in the three countries (Reitel 2006).




Figure 7. Basel's boundaries in 1882 (a), 1957 (b), and 1994 (c). The boundaries of each urban agglomeration result from the successive application of a positive and a negative buffer the width of which is the distance threshold of Fig. 6. Data source: Bundesamt für Landestopographie, 3084 Wabern, Germany. Digitalized by Research team 'Image, Ville, Environnement', Strasbourg, France.

Conclusion

A fractal-based method is proposed for identifying the morphological boundaries of urban agglomerations by detecting multiscale changes in their built morphologies. A dilation of each individual building is first applied; then a distance threshold is detected for the dilation curve, leading to a characterization of its urban envelope. This distance threshold corresponds to the dilation step at which distances separating buildings no longer exhibit the same fractal (or multifractal) behavior. Thus, the traditional morphological criterion (the maximum distance threshold between neighboring buildings) is superseded by a multiscale definition of morphological connectivity. Theoretical and real-world examples illustrate the operability of this method.

While the proposed method can be applied to any metropolitan area regardless of its size or characteristics, significant thresholds may or may not be detected. If built elements are uniformly distributed in space, no urban agglomeration is distinguishable because a dilation threshold can be identified only if local variations in distances separating buildings change dramatically at some scale. In contrast, if local variation of the distances separating buildings is always the same across scales, or if distances separating buildings do not vary, no threshold appears.

A threshold identified for an agglomeration may vary with the geographic extent of a study area. Any such variation is interesting because it may reveal hidden features of the spatial organization of that study area. This contention is consistent with Openshaw's (1984) idea that the MAUP (Modifiable Areal Unit Problem) is not one that has to be solved using sophisticated means, but is a useful tool for exploring the multiscale structure of a phenomenon (Lam and Quattrochi 1992). Moreover, besides identifying an urban boundary for just one (either monocentric or polycentric) agglomeration, the method could be applied to larger spaces to search for morphological thresholds common to several metropolitan areas, or to identify differences between the thresholds found for a set of metropolitan areas and the threshold found for a wider study area encompassing that set of individual metropolitan areas. This approach could bridge the gap between studies of individual cities and of systems of cities. The method proposed could further our understanding of urban growth and the changes affecting urban settlements. This potential now needs to be tested more thoroughly over a wider range of case studies.

Note

1 Mandelbrot defines multifractality differently. He builds multifractal patterns by introducing several reduction factors into the same generator (Mandelbrot 1982). Such multifractal patterns are characterized by several fractal dimensions that do not change regardless of the range of scales. The two points of view (Mandelbrot's and that expressed in this paper), however, are not mutually exclusive.

References

- Abed, J., and I. Kaysi. (2003). "Identifying Urban Boundaries: Application of Remote Sensing and GIS Technologies." *Canadian Journal of Civil Engineering* 30(6), 992-9.
- Akaike, H. (1974). "A New Look at the Statistical Model Identification." *IEEE Transactions on Automatic Control* 19(6), 716–23.
- Arlinghaus, S. (1985). "Fractals Take a Central Place." *Geografiska Annaler. Series B, Human Geography* 67(2), 83-8.
- Arlinghaus, S. (1987). "Essays On Mathematical Geography II". Institute of Mathematical Geography, Monographs Series, Ann Arbor, Michigan.
- Arlinghaus, S.L., and J. D. Nystuen. (1990). "Geometry of Boundary Exchanges." *Geographical Review* 80(1), 21-31.
- Bailly, E. (1996). "Position de recherche sur une méthode de détermination d'un contour urbain. (Research Report on a Method for Determining an Urban Outline)." *Cybergeo: European Journal of Geography* 10. <http://www.cybergeo.eu/index2235.html>
- Batty, M., and K. Kim. (1992). "Form Follows Function: Reformulating Urban Population Density Function." *Urban Studies* 29(7), 1043-70.
- Batty, M., and P. Longley. (1986). "The Fractal Simulation of Urban Structure." *Environment and Planning A* 18, 1143-79.
- Batty, M., and Y. Xie. (1996). "Preliminary Evidence for a Theory of the Fractal City." *Environment and Planning A* 28, 1745-62.
- Batty, M., and X. Xie. (1999). "Self-Organized Criticality and Urban Development." *Discrete Dynamics in Nature and Society* 3, 109-24.
- Benediktsson, J., M. Pesaresi, and K. Arnason. (2003). "Classification and Feature Extraction for Remote Sensing Images from Urban Areas Based on Morphological Transformations." *IEEE Transactions on Geoscience and Remote Sensing* 41(9), 1940-9.
- Benguigui, L., E. Blumenfeld-Lieberthal, and D. Czamanski. (2006). "The Dynamics of the Tel Aviv Morphology." *Environment and Planning B: Planning and Design* 33, 269-84.
- Benguigui, L., and D. Czamanski. (2004). "Simulation Analysis of the Fractality of Cities." *Geographical Analysis* 36(1), 69-84.
- Benguigui, L., D. Czamanski, M. Marinov, and Y. Portugali. (2000). "When and Where Is a City Fractal?"

- Environment and Planning B: Planning and Design* 27, 507-19.
- Bouligand, G. (1929). "Sur la notion d'ordre de mesure d'un ensemble plan." *Bulletin des Sciences Mathématiques* II-53, 185-92.
- Caruso, G. (2001). "Peri-Urbanisation: the Situation in Europe - A Bibliographical Note and Survey of Studies In the Netherlands, Belgium, Great Britain, Germany, Italy and the Nordic Countries." Report prepared for the DATAR (with contributions from J. Cavailhès, D. Peeters, M. Rounsevell, and I. Thomas), Ministère de l'Aménagement du Territoire et de l'Environnement, France.
- Champion, A. G. (2001). "A Changing Demographic Regime and Evolving Polycentric Urban Regions: Consequences for the Size, Composition and Distribution of City Populations." *Urban Studies* 38(4), 657-77.
- Champion, A. G., and G. Hugo. (2004). *New Forms of Urbanization, Beyond the Urban-Rural Dichotomy*. edited by A. G. Champion, and G. Hugo. Ashgate, Aldershot, Hants.
- Chaudhry, O., and W. Mackaness. (2008). "Automatic Identification of Urban Settlement Boundaries for Multiple Representation Databases." *Computers, Environment and Urban Systems* 32, 95-109.
- De Keersmaecker, M.-L., P. Frankhauser, and I. Thomas. (2003). "Using Fractal Dimensions for Characterizing Intra-Urban Diversity. The Example of Brussels." *Geographical Analysis* 35(4), 310-28.
- Donnay, J.-P. (1994). "Agglomérations morphologiques et fonctionnelles, l'apport de la télédétection urbaine." *Acta Geographica Lovaniensia* 34, 191-9.
- Donnay, J.-P., M. Barnsley, and P. Longley. (2001). *Remote Sensing and Urban Analysis*. London, U.K.:Taylor and Francis.
- Dujardin, C., I. Thomas, and H. Tulkens. (2007). "Quelles frontières pour Bruxelles ? Une mise à jour." *Reflets et perspectives de la vie économique* XLVI(2-3), 155-76.
- Fotheringham, A.S., M. Batty, and P. A. Longley. (1989). "Diffusion-Limited Aggregation and the Fractal Nature of Urban Growth." *Papers of the Regional Science Association* 67, 55-69.
- Frankhauser, P. (1988). "Fractal Aspects of Urban Systems". *Sonderforschungsbereich 230 "Natürliche Konstruktionen"* 1, 67-76.
- Frankhauser, P. (1994). *La fractalité des structures urbaines*. Paris: Anthropos.
- Frankhauser, P., and C. Tannier. (2005). "A Multiscale Morphological Approach for Delimiting Urban Areas." Working paper presented at the 9th Conference on Computers in Urban Planning and Urban Management, London, U.K. <http://128.40.111.250/cupum/searchpapers/papers/paper392.pdf>
- Goodchild, M. (1980). "Fractals and the Accuracy of Geographical Measures." *Mathematical Geology* 12(2), 85-98.

- Goodchild, M., and D. Mark. (1987). "The Fractal Nature of Geographic Phenomena." *Annals of the Association of American Geographers* 77(2), 265-78.
- Guérois, M. (2003). "Les formes des villes européennes vues du ciel - Une contribution de l'image CORINE Land Cover à la comparaison morphologique des grandes villes d'Europe occidentale". PhD thesis, directed by D. Pumain, University of Paris I. <http://halshs.archives-ouvertes.fr/docs/00/04/59/86/PDF/tel-00004303.pdf>
- Lam, N. (1990). "Description and Measurement of Landsat TM Images Using Fractals." *Photogrammetric Engineering and Remote Sensing* 56(2), 187-95.
- Lam, N., and D. Quattrochi. (1992). "On the Issues of Scale, Resolution, and Fractal Analysis in the Mapping Sciences." *Professional Geographer* 44(1), 88-98.
- Le Gléau, J., D. Pumain, and T. Saint-Julien (1997). "Towns of Europe: to Each Country its Definition." *INSEE Studies* 6, 1-14.
- Longley, P., and M. Batty. (1989). "On the Fractal Measurement of Geographical Boundaries." *Geographical Analysis* 21(1), 47-67.
- Lowe, D. (1989). "Organization of Smooth Image Curves at Multiple Scales." *International Journal of Computer Vision* 3, 119-30.
- Luyten, S., and E. Van Hecke. (2007). "De Belgische Stadsgewesten 2001." *Statistics Belgium Working Paper* 14.
- Mandelbrot, B. (1982). *The Fractal Geometry of Nature*. San Francisco: Freeman.
- Minkowski, H. (1903). "Volumen und Oberfläche." *Mathematische Annalen* 57, 447-95.
- Openshaw, S. (1984). *The Modifiable Areal Unit Problem*. Concepts and Techniques in Modern Geography 38, Geo Books, Norwich (Norfolk), U.K.
- Parr, J. (2007). "Spatial Definitions of the City: Four Perspectives." *Urban studies* 44(2), 381-92.
- Pesaresi, M., and I. A. Benediktsson. (2000). "Image Segmentation Based on the Derivative of the Morphological Profile". In *Mathematical Morphology and its Application to Image and Signal Processing*, 179-88.
- Reitel, B. (2006). "Governance in Cross-Border Agglomerations in Europe – The Examples of Basel and Strasbourg." *Europa Regional* 14(1), 9-21.
- Schwarz, G. (1978). "Estimating the Dimension of a Model." *Annals of Statistics* 6(2), 461–464.
- Shen, G. (2002). "Fractal Dimension and Fractal Growth of Urbanized Areas." *International Journal of Geographical Information Science* 16(5), 437–519.
- Tannier, C., and D. Pumain. (2005). "Fractals in Urban Geography: a Theoretical Outline and an Empirical Example." *Cybergeo: European Journal of Geography* 307. <http://www.cybergeo.eu/index3275.html>

- Thomas, I., P. Frankhauser, and C. Biernacki. (2008). "The Morphology of Built-Up Landscapes in Wallonia (Belgium): a Classification Using Fractal Indices." *Landscape and Urban Planning* 84, 99–115.
- Thomas, I., P. Frankhauser, and M. De Keersmaecker. (2007). "Fractal Dimension versus Density of the Built-up Surfaces in the Periphery of Brussels." *Papers in Regional Science* 86(2), 287-307.
- Treitz, P., P. Howarth, and P. Gong. (1992). "Application of Satellite and GIS Technologies for Land-Cover and Land-Use Mapping at the Rural–Urban Fringe: a Case Study." *Photogrammetric Engineering and Remote Sensing* 58, 439–448.
- Weber, C. (2001). "Urban Agglomeration Delimitation Using Remote Sensing". In *Remote Sensing and Urban Analysis*, 145–159, edited by J. Donnay, M. Barnsley, and P. Longley. London, U.K.: Taylor and Francis.
- White, R., and G. Engelen. (1993). "Cellular Automata and Fractal Urban Form: a Cellular Modelling Approach to the Evolution of Urban Land-Use Patterns." *Environment and Planning A* 25(8), 1175-99.
- White, R., and G. Engelen. (1994). "Urban Systems Dynamics and Cellular Automata: Fractal Structures between Order and Chaos." *Chaos, Solitons and Fractals* 4(4), 563-83.