



HAL
open science

Specification tests of parametric dynamic conditional quantiles

J. Carlos Escanciano, Carlos Velasco

► **To cite this version:**

J. Carlos Escanciano, Carlos Velasco. Specification tests of parametric dynamic conditional quantiles. *Econometrics*, 2010, 159 (1), pp.209. 10.1016/j.jeconom.2010.06.003 . hal-00732534

HAL Id: hal-00732534

<https://hal.science/hal-00732534>

Submitted on 15 Sep 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Specification tests of parametric dynamic conditional quantiles

J. Carlos Escanciano, Carlos Velasco

PII: S0304-4076(10)00142-9

DOI: 10.1016/j.jeconom.2010.06.003

Reference: ECONOM 3386

To appear in: *Journal of Econometrics*

Received date: 5 August 2008

Revised date: 8 June 2010

Accepted date: 20 June 2010



Please cite this article as: Carlos Escanciano, J., Velasco, C., Specification tests of parametric dynamic conditional quantiles. *Journal of Econometrics* (2010), doi:10.1016/j.jeconom.2010.06.003

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

SPECIFICATION TESTS OF PARAMETRIC DYNAMIC CONDITIONAL QUANTILES*

J. CARLOS ESCANCIANO[†]

Indiana University

CARLOS VELASCO[‡]

Universidad Carlos III de Madrid

June 8, 2010

Abstract

This article proposes omnibus specification tests of parametric dynamic quantile models. Contrary to the existing procedures, we allow for a flexible specification, where a possibly continuum of quantiles are simultaneously specified under fairly weak conditions on the serial dependence in the underlying data generating process. Since the null limit distribution of tests is not pivotal, we propose a subsampling approximation of the asymptotic critical values. A Monte Carlo study shows that the asymptotic results provide good approximations for small sample sizes. Finally, an application suggests that our methodology is a powerful alternative to standard backtesting procedures in evaluating market risk.

Keywords and Phrases: Omnibus tests; Conditional quantiles; Nonlinear time series; Empirical processes; Quantile processes; Subsampling; Value-at-Risk; Tail Risk.

JEL classification: C12, C22.

*We would like to thank Roger Koenker and participants at the Econometrics in Rio Conference for helpful comments. Also, we would like to thank one referee, the associate editor and the joint editor for very helpful comments and suggestions. All errors are our own. Research funded by the Spanish Plan Nacional de I+D+I, reference number SEJ2007-62908.

[†]Economics Department, Indiana University, 100 S. Woodlawn, 47401, Bloomington, Indiana, USA; E-mail: jescanci@iu.edu. Parts of this paper were written while J. Carlos Escanciano was visiting the Cowles Foundation at Yale University and the Economics Department at Cornell University, whose hospitality is gratefully acknowledged. This author thanks Professor Peter C.B. Phillips and Professor Yongmiao Hong the possibility of these visits.

[‡]Corresponding author: Departamento de Economía. Universidad Carlos III de Madrid. Calle Madrid 126, 28903 Getafe (Madrid), Spain. E-mail: carlos.velasco@uc3m.es

1. INTRODUCTION

Quantile regression is a powerful alternative to least squares regression in a wide range of econometric applications that vary from labor economics or demand analysis to finance; see the special issue of *Empirical Economics* (2001, vol .26) and the references therein. Rather than relying on a single measure of conditional location, the quantile regression approach allows the researcher to explore a continuous range of conditional quantile functions, thereby providing a more complete and flexible analysis of the conditional dependence structure of the variables under consideration. A researcher interested in the whole conditional distribution will consider the specification of the conditional quantile at all quantile levels and some diagnostic on its global suitability. As a matter of fact, conditional Goodness-of-fit tests are of paramount importance in econometrics and finance, see e.g. Andrews (1997) and Corradi and Swanson (2006). On the other hand, a risk manager will not be interested in the whole Profit&Loss account's distribution but mainly in its left tail, and hence she will consider a set of small values of quantile levels, usually 1% or 5% as recommended by the Basel Accord (1996a). Obviously, one can envision many situations in economics where the interest is in the lower or upper parts of the distribution; see, for instance, studies of unemployment duration (e.g. Koenker and Xiao (2002) and references therein), and wage inequalities (e.g. Machado and Mata, 2005). For these various situations parametric quantile regressions have been shown to be a useful and flexible modelling strategy.

It is well-known, however, that inference procedures within parametric quantile models depend crucially on the validity of the specified parametric functional forms for the range of quantiles under consideration. For instance, the counterfactual decomposition described in Machado and Mata (2005), that has been recently used in many studies to analyze the gender gap in log wages across the distribution (see e.g. Albrecht, van Vuuren and Vroman, 2007), and the martingale transform methods in Koenker and Xiao (2002) depend crucially on the linear quantile specification. Therefore, it is important to develop powerful tests for the correct specification of parametric conditional quantiles over a possibly continuous range of quantiles of interest and under fairly general conditions on the underlying DGP. This is the main purpose of the present paper.

More precisely, suppose we observe a real-valued dependent variable Y_t , and the explanatory vector $I_{t-1} = (W'_{t-1}, Z'_t)' \in \mathbb{R}^d$, $d = s + m$, where $Z_t \in \mathbb{R}^m$, $m \in \mathbb{N}$, is an observable random vector (r.v.) and $W_{t-1} = (Y_{t-1}, \dots, Y_{t-s})' \in \mathbb{R}^s$, where A' denotes the matrix transpose of A . We assume throughout the article that the time series process $\{(Y_t, Z'_t)' : t = 0, \pm 1, \pm 2, \dots\}$, defined on the probability space (Ω, \mathcal{A}, P) , is strictly stationary and ergodic. Assuming that the conditional distribution of Y_t given I_{t-1} is continuous, we define the α -th conditional quantile of Y_t given I_{t-1}

as the measurable function q_α satisfying the conditional restriction

$$P(Y_t \leq q_\alpha(I_{t-1}) \mid I_{t-1}) = \alpha, \text{ almost surely (a.s.), } \alpha \in [0, 1]. \quad (1)$$

In *parametric* quantile regression modeling one assumes the existence of a family of functions $\mathcal{M} = \{m(\cdot, \theta(\alpha)) : \theta(\cdot) : \mathcal{T} \rightarrow \Theta \subset \mathbb{R}^p\}$, where \mathcal{T} is a compact set which comprises the range of quantiles of interest, $\mathcal{T} \subset [0, 1]$, and one proceeds to make inference on θ or to test if $q \in \mathcal{M}$, i.e., if there exists some $\theta_0 : \mathcal{T} \rightarrow \Theta$ such that $m(\cdot, \theta_0(\alpha)) = q_\alpha(\cdot)$ a.s. for all $\alpha \in \mathcal{T}$.¹

Leading examples of specifications \mathcal{M} are the Linear Quantile Regression (LQR) model

$$m(I_{t-1}, \theta_0(\alpha)) \equiv m(Z_t, \theta_0(\alpha)) = Z_t' \theta_0(\alpha), \quad \alpha \in \mathcal{T},$$

with the *location-scale* regression model as the prominent example, in which $\theta_0(\alpha) = (\beta_0, \gamma_0 F_0^{-1}(\alpha)) \in \Theta \subset \mathbb{R}^p$, $m = p$, and where $F_0^{-1}(\alpha)$ denotes a univariate quantile function, see, e.g., Koenker and Xiao (2002), or the Linear Quantile Autoregression model of order s (LQAR(s)), where

$$m(I_{t-1}, \theta_0(\alpha)) \equiv m(W_{t-1}, \theta_0(\alpha)) = \theta_{01}(\alpha) + W_{t-1}' \theta_{02}(\alpha), \quad \theta_0(\alpha) = (\theta_{01}(\alpha), \theta_{02}'(\alpha))',$$

which results, for instance, from the random coefficient model

$$Y_t = \theta_{01}(U_t) + W_{t-1}' \theta_{02}(U_t), \quad (2)$$

where $\theta_{01}(\cdot)$ and $\theta_{02}(\cdot)$ are such that the right hand side of (2) is monotone increasing in U_t , and $\{U_t\}$ are independent and identically distributed (*iid*) $U[0, 1]$ random variables; see Koenker and Xiao (2006) for inferences on the LQAR(s) model.

Much effort has been devoted to inferences on $\theta_0(\alpha)$ for the aforementioned models based on the associated quantile processes $Q_n(\alpha) := \sqrt{n}(\theta_n(\alpha) - \theta_0(\alpha))$, for $\theta_n(\alpha)$ a \sqrt{n} -consistent estimator of $\theta_0(\alpha)$. It is well-known, however, that inferences based on $Q_n(\alpha)$ will heavily depend on the correct specification of the parametric quantile regression model. Although there exist some works on quantile regression model checks, to the best of our knowledge no consistent test for $q \in \mathcal{M}$ has been proposed. The existing literature has been mostly limited to *iid* observations, linear models, and more importantly to a fixed quantile level $\alpha \equiv \alpha_0 \in (0, 1)$. In particular, Zheng (1998) has proposed a quantile regression specification test based on kernel smoothing estimators of the conditional moment $E[1(Y_t \leq m(I_{t-1}, \theta_0(\alpha_0))) - \alpha_0 \mid I_{t-1}]$; see also Horowitz and Spokoiny (2002) for the median function (i.e., $\alpha_0 = 0.5$). Recently, Whang (2005), using empirical likelihood methods, has proposed a specification test for quantile regression and censored quantile regression for *iid* data.

¹We can actually take $\mathcal{T} = [0, 1]$ in our theory provided the centered estimator $\sqrt{n}(\theta_n - \theta_0)$ is asymptotically tight on the whole interval $[0, 1]$. To the best of our knowledge, such result is, however, not available in the literature for most popular estimators. Thus, we do not pursue such generality in this paper and we restrict our analysis to $\mathcal{T} \subset [0, 1]$, in accordance with the econometrics literature.

Tests based on smoothers usually have known asymptotic null distributions after an appropriate choice of the bandwidth sequence, but they are not consistent against Pitman's local alternatives.

Using an integrated approach, Bierens and Ginther (2001) proposed a diagnostic test for a linear quantile regression. These authors consider *iid* observations and do not take into account the uncertainty due to parameter estimation. Their test is consistent against $n^{-1/2}$ local alternatives, with n the sample size, but it relies on an upper bound on the asymptotic critical value, which might be too conservative. To solve this deficiency, Whang (2006) considered a subsampling approach to approximate the asymptotic critical values. Koul and Stute (1999) introduced asymptotic pivotal tests for parametric conditional quantiles of first-order nonlinear autoregressive processes. To obtain the pivotal property of the test they use a martingale transform (cf. Khmaladze, 1981). Alternatively, He and Zhu (2003) developed a bootstrap-based test for linear and nonlinear quantile regressions. Our paper also contributes to this literature of specification tests for a unique quantile, since our methods trivially apply to the unique quantile case in a more general framework than the previously cited works. By extending the scope of conditional quantile specifications to a, possibly, continuum of quantiles we provide a new and flexible specification procedure.²

In the present article we propose omnibus tests for $q. \in \mathcal{M}$ that are valid for general linear and nonlinear quantile models under time series. Our tests are based on the fact that $q. \in \mathcal{M}$ is characterized by the *infinite* set of conditional moment restrictions

$$E[1(Y_t \leq m(I_{t-1}, \theta_0(\alpha))) - \alpha \mid I_{t-1}] = 0 \text{ a.s. for some } \theta_0(\cdot) : \mathcal{T} \rightarrow \Theta \subset \mathbb{R}^p, \forall \alpha \in \mathcal{T}. \quad (3)$$

The proposed tests are functionals of a quantile-marked empirical process that characterizes condition (3). The asymptotic theory is largely complicated by the fact that (3) involves an infinite number of conditional moment restrictions, indexed by $\alpha \in \mathcal{T}$. We solve this technical difficulty using delicate weak convergence results for empirical processes under martingale conditions. It turns out that the asymptotic null distributions of test statistics depend on the specification under the null and the DGP. Therefore, we propose to implement the test with the assistance of the subsampling.

The rest of the article is organized as follows. In Section 2 we introduce the quantile-marked process, which is the basis upon which the new test statistics for testing (3) are developed. We study the asymptotic distribution of the proposed tests under the null, fixed and local alternatives. In Section 3 a subsampling procedure for approximating the asymptotic null distribution of tests is considered. In Section 4 we present a simulation exercise assessing the finite-sample performance of

²During the revision of this paper, one referee pointed out an unpublished PhD dissertation by Nejmeldoen (2003) proposing specification tests for a continuum of quantiles in a linear quantile model. Nejmeldoen's (2003) results require high-level conditions. In particular, his Assumption (E) directly assumes that a certain empirical process is stochastic equicontinuous, which is certainly the most difficult part and the bulk in the development of our specification tests. On the contrary, we prove the stochastic equicontinuity condition under a set of primitive regularity assumptions and for a more general setting than that considered in Nejmeldoen (2003).

tests. Section 5 summarizes the results of an application to some European stock indexes, showing that our methodology can serve as a powerful and flexible alternative to standard backtesting procedures in evaluating market risk. Finally, Section 6 concludes. Proofs are deferred to an appendix. Throughout the article A^c and $|A|$ denote the complex conjugate and Euclidean norm of A , respectively. In the sequel C is a generic constant that may change from one expression to another. All limits are taken as $n \rightarrow \infty$.

2. TEST STATISTICS AND ASYMPTOTIC THEORY

We aim to test the null hypothesis

$$H_0 : E[\Psi_\alpha(Y_t - m(I_{t-1}, \theta_0(\alpha))) | I_{t-1}] = 0 \text{ a.s. for some } \theta_0 \in \mathcal{B} \text{ and for all } \alpha \in \mathcal{T},$$

against the nonparametric alternatives

$$H_A : P(E[\Psi_\alpha(Y_t - m(I_{t-1}, \theta(\alpha))) | I_{t-1}] \neq 0) > 0, \text{ for some } \alpha \in \mathcal{T} \text{ and for all } \theta(\alpha) \in \Theta \subset \mathbb{R}^p,$$

where $\Psi_\alpha(\varepsilon) = 1(\varepsilon \leq 0) - \alpha$, and \mathcal{B} is a family of uniformly bounded functions from \mathcal{T} to $\Theta \subset \mathbb{R}^p$. To simplify notation denote $\Psi_{\alpha,t}(\theta) \equiv \Psi_\alpha(Y_t - m(I_{t-1}, \theta))$ and $m_{t-1}(\theta) \equiv m(I_{t-1}, \theta)$. Note that under H_0 (and a mild continuity condition), $m_{t-1}(\theta_0)$ is identified as the α -th quantile of the conditional distribution of Y_t given I_{t-1} , for all $\alpha \in \mathcal{T}$. Testing for H_0 is a challenging testing problem since it involves an infinite number of non-smooth conditional moments parameterized by $\alpha \in \mathcal{T}$.

Our first aim is to characterize H_0 by the infinite number of unconditional moment restrictions

$$E[\Psi_{\alpha,t}(\theta_0) \exp(ix' I_{t-1})] = 0, \forall x \in \mathbb{R}^d, \text{ for some } \theta_0 \in \mathcal{B} \text{ and for all } \alpha \in \mathcal{T}, \quad (4)$$

where $i = \sqrt{-1}$ is the imaginary unit; see Bierens (1982). Instead of the exponential function we may also use, for instance, any of the parametric families considered in Escanciano (2006). Unreported simulations suggested that the exponential function performs favorably to other choices such as indicator functions. These simulations can be obtained from the authors upon request.

Given a sample $\{(Y_t, I_{t-1}')' : 1 \leq t \leq n\}$ and a parameter value $\theta \in \mathcal{B}$, we consider the quantile-marked empirical process indexed by $x \in \mathbb{R}^d$, $\alpha \in \mathcal{T}$ and $\theta \in \mathcal{B}$,

$$S_n(x, \alpha, \theta) := n^{-1/2} \sum_{t=1}^n \Psi_{\alpha,t}(\theta) \exp(ix' I_{t-1}).$$

Associated to S_n are the quantile-marked *error* and *residual* processes, respectively, defined by

$$R_n(x, \alpha) \equiv S_n(x, \alpha, \theta_0) \quad \text{and} \quad R_n^1(x, \alpha) \equiv S_n(x, \alpha, \theta_n),$$

for a \sqrt{n} -consistent estimator $\theta_n(\alpha)$ of $\theta_0(\alpha)$, say. The null hypothesis is likely to hold when the process $R_n^1(x, \alpha)$ is close to zero for almost all $(x', \alpha)' \in \mathbb{R}^d \times \mathcal{T}$.

The most popular estimator of θ_0 is the Quantile Regression Estimator (QRE), initially proposed by Koenker and Basset (1978) for linear models, and subsequently generalized to other frameworks by numerous authors, see references below. The QRE is defined as any solution $\theta_{KB,n}(\alpha)$ minimizing

$$\theta \mapsto \sum_{t=1}^n \rho_\alpha(Y_t - m(I_{t-1}, \theta))$$

with respect to $\theta \in \Theta \subset \mathbb{R}^p$, where $\rho_\alpha(\varepsilon) = -\Psi_\alpha(\varepsilon)\varepsilon$. Koenker and Park (1996) discussed the existence of $\theta_{KB,n}(\alpha)$ and an interior point algorithm for its computation.

Basset and Koenker (1978) proved the consistency and asymptotic normality of $\theta_{KB,n}(\alpha)$ in the linear regression model, including the least absolute deviation estimator, see also Pollard (1991). Moreover, the asymptotic theory for $\sqrt{n}(\theta_{KB,n}(\cdot) - \theta_0(\cdot))$ as a process indexed by the parameter $\alpha \in \mathcal{T}$, has been considered, among others, in Gutenbrunner and Jurečková (1992) and Gutenbrunner, Jurečková, Koenker and Portnoy (1993) for linear models, in Koul and Saleh (1994) and Jurečková and Hallin (1999) for linear autoregressions, and in Mukherjee (1999) for nonlinear autoregressions. For early contributions see Portnoy (1984). In the present article we do not restrict ourselves to $\theta_{KB,n}$ and we consider any estimator θ_n satisfying some mild conditions, see A4 below. For instance, our results apply to the Quasi-Maximum Likelihood Estimator in Komunjer (2005), under suitable regularity conditions.

The process R_n^1 is a mapping from (Ω, \mathcal{A}, P) with values in $\ell^\infty(\Pi)$, where $\ell^\infty(\Pi)$ is the space of all complex-valued functions that are uniformly bounded on Π , with $\Pi := \Upsilon \times \mathcal{T}$, and Υ a generic compact subset of \mathbb{R}^d containing the origin. The space $\ell^\infty(\Pi)$ is furnished with the supremum metric, say d_∞ , and let \mathcal{B}_{d_∞} be the corresponding Borel σ -algebra. Let \implies denote weak convergence on $(\ell^\infty(\Pi), \mathcal{B}_{d_\infty})$ in the sense of J. Hoffmann-Jørgensen, see, e.g., Dudley (1999, p. 94), or Definition 1.3.3 in van der Vaart and Wellner (1996). Since Υ is generic, \implies is indeed weak convergence on compacta.

After (4), test statistics are based on a distance from the standardized sample analogue of $E[\Psi_{\alpha,t}(\theta_0) \exp(ix' I_{t-1})]$ to zero, i.e., on a norm of R_n^1 , say $\Gamma(R_n^1)$. A popular norm is the Cramér-von Mises (CvM) functional

$$CvM_n := \int_{\Pi} |R_n^1(x, \alpha)|^2 d\Phi(x) dW(\alpha), \quad (5)$$

where Φ and W are some integrating measures on Υ and \mathcal{T} , respectively. Other continuous (with respect to d_∞) functionals Γ from $\ell^\infty(\Pi)$ to \mathbb{R} are of course possible. Our simulations suggested that CvM_n outperforms other alternative functionals such as the Kolmogorov–Smirnov-type functional

$$KS_n := \sup_{\alpha \in \mathcal{T}} \int_{\Upsilon} |R_n^1(x, \alpha)|^2 d\Phi(x). \quad (6)$$

Therefore, we focus in the rest of paper on CvM_n . We reject the null hypothesis H_0 for “large”

values of CvM_n . Practical issues about the computation of the test statistic CvM_n are deferred to Section 4.

2.1 Asymptotic null distribution.

In this subsection we establish the limit distribution of the quantile-marked empirical process R_n^1 under the null hypothesis H_0 . The null limit distributions of the tests are the limit distributions of some continuous functionals of R_n^1 . To derive asymptotic results we consider the following notation and assumptions. Throughout the paper the family \mathcal{B} , in which the parameter θ_0 takes values, is endowed with the sup norm, i.e., $\|\theta\|_{\mathcal{B}} = \sup_{\alpha \in \mathcal{T}} |\theta(\alpha)|$. Let, for each $t \in \mathbb{Z}$, $\mathcal{F}_t = \sigma(I'_t, I'_{t-1}, \dots)$, be the σ -field generated by the information set obtained up to time t . Define for each $t \in \mathbb{Z}$, the quantile “innovation” $\varepsilon_{t,\alpha} := Y_t - q_\alpha(I_{t-1})$ and the parametric quantile “error” $e_t(\theta) \equiv e_t(\theta(\alpha)) := Y_t - m(I_{t-1}, \theta(\alpha))$. Define also the family of conditional distributions $F_x(y) := P(Y_t \leq y \mid I_{t-1} = x)$. Let f_x be the density function of the cumulative distribution function (cdf) F_x . In particular, $f_{I_{t-1}}(y)$ denotes the density of Y_t given I_{t-1} , evaluated at y . Let $N_{[\cdot]}(\delta, \mathcal{G}, \|\cdot\|)$ be the δ -bracketing number of a class of functions \mathcal{G} with respect to a norm $\|\cdot\|$, i.e., the smallest number r such that there exist f_1, \dots, f_r and $\Delta_1, \dots, \Delta_r$ such that $\max_{1 \leq i \leq r} \|\Delta_i\| < \delta$ and for all $f \in \mathcal{G}$, there exists a $1 \leq i \leq r$ such that $\|f - f_i\| < \Delta_i$, see Definition 2.1.6 in van der Vaart and Wellner (1996).

Assumption A1:

A1(a): $\{(Y_t, Z'_t)' : t = 0, \pm 1, \pm 2, \dots\}$ is a strictly stationary and ergodic process. Under H_0 , $\{\Psi_{\alpha,t}(\theta_0), \mathcal{F}_t\}$ is a martingale difference sequence for all $\alpha \in \mathcal{T}$.

A1(b): The parametric family $m(\cdot, \theta_0(\alpha))$ is nondecreasing in α a.s..

A1(c): $E[|I_0|^2] < C$.

A1(d): The family of distributions functions $\{F_x, x \in \mathbb{R}^d\}$ has Lebesgue densities $\{f_x, x \in \mathbb{R}^d\}$ that are uniformly bounded away from zero for the quantiles of interest, i.e., $\inf_{x \in \mathbb{R}^d, \alpha \in \mathcal{T}} |f_x(m(x, \theta(\alpha)))| \geq C > 0$, satisfy

$$\sup_{x \in \mathbb{R}^d, y \in \mathbb{R}} |f_x(y)| \leq C,$$

and are equicontinuous: for every $\epsilon > 0$ there exists a $\delta > 0$ such that

$$\sup_{x \in \mathbb{R}^d, |y-z| \leq \delta} |f_x(y) - f_x(z)| \leq \epsilon.$$

Assumption A2: For each $\theta_1 \in \mathcal{B}$,

A2(a): There exists a vector of functions $g_{t-1} : \Theta \rightarrow \mathbb{R}^q$ such that $g_{t-1}(\theta_1(\alpha))$ is \mathcal{F}_{t-1} -measurable for each $t \in \mathbb{Z}$, and satisfies, for all $k < \infty$,

$$\sup_{1 \leq t \leq n, \|\theta_1 - \theta_2\|_{\mathcal{B}} \leq kn^{-1/2}} n^{1/2} \|m_{t-1}(\theta_2) - m_{t-1}(\theta_1) - (\theta_2 - \theta_1)' g_{t-1}(\theta_1)\|_{\mathcal{B}} = o_P(1)$$

A2(b): For all sufficiently small $\delta > 0$,

$$E \left[\sup_{\|\theta_1 - \theta_2\|_{\mathcal{B}} \leq \delta} |1(Y_t \leq m_{t-1}(\theta_1(\alpha))) - 1(Y_t \leq m_{t-1}(\theta_2(\alpha)))| \right] \leq C\delta, \quad \forall \alpha \in \mathcal{T}, \text{ and}$$

$$E \left[\sup_{|\alpha_1 - \alpha_2| \leq \delta} |m_{t-1}(\theta_1(\alpha_1)) - m_{t-1}(\theta_1(\alpha_2))| \right] \leq C\delta.$$

A2(c): Uniformly in $\alpha \in \mathcal{T}$, $E |g_{t-1}(\theta_1(\alpha))|^2 < \infty$, and uniformly in $(x', \alpha)' \in \Pi$,

$$\left| \frac{1}{n} \sum_{t=1}^n g_{t-1}(\theta_0(\alpha)) \exp(ix' I_{t-1}) f_{I_{t-1}}(m_{t-1}(\theta_0)) - E [g_{t-1}(\theta_0(\alpha)) \exp(ix' I_{t-1}) f_{I_{t-1}}(m_{t-1}(\theta_0))] \right| = o_P(1).$$

Assumption A3: The parametric space Θ is compact in \mathbb{R}^p . The true parameter $\theta_0(\alpha)$ belongs to the interior of Θ for each $\alpha \in \mathcal{T}$, and $\theta_0 \in \mathcal{B}$. The class \mathcal{B} satisfies

$$\int_0^\infty (\log(N_{[\cdot]}(\delta^2, \mathcal{B}, \|\cdot\|_{\mathcal{B}})))^{1/2} d\delta < \infty.$$

Assumption A4: The estimator θ_n satisfies that $P(\theta_n \in \mathcal{B}) \rightarrow 1$ as $n \rightarrow \infty$, and the following asymptotic expansion under H_0 , uniformly in $\alpha \in \mathcal{T}$,

$$Q_n(\alpha) = \sqrt{n}(\theta_n(\alpha) - \theta_0(\alpha)) = \frac{1}{\sqrt{n}} \sum_{t=1}^n l_\alpha(Y_t, I_{t-1}, \theta_0(\alpha)) + o_P(1),$$

where $l_\alpha(\cdot)$ is such that $E[l_\alpha(Y_1, I_0, \theta_0(\alpha))] = 0$, $L_\alpha(\theta_0(\alpha)) = E[l_\alpha(Y_1, I_0, \theta_0(\alpha))]'_\alpha(Y_1, I_0, \theta_0(\alpha))$ exists and is positive definite, and $E[l_\alpha(Y_t, I_{t-1}, \theta_0(\alpha))\Psi_\alpha(Y_s - m(I_{s-1}, \theta_0(\alpha)))] = 0$ if $t \neq s$. Furthermore, as a process in $\ell^\infty(\mathcal{T})$, $Q_n(\alpha)$ converges weakly to a Gaussian process $Q(\cdot)$ with zero mean and covariance function

$$K_Q(\alpha_1, \alpha_2) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{t=1}^n \sum_{s=1}^n E[l_{\alpha_1}(Y_t, I_{t-1}, \theta_0(\alpha_1)) l_{\alpha_2}(Y_s, I_{s-1}, \theta_0(\alpha_2))].$$

Assumption A1(a) is standard in the time series model checks literature, see, e.g., Koul and Stute (1999). A1(b) is natural in the present context. A1(c) is needed to prove the equicontinuity of the limit process of R_n and can be avoided if we replace $\exp(ix' I_{t-1})$ by $\exp(ix' \phi(I_{t-1}))$, with $\phi(\cdot)$ a one-to-one bounded mapping (see e.g. Bierens and Ginther, 2001). A1(d) is necessary for the tightness of the process R_n^1 and is required in Koul and Stute (1999). Assumptions A2(a)-A2(c) are classical in inference about nonlinear models, see Koul's (2002) monograph. A2 is satisfied for all models considered in the literature under mild moment assumptions, e.g. LQR and LQAR models. Sufficient conditions for A3 can be found in van der Vaart and Wellner (1996), see e.g. their Theorem 2.7.5 for monotone classes of functions, which applies to LQAR models. A4 has been established in the literature under a variety of conditions and different models and DGP's, see, for instance, Theorem 1 in Gutenbrunner and Jurečková (1992) or Theorem 3.2 in Mukherjee (1999). For nonlinear models

with *iid* innovations $(\varepsilon_t)_{t \in \mathbb{Z}}$ distributed as F_ε , Mukherjee (1999) proved A4 for $\theta_{KB,n}(\alpha)$. Under some mild additional assumptions, including that $\Sigma_{\theta_0(\alpha)} := E[g(I_1, \theta_0(\alpha))g(I_1, \theta_0(\alpha))']$ exists and is positive definite, Mukherjee (1999) showed that A4 holds for the QRE under H_0 with

$$l_\alpha(Y_t, I_{t-1}, \theta_0(\alpha)) = -\frac{\Sigma_{\theta_0(\alpha)}^{-1}g(I_{t-1}, \theta_0(\alpha))\Psi_\alpha(\varepsilon_t)}{q(\alpha)},$$

where $q(\alpha) = f_\varepsilon(F_\varepsilon^{-1}(\alpha))$ is the reciprocal of the sparsity function and f_ε is the density of F_ε . The quantile limit process $Q(\cdot)$ in that case is $\Sigma_{\theta_0(\cdot)}^{-1}W(\cdot)/q(\cdot)$, where $W(\cdot)$ denotes a vector of p independent Brownian bridges on \mathcal{T} .

We now establish the limit distribution of R_n . Under A1(a) and H_0 , because $R_n(v)$ is a zero-mean square-integrable martingale for each $v = (x', \alpha)' \in \Pi$, using a suitable Central Limit Theorem (CLT) for stationary ergodic martingale difference sequences, cf. Billingsley (1961), we have that the finite-dimensional distributions of R_n converge to those of a multivariate normal distribution with a zero mean vector and variance-covariance matrix given by the covariance function

$$K_\infty(v_1, v_2) = (\alpha_1 \wedge \alpha_2 - \alpha_1 \alpha_2)E[\exp(i(x_1 - x_2)'I_0)], \quad (7)$$

where, henceforth, $v_1 = (x_1', \alpha_1)'$ and $v_2 = (x_2', \alpha_2)'$ represent generic elements of Π , and \wedge denotes the minimum, i.e., $a \wedge b = \min\{a, b\}$. The next result is an extension of the convergence of the finite-dimensional distributions of R_n to weak convergence in the space $\ell^\infty(\Pi)$. We stress that no mixing conditions are required for the weak convergence to hold.

THEOREM 1: *Under the null hypothesis H_0 and Assumptions A1(a-c)*

$$R_n \Longrightarrow R_\infty,$$

where R_∞ is a Gaussian process with zero mean and covariance function (7).

In practice, θ_0 is unknown and has to be estimated from a sample $\{(Y_t, I_{t-1})' : 1 \leq t \leq n\}$ by an estimator θ_n . When we replace θ_0 in R_n by θ_n , resulting in R_n^1 , we need to investigate how the estimation error will affect the asymptotic properties of R_n^1 . The next result shows this effect on the asymptotic null distribution of R_n^1 . Define the function

$$G(x, \theta_0(\alpha)) := E[g_{t-1}(\theta_0(\alpha))f_{I_{t-1}}(m_{t-1}(\theta_0))\exp(ix'I_{t-1})], \quad x \in \Upsilon, \alpha \in \mathcal{T}.$$

THEOREM 2: *Under the null hypothesis H_0 and Assumptions A1-A4*

$$\sup_{x \in \Upsilon, \alpha \in \mathcal{T}} \left| R_n^1(x, \alpha) - R_n(x, \alpha) + G'(x, \theta_0(\alpha))n^{-1/2} \sum_{t=1}^n l_\alpha(Y_t, I_{t-1}, \theta_0(\alpha)) \right| = o_P(1).$$

As a consequence, we obtain the following corollary.

COROLLARY 1: *Under the assumptions of Theorem 2*

$$R_n^1 \implies R_\infty^1,$$

where $R_\infty^1(\cdot) = R_\infty(\cdot) - G'(\cdot, \theta_0(\cdot))Q(\cdot)$ (in distribution).

Now, using the last corollary and the Continuous Mapping Theorem (CMT) we obtain the asymptotic null distribution of continuous functionals such as CvM_n and KS_n .

COROLLARY 2: *Under the assumptions of Theorem 2, for any continuous functional $\Gamma(\cdot)$ from $\ell^\infty(\Pi)$ to \mathbb{R} ,*

$$\Gamma(R_n^1) \xrightarrow{d} \Gamma(R_\infty^1).$$

2.2 Consistency and Pitman's local alternatives.

In this section we study the consistency properties of tests based on functionals $\Gamma(R_n^1)$. First, we show that these tests are consistent against all fixed alternatives provided a mild regularity condition is satisfied.

Assumption A5: Under H_A , (i) there exists a $\theta_1 \in \mathcal{B}$ such that $\|\theta_n - \theta_1\|_{\mathcal{B}} = o_P(1)$; (ii) $E[\Psi \cdot (e_t(\theta_1(\cdot))) \exp(i \cdot I_{t-1})]$ is different from zero in a subset with positive Lebesgue measure on Π .

See Kim and White (2003) for conditions on $\theta_{KB,n}$ to satisfy Assumption A5(i), see also Section 3 in Angrist, Chernozhukov and Fernández-Val (2006). A sufficient condition for A5(ii) is that I_{t-1} is bounded. Notice that this condition always holds if we replace I_{t-1} by $\phi(I_{t-1})$, with ϕ a one-to-one bounded mapping, as in Bierens and Ginther (2001). Henceforth, almost sure convergence of nonmesurable maps is understood, as usual, as outer almost sure convergence, see van der Vaart and Wellner (1996) for definitions.

THEOREM 3: *Under the alternative hypothesis H_A and Assumptions A1, A2, A3 and A5,*

$$n^{-1/2} R_n^1(\cdot) \xrightarrow{a.s.} E[\Psi \cdot (e_t(\theta_1(\cdot))) \exp(i \cdot I_{t-1})].$$

A consequence of Theorem 3 and the CMT is that (under the same set of assumptions),

$$\int_{\Pi} \left| n^{-1/2} R_n^1(x, \alpha) \right|^2 d\Phi(x) dW(\alpha) \xrightarrow{P} \int_{\Pi} |E[\Psi_\alpha(e_t(\theta_1(\alpha))) \exp(ix' I_{t-1})]|^2 d\Phi(x) dW(\alpha) > 0,$$

provided that Φ and W are absolute continuous with respect to the Lebesgue measure on Π . In such a situation, the test statistic CvM_n will diverge to $+\infty$ under any fixed alternative, and the test will be consistent against all directions in the alternative hypothesis.

Now we analyze the asymptotic distribution of R_n^1 under a sequence of local alternatives converging to null at a parametric rate $n^{-1/2}$. We consider the DGP generating the local alternatives

$$H_{A,n} : E[\Psi_\alpha(Y_t - m_{t-1}(\theta_0)) | \mathcal{F}_{t-1}] = \frac{a_\alpha(I_{t-1})}{n^{1/2}} \text{ a.s. for some } \theta_0 \in \mathcal{B} \text{ and for all } \alpha \in \mathcal{T}, \quad (8)$$

where the function $a_\alpha(\cdot) : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfies the following assumption.

Assumption A6: $a_\alpha(\cdot)$ is such that $E \sup_{\alpha \in \mathcal{T}} |a_\alpha(I_{t-1})| < \infty$. There exists a \mathcal{F}_{t-1} -measurable r.v. C_{t-1} with $E[C_{t-1}^2] < \infty$, such that for all $t \in \mathbb{Z}$ and for all $\alpha_1, \alpha_2 \in \mathcal{T}$,

$$|a_{\alpha_1}(I_{t-1}) - a_{\alpha_2}(I_{t-1})| \leq C_{t-1} |\alpha_1 - \alpha_2|, \text{ a.s..}$$

To derive the next result we need the following assumption on the behaviour of the estimator under the local alternatives.

Assumption A4': The estimator $\theta_n(\alpha)$ satisfies the following asymptotic expansion under $H_{A,n}$, uniformly in α ,

$$\sqrt{n}(\theta_n(\alpha) - \theta_0(\alpha)) = \xi_a(\alpha) + \frac{1}{\sqrt{n}} \sum_{t=1}^n l_\alpha(Y_t, I_{t-1}, \theta_0(\alpha)) + o_P(1),$$

where the function $l_\alpha(\cdot)$ is as in A4 and $\xi_a(\alpha) \in \mathbb{R}^p$ for each $\alpha \in \mathcal{T}$.

Assumption A4' holds for most estimators considered in the literature. For instance, in the nonlinear time series context of Mukherjee (1999), the corresponding term $\xi_a(\alpha)$ to $\theta_{KB,n}(\alpha)$ is

$$\xi_a(\alpha) = -q^{-1}(\alpha) \Sigma_{\theta_0(\alpha)}^{-1} E[f_{I_{t-1}}(m_{t-1}(\theta_0)) g_{t-1}(\theta_0) a_\alpha(I_{t-1})].$$

The shift in charge of local power against alternatives in $H_{A,n}$ is given by

$$D_a(x, \theta_0(\alpha), \alpha) := E[a_\alpha(I_0) \exp(ix' I_0)] - \xi'_a(\alpha) G(x, \theta_0(\alpha)).$$

THEOREM 4: *Under the local alternatives (8), Assumptions A1-A3, A6 and A4'*

$$R_n^1 \implies R_\infty^1 + D_a,$$

where R_∞^1 is the process defined in Theorem 2.

It is not difficult to show that

$$D_a \equiv 0 \text{ a.e.} \iff a_\alpha(I_{t-1}) = \xi'_\alpha(\alpha)g(I_{t-1}, \theta_0(\alpha)) \text{ for all } \alpha \in \mathcal{T} \text{ a.s..}$$

Therefore, for directions $a_\alpha(\cdot)$ not collinear to the score $g(I_{t-1}, \theta_0(\alpha))$, the shift function D_a is non-trivial and test statistics based on $\Gamma(R_n^1)$ for a symmetric functional Γ are asymptotically strictly unbiased against the local alternatives (8); see Escanciano (2009) for a related formal proof of this statement.

3. SUBSAMPLING APPROXIMATION

We have seen in the previous section that the asymptotic null distribution of continuous functionals of R_n^1 depends in a complex way on the DGP and the specification under the null. Therefore, critical values for the test statistics can not be tabulated for general cases. In this section we overcome this problem with the assistance of the subsampling methodology. Resampling methods have been extensively used in the literature of quantile regression models, see, e.g., Hahn (1995), Horowitz (1998), Biliias, Chen and Ying (2000), Sakov and Bickel (2000) or He and Hu (2002). These articles consider *iid* sequences. When time series are involved the bootstrap approximation becomes more challenging. Subsampling is a powerful resampling scheme that allows an asymptotically valid inference under very general conditions on the DGP, see the monograph by Politis, Romano and Wolf (1999). Chernozhukov (2002) and Whang (2006) considered subsampling approximation for LQR model checks. In this section we apply the subsampling methodology to approximate the critical values of continuous functionals of R_n^1 . With an abuse of notation we write the test statistic as a function of the data $\{X_t = (Y_t, Z'_{t+1})' : t = 0, \pm 1, \pm 2, \dots\}$, $\Gamma(R_n^1) = \Gamma(R_n^1(X_1, \dots, X_n))$. Let $G_n^\Gamma(w)$ be the test statistic's cdf,

$$G_n^\Gamma(w) = P(\Gamma(R_n^1) \leq w).$$

We describe the subsampling approximation for the time series case; see the aforementioned references for *iid* sequences. Let $\Gamma(R_{b,i}^1) = \Gamma(R_b^1(X_i, \dots, X_{i+b-1}))$ be the test statistic computed with the subsample (X_i, \dots, X_{i+b-1}) of size b . We note that each subsample of size b (taken without replacement from the original data) is indeed a sample of size b from the true DGP. Hence, it is clear that one can approximate the sampling distribution $G_n^\Gamma(w)$ using the distribution of the values of $\Gamma(R_{b,i}^1)$ computed over the $n - b + 1$ different subsamples of size b (or the $\binom{n}{b}$ different subsamples of size b in the cross-section case). That is, we approximate $G_n^\Gamma(w)$ by

$$G_{n,b}^\Gamma(w) = \frac{1}{n - b + 1} \sum_{i=1}^{n-b+1} 1(\Gamma(R_{b,i}^1) \leq w), \quad w \in [0, \infty). \quad (9)$$

Let $c_{n,1-\tau,b}^\Gamma$ be the $(1-\tau)$ -th sample quantile of $G_{n,b}^\Gamma(w)$, i.e.,

$$c_{n,1-\tau,b}^\Gamma = \inf\{w : G_{n,b}^\Gamma(w) \geq 1 - \tau\}.$$

Thus, our subsampling tests reject the null hypothesis if $\Gamma(R_n^1) > c_{n,1-\tau,b}^\Gamma$. Let $c_{1-\tau}^\Gamma$ be the $(1-\tau)$ -th quantile of $G_\infty^\Gamma(w) = P(\Gamma(R_\infty^1) \leq w)$. To justify theoretically this resampling approximation we need an additional assumption on the serial dependence of the DGP. Define the α -mixing coefficients as

$$\alpha(m) = \sup_{n \in \mathbb{Z}} \sup_{B \in \mathcal{F}_n, A \in \mathcal{P}_{n+m}} |P(A \cap B) - P(A)P(B)|, \quad m \geq 1$$

where the σ -fields \mathcal{F}_n and \mathcal{P}_n are $\mathcal{F}_n := \sigma(X_t, t \leq n)$ and $\mathcal{P}_n := \sigma(X_t, t \geq n)$, respectively, with $X_t = (Y_t, Z'_{t+1})'$.

Assumption A6: $\{X_t = (Y_t, Z'_{t+1})' : t = 0, \pm 1, \pm 2, \dots\}$ is a strictly stationary strong mixing process with α -mixing coefficients satisfying

$$\sum_{m=1}^n \alpha(m) = o(n).$$

The mixing assumption in A6 is sufficient but not necessary for the validity of the subsampling, see Politis, Romano and Wolf (1999). This subsampling procedure allows us to approximate the asymptotic critical values of the tests based on $\Gamma(R_n^1)$. The next result justifies theoretically the subsampling approximation. Its proof follows closely that of Theorem 2 in Whang (2006).

THEOREM 5: *Assume Assumptions A1-A6 and that $b/n \rightarrow 0$ and $b \rightarrow \infty$ as $n \rightarrow \infty$. Then,*

(i) *Under the null hypothesis H_0 ,*

$$c_{n,1-\tau,b}^\Gamma \xrightarrow{P} c_{1-\tau}^\Gamma.$$

and

$$P(\Gamma(R_n^1) > c_{n,1-\tau,b}^\Gamma) \rightarrow \tau.$$

(ii) *Under any fixed alternative hypothesis*

$$P(\Gamma(R_n^1) > c_{n,1-\tau,b}^\Gamma) \rightarrow 1.$$

(iii) *Under the local alternatives (8),*

$$P(\Gamma(R_n^1) > c_{n,1-\tau,b}^\Gamma) \rightarrow P(\Gamma(R_\infty^1 + D_a) > c_{1-\tau}^\Gamma).$$

Theorem 5 implies that the proposed subsampling tests have a correct asymptotic level, are consistent and are able to detect alternatives tending to the null at the parametric rate $n^{-1/2}$. An appealing property of our subsampling tests is that they do not need estimation of the nonparametric (conditional) sparsity function $1/f_{I_{t-1}}(m_{t-1}(\theta_0))$, which results in a substantial simplification of the tests.

In practice, the empirical size and power of the tests depend on the choice of the parameter b . For this choice the reader is referred to Politis, Romano and Wolf (1999) or Sakov and Bickel (2000). In the present article, we follow the suggestion of Sakov and Bickel (2000) and we chose $b = \lfloor kn^{2/5} \rfloor$, where $\lfloor \cdot \rfloor$ denotes the integer part, which yields the optimal minimax accuracy under certain conditions. Section 5 below shows that this resampling procedure provides good approximations in finite samples for a variety of values for k .³

To implement the subsampling test we follow the suggestion of Chernozhukov and Fernández-Val (2005) and compute the subsampling critical value with recentered subsampling statistic $\Gamma(R_{b,i}^1) - \Gamma(b^{1/2}n^{-1/2}R_n^1)$. This centering leads to better power properties of the resulting subsampling test. An alternative possible centering is $\Gamma(R_{b,i}^1 - b^{1/2}n^{-1/2}R_n^1)$. The latter performs similarly to the former in our simulations, and hence it is omitted.

4. FINITE SAMPLE PERFORMANCE

We investigate in this section by means of Monte Carlo experiments the finite sample performance of the proposed test based on CvM_n and we compare it with some related tests. The aim is to provide evidence of the good finite-sample performance of the new test statistic.

We describe our simulation setup. Following Escanciano and Velasco (2006) and references therein, we choose $\Phi(\cdot)$ equal to the d -variate standard normal random vector⁴. In the first experiment, we consider as W a uniform discrete distribution over a grid of \mathcal{T} in $m = 20$ equidistributed points from $\epsilon = 0.1$ to $1 - \epsilon = 0.9$. Denote by $\mathcal{T}_m = \{\alpha_j\}_{j=1}^m$ the points in the grid, with $\epsilon = \alpha_1 < \dots < \alpha_m = 1 - \epsilon$. Let W_{exp} be the $n \times n$ matrix with elements $w_{\text{exp},t,s} = \exp(-\frac{1}{2} |I_{t-1} - I_{s-1}|^2)$ and let Ψ be the $n \times m$ matrix with elements $\psi_{ij} = \Psi_{\alpha_j}(Y_i - m(I_{i-1}, \theta_n))$. Hence, the CvM test statistic is computed as

$$CvM_n = \frac{1}{nm} \sum_{j=1}^m \psi'_{\cdot j} W_{\text{exp}} \psi_{\cdot j}, \quad (10)$$

where $\psi_{\cdot j}$ denotes the j column of Ψ . Therefore, the computation of CvM_n is straightforward. Our theory allows for $m \rightarrow \infty$ as $n \rightarrow \infty$ and the $\{\alpha_j\}_{j=1}^m$ generated independently from a distribution on \mathcal{T} , but for simplicity in the computations we have considered m fixed and $\{\alpha_j\}_{j=1}^m$ deterministic throughout this section.

We compare our omnibus test with other related tests. The first test is the following continuous

³Our experience suggests that values of k in the range $k = 4, 5$ and 6 provide good approximation for finite sample distributions.

⁴Strictly speaking our present theory does not allow to integrate in the whole \mathbb{R}^d in the CvM test, but our theory can be easily adapted, see e.g. Escanciano's (2006) Hilbert space approach, to allow for the present definition of the CvM test. In any case, there is no practical difference.

functional of $R_n^1(\cdot)$,

$$\begin{aligned} K_n &:= \int_{\Pi} |R_n^1(x, \alpha)|^2 d\delta_0(x) dW(\alpha) \\ &= \int_{\mathcal{T}} |R_n^1(0, \alpha)|^2 dW(\alpha), \end{aligned}$$

where δ_0 is the delta-Dirac measure at zero. For the present situation

$$K_n = \frac{1}{m} \sum_{j=1}^m K_{n, \alpha_j}^2, \quad (11)$$

where

$$K_{n, \alpha} := \frac{1}{\sqrt{n}} \sum_{t=1}^n \Psi_{\alpha}(Y_t - m(I_{t-1}, \theta_n)). \quad (12)$$

The test statistic K_n can be considered as an extension (to simultaneous α 's) of the traditional unconditional backtest proposed by Kupiec (1995), which is extensively used in the risk management literature; see Berkowitz, Christoffersen and Pelletier (2006) and Escanciano and Olmo (2010) for a review of this literature. Note that the asymptotic theory for K_n follows directly from our results by an application of the CMT.

Likewise, an extended version of Christoffersen's (1998) conditional or independence test can be computed as

$$C_n = \frac{1}{m} \sum_{j=1}^m C_{n, \alpha_j}^2, \quad (13)$$

where

$$C_{n, \alpha} = \frac{1}{\sqrt{n}} \sum_{t=2}^n \Psi_{\alpha}(Y_t - m(I_{t-1}, \theta_n)) \Psi_{\alpha}(Y_{t-1} - m(I_{t-2}, \theta_n)). \quad (14)$$

For further motivation of the test statistics K_n and C_n , see our application in Section 5.

We implement tests based on K_n and C_n using the subsampling approximation described in Section 3. The subsampling approximation is convenient as it accounts for estimation effects that affect the asymptotic distribution of K_n and C_n (cf. Escanciano and Olmo, 2009).

Motivated by our subsequent application, in the first experiment we entertain a Gaussian GARCH(1,1) model with AR(1) conditional mean for Y_t , leading to the quantile model

$$\begin{aligned} m(I_{t-1}, \theta_0(\alpha)) &= \mu_0 + \mu_1 Y_{t-1} + \sigma_t \Phi_{\varepsilon}^{-1}(\alpha), \\ \text{with } \sigma_t^2 &= \eta_{00} + \eta_{10}(Y_t - \mu_0 - \mu_1 Y_{t-1})^2 + \eta_{20} \sigma_{t-1}^2, \end{aligned} \quad (15)$$

where $\Phi_{\varepsilon}^{-1}(\alpha)$ is the α -quantile of the standard Gaussian error distribution and the parameters $\beta_0 = (\mu_0, \mu_1, \eta_{00}, \eta_{10}, \eta_{20})'$ are estimated by Quasi-Maximum Likelihood (QML). Here $\theta_0(\alpha) = \beta_0$, that is, we assume that the innovation's quantile is known and given by $\Phi_{\varepsilon}^{-1}(\alpha)$. The results with estimated innovation's quantiles are quantitatively the same as with known quantiles, and hence they are omitted. We consider the following data generating processes:

NULL: AR(1)-GARCH(1,1) model: $Y_t = 0.053 - 0.092Y_{t-1} + \varepsilon_t$,

ALT1: ARMA(1,1)-GARCH(1,1) model: $Y_t = 0.053 - 0.092Y_{t-1} + 0.7\varepsilon_{t-1} + \varepsilon_t$.

ALT2: TAR model: $Y_t = 0.6Y_{t-1} + \varepsilon_t$ if $Y_{t-1} \leq 1$ and $Y_t = -0.5Y_{t-1} + \varepsilon_t$ if $Y_{t-1} > 1$.

ALT3: SV model: $Y_t = 0.053 - 0.092Y_{t-1} + \exp(h_t)u_t$, $h_t^2 = 0.936h_{t-1}^2 + 0.32u_t$.

ALT4: Bilinear model (BIL): $Y_t = 0.6Y_{t-1} + 0.7u_{t-1}Y_{t-2} + u_t$.

ALT5: Non-Linear Moving Average model (NLMA): $Y_t = 0.8u_{t-1}^2 + u_t$.

ALT6: AR(1)-GARCH(1,1) model: $Y_t = 0.053 - 0.092Y_{t-1} + v_t$.

In models NULL and ALT1-ALT2, $\varepsilon_t = \sigma_t u_t$, with σ_t as in (15) with $(\eta_{00}, \eta_{10}, \eta_{20}) = (0.013, 0.104, 0.880)$ and $\{u_t\}$ iid $N(0, 1)$ variates. These parameter values are motivated from our application. Also, in ALT6 $v_t = \sigma_t w_t$ with σ_t as before and $\{w_t\}$ a standardized sequence of Student-t innovations with 5 degrees of freedom. Model NULL is frequently used in risk management, whereas ALT1-ALT6 are common linear and nonlinear models used in the time series literature.

We first consider two sample sizes $n = 100$ and $n = 300$ and a quantile interval $[0.1, 0.9]$. As the number of subsamples, we follow the suggestion of Sakov and Bickel (2000) and we chose $b = \lfloor kn^{2/5} \rfloor$, with several choices of k . We consider $k = 3, 4$ and 5 . These values correspond to $b = 18, 25$ and 31 for $n = 100$ and $b = 29, 39$ and 48 for $n = 300$, respectively. We set the number of Monte Carlo repetitions to 1,000 throughout the paper. Therefore, the maximal simulation standard error is $\max_{0 \leq p \leq 1} \sqrt{p(1-p)/1000} \approx 0.016$. In all experiments, the nominal probability of rejecting a correct null hypothesis is 0.05. The conclusions with other nominal values are similar.

In Table I we report the rejection frequencies of the tests based on CvM_n , K_n and C_n for models NULL and ALT1-ALT6. The empirical size performance is satisfactory for CvM_n and K_n , but for C_n there are some size distortions, with increasing overrejections as k increases. As for the empirical power, we observe that none of the test dominates the others uniformly in all alternatives. The test based on CvM_n has, however, an omnibus power performance, having the highest rejection frequencies for ALT2, ALT4 and ALT5, and performing satisfactorily for the rest of alternatives. In any of the alternatives CvM_n presents the lowest rejections. The tests based K_n and C_n have good power for some alternatives but also poor power for others, since they are directional in nature. The unconditional backtest has good power against ALT3 and ALT6, but low power against the rest of alternatives, whereas the conditional test C_n is good in detecting the linear alternative ALT1. For ALT6, since the resulting errors are *iid*, the test based on C_n is expected to have low power, which is consistent with our Monte Carlo results for this alternative. The rejection frequencies increase

as n increases, specially with the omnibus test based on CvM_n , showing consistency against these fixed alternatives.

Please insert Table I about here.

In unreported simulations, we observed that the CvM test statistic CvM_n had higher power than the Kolmogorov–Smirnov-type test KS_n in (6), at least for the alternatives considered. We also observed that uncentered tests led to less powerful procedures than centered test, in accordance with the theoretical results in Chernozhukov and Fernández-Val (2005). Finally, simulations using the indicator weight function $1(I_{t-1} \leq x)$, instead of $\exp(ix'I_{t-1})$, confirmed that exponential-based tests have higher power than indicator-based tests for the alternatives considered. Unreported simulations showing all these issues can be obtained from the authors upon request.

In the second experiment we focus on the left tail of the conditional distribution. We consider as W a uniform discrete distribution over a grid of \mathcal{T} in $m = 20$ equidistributed points from 0.05 to 0.2. The sample sizes are $n = 100$ and $n = 500$. Again, we consider $b = \lfloor kn^{2/5} \rfloor$, now with $k = 6, 7$ and 8, and 1,000 Monte Carlo repetitions.

In Table II we report the rejection frequencies of the tests based on CvM_n , K_n and C_n for models NULL, ALT2, ALT4, ALT6 and ALT8. The new ALT8 is an alternative where the model generated is NULL but the fitted model is an unconditional quantile, i.e. we specify $m(I_{t-1}, \theta_0(\alpha)) = \theta_0(\alpha) = F_Y^{-1}(\alpha)$, with $\theta_0(\alpha) = F_Y^{-1}(\alpha)$ estimated by the unconditional empirical α -quantile. The unconditional quantile model is very popular in the risk management literature and is known there as the Historical Simulation model.

The empirical size performance is similar to that of the previous simulations. The test statistics CvM_n and K_n have a satisfactory empirical size but C_n leads to overrejections when $n = 100$. However, these overrejections are corrected when $n = 500$. As for the empirical power, we observe that CvM_n outperforms the rest of tests for all alternatives but for ALT4, for which K_n performs the best. The conditional test C_n has low power against these alternatives, and in particular, no power against alternatives ALT4 and ALT6. For ALT8, since the resulting fitted hits satisfy the unconditional hypothesis $E[1(Y_t \leq F_Y^{-1}(\alpha))] = \alpha$, K_n is expected to have no power, which is consistent with our results for K_n against this alternative.

Please insert Table II about here.

This limited simulation study suggests that even with relative small sample sizes the subsampling tests exhibit fairly good size accuracy and power performance. The omnibus nature of the test statistic CvM_n is confirmed in our simulations. Extensions of well known tests in the risk management literature have been provided, and we have seen that they may perform well for certain alternatives

and not so well for others, so these more directional tests may well complement CvM_n in practical situations.

5. APPLICATION TO MARKET RISK MANAGEMENT

The quantification of market risk for derivative pricing, portfolio optimization and pricing risk purposes has generated a large amount of theoretical and practical work. One of the implications of the creation of the Basel Committee on Banking Supervision was the implementation of Value-at-Risk (VaR) as the standard tool for measuring market risk. In financial terms, VaR is the maximum loss on a trading portfolio for a period of time given a confidence level. In statistical terms, VaR is the conditional quantile of the conditional distribution of returns on the portfolio given agent's information set. Nowadays, VaR has become a standard risk measure due its universality, conceptual simplicity and easy computation and evaluation.

The evaluation of VaR measures has become of paramount importance in risk management. In fact, for banks with sufficiently highly developed risk management systems the implementation of VaR techniques was a priori the only restriction set by the Basel Accord (1996a) for computing capital reserves. Thus, in order to monitor and assess the accuracy and quality of the different VaR forecasts techniques the Basel Accord (1996a) and the Amendment of Basel Accord (1996b) developed a diagnostic testing procedure that was denominated *backtesting*. To explain formally what backtesting is, let us consider the following implication of (1),

$$E[\Psi_{\alpha,t}(\theta_0) \mid \tilde{I}_{t-1,\alpha}(\theta_0)] = 0, \text{ a.s. for some } \theta_0(\alpha) \in \Theta \text{ and some } \alpha \in (0,1), \quad (16)$$

where $\tilde{I}_{t-1,\alpha}(\theta_0) := (\Psi_{\alpha,t-1}(\theta_0), \Psi_{\alpha,t-2}(\theta_0), \dots)'$. The popularity of condition (16) is mostly due to the discrete character and ease of interpretation of the variables $\{H_{t,\alpha}(\theta_0)\}$, with $H_{t,\alpha}(\theta_0) = 1(Y_t \leq m(I_{t-1}, \theta_0(\alpha)))$, which are the so-called *hits* or *exceedances*. In the VaR literature, the satisfaction of condition (16) has been taken as the criteria for the out-of-sample evaluation of VaR forecasts, leading to the so-called unconditional backtesting (i.e. tests for $E[H_{t,\alpha}(\theta_0)] = \alpha$) and conditional tests of independence (i.e. tests for $\{H_{t,\alpha}(\theta_0)\}$ being *iid*).

The unconditional backtest is carried out with the so-called Kupiec-test statistic (cf. Kupiec, 1995), given in (12) for marginal quantiles. We have proposed in this paper an extension of this test to multiple quantiles as in (11). Similarly, we have generalized Christoffersen's (1998) conditional test statistic in (14) to multiple quantiles in (13).

Then, the alternative methodology proposed in this paper to the mentioned classical backtesting methods overcomes some of their important deficiencies. First, it is important to stress that tests based on R_n^1 , such as CvM_n , are expected to be more powerful than standard backtesting techniques. This is so because we incorporate more (possibly nonlinear) information in the test

statistic. In particular, the unconditional marginal backtest coincides with $R_n^1(0, \alpha)$, whereas we exploit a continuum number of x 's, thereby leading to a more powerful test. This is confirmed in the applications below. Second, by using only one quantile level, VaR only tells us the most we can lose if a tail event does not occur; if a tail event does occur, we can expect to lose more than the VaR, but the VaR itself gives us no indication of how much that might be. Therefore, two positions can have the same VaR and yet have very different risk exposures. This is the so-called *tail risk* problem in VaR. Our methodology solves this deficiency by taking a larger, possibly infinite, number of quantiles in the tail, thereby giving a more complete picture of the underlying risk exposure and leading to a better understanding of the fitting properties of the associated risk model.

Indeed, our methodology can be seen as a general framework to analyze market risk. For instance, there is now an important growing literature in finance, proposing the Conditional Expected Shortfall (CES) as an alternative to the VaR for measuring market risk in financial data. This measure of risk is defined as

$$\rho_{W_{t-1}, \alpha}(Y_t) = \alpha^{-1} \int_0^\alpha q_\nu(I_{t-1}) d\nu, \quad \alpha \in (0, 1). \quad (17)$$

Therefore, in modeling CES the interest is only in the range of quantiles $[0, \alpha]$ and not on the whole conditional distribution; see Escanciano and Mayoral (2008) for discussion of parametric CES models. Hence, the methods proposed in this section can be also seen as model specification tools of CES models.

In this section, we compare the new methodology with the standard backtesting techniques discussed previously. For simplicity in the arguments, we only consider in-sample comparisons. The extension to out-of-sample exercises poses no extra difficulties, and hence it is omitted. The data sets we consider are daily closed European stock indexes returns from the Frankfurt DAX Index (DAX) and the London FTSE-100 Index (FTSE) from 1 January 2003 to 9 June 2008, with a total of $n = 1417$ observations. We consider the returns of the indexes obtained as the log differences of the data.

We entertain a Gaussian AR(1)-GARCH(1,1) model as in (15), with parameters estimated by QML. We also considered other specifications, like pure Gaussian GARCH(1,1) and Student-t GARCH(1,1) models with degrees of freedom estimated by MLE, and we obtained similar conclusions. For the sake of exposition we omit these alternative specifications.

The Basel Accord (1996a) and the Amendment of Basel Accord (1996b) recommend to carry out backtesting procedures with quantile levels $\alpha = 0.01$ or $\alpha = 0.05$. Here, we take as \mathcal{T} a grid of $m = 10$ equidistributed points $\{\alpha_j\}_{j=1}^m$ from $\alpha_1 = 0.01$ to $\alpha_m = 0.10$, in intervals of length 0.01, covering the region recommended by Basel Accord (1996a). We apply our CvM test in (5) with $I_{t-1} = (Y_{t-1}, Y_{t-d})$ for $d = 1$ and 2, and denote the corresponding test statistic by $CvM_{n,d}$. We

compute $CvM_{n,d}$ following (10), and K_n and C_n following (11) and (13), respectively. In Table III we report the subsampling p -values for several choices of k in $b = \lfloor kn^{2/5} \rfloor$.

Please insert Table III about here.

We can draw several conclusions from the results of Table III. First, our results indicate that the AR(1)-GARCH(1,1) model with Gaussian innovations is not able to adequately fit the tails of these stock returns. The omnibus test based on $CvM_{n,d}$ strongly rejects this model for all stocks at reasonable significance levels. Second, the cumulative conditional backtest statistic C_n has rather low power and indeed, it is not able to detect any of these alternatives. This result is consistent with other finite sample studies using marginal (i.e. for fixed α) versions of this test, see Escanciano and Olmo (2010). Third, the unconditional test K_n also detects these alternatives, although the corresponding p -values are larger than those for $CvM_{n,d}$.

We complement the previous analysis with the marginal tests for each $\alpha_j, j = 1, \dots, m$, in Figures 1 and 2 for the subsampling size $b = \lfloor kn^{2/5} \rfloor$ with $k = 5$, i.e., $b = 91$. We take $d = 1$ in the information set for all stocks. We observe that the conditional marginal tests $C_{n,\alpha}$ are more sensitive to α than the other tests. We reject the correct specification hypothesis for the DAX index for most values of α with the omnibus test, but not with conditional and unconditional backtests. For a risk manager applying classical backtesting techniques at the recommended level $\alpha = 0.05$, the risk model provided by the AR(1)-GARCH(1,1) would seem appropriate. Using our more powerful test, he or she would conclude that this is not the case. In particular, this model fails to fit quantiles in the range $\alpha \in \{0.01, 0.02, 0.05\}$.

Please insert Figure 1 and Figure 2 about here.

Figure 2 reveals that the rejection for the FTSE index of our CvM test is due to the misspecification of the conditional quantiles at low levels $[0.01, 0.05]$. For this data set, the traditional unconditional backtests is very sensitive to α , giving very different answers if $\alpha = 0.05$ or $\alpha = 0.07$. Overall, we observe that marginal tests are in general sensitive to the choice of α , which provides motivation for our cumulative measures of fit when the whole tail is of interest.

For a better understanding of the cause of rejection, we report in Table IV the number of violations $Viol_\alpha = \sum_{t=1}^n H_{t,\alpha}(\theta_n)$ for each $\alpha_j, j = 1, \dots, m$, as well as the number of expected violations $EViol_\alpha = n \cdot \alpha$. We observe that in all cases with the DAX and FTSE indexes the number of violations is higher than its expected value, indicating fatter tails than the Gaussian AR-GARCH model, especially within the quantile region $[0.01, 0.05]$. Unreported simulations with a Student-t distribution showed that an AR(1)-GARCH(1,1) model with Student-t innovations is still not able to fit the tails of these data sets, although the number of violations is reduced considerably in all cases. We omit these additional simulations for the sake of space.

Please insert Table IV about here.

CONCLUSIONS

Econometric modeling often requires the specification of conditional quantile models for a range of quantiles of the conditional distribution. For the evaluation of models for quantile regression we propose and justify a general and flexible method which compares favorably with single quantile techniques and ad-hoc tests. We have shown in this paper that our test has higher power than the standard (marginal) unconditional and conditional backtesting commonly used by banks and regulators to assess dynamic parametric VaR estimates. In particular, we find that the standard conditional backtesting procedure has rather low power in detecting misspecifications of an AR(1)-GARCH(1,1) VaR model for two major European stock indexes. Our methods provide flexible and powerful tools that can be used by practitioners to assess the plausibility of standard market risk models.

APPENDIX. PROOFS

First, we shall state a weak convergence theorem which is an extension of Theorem A1 in Delgado and Escanciano (2007) and that is of independent interest. Let for each $n \geq 1$, $I'_{n,0}, \dots, I'_{n,n-1}$, be an array of random vectors in \mathbb{R}^p , $p \in \mathbb{N}$, and $Y_{n,1}, \dots, Y_{n,n}$, be an array of real random variables (r.v.'s). Denote by $(\Omega_n, \mathcal{A}_n, P_n)$, $n \geq 1$, the probability space in which all the r.v.'s $\{Y_{n,t}, I'_{n,t-1}\}_{t=1}^n$ are defined. Let $\mathcal{F}_{n,t}$, $0 \leq t \leq n$, be a double array of sub σ -fields of \mathcal{A}_n such that $\mathcal{F}_{n,t-1} \subset \mathcal{F}_{n,t}$, $t = 1, \dots, n$ and such that for each $n \geq 1$ and each $\gamma \in \mathcal{H}$,

$$E[w(Y_{n,t}, I_{n,t-1}, \gamma) \mid \mathcal{F}_{n,t-1}] = 0 \text{ a.s.}, \quad 1 \leq t \leq n, \forall n \geq 1. \quad (18)$$

Moreover, we shall assume that $\{w(Y_{n,t}, I_{n,t-1}, \gamma), \mathcal{F}_{n,t}, 0 \leq t \leq n\}$ is a square-integrable martingale difference sequence for each $\gamma \in \mathcal{H}$, that is, (18) holds, $Ew^2(Y_{n,t}, I_{n,t-1}, \gamma) < \infty$ and $w(Y_{n,t}, I_{n,t-1}, \gamma)$ is $\mathcal{F}_{n,t}$ -measurable for each $\gamma \in \mathcal{H}$ and $\forall t, 1 \leq t \leq n, \forall n \in \mathbb{N}$. The following result gives sufficient conditions for the weak convergence of the empirical process

$$\alpha_{n,w}(\gamma) = n^{-1/2} \sum_{t=1}^n w(Y_{n,t}, I_{n,t-1}, \gamma), \quad \gamma \in \mathcal{H}.$$

Under mild conditions the empirical process $\alpha_{n,w}$ can be viewed as a mapping from Ω_n to $\ell^\infty(\mathcal{H})$, the space of all complex-valued functions that are uniformly bounded on \mathcal{H} , with \mathcal{H} a generic metric space. The weak convergence theorem that we present here is founded on results by Levental (1989), Bae and Levental (1995) and Nishiyama (2000). In Theorem A1 in Delgado and Escanciano (2007)

\mathcal{H} was finite-dimensional, but here we allow for an infinite-dimensional \mathcal{H} . The proof of theorem does not change by this possibility, however.

An important role in the weak convergence theorem is played by the conditional quadratic variation (CV) of the empirical process $\alpha_{n,w}$ on a finite partition $\mathcal{B} = \{H_k; 1 \leq k \leq N\}$ of \mathcal{H} , which is defined as

$$CV_{n,w}(\mathcal{B}) = \max_{1 \leq k \leq N} n^{-1} \sum_{t=1}^n E \left[\sup_{\gamma_1, \gamma_2 \in H_k} |w(Y_{n,t}, I_{n,t-1}, \gamma_1) - w(Y_{n,t}, I_{n,t-1}, \gamma_2)|^2 \mid \mathcal{F}_{n,t-1} \right]. \quad (19)$$

Then, for the weak convergence theorem we need the following assumptions.

W1: For each $n \geq 1$, $\{(Y_{n,t}, I_{n,t-1})' : 1 \leq t \leq n\}$ is a strictly stationary and ergodic process.

The sequence $\{w(Y_{n,t}, I_{n,t-1}, \gamma), \mathcal{F}_{n,t}, 1 \leq t \leq n\}$ is a square-integrable martingale difference sequence for each $\gamma \in \mathcal{H}$. Also, there exists a function $C_w(\gamma_1, \gamma_2)$ on $\mathcal{H} \times \mathcal{H}$ to \mathbb{R} such that uniformly in $(\gamma_1, \gamma_2) \in \mathcal{H} \times \mathcal{H}$

$$n^{-1} \sum_{t=1}^n w(Y_{n,t}, I_{n,t-1}, \gamma_1) w^c(Y_{n,t}, I_{n,t-1}, \gamma_2) = C_w(\gamma_1, \gamma_2) + o_{P_n}(1).$$

W2: The family $w(Y_{n,t}, I_{n,t-1}, \gamma)$ is such that $\alpha_{n,w}$ is a mapping from Ω_n to $\ell^\infty(\mathcal{H})$ and for every $\delta > 0$ there exists a finite partition $\mathcal{B}_\delta = \{H_k; 1 \leq k \leq N_\delta\}$ of \mathcal{H} , with N_δ being the number of elements of such partition, such that

$$\int_0^\infty \sqrt{\log(N_\delta)} d\delta < \infty \quad (20)$$

and

$$\sup_{\delta \in (0,1) \cap \mathbb{Q}} \frac{CV_{n,w}(\mathcal{B}_\delta)}{\delta^2} = O_{P_n}(1). \quad (21)$$

Let $\alpha_{\infty,w}(\cdot)$ be a Gaussian process with zero mean and covariance function given by $C_w(\gamma_1, \gamma_2)$. We are now in position to state the following

THEOREM A1: *If Assumptions W1 and W2 hold, then it follows that*

$$\alpha_{n,w} \implies \alpha_{\infty,w} \text{ in } \ell^\infty(\mathcal{H}).$$

PROOF OF THEOREM A1: Theorem A1 in Delgado and Escanciano (2007).

COROLLARY A1: *Assuming that W1 holds for $w(Y_{n,t}, I_{n,t-1}, v) = \Psi_\alpha(Y_{n,t-m}(I_{n,t-1}, \theta_0(\alpha))) \exp(ix' I_{n,t-1})$, $v = (x', \alpha)' \in \Pi$, A1(b) and that*

$$n^{-1} \sum_{t=1}^n |I_{n,t-1}|^2 = O_{P_n}(1),$$

then the weak convergence of Theorem A1 holds.

PROOF OF COROLLARY A1: We shall apply Theorem A1. Let us define the metric

$$d(v_1, v_2) := \sqrt{|\alpha_1 - \alpha_2|^2 + |x_1 - x_2|^2}, \quad v_1, v_2 \in \Pi.$$

Then, we define an δ -bracket as an interval $[v_1, v_2]$ such that $v_1 \leq v_2$ and $d(v_1, v_2) \leq \delta$. The bracketing number $N(\delta, \Pi, d)$ is the minimum number of δ -brackets needed to cover Π . Then, it is easy to show that

$$\int_0^\infty \sqrt{\log(N(\delta, \Pi, d))} d\delta < \infty$$

holds. It remains to show that (21) holds. Consider a partition $\mathcal{B}_\delta = \{H_k; 1 \leq k \leq N(\delta, \Pi, d) \equiv N_\delta\}$ of Π in δ -brackets $H_k = [v_k, \bar{v}_k]$, with $v_k = (\underline{x}'_k, \underline{\alpha}_k)'$ and $\bar{v}_k = (\bar{x}'_k, \bar{\alpha}_k)'$, $\underline{x}_k \leq \bar{x}_k$ and $\underline{\alpha}_k \leq \bar{\alpha}_k$. Define $\varepsilon_{n,t}(\alpha) = Y_{n,t} - m(I_{n,t-1}, \theta_0(\alpha))$. Then, by simple algebra and the monotonicity of $1(\varepsilon_{n,t}(\alpha) \leq 0)$ due to A1(b), $CV_{n,w}(\mathcal{B}_\delta)$ in (19) is bounded by

$$\begin{aligned} & 2 \max_{1 \leq k \leq N_\varepsilon} n^{-1} \sum_{t=1}^n E \left[\sup_{v_1, v_2 \in H_k} |1(\varepsilon_{n,t}(\alpha_1) \leq 0) - \alpha_1 - 1(\varepsilon_{n,t}(\alpha_2) \leq 0) + \alpha_2|^2 \mid \mathcal{F}_{n,t-1} \right] \\ & + 2 \max_{1 \leq k \leq N_\varepsilon} n^{-1} \sum_{t=1}^n \left[\sup_{v_1, v_2 \in H_k} |\exp(ix'_1 I_{n,t-1}) - \exp(ix'_2 I_{n,t-1})|^2 \right] \\ & \leq C \max_{1 \leq k \leq N_\varepsilon} \left\{ |\underline{\alpha}_k - \bar{\alpha}_k| + |\underline{x}_k - \bar{x}_k|^2 n^{-1} \sum_{t=1}^n |I_{n,t-1}|^2 \right\}. \end{aligned}$$

Hence, (21) holds for the partition \mathcal{B}_δ . Therefore, W2 of Theorem A1 holds and the corollary is proved. \square

PROOF OF THEOREM 1. Follows from Corollary A1. \square

THEOREM A2. Assume Assumptions A1(c-d), A2, A3, and that there exists a $\theta_1 \in \mathcal{B}$ such that $\|\theta_n - \theta_1\|_{\mathcal{B}} = o_P(1)$. Then, uniformly in $(x', \alpha)' \in \Pi$,

$$\begin{aligned} R_n^1(x, \alpha) &= \frac{1}{\sqrt{n}} \sum_{t=1}^n \{\Psi_{\alpha,t}(\theta_1) - E[\Psi_{\alpha,t}(\theta_1) \mid \mathcal{F}_{t-1}]\} \exp(ix' I_{t-1}) \\ &+ \frac{1}{\sqrt{n}} \sum_{t=1}^n \{E[\Psi_{\alpha,t}(\theta) \mid \mathcal{F}_{t-1}]_{\theta=\theta_n} - E[\Psi_{\alpha,t}(\theta_1) \mid \mathcal{F}_{t-1}]\} \exp(ix' I_{t-1}) \\ &+ \frac{1}{\sqrt{n}} \sum_{t=1}^n E[\Psi_{\alpha,t}(\theta_1) \mid \mathcal{F}_{t-1}] \exp(ix' I_{t-1}) - E[E[\Psi_{\alpha,t}(\theta_1) \mid \mathcal{F}_{t-1}] \exp(ix' I_{t-1})] \\ &+ \sqrt{n} E[E[\Psi_{\alpha,t}(\theta_1) \mid \mathcal{F}_{t-1}] \exp(ix' I_{t-1})] + o_P(1). \end{aligned} \tag{22}$$

PROOF OF THEOREM A2: Write $w_{t-1}(v, \theta) := \{\Psi_{\alpha,t}(\theta) - E[\Psi_{\alpha,t}(\theta) \mid \mathcal{F}_{t-1}]\} \exp(ix' I_{t-1})$. First we shall show that the process

$$S_n(v, \theta) = \frac{1}{\sqrt{n}} \sum_{t=1}^n w_{t-1}(v, \theta)$$

is asymptotically tight with respect to $(v, \theta) \in \mathcal{W} = \Pi \times \mathcal{B}$.

Let us define the class $\mathcal{K} = \{w(v, \theta) : (v, \theta) \in \mathcal{W}\}$. Denote $X_{t-1, \infty} = (I_{t-1}, I_{t-2}, \dots)'$. Let $\mathcal{B}_\delta = \{B_k; 1 \leq k \leq N_\delta \equiv N_{\square}(\delta, \mathcal{K}, \|\cdot\|_2)\}$, with $B_k = [\underline{w}_k(Y_t, X_{t-1, \infty}), \bar{w}_k(Y_t, X_{t-1, \infty})]$, be a partition of \mathcal{K} in δ -brackets with respect to $\|\cdot\|_2$, where $\|\cdot\|_2$ denotes the L_2 norm of random variables, i.e., $\|X\|_2 = (E[X^2])^{1/2}$.

Conditions A1(c-d) and A2 imply that for a sufficiently small $\delta > 0$,

$$\begin{aligned} & \left\| \sup_{\substack{(v_2, \theta_2) \in \mathcal{W}: d(v_1, v_2) \leq \delta \\ \|\theta_1 - \theta_2\|_{\mathcal{B}} \leq \delta}} |w_{t-1}(v_1, \theta_1) - w_{t-1}(v_2, \theta_2)| \right\|_2 \\ & \leq C \left\| \sup_{\substack{(v_2, \theta_2) \in \mathcal{W}: d(v_1, v_2) \leq \delta \\ \|\theta_1 - \theta_2\|_{\mathcal{B}} \leq \delta}} |\Psi_{\alpha_1, t}(\theta_1) - \Psi_{\alpha_2, t}(\theta_2)| \right\|_2 + C\delta \\ & \leq C \left\| \sup_{|\alpha_1 - \alpha_2| \leq \delta} |1(Y_t \leq m_{t-1}(\theta_1(\alpha_1))) - 1(Y_t \leq m_{t-1}(\theta_1(\alpha_2)))| \right\|_2 \\ & \quad + C \left(E \left[\sup_{\|\theta_1 - \theta_2\|_{\mathcal{B}} \leq \delta} |1(Y_t \leq m_{t-1}(\theta_1(\alpha))) - 1(Y_t \leq m_{t-1}(\theta_2(\alpha)))| \right] \right)^{1/2} + C\delta \\ & \leq C\delta^{1/2}. \end{aligned}$$

Theorem 3 in Chen et al. (2003) and A3 yield that (20) holds for such partition. Therefore, by similar arguments as in Corollary A1, (21) follows, and condition W2 of Theorem A1 holds. The asymptotically tightness of $S_n(v, \theta)$ is then proved. As a result,

$$\sup_{v \in \Pi} |S_n(v, \theta_n) - S_n(v, \theta_1)| = o_P(1),$$

which can be rewritten as

$$\begin{aligned} R_n^1(\cdot) &= \frac{1}{\sqrt{n}} \sum_{t=1}^n \{\Psi_{\alpha, t}(\theta_1) - E[\Psi_{\alpha, t}(\theta_1) | \mathcal{F}_{t-1}]\} \exp(ix' I_{t-1}) \\ & \quad + \frac{1}{\sqrt{n}} \sum_{t=1}^n E[\Psi_{\alpha, t}(\theta) | \mathcal{F}_{t-1}]_{\theta=\theta_n} + o_P(1), \end{aligned}$$

from which (22) follows. \square

PROOF OF THEOREM 2: Under the null $\theta_1 = \theta_0$ and $E[\Psi_{\alpha}(e_t(\theta_0)) | \mathcal{F}_{t-1}] = 0$ a.s.. From the expansion in (22), it follows that, uniformly in $v \in \Pi$,

$$\begin{aligned} R_n^1(\cdot) &= \frac{1}{\sqrt{n}} \sum_{t=1}^n \Psi_{\alpha, t}(\theta_0) \exp(ix' I_{t-1}) \\ & \quad + \frac{1}{\sqrt{n}} \sum_{t=1}^n \{E[\Psi_{\alpha, t}(\theta) | \mathcal{F}_{t-1}]_{\theta=\theta_n} - E[\Psi_{\alpha, t}(\theta_0) | \mathcal{F}_{t-1}]\} \exp(ix' I_{t-1}) + o_P(1) \\ &= R_n(\cdot) + \frac{1}{\sqrt{n}} \sum_{t=1}^n \{F_{I_{t-1}}(m(I_{t-1}, \theta_n)) - F_{I_{t-1}}(m_{t-1}(\theta_0))\} \exp(ix' I_{t-1}) + o_P(1). \end{aligned}$$

Now, from A1(d) and Koul and Stute (1999, pp. 228-229), uniformly in $v \in \Pi$,

$$\begin{aligned} & \frac{1}{\sqrt{n}} \sum_{t=1}^n \{F_{I_{t-1}}(m(I_{t-1}, \theta_n)) - F_{I_{t-1}}(m_{t-1}(\theta_0))\} \exp(ix' I_{t-1}) \\ &= \sqrt{n}(\theta_n - \theta_0) \frac{1}{n} \sum_{t=1}^n g(I_{t-1}, \theta_0) f_{I_{t-1}}(m_{t-1}(\theta_0)) \exp(ix' I_{t-1}) + o_P(1). \end{aligned}$$

This together with Theorem 1, A2(c) and A4 proves the theorem. \square

PROOF OF THEOREM 3: Let $\mathcal{W} = \Pi \times \mathcal{B}$. Let $w = (x', \alpha, \theta'(\cdot))'$ be a general element of \mathcal{W} . The space \mathcal{W} is endowed with the metric

$$\rho(w_1, w_2) = |x_1 - x_2| + |\alpha_1 - \alpha_2| + \sup_{\alpha \in T} |\theta_1(\alpha) - \theta_2(\alpha)|,$$

where $w_1 = (x'_1, \alpha_1, \theta'_1(\cdot))'$ and $w_2 = (x'_2, \alpha_2, \theta'_2(\cdot))'$ belong to \mathcal{W} . Let $B(w, \delta)$ be the open ball of radius δ around w , i.e., $B(w, \delta) = \{w_1 \in \mathcal{W} : \rho(w_1, w) < \delta\}$. Note that A1-A3 yield that for each $w = (x', \alpha, \theta'(\cdot))' \in \mathcal{W}$ it holds that

$$\lim_{\delta \rightarrow 0} E \left[\sup_{w_1 \in B(w, \delta)} |\Psi_{\alpha_1, t}(\theta_1(\alpha_1)) \exp(ix'_1 I_{t-1}) - \Psi_{\alpha, t}(\theta(\alpha)) \exp(ix' I_{t-1})|^2 \right] = 0.$$

Therefore, $E[\Psi_{\alpha, t}(\theta_1(\alpha)) \exp(ix' I_{t-1})]$ is a continuous function of $v = (x', \alpha)'$. Therefore, a uniform version of the Ergodic Theorem

$$\sup_{\theta \in \mathcal{B}} \sup_{v \in \Pi} \left| \frac{1}{n} \sum_{t=1}^n [\Psi_{\alpha, t}(\theta(\alpha)) \exp(ix' I_{t-1}) - E[\Psi_{\alpha, t}(\theta(\alpha)) \exp(ix' I_{t-1})]] \right| = o_P(1).$$

Hence, from the last display and A5

$$\sup_{v \in \Pi} \left| \frac{1}{n} \sum_{t=1}^n [\Psi_{\alpha, t}(\theta_n(\alpha)) \exp(ix' I_{t-1}) - E[\Psi_{\alpha, t}(\theta_1(\alpha)) \exp(ix' I_{t-1})]] \right| = o_P(1).$$

and the function $E[\Psi_{\alpha, t}(\theta_1(\alpha)) 1(I_{t-1} \leq \cdot)]$ is different from zero in a subset with positive Lebesgue measure on Π . \square

PROOF OF THEOREM 4: The proof follows from Theorem A2 and Assumptions A5 and A6 jointly with A4' in a routine fashion, and then, it is omitted. \square

PROOF OF THEOREM 5. The proof follows the same steps as Theorems 2, 3 and 4 of Whang (2006), using our Corollary 1 instead of his Lemma 3, and then, it is omitted. \square

REFERENCES

- Andrews, D. W. K., 1997. A conditional Kolmogorov test. *Econometrica* 65, 1097-1128.
- Albrecht, J., van Vuuren, A. and Vroman, S., 2007. Counterfactual distributions with sample selection adjustments: econometric theory and an application to the Netherlands. Working Papers guecon-wpa 07-07-06, Georgetown University, Department of Economics.
- Angrist, J. Chernozhukov, V. and Fernández-Val, I. 2006. Quantile regression under misspecification, with an application to the U.S. wage structure. *Econometrica* 74, 539-563.
- Bae, J. and Levental, S., 1995. Uniform CLT for Markov chains and its invariance principle: a martingale approach. *Journal of Theoretical Probability* 8, 549-570.
- Basel Committee on Banking Supervision, 1996a. Amendment to the capital accord to incorporate market risks. Bank for International Settlements.
- Basel Committee on Banking Supervision, 1996b. Supervisory framework for the use of backtesting in conjunction with the internal models approach to market risk capital requirements. Bank for International Settlements.
- Bassett, G. and Chen, H.L., 2001. Portfolio style: *return*-based attribution using regression quantiles. *Empirical Economics* 26, 293-305.
- Bassett, Jr., G., and Koenker, R., 1978. Asymptotic theory of least absolute error regression. *Journal of the American Statistical Association* 73, 618-622.
- Berkowitz, J., Christoffersen, P., and Pelletier, D. 2006. Evaluating Value-at-Risk models with desk-level data. Working Paper Series, 010 Department of Economics, North Carolina State University.
- Bierens, H.J., 1982. Consistent model specification tests. *Journal of Econometrics* 20, 105-134.
- Bierens, H., and Ginther, D.K., 2001. Integrated conditional moment testing of quantile regression models. *Empirical Economics* 26, 307-324.
- Bilias, Y., Chen, S. and Ying, Z., 2000. Simple resampling methods for the censored regression quantiles. *Journal of Econometrics* 99, 373-386.
- Billingsley, P., 1961. The Lindeberg-Levy theorem for martingales. *Proceedings of the American Mathematical Society* 12, 788-792.
- Chen, X., O. Linton, and I. van Keilegom, 2003. Estimation of semiparametric models when the criterion function is not smooth. *Econometrica* 71, 1591-1608.
- Chernozhukov, V., 2002. Inference on the quantile regression process, an alternative. Working article 02-12 February, MIT www.ssrn.com.
- Chernozhukov, V. and Fernandez-Val, I., 2005. Subsampling inference on quantile regression processes.

- Sankhya 67, 253-276.
- Christoffersen, P., 1998. Evaluating interval forecasts. *International Economic Review* 39, 841-862.
- Corradi, V. and N.R. Swanson 2006. Predictive density and conditional confidence intervals accuracy tests. *Journal of Econometrics* 135, 187-228.
- Delgado, M.A. and Escanciano, J.C. 2007. Nonparametric tests for conditional symmetry in dynamic models. *Journal of Econometrics* 127, 652-682.
- Dudley, R.M. 1999. *Uniform Central Limit Theorems*. Cambridge. University Press, Massachusetts.
- Escanciano, J. C. 2006. Goodness-of-fit tests for linear and non-linear time series models. *Journal of the American Statistical Association* 101, 531-541.
- Escanciano, J. C. 2009. On the Lack of Power of Omnibus Specification Tests. *Econometric Theory* 25, 162-194.
- Escanciano, J. C. and Mayoral, S. 2008. Semiparametric estimation of dynamic conditional expected shortfall models. *The International Journal of Monetary Economics and Finance* 1, 106-120.
- Escanciano, J. C. and Olmo, J. 2010. Backtesting VaR with estimation risk. *Journal of Business and Economics Statistics* 28, 36-51.
- Escanciano, J.C. and Velasco, C. 2006. Generalized spectral tests for the martingale difference hypothesis. *Journal of Econometrics* 134, 151-185.
- Gutenbrunner, C. and Jurečková, J. 1992. Regression rank scores and regression quantiles. *The Annals of Statistics* 20, 305-329.
- Gutenbrunner, C., Jurečková, J., Koenker, R. and Portnoy, S. 1993. Test of linear hypotheses based on regression rank scores. *Journal of Nonparametric Statistics* 2, 307-331.
- Hahn, J. 1995. Bootstrapping quantile regression estimators. *Econometric Theory* 11, 105-121.
- He, X., and Hu, F. 2002. Markov chain marginal bootstrap. *Journal of the American Statistical Association* 97, 783-795.
- He, X. and Zhu, L.X 2003. A Lack-of-Fit test for quantile regression. *Journal of the American Statistical Association* 98, 1013-1022.
- Horowitz, J.L. 1998. Bootstrap methods for median regression models. *Econometrica* 66, 1327-1351.
- Horowitz, J.L. and Spokoiny, V.G. 2002. An adaptive, rate-optimal test of linearity for median regression models. *Journal of the American Statistical Association* 97, 822-835.
- Jurečková, J. and Hallin, M. 1999. Optimal tests for autoregressive models based on autoregression rank scores. *The Annals of Statistics* 27, 1385-1414.
- Jurečková, J. and Prochazka, B. 1994. Regression quantiles and trimmed least squares estimation in

- nonlinear regression models. *Journal of Nonparametric Statistics* 3, 201–222.
- Khmaladze, E.V. 1981. Martingale approach to the goodness of fit tests. *Theory of Probability and Its Applications* 26, 246-265.
- Kim, T.H. and White, H. 2003. Estimation, inference, and specification testing for possibly misspecified quantile regressions. in T. Fomby and R.C. Hill, eds., *Maximum Likelihood Estimation of Misspecified Models: Twenty Years Later*. New York: Elsevier, 107-132.
- Koenker, R. and Bassett, G. 1978. Regression quantiles. *Econometrica* 46, 33-50.
- Koenker, R. and Machado, J.A. 1999. Goodness of fit and related inference processes for quantile regression. *Journal of the American Statistical Association* 94, 1296-1310.
- Koenker, R. and Park, B.J. 1996. An interior point algorithm for nonlinear quantile regression. *Journal of Econometrics* 71, 265-285.
- Koenker, R. and Xiao, Z. 2002. Inference on the quantile regression process. *Econometrica* 70, 1583–1612.
- Koenker, R. and Xiao, Z. 2006. Quantile autoregression. *Journal of the American Statistical Association* 101, 980-990.
- Komunjer, I. 2005. Quasi-maximum likelihood estimation for conditional quantiles. *Journal of Econometrics* 128, 137-164.
- Koul, H. L. 2002. *Weighted empirical processes in dynamic nonlinear models*. 2nd ed, *Lecture Notes in Statistics*, Vol. 166, Springer.
- Koul, H.L., and Saleh, A.K.M.E 1994. Autoregression quantile and related rank scores processes. *The Annals of Statistics* 23, 670-689.
- Koul, H.L., and Stute W. 1999. Nonparametric model checks for time series. *The Annals of Statistics* 27, 204-236.
- Kupiec, P. 1995. Techniques for verifying the accuracy of risk measurement models. *Journal of Derivatives* 3, 73-84.
- Levental, S. 1989. A uniform CLT for uniformly bounded families of martingale differences. *Journal of Theoretical Probability* 2, 271-287.
- Machado, J.A.F. and Mata, J. 2005. Counterfactual decomposition of changes in wage distributions using quantile regression. *Journal of Applied Econometrics* 20, 445-65.
- Mukherjee, K. 1999. Asymptotics of quantiles and rank scores in nonlinear time series. *Journal of Time Series Analysis* 20, 173-192.
- Nishiyama, Y. 2000. Weak convergence of some classes of martingales with jumps. *The Annals of Probability* 28, 685-712.

- Pollard, D. 1991. Asymptotics for least absolute deviation regression estimators. *Econometric Theory* 7, 186-199.
- Politis, D. N., Romano, J.P. and Wolf, M. 1999. *Subsampling*. Springer-Verlag, New York.
- Portnoy, S. 1984. Tightness of the sequence of empiric cdf processes defined from regression fractiles. In *Robust and Nonlinear Time Series Analysis*, eds. J. Franke, W. Hardle, and D. Martin, Springer-Verlag, New York.
- Sakov, A. and Bickel, P. 2000. An Edgeworth expansion for the m out of n bootstrap median. *Statistical and Probability Letters* 49, 217-223.
- van der Vaart, A.W., and Wellner, J.A. 1996. *Weak convergence and empirical processes*. Springer. New York.
- Whang, Y-J. 2006. Consistent specification tests for quantile regression models. in *Frontiers of Analysis and Applied Research: Essays in Honor of Peter C.B. Phillips*, Cambridge University Press.
- Whang, Y-J. 2005. Smoothed Empirical Likelihood Methods for Quantile Regression Models. *Econometric Theory* 22, 173-205.
- Zheng, J.X. 1998. A consistent nonparametric test of parametric regression models under conditional quantile restrictions. *Econometric Theory* 14, 123-138.

TABLE I: Empirical size and power Rejection Frequencies).

n	k	Test	NULL	ALT1	ALT2	ALT3	ALT4	ALT5	ALT6
100	3	CvM_n	3.6	23.9	29.9	44.1	32.0	74.6	18.1
		K_n	3.4	2.8	5.0	62.5	7.7	24.1	22.9
		C_n	9.4	27.4	21.6	25.5	9.3	46.2	12.1
	4	CvM_n	3.7	25.9	29.4	40.6	35.2	73.4	20.7
		K_n	3.7	3.0	4.9	59.1	8.2	20.4	26.3
		C_n	10.6	30.8	24.9	24.8	10.7	46.3	13.4
	5	CvM_n	7.1	32.0	35.0	42.3	43.4	76.8	23.2
		K_n	5.6	4.3	6.1	59.7	9.2	22.0	28.1
		C_n	13.6	37.9	29.0	26.9	11.7	51.9	15.0
300	3	CvM_n	6.3	69.5	84.1	60.9	85.4	100.0	57.0
		K_n	5.6	3.3	20.5	87.3	4.7	81.2	72.1
		C_n	9.3	63.5	50.2	28.5	11.0	89.4	12.8
	4	CvM_n	7.7	70.3	82.0	55.2	85.6	100.0	60.0
		K_n	7.3	4.7	20.5	81.8	5.7	79.7	74.5
		C_n	10.1	66.0	52.7	27.4	11.5	89.1	14.4
	5	CvM_n	7.8	71.7	81.8	51.2	85.9	99.8	61.8
		K_n	8.1	5.9	21.4	78.7	5.9	78.3	75.1
		C_n	10.9	57.1	53.8	26.6	11.3	90.5	15.2

5% of significance level. $\alpha \in [0.1, 0.9]$. $b = \lfloor kn^{2/5} \rfloor$. Maximal simulation s.e.=1.6%

TABLE II: Empirical size and power Rejection Frequencies).

n	k	Test	NULL	ALT2	ALT4	ALT6	ALT8
100	6	CvM_n	2.3	30.1	13.8	9.8	46.1
		K_n	2.6	7.8	7.8	12.6	0.0
		C_n	10.2	19.2	7.1	6.4	25.5
	7	CvM_n	3.6	35.7	17.7	11.7	54.4
		K_n	3.5	9.4	10.0	14.0	0.0
		C_n	12.3	22.4	7.8	6.6	26.4
	8	CvM_n	5.9	42.9	24.3	17.8	66.8
		K_n	5.7	12.9	12.1	19.5	0.0
		C_n	15.2	27.1	10.2	10.0	32.5
500	6	CvM_n	5.0	89.9	37.4	58.5	65.8
		K_n	5.2	29.4	8.5	82.9	0.0
		C_n	5.7	36.5	6.3	7.1	29.3
	7	CvM_n	5.6	91.5	40.7	60.5	66.9
		K_n	6.4	29.8	9.8	83.0	0.0
		C_n	6.3	38.9	6.9	7.8	27.4
	8	CvM_n	6.7	93.2	44.6	62.3	68.3
		K_n	8.6	30.0	11.7	83.7	0.0
		C_n	6.8	41.0	7.4	8.5	29.4

5% of significance level. $\alpha \in [0.05, 0.2]$. $b = \lfloor kn^{2/5} \rfloor$. Maximal simulation s.e.=1.6%

TABLE III: Aggregated tests: Subsampling p -values.

n=1471	k	$CvM_{n,1}$	$CvM_{n,2}$	K_n	C_n
DAX	4	0.012	0.026	0.092	0.468
	5	0.043	0.048	0.087	0.478
	6	0.042	0.068	0.096	0.500
FTSE	4	0.000	0.003	0.009	0.976
	5	0.000	0.000	0.003	0.994
	6	0.001	0.000	0.012	0.990

ACCEPTED MANUSCRIPT

TABLE IV: Number of violations ($Viol_\alpha$) and expected violations ($EViol_\alpha$)

α_j	$EViol_\alpha$	DAX		FTSE	
		$Viol_\alpha$	$Viol_\alpha$	$Viol_\alpha$	$Viol_\alpha$
0.01	14.17	25	29		
0.02	28.34	43	48		
0.03	42.51	52	62		
0.04	56.68	59	69		
0.05	70.85	84	82		
0.06	85.02	97	89		
0.07	99.19	109	100		
0.08	113.36	126	115		
0.09	127.53	142	129		
0.10	141.70	158	145		




Figure 1. Subsampling p-values for $CvM_{n,1,\alpha}$ test (solid line), unconditional backtest $K_{n,\alpha}$ (dashed line), and the conditional backtest $C_{n,\alpha}$ (dotted line) as a function of α . Subsample size $b = 91$.

ACCEPTED MANUSCRIPT




Figure 2. Subsampling p-values for $CvM_{n,1,\alpha}$ test (solid line), unconditional backtest $K_{n,\alpha}$ (dashed line), and the conditional backtest $C_{n,\alpha}$ (dotted line) as a function of α . Subsample size $b = 91$.

ACCEPTED MANUSCRIPT