



# Coastal Uplift along the Eastern Black Sea Coast: New Marine Terrace Data from Eastern Pontides, Trabzon (Turkey) and a Review

Serdar Keskin, Kevin Pedroja, Osman Bektaş

## ► To cite this version:

Serdar Keskin, Kevin Pedroja, Osman Bektaş. Coastal Uplift along the Eastern Black Sea Coast: New Marine Terrace Data from Eastern Pontides, Trabzon (Turkey) and a Review. *Journal of Coastal Research*, 2011, 27 (6A), pp.63-73. 10.2112/JCOASTRES-D-09-00058.1 . hal-00735667

**HAL Id: hal-00735667**

**<https://hal.science/hal-00735667>**

Submitted on 23 Sep 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Coastal Uplift along the Eastern Black Sea Coast: New Marine Terrace Data from Eastern Pontides, Trabzon (Turkey) and a Review

Serdar Keskin<sup>†§</sup>, Kevin Pedoja<sup>‡\*</sup>, and Osman Bektaş<sup>§</sup>

<sup>†</sup>General Directorate of Mineral Research  
and Exploration  
The Mineral Etude and Research  
Department  
Ankara, Turkey 06520  
serdarkeskin@mta.gov.tr

<sup>‡</sup>UMR 6143 “Morphodynamique  
Continentale et Côtière” (M2C)  
Université de Caen  
2-4 rue des Tilleuls  
14000 Caen, France  
kevin.pedoja@unicaen.fr

<sup>§</sup>Karadeniz Technical University  
Engineering Faculty  
Department of Geology  
Trabzon, Turkey 61080

## ABSTRACT

An emerged, Quaternary, marine terrace sequence has been investigated near Trabzon, Turkey, along a ~20-km-long stretch of the coast of the southeastern Black Sea. The sequence includes seven principal marine terraces. The terraces are frequently compound and include up to three second-order marine terraces. The upper shoreline angles of the main terraces were found at the following maximum elevations above mean sea level:  $3 \pm 0.5$  m (TH),  $12 \pm 3$  m (T1),  $36 \pm 2$  m (T2),  $79 \pm 9$  m (T3),  $120 \pm 3$  m (T4),  $138 \pm 10$  m (T5), and  $260 \pm 25$  m (T6). Fossil bivalves and gastropods from the TH, T1, T2, and T3 deposits have been dated by Electron Spin Resonance (ESR). The results show that the ages of the deposits from the TH, T1, T2, and T3 terraces are  $5.141 \pm 0.294$  ka,  $124.8 \pm 26.0$  ka,  $292.5 \pm 49.8$  ka, and  $407.998 \pm 67.475$  ka, respectively. Consequently, we correlate TH, T1, T2, and T3 to marine isotope stage (MIS) 1, 5e, 9, and 11, which correspond to ~5, ~123, ~321, and ~400 ka highstands, respectively. Estimated uplift rates deduced from the elevations of T1, T2, and T3 are  $0.07 \pm 0.05$ ,  $0.10 \pm 0.02$ ,  $0.17 \pm 0.03$  mm/y or m/ka. These results show nonsteady, long-term uplift rates. Extrapolation of the oldest uplift rate (*i.e.*, determined on the highest dated terrace) shows that in the region of Trabzon, Turkey, coastal, positive, vertical deformations are recorded at ~2 Ma, which corresponds to the extrapolated age of the highest terrace in the sequence. Comparison with other sequences on the Black Sea coast reveals a rather weak uplift in this zone, which can be considered significative for the recent uplift of the Eastern Pontides.

**ADDITIONAL INDEX WORDS:** *Marine terrace, coastal uplift, Electron Spin Resonance, Eastern Pontides, Eastern Black Sea.*

## INTRODUCTION

The emerged sequence of Quaternary marine terraces near Trabzon, Turkey, lies on the northern zone of the Eastern Pontides orogenic belt in a context of plate collision. In this zone, studies on marine terraces began as early as the 1940s (Ardel, 1943; Erol, 1952; Ertek and Aytac, 2001; Keskin, 2007; Solmaz, 1990; Yilmaz *et al.*, 1997, 1998, 2005). The sequence of paleoshorelines, which increase in altitude from a few meters up to 260 m in a short distance, has long been interpreted as a consequence of uplift (Ardel, 1943; Erol, 1952; Yilmaz *et al.*, 2005) but surprisingly, previous studies did not focus on determining uplift rates. Consequently, no estimation of the Quaternary uplift of the Eastern Pontides orogenic belt was proposed through coastal morphotectonic studies. The main purposes of this work are to (1) refine the geomorphological data on marine terraces in this zone, (2) correlate the marine

terraces's shoreline angle to the Pleistocene highstands through Electron Spin Resonance (ESR) dating of the terrace deposits, (3) calculate the average uplift rates, and (4) compare the average uplift rates of the Trabzon, Turkey, area with uplift rates calculated through similar studies at other sites along the Eastern Black Sea coast.

## SETTINGS

### Geodynamics

The Black Sea is located between the Eurasian plate in the north and the African–Arabian plates in the south, within the Anatolian sector of the Alpine–Himalayan orogenic system (Figure 1). Around the Black Sea, this orogenic system includes the Pontides, the Greater Caucasus, and the Crimean Mountains. Plate models (DeMets *et al.*, 1989) and recent geodetic data (Reilinger *et al.*, 1997; Smith *et al.*, 1994) indicate that in the surrounding region, northward-moving African and Arabian plates collide with the Eurasian plate. As a consequence, the Anatolian block moves westward. Within this tectonic framework, the present-day tectonics of the Black Sea has been a puzzle (Tari *et al.*, 2000). The Black Sea basin is

\* Corresponding author.




Figure 1. Geodynamical context of the Eastern Black Sea. Map sketch redrawn from Smith *et al.*, 1994; Reilinger *et al.*, 1997; and Tari *et al.*, 2000.

divided into two parts, the Western and Eastern basins, separated by the mid-Black Sea ridge (Figure 1). The origin of the Black Sea has long been discussed, and it has been proposed that the entire Black Sea was rifted starting during Aptian-Albian period (Görür, 1993), but the age of the rifting for the Eastern Black Sea is less well known (Spadini *et al.*, 1996). Following the closure of the Neotethys, the Black Sea began to close during the Eocene-Oligocene age, and the Eastern Black Sea continued to close from the Miocene to the present. Seismic activity within the circum-Black Sea has been low to moderate during the 20th century. The area studied lies in the Eastern Pontides (NE Turkey) orogenic belt, which extends along the southeastern coast of the Black Sea region. This belt is divided into three subgroups from north to south, depending on lithological assemblages and facies (Figure 2A) (Bektaş *et al.*, 1999; Eyüboğlu *et al.*, 2006). The fault zones are oriented NW-SE, E-W, and NE-SW (Figure 2A). The northern zone (the magmatic arc), including the study area, is characterized by a dominance of subduction-related, upper Cretaceous and Tertiary volcanic rocks and by upper Cretaceous and Eocene arc granites.

The evolution of the Eastern Black Sea basin has been controlled by subsidence (Görür *et al.*, 1993; Nikishin *et al.*, 2003; Yılmaz *et al.*, 1997). Görür *et al.* (1993) and Yılmaz *et al.* (1997) emphasized that Pontides underwent regional subsidence and local uplift. Nikishin *et al.* (2003) proposed that the




Figure 2. Geology of the Eastern Pontides and Trabzon, Turkey, area. (A) Main tectonic features and tectonic zones of the Eastern Pontides: (1) Paleozoic metamorphic basement, (2) paleozoic granites, (3) serpentinite, (4) undifferentiated Mesozoic and Cenozoic rocks, (5) platform carbonates, (6) primarily Mesozoic sedimentary rocks, (7) Cretaceous and Eocene arc volcanics, (8) Upper Cretaceous and Eocene arc granites, (9) caldera or dome, (10) orthogonal drape and drag folds, (11) fault, (12) thrust fault, (13) normal fault, and (14) study area. Abbreviations: NAF = North Anatolian Fault, NEAF = Northeast Anatolian Fault (*per* Eyüboğlu *et al.*, 2006). (B) Geological map of the study area.

regional compression, folding, and thrusting started during the late Eocene-Oligocene, and the general uplift of the recent mountains that surround the Black Sea began nearly 10 million years ago. In the Transcaucasus and Turkish regions, large-scale uplift of the mountains started during the Oligocene, but an acceleration of the deformation started during the Late Sarmatian (~10 Ma) (Milanovsky, 1968, 1991). Milanovsky (1968, 1991) also suggested that in the Eastern Black Sea basin, tectonic subsidence rate were 70–80 m/Ma, 20–30 m/Ma, and 85 m/Ma during the Late Cretaceous-Eocene, the Oligocene-Miocene, and the Plio-Quaternary, respectively.

## Geology

In the study area (Figure 2B), six different units (excluding marine terraces) are distinguished (Yılmaz *et al.*, 1998): (1) the Upper Cretaceous-Paleocene Bakirköy Formation, (2) the Eocene-Neogene Kabaköy Formation, (3) the Miocene Kaymakli Formation, (4) the Pliocene Besirli Formation, (5) the Pliocene-Quaternary clays, and (6) the Quaternary alluvium. All units are briefly described in the next paragraph.

The Bakirköy Formation is composed of Upper Cretaceous-Paleogene age sandy limestone, red biomicrite, and marl intercalation (Eyüboğlu *et al.*, 2006). The Cenomanian to early the Tertiary period, especially the Eocene, is characterized by

the evolution of the sedimentary rocks in the southern zone and represented by dense volcanic activity in the northern zone. The Neogene (Mio–Pliocene) volcanic activity that formed the Bakirköy Formation cut all units in the region. The Miocene Kaymakli Formation overlies these volcanic rocks and was first described as Pontian clays by Özsayar (1971). This formation contains sandy, clayey siltstone. The Pliocene Besirli Formation is an outcropping near the seashore and is represented by basaltic and andesitic agglomerate, tuff, claystone–sandstone–marl intercalations, coarse-grained sandstone, loosely cemented conglomerate, and breccias. The Pliocene Besirli Formation and the Eocene–Neogene Kabaköy Formation (Yilmaz *et al.*, 1998) are uncomfortably overlaid by the Quaternary marine terraces and alluvium.

### Quaternary Sea-Level History of the Black Sea

Throughout geological time, the Black Sea experienced major sea-level changes, accompanied by environmental modifications, such as geomorphological reshaping (Panin and Popescu, 2007).

The Pleistocene epoch was characterized by alternations between a flow-type and a semiclosed water body. In a broad sense, marine transgressions occurred during the interglacials, whereas during the glacial periods, the basin became a flow-through, brackish water body (Algan *et al.*, 2007). During the Quaternary glaciations and delectation, the induced sea-level fluctuations endowed the restricted channels of the Bosphorus and Dardanelle with the power to control the Black Sea's connection to the Mediterranean Sea (Panin and Popescu, 2007). When global sea level dropped below the Bosphorus sill, the water level in the Black Sea varied according to regional conditions, independent of those in the World Ocean (Panin and Popescu, 2007). One important consequence of these lowstands was the interruption of Mediterranean inflow to the Black Sea, which, lacking any saline contribution gradually became a giant brackish to freshwater lake (Panin and Popescu, 2007). As noted by Zubakov (1988), the number and ages of glacial–interglacial cycles in continental Europe are identical to the climatic cycles in the Black Sea and Mediterranean Sea. During marine events, Mediterranean mollusks penetrated into the Black Sea and up to the Azov Sea (*e.g.*, Zubakov, 1988). Thus, the fluctuations of the Black Sea levels were effectively controlled by the global climate changes, with the higher sea levels corresponding to the increased inflow of water during the milder and wetter interstadial episodes and the lower levels occurring during the colder and drier stages (Arslanova, Dolukhanov, and Gei, 2007). For example, during the Last Interglacial (marine isotope stage [MIS] 5), eustatic rise in sea level allowed saline Mediterranean water to penetrate the Pontic basin (Panin and Popescu, 2007). This phase, locally named the Karangatian, introduced marine conditions that lasted about 60 ka (from 125 to ~65 ka BP). Marine terraces of this age (see “Marine Terraces” below) have been found around the Black Sea basin at elevations as high as 10 m on the Kerch and Crimea peninsulas (Panin and Popescu, 2007; Zubakov, 1988) and up to 48 m along the Caucasus coast because of the tectonic uplift (Nesmeyanov, 1995).

The sea level during the Black Sea's history has been rather

complicated and heavily discussed; there is still little consensus on the Black Sea variations in level, and different scenarios have been proposed to explain the evolution of the Black Sea since the Last Glacial Maximum (*e.g.*, Giosan, Filip, and Constatinescu, 2009; Hiscott *et al.*, 2007; Pirazzoli, 1991; Ryan and Pitman, 1999). The main conclusion in the latest review (Panin and Popescu, 2007) on this subject is that more extensive studies are needed to fill the remaining gaps in our knowledge about Late Pleistocene and Holocene sea-level changes. It must be pointed out here, that our study concentrated on the highstands of the Black Sea and the sea-level history of the Black Sea during the interglacials is not controversial. All authors, to our knowledge, agree on the connection of the Black Sea to the Mediterranean Sea through the Marmara Sea.

The maximum rise in the Black Sea (3–5 m above current level) took place 3.5–4 ka ago during the Subboreal period (Panin and Popescu, 2007). At this time, the so-called Old Black Sea terrace was formed (Panin and Popescu, 2007). A rapid lowering of the water level by –5 to –8 m followed, coeval with the first Greek colonization of the Black Sea coast. A new, short-lived ingression of the sea to a stand of +1 to +3 m then occurred. By about the 10th century AD, the level of the Black Sea experienced a decline of 1–2 m, then a slow rise, which continues today (Panin and Popescu, 2007).

In any case, most researchers agree that (1) all fauna encountered in the marine terrace deposits of the Trabzon area are characteristics of saline water (James Nebelsick, personal communication), (2) the ages provided by ESR dating on fossil bivalves and gastropods yield ages very close to the well-known, worldwide, Quaternary sea-level highstands, and (3) in the Black Sea, sea levels during past interglacials are not controversial. Authors agree on a connection of the Black Sea to the Mediterranean Sea through the Marmara Sea (see above). Therefore, we conclude that the marine terrace of the Trabzon area can be correlated to worldwide, major highstands.

## METHODS

### Marine Terraces

On uplifting coasts, marine terrace sequences are sometimes represented by flights of terraces up to hundreds of meters above mean sea level (Better Bull, 2008). Each terrace corresponds to a gently seaward dipping erosional or depositional platform backed by a relict sea cliff. The shoreline angle (*i.e.*, the intersection of the relict platform and the relict sea cliff) provides a good approximation of the location and elevation of a former shoreline and hence a relative sea level (Bull, 1985; Pedoja *et al.*, 2008). Most researchers now view marine terraces as being formed during the separated highstands of interglacial stages, which are correlated to marine oxygen-isotopic stages. The total displacement of the shoreline, relative to the age of the associated interglacial stage, allows us to calculate the mean uplift rate (Lajoie *et al.*, 1986) through the following classical equation:  $U = (E - e)/A$ , where  $U$  is the total uplift rate of each marine terrace,  $E$  is the present-day elevation of the marine terrace,  $e$  is the eustatic sea level of the considered MIS, and  $A$  is the age of the marine terrace






Figure 3. Marine terrace sequence in Trabzon, Turkey. The contour line is at 10-m intervals. (A) General repartition. (B) Zoom-in view of Yildizli. (C) Zoom-in view of Yalı. (D) Zoom-in view of Ganita. (E) Zoom-in view of the Trabzon Airport area.

corresponding to the age of the MIS considered for the carving of the shoreline angle. In other word, dating of the marine terrace deposits allows discrete correlation to an interglacial stage, the latter being itself correlated to an MIS (see, *e.g.*, Lajoie *et al.*, 1986).

The best-represented terrace worldwide is the one correlated to the Last Interglacial maximum (MIS 5e) (Hearty and Kindler, 1995; Johnson and Libbey, 1997; Pedoja *et al.*, 2006a, 2006b, 2006c, 2008). The age of MIS 5e is astronomically fixed to range from 130 to 116 ka (Kukla *et al.*, 2002) but is demonstrated to range from 134 to 113 ka in Hawaii and the Barbados, with a peak from 128 to 116 ka on tectonically stable coastlines (Muhs, Simmons, and Steinke, 2002; Shackleton *et al.*, 2003). In our calculations, we used a mean of  $123 \pm 11$  ka. Older marine terraces well represented in worldwide sequences are those related to MIS 9 (303–339 ka) and 11 (362–423 ka) (Imbrie *et al.*, 1984). For MIS 9 and 11, we used a value of  $321 \pm 18$  ka and  $400 \pm 20$ . Compilations show that the sea level was  $3 \pm 3$  m higher during mean isotope substage (MISS) 5e, MIS 9, and MIS 11 than at present and  $-1 \pm 1$  m, compared to the present level, during MIS 7 (Hearty and Kindler, 1995; Zazo, 1999). Consequently, MIS 7 (180–240 ka) (Imbrie *et al.*, 1984) marine terraces are less pronounced and sometimes absent (Zazo, 1999). Because the elevations of these terraces are higher than the uncertainties in paleoeustatic sea levels for the Holocene and Late Pleistocene, these uncertainties have no effect on the overall interpretation.

Field surveys and geological and geomorphological study of the marine terraces on the Trabzon coast (Figures 3, 4, and 5)




Figure 4. Altitudes of the marine terraces along the Trabzon, Turkey, coast. All altitudes are given as above mean sea level (amsl). See Figure 3 for the location of transect.

were carried out between 2004 and 2008. The marine terraces were named TH for the Holocene terrace and T1, T2, T3, T4, T5, and T6 (from lowest to highest) (Figures 3, 4, and 6) for the Pleistocene terraces. Elevations of the shoreline angle of the marine terraces were determined using a Global Positioning System (GPS), altimeter measurements, and ArcGIS software. All altitudes are given in meters above mean sea level (amsl) with an associated margin of error, which depends mainly on shoreline preservation. The vertical and horizontal distribution of the highest terraces and the associated deposits are not always well identified in the field or on aerial photographs. Consequently, in some cases, Geographic Information System (GIS) mapping was used to determine the distribution and elevation of marine terraces (Figures 3 and 4). Mollusk shell samples, mainly bivalves and gastropods fragments, were collected from deposits in TH, T1, T2, and T3 marine terraces for dating. As noted before, all the fauna identified are characteristics of saline water. The fauna is present as fragments, with the exception of T2 and T3 deposits, where we found *Chione* sp. (T2) and *Mytilus* sp. and *Ostrea* sp. (T3).

### ESR Dating

During the past 20 y, the ESR method has improved considerably in dating littoral deposits (coral reefs, barrier beach systems, and aeolianites) (Jonas, 1997). The ESR ages of Quaternary mollusk shells not only enable the chronological differentiation of the MIS 1, 5, 7, 9, 11 but also, sometimes, the substages of the main interglacials MIS 5a, 5c, 5e, 9, and 11




Figure 5. Illustration of the marine terraces on the Trabzon, Turkey, coast. (A) Interpreted panorama. Photo taken west of Trabzon city. (B) Detail of the Holocene terrace in Ganita. (C) Marine terrace T4 deposits overlying the Pliocene Besirli formation in Erdogdu. (D) Details of the *in situ* Holocene marine deposits on the TH terrace in the Yildizli area. Note the typical marine terrace deposits consisting of boulders and broken shells in a sandy matrix. (E) Details of the T2 terrace deposits as studied in the laboratory.

(Schellmann and Radtke, 2000; 2004) (Table 1). The temporal resolution of ESR dating on mollusk shells is within a mean age error of 10–15% or less (Schellmann *et al.*, 2004). To calculate the ESR age of mollusk shells, two parameters are needed: the equivalent dose and the total dose or annual dose rate (Tables 2 and 3). The equivalent dose is calibrated by construction of so-called growth curves. The total dose combines the cosmic dose rate and the  $\beta$  and  $\gamma$  doses from the potassium (K), thorium (Th), and uranium (U) concentrations in the sediment and the  $\alpha$ ,  $\beta$ , and  $\gamma$  doses from the U concentrations in the mollusk shells. All ESR measurements were carried out at the Geography Department, University of Cologne, and performed with a Bruker 400 ESP spectrometer, using the multiple aliquot additive method for the equivalent dose determination (see Tables 2 and 3). The radionuclide content of K, Th, and U was determined by inductively coupled plasma mass spectroscopy (ICP-MS) to calculate the annual dose rate. Equivalent doses were calculated using the FITSIM software program, and all ESR ages calculated by DATA 7 program. The ESR age is calculated dividing the equivalent dose rate by the total dose rate (Table 1).




Figure 6. Synthetic marine terrace sequence of the Trabzon, Turkey, area.

## RESULTS

### Marine Terrace Sequence of the Trabzon Region

A well-developed marine terrace sequence in the Trabzon region has been studied in detail. The sequence has seven main terraces, but they are frequently compound: two or three “second-order” terraces, separated by small, intermediate shoreline angles. In total, there are as many as 13 terraces, including the second-order ones (Figures 3, 4, and 6). The sequence can be described as two narrow, low-lying terraces (TH and T1) overlooked by higher compound levels (T2, T3, T4, T5, and T6) (Figure 6). Although the highest terraces (T4, T5, and T6) are distinguishable in the study area (in Bahcecik, Erdogdu, Yenicuma, and Boztepe sectors) (Figures 3 and 4), they have been strongly affected by (1) fluvial processes, (2) pedogenetization, and (3) anthropization (mainly farming).

In the Trabzon area, marine terrace deposits are composed of less than 2 m of sand, clay, pebbles, blocks, and boulders, mostly composed of sedimentary and volcanic rocks (andesite and basalt) from the substratum (the Pliocene Besirli Formation and Eocene–Neogene Kabaköy Formation). These deposits contain common fossil bivalves and gastropods, whole or broken. They are overlaid by brownish–dark brownish soil. Deposits are very similar from one terrace to another.

The TH terrace is well observed in the Yildizli and Ganita district (Figures 3A, 3B, 3D, 4, 5B, and 5D). It is present along an ~3-km stretch of coast between Yildizli and Akyazi and ~2 km between the towns of Trabzon and Kemer kaya. In this zone, the marine terrace is ~0.2 km wide, and its shoreline angle is present at an altitude of  $3 \pm 1$  m (Figure 6). In two outcrops located on the Yildizli point, we sampled fossil seashells for ESR dating (samples 1 and 2 in Figure 3B, laboratory numbers K-5339 and K-5341). A general view of the TH in Ganita is provided in Figure 5B, and an example of an *in situ* deposit of TH is provided in Figure 5D.

The T1 terrace is clearly recognizable and observable in the Yildizli, Yali, and Bostanci areas and in the area of the Trabzon Airport (Figures 3A–E and 4). The T1 is present along a 6-km stretch of coast between Yildizli and Besirli and along a 4-km stretch of coast between the city center and the airport. Its shoreline angle is present at a maximum altitude of  $12 \pm 3$  m amsl (Figure 6), and the terrace is nearly 1 km wide. In an

Table 1. Chronostratigraphical interpretation of the dated marine terraces in the Trabzon, Turkey, area.

Location	Sample	Latitude	Longitude	Elevation (m)	ESR age (ka)	MIS or MISS assigned to the terrace
Fatih	8	39°42'19.20" N	41°0'3.22" E	60 ± 2	407.998 ± 67.475	11
Fatih	9	39°42'20.20" N	40°59'58.67" E	67 ± 9	416.150 ± 92.669	11
Gazipasa	10	39°43'41.57" N	41°0'12.02" E	70 ± 9	395.099 ± 30.830	11
Yildizli	X-2a	39°37'15.65" N	41°0'30.65" E	30 ± 3	292.5 ± 49.8	9
KTÜ Sahil Tes.	X-42	39°46'39.48" N	41°0'3.81" E	12 ± 3	124.8 ± 26.0	5e
Yildizli-DSİ	2	39°37'33.58" N	41°0'26.84" E	3 ± 1	5.141 ± 294	1
Yildizli-DSİ	1	39°37'31.59" N	41°0'26.00" E	3 ± 1	5.008 ± 533	1

Abbreviations: MIS = marine isotope stage, MISS = marine isotope substage, ESR = Electron Spin Resonance, KTÜ = Karadeniz Technical University, Tes.

outcrop located near the airport, we sampled some fossils seashells for ESR dating (sample X-42 in Figure 3E, laboratory number K-5312 in Table 2).

The T2 terrace is best observed in Yıldizli, Degirmendere, the Trabzon Airport area, İnönü, Fatih, and around the Trabzon Stadium. The T2 is present along a 7-km-long stretch of coast between Trabzon and Toklu and also along a 4-km stretch of coast between the Trabzon Airport and the city center (Figures 3A–E and 4). This terrace is generally ~2 km wide. The T2 is a compound terrace that includes three second-order terraces: T2', T2'', and T2''' (Figure 6). From GIS data (Figures 3 and 6), we calculated that the shoreline angles of T2', T2'', and T2''' are, respectively, at  $20 \pm 2$ ,  $30 \pm 3$ , and  $36 \pm 2$  m, and we consider the highest shoreline angle ( $36 \pm 2$  m) as characteristic of the T2 marine terrace. In an outcrop located on the Yıldizli point, we sampled fossils seashells for ESR dating (sample X-2a in Figure 3A, laboratory number K-5311 in Table 2) and recognized bivalves, such as *Chione* sp. (for an example of the T2 deposits, see Figure 5E).

The T3 terrace is well observed in Yenice, Ayasofya, Fatih, İnönü, Gazipasa, AKL, and Degirmendere sectors (Figures 3A and C–E). The terrace is present along a 5-km stretch of coast between Toklu and Cömlekci and along a 4-km stretch of coast between Trabzon Airport and Degirmendere. Similar to T2, T3 is a compound terrace that includes at least three second-order marine terraces: T3', T3'', and T3'''. The T3', T3'', and T3''' terraces were calculated to be at altitudes of  $60 \pm 2$ ,  $67 \pm 9$ , and  $70 \pm 9$  m, respectively, by GIS and field measurements (Figure 6). In section F, the T3''' shoreline angle was measured at the maximum altitude of  $80 \pm 2$  m amsl (Figure 4). In three outcrops, two near Fatih and one near Gazipasa, we sampled some fossil seashells for ESR dating (samples 8–10 in Figures 3C and D, laboratory numbers K-5335, K-5336, and K-5347 in Table 3).

The T4 terrace is well observed in Bahcecik, Besirli, Yenice, and Erdogdu (Figures 3A, C, and E). It is present along an 8-km stretch of coast between Besirli and the Karadeniz Technical University campus. In this zone, the T4 is 1 km wide. Similar to T2 and T3, T4 is a compound terrace that includes at least three second-order marine terraces: T4', T4'', and T4''' (Figure 6). The shoreline angle of T4' is present at  $102 \pm 5$  m in section H. The shoreline angles of T4'' and T4''' were found, respectively, to be at  $110 \pm 5$  m and  $120 \pm 3$  m in section H (Figures 4 and 6). A photograph of the T4 deposits (Figure 5C) shows it to be lying uncomfortably on the Besirli Formation.

The T5 terrace is well observed in the Bostanci, AKL, and Bahcecik districts (Figures 3A, C, and E). It is present along a 5-km stretch of coast between the Bahcecik and Bostanci districts, where it is ~1 km wide. The shoreline angle of the T5 has been measured at  $127 \pm 5$  m amsl in section E. Based on GIS analysis, the shoreline angle of the T5 is present at  $138 \pm 10$  m in section G and at  $135 \pm 5$  m amsl in section H (Figure 4).

The T6 terrace is clearly observed in Boztepe, Bostanci, Bahcecik, and Cezaevi districts (Figures 3A, C, D, and E). It is present along a 6-km stretch of coast between Bahcecik and Bostanci. Based on GIS analysis and field survey, the shoreline angle of the T6 is present at an altitude of  $260 \pm 25$  m (Figure 6).

## Dating and Chronostratigraphical Interpretation

Fossil bivalves and gastropod were collected from the deposits of TH, T1, T2, and T3 to obtain geochronological measurements through the ESR method. Two samples collected in the TH deposits in the Yıldizli area gave corrected ESR ages of  $5.008 \pm 0.533$  ka and  $5.141 \pm 0.294$  ka (Figure 3B; Table 3). Consequently, we correlated TH to the Holocene (MIS

Table 2. Dosimetric parameters and Electron Spin Resonance (ESR) ages obtained from marine shells.

Laboratory No.	Sample No.	K (ext) (%) <sup>1</sup>	Th (ext) (ppm) <sup>1</sup>	U (ext) (ppm) <sup>1</sup>	U (int) (ppm) <sup>1</sup>	Water Content (%)	Depth (m)	Total Dose Rate <sup>2,3</sup>		
								Gy/ka	D <sub>e</sub> (Gy)	ESR Age <sup>4</sup> (ka)
K-5312	X-42	0.63	2.57	2.41	1.48	15 ± 10	4.5 ± 1	0.977 ± 0.08	121.9 ± 23.2	124.8 ± 26.0
K-5311	X-2a	0.74	4.55	1.4	1.43	10 ± 5	0.2 ± 0.1	1.230 ± 0.08	359.9 ± 56.7	292.5 ± 49.8

<sup>1</sup>Uncertainties on ICP-MS analyses of U, Th, and K are taken to be ±1%.

<sup>2</sup>The total dose combines the cosmic dose rate; the β and γ doses from the K, Th, and U concentrations in the sediment; and the α, β, and γ doses from the U concentrations in the shells. Sample thickness was 4–5 mm, the thickness removed was 0.4–1 mm (age calculation with β-attenuation).

<sup>3</sup>D<sub>e</sub> = equivalent dose (calculated via the program FIT-SIM [R. Grün]).

<sup>4</sup>Ages were calculated via the program DATA 7 (R. Grün).



Table 3. Dosimetric parameters and Electron Spin Resonance (ESR) ages obtained from marine shells.

Laboratory No.	Sample No.	Th (ppm) <sup>1</sup>	U (ppm) <sup>1</sup>	K (%) <sup>1</sup>	Thickness (mm)	Water content (%)	Depth (m)	D <sub>e</sub> <sup>2</sup> (Gy)	ESR Age <sup>3</sup> (ka)
K-5335	8	5.68	1.06	0.63	1.28/0.84 (0.44)	7	10	463.51 ± 69.91	407.998 ± 67.475
K-5336	9	2.72	0.76	0.45	1.46/1.15 (0.31)	4	0.3	463.51 ± 69.91	416.150 ± 92.669
K-5337	10	2.93	0.72	0.68	1.26/1.10 (0.16)	8	0.25	379.72 ± 12.49	395.099 ± 30.830
K-5339	1	3.77	0.83	1.33	1.20/0.96 (0.24)	12.59	8–8.5	5.02 ± 0.47	5.008 ± 533
K-5341	2	3.93	0.83	1.33	1.18/1.02 (0.16)	8.92	9	5.35 ± 0.16	5.141 ± 294

<sup>1</sup>Uncertainties on ICP-MS analyses of U, Th, and K are taken to be ±1%.<sup>2</sup>D<sub>e</sub> = equivalent dose (calculated *via* the program FIT-SIM [R. Grün]).<sup>3</sup>Ages were calculated *via* the program DATA 7 [R. Grün].

1) (Table 1). Sample X-2a, taken from the T1 deposits, gave an ESR age of  $124.8 \pm 26.0$  ka (Figure 3B; Table 2). Consequently, we propose to correlate this well-preserved marine terrace to the Last Interglacial maximum (MISS 5e,  $123 \pm 11$  ka) (Table 1). Sample X-42, collected from T2' deposits in the university area (Figure 3E), gave an ESR age of about  $292.5 \pm 49.8$  ka (Table 2). This result leads us to propose a correlation of the upper shoreline of T2 with MIS 9 (303–339 ka) (Table 1). Samples 8, 9, and 11, taken from T3', T3'', and T3''' deposits, yielded ESR ages of  $395.099 \pm 30.830$  ka,  $407.998 \pm 67.475$  ka, and  $416.150 \pm 92.669$  ka, respectively. Consequently, we propose to correlate the highest T3 shoreline angle with MIS 11 (362–423 ka) and probably with an early substage of this complex interglacial period.

## DISCUSSION

### Uplift Rates and Extrapolation

Because the maximum rise in the Black Sea (3–5 m above current level) took place 3.5–4 ka ago, during the Subboreal, and given the present elevation of the TH, we did not calculate the uplift rate for this period. In other words, the elevation of the TH can be explained by eustatic variation alone.

The uplift rate for each dated Pleistocene shoreline was calculated using parameters shown in Table 4. The T1, T2, and T3 altitudes and age lead us to determine uplift rates of  $0.07 \pm 0.05$ ,  $0.10 \pm 0.02$ , and  $0.17 \pm 0.03$  mm/y or m/ka for MIS 5e, 7, and 9, respectively (Table 4, Figure 7). The low elevation of the terraces, the uncertainties about the duration, and the eustatic sea level of Last Interglacials (Siddall *et al.*, 2006) create a notable margin of error, particularly for the T1 terrace (Table 4). Nevertheless, mean value reflects the sequence morphostratigraphy, *i.e.*, two low, well-preserved terraces below higher, compound levels (T2, T3, T4, T5, and T6). Therefore, the variation in the mean uplift rate can be considered representative of the evolution of the deformation. A slight decrease was observed. Relationships were observed

between marine terrace development and nonsteady, long-term uplift rates, more drastically in Futuna (Neef and McCulloch, 2003), and, more recently, in Chile (Saillard *et al.*, 2009). Generally these variations are interpreted as pulses of tectonic activity.

Nevertheless, to have an approximation of the duration of the uplift, we extrapolated the T3 uplift rates, the oldest uplift, to propose “ages” for the oldest terraces (T4, T5, and T6). As stated in a recent synthesis (Siddall *et al.*, 2006), we consider eustatic sea levels older than MIS 11 to be similar to those present now. This means that the actual altitude of the shoreline corresponds roughly to its displacement (Figure 7). Following that idea, the highest shoreline angles of the three upper marine terraces were carved during the following time intervals: 680–910 ka (T4), 750 ka to ~1.1 Ma (T5), and 1.4–2.1 Ma (T6). As a preliminary result, we propose that the entire marine terrace sequence in the Trabzon region is from the Quaternary period.

Given that finding, the observed uplift has been rather slow in comparison to other orogenic belts: in the Caucasus (see below and Table 5) or in Morocco, on the sea side of the Atlas range (Meghraoui *et al.*, 1998). In Morocco, north of Agadir, the MIS 5e terrace is found at elevations ranging from 18 to  $28 \pm 1$  m amsl. Following the calculation we used for the Eastern Black Sea, we calculate a maximum uplift rate of  $0.20 \pm 0.05$  mm/y or m/ka.

### Origin of the Uplift

The marine terrace in Trabzon reveals a rather slow, Quaternary uplift of the southeastern Black Sea shore. From our point of view, these deformations have to be related to the Pontides orogenesis. More precisely, in the studied area, three distinctive lineaments, trending NE–SW, NW–SE, and E–W (Figure 2A), are present and are probably responsible for the uplift of the marine terrace sequence. At this stage, it must be pointed out that these faults are not directly observed in the

Table 4. Measured shoreline angles and calculated uplift rates for marine terraces T1, T2, and T3.

Terrace	MIS	E (m)		A (ka)		e (m)		D (m)		U (mm/y, m/ka)	
		E	MoE	A	MoE	e	MoE	D	MoE	U	MoE
T1	5e	12	3	123	11	3	3	9	6	0.07	0.05
T2	9	36	2	321	18	3	3	33	5	0.1	0.02
T3	11	70	9	400	20	3	3	67	12	0.17	0.03

Abbreviations: MIS = marine isotope stage, MoE = margin of error.






Figure 7. Graphic representation of the uplift rates deduced from the marine terraces along the Trabzon, Turkey, coast.

field because of urbanization and vegetation, but they are seen on aerial photos and geomorphological maps. We suggest that the marine terrace of Trabzon reveals the vertical motion rates of active faults and, more particularly, of the northern fault; however, at this stage in our knowledge, we can not exclude the possibility that the uplift is caused by local compression associated with a restraining bend in the eastern segment of the North Anatolian Fault. To improve upon our hypothesis, other geomorphic markers need to be studied directly to the south of our study area. In a broad sense, we propose that the primary cause of the uplift is the local compression associated with the convergence of the African–Arabian plates with the Eurasian plate. The variation in the uplift rates may be a consequence of tectonic pulses during the ongoing Eastern Pontides orogenesis.

### Comparison with Other Marine Terrace Sequence along the Eastern Black Sea Shore

Other studies of marine terraces have been conducted along the Eastern Black Sea coast (Figures 1 and 8). Marine terrace sequences have been studied in Russia and in the Ukraine (Nesmeyanov, 1995; Papin and Popescu, 2007; Zubakov, 1988), where the deformations are related to the Caucasus orogenesis. On the Black Sea shore of Turkey, a sequence was briefly described on the western part of the Sinop Peninsula, where it is related to the Central Pontides orogenesis (Figures 1 and 8) (Ertek and Aytaç, 2001).

To compare the deformation rates of these different zones, we propose to calculate the homogeneous uplift rates deduced from the marine terrace correlated to the Last Interglacial maximum. Consequently, we use the same values for  $e$  and  $A$  (see the equation in the “Methods” section). In other words, we propose to establish the Eemian (or Karangatian) marine terrace as tectonical benchmark, as has already been done in Italy (Antonioli *et al.*, 2006; Bordini and Valensise, 1998). In this synthesis, we excluded data from the Marmara Sea (Yaltırak *et al.*, 2002) that corresponds to a different geodynamical context. In fact, this zone has complicated, active tectonics, influenced by the escape of the Anatolian block, the northward motion of the African plate, and the Aegean extension, leading to a global extensional regime (*e.g.*, Tari *et al.*, 2000).

The results of these calculations are presented in Figure 8 and Table 5. Around the Eastern Black Sea, altitudes of MIS 5e

shoreline angles range from  $6 \pm 2$  mm/a to  $48 \pm 3$  mm/a in the Taman Peninsula and the Tuapse Loo area, respectively (Table 5). Consequently, uplift rates deduced from the elevation of the MIS 5e terrace range from  $0.02 \pm 0.04$  mm/a to  $0.37 \pm 0.08$  mm/a (Figure 8; Table 5). The most striking result is that the Trabzon marine terrace sequence reveals one of the weakest uplifts of the Eastern Black Sea shore ( $0.07 \pm 0.06$  mm/a for MIS 5e). At a regional scale, the highest coastal uplift rates are found on the coasts bordering the Greater Caucasus and the Central Pontides (Sinop Peninsula). The greatest variability is observed on the coasts bordering the Greater Caucasus, where the marine terrace sequence is defined by arching (Nesmeyanov, 1995). This fact should be interpreted as the difference in the intensity of orogenesis between the Greater Caucasus and the Eastern Pontides. No marine terrace sequences have been described, to our knowledge, in the Lower Caucasus (*e.g.*, near Batumi in Figure 1), and data are lacking in Abkhazia (Figures 1 and 8). In other words, studies of the Quaternary coastal uplift of the Eastern Black Sea are still in the preliminary stages, and more work has to be done (by us or other geologists) to complete the puzzle and to provide a coherent, regional interpretation.

### CONCLUSION

- (1) A Pleistocene marine terrace sequence borders the Eastern Black Sea in the vicinity of Trabzon, Turkey, at elevations ranging from  $3 \pm 1$  m up to  $260 \pm 25$  m. The ESR dates on *in situ* shells extracted from shallow marine deposits at key localities show that these sediments accumulated between MIS 11 and the Holocene, with the best-developed successions occurring during the interglacial oxygen isotopic stages 11, 9, 5e, and 1. Older terraces are present and have extrapolated ages corresponding to the Early Pleistocene (2 Ma).
- (2) The ages of the raised terraces and shoreline deposits document that the southeastern Black Sea shore, north of Northeast Anatolian Fault Zone, has been experiencing maximal regional uplift of  $0.017 \pm 0.03$  mm/y or m/ka. We propose that this local uplift owes its origin to the Pontides orogenesis as a consequence of the convergence of the African–Arabian plate with the Eurasian plate.
- (3) Comparison of the uplift rates in the Trabzon area with other places around the Black Sea shows that the vertical reaction of this portion of coast is quite limited. The Caucasus orogenesis seems to have had more of an effect on the Quaternary coastal deformation than on the Pontides orogenesis.
- (4) The Quaternary shoreline displacements around the Black Sea in general, and eastern Turkey in particular, are still poorly known. New ESR data allow us to revise and reinterpret, in one place, the literature on this topic, taking into account only those markers that allow precise estimates of the altitude and age of the sea-level highstands. Several sequences of Quaternary marine terrace, with associated deposits, can be observed along the Black Sea coast of Turkey and need further attention to refine our knowledge of the Pontides orogenesis in

Table 5. *Synthesis on MIS 5e terraces along the Eastern Black Sea coast.*

No.	Country	Latitude	Longitude	Latitude	Longitude	Name	No. of Sequences Described	Min. Length of Coastal Stretch (km)	Min. Dimension of the Sequences (km)	Max. No. of Palaeoshoreline	Max. Altitude of Sequence	Max. Altitude of MIS 5e with MoE (m amsl)	Chrono-stratigraphy	Uplift Rates		Dating Method	Dating performed on	Reference
														Rate	MoE			
1	Ukraine	44°28'0.00" N	34°7'0.00" E	44°28'0.00" N	34°7'0.00" E	Crimea	1	30	ng	4	10	10	Holocene, MIS 5e	0.06	0.05	*	*	Panin & Popescu, 2007
2	Russia	45°22'0.00" N	36°28'0.00" E	45°22'0.00" N	36°28'0.00" E	Kerch	3	30	ng	4	16	10	Holocene, MIS 5e	0.06	0.05	U/Th	Shells	Panin & Popescu, 2007; Zubakov, 1988
3	Russia	45°10'52.89" N	36°38'51.74" E	45°10'52.89" N	36°38'51.74" E	Taman	3	58	ng	4	16	6	Holocene, MIS 5e	0.02	0.04	U/Th	Shells	Panin & Popescu, 2007; Zubakov, 1988
4	Russia	44°53'28.23" N	37°19'29.04" E	44°30'3.48" N	38°7'33.69" E	Anapa-Gelendzhik	5	91	ng	7	175	25	MIS 5e and lower Pleistocene	0.18	0.08	Fauna	shells	Nesmeyanov, 1995
5	Russia	44°22'46.38" N	38°22'53.13" E	44°22'46.38" N	38°22'53.13" E	Pshadsk	6	28	ng	7	175	22	MIS 5e and lower Pleistocene	0.15	0.06	Fauna	shells	Nesmeyanov, 1995
6	Russia	44°18'49.45" N	38°42'1.10" E	44°18'49.45" N	38°42'1.10" E	Dzhulgga	8	30	ng	6	210	30	MIS 5e and lower Pleistocene	0.22	0.09	Fauna	shells	Nesmeyanov, 1995
7	Russia	44°0'3.91" N	39°11'22.13" E	44°0'3.91" N	39°11'22.13" E	Tuapse-Loo	1	60	3 × 1.2	6	135 ± 10	48	MIS 5e and lower Pleistocene	0.37	0.08	Fauna	shells	Nesmeyanov, 1995
8	Russia	43°35'53.94" N	39°42'59.43" E	43°35'53.94" N	39°42'59.43" E	Sochi	7	11	ng	9	120	27.5	MIS 5e and lower Pleistocene	0.20	0.06	Fauna	shells	Nesmeyanov, 1995
9	Russia	43°25'31.16" N	39°55'15.80" E	43°25'31.16" N	39°55'15.80" E	Adler	6	10	ng	3	120	20	MIS 5e and lower Pleistocene	0.14	0.08	Fauna	shells	Nesmeyanov, 1995
10	Turkey	41°0'21.62" N	39°37'52.88" E	41°0'0.48" N	39°46'10.31" E	Trabzon	1	12	5 × 3	7	260 ± 25	12	Holocene, MIS 5e, 9, 11	0.07	0.06	ESR	Shells	This study
11	Turkey	42°5'24.48" N	34°57'4.40" E	42°5'24.48" N	34°57'4.40" E	Sinop West	1	ng	ng	ng	40	5	MIS 5e	0.30	0.09	*	*	Ertek & Aytac, 2001

Abbreviations: MIS = marine isotope stage, No. = number, MoE = margin of error, Min. = minimum, Max. = maximum.




Figure 8. Structural framework and coastal uplift along the Eastern Black Sea Coast.

particular, and the Quaternary coastal deformation in general.

## ACKNOWLEDGMENTS

We are indebted to Steve Kershaw for his careful revision of the manuscript and to colleagues in the Department of Geodesy and Photogrammetry Engineering at Karadeniz Technical University for digitizing the map and preparing the GIS data. This work was supported by funds from Karadeniz Technical University research grants (2005.112.05.2 and 2006.118.001.2). All ESR dating was performed by Ulrich Radtke in the Geography Department at the University of Cologne.

## LITERATURE CITED

- Algan, A.; Ergin, M.; Keskin, S.; Gökaşan, E.; Alpar, B.; Ongan, D., and Kirci-Elmas, E., 2007. Sea-level changes during the late Pleistocene-Holocene on the southern shelves of the Black Sea. In: Yanko-Hombach, Y.; Gilbert, A.S.; Panin, N., and Dolukhanov, P. (eds.) *The Black Sea Flood Question: Changes in Coastline, Climate, and Human Settlement*. Dordrecht, The Netherlands: Springer, pp. 603–631
- Antonoli, F.; Kershaw, S.; Renda, P.; Rust, D.; Belluomini, G.;

- Cerasoli, M.; Radtke, U., and Silenzi, S., 2006. Elevation of the last interglacial highstand in Sicily (Italy): a benchmark of coastal tectonics. *Quaternary International*, 145–146, 3–18.
- Ardel, A., 1943. Trabzon ve Civarının Morfolojisi Üzerine Gözlemler. *Türk Coğrafya Dergisi*, 1, 71–82. [In Turkish]
- Arslanova, K.H.A.; Dolukhanov, P.M., and Gei, N.A., 2007. Climate, Black Sea levels and human settlements in Caucasus Littoral 50,000–9000 BP. *Quaternary International*, 167–168, 121–127.
- Bektaş, O.; Şen, C.; Atıcı, Y., and Köprübaşı, N., 1999. Migration of the upper cretaceous subduction related volcanism towards the back-arc basin of the eastern Pontide magmatic arc (NE Turkey). *Geological Journal*, 34, 95–106.
- Bordonì, P. and Valensise, G., 1998. Deformation of the 125 ka marine terrace in Italy: tectonic implications. *Geological Society, London, Special Publications*, 146, 71–110.
- Bull, W.B., 1985. Correlation of flights of global marine terraces. In: Morisawa, M. and Hack, J. (eds.), *Tectonic Geomorphology: Proceedings of the 15th Annual Geomorphology Symposium*. Binghamton, New York: State University of New York, pp. 129–152.
- De Mets, C.; Gordon, R.G.; Argus, D.F., and Stein, S., 1989. Current plate motions. *Geophysical Journal*, 101, 425–478.
- Erol, O., 1952. Trabzon sekileri hakkında bir not. *Dil ve Tarih-Coğrafya Fakültesi Dergisi*, 10, 125–136. [In Turkish]
- Ertek, T.A. and Aytaç, A., 2001. Karadeniz kıyılarımızda denizel taraçaların korelasyonu [The Marine Terraces Correlation on the Black Sea Coasts of Turkey]. In: Kuvaterner Sempozyumu 3, (İTÜ, İstanbul, Turkey), pp. 71–74. [In Turkish]
- Eyüboğlu, Y.; Bektaş, O.; Seren, A.; Nafiz, M.; Jacoby, W.R., and Özer, R., 2006. Three-directional extensional deformation and formation of the Liassic rift basins in the eastern Pontides (NE Turkey). *Geologica Carpathica*, 57, 5, 337–346.
- Giosan, L.; Filip, F., and Constatinescu, S., 2009. Was the Black Sea catastrophically flooded in the early Holocene? *Quaternary Science Reviews*, 28, pp. 1–6.
- Görür, N.; Tüysüz, Z.O.; Aykol, A.; Sakinc, M.; Yigitbac, E., and Akkok, R., 1993. Cretaceous red pelagic carbonates of northern Turkey: their place in the opening history of the Black Sea. *Eclogae Geologicae Helvetica*, 86, 819–838.
- Hearty, P.J. and Kindler, P., 1995. Sea-level highstand chronology from stable carbonate platforms (Bermuda and the Bahamas). *Journal of Coastal Research*, 11, 675–689.
- Hiscott, R. N.; Aksu, A.E.; Mudie, P.J.; Marret, F.; Abrajano, T.; Kaminski, M.A.; Evans, J.; Çakiroğlu, A.I., and Yasar, D., 2007. A gradual drowning of the southwestern Black Sea shelf: evidence for a progressive rather than abrupt Holocene reconnection with the eastern Mediterranean Sea through the Marmara Sea gateway. *Quaternary International*, 167–168, 9–34.
- Imbrie, J.; Hays, J.D.; Martinson, D.G.; McIntyre, A.; Mix, A.L.; Morley, J.J.; Pisias, N.G.; Prell, W., and Shackleton, N.J., 1984. The orbital theory of Pleistocene climate: support from a revised chronology of the marine  $^{18}\text{O}$  record. In: Berger, A.; Imbrie, J.; Hays, J.; Kukla, G., and Saltzman, B. (eds). *Milankovitch and Climate*. Dordrecht, The Netherlands: Reidel Publications, pp. 269–305.
- Johnson, M.E. and Libbey, L., 1997. Global review of upper Pleistocene (substage 5e) rocky shores: tectonic segregation, substrate variation, and biological diversity. *Journal of Coastal Research*, 13, 297–307.
- Jonas, M., 1997. Concepts and methods of ESR dating. *Radiation Measurements*, 27, 943–973.
- Keskin, S., 2007. Güneydoğu (GD) Karadeniz Sahil Kesiminin (Trabzon Yöresi) Denizel Taraçaları ve Aktif Tektoniği. Trabzon, Turkey: Fen Bilimleri Enstitüsü, Karadeniz Technical University, Master's thesis, 113p. [In Turkish]
- Kukla, G.J.; Michael, L.B.; de Beaulieu, J.-L.; Bond, G.; Broecker, W.S.; Cleveringa, P.; Gavin, J.E.; Herbert, T.D.; Imbrie, J.; Jouzel, J.; Keigwin, L.D.; Knudsen, K.-L.; McManus, J.F.; Merkt, J.; Muhs, D.R.; Müller, H.; Poore, R.Z.; Porter, S.C.; Seret, G.; Shackleton, N.J.; Turner, C.; Tzedakis, P.C., and Winograd, I.J., 2002. Last interglacial climates. *Quaternary Research*, 58, 2–13.
- Lajoie, K.R., 1986. Coastal tectonics. In: Wallace, R.E. (ed.). *Active Tectonics: Impact on Society*. Washington, DC: National Academy Press, pp. 95–124.
- Meghraoui, M.; Outtani, F.; Choukri, A., and De Lamotte, D.F., 1998. Coastal tectonics across the south atlas thrust front and the Agadir active zone, Morocco. *Geological Society Special Publications*, 146, 239–253.
- Milanovsky, E.E., 1968. Neotectonics of the Caucasus. Nedra, Moscow. 683 pp., in Russian.
- Milanovsky, E.E., 1991. *Geology of the USSR, Part 3*. Moscow, Russia: Moscow University Press. 272p. [In Russian]
- Muhs, D.R.; Simmons, K.R., and Steinke, B., 2002. Timing and warmth of the last interglacial period: new U-series evidence from Hawaii and Bermuda and a new fossil compilation for North America. *Quaternary Science Reviews*, 21, 1355–1383.
- Neef, G. and McCulloch M.T., 2003. Pliocene–Quaternary history of Futuna Island, south Vanuatu, southwest Pacific. *Australian Journal of Earth Sciences*, 48, 805–814.
- Nesmeyanov, S.A., 1995. Pleistocene deformation of Black Sea terraces along the Caucasus coast. *Geotectonics*, 29, 3, 259–269.
- Nikishin A.M.; Korotaev, M.V.; Ershov A.V., and Brunet, M.-F., 2003. The Black Sea basin: tectonic history and Neogene–Quaternary rapid subsidence modelling. *Sedimentary Geology*, 156, 149–168.
- Özsayar, T., 1971. Paläontologie und Geologie des Gebietes Östlich Trabzon (Anatolien). Ph.D. thesis. *Giesener Geologische Schriften*, Heft 1, 138p. [In Turkish]
- Panin, N. and Popescu, I., 2007. The northwestern Black Sea: climatic and sea-level changes in the late Quaternary. In: Yanko-Hombach, Y.; Gilbert, A.S.; Panin, N., and Dolukhanov, P. (eds.) *The Black Sea Flood Question: Changes in Coastline, Climate, and Human Settlement*. Dordrecht, The Netherlands: Springer, pp. 387–404.
- Pedoja, K.; Bourgeois, J.; Pinegina, T., and Higman, B., 2006a. Does Kamchatka belong to North America? an extruding Okhotsk block suggested by coastal neotectonics of the Ozernoi Peninsula, Kamchatka, Russia. *Geology*, 34(5), 353–356.
- Pedoja, K.; Dumont, J.F.; Lamothe, M.; Ortlieb, L.; Collot, J.-Y.; Ghaleb, B.; Auclair, M.; Alvarez, V., and Labrousse, B., 2006b. Plio Quaternary uplift of the Manta Peninsula and La Plata Island and the subduction of the Carnegie Ridge, central coast of Ecuador." *Journal of South American Earth Sciences*, 22(1–2), 1–21.
- Pedoja, K.; Ortlieb, L.; Dumont, J.F.; Lamothe M.; Ghaleb, B.; Auclair, M., and Labrousse, B., 2006c. Quaternary coastal uplift along the Talara Arc (Ecuador, Northern Peru) from new marine terrace data. *Marine Geology*, 228, 73–91.
- Pedoja, K.; Shen, J.W.; Pedoja, K.; Shen, J.-W.; Kershaw, S., and Tang, C., 2008. Coastal Quaternary morphologies on the northern coast of the South China Sea, China, and their implications for current tectonic models: a review and preliminary study. *Marine Geology*, 255, 103–117.
- Pirazzoli, P.A., 1991. *World Atlas of Holocene Sea-Level Changes*. Amsterdam, The Netherlands: Elsevier, 300p.
- Reilinger, R.E.; McClusky, S.C., and Souter, B.J., 1997. Preliminary estimates of plate convergence in the Caucasus collision zone from GPS measurements. *Geophysical Research Letters*, 24, 1815–1818.
- Ryan, W.B.F. and Pitman III, W.C., 1999. Noah's Flood: the New Scientific Discoveries about the Event that Changed History. New York: Simon and Schuster, 319p.
- Saillard, M.; Hall, S.R.; Audina, L.; Farber, D.L.; Hérail, G.; Martinod, J.; Regard, V.; Finkel, R.C., and Bondoux, F., 2009. Non-steady long-term uplift rates and Pleistocene marine terrace development along the Andean margin of Chile (31°S) inferred from  $^{10}\text{Be}$  dating. *Earth and Planetary Science Letters*, 227(1–2), 50–63.
- Schellmann, G. and Radtke, U., 2000. ESR dating stratigraphically well-constrained marine terraces along the Patagonian Atlantic coast (Argentina). *Quaternary International*, 68–71, 261–273.
- Schellmann, G. and Radtke, U., 2004. *The Marine Quaternary of Barbados*. Cologne, Germany: Kölner Geographische Arbeiten. 137p.
- Schellmann, G.; Radtke, U.; Potter, E.-K.; Esat, T.M., and McCulloch, M.T., 2004. Comparison of ESR and TMS U/Th dating of marine isotope stage (MIS) 5a, 5c, and 5e coral from Barbados—implications for palaeo sea-level changes in the Caribbean. *Quaternary International*, 120, 41–50.

- Shackleton, N.J.; Sanchez-Goni, M.F.; Pailler, D., and Lancelot, Y., 2003. Marine isotope substage 5e and the Eemian interglacial. *Global and Planetary Change*, 36, 151–155.
- Siddall, M.; Chappell, J., and Potter, E.K., 2006. Eustatic sea level during past interglacials. In: Sirocko, F.; Claussen, M.; Litt, T., and Sánchez-Goni, M.F. (eds.). *Developments in Quaternary Science*, Volume 7: *The Climate of Past Interglacials*. Elsevier: 75–92.
- Smith, D.E.; Kolenkiewicz, R.J.; Robbins, W.; Dunn, P.J., and Torrence, M.H., 1994. Horizontal crustal motion in the central and eastern Mediterranean inferred from satellite laser ranging measurements, *Geophysical Research Letters*, 21, 1979–1982.
- Solmaz, F., 1990. *Vakfikebir-Yomra Arası Kıyı Şeridinin Morfolojisi ve Taraçalar*. Istanbul, Turkey: Istanbul University, Sos. Bil. Ens., Yüksek Lisans Tezi. [In Turkish]
- Spadini, G.; Robinson, A., and Cloetingh, S., 1996. Western versus Eastern Black Sea tectonic evolution: pre-rift lithospheric controls on basin formation. *Tectonophysics Dynamics of Extensional Basins and Inversion Tectonics*, 266(1–4), 139–154
- Tari, E.; Sahin, M.; Barka, A.; Reilinger, R.; King, R.W.; McClusky, S., and Prilepin, M., 2000. Active tectonics of the Black Sea with GPS. *Earth Planets and Space*, 52, 747–751.
- Yaltirak, C.; Saking, M.; Aksu, A.E.; Hiscott, R.N.; Galeb, B., and Ulgen, U.B., 2002. Late Pleistocene uplift history along the southwestern Marmara Sea determined from raised coastal deposits and global sea-level variations. *Marine Geology*, 190(1–2), 283–305.
- Yilmaz, B.S.; Güç, A.R.; Gülibrahimoğlu, İ.; Yazici, E.N.; Konak, O.; Yaprak, S., and Köse, Z., 1998. *Trabzon İlinin Çevre Jeolojisi ve Doğal Kaynakları*. Ankara, Turkey: Maden Tetkik ve Arama Genel Müdürlüğü.
- Yilmaz, C.; Şen, C.; Şener, Ş.; Kandemir, R.; Karsli, O., and Bak T.K., 2005. Trabzon Kıyı Bölgesinin Pliyo-Kuvaterner Stratigrafisi. *TURQUA V: Türkiye Kuvaterner Sempozyumu* (Istanbul Technical University, Turkey), pp. 111–117.
- Yilmaz, Y.; Tüysüz, O.; Yiğitbaş, E.; Genç, Ş.C., and Sengör, A.M.C., 1997. Geology and tectonic evolution of the Pontides. In: Robinson, A.G. (ed.). *Regional and Petroleum Geology of the Black Sea and Surrounding Region*. Tulsa, Oklahoma: American Association of Petroleum Geologists Memoir 68, pp. 183–226.
- Zazo, C., 1999. Interglacial sea levels. *Quaternary International*, 55, 101–113.
- Zubakov, V.A., 1988. Climatostratigraphic scheme of the Black Sea Pleistocene and its correlation with the oxygen—oxygen isotope scale and glacial event. *Quaternary Research*, 29, 1–24.