



HAL
open science

On Harnack inequalities and optimal transportation

Dominique Bakry, Ivan Gentil, Michel Ledoux

► **To cite this version:**

Dominique Bakry, Ivan Gentil, Michel Ledoux. On Harnack inequalities and optimal transportation. 2012. hal-00742654v4

HAL Id: hal-00742654

<https://hal.science/hal-00742654v4>

Preprint submitted on 25 Oct 2012 (v4), last revised 8 Jul 2013 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON HARNACK INEQUALITIES AND OPTIMAL TRANSPORTATION

D. BAKRY, I. GENTIL, M. LEDOUX

University of Toulouse and University of Lyon, France

Abstract. – We develop connections between Harnack inequalities for the heat flow of diffusion operators with curvature bounded from below and optimal transportation. Through heat kernel inequalities, a new isoperimetric-type Harnack inequality is emphasized. Commutation properties between the heat and Hopf-Lax semigroups are developed consequently, providing direct access to the heat flow contraction property along Wasserstein distances.

1. Introduction

Harnack inequalities classically provide strong tools towards regularity properties of solutions of partial differential equations and heat kernel bounds. A renowned result on the topic is the parabolic inequality by P. Li and S.-T. Yau [L-Y]

$$(1) \quad \frac{|\nabla P_t f|^2}{(P_t f)^2} - \frac{\Delta P_t f}{P_t f} \leq \frac{n}{2t}$$

for the heat semigroup $(P_t)_{t \geq 0}$ on an n -dimensional Riemannian manifold (M, g) with non-negative Ricci curvature, and every $t > 0$ and smooth positive function $f : M \rightarrow \mathbb{R}$. By integration along geodesics, it yields the Harnack inequality

$$(2) \quad P_t f(x) \leq P_{t+s} f(y) \left(\frac{t+s}{t} \right)^{n/2} e^{d^2/4s}$$

for $f : M \rightarrow \mathbb{R}$ non-negative and $t, s > 0$, where $d = d(x, y)$ is the Riemannian distance between $x, y \in M$. The results (1) and (2) admit versions for any lower bound on the Ricci curvature (cf. [L-Y], [D]). A heat flow proof of (1), in the spirit of the arguments developed in this work, has been provided in [B-L2].

In the context of diffusion operators, the Harnack inequality (2) may actually lose its relevance due to the infinite-dimensional feature of some models. Let $L = \Delta - \nabla V \cdot \nabla$ be a diffusion operator on a smooth complete connected Riemannian manifold (M, g) , where $V : M \rightarrow \mathbb{R}$ is a smooth potential, with associated Markov semigroup $(P_t)_{t \geq 0}$ and invariant and symmetric measure $d\mu = e^{-V} dx$ (where dx is the Riemannian volume

element). A notion of curvature-dimension $CD(\rho, m)$, $\rho \in \mathbb{R}$, $m \geq 1$, of such operators L has been introduced by D. Bakry and M. Émery [B-É] (cf. [B], [Ba-G-L]), under the name of Γ_2 criterion, through the Bochner-type inequality

$$(3) \quad \frac{1}{2} L(|\nabla f|^2) - \nabla f \cdot \nabla Lf \geq \rho |\nabla f|^2 + \frac{1}{m} (Lf)^2$$

for any smooth $f : M \rightarrow \mathbb{R}$. (The Γ_2 operator in this context is precisely the expression on the right-hand of (3).) For example, by the standard Bochner formula from Riemannian geometry, the Laplace operator Δ on an n -dimensional Riemannian manifold with Ricci curvature bounded below by ρ satisfies $CD(\rho, m)$ with $m \geq n$. On the other hand, on $M = \mathbb{R}^n$ with V the quadratic potential, the associated Ornstein-Uhlenbeck operator L is intrinsically of infinite dimension $m = \infty$ since (3) cannot hold for some $\rho \in \mathbb{R}$ with m finite. (It actually holds in this example with $\rho = 1$ and $m = \infty$.) In particular a Harnack inequality (2) cannot hold in this case, as well as in further similar infinite-dimensional models. Note that, when $m = \infty$, again by the Bochner formula, the curvature condition $CD(\rho, \infty)$, $\rho \in \mathbb{R}$, amounts to the local geometric lower bound

$$(4) \quad \text{Ric} + \text{Hess}(V) \geq \rho$$

(as symmetric matrices) uniformly over the manifold.

To circumvent the drawbacks attached to the case $m = \infty$, F.-Y. Wang introduced in [W1] (see also [W2]) a new form of Harnack inequalities for infinite dimensional diffusion operators of the type $L = \Delta - \nabla V \cdot \nabla$ (and more general ones). Namely, Wang's Harnack inequalities indicate that, under the curvature condition $CD(\rho, \infty)$ (equivalent to (4)), for every (smooth) non-negative function f on M , every $t > 0$, every $\alpha > 1$, and every $x, y \in M$,

$$(5) \quad (P_t f(x))^\alpha \leq P_t(f^\alpha)(y) e^{\alpha d^2/2(\alpha-1)\sigma(t)}$$

where $d = d(x, y)$ the Riemannian distance between x and y and where $\sigma(t) = \frac{1}{\rho} (e^{2\rho t} - 1)$ ($= 2t$ if $\rho = 0$). The proof of (5) is based on the interpolation

$$P_s((P_{t-s}f)^\alpha)(x_s), \quad s \in [0, t],$$

along a geodesic $(x_s)_{s \in [0, t]}$ joining x to y together with the commutation, for all $t \geq 0$ and smooth $g : M \rightarrow \mathbb{R}$,

$$(6) \quad |\nabla P_t g| \leq e^{-\rho t} P_t(|\nabla g|)$$

as an equivalent formulation of the curvature lower bound $CD(\rho, \infty)$. In a sense, the gradient bound (6) is the counterpart of the Li-Yau inequality (1) in this context.

In the limit as $\alpha \rightarrow \infty$, a log-Harnack inequality

$$(7) \quad P_t(\log f)(x) \leq \log P_t f(y) + \frac{d^2}{2\sigma(t)}$$

also holds (cf. [Bo-G-L], [W3]). It was further shown in [W1], [W3] that either (5) (for one $\alpha > 1$) or (7) implies back, as $t \rightarrow 0$, the curvature condition $CD(\rho, \infty)$ (in its infinitesimal form (4)).

The aim of this work is two-fold. We will first show how the previous infinite-dimensional Harnack inequalities may actually be seen as consequences of the Lipschitz property of a suitable functional of isoperimetric type. On the basis of this observation, we establish next a kind of isoperimetric-type Harnack inequality. These results naturally lead to develop connections between Harnack inequalities and mass transportation. In particular, we establish a commutation property between the heat semigroup and the Hopf-Lax infimum-convolution semigroup at the root of the Wasserstein contraction property along the heat flow.

Two observations are actually at the starting point of this work. For simplicity in the discussion below, let us restrict ourselves to $CD(0, \infty)$ ($\rho = 0$).

First, the gradient bound (6) (and thus the curvature condition $CD(0, \infty)$) is known to imply (to be equivalent) to logarithmic Sobolev inequalities under P_t , in particular in reverse form

$$(8) \quad t \frac{|\nabla P_t f|^2}{P_t f} \leq P_t(f \log f) - P_t f \log P_t f$$

for every smooth $f > 0$ and every $t > 0$ (cf. [Ba-G-L]). Inequalities like the preceding one are understood point-wise throughout this work. Now, as was noticed by M. Hino [H], the latter ensures that whenever $0 < f \leq 1$ and $\psi = (\log(1/P_t f))^{1/2}$, then

$$|\nabla \psi|^2 \leq \frac{1}{2t}.$$

In other words, ψ is Lipschitz with Lipschitz coefficient less than or equal to $(2t)^{-1/2}$. In particular, for every $x, y \in M$,

$$\sqrt{\log \frac{1}{P_t f(x)}} \leq \sqrt{\log \frac{1}{P_t f(y)}} + \frac{d}{\sqrt{2t}}$$

where we recall that $d = d(x, y)$. After some work, it may then be shown that for each $\varepsilon > 0$, there exists $C(\varepsilon) > 0$ such that

$$(P_t f(x))^2 \leq C(\varepsilon) P_t(f^2)(y) e^{d^2/4(1+\varepsilon)t},$$

that is as close as possible to (5) (for $\alpha = 2$).

It should be mentioned that it is precisely the dimensional version of the reverse logarithmic Sobolev inequality (8) which has been used in [B-L2] to provide a monotonicity proof of the Li-Yau parabolic inequality (1). We will exploit this information toward dimensional statements in Section 5 below. For further dimensional Harnack inequalities under the curvature-dimension condition $CD(\rho, m)$, comparing in particular different times, see [W4].

The second observation at the starting point of this investigation is the links between Harnack-type inequalities and optimal transportation already put forward in [Bo-G-L] where semigroup tools were developed towards a proof of the Otto-Villani HWI inequality [O-V] (cf. [V1], [V2]). We briefly recall the basic step. Namely, the log-Harnack inequality (7) may be translated equivalently as

$$(9) \quad P_t(\log f) \leq Q_{2t}(\log P_t f)$$

where $(Q_s)_{s>0}$ is the Hopf-Lax infimum-convolution semigroup

$$Q_s \varphi(x) = \inf_{y \in M} \left[\varphi(y) + \frac{d(x, y)^2}{2s} \right], \quad x \in M, \quad s > 0.$$

Assume now that μ is a probability measure and let $f > 0$ be a probability density with respect to μ . Then, by time reversibility and (9) applied to $P_t f$, $t > 0$,

$$\int_M P_t f \log P_t f d\mu = \int_M f P_t(\log P_t f) d\mu \leq \int_M f Q_{2t}(\log P_{2t} f) d\mu.$$

Now $\int_M \log P_{2t} f d\mu \leq 0$ by Jensen's inequality. Hence, combining with the scaling properties of $(Q_s)_{s>0}$,

$$\int_M P_t f \log P_t f d\mu \leq \frac{1}{2t} \left[\int_M Q_1 \varphi f d\mu - \int_M \varphi d\mu \right]$$

where $\varphi = 2t \log P_{2t} f$. By the Kantorovich-Rubinstein dual description of the quadratic Wasserstein distance $W_2(\nu, \mu)$ between the probability measures $d\nu = f d\mu$ and μ on M (cf. e.g. [V1]), the latter amounts to

$$(10) \quad \int_M P_t f \log P_t f d\mu \leq \frac{1}{2t} W_2^2(\nu, \mu).$$

Recall that the (quadratic) Wasserstein distance $W_2(\nu, \mu)$ is defined by

$$W_2(\nu, \mu) = \left(\int_{M \times M} \frac{1}{2} d(x, y)^2 d\pi(x, y) \right)^{1/2}$$

where the infimum is taken over all couplings π with respective marginals ν and μ .

Note that the preceding argument similarly yields, for every $t \geq 0$,

$$(11) \quad t \int_M P_t f \log P_t f d\mu \leq \frac{1}{2} W_2^2(f\mu, g\mu) + t \int_M g \log g d\mu$$

where g is a further probability density with respect to μ , and where, for simplicity here, $f\mu$ and $g\mu$ denote the probability measures $f d\mu$ and $g d\mu$. Indeed, write in the preceding notation that

$$t \int_M P_t f \log P_t f d\mu \leq \frac{1}{2} \left[\int_M Q_1 \varphi f d\mu - \int_M \varphi g d\mu \right] + t \int_M g \log P_{2t} f d\mu.$$

Since by convexity $\int_M g \log P_{2t} f d\mu \leq \int_M g \log g d\mu$, the claim follows.

Now (10) is actually the major step in the semigroup proof of the HWI inequality of [O-V] under the curvature condition $CD(0, \infty)$. Namely, the classical heat flow interpolation scheme (cf. [B], [Ba-G-L]) indicates that, for every suitable probability density $f : M \rightarrow \mathbb{R}$ and every $t \geq 0$,

$$\begin{aligned} \int_M f \log f d\mu - \int_M P_t f \log P_t f d\mu &= - \int_0^t \left(\frac{d}{ds} \int_M P_s f \log P_s f d\mu \right) ds \\ &= \int_0^t \int_M \frac{|\nabla P_s f|^2}{P_s f} d\mu ds. \end{aligned}$$

Since $|\nabla P_s f| \leq P_s(|\nabla f|)$ according to (6),

$$\frac{|\nabla P_s f|^2}{P_s f} \leq P_s \left(\frac{|\nabla f|^2}{f} \right)$$

by the Cauchy-Schwarz inequality along the Markov kernel P_s . Therefore,

$$\int_M f \log f d\mu \leq \int_M P_t f \log P_t f d\mu + t \int_M \frac{|\nabla f|^2}{f} d\mu.$$

Together thus with (10), optimization in $t > 0$ yields

$$\int_M f \log f d\mu \leq W_2(\nu, \mu) \left(2 \int_M \frac{|\nabla f|^2}{f} d\mu \right)^{1/2}$$

which is the announced HWI inequality, connecting Entropy, Wasserstein distance and Fisher Information. Similar arguments may be developed under $CD(\rho, \infty)$ for any $\rho \in \mathbb{R}$ to yield the full formulation of Otto-Villani's HWI inequality (cf. [Bo-G-L], [Ba-G-L]). Note that together with (11), the argument also recovers the known inequality (cf. e.g. [CE])

$$\int_M f \log f d\mu \leq W_2(f\mu, g\mu) \left(2 \int_M \frac{|\nabla f|^2}{f} d\mu \right)^{1/2} + \int_M g \log g d\mu$$

for probability densities f and g with respect to μ .

For the matter of comparison, it might be worthwhile mentioning that the recent Kuwada lemma (see [G-K-O]) develops similar arguments towards the inequality

$$(12) \quad W_2^2(P_t f\mu, f\mu) \leq \frac{t}{2} \left[\int_E f \log f d\mu - \int_E P_t f \log P_t f d\mu \right]$$

for any probability density f with respect to μ and any $t \geq 0$. Indeed, for $\varphi : E \rightarrow \mathbb{R}$ bounded and measurable,

$$\begin{aligned} \int_M Q_1 \varphi P_t f d\mu - \int_M \varphi f d\mu &= \int_0^1 \left(\frac{d}{ds} \int_M Q_s \varphi P_{st} f d\mu \right) ds \\ &= \int_0^1 \int_M \left[-\frac{1}{2} |\nabla Q_s \varphi|^2 P_{st} f + t Q_s \varphi \mathbf{L} P_{ts} f \right] d\mu ds \\ &= \int_0^1 \int_M \left[-\frac{1}{2} |\nabla Q_s \varphi|^2 P_{st} f - t \nabla Q_s \varphi \cdot \nabla P_{st} f \right] d\mu ds \end{aligned}$$

by the fact that the Hopf-Lax semigroup solves the standard Hamilton-Jacobi equation and by integration by parts. Next, by the Cauchy-Schwarz inequality,

$$\int_M Q_1 \varphi P_t f d\mu - \int_M \varphi f d\mu \leq \frac{t^2}{2} \int_0^1 \int_M \frac{|\nabla P_{st} f|^2}{P_{st} f} d\mu ds$$

which yields (12) by the Kantorovich-Rubinstein duality and integration.

With respect to (10), the Kuwada inequality (12) somewhat works in the other direction. The point is that while (10) leads to the HWI inequality, the Kuwada inequality (10) has been identified in [G-L] at the root of the Otto-Villani theorem [O-V] (cf. [Bo-G-L], [V1], [V2]) connecting logarithmic Sobolev inequalities to transportation cost inequalities.

On the basis of these two early observations, the purpose of this work is, as announced, to develop a synthetic and refined treatment of Harnack-type inequalities for diffusion operators with curvature bounded from below and of their connections with transportation cost inequalities. The various contributions of this work are summarized as follows.

In Section 2, we provide a direct treatment of Wang's Harnack inequalities (5) and (7) relying on an improved, isoperimetric-type version, of the reverse logarithmic Sobolev inequality (8) along the heat flow.

This reverse isoperimetric-type inequality in turn implies a new isoperimetric version of Harnack inequalities emphasized in Section 3. For example, under non-negative curvature, it yields that for any say closed set A in M , any $t > 0$ and any $x, y \in M$,

$$(13) \quad P_t(\mathbf{1}_A)(x) \leq P_t(\mathbf{1}_{A_d})(y)$$

where $d = d(x, y)$ and A_ε is the ε -neighborhood of A in the metric d . This result seems to be new even for the standard heat flow operator on a Riemannian manifold.

A direct consequence of (13) is the commutation

$$(14) \quad P_t(Q_s) \leq Q_s(P_t), \quad t, s > 0,$$

between the heat and Hopf-Lax semigroups (under non-negative curvature) which we emphasize in Section 4. This commutation was actually used earlier by K. Kuwada [K] in the study of the duality of gradient estimates at the root of the contraction property of the Wasserstein distance along the heat flow

$$W_2(\mu_t, \nu_t) \leq W_2(\mu_0, \nu_0)$$

where $d\mu_t = P_t f d\mu$ and $d\nu_t = P_t g d\mu$, $t \geq 0$, f, g probability densities with respect to μ , a result first established in [C-MC-V], [vR-S] and [O-W] (see also [W2]).

The commutation property (14) may actually be reached in several ways, and Section 5 presents a variety of methods depending on the underlying context, including the original approach of [K]. This section further includes dimensional version of the commutation property together with the corresponding Wasserstein contractions.

In the last Section 6, we briefly discuss some connections between the material presented here and recent developments, following [A-G-S3], around the Evolutionary Variational Inequality (EVI) expressing in the preceding notation that

$$W_2^2(\mu_t, \nu_0) + t \int_M P_t f \log P_t f d\mu \leq W_2^2(\mu_0, \nu_0) + t \int_M g \log g d\mu.$$

This property actually connects the Γ_2 Bakry-Émery $CD(\rho, \infty)$ curvature condition ([B-É], [B], [Ba-G-L]), expressed by the commutation (6), with the curvature bound in the sense of Lott-Villani-Sturm in metric measure spaces as convexity of relative entropy along the geodesics of optimal transportation ([L-V], [S1], [S2], [V2]). The recent main achievement by L. Ambrosio, N. Gigli and G. Savaré [A-G-S3] actually provides a link between the Γ_2 and Lott-Villani-Sturm curvature lower bounds in the class of the Riemannian energy measure spaces through the EVI. In Section 6, we sketch, following [A-G-S3], the principle of proof of the EVI in a smooth setting, for comparison with some of the tools developed here.

For simplicity in the exposition, the results of this work are presented in the weighted Riemannian setting, for thus diffusion operators $L = \Delta - \nabla \cdot \nabla V$ on a complete connected Riemannian manifold (M, g) with invariant and reversible measure $d\mu = e^{-V} dx$ (not necessarily a probability measure) where $V : M \rightarrow \mathbb{R}$ is a smooth potential. Integration by parts with respect to L is expressed by $\int_M f(-Lg) d\mu = \int_M \nabla f \cdot \nabla g d\mu$ for smooth functions $f, g : M \rightarrow \mathbb{R}$. The associated curvature condition $CD(\rho, \infty)$, $\rho \in \mathbb{R}$, is expressed equivalently by (4) as the infinitesimal version of the Bochner-type inequality (3). It amounts to the standard lower bound on the Ricci curvature for the Laplace operator Δ on (M, g) . The curvature condition $CD(\rho, \infty)$ is also equivalent to the gradient bound (6) which is, in an essential manner, the only way the curvature condition will be used throughout this work.

Most of the results below actually extend to the framework of a Markov diffusion triple (E, μ, Γ) in the sense of [B], [Ba-G-L], consisting of a state space E equipped with a diffusion semigroup $(P_t)_{t \geq 0}$ with infinitesimal generator L and carré du champ operator Γ and invariant and symmetric σ -finite measure μ . In the weighted Riemannian context, $\Gamma(f, f) = |\nabla f|^2$ for smooth functions. In this setting, the abstract curvature condition $CD(\rho, \infty)$, $\rho \in \mathbb{R}$, stems from the Bochner-type inequality (3) (with $m = \infty$) which is by now classically referred to as the Γ_2 criterion going back to [B-É] (see [B], [Ba-G-L]). The condition $CD(\rho, \infty)$ is equivalent to the gradient bound (6)

$$\sqrt{\Gamma(P_t f)} \leq e^{-\rho t} P_t(\sqrt{\Gamma(f)})$$

for every $t \geq 0$ and every f in a suitable algebra of functions. The state space E may be endowed with an intrinsic distance d for which Lipschitz functions f are such that $\Gamma(f)$ is bounded (μ -almost everywhere). We refer to [Ba-G-L] for details on this general setting. Note that at the level of the local inequalities along the semigroup, the setting covers the example of a generator $L = \Delta + Z$ for some smooth vector field Z on a manifold M as developed in [W1], [W2], [W3].

2. Reverse isoperimetry and Wang's Harnack inequalities

In this section, we address a direct proof of Wang's Harnack inequalities (5) and (7) along the Hino argument on the basis of a reinforced family of heat kernel inequalities first emphasized in [B-L1].

Denote by $I : [0, 1] \rightarrow \mathbb{R}_+$ the Gaussian isoperimetric function defined by $I = \varphi \circ \Phi^{-1}$ where

$$\Phi(x) = \int_{-\infty}^x e^{-u^2/2} \frac{du}{\sqrt{2\pi}}, \quad x \in \mathbb{R},$$

and $\varphi = \Phi'$. The function I satisfies the basic differential equality $II' = -1$.

The following statement, as a kind of reverse isoperimetric-type inequality, was first put forward in [B-L1]. We enclose a proof for completeness.

Proposition 2.1. *Under the curvature condition $CD(\rho, \infty)$ for some $\rho \in \mathbb{R}$, for every smooth function f on M with values in $[0, 1]$ and every $t \geq 0$,*

$$(15) \quad [I(P_t f)]^2 - [P_t(I(f))]^2 \geq \sigma(t) |\nabla P_t f|^2$$

where $\sigma(t) = \frac{1}{\rho} (e^{2\rho t} - 1)$ ($= 2t$ if $\rho = 0$).

Proof. By the heat flow interpolation, write

$$[I(P_t f)]^2 - [P_t(I(f))]^2 = - \int_0^t \frac{d}{ds} [P_s(I(P_{t-s} f))]^2 ds.$$

Now, by the chain rule for the diffusion operator L ,

$$\begin{aligned} - \frac{d}{ds} [P_s(I(P_{t-s} f))]^2 &= -2P_s(I(P_{t-s} f)) P_s \left(L I(P_{t-s} f) - I'(P_{t-s} f) L P_{t-s} f \right) \\ &= -2P_s(I(P_{t-s} f)) P_s (I''(P_{t-s} f) |\nabla P_t f|^2) \\ &= 2P_s(I(P_{t-s} f)) P_s \left(\frac{|\nabla P_t f|^2}{I(P_{t-s} f)} \right) \end{aligned}$$

where we used that $II' = -1$ in the last step. Since P_s is given by a kernel, it satisfies a Cauchy-Schwarz inequality, and hence

$$P_s(Y) P_s \left(\frac{X^2}{Y} \right) \geq [P_s(X)]^2, \quad X, Y \geq 0.$$

Hence, with $X = |\nabla P_t f|$ and $Y = I(P_{t-s} f)$,

$$[I(P_t f)]^2 - [P_t(I(f))]^2 \geq 2 \int_0^t [P_s(|\nabla P_{t-s} f|)]^2 ds.$$

By the gradient bound (6), $P_s(|\nabla g|) \geq e^{\rho s} |\nabla P_s g|$. With $g = P_{t-s} f$, it follows that

$$[I(P_t f)]^2 - [P_t(I(f))]^2 \geq 2 \int_0^t e^{2\rho s} ds |\nabla P_t f|^2$$

which is the result. \square

For the comparison with the Hino observation mentioned in the introduction, note that the inequality (15) of Proposition 2.1 implies the reverse logarithmic Sobolev inequality (8) by applying it to εf and letting $\varepsilon \rightarrow 0$.

As announced, we next show how Proposition 2.1 actually covers Wang's Harnack inequalities recalled in the introduction. Namely, in terms of gradient bounds, Proposition 2.1 indicates that for every (smooth) f with values in $[0, 1]$, $\Phi^{-1} \circ P_t f$ is $\sigma(t)^{-1/2}$ -Lipschitz, $t > 0$. In particular, for $x, y \in M$ and $d = d(x, y)$ the Riemannian distance between x and y ,

$$(16) \quad \Phi^{-1} \circ P_t f(x) \leq \Phi^{-1} \circ P_t f(y) + \frac{d}{\sqrt{\sigma(t)}}.$$

This Lipschitz property actually entails most of the inequalities emphasized in this work.

Towards a first illustration, set $\delta = d/\sqrt{\sigma(t)}$, so that (16) reads as

$$P_t f(x) \leq \Phi(\Phi^{-1} \circ P_t f(y) + \delta).$$

Apply now this inequality to $\mathbf{1}_{\{f \geq a\}}$, $a \geq 0$, for a non-negative (measurable) function f on M . Denoting by λ the distribution of f under P_t at the point y ,

$$P_t(\mathbf{1}_{\{f \geq a\}})(x) \leq \Phi\left(\Phi^{-1}(\lambda([a, \infty))) + \delta\right).$$

Integrating in $a \geq 0$ and using Fubini's theorem, denoting by $d\gamma(u) = e^{-|u|^2/2} \frac{du}{\sqrt{2\pi}}$ the standard Gaussian distribution on the line,

$$P_t f(x) \leq \int_0^\infty \int_{-\infty}^{\Phi^{-1}(\lambda([a, \infty))) + \delta} d\gamma(u) da = \int_{-\infty}^\infty \left(\int_0^\infty \mathbf{1}_{\{u \leq \Phi^{-1}(\lambda([a, \infty))) + \delta\}} da \right) d\gamma(u).$$

Change u into $u + \delta$ to get

$$P_t f(x) \leq e^{-\delta^2/2} \int_{-\infty}^\infty e^{-\delta u} \left(\int_0^\infty \mathbf{1}_{\{\Phi(u) \leq \lambda([a, \infty))\}} da \right) d\gamma(u).$$

Changing u into $-u$ and denoting by F the distribution function of λ , it follows that

$$P_t f(x) \leq e^{-\delta^2/2} \int_{-\infty}^\infty e^{\delta u} \left(\int_0^\infty \mathbf{1}_{\{F(a) \leq \Phi(u)\}} da \right) d\gamma(u).$$

After the further change of variables $v = \Phi(u)$,

$$P_t f(x) \leq e^{-\delta^2/2} \int_0^1 e^{\delta \Phi^{-1}(v)} \left(\int_0^\infty \mathbf{1}_{\{F(a) \leq v\}} da \right) dv.$$

The next statement summarizes the conclusion reached so far.

Theorem 2.2. *Under the curvature condition $CD(\rho, \infty)$ for some $\rho \in \mathbb{R}$, for every non-negative measurable function f on M , every $t > 0$ and every $x, y \in M$,*

$$P_t f(x) \leq e^{-\delta^2/2} \int_0^\infty e^{\delta \Phi^{-1} \circ F(s)} s dF(s)$$

where $\delta = d(x, y)/\sqrt{\sigma(t)}$ and F is the distribution function of f under P_t at the point y .

Theorem 2.2 appears at the root of the various Harnack inequalities in this context. It is however not expressed in a very tractable form. But it easily implies known ones. For example, by Cauchy-Schwarz,

$$\begin{aligned} \int_0^\infty e^{\delta \Phi^{-1} \circ F(s)} s dF(s) &\leq \left(\int_0^\infty e^{2\delta \Phi^{-1} \circ F(s)} dF(s) \right)^{1/2} \left(\int_0^\infty s^2 dF(s) \right)^{1/2} \\ &\leq e^{\delta^2} (P_t(f^2)(y))^{1/2} \end{aligned}$$

since

$$\int_0^\infty e^{2\delta \Phi^{-1} \circ F(s)} dF(s) = \int_0^1 e^{2\delta \Phi^{-1}(v)} dv = \int_{-\infty}^\infty e^{2\delta u} d\gamma(u) = e^{2\delta^2}.$$

The preceding therefore yields Wang's Harnack inequality (5) for $\alpha = 2$,

$$P_t f(x)^2 \leq P_t(f^2)(y) e^{d^2/\sigma(t)}.$$

By Hölder's inequality rather than Cauchy-Schwarz, one obtains the whole family of inequalities (5) with $\alpha > 1$. Using the entropic inequality yields similarly the log-Harnack inequality (7) (which is also obtained as $\alpha \rightarrow \infty$). With respect to Wang's original argument, the proof here avoids interpolation along geodesics (although the length space property is required to move from (15) to (16)).

3. Isoperimetric-type Harnack inequalities

As announced, the basic Lipschitz inequality (16) may be seen at the origin of a number of inequalities of interest, and this section develops further consequences in combination with isoperimetric bounds for heat kernel measures. To this task, recall first the isoperimetric comparison theorem for heat kernel measures under curvature bounds of [B-L1]. Recall I the Gaussian isoperimetric function.

Theorem 3.1. *Under the curvature condition $CD(\rho, \infty)$ for some $\rho \in \mathbb{R}$, for every smooth function f on M with values in $[0, 1]$ and every $t \geq 0$,*

$$I(P_t f) \leq P_t \left(\sqrt{I^2(f) + \rho(t) |\nabla f|^2} \right)$$

where $\rho(t) = \frac{1}{\rho} (1 - e^{-2\rho t})$ ($= 2t$ if $\rho = 0$).

As developed in [B-L1] (cf. also [Ba-G-L]), this result is an isoperimetric comparison theorem expressing that the isoperimetric profile of the heat kernel measures is bounded below by the Gaussian isoperimetric function (up to a scaling depending on t and ρ). More precisely, for any $y \in M$, any measurable set $A \subset M$, any $t > 0$ and any $\varepsilon > 0$,

$$(17) \quad P_t(\mathbf{1}_{A_\varepsilon})(y) \geq \Phi\left(\Phi^{-1}(P_t(\mathbf{1}_A)(y)) + \frac{\varepsilon}{\sqrt{\rho(t)}}\right)$$

where A_ε is the ε -neighborhood of A in the distance d .

Applied to $f = \mathbf{1}_A$, the Lipschitz property (16) ensures on the other hand that, for any measurable set $A \subset M$, and again with $\delta = d/\sqrt{\sigma(t)}$,

$$(18) \quad P_t(\mathbf{1}_A)(x) \leq \Phi\left(\Phi^{-1}(P_t(\mathbf{1}_A)(y)) + \delta\right).$$

The combination of (17) and (18) together with the fact that $\frac{\rho(t)}{\sigma(t)} = e^{-2\rho t}$ then yields the following isoperimetric-type Harnack inequality.

Theorem 3.2. *Under the curvature condition $CD(\rho, \infty)$ for some $\rho \in \mathbb{R}$, for every measurable set A in M , every $t \geq 0$ and every $x, y \in M$,*

$$P_t(\mathbf{1}_A)(x) \leq P_t(\mathbf{1}_{A_{d_t}})(y)$$

where $d_t = e^{-\rho t}d(x, y) = e^{-\rho t}d$. In particular, when $\rho = 0$,

$$P_t(\mathbf{1}_A)(x) \leq P_t(\mathbf{1}_{A_d})(y).$$

4. The commutation property and contraction in Wasserstein distance

The isoperimetric-type Harnack inequality of Theorem 3.2 has several consequences of interest in terms of commutation properties between the heat and the Hopf-Lax semigroups developed in this section.

To start with, let $f : M \rightarrow \mathbb{R}$ be Lipschitz with Lipschitz constant L . Then, if $A = \{f \geq a\}$ ($f \geq 0$ for simplicity), $A_{d_t} \subset \{f \geq a - Ld_t\}$. Therefore, by Theorem 3.2, under $CD(\rho, \infty)$, for every $x, y \in M$ and $t \geq 0$,

$$P_t(\mathbf{1}_{\{f \geq a\}})(x) \leq P_t(\mathbf{1}_{\{f + Ld_t \geq a\}})(y).$$

Integrating in $a \geq 0$,

$$P_t f(x) \leq P_t f(y) + Ld_t.$$

In other words, $P_t f$ is Lipschitz with Lipschitz constant less than $e^{-\rho t}L$, a result by M.-K. von Renesse and K.-Th. Sturm [vR-S].

The same argument may actually be used to reach a more precise property which in turn entails the contraction property of the heat flow with respect to Wasserstein metrics.

The key is expressed in the commutation between the heat and Hopf-Lax semigroups. Recall the Hopf-Lax infimum-convolution semigroup (cf. [E], [V1], [V2])

$$Q_s f(x) = \inf_{y \in M} \left[f(y) + \frac{d(x, y)^2}{2s} \right], \quad x \in M, \quad s > 0.$$

This property was actually used first by K. Kuwada [K] in the analysis of gradient bounds and Wasserstein contractions (see Corollary 4.2 below). The proof in [K], developed in the context of length spaces and actually for more general costs, relies on an interpolation along geodesics and the use of the Hamilton-Jacobi equation (see the first alternate proof in Section 5 below).

Theorem 4.1. *Under the curvature condition $CD(\rho, \infty)$ for some $\rho \in \mathbb{R}$, for any $t, s > 0$ and any measurable $f : M \rightarrow \mathbb{R}$,*

$$(19) \quad P_t(Q_s f) \leq Q_{e^{2\rho t}s}(P_t f).$$

Proof. Let f be non-negative on M . It is enough by homogeneity to consider $s = 1$. As above, apply Theorem 3.2 to $A = \{Q_1 f \geq a\}$ for every $a \geq 0$. Since $A_{d_t} \subset \{f + d_t^2/2 \geq a\}$,

$$P_t(\mathbf{1}_{\{Q_1 f \geq a\}})(x) \leq P_t(\mathbf{1}_{\{f + d_t^2/2 \geq a\}})(y)$$

for all $x, y \in M$ and $t \geq 0$. Integrating in $a \geq 0$ yields

$$P_t(Q_1 f)(x) \leq P_t f(y) + \frac{d_t^2}{2}.$$

Taking then the infimum in $y \in M$ yields the result by definition of the infimum-convolution Q_1 . \square

The infimum-convolution semigroup $(Q_s)_{s>0}$ being solution of the Hamilton-Jacobi equation $\partial_s u = -\frac{1}{2}|\nabla u|^2$ with initial condition $u(0, \cdot) = f$, the commutation property (19) implies by a Taylor expansion at $s = 0$ that $|\nabla P_t f|^2 \leq e^{-2\rho t} P_t(|\nabla f|^2)$ for every $t \geq 0$. This gradient bound, weaker than (6), is still equivalent to the curvature bound $CD(\rho, \infty)$, providing therefore a converse to Theorem 4.1. In particular also, the isoperimetric Harnack inequality from Theorem 3.2 is actually equivalent to the curvature condition $CD(\rho, \infty)$.

As announced, it immediately follows from the commutation property (19) of Theorem 4.1 that the Wasserstein distance W_2 is contractive along the semigroup $(P_t)_{t \geq 0}$, an observation due to K. Kuwada [K]. The Wasserstein contraction property in this context is going back to [vR-S], with a stochastic proof, and to [O-W], with an Eulerian point of view, in a Riemannian framework (further discussed below in Section 6). The proof presented here on the basis of Theorem 4.1 extends to the abstract Markov semigroup setting of [B], [Ba-G-L]. The measure μ is assumed here to be a probability measure.

Corollary 4.2. *Under the curvature condition $CD(\rho, \infty)$ for some $\rho \in \mathbb{R}$, for any $t \geq 0$,*

$$(20) \quad W_2(\mu_t, \nu_t) \leq e^{-2\rho t} W_2(\mu_0, \nu_0)$$

where $d\mu_t = P_t f d\mu$ and $d\nu_t = P_t g d\mu$ for probability densities f, g with respect to the probability measure μ .

Proof. For any bounded measurable $\varphi : M \rightarrow \mathbb{R}$, by time reversibility and the commutation property (19),

$$\begin{aligned} \int_M Q_1 \varphi P_t f d\mu - \int_M \varphi P_t g d\mu &= \int_M P_t(Q_1 \varphi) f d\mu - \int_M P_t \varphi g d\mu \\ &\leq \int_M Q_{e^{2\rho t}}(P_t \varphi) f d\mu - \int_M P_t \varphi g d\mu \\ &\leq e^{-2\rho t} \left[\int_M Q_1(e^{2\rho t} P_t \varphi) f d\mu - \int_M e^{2\rho t} P_t \varphi g d\mu \right] \\ &\leq e^{-2\rho t} W_2^2(\mu_0, \nu_0) \end{aligned}$$

where the last step follows from the Kantorovich-Rubinstein dual description of the Wasserstein distance W_2 . The proof is complete. \square

By adapting Theorem 4.1 to costs $d(x, y)^p$, the same argument works for any Wasserstein distance W_p , $1 \leq p < \infty$ (cf. [K]), extending the contraction property of Corollary 4.2 to this class. More general Wasserstein functionals associated to further transportation costs may be considered similarly. Note that one main conclusion of the work [vR-S] by M.-K. von Renesse and K.-Th. Sturm is the equivalence of the Wasserstein contraction of Corollary 4.2 with the curvature bound. In [K], K. Kuwada established the equivalence of the Wasserstein contraction property for the cost $d(x, y)^p$ with the bound (6) with power q on the gradient, where $q > 1$ and $p < \infty$ are dual exponents. Actually, reading backwards the proof of Corollary 4.2, the contraction property (20) indicates that for all $\varphi : M \rightarrow \mathbb{R}$ bounded and measurable,

$$\int_M P_t(Q_1 \varphi) f d\mu - \int_M P_t \varphi g d\mu \leq \int_M Q_1 \varphi P_t f d\mu - \int_M \varphi P_t g d\mu \leq e^{-2\rho t} W_2^2(\mu_0, \nu_0).$$

Now, if $x, y \in M$ and if f and g are densities with respect to μ such that $d\mu_0 = f d\mu$ and $d\nu_0 = g d\mu$ approach Dirac masses at x and y respectively (for example by heat kernel approximations), the preceding yields

$$P_t(Q_1 \varphi)(x) - P_t \varphi(y) \leq e^{-2\rho t} \frac{1}{2} d(x, y)^2,$$

that is exactly the commutation property (19). As we have seen, the latter ensures in turn the curvature condition $CD(\rho, \infty)$.

5. Alternate proofs of the commutation property

In this last section, we briefly outline alternate proofs of the basic commutation property (19) of Theorem 4.1. For simplicity in the notation and the exposition, we only consider $\rho = 0$ below. Each proof involves at some point specific properties and may thus be adapted to more general settings accordingly.

First alternate proof. This proof is the original argument by K. Kuwada [K]. It requires the use of geodesics and the Hopf-Lax formula as solution of the Hamilton-Jacobi equation. Consider

$$\phi(s) = P_t(Q_s f)(x_s), \quad 0 \leq s \leq 1,$$

where $(x_s)_{s \in [0,1]}$ is a constant speed curve joining $x_0 = y$ to $x_1 = x$ in M . Set as above $d = d(x, y)$. Then, by the gradient bound (6) under $CD(0, \infty)$,

$$\begin{aligned} \phi'(s) &= P_t \left(-\frac{1}{2} |\nabla Q_s f|^2 \right) (x_s) + \nabla P_t(Q_s f)(x_s) \cdot \dot{x}_s \\ &\leq P_t \left(-\frac{1}{2} |\nabla Q_s f|^2 \right) (x_s) + d |\nabla P_t(Q_s f)(x_s)| \\ &\leq P_t \left(-\frac{1}{2} |\nabla Q_s f|^2 \right) + d P_t(|\nabla Q_s f|) \\ &\leq \frac{d^2}{2}. \end{aligned}$$

Hence

$$P_t(Q_1 f)(x) - P_t f(y) = \phi(1) - \phi(0) = \int_0^1 \phi'(s) ds \leq \frac{d^2}{2}$$

which is the result.

Second alternate proof. This second alternate proof also uses the Hopf-Lax infimum-convolution semigroup as solution of the Hamilton-Jacobi equation, and relies on the hypercontractivity property along the heat flow recently put forward in [B-B-G]. Namely, by the log-Harnack inequality (7), for every $v > 0$,

$$P_t(Q_1 f) \leq \frac{1}{v} Q_{2t}(\log P_t(e^{vQ_1 f})).$$

Under non-negative curvature, it is shown in [B-B-G] that for every $\psi : M \rightarrow \mathbb{R}$ and $t > 0$,

$$\log P_t(e^{Q_{2t}\psi}) \leq P_t \psi.$$

With $v = 1/2t$ and $\psi = f/2t$, the conclusion immediately follows by homogeneity of the infimum-convolutions.

Third alternate proof. This proof may be obtained by linear approximations of the Hamilton-Jacobi equation (vanishing viscosity method) along the lines of [Bo-G-L]. Following the notation therein, let for every $\varepsilon > 0$, the approximated Hopf-Lax semigroup

$$Q_t^\varepsilon f = -2\varepsilon \log P_t(e^{-f/2\varepsilon})$$

solution of the equation

$$\partial_t u = \varepsilon Lu - \frac{1}{2} |\nabla u|^2.$$

Moreover, $\lim_{\varepsilon \rightarrow 0} Q_t^\varepsilon f = Q_t f$. Dealing with

$$\phi(s) = P_s(Q_1^\varepsilon(P_{t-s}f)), \quad 0 \leq s \leq t,$$

shows that

$$\phi'(s) = 2\varepsilon P_s \left(\frac{1}{P_\varepsilon g} \left[\frac{|\nabla P_\varepsilon g|^2}{P_\varepsilon g} - P_\varepsilon \left(\frac{|\nabla g|^2}{g} \right) \right] \right)$$

where $g = e^{-P_{t-s}f/2\varepsilon}$. Under the gradient bound (6), $\phi'(s) \leq 0$ which yields that

$$P_t(Q_1^\varepsilon f) \leq Q_1^\varepsilon(P_t f).$$

In the limit as $\varepsilon \rightarrow 0$, the announced commutation property follows.

One benefit of the third alternate proof is that it may be developed similarly on the reinforced curvature-dimension condition $CD(0, m)$ with a finite-dimensional parameter m , for example with $m = n$ for the Laplace operator on an n -dimensional Riemannian manifold with non-negative Ricci curvature (cf. [B], [Ba-G-L]). We sketch the argument. The local logarithmic Sobolev inequalities of [B-L2] (see also [Ba-G-L]) under $CD(0, m)$ ensure after linearization that, for any $t > 0$, any non-negative smooth function $g : M \rightarrow \mathbb{R}$ and any $c > 0$,

$$c \frac{|\nabla P_t g|^2}{P_t g} - P_t \left(\frac{|\nabla g|^2}{g} \right) \leq (c-1)P_t(Lg) + \frac{m}{2t} (\sqrt{c}-1)^2 P_t g.$$

Arguing as previously in the $CD(0, \infty)$ case yields that for any f and $t, s > 0$,

$$P_t(Q_1^\varepsilon f) \leq Q_1^\varepsilon(P_s f) + m(\sqrt{t} - \sqrt{s})^2$$

and similarly in the limit as $\varepsilon \rightarrow 0$.

Applied to the Wasserstein contraction, the latter shows that, under $CD(0, m)$ and in the notation of Corollary 4.2,

$$(21) \quad W_2^2(\mu_t, \nu_s) \leq W_2^2(\mu_0, \nu_0) + m(\sqrt{t} - \sqrt{s})^2.$$

This inequality covers (20), however only when $s = t$. Note that when $s = 0$ and $\mu_t = \nu_t$, then $W_2(\mu_t, \mu_0) \leq \sqrt{mt}$ which describes a classical behavior of Brownian motion in Euclidean space. Further Wasserstein contraction properties under curvature-dimension condition are emphasized in [W4].

It should be mentioned in addition that, following the argument at the end of Section 4, the contraction inequality implies back the commutation

$$(22) \quad P_t(Q_1 f) \leq Q_1(P_s f) + m(\sqrt{t} - \sqrt{s})^2$$

for all $f : M \rightarrow \mathbb{R}$ and $t, s > 0$. Now, given $a \in \mathbb{R}$ to be specified, set $t = (1 + \varepsilon a)s$ for $\varepsilon > 0$ small enough. By homogeneity, the latter yields

$$P_{(1+a\varepsilon)s}(Q_\varepsilon f) \leq Q_\varepsilon(P_s f) + \frac{ma^2\varepsilon s}{(\sqrt{1+a\varepsilon}+1)^2}.$$

A Taylor expansion at $\varepsilon = 0$ then shows that

$$asP_s(Lf) - \frac{1}{2}P_s(|\nabla f|^2) \leq -\frac{1}{2}|\nabla P_s f|^2 + \frac{ma^2s}{4}.$$

For $a = \frac{2}{m}LP_s f$, it follows that

$$|\nabla P_s f|^2 \leq P_s(|\nabla f|^2) - \frac{2s}{m}(LP_s f)^2.$$

This inequality, holding (pointwise) for every (smooth) f and every $s > 0$, is known to be equivalent to the curvature-dimension condition $CD(0, m)$ (cf. (12) in [B-L2]). As such, the Wasserstein contraction (21), as well as the dimensional commutation (22), are also equivalent to $CD(0, m)$.

6. Links with the Evolutionary Variational Inequality

To conclude this work, we briefly describe some of the connections between the preceding material and recent contributions around the so-called Evolutionary Variational Inequality (EVI). As mentioned in the introduction, the EVI actually connects the curvature condition $CD(\rho, \infty)$ in the sense of the Γ_2 operator of [B-É] (see [B], [Ba-G-L]), expressed here through the commutation (6), with the curvature bound in the sense of Lott-Villani-Sturm in metric measure spaces as convexity of relative entropy along the geodesics of optimal transportation ([L-V], [S1], [S2], [V2]). See [A-G-S2], [A-G-S3].

The purpose of this short paragraph is to describe the main ideas at the root of the EVI following the recent main development [A-G-S3] by L. Ambrosio, N. Gigli and G. Savaré. In this work, the authors actually establish the EVI in the extended class of Riemannian energy measure spaces, providing there a complete link between the Bakry-Émery Γ_2 and Lott-Villani-Sturm curvatures (the implication from Lott-Villani-Sturm to Γ_2 was achieved in [A-G-S2]). With respect to [A-G-S3], we only concentrate here on the main principle of proof in the simplified framework of weighted Riemannian manifolds, the main achievement of [A-G-S3] being actually to perform the argument in a much larger class of non-smooth spaces together with a rather involved analysis. The guideline of this investigation is the Eulerian approach of [O-W] and [D-S] but the non-smooth structure causes a lot of technical problems. We nevertheless found it useful to outline the argument, avoiding all the regularity issues, in the context of this paper, to illustrate the general principle and the links with the material of the previous sections. We of course refer to [A-G-S3] for a complete rigorous investigation.

For simplicity thus, we deal with the weighted Riemannian framework of the preceding sections with $d\mu = e^{-V}dx$ a probability measure, and restrict ourselves to the

non-negative curvature assumption $CD(0, \infty)$ expressed in the form of the commutation property (6) with $\rho = 0$. The case of arbitrary $\rho \in \mathbb{R}$ is easily adapted along the same lines (cf. [A-G-S3]). Recall that we are using the cost $\frac{1}{2}d(x, y)^2$ in the Wasserstein distance W_2 .

Let f and g be probability densities with respect to the probability measure μ . The Evolutionary Variational Inequality (EVI) indicates that under $CD(0, \infty)$, for any $t > 0$,

$$(23) \quad W_2^2(\mu_t, \nu_0) + t \int_M P_t f \log P_t f d\mu \leq W_2^2(\mu_0, \nu_0) + t \int_M g \log g d\mu$$

where $d\mu_t = P_t f d\mu$, $d\nu_t = P_t g d\mu$. In the limit as $t \rightarrow 0$, together with the semigroup property,

$$(24) \quad \frac{d}{dt} W_2^2(\mu_t, \nu_0) \leq \int_M g \log g d\mu - \int_M P_t f \log P_t f d\mu$$

(the derivative being understood in an extended sense).

The material described in the preceding sections gets close to (23), however not quite. Indeed, the conjunction of (11) and of the Wasserstein contraction (20) (for $\rho = 0$) yields

$$W_2^2(\mu_t, \nu_t) + t \int_M P_t f \log P_t f d\mu \leq \frac{3}{2} W_2^2(\mu_0, \nu_0) + t \int_M g \log g d\mu$$

which is not directly comparable to (23) but which, in any case, is useless in the limit as $t \rightarrow 0$. To reach EVI, more on optimal transportation is actually required.

One key step in this regard is the existence of curves of probability densities h_s , $s \in [0, 1]$, with respect to μ interpolating between $h_0 = g$ and $h_1 = f$, assumed to be smooth both in space and s , such that for every smooth function ψ on M ,

$$(25) \quad \int_M \dot{h}_s \psi d\mu \leq W_2^2(\mu_0, \nu_0) + \frac{1}{2} \int_M |\nabla \psi|^2 h_s d\mu.$$

Such curves are naturally provided by optimal transportation, and arise for example in the Benamou-Brenier dynamical description of the Wasserstein distance [B-B] (cf. [A-G-S1], [Vi1], [Vi2]). To illustrate at a mild level such curves, and in $M = \mathbb{R}^n$ for the simplicity of the notation (the manifold case being similar at the expense of further Riemannian technology, cf. [V2]), consider the Brenier map $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ pushing forward $d\mu_0 = f d\mu$ to $d\nu_0 = g d\mu$ and providing optimal transportation in the sense of the Wasserstein distance W_2 as

$$(26) \quad W_2^2(\mu_0, \nu_0) = \frac{1}{2} \int_{\mathbb{R}^n} |x - T(x)|^2 f(x) d\mu(x)$$

(cf. [A-G-S1], [V1], [V2]...). Consider then the geodesics $T_s = s \text{Id} + (1-s)T$, $s \in [0, 1]$, of optimal transportation. If h_s denotes the density with respect to μ of the pushforward measure of $d\mu_0 = f d\mu$ by T_s (so that $h_0 = g$ and $h_1 = f$) assumed to be smooth both in space and s , it is easily checked that for every smooth function ψ on \mathbb{R}^n ,

$$\int_{\mathbb{R}^n} \dot{h}_s \psi d\mu = \int_{\mathbb{R}^n} (x - T(x)) \cdot \nabla \psi(T_s(x)) f(x) d\mu(x)$$

yielding (25) by the quadratic inequality and (26). Of course, the existence of such curves h_s , $s \in [0, 1]$, satisfying (25) in a non-smooth setting is a delicate issue carefully investigated in [A-G-S3].

On the basis of (25), the EVI (23) is then analyzed by a suitable coupling between the heat kernel and optimal transportation parametrizations. Precisely, the expressions

$$(27) \quad \int_M Q_1 \varphi P_t f d\mu - \int_M \varphi g d\mu + t \left(\int_M P_t f \log P_t f d\mu - \int_M g \log g d\mu \right)$$

for any smooth φ on M may be represented as

$$\int_0^1 \left(\frac{d}{ds} \int_M Q_s \varphi P_{st} h_s d\mu + t \frac{d}{ds} \int_M P_{st} h_s \log P_{st} h_s d\mu \right) ds.$$

Now, again under suitable smoothness assumptions not detailed here, by the Hamilton-Jacobi equation and integration by parts,

$$\begin{aligned} \frac{d}{ds} \int_M Q_s \varphi P_{st} h_s d\mu &= -\frac{1}{2} \int_M |\nabla Q_s \varphi|^2 P_{st} h_s d\mu + \int_M \dot{h}_s P_{st} (Q_s \varphi) d\mu \\ &\quad + t \int_M Q_s \varphi \mathbb{L} P_{st} h_s d\mu \\ &= -\frac{1}{2} \int_M |\nabla Q_s \varphi|^2 P_{st} h_s d\mu + \int_M \dot{h}_s P_{st} (Q_s \varphi) d\mu \\ &\quad - t \int_M \nabla Q_s \varphi \cdot \nabla P_{st} h_s d\mu. \end{aligned}$$

On the other hand,

$$\begin{aligned} \frac{d}{ds} \int_M P_{st} h_s \log P_{st} h_s d\mu &= \int_M [P_{st} \dot{h}_s + t \mathbb{L} P_{st} h_s] \log P_{st} h_s d\mu \\ &= \int_M P_{st} \dot{h}_s \log P_{st} h_s d\mu - t \int_M \nabla P_{st} h_s \cdot \nabla (\log P_{st} h_s) d\mu \end{aligned}$$

where we used that $\frac{d}{ds} P_{st} h_s = P_{st} \dot{h}_s + t \mathbb{L} P_{st} h_s$ and $\int_M P_{st} \dot{h}_s d\mu = \int_M \dot{h}_s d\mu = 0$.

From these expressions, it is easily checked that the sum

$$\frac{d}{ds} \int_M Q_s \varphi P_{st} h_s d\mu + t \frac{d}{ds} \int_M P_{st} h_s \log P_{st} h_s d\mu$$

may be rearranged as

$$\begin{aligned} -\frac{1}{2} \int_M |\nabla (Q_s \varphi + t \log P_{st} h_s)|^2 P_{st} h_s d\mu - \frac{t^2}{2} \int_M \frac{|\nabla P_{st} h_s|^2}{P_{st} h_s} d\mu \\ + \int_M \dot{h}_s P_{st} (Q_s \varphi + t \log P_{st} h_s) d\mu. \end{aligned}$$

Forgetting the term $\frac{t^2}{2} \int_M \frac{|\nabla P_{st} h_s|^2}{P_{st} h_s} d\mu$ (which is anyway of the order $o(t)$ in the limit (24)), this quantity is upper-bounded by

$$-\frac{1}{2} \int_M P_{st} (|\nabla (Q_s \varphi + t \log P_{st} h_s)|^2) h_s d\mu + \int_M \dot{h}_s P_{st} (Q_s \varphi + t \log P_{st} h_s) d\mu$$

where we used also symmetry of the semigroup. Now, by the curvature condition in the form of the commutation (6), the latter is further upper-bounded by

$$-\frac{1}{2} \int_M |\nabla P_{st}(Q_s \varphi + t \log P_{st} h_s)|^2 h_s d\mu + \int_M \dot{h}_s P_{st}(Q_s \varphi + t \log P_{st} h_s) d\mu$$

With $\psi = P_{st}(Q_s \varphi + t \log P_{st} h_s)$, (25) implies that this expression is precisely bounded above by $W_2^2(\mu_0, \nu_0)$. Integrating in s from 0 to 1 and taking the supremum over all φ then yields the announced EVI (23).

As mentioned above, the preceding argument is inspired by the Eulerian calculus developed by F. Otto and M. Westdickenberg [O-W] in their approach of the contraction property (20). Namely, if the parametrization does not involve the heat flow, consider for $\varphi : M \rightarrow \mathbb{R}$ smooth enough,

$$\int_M Q_1 \varphi P_t f d\mu - \int_M \varphi P_t g d\mu = \int_0^1 \left(\frac{d}{ds} \int_M Q_s \varphi P_t h_s d\mu \right) ds.$$

Since, as above,

$$\frac{d}{ds} \int_M Q_s \varphi P_t h_s d\mu = -\frac{1}{2} \int_M |\nabla Q_s \varphi|^2 P_t h_s d\mu + \int_M \dot{h}_s P_t(Q_s \varphi) d\mu,$$

by time reversibility and the gradient bound (6),

$$\begin{aligned} \frac{d}{ds} \int_M Q_s \varphi P_t h_s d\mu &= -\frac{1}{2} \int_M P_t(|\nabla Q_s \varphi|^2) h_s d\mu + \int_M \dot{h}_s P_t(Q_s \varphi) d\mu \\ &\leq -\frac{1}{2} \int_M |\nabla P_t(Q_s \varphi)|^2 h_s d\mu + \int_M \dot{h}_s P_t(Q_s \varphi) d\mu. \end{aligned}$$

Using (25) then yields

$$\int_M Q_1 \varphi P_t f d\mu - \int_M \varphi P_t g d\mu \leq W_2^2(\mu_0, \nu_0),$$

that is, after taking the supremum in φ , the contraction property (20) of Corollary 4.2.

Acknowledgement. We are grateful to L. Ambrosio, N. Gigli and G. Savaré for helpful discussions on the EVI and for pointing out relevant references, and to A. Guillin for sharing with us his simple proof from the Wasserstein contraction to the curvature condition at the end of Section 4.

REFERENCES

- [A-G-S1] L. AMBROSIO, N. GIGLI, G. SAVARÉ. Gradient flows in metric spaces and in the space of probability measures. Lectures in Mathematics ETH Zürich. Birkhäuser (2005).
- [A-G-S2] L. AMBROSIO, N. GIGLI, G. SAVARÉ. Metric measures spaces with Riemannian Ricci curvature bounded from below (2011).
- [A-G-S3] L. AMBROSIO, N. GIGLI, G. SAVARÉ. Bakry-Émery curvature-dimension condition and Riemannian Ricci curvature bounds (2012).
- [B] D. BAKRY. L'hypercontractivité et son utilisation en théorie des semigroupes. Ecole d'Eté de Probabilités de Saint-Flour. Lecture Notes in Math. 1581, 1–114 (1994). Springer.
- [B-B-G] D. BAKRY, F. BOLLEY, I. GENTIL. Dimension dependent hypercontractivity for Gaussian kernels. Probab. Theory Related Fields, to appear (2012).
- [B-É] D. BAKRY, M. ÉMERY. Diffusions hypercontractives. Séminaire de Probabilités XIX, 1983/84, Lecture Notes in Math. 1123, 177–206. Springer (1985).
- [Ba-G-L] D. BAKRY, I. GENTIL, M. LEDOUX. Analysis and geometry of Markov diffusion operators. Forthcoming monograph (2013).
- [B-L1] D. BAKRY, M. LEDOUX. Lévy-Gromov's isoperimetric inequality for an infinite-dimensional diffusion generator. Invent. Math. 123, 259–281 (1996).
- [B-L2] D. BAKRY, M. LEDOUX. A logarithmic Sobolev form of the Li-Yau parabolic inequality. Rev. Mat. Iberoam. 22, 683–702 (2006).
- [B-B] J.-D. BENAMOU, Y. BRENIER. A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem. Numer. Math. 84, 375–393 (2000).
- [Bo-G-L] S. BOBKOV, I. GENTIL, M. LEDOUX. Hypercontractivity of Hamilton-Jacobi equations. J. Math. Pures Appl. 80, 669–696 (2001).
- [C-MC-V] J. CARRILLO, R. MCCANN, C. VILLANI. Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates. Rev. Mat. Iberoamericana 19, 971–1018 (2003).
- [CE] D. CORDERO-ERAUSQUIN. Some applications of mass transport to Gaussian-type inequalities. Arch. Ration. Mech. Anal. 161, 257–269 (2002).
- [D-S] S. DANERI, G. SAVARÉ. Eulerian calculus for the displacement convexity in the Wasserstein distance. SIAM J. Math. Anal. 40, 1104–1122 (2008).
- [D] E. B. DAVIES. Heat kernels and spectral theory. Cambridge Tracts in Mathematics 92. Cambridge (1989).
- [E] L. EVANS. Partial differential equations. Graduate Studies in Mathematics 19. American Mathematical Society (1998).
- [G-K-O] N. GIGLI, K. KUWADA, S.-I. OHTA. Heat flow on Alexandrov spaces. Comm. Pure Appl. Math., to appear (2012).
- [G-L] N. GIGLI, M. LEDOUX. From log Sobolev to Talagrand: a quick proof. Discrete and Continuous Dynamical Systems, to appear (2012).
- [H] M. HINO. On short time asymptotic behavior of some symmetric diffusions on general state spaces. Potential Anal. 16, 249–264 (2002).
- [K] K. KUWADA. Duality on gradient estimates and Wasserstein controls. J. Funct. Anal. 258, 3758–3774 (2010).
- [L-Y] P. LI, S.-T. YAU. On the parabolic kernel of the Schrödinger operator. Acta Math. 156, 153–201 (1986).

- [L-V] J. LOTT, C. VILLANI. Ricci curvature for metric-measure spaces via optimal transport. *Ann. Math.* 169, 903–991 (2009).
- [O-V] F. OTTO, C. VILLANI. Generalization of an inequality by Talagrand, and links with the logarithmic Sobolev inequality. *J. Funct. Anal.* 173, 361–400 (2000).
- [O-W] F. OTTO, M. WESTDICKENBERG. Eulerian calculus for the contraction in the Wasserstein distance. *SIAM J. Math. Anal.* 37, 1227–1255 (2005).
- [vR-S] M.-K. VON RENESSE, K.-T. STURM. Transport inequalities, gradient estimates, entropy and Ricci Curvature. *Comm. Pure Appl. Math.* 68, 923–940 (2005).
- [S1] K.-T. STURM. On the geometry of metric measure spaces I. *Acta Math.* 196, 65–131 (2006).
- [S1] K.-T. STURM. On the geometry of metric measure spaces I. *Acta Math.* 196, 133–177 (2006).
- [V1] C. VILLANI. Topics in optimal transportation. Graduate Studies in Mathematics 58. American Mathematical Society (2003).
- [V2] C. VILLANI. Optimal transport. Old and new. *Grundlehren der Mathematischen Wissenschaften* 338. Springer (2009).
- [W1] F.-Y. WANG. Logarithmic Sobolev inequalities on noncompact Riemannian manifolds. *Probab. Theor. Rel. Fields* 109, 417–424 (1997).
- [W2] F.-Y. WANG. Functional inequalities, Markov properties and spectral theory. Science Press (2005).
- [W3] F.-Y. WANG. Harnack inequalities on manifolds with boundary and applications. *J. Math. Pures Appl.* 94, 304–321 (2010).
- [W4] F.-Y. WANG. Equivalent semigroup properties for the curvature-dimension condition. *Bull. Sci. Math.* 135, 803–815 (2011).

D. B., M. L.: INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UNIVERSITÉ DE TOULOUSE, 31062 TOULOUSE, FRANCE, AND INSTITUT UNIVERSITAIRE DE FRANCE, bakry, ledoux@math.univ-toulouse.fr

I. G.: LABORATOIRE CAMILLE JORDAN, UNIVERSITÉ DE LYON, 69622 LYON, FRANCE, gentil@math.univ-lyon1.fr