



**HAL**  
open science

# Complements to the paper " $W^{1,1}$ -maps with values into $S^1$ "

Haïm Brezis, Petru Mironescu, Augusto C. Ponce

► **To cite this version:**

Haïm Brezis, Petru Mironescu, Augusto C. Ponce. Complements to the paper " $W^{1,1}$ -maps with values into  $S^1$ ". 2004. hal-00747667

**HAL Id: hal-00747667**

**<https://hal.science/hal-00747667>**

Preprint submitted on 31 Oct 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**COMPLEMENTS TO THE PAPER**  
**“ $W^{1,1}$ -MAPS WITH VALUES INTO  $S^1$ ”**  
**07/05/04**

HAÏM BREZIS<sup>(1),(2)</sup>, PETRU MIRONESCU<sup>(3)</sup> AND AUGUSTO C. PONCE<sup>(1),(2)</sup>

The purpose of these notes is to complement some of our results in [BMP]. We also establish some of the claims we stated there without proof.

**A. Extending Theorem 10 to other seminorms in  $W^{1,1}(\Omega; S^1)$ .**

In view of Theorem 10, it is natural to introduce the following quantity

$$\rho(P, N) = \frac{1}{2\pi} \text{Inf} \left\{ [g]_{W^{1,1}} ; g \in W^{1,1}(\Omega; S^1), T(g) = 2\pi(\delta_P - \delta_N) \right\}.$$

Here,  $[ \ ]_{W^{1,1}}$  is a general given semi-norm on  $W^{1,1}(\Omega; \mathbb{R}^2)$  equivalent to  $| \ |_{W^{1,1}}$ . We require from  $[ \ ]_{W^{1,1}}$  some structural properties :

(P1)  $[\alpha g]_{W^{1,1}} = [g]_{W^{1,1}}, \forall g \in W^{1,1}(\Omega; \mathbb{R}^2), \forall \alpha \in S^1 ;$

(P2)  $[\bar{g}]_{W^{1,1}} = [g]_{W^{1,1}}, \forall g \in W^{1,1}(\Omega; \mathbb{R}^2) ;$

(P3)  $[gh]_{W^{1,1}} \leq \|g\|_{L^\infty} [h]_{W^{1,1}} + \|h\|_{L^\infty} [g]_{W^{1,1}}, \forall g, h \in W^{1,1}(\Omega; \mathbb{R}^2) \cap L^\infty.$

It follows from (P3) that  $\rho$  is a distance.

[We refer the reader to [Mi] for the solution of the exercise given at the end of Section 3.]

Alternatively, we may define  $\rho$  starting from maps in  $\mathcal{R}$  :

**Lemma A1.** *We have*

$$\rho(P, N) = \frac{1}{2\pi} \text{Inf} \left\{ [g]_{W^{1,1}} \left| \begin{array}{l} g \in C^\infty(\Omega \setminus \{P, N\}; S^1) \cap W^{1,1}, \\ \deg(g, P) = +1, \deg(g, N) = -1 \end{array} \right. \right\}.$$

*Proof.* It suffices to prove that, for  $g \in W^{1,1}(\Omega; S^1)$  such that  $T(g) = 2\pi(\delta_P - \delta_N)$ , we may find a sequence  $(g_n) \subset \mathcal{R}$  such that  $T(g_n) = 2\pi(\delta_P - \delta_N)$  and  $g_n \rightarrow g$

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

in  $W^{1,1}$ . Note that the desired conclusion is invariant with respect to orientation-preserving diffeomorphisms of  $\Omega$ . Therefore, we may assume that  $\Omega = S^2$  and that  $P, N$  are respectively the North and the South pole of  $S^2$ . Let  $h(x, y, z) = \frac{(x, y)}{|(x, y)|}$ . Then  $h \in \mathcal{R}$ ,  $T(h) = 2\pi(\delta_P - \delta_N)$ . It follows that  $k = g\bar{h} \in Y$ . Thus, we may find a sequence  $(k_n) \subset C^\infty(\Omega; S^1)$  such that  $k_n \rightarrow k$  in  $W^{1,1}$ . Set  $g_n = hk_n$ . Then  $g_n \in \mathcal{R}$ ,  $T(g_n) = 2\pi(\delta_P - \delta_N)$  and  $g_n \rightarrow g$  in  $W^{1,1}$ .

Another simple property of  $\rho$  is

$$(A1) \quad C_1 d(P, N) \leq \rho(P, N) \leq C_2 d(P, N)$$

for some  $C_1, C_2 > 0$  independent of  $P, N$ . This follows from the equivalence of  $|\cdot|_{W^{1,1}}$  and  $[\cdot]_{W^{1,1}}$ .

Part of Theorem 10 holds in this more general setting :

**Theorem A1.** *Let  $P_i, N_i \in \Omega$  be such that*

$$\sum \rho(P_i, N_i) < \infty \quad (\text{or, equivalently, } \sum d(P_i, N_i) < \infty).$$

Set

$$\tilde{L} = \text{Inf} \left\{ \sum \rho(\tilde{P}_j, \tilde{N}_j) ; \sum (\delta_{\tilde{P}_j} - \delta_{\tilde{N}_j}) = \sum (\delta_{P_i} - \delta_{N_i}) \right\}.$$

Then

$$\frac{1}{2\pi} \text{Inf} \left\{ [g]_{W^{1,1}} ; g \in W^{1,1}(\Omega; S^1), T(g) = 2\pi \sum (\delta_{P_i} - \delta_{N_i}) \right\} \leq \tilde{L}.$$

*Proof.* Let  $\varepsilon > 0$  and let  $\tilde{P}_j, \tilde{N}_j$  be such that

$$\sum (\delta_{\tilde{P}_j} - \delta_{\tilde{N}_j}) = \sum (\delta_{P_i} - \delta_{N_i}) \quad \text{and} \quad \sum \rho(\tilde{P}_j, \tilde{N}_j) \leq \tilde{L} + \varepsilon.$$

By definition, for each  $j$  we may find some  $g_j \in W^{1,1}(\Omega; S^1)$  such that

$$T(g_j) = 2\pi(\delta_{\tilde{P}_j} - \delta_{\tilde{N}_j})$$

and

$$[g_j]_{W^{1,1}} \leq 2\pi\rho(\tilde{P}_j, \tilde{N}_j) + \frac{\varepsilon}{2^j}.$$

We claim that there is a sequence  $k_n \rightarrow \infty$  such that  $\prod_{j=1}^{k_n} g_j \xrightarrow{n} g$  in  $W^{1,1}$  for some  $g \in W^{1,1}(\Omega; S^1)$ . By Lemma 1, this implies that

$$T(g) = 2\pi \sum_j (\delta_{\tilde{P}_j} - \delta_{\tilde{N}_j}) = 2\pi \sum_i (\delta_{P_i} - \delta_{N_i}).$$

Using (P3), we will also have  $[g]_{W^{1,1}} \leq 2\pi\tilde{L} + \varepsilon$ . Therefore, the conclusion of Theorem A1 follows if we prove the existence of the sequence  $(k_n)$ . We adapt below an argument used in [BBM2]. Set  $H = \sum_{j \geq 1} |\nabla g_j| \in L^1$  (this uses the equivalence of  $|\cdot|_{W^{1,1}}$  and  $[\cdot]_{W^{1,1}}$ ). Since  $|\nabla(\Pi_{j=1}^k g_j)| \leq H$ ,  $\forall k$ , we may find a sequence  $k_n \rightarrow \infty$  and a map  $g \in BV(\Omega; S^1)$  such that  $h_n = \Pi_{j=1}^{k_n} g_j \rightarrow g$  a.e. Then, for  $m > n$ , we have

$$\begin{aligned} |h_m - h_n|_{W^{1,1}} &= |h_n(h_m \bar{h}_n - 1)|_{W^{1,1}} \\ &\leq |h_m \bar{h}_n|_{W^{1,1}} + \|(1 - h_m \bar{h}_n) \nabla h_n\|_{L^1} \\ &\leq \sum_{j=k_n+1}^{k_m} \|\nabla g_j\|_{L^1} + \|(1 - h_m \bar{h}_n) H\|_{L^1} = A_{m,n} + B_{m,n}. \end{aligned}$$

Let  $0 < \delta < 1$ . Then, clearly,  $A_{m,n} < \delta$  provided  $m, n$  are sufficiently large. On the other hand,

$$B_{m,n} \leq \delta \|H\|_{L^1} + 2 \int_{\{x: |1 - h_m(x) \bar{h}_n(x)| \geq \delta\}} |H|.$$

Note that

$$\{x; |1 - h_m(x) \bar{h}_n(x)| \geq \delta\} \subset \left\{x; |g(x) - h_m(x)| \geq \frac{\delta}{3}\right\} \cup \left\{x; |g(x) - h_n(x)| \geq \frac{\delta}{3}\right\}.$$

Since  $h_n \rightarrow g$  a.e., we find that  $B_{m,n} \leq \delta(\|H\|_{L^1} + 1)$ , provided  $m, n$  are sufficiently large. Therefore,  $(h_n)$  is a Cauchy sequence in  $W^{1,1}$  and converges to the above  $g$  in  $W^{1,1}$ .

It is not clear whether the reverse inequality in Theorem A1 is valid in general :

**Open Problem 8.** Let  $P_i, N_i \in \Omega$  be such that  $T(g) = 2\pi \sum_i (\delta_{P_i} - \delta_{N_i})$ . Is it true that

$$[g]_{W^{1,1}} \geq 2\pi\tilde{L} ?$$

Note that, by definition, the answer is yes if  $T(g) = 2\pi(\delta_P - \delta_N)$ .

## B. Proof of Theorems 1', 3', and 5'.

*Proof of Theorem 3'.* Let us first assume that  $g \in C^\infty(\bar{\Omega} \setminus \{a_1, \dots, a_k\}; S^1) \cap W^{1,1}$ . It is then easy to see that

$$\langle T(g), \zeta \rangle = 2\pi \sum_{j=1}^k d_j \zeta(a_j) + \int_{\partial\Omega} (g \wedge g_\tau) \zeta, \quad \forall \zeta \in \text{Lip}(\Omega; \mathbb{R}),$$

where  $d_j$  denotes the topological degree of  $g$  with respect to any small circle centered at  $a_j$ . In particular,

$$(B1) \quad \langle T(g), \zeta \rangle = 2\pi \sum_{j=1}^k d_j \zeta(a_j), \quad \forall \zeta \in W_0^{1,\infty}(\Omega).$$

Note that, in general,  $\sum_j d_j \neq 0$ . This means that we do not have necessarily the same number of positive and negative points as before. In order to compensate this, we insert points from  $\partial\Omega$  into (B1). Since  $\zeta = 0$  on  $\partial\Omega$ , equality in (B1) remains true. We can then relabel the points  $a_j$  as  $P_1, \dots, P_\ell, N_1, \dots, N_\ell$ , according to their multiplicity  $d_j$ , so that (B1) becomes

$$T(g) = 2\pi \sum_{j=1}^{\ell} (\delta_{P_j} - \delta_{N_j}) \quad \text{in } W_0^{1,\infty}(\Omega).$$

For a general  $g \in W^{1,1}(\Omega; S^1)$ , we argue by density using Lemma 2 to conclude that

$$T(g) = 2\pi \sum (\delta_{P_i} - \delta_{N_i}) \quad \text{in } [W_0^{1,\infty}(\Omega)]^*.$$

Note that  $d_\Omega$  induces a metric on the space  $\overline{\Omega}/\partial\Omega$ , where  $\partial\Omega$  is identified with a single point. Moreover, Lipschitz functions  $\zeta$  on  $\overline{\Omega}/\partial\Omega$  with  $|\zeta|_{\text{Lip}} \leq 1$  and  $\zeta(\partial\Omega) = 0$  correspond to elements in  $W_0^{1,\infty}(\Omega)$  such that  $\|\nabla\zeta\|_{L^\infty} \leq 1$ . Applying Lemma 12' in [BBM2] to  $\overline{\Omega}/\partial\Omega$ , we obtain

$$L(g) = \text{Inf} \sum_i d_\Omega(P_i, N_i).$$

*Remark B1.* The main new feature when  $\Omega$  is a bounded domain in  $\mathbb{R}^2$  is that a minimal connection is made of segments from a positive singularity  $P_i$  to some negative  $N_j$ , but we can also have line segments joining the singularities  $P_i, N_i$  to the boundary  $\partial\Omega$ . This is the analog of Example 3 in [BCL].

*Proof of Theorem 1'.* The proof of  $E(g) = E_{\text{rel}}(g)$  is exactly the same as in Proposition 2 and we shall omit it.

We are left to show that

$$(B2) \quad E(g) = \int_{\Omega} |\nabla g| + 2\pi L(g).$$

Let  $\varphi \in BV(\Omega; \mathbb{R})$  be such that  $g = e^{i\varphi}$ . Using Vol'pert's chain rule as in the proof of Lemma 5, we have

$$(B3) \quad |D\varphi|_{\mathcal{M}(\Omega)} = |g|_{W^{1,1}} + |g \wedge \nabla g - D\varphi|_{\mathcal{M}(\Omega)}.$$

We claim that

$$(B4) \quad |g \wedge \nabla g - D\varphi|_{\mathcal{M}(\Omega)} \geq 2\pi L(g).$$

In fact, for every  $\zeta \in C_0^\infty(\Omega)$  such that  $\|\nabla\zeta\|_{L^\infty} \leq 1$ ,

$$|g \wedge \nabla g - D\varphi|_{\mathcal{M}(\Omega)} \geq \int_{\Omega} (g \wedge \nabla g) \cdot \nabla^\perp \zeta - \int_{\Omega} D\varphi \cdot \nabla^\perp \zeta = \langle T(g), \zeta \rangle.$$

Taking the supremum with respect to  $\zeta$ , we conclude that (B4) holds.

Inequality  $\geq$  in (B2) follows immediately from (B3) and (B4).

We now establish  $\leq$  in (B2). Let us assume for the moment that  $g$  is smooth outside finitely many points  $a_1, \dots, a_k$ , and that  $g$  has topological degree  $\pm 1$  at each one of those points. Let  $\mathcal{C}$  be a minimal connection between those points with respect to the distance  $d_\Omega$ .

Note that on any closed curve contained in  $\Omega \setminus \mathcal{C}$ ,  $g$  has zero topological degree. We conclude that  $g$  has a smooth lifting  $\varphi$  on  $\Omega \setminus \mathcal{C}$ . Moreover, as we cross any one of the line segments of  $\mathcal{C}$ ,  $\varphi$  jumps by  $2\pi$ . Thus,  $\varphi \in BV(\Omega; \mathbb{R})$  and

$$\int_{\Omega} |D\varphi| = \int_{\Omega} |\nabla g| + 2\pi|\mathcal{C}| = \int_{\Omega} |\nabla g| + 2\pi L(g).$$

We can now argue by density, using Lemma 2, to conclude that for any  $g \in W^{1,1}(\Omega; S^1)$  there exists  $\varphi \in BV(\Omega; \mathbb{R})$  such that  $g = e^{i\varphi}$  a.e. in  $\Omega$  and

$$(B5) \quad \int_{\Omega} |D\varphi| \leq \int_{\Omega} |\nabla g| + 2\pi L(g).$$

This concludes the proof of the theorem.

*Proof of Theorem 5'.* Using exactly the same argument as in the proof of Proposition 3, we have

$$(B6) \quad |\langle \text{Det}(\nabla g), \zeta \rangle| \leq TV(g) \|\zeta\|_{L^\infty}, \quad \forall \zeta \in C_0^\infty(\Omega).$$

Thus, if  $TV(g) < \infty$ , then  $\text{Det}(\nabla g) \in \mathcal{M}(\Omega)$ . We now apply Proposition 3.2 in [S] (see also [P]) to the quotient space  $\bar{\Omega}/\partial\Omega$ . We conclude that there exist distinct points  $a_1, \dots, a_k \in \Omega$  and nonzero integers  $d_1, \dots, d_k$  such that

$$(B7) \quad \text{Det}(\nabla g) = \pi \sum_{j=1}^k d_j \delta_{a_j}.$$

We now define

$$h(x) := \left( \frac{x - a_1}{|x - a_1|} \right)^{-d_1} \cdots \left( \frac{x - a_k}{|x - a_k|} \right)^{-d_k} g(x) \quad \text{for a.e. } x \in \Omega.$$

Clearly,  $\text{Det}(\nabla h) = 0$  in  $\mathcal{D}'(\Omega)$ . It follows from the analog of Theorem 7 for domains in  $\mathbb{R}^2$  (see also [D]) that  $h$  has a lifting in  $W^{1,1}$ . In other words, we can find  $\varphi \in W^{1,1}(\Omega; \mathbb{R})$  such that  $h = e^{i\varphi}$  a.e. in  $\Omega$ . We then conclude that

$$g(x) = \left( \frac{x - a_1}{|x - a_1|} \right)^{d_1} \cdots \left( \frac{x - a_k}{|x - a_k|} \right)^{d_k} e^{i\varphi(x)} \quad \text{for a.e. } x \in \Omega.$$

Arguing as in the proof of Theorem 5, this implies that

$$(B8) \quad TV(g) \leq \pi \sum_{j=1}^k |d_j|.$$

The reverse inequality already follows from (B6). We then conclude that (6.8) holds.

Conversely, if  $\text{Det}(\nabla g) \in \mathcal{M}(\Omega)$ , then (B7) holds. The above argument then shows that  $TV(g) < \infty$  and

$$TV(g) = |\text{Det}(\nabla g)|_{\mathcal{M}} = \pi \sum_{i=1}^k |d_i|.$$

### C. Proof of Theorem 3''''.

Theorem 3'''' follows immediately from Theorem 3 and the next

**Lemma C1.** *Given  $g \in W^{1,1}(S^2; \Gamma)$ , we define*

$$g_j := \frac{g - a_j}{|g - a_j|} \in W^{1,1}(S^2; S^1).$$

*Then,*

$$(C1) \quad \text{Det}(\nabla g) = \frac{1}{\pi} \sum_j |A_j| \text{Det}(\nabla g_j) \quad \text{in } \mathcal{D}'(S^2).$$

The proof of Lemma C1 relies on the following

**Lemma C2.** *For any  $u \in W^{1,1}(S^1; \Gamma)$ , we have*

$$(C2) \quad \frac{1}{2} \int_{S^1} u \wedge u_\tau = \sum_j |A_j| \deg \frac{u - a_j}{|u - a_j|}.$$

*Proof of Lemma C2.*

*Step 1.*  $\Gamma$  is a simple curve.

It is well-known that (C2) holds if  $u \in C^1(S^1; \Gamma)$  (see, e.g., [N]). By approximation, we conclude that (C2) is also true for any  $u \in W^{1,1}(S^1; \Gamma)$ .

*Step 2.* Assume  $\Gamma$  has finitely many self-intersections, say  $q_1, \dots, q_k$ . Since  $u$  is continuous, the set

$$S^1 \setminus u^{-1}(\{q_1, \dots, q_k\})$$

is open and can be written as a countable union of open arcs in  $S^1$ . Let  $\alpha_1$  be such an arc. It is easy to see that we can select disjoint arcs  $\alpha_2, \dots, \alpha_j$  (oriented anticlockwise) such that  $u$  at the positive endpoint of  $\alpha_i$  coincides with the value of  $u$  at the negative endpoint of  $\alpha_{i+1}$  for  $i = 1, \dots, j$ , with the convention that  $\alpha_{j+1} = \alpha_1$ . By removing arcs from this list if necessary, we can assume that each point  $q_i$  appears only twice in the list

$$\{u(\partial\alpha_1), \dots, u(\partial\alpha_j)\}.$$

This construction induces a function  $\tilde{u} \in W^{1,1}(S^1; \Gamma)$  such that

- (a)  $\tilde{u} = u$  on  $\alpha_1 \cup \dots \cup \alpha_j$  ;
- (b)  $\tilde{u}$  is locally constant on  $S^1 \setminus \alpha_1 \cup \dots \cup \alpha_j$ .

By construction,  $\tilde{u}(S^1)$  is a subset of a Jordan curve  $\tilde{\Gamma}$  contained in  $\Gamma$ . Let  $A_{t_1}, \dots, A_{t_\ell}$  be the components of  $\mathbb{R}^2 \setminus \tilde{\Gamma}$  enclosed by  $\tilde{\Gamma}$ . By our first step, we have

$$(C3) \quad \frac{1}{2} \int_{S^1} \tilde{u} \wedge \tilde{u}_\tau = |A_{t_1} \cup \dots \cup A_{t_\ell}| \deg \frac{\tilde{u} - a}{|\tilde{u} - a|}.$$

for some point  $a$  inside  $\tilde{\Gamma}$ . Note, however, that

$$\deg \frac{\tilde{u} - a}{|\tilde{u} - a|} = \deg \frac{\tilde{u} - a_{t_s}}{|\tilde{u} - a_{t_s}|}, \quad \forall s = 1, \dots, \ell ;$$

moreover,

$$\deg \frac{\tilde{u} - a_j}{|\tilde{u} - a_j|} = 0, \quad \text{if } j \notin \{t_1, \dots, t_\ell\}.$$



We can thus rewrite (C3) as

$$(C4) \quad \frac{1}{2} \int_{S^1} \tilde{u} \wedge \tilde{u}_\tau = \sum_{s=1}^{\ell} |A_{t_s}| \deg \frac{\tilde{u} - a_{t_s}}{|\tilde{u} - a_{t_s}|} = \sum_j |A_j| \deg \frac{\tilde{u} - a_j}{|\tilde{u} - a_j|}.$$

We can proceed with the construction of  $\tilde{u}$  and “decompose”  $u \in W^{1,1}(S^1; \Gamma)$  as  $\tilde{u}_1, \tilde{u}_2, \dots$  so that

- (a)  $\tilde{u}_i \in W^{1,1}(S^1; \Gamma)$  and  $\tilde{u}_i(S^1)$  is contained in some Jordan curve for every  $i$  ;
- (b)  $u_\tau = \tilde{u}_{1\tau} + \tilde{u}_{2\tau} + \dots$  in  $S^1$  ;
- (c)  $\tilde{u}_i$  coincides with  $u$  on finitely many arcs in  $S^1 \setminus u^{-1}(\{q_1, \dots, q_k\})$  and  $\tilde{u}_i$  is locally constant outside those arcs.

By (C4), we have

$$\frac{1}{2} \int_{S^1} \tilde{u}_i \wedge \tilde{u}_{i\tau} = \sum_j |A_j| \deg \frac{\tilde{u}_i - a_j}{|\tilde{u}_i - a_j|}, \quad \forall i.$$

Note that, by (b) and (c),

$$\int_{S^1} u \wedge u_\tau = \sum_i \int_{S^1} \tilde{u}_i \wedge \tilde{u}_{i\tau}.$$

For the same reason,

$$\deg \frac{u - a_j}{|u - a_j|} = \sum_i \deg \frac{\tilde{u}_i - a_j}{|\tilde{u}_i - a_j|}, \quad \forall j.$$

We conclude that (C2) holds.

*Proof of Lemma C1.* Let  $g \in W^{1,1}(S^2; \Gamma)$ . By the coarea formula (see [BBM2]), we have

$$(C5) \quad \langle \text{Det}(\nabla g), \zeta \rangle = \int_{\mathbb{R}} \left( \int_{\Sigma_\lambda} g \wedge g_\tau \right) d\lambda,$$

where  $\zeta \in C^\infty(S^2)$ , and  $\Sigma_\lambda = \{x \in S^2; \zeta(x) = \lambda\}$  is equipped with the appropriate orientation, whenever  $\lambda$  is a regular value of  $\zeta$ .

Recall that, for a.e.  $\lambda \in \mathbb{R}$ ,  $g|_{\Sigma_\lambda}$  belongs to  $W^{1,1}$ . Applying Lemma C2 to  $g|_{\Sigma_\lambda}$  for such  $\lambda$ s we get

$$\int_{\Sigma_\lambda} g \wedge g_\tau = 2 \sum_j |A_j| \deg g_j = \frac{1}{\pi} \sum_j |A_j| \int_{\Sigma_\lambda} g_j \wedge g_{j\tau}.$$

Integrate both sides of the identity above with respect to  $\lambda$ . Using (C5), we conclude that

$$\langle \text{Det}(\nabla g), \zeta \rangle = \frac{1}{\pi} \sum_j |A_j| \langle \text{Det}(\nabla g_j), \zeta \rangle, \quad \forall \zeta \in C^\infty(S^2).$$

This establishes (C1).

We also call the attention of the reader to the following analog of Lemma 12' in [BBM2] :

**Proposition C1.** *Let  $X$  be a metric space. Given two sequences  $(P_i), (N_i)$  in  $X$  and nonnegative numbers  $\alpha_i$  such that  $\sum_i \alpha_i d(P_i, N_i) < \infty$ , let*

$$(C6) \quad T = \sum_i \alpha_i (\delta_{P_i} - \delta_{N_i}) \quad \text{in } [\text{Lip}(X)]^*.$$

Define

$$L = \sup_{\substack{\zeta \in \text{Lip}(X) \\ |\zeta|_{\text{Lip}} \leq 1}} \langle T, \zeta \rangle.$$

Then,

$$(C7) \quad L = \inf \sum_i \alpha_i d(P_i, N_i),$$

where the infimum is taken over all sequences  $(P_i), (N_i)$  in  $X$  and numbers  $\alpha_i \geq 0$  such that (C6) holds.

*Proof.* Let us denote by  $\tilde{L}$  the infimum in (C7). Clearly,  $L \leq \tilde{L}$ . We now establish the reverse inequality.

Let  $\varepsilon > 0$ . We take  $k \geq 1$  sufficiently large so that

$$\sum_{i>k} \alpha_i d(P_i, N_i) < \varepsilon.$$

Without loss of generality, we can assume that each  $\alpha_i$  is rational for  $i = 1, \dots, k$ . We then choose an integer  $J \geq 1$  sufficiently large so that  $J\alpha_i$  is an integer for every  $i = 1, \dots, k$ . Write the points  $P_i, N_i$  as  $p_1, p_2, \dots$  and  $n_1, n_2, \dots$ , with multiplicity  $J\alpha_i$ . It follows from Lemma 4.2 in [BCL] that we can find  $\zeta_0 \in \text{Lip}(X)$ , with  $|\zeta_0|_{\text{Lip}} \leq 1$ , such that, after relabeling the points  $n_j$  if necessary, we have :

$$\sum_{i=1}^k J\alpha_i [\zeta_0(P_i) - \zeta_0(N_i)] = \sum_j [\zeta_0(p_j) - \zeta_0(n_j)] = \sum_j d(p_j, n_j).$$

Thus,

$$\begin{aligned}
L &\geq \sum_{i=1}^{\infty} \alpha_i [\zeta_0(P_i) - \zeta_0(N_i)] \\
&\geq \sum_{i=1}^k \alpha_i [\zeta_0(P_i) - \zeta_0(N_i)] - \varepsilon \\
&= \frac{1}{J} \sum_j d(p_j, n_j) - \varepsilon \geq \sum_j \frac{1}{J} d(p_j, n_j) + \sum_{i>k} \alpha_i d(P_i, N_i) - 2\varepsilon.
\end{aligned}$$

Note that

$$T = \sum_j \frac{1}{J} (\delta_{p_j} - \delta_{n_j}) + \sum_{i>k} \alpha_i (\delta_{P_i} - \delta_{N_i}).$$

We conclude that  $L \geq \tilde{L} - 2\varepsilon$ . Since  $\varepsilon > 0$  was arbitrary, this implies  $L \geq \tilde{L}$ . Thus,  $L = \tilde{L}$  as claimed.

#### D. Proof of Theorems 1'' and 2''.

We begin with a few preliminary results

**Lemma D1.** *Given  $\varepsilon > 0$ , let  $\Phi_\varepsilon : \bar{B}^N \rightarrow \bar{B}^N$  be defined as*

$$\Phi_\varepsilon(x) = \begin{cases} 0 & \text{if } |x| \leq \varepsilon, \\ \frac{|x| - \varepsilon}{1 - \varepsilon} \frac{x}{|x|} & \text{if } \varepsilon < |x| \leq 1. \end{cases}$$

Then, for every  $f \in C^\infty(\bar{B}^N; \mathbb{R}^M)$ , we have

$$f \circ \Phi_\varepsilon \rightarrow f \quad \text{in } W^{1,N}(B^N).$$

*Proof.* Given  $x \in B^N$ ,  $|x| \geq \varepsilon$ , let  $r = |x|$  and  $\psi_\varepsilon(r) = \frac{r - \varepsilon}{(1 - \varepsilon)r}$ . Using this notation, we have

$$\Phi'_\varepsilon(x) = \psi_\varepsilon(r) \text{Id} + \left( \frac{x_i x_j}{r} \psi'_\varepsilon(r) \right).$$

Since  $\psi'_\varepsilon(r) = \frac{\varepsilon}{(1 - \varepsilon)r^2}$ ,

$$\left| \frac{x_i x_j}{r} \psi'_\varepsilon(r) \right| \leq \frac{C\varepsilon}{r}.$$

Moreover,

$$|\Phi_\varepsilon(x) - x| \leq C\varepsilon \quad \text{and} \quad |\psi_\varepsilon(r) - 1| \leq \frac{C\varepsilon}{r}.$$

We then have

$$\begin{aligned}
 |\nabla f_\varepsilon(x) - \nabla f(x)| &= |{}^t\Phi'_\varepsilon(x) \nabla f(\Phi_\varepsilon(x)) - \nabla f(x)| \\
 &\leq |\nabla f(\Phi_\varepsilon(x)) - \nabla f(x)| + |\nabla f(\Phi_\varepsilon(x))| |\text{Id} - {}^t\Phi'_\varepsilon(x)| \\
 &\leq C|\Phi_\varepsilon(x) - x| + C|\psi_\varepsilon(r) - 1| + \frac{C\varepsilon}{r} \leq \frac{C\varepsilon}{r}
 \end{aligned}$$

for  $|x| \geq \varepsilon$ . Therefore,

$$\int_{B^N} |\nabla f_\varepsilon - \nabla f|^N \leq C\varepsilon^N \int_{\varepsilon \leq |x| \leq 1} \frac{dx}{|x|^N} + \int_{|x| < \varepsilon} |\nabla f|^N \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

Next, we establish the following

**Lemma D2.** *Given  $f \in C^\infty(\bar{B}^N \times [0, 1]; \mathbb{R}^M)$ , let*

$$f_\varepsilon(x, t) = f(\Phi_\varepsilon(x), t).$$

*Then*

$$f_\varepsilon \rightarrow f \quad \text{in } W^{1,N}(B^N \times [0, 1]).$$

*Proof.* Note that

$$\frac{\partial f_\varepsilon}{\partial t}(x, t) - \frac{\partial f}{\partial t}(x, t) = \begin{cases} \frac{\partial f}{\partial t}(0, t) - \frac{\partial f}{\partial t}(x, t) & \text{if } |x| < \varepsilon, \\ \frac{\partial f}{\partial t}(\Phi_\varepsilon(x), t) - \frac{\partial f}{\partial t}(x, t) & \text{if } |x| \geq \varepsilon. \end{cases}$$

Thus, since  $|\Phi_\varepsilon(x) - x| \leq C\varepsilon$ , we have

$$\left| \frac{\partial f_\varepsilon}{\partial t} - \frac{\partial f}{\partial t} \right| \leq C\varepsilon.$$

The result now immediately follows from Lemma D1.

As a consequence of Lemma D2 above, any map  $f \in C^\infty(\bar{B}^N \times [0, 1]; S^k)$  can be approximated in  $W^{1,N}$  by maps  $f_\varepsilon$  such that

$$f_\varepsilon = f \quad \text{on } S^{N-1} \times [0, 1] \quad \text{and} \quad f_\varepsilon(x, t) = g_\varepsilon(t) \quad \text{if } |x| \leq \varepsilon.$$

**Lemma D3.** *Given  $g \in C^\infty([0, 1]; \mathbb{R}^M)$ , let*

$$f(x, t) = g(t) \quad \forall (x, t) \in \bar{B}^N \times [0, 1].$$

*Then, there exists a sequence  $(f_\varepsilon)$  in  $W^{1,N}$  such that*

$$f_\varepsilon(x, t) = \text{const} \quad \text{if } |x| \leq \varepsilon, \quad f_\varepsilon(x, t) = g(t) \quad \text{if } |x| = 1,$$

*and*

$$f_\varepsilon \rightarrow f \quad \text{in } W^{1,N}(B^N \times [0, 1]).$$

*Proof.* Let  $\zeta_\varepsilon : \bar{B}^N \rightarrow \mathbb{R}$  be given by

$$\zeta_\varepsilon(x) = \begin{cases} \frac{\log \frac{|x|}{\varepsilon}}{\log \frac{1}{\varepsilon}} & \text{if } \varepsilon \leq |x| \leq 1, \\ 0 & \text{if } |x| < \varepsilon. \end{cases}$$

It is easy to see that

$$\int_{B^N} |\nabla \zeta_\varepsilon|^N \rightarrow 0 \quad \text{and} \quad \int_{B^N} |\zeta_\varepsilon - 1|^N \rightarrow 0.$$

Since  $N \geq 2$ , there exists  $Q \in S^N$  such that  $Q \notin g([0, 1])$ . Let  $\Psi : S^N \setminus \{Q\} \rightarrow \mathbb{R}^N$  denote the stereographic projection. Set  $F = \Psi \circ g$ . Clearly, in order to establish the lemma, it suffices to approximate  $F$  in  $W^{1,N}$  by a sequence  $F_\varepsilon$  such that  $|F_\varepsilon| \leq C$ ,

$$F_\varepsilon(x, t) = \text{const} \quad \text{if } |x| \leq \varepsilon \quad \text{and} \quad F_\varepsilon(x, t) = F(x, t) \quad \text{if } |x| = 1.$$

Set  $G(t) = \Psi \circ g(t)$  and  $F_\varepsilon(x, t) = \zeta_\varepsilon(x)G(t)$ . We then have

$$\iint |\nabla_x F_\varepsilon - \nabla F|^N dx dt = \int |\nabla \zeta_\varepsilon|^N \int |G|^N \rightarrow 0$$

and

$$\iint |\partial_t F_\varepsilon - \partial_t F|^N dx dt = \int |1 - \zeta_\varepsilon|^N \int |G'|^N \rightarrow 0.$$

The proof of Lemma D3 is complete.

We conclude from Lemma D3 that, given any  $u \in W^{1,N-1}(S^N; S^{N-1})$ ,  $N \geq 3$ , there exists a sequence  $(u_n)$  such that  $u_n \rightarrow u$  in  $W^{1,N}$ , where each  $u_n$  satisfies the following properties :

- (i)  $u_n$  has a finite number of point singularities  $P_i, N_i$  ;

- (ii)  $u_n$  is homogeneous of degree 0 in a neighborhood of each singularity ;
- (iii)  $u_n \equiv Q$  in some conic neighborhood of a geodesic joining  $P_i$  and  $N_i$ .

In fact, (ii) holds since every topological singularity may be approximated by homogeneous singularities (see Lemma E.5 below). We then apply Lemmas D.2 and D.3 to obtain property (iii).

We shall say that a map is *good* if it satisfies properties (i)–(iii).

*Proof of Theorem 2''.* Our goal is to show that, for every  $u \in W^{1,N-1}(S^N; S^{N-1})$ ,

$$(D1) \quad \inf_{v \in C^\infty(S^N; S^{N-1})} \int_{S^N} |D(u) - D(v)| = \sigma_N L(u).$$

*Proof of “ $\geq$ ”.* For every  $\zeta \in \text{Lip}(S^N)$  such that  $\|\nabla \zeta\|_{L^\infty} \leq 1$ , we have

$$\int_{S^N} |D(u) - D(v)| \geq \int_{S^N} [D(u) - D(v)] \cdot \nabla \zeta = \int_{S^N} D(u) \cdot \nabla \zeta = \langle T(u), \zeta \rangle.$$

Taking the supremum with respect to  $\zeta$ , we get

$$\int_{S^N} |D(u) - D(v)| \geq \sigma_N L(u) \quad \forall v \in C^\infty(S^N; S^{N-1}).$$

*Proof of “ $\leq$ ”.* Assume  $u$  is a good map. We shall assume for simplicity that  $u$  has a single dipole  $P, N$ . Given  $\varepsilon > 0$ , let  $U_\varepsilon : S^{N-1} \rightarrow S^{N-1}$  be such that

$$U_\varepsilon \equiv Q \quad \text{if } |x - P| \geq \frac{\varepsilon}{2} \quad \text{and} \quad \int_{S^{N-1}} |\nabla_T U_\varepsilon|^{N-1} \leq \sigma_N (N-1)^{\frac{N-1}{2}} + \varepsilon.$$

The existence of such map is established in [BCL, Section VIII].

Let  $W$  denote the  $\varepsilon$ -conic neighborhood of the geodesic segment joining  $P$  and  $N$ . We decompose  $W$  as  $W = W_1 \cup W_2$ , where  $W_1$  is the cylindrical part of  $W$ , and  $W_2$  is the union of the two conic caps. We then define  $u_\varepsilon$  as

$$u_\varepsilon(x) = u(x) \quad \text{if } x \notin W, \quad u_\varepsilon(x) = U_\varepsilon(x) \quad \text{if } x \in W_1$$

and  $u_\varepsilon$  is extended by homogeneity of degree 0 in  $W_2$ . We then have

$$\int_{S^N} |D(u) - D(u_\varepsilon)| = \int_{W_1} |D(U_\varepsilon)| + o(1).$$

In [BCL, Section VIII], it is proved that

$$|D(U_\varepsilon)| = |\nabla_T U_\varepsilon| \leq \frac{1}{(N-1)^{\frac{N-1}{2}}} |\nabla_T U_\varepsilon|^{N-1}.$$

We conclude that

$$(D2) \quad \lim_{\varepsilon \rightarrow 0} \int_{S^N} |D(u) - D(u_\varepsilon)| \leq \sigma_N L(u).$$

On the other hand, since  $u_\varepsilon$  has only singularities of degree 0, Hopf's theorem implies that  $u_\varepsilon \in \overline{C^\infty(S^N; S^{N-1})}^{W^{1,N-1}}$ . It then follows from (D2) that inequality “ $\leq$ ” in (D1) holds, at least when  $u$  is a good map.

We now establish  $\leq$  in (D1) for any map  $u \in W^{1,N-1}(S^N; S^{N-1})$ . Let  $(u_n)$  be a sequence of good maps such that  $u_n \rightarrow u$  in  $W^{1,N-1}$ . For each  $n \geq 1$ , we have just shown that there exists  $v_n : S^N \rightarrow S^{N-1}$  smooth such that

$$\int_{S^N} |D(u_n) - D(v_n)| \leq \sigma_N L(u_n) + \frac{1}{n}.$$

Thus

$$\int_{S^N} |D(u) - D(v_n)| \leq \int_{S^N} |D(u) - D(u_n)| + \sigma_N L(u_n) + \frac{1}{n} = \sigma_N L(u) + o(1),$$

which gives the desired result.

*Proof of Theorem 1''.* We want to show that

$$(D3) \quad E_{\text{rel}}(u) = \int_{S^N} |\nabla u|^{N-1} + (N-1)^{\frac{N-1}{2}} \sigma_N L(u).$$

*Proof of “ $\leq$ ”.* It suffices to establish the result for good maps. In fact, if  $u_n \rightarrow u$  in  $W^{1,N-1}$ , then

$$E_{\text{rel}}(u) \leq \liminf_{n \rightarrow \infty} E_{\text{rel}}(u_n),$$

while the right-hand side of (D3) is continuous with respect to the strong topology in  $W^{1,N-1}$ . Thus, we may assume that  $u$  is good and we can proceed exactly as in the proof of Theorem 2''. We shall leave the details to the reader.

*Proof of “ $\geq$ ”.* As in [BCL, Section VIII], we have

$$(D4) \quad |(w \cdot v_2 \wedge \cdots \wedge v_N, w \cdot v_1 \wedge v_3 \wedge \cdots \wedge v_N, \dots)| \leq \frac{|w| \left( \sum_j |v_j|^2 \right)^{\frac{N-1}{2}}}{(N-1)^{\frac{N-1}{2}}}$$

for every  $w, v_1, \dots, v_N \in \mathbb{R}^N$ .

On the other hand, given a sequence  $(u_n) \subset C^\infty(S^N; S^{N-1})$  such that  $(u_n)$  is bounded in  $W^{1,N-1}$  and  $u_n \rightarrow u$  a.e., we have

$$(D5) \quad \int_{S^N} |\nabla u_n|^{N-1} = \int_{S^N} |\nabla u|^{N-1} + \int_{S^N} |\nabla u_n - \nabla u|^{N-1} + o(1).$$

Let  $\zeta \in \text{Lip}(S^N)$  be such that  $\|\nabla \zeta\|_{L^\infty} \leq 1$ . Applying (D4) to  $w = u_n$  and  $v = \nabla u_n - \nabla u$ , we get

$$\frac{\int_{S^N} |\nabla u_n - \nabla u|^{N-1}}{(N-1)^{\frac{N-1}{2}}} \geq \sum_{\sigma} \varepsilon_{\sigma} \int_{S^N} u_n \cdot (u_{nx_{\sigma_2}} - u_{x_{\sigma_2}}) \wedge \cdots \wedge (u_{nx_{\sigma_N}} - u_{x_{\sigma_N}}) \zeta_{x_{\sigma_1}},$$

where the sum is taken with respect to all permutations  $\sigma$  of  $N$  elements, and  $\varepsilon_{\sigma} = \pm 1$  denotes the sign of  $\sigma$ . Thus,

$$(D6) \quad \int_{S^N} |\nabla u_n - \nabla u|^{N-1} \geq -(N-1)^{\frac{N-1}{2}} \int_{S^N} D(u) \cdot \nabla \zeta + R_n(\zeta).$$

Assume for the moment that  $R_n(\zeta) \rightarrow 0$ . It follows from (D5) and (D6) that

$$\liminf_{n \rightarrow \infty} \int_{S^N} |\nabla u_n|^{N-1} \geq \int_{S^N} |\nabla u|^{N-1} + (N-1)^{\frac{N-1}{2}} \langle T(u), \zeta \rangle.$$

Taking the supremum over  $\zeta$ , we obtain “ $\leq$ ” in (D3).

Thus, in order to conclude the proof of Theorem 1'', we need to show that  $R_n(\zeta) \rightarrow 0$ . Since each  $u_n$  is smooth,

$$\int_{S^N} D(u_n) \cdot \nabla \zeta = 0.$$

It then follows that

$$R_n(\zeta) = S_n(\zeta) + o(1),$$

where  $S_n(\zeta)$  is a sum of integrals of the form

$$I_n = \pm \int_{S^N} u_n \cdot v_{nx_{i_1}} \wedge \cdots \wedge v_{nx_{i_k}} \wedge u_{x_{j_1}} \wedge \cdots \wedge u_{x_{j_l}} \zeta_{x_t},$$

where  $k + l = N - 1$  and  $v_n = u_n - u$ . It is important to notice that  $k \geq 1$  and  $l \geq 1$ . Since

$$(u_n - u) u_{x_{j_1}} \wedge \cdots \wedge u_{x_{j_l}} \rightarrow 0 \quad \text{strongly in } L^{\frac{N-1}{l}},$$



we may replace  $I_n$  by

$$\pm \int_{S^N} u \cdot v_{nx_{i_1}} \wedge \cdots \wedge v_{nx_{i_k}} \wedge u_{x_{j_1}} \wedge \cdots \wedge u_{x_{j_l}} \zeta_{x_t}.$$

We can now formally integrate by parts to write  $S_n(\zeta)$  as a sum of integrals of the form

$$\pm \int_{S^N} u_{x_{i_1}} \cdot v_n \wedge v_{nx_{i_2}} \wedge \cdots \wedge v_{nx_{i_k}} \wedge u_{x_{j_1}} \wedge \cdots \wedge u_{x_{j_l}} \zeta_{x_t}.$$

Such integration by parts can be easily justified by approximation ; note that all the second derivatives are cancelled by symmetry.

Since

$$v_n u_{x_{j_1}} \wedge \cdots \wedge u_{x_{j_l}} \rightarrow 0 \quad \text{strongly in } L^{\frac{N-1}{l+1}},$$

we conclude that

$$R_n(\zeta) = S_n(\zeta) + o(1) = o(1).$$

The proof of Theorem 1' is completed.

### E. Proof of Theorem 5''.

Throughout this section, we assume that  $\Omega = \partial G$ , where  $G$  is a domain in  $\mathbb{R}^N$  ; more generally,  $\Omega$  could be any smooth domain (with boundary) in  $\partial G$ . We start with

**Lemma E1.** *Let  $0 < \sigma < \infty$ ,  $1 \leq q < \infty$ . Then*

$$\overline{C^\infty(\Omega; S^{N-1}) \cap W^{\sigma,q}}^{W^{\sigma,q}} = \overline{C^\infty(\overline{\Omega}; S^{N-1})}^{W^{\sigma,q}}.$$

*In other words, for each map  $u \in C^\infty(\Omega; S^{N-1}) \cap W^{\sigma,q}$ , there is a sequence  $(u_n) \subset C^\infty(\overline{\Omega}; S^{N-1})$  such that  $u_n \rightarrow u$  in  $W^{\sigma,q}$ .*

*Proof.* Let, for  $t > 0$  sufficiently small,  $\Omega_t = \{x \in \Omega ; d(x, \partial\Omega) > t\}$ . Consider, for any such  $t$ , a diffeomorphism  $\Phi_t : \Omega \rightarrow \Omega_t$  such that

$$\|D^k \Phi - D^k \text{Id}\|_{L^\infty} \leq C_k t, \quad k = 0, 1, 2, \dots$$

and set, for  $u \in C^\infty(\Omega; S^{N-1})$ ,  $u^t = u \circ \Phi_t$ . Then

$$u^t \in C^\infty(\overline{\Omega}; S^{N-1}) \quad \text{and} \quad u^t \rightarrow u \quad \text{in } W^{\sigma,q}.$$

**Lemma E2.** *Let  $u \in W^{1,N}(\Omega; S^{N-1})$ . Then*

$$L(u) = 0 \quad \Longleftrightarrow \quad u \in \overline{C^\infty(\Omega; S^{N-1}) \cap W^{1,N}}^{W^{1,N}}.$$

Here,

$$\langle T(u), \zeta \rangle = - \int_{\Omega} D(u) \cdot \nabla \zeta,$$

where  $\zeta \in W^{1,\infty}(\Omega)$  and  $\zeta$  is constant on each connected component of  $\partial\Omega$ ;  $L(u)$  is computed accordingly. When  $N = 3$ , this result is due to Bethuel [B1]; the same proof yields Lemma E2.

**Lemma E3.** *Let  $1 \leq p < N$ . For  $g \in W^{1,p}(S^{N-1}; S^{N-1})$ , set  $\tilde{g}(x) = g\left(\frac{x}{|x|}\right)$  for  $x \in B^N$ . Then  $\tilde{g} \in W^{1,p}$  and the map*

$$g \in W^{1,p}(S^{N-1}; S^{N-1}) \quad \longmapsto \quad \tilde{g} \in W^{1,p}(B^N; S^{N-1})$$

*is continuous and verifies  $|\tilde{g}|_{W^{1,p}} \leq C|g|_{W^{1,p}}$ .*

*Proof.* Trivial computation.

**Lemma E4.** *In the definition of TV, we may replace  $C^\infty(\bar{\Omega}; \mathbb{R}^N)$ -maps by maps in  $\text{Lip}(\bar{\Omega}; \mathbb{R}^N)$ .*

*Proof.* Clear, by approximation.

**Lemma E5.** *Let  $N \geq 2$ ,  $N - 1 \leq p < N$ , and let  $u \in W^{1,p}(\Omega; S^{N-1})$ . Fix  $a_1, \dots, a_k \in \Omega$  and define, for  $\rho > 0$  sufficiently small,*

$$u_\rho(x) = \begin{cases} u(x), & \text{if } d(x, \{a_1, \dots, a_k\}) \geq \rho \\ u(y), & \text{if, for some } j, d(x, a_j) < \rho \text{ and} \\ & \text{\textit{x lies on the geodesic segment from}} \\ & \text{\textit{a}_j \textit{ to y, where } } d(a_j, y) = \rho \end{cases}$$

*(here,  $d$  denotes the geodesic distance in  $S^N$ .)*

*Then  $u_{\rho_n} \rightarrow u$  for some sequence  $\rho_n \rightarrow 0$ .*

*Proof.* For simplicity, we may assume  $\Omega$  is flat near each  $a_j$ . Then the definition of  $u_\rho$  becomes

$$u_\rho(x) = \begin{cases} u(x) & \text{if } d(x, \{a_1, \dots, a_k\}) \geq \rho, \\ u\left(\rho \frac{x - a_j}{|x - a_j|}\right) & \text{if } |x - a_j| < \rho. \end{cases}$$

Alternatively, denoting by  $u_j^\rho(y) = u(a_j + \rho y)$ ,  $y \in S^{N-1}$ , then

$$u_\rho(x) = u_j^\rho \left( \frac{x - a_j}{|x - a_j|} \right) \quad \text{if } |x - a_j| < \rho.$$

Assume, for simplicity, that there is only one singularity, say  $a_1 = 0$ .

Let, for  $n \geq 1$ ,  $\rho_n$  be such that  $\frac{1}{2n} \leq \rho_n \leq \frac{1}{n}$  and

$$\frac{1}{2n} \int_{S_{\rho_n}} |\nabla u|^p \leq \int_{\frac{1}{2n}}^{\frac{1}{n}} \left( \int_{|x|=\rho} |\nabla u|^p \right) d\rho = \int_{\frac{1}{2n} \leq |x| \leq \frac{1}{n}} |\nabla u|^p.$$

Then

$$\int_{B_{\rho_n}} |\nabla u_{\rho_n}|^p \leq C_{N,p} \rho_n \int_{S_{\rho_n}} |\nabla u|^p \leq C \int_{\frac{1}{2n} \leq |x| \leq \frac{1}{n}} |\nabla u|^p \xrightarrow{n \rightarrow \infty} 0.$$

Thus

$$\begin{aligned} \int_{\Omega} |\nabla u_{\rho_n} - \nabla u|^p &= \int_{B_{\rho_n}} |\nabla u_{\rho_n} - \nabla u|^p \\ &\leq C \left( \int_{B_{\rho_n}} |\nabla u_{\rho_n}|^p + \int_{B_{\rho_n}} |\nabla u|^p \right) \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

*Proof of Theorem 5''.* The equivalence

$$TV(u) < \infty \quad \Longleftrightarrow \quad \text{Det}(\nabla u) \quad \text{is a measure}$$

is established as in the proof of Theorem 5. As already noted,  $TV(u) < \infty$  implies

$$\text{Det}(\nabla u) = \frac{\sigma_N}{N} \sum_{\text{finite}} (\delta_{P_i} - \delta_{N_i}).$$

Let  $a_1, \dots, a_k$  be the collection of points  $P_i, N_i$ . Given  $n \geq 1$ , let

$$\Omega^n = \left\{ x \in \Omega ; d(x, \{a_1, \dots, a_k\}) > \frac{1}{n} \right\}$$

and

$$A^n = \left\{ x \in \Omega ; \frac{1}{n} \leq d(x, \{a_1, \dots, a_k\}) \leq \frac{2}{n} \right\}.$$

Consider a sequence  $(u_n^m) \subset C^\infty(\overline{\Omega^n}; S^{N-1})$  such that  $u_n^m \rightarrow u$  in  $W^{1,N-1}$  as  $m \rightarrow \infty$  (clearly,  $L(u|_{\Omega_n}) = 0$ ). There is some  $\rho_n \in \left(\frac{1}{n}, \frac{2}{n}\right)$  such that, up to a subsequence in  $m$ ,

- (i)  $u_n^m|_{\Sigma_{\rho_n}} \rightarrow u|_{\Sigma_{\rho_n}}$  in  $W^{1,N-1}(\Sigma_{\rho_n})$  as  $m \rightarrow \infty$ ,
- (ii)  $\int_{\Sigma_{\rho_n}} |\nabla u|^N \leq Cn \int_{A_n} |\nabla u|^N$ .

Here,  $\Sigma_\rho = \{x; d(x, \{a_1, \dots, a_k\}) = \rho\}$ .

Extend  $u_n^m$  to  $\Omega$  as in Lemma 5 (by homogeneity of degree 0); let  $(u_n^m)_{\rho_n} = \tilde{u}_n^m$  be this extension. By (i) and Lemma E5, we have  $\tilde{u}_n^m \rightarrow u_{\rho_n}$  in  $W^{1,N-1}(\Omega)$  as  $m \rightarrow \infty$ . By (ii) and Lemma E3, we have  $u_{\rho_n} \rightarrow u$  in  $W^{1,N-1}(\Omega)$ . Thus we may find a sequence  $(v_n) \subset \text{Lip}_{\text{loc}}(\Omega \setminus \{a_1, \dots, a_k\}; S^{N-1})$  such that

- (a)  $v_n$  is homogeneous of degree 0 near each  $a_j$ ;
- (b)  $v_n \rightarrow u$  in  $W^{1,N-1}(\Omega)$ ;
- (c) near each  $a_j$ , the degree of  $v_n$  is  $d_j$ .

Assertion (c) follows from (i), the continuity of degree of maps from  $S^{N-1}$  into  $S^{N-1}$  for  $W^{1,N-1}$  convergence and the following clear fact

$$T(v_n) = \sigma_N \sum_j \deg(v_n, a_j) \delta_{a_j} \rightarrow T(u) = \sigma_N \sum_j d_j \delta_{a_j}.$$

For the remaining part of the proof, assume for simplicity that there is only one singularity  $a = 0$  of degree  $d > 0$  and that  $\Omega$  is flat near  $a$ . Let  $\rho_n$  be such that  $v_n$  is homogenous of degree 0 in  $B_{\rho_n}(0)$ . Here,  $B_\rho(0)$  is a ball in  $\Omega$  centered at  $a = 0$ . Fix  $d$  distinct points  $p_1, \dots, p_d$  in  $B_1$ .

Let  $\varepsilon > 0$  be sufficiently small. For  $w \in \text{Lip}(\partial B_1; S^{N-1})$  ( $B_1$  is the unit ball in  $\mathbb{R}^N$ ), with  $\deg w = d$ , let  $\tilde{w} : B_1 \setminus \bigcup_j B_\varepsilon(p_j) \rightarrow S^{N-1}$  be such that  $\tilde{w}|_{\partial B_1} = w$ ,  $\tilde{w}(x) = \frac{x - p_j}{|x - p_j|}$  if  $|x - p_j| = \varepsilon$ ,  $j = 1, \dots, d$  and  $\tilde{w}$  is a Lipschitz function. (Such a map exists, since  $\deg w = d$ .) We then extend  $\tilde{w}$  to  $B_1$  by setting  $\tilde{w}(x) = \frac{x - p_j}{\varepsilon}$ , if  $|x - p_j| \leq \varepsilon$ . Thus  $\tilde{w}$  is still Lipschitz.

Define, for  $0 < \rho < \rho_n$ ,

$$v_{n,\rho}(x) = \begin{cases} v_n(x), & \text{if } d(x, 0) > \rho \\ \tilde{w}_n\left(\frac{x}{\rho}\right), & \text{if } d(x, 0) \leq \rho \end{cases};$$

here,  $w_n(x) = v_n(\rho_n x)$ , if  $|x| = 1$ . Lemma E5 yields

$$v_{n,\rho} \rightarrow v_n \quad \text{in } W^{1,N-1} \quad \text{as } \rho \rightarrow 0.$$

Clearly, by the definition of  $v_{n,\rho}$ , we have

$$\int_{\Omega} |\text{Jac } v_{n,\rho}| = \omega_N d.$$

Considering now the case of several singularities, we obtain by a diagonal procedure a sequence  $(w_n) \subset \text{Lip}(\Omega; \mathbb{R}^N)$  such that

$$w_n \rightarrow u \quad \text{in } W^{1,N-1} \quad \text{and} \quad \int_{\Omega} |\text{Jac } w_n| = \omega_N \sum_j |d_j|.$$

### F. Proof of Theorems 5'''.

We start with the following well-known

**Lemma F1.** *Let  $1/p < s < 1$  and  $1 < p < \infty$ . Given  $u \in W_0^{1,p}(\mathbb{R}_+^N)$ , let*

$$\tilde{u}(x) = \begin{cases} u(x) & \text{if } x_N > 0, \\ 0 & \text{if } x_N \leq 0. \end{cases}$$

*Then,  $u \mapsto \tilde{u}$  is a continuous mapping from  $W_0^{1,p}(\mathbb{R}_+^N)$  into  $W^{1,p}(\mathbb{R}^N)$ .*

*Proof.* By density, it suffices to deal with  $u \in C_0^\infty(\mathbb{R}_+^N)$ . Using the Besov seminorm, we have

$$(F1) \quad |\tilde{u}|_{W^{s,p}}^p \sim |u|_{W^{s,p}}^p + \|u\|_{L^p}^p + \int_0^\infty \int_{\mathbb{R}} \int_{\mathbb{R}^{N-1}} \frac{|\tilde{u}(x', t + \tau) - \tilde{u}(x', t)|^p}{\tau^{1+sp}} dx' dt d\tau.$$

Denote by  $I$  the last term in the right-hand side of (F1). Clearly, it suffices to estimate  $I$ :

$$(F2) \quad I \sim \int_{\mathbb{R}^{N-1}} \int_0^\infty \int_0^\infty \frac{|u(x', t) - u(x', \sigma)|^p}{|t - \sigma|^{1+sp}} d\sigma dt dx' + \int_{\mathbb{R}^{N-1}} \int_0^\infty \frac{|u(x', t)|^p}{t^{sp}} dt dx'.$$

It then suffices to estimate the last term in (F2). By Fubini, it suffices to consider the 1-dimensional integral

$$\int_0^\infty \frac{|u(t)|^p}{t^{sp}} dt, \quad u \in C_0^\infty(0, \infty).$$

This is a trivial consequence of the following

**Lemma F2.** *Let  $1/p < s < 1$  and  $1 < p < \infty$ . Given  $u \in C_0^\infty(0, \infty)$ , we have*

$$(F3) \quad \int_0^\infty \frac{|u(t)|^p}{t^{sp}} dt \leq C \int_0^\infty \int_0^\infty \frac{|u(x) - u(y)|^p}{|x - y|^{1+sp}} dx dy.$$

*Proof.* We first point out that both integrals are finite. Given  $0 < \alpha < \beta < 1$ , we have

$$|u(t)| \leq |u(x)| + |u(t) - u(x)| \quad \forall x \in [\alpha t, \beta t] =: I_t.$$

Thus,

$$|u(t)|^p \leq 2|u(x)|^p + C|u(t) - u(x)|^p \quad \forall x \in I_t.$$

Integrating over  $x \in I_t$ , we get

$$(F4) \quad |u(t)|^p \leq \frac{2}{(\beta - \alpha)t} \int_0^\infty |u(x)|^p dx + \frac{C}{(\beta - \alpha)t} \int_{I_t} |u(t) - u(x)|^p dx.$$

Since  $\alpha, \beta < 1$ , we have  $|t - x| \sim t$  for every  $x \in I_t$ . Integrating (F4) with respect to  $t$  we then have

$$(F5) \quad \begin{aligned} \int_0^\infty \frac{|u(t)|^p}{t^{sp}} dt &\leq \frac{2}{\beta - \alpha} \int_0^\infty \int_{I_t} \frac{|u(x)|^p}{t^{sp}} dx dt + \frac{C}{\beta - \alpha} \int_0^\infty \int_0^\infty \frac{|u(t) - u(x)|^p}{|t - x|^{1+sp}} dx dt \\ &= \frac{2(\beta^{sp} - \alpha^{sp})}{sp(\beta - \alpha)} \int_0^\infty \frac{|u(x)|^p}{x^{sp}} dx + \frac{C}{\beta - \alpha} \int_0^\infty \int_0^\infty \frac{|u(t) - u(x)|^p}{|t - x|^{1+sp}} dx dt. \end{aligned}$$

Note that  $\frac{\beta^{sp} - \alpha^{sp}}{sp(\beta - \alpha)} = \gamma^{sp-1}$  for some  $\gamma \in [\alpha, \beta]$ . Thus, by taking  $\beta \ll 1/2$  sufficiently small, we have

$$\frac{2(\beta^{sp} - \alpha^{sp})}{sp(\beta - \alpha)} \leq \frac{1}{2}.$$

With such choice, (F3) trivially follows from (F5).

**Lemma F3.** *Let  $\omega_1, \Omega$  be two smooth domains,  $\omega_1 \subset\subset \Omega$ . Set  $\omega_2 = \Omega \setminus \bar{\omega}_1$ . Assume that  $u_n \rightarrow u$  in  $W^{s,p}(\omega_1)$  and  $v_n \rightarrow v$  in  $W^{s,p}(\omega_2)$ , with  $\text{tr } u_n = \text{tr } v_n$  on  $\partial\omega_1$ . Let*

$$w_n = \begin{cases} u_n & \text{in } \omega_1 \\ v_n & \text{in } \omega_2 \end{cases} \quad \text{and} \quad w = \begin{cases} u & \text{in } \omega_1 \\ v & \text{in } \omega_2 \end{cases}.$$

Then

$$w_n \rightarrow w \quad \text{in } W^{s,p}(\Omega).$$

*Proof.* It suffices to show that

$$\|w\|_{W^{s,p}(\Omega)} \leq C(\|u\|_{W^{s,p}(\omega_1)} + \|v\|_{W^{s,p}(\omega_2)}).$$

Let  $\eta = \operatorname{tr} u = \operatorname{tr} v$ . By the standard trace theory,

- (i)  $\|\eta\|_{W^{s-1/p,p}} \leq C\|u\|_{W^{s,p}(\omega_1)}$  ;
- (ii)  $\|\eta\|_{W^{s-1/p,p}} \leq C\|v\|_{W^{s,p}(\omega_2)}$  ;
- (iii) there exists an extension  $g \in W^{s,p}(\Omega)$  of  $\eta$  to  $\omega_1$  and  $\omega_2$  such that

$$\|g\|_{W^{s,p}(\Omega)} \leq C\|\eta\|_{W^{s-1/p,p}}.$$

Let

$$\tilde{w} = \begin{cases} u - g & \text{in } \omega_1, \\ v - g & \text{in } \omega_2. \end{cases}$$

By Lemma F1,  $\tilde{w} \in W^{s,p}(\Omega)$  and

$$\|\tilde{w}\|_{W^{s,p}(\Omega)} \leq C(\|u\|_{W^{s,p}(\omega_1)} + \|v\|_{W^{s,p}(\omega_2)}).$$

Since  $w = \tilde{w} + g$  a.e. in  $\Omega$ , Lemma F.3 follows.

In the sequel, we shall denote by  $C$  the cube  $(-1, 1)^N$ . Let  $\|x\|_\infty = \operatorname{Max}_i \{|x_i|\}$ .

**Lemma F4 (Brezis-Mironescu [BM1]).** *Let  $0 < s < 1$  and  $1 < p < \infty$ , with  $sp < N$ . Given  $f \in W^{s,p}(\partial C)$ , set  $\tilde{f}(x) = f(x/\|x\|_\infty)$ ,  $x \in C$ . Then,  $\tilde{f} \in W^{s,p}(C)$  and the mapping*

$$f \mapsto \tilde{f}$$

*is continuous from  $W^{s,p}(\partial C)$  into  $W^{s,p}(C)$ .*

We refer the reader to [BM1, Lemma D.1] for a proof of Lemma F4.

We denote by  $C_\varepsilon = (-\varepsilon, \varepsilon)^N$ . The following lemma is a variant of the approximation procedure in [BM2]:

**Lemma F5.** *Let  $0 < s < 1$  and  $1 < p < \infty$ , with  $sp < N$ . Let  $f \in W^{s,p}(\mathbb{R}^N)$ . Given  $\varepsilon > 0$  and  $Q \in C_\varepsilon$ , set*

$$f_{\varepsilon,Q}(x) = \begin{cases} f(x) & \text{if } x \notin C_\varepsilon(Q), \\ f(\pi_{\varepsilon,Q}(x)) & \text{if } x \in C_\varepsilon(Q), \end{cases}$$

*where  $\pi_{\varepsilon,Q}(x) = Q + \varepsilon \frac{x-Q}{\|x-Q\|_\infty}$  is the projection of  $x$  to  $\partial C_\varepsilon(Q)$ , with respect to  $Q$ . Then, there exist  $\varepsilon_n \rightarrow 0$  and  $Q_n \in C_{\varepsilon_n/2}$  such that*

$$f_{\varepsilon_n, Q_n} \rightarrow f \quad \text{in } W^{s,p}(\mathbb{R}^N).$$

*Proof.* Set  $g_{\varepsilon,Q} = f_{\varepsilon,Q} - f$ . We have

$$\begin{aligned}
 |g_{\varepsilon,Q}|_{W^{s,p}}^p &= 2 \int_{\mathbb{R}^N \setminus C_\varepsilon(Q)} dy \int_{C_\varepsilon(Q)} \frac{|f_{\varepsilon,Q}(x) - f(x)|^p}{|x-y|^{N+sp}} dx + \\
 &\quad + \int_{C_\varepsilon(Q)} \int_{C_\varepsilon(Q)} \frac{|g_{\varepsilon,Q}(x) - g_{\varepsilon,Q}(y)|^p}{|x-y|^{N+sp}} dx dy \\
 &\sim \int_{C_\varepsilon(Q)} \frac{|f_{\varepsilon,Q}(x) - f(x)|^p}{d(x, \partial C_\varepsilon(Q))^{sp}} dx + \\
 &\quad + \int_{C_\varepsilon(Q)} \int_{C_\varepsilon(Q)} \frac{|f_{\varepsilon,Q}(x) - f_{\varepsilon,Q}(y)|^p}{|x-y|^{N+sp}} dx dy + o(1) \\
 &=: I_{\varepsilon,Q} + J_{\varepsilon,Q} + o(1).
 \end{aligned}$$

It suffices to show that

$$\int_0^1 \frac{d\varepsilon}{\varepsilon} \left\{ \int_{C_\varepsilon} (I_{\varepsilon,Q} + J_{\varepsilon,Q}) dQ \right\} < \infty.$$

The proof of

$$\int_0^1 \frac{d\varepsilon}{\varepsilon} \int_{C_\varepsilon} J_{\varepsilon,Q} dQ < \infty$$

may be found in [BM2, Appendix A]. Next,

$$\int_0^1 \frac{d\varepsilon}{\varepsilon} \int_{C_\varepsilon} I_{\varepsilon,Q} dQ \leq \int_0^1 \frac{d\varepsilon}{\varepsilon} \int_{C_{2\varepsilon}} dx \int_{C_\varepsilon(x)} \frac{|f_{\varepsilon,Q}(x) - f(x)|^p}{d(x, \partial C_\varepsilon(Q))^{sp}} dQ.$$

We now make the change of variables  $Q = x - y$ ,  $y \in C_\varepsilon$ . We get

$$\int_0^1 \frac{d\varepsilon}{\varepsilon} \int_{C_\varepsilon} I_{\varepsilon,Q} dQ \leq \int_0^1 \frac{d\varepsilon}{\varepsilon} \int_{C_{2\varepsilon}} dx \int_{C_\varepsilon} \frac{|f(x) - f(x - y + \varepsilon \frac{y}{\|y\|})|^p}{|y - \varepsilon \frac{y}{\|y\|}|^{sp}} dy.$$

Let  $z = -y + \varepsilon \frac{y}{\|y\|}$ , whose Jacobian is  $O((\frac{\varepsilon}{\|z\|^{N-1}}))$  (see [BM2]). We then have

$$\begin{aligned}
 \int_0^1 \frac{d\varepsilon}{\varepsilon} \int_{C_\varepsilon} I_{\varepsilon,Q} dQ &\leq C \int_0^1 \frac{d\varepsilon}{\varepsilon} \int_{C_{2\varepsilon}} dx \int_{C_\varepsilon} \frac{|f(x) - f(x+z)|^p}{|z|^{N+sp-1}} \varepsilon^{N-1} dz \\
 &\leq C \int_{C_2} dx \int_{C_1} \frac{|f(x) - f(x+z)|^p}{|z|^{N+sp-1}} dz \int_{|z|}^1 \frac{d\varepsilon}{\varepsilon^2} \\
 &\leq \iint \frac{|f(x) - f(x+z)|^p}{|z|^{N+sp}} dx dz < \infty.
 \end{aligned}$$



This concludes the proof of Lemma F5.

Given  $N - 1 < p \leq N$ , any map  $u \in W^{(N-1)/p,p}(S^N; S^{N-1})$  has a harmonic extension  $U \in W^{N/p,p}(B^{N+1}; \mathbb{R}^N) \subset W^{1,N}$ . We then define  $T(u)$  as

$$\langle T(u), \zeta \rangle = - \sum_{j=1}^N \int_{B^{N+1}} \det(U_{x_1}, \dots, U_{x_j}, u, U_{x_{j+1}}, \dots, U_{x_{N+1}}) \xi_{x_j},$$

where  $\zeta \in \text{Lip}(S^N)$  and  $\xi$  is any extension of  $\zeta$  to  $B^{N+1}$ . One can see that this definition is independent of the extension  $\xi$ . Let

$$L(u) = \frac{1}{\sigma_N \|\nabla \zeta\|_{L^\infty} \leq 1} \text{Max} \langle T(u), \zeta \rangle.$$

We have the following

**Lemma F6.** *Assume  $N \geq 2$ . Let  $1 < p < \infty$  and  $u \in W^{(N-1)/p,p}(S^N; S^{N-1})$ . If  $T(u) = 0$ , then  $u \in \overline{C^\infty(S^N; S^{N-1})}^{W^{(N-1)/p,p}}$ .*

*Proof.*

*Case 1.* Proof of the lemma if  $N \geq 3$ .

Note that good maps are dense in  $W^{1,N-1}(S^N; S^{N-1})$  and, by interpolation, in  $W^{(N-1)/p,p}(S^N; S^{N-1})$ . Thus, it suffices to show that if  $u$  is a good map, then there exists  $v \in C^\infty(S^N; S^{N+1})$  such that

$$\|u - v\|_{W^{1,N-1}} \leq CL(u),$$

which can be done by a dipole construction. By interpolation, we obtain Lemma F6.

*Case 2.* Proof of the lemma if  $N = 2$ .

The interpolation argument does not work in this case. However, for any map  $u \in \mathcal{R}$ , the dipole construction in [BBM2] gives a sequence  $(v_n)$  such that

$$T(v_n) = T(u), \quad |v_n|_{W^{1/p,p}}^p \leq CL(u), \quad \text{and} \quad v_n \rightarrow 1 \quad \text{a.e.}$$

Clearly,  $u\bar{v}_n \in \overline{C^\infty}^{W^{1/p,p}}$  and

$$\begin{aligned} |u - u\bar{v}_n|_{W^{1/p,p}}^p &\leq 2|1 - \bar{v}_n|_{W^{1/p,p}}^p + K \iint |1 - \bar{v}_n(x)|^p \frac{|u(x) - u(y)|^p}{|x - y|^2} dx dy \\ &\leq 2CL(u) + o(1). \end{aligned}$$

Using the density of  $\mathcal{R}$  in  $W^{1/p,p}(S^2; S^1)$ , we obtain the desired result.

We also point out the following extension of Lemma F6 whose proof is left to the reader :

**Lemma F7.** *Assume  $N - 1 < p \leq N$ . Let  $\Omega$  be a smooth subdomain of  $S^N$ . For any  $u \in W^{(N-1)/p,p}(S^N; S^{N-1})$ ,  $T(u|_\Omega)$  is well-defined when computed against Lipschitz functions which are constant on each connected component of  $\partial\Omega$ . If  $T(u|_\Omega) = 0$ , then  $u|_\Omega \in \overline{C^\infty(\overline{\Omega}; S^{N-1})}^{W^{(N-1)/p,p}}$ .*

A key ingredient in the proof of Theorem 5''' is the following

**Proposition F1.** *Assume  $N - 1 < p \leq N$ . Let  $u \in W^{(N-1)/p,p}(S^N; S^{N-1})$  be such that*

$$T(u) = \sigma_N \sum_{\text{finite}} d_i \delta_{M_i}.$$

*Then, there exist  $(u_n) \subset W^{(N-1)/p,p}(S^N; S^{N-1})$ ,  $\varepsilon_n \rightarrow 0$ , and  $M_i^n \rightarrow M_i$  such that*

- (i)  $u_n \in C^\infty(S^N \setminus \bigcup_i C_{\varepsilon_n}(M_i^n); S^{N-1})$  ;
- (ii)  $u_n$  is homogeneous of degree 0 on each cube  $C_{\varepsilon_n}(M_i^n)$  ;
- (iii)  $T(u_n) = \sigma_N \sum_{\text{finite}} d_i \delta_{M_i^n}$  ;
- (iv)  $u_n \rightarrow u$  in  $W^{(N-1)/p,p}$ .

*Proof.* We first observe that the set of good pairs  $(\varepsilon, Q)$ , in the sense of Lemma F5, is “fat”. More precisely, there exists a sequence  $\varepsilon_n \rightarrow 0$  such that

$$\frac{|\{Q \in C_{\varepsilon_n/4} ; |f_{\varepsilon_n, Q} - f|_{W^{(N-1)/p,p}} < 1/n\}|}{|C_{\varepsilon_n/4}|} \geq \frac{1}{2}.$$

In particular,  $\{Q \in C_{\varepsilon_n/4} ; |f_{\varepsilon_n, Q} - f|_{W^{(N-1)/p,p}} < 1/n\}$  intersects the complement of any null set of  $C_{\varepsilon_n/4}$ . For  $n \geq 1$  fixed, consider

$$v_n = u|_{S^N \setminus \bigcup_i C_{\varepsilon_n/4}(M_i)}.$$

Then,  $T(v_n) = 0$ , so that there exists a sequence  $(v_n^k)_k$ ,

$$v_n^k \in C^\infty(S^N \setminus \bigcup_i C_{\varepsilon_n/4}(M_i); S^{N-1}) \quad \forall k \geq 1,$$

such that

$$v_n^k \rightarrow v_n \quad \text{in } W^{(N-1)/p,p}(S^N \setminus \bigcup_i C_{\varepsilon_n/4}(M_i)) \quad \text{as } k \rightarrow \infty.$$

By Fubini, for a.e.  $Q \in C_{\varepsilon_n/4}$ , we have

$$v_n^k|_{\bigcup_i \partial C_{\varepsilon_n}(Q+M_i)} \rightarrow u|_{\bigcup_i \partial C_{\varepsilon_n}(Q+M_i)} \quad \text{in } W^{(N-1)/p,p} \quad \text{as } k \rightarrow \infty.$$

By Lemmas F3 and F4, for any such  $Q$  we have

$$\tilde{v}_n^k \rightarrow \tilde{u}_n \quad \text{in } W^{(N-1)/p,p} \quad \text{as } k \rightarrow \infty,$$

where  $\tilde{v}_n^k$  (resp.,  $\tilde{u}_n$ ) is  $v_n^k$  (resp.,  $u$ ) extended by homogeneity of degree 0 on each cube  $C_{\varepsilon_n}(Q + M_i)$ . By Lemma F5, we can choose  $Q = Q_n$  such that  $\tilde{u}_n \rightarrow u$  in  $W^{(N-1)/p,p}$ . Applying a diagonalization argument,  $u_n$  may be taken among  $(\tilde{u}_n^k)$ . We only need to show that (iii) holds.

Note that  $\tilde{v}_n^k$  is locally Lipschitz on  $S^N \setminus \bigcup_i M_i^n$ , where  $M_i^n = Q_n + M_i$ . Thus,

$$T(\tilde{v}_n^k) = \sigma_N \sum_{\text{finite}} \tilde{d}_i \delta_{M_i^n}.$$

Since  $T(u_n) \rightarrow T(u)$ , for  $n$  large enough we have (iii). This concludes the proof of Proposition F1.

We may now present the

*Proof of Theorem 5'''.*

*Step 1.* If  $TV(u) < \infty$ , then  $\text{Det}(\nabla u)$  is a measure and

$$(F6) \quad |\text{Det}(\nabla u)|_{\mathcal{M}} \leq TV(u).$$

Clearly, we may replace  $u_n$  in the definition of  $TV(u)$  by

$$\tilde{u}_n = \begin{cases} u_n & \text{if } |u_n| \leq 1, \\ \frac{u_n}{|u_n|} & \text{if } |u_n| > 1. \end{cases}$$

We may thus assume that  $|u_n| \leq 1$ . Since  $u_n \rightarrow u$  in  $W^{(N-1)/p,p}$  and  $|u_n| \leq 1$ , we have

$$\langle T(u_n), \zeta \rangle \rightarrow \langle T(u), \zeta \rangle$$

for every  $\zeta \in \text{Lip}(S^N; S^{N-1})$ . In addition,

$$\langle T(u_n), \zeta \rangle = N \int_{S^{N-1}} \text{Jac } u_n \zeta \leq TV(u) \|\zeta\|_{L^\infty} + o(1).$$

Thus,  $T(u)$  is a measure and (F6) holds.

*Step 2.* If  $\text{Det}(\nabla u)$  is a measure, then  $TV(u) < \infty$  and

$$(F7) \quad \frac{1}{\omega_n} TV(u) = \text{number of topological singularities of } u.$$

By Proposition F1, it suffices to compute  $TV(u)$ , where  $u$  is smooth outside finitely many (disjoint) cubes and  $u$  is homogeneous of degree 0 inside each one of these cubes. By (F6), we have “ $\geq$ ” in (F7). It then suffices to show the reverse inequality.

Note that  $u \in W^{1,q}(S^N; S^{N-1})$  for every  $q < N$ . As in the proof of the case  $W^{1,N-1}(S^N; S^{N-1})$ , we can find  $(u_n) \subset C^\infty$ ,  $u_n \rightarrow u$  in  $W^{1,q}$ , with

$$\frac{1}{\omega_N} \int_{S^{N-1}} |\text{Jac } u_n| = \text{number of topological singularities of } u.$$

For  $N-1 < q < N$ , we have  $W^{1,q} \cap L^\infty \subset W^{(N-1)/p,p}$ , so that  $u_n \rightarrow u$  in  $W^{(N-1)/p,p}$ . We conclude that (F7) holds.

## G. Proofs of Theorems 14–17.

We start by establishing the precise value of  $E(g)$  :

**Lemma G1.** *Let  $g \in BV(I; S^1)$  and let  $A$  be the set of jump points of  $g$ . Then*

$$(G1) \quad E(g) = |\dot{g}_d| + \sum_{a \in A} d_{S^1}(g(a+), g(a-)).$$

*Proof.* Let  $\varphi \in BV(I; \mathbb{R})$  be any lifting of  $g$ . We claim that

$$(G2) \quad |\dot{\varphi}_d|_{\mathcal{M}(I)} = |\dot{g}_d|_{\mathcal{M}(I)}.$$

Indeed, recall that, by the chain rule, we have

$$\dot{\varphi}_d = -i\bar{g}\dot{g}_d.$$

Set  $\nu = \dot{g}_d$  and  $\mu = |\nu|$ . Then there is some  $k \in L^\infty((I, d\mu); S^1)$  such that  $\nu = k\mu$ . Since  $\nu$  is diffuse and  $-i\bar{g} \in BV$ , we have  $-ig \in L^\infty((I, d\mu); S^1)$ , and thus  $\dot{\varphi}_d = \ell\mu$ , where  $\ell = -i\bar{g}k \in L^\infty((I, d\mu); S^1)$ . It follows that

$$|\dot{\varphi}_d|_{\mathcal{M}(I)} = \text{Sup}_{\substack{\zeta \in C_0(I; \mathbb{C}) \\ |\zeta| \leq 1}} \langle \dot{\varphi}_d, \zeta \rangle = \text{Sup}_{\substack{\zeta \in C_0(I; \mathbb{C}) \\ |\zeta| \leq 1}} \langle \mu, \ell\zeta \rangle = \langle \mu, |\ell| \rangle = |\mu|_{\mathcal{M}(I)} = |\dot{g}_d|_{\mathcal{M}(I)}.$$

Let now  $B$  denote the set of the jump points of  $\varphi$ ; clearly,  $B \supset A$ . For each  $a \in A$ , we have

$$\begin{cases} g(a+) = e^{i\varphi(a+)}, \\ g(a-) = e^{i\varphi(a-)}, \end{cases}$$

so that

$$|\varphi(a+) - \varphi(a-)| \geq d_{S^1}(g(a+), g(a-)).$$

If  $a \in B \setminus A$ , then  $e^{i\varphi(a+)} = e^{i\varphi(a-)}$ ; thus

$$|\varphi(a+) - \varphi(a-)| \geq 2\pi.$$

Consequently,

$$\begin{aligned} |\dot{\varphi}|_{\mathcal{M}(I)} &= |\dot{\varphi}_d|_{\mathcal{M}(I)} + \sum_{a \in A} |\varphi(a+) - \varphi(a-)| + \sum_{a \in B \setminus A} |\varphi(a+) - \varphi(a-)| \\ \text{(G3)} \quad &\geq |\dot{g}_d| + \sum_{a \in A} d_{S^1}(g(a+), g(a-)) + 2\pi \text{card}(B \setminus A) \\ &\geq |\dot{g}_d| + \sum_{a \in A} d_{S^1}(g(a+), g(a-)). \end{aligned}$$

Equality holds if and only if

$$|\varphi(a+) - \varphi(a-)| = d_{S^1}(g(a+), g(a-)) \quad \forall a \in A \quad \text{and} \quad B = A.$$

This proves “ $\geq$ ” in (G1).

In order to prove “ $\leq$ ”, we split  $A$  as  $A = A_1 \cup A_2$ , where

$$A_1 = \{a \in A; |g(a+) - g(a-)| = 2\} \quad \text{and} \quad A_2 = \{a \in A; |g(a+) - g(a-)| < 2\}.$$

If  $a \in A_2$  we may define a *signed* distance

$$\delta_{S^1}(g(a+), g(a-)) = \arg \left( \frac{g(a+)}{g(a-)} \right).$$

Here,  $\arg$  stands for the argument in  $(-\pi, \pi)$ . Set

$$\text{(G4)} \quad \mu = -i\bar{g}\dot{g}_d + \pi \sum_{a \in A_1} \delta_a + \sum_{a \in A_2} \delta_{S^1}(g(a+), g(a-)) \delta_a.$$

We claim that  $\mu$  is a measure. Indeed,  $A_1$  is finite, since  $g \in BV$ . On the other hand,

$$\text{Sup}_{a \in A_2} |g(a+) - g(a-)| = d < 2$$

(again since  $g \in BV$ ). Thus there is some  $C > 0$  such that

$$|\delta_{S^1}(g(a+), g(a-))| \leq C|g(a+) - g(a-)|.$$

It follows that

$$\sum_{a \in A_2} |\delta_{S^1}(g(a+), g(a-))| \leq C \sum_{a \in A_2} |g(a+) - g(a-)| < \infty.$$

Assume that  $I = (0, \alpha)$  for some  $\alpha > 0$  and set  $\varphi_0(x) = \mu((0, x))$ ,  $x \in I$ . We claim that, up to a constant,  $\varphi_0$  is a lifting of  $g$  and that  $|\dot{\varphi}_0|_{\mathcal{M}(I)} = E(g)$ .

Indeed, using the chain rule for a product we have

$$(G5) \quad \overline{ge^{-i\varphi_0}} = e^{-i\varphi_0} \dot{g}_d - ig e^{-i\varphi_0} (\dot{\varphi}_0)_d + \sum_{a \in A} (ge^{-i\varphi_0}(a+) - ge^{-i\varphi_0}(a-)) \delta_a.$$

Here, we have used the fact that  $\varphi$  is continuous outside  $A$ . For  $a \in A$ , we have

$$\varphi_0(a+) = \varphi_0(a-) + \mu(\{a\}),$$

so that

$$e^{-i\varphi_0(a+)} = e^{-i\varphi_0(a-)} \frac{g(a-)}{g(a+)},$$

by our definition of  $\mu$ . Thus the sum in (G5) vanishes. On the other hand,

$$(\dot{\varphi}_0)_d = -i\bar{g}\dot{g}_d,$$

so that

$$e^{-i\varphi_0} \dot{g}_d - ig e^{-i\varphi_0} (\dot{\varphi}_0)_d = e^{-i\varphi_0} (\dot{g}_d - \dot{g}_d) = 0.$$

Thus, there is some  $C \in \mathbb{C}$  such that  $\varphi = \varphi_0 + C$  is a lifting of  $g$ .

On the other hand,

$$\begin{aligned} |\dot{\varphi}|_{\mathcal{M}(I)} &= |\dot{\varphi}_0|_{\mathcal{M}(I)} = |-i\bar{g}\dot{g}_d|_{\mathcal{M}(I)} + \pi \operatorname{card}(A_1) + \sum_{a \in A_2} |\delta_{S^1}(g(a+), g(a-))| \\ &= |\dot{g}_d|_{\mathcal{M}(I)} + \sum_{a \in A} d_{S^1}(g(a+), g(a-)). \end{aligned}$$

The proof of Lemma G1 is complete.

*Proof of Theorem 14.* We shall prove a slightly stronger assertion, which implies all the properties stated in the theorem. Namely, a lifting  $\varphi \in BV(I; \mathbb{R})$  of  $g$  is a canonical lifting if and only if

- (i)  $\varphi$  and  $g$  have the same jump sets ;
- (ii) for  $a \in A_1$ , we have  $\varphi(a+) - \varphi(a-) = \pm\pi$  ;
- (iii) for  $a \in A_2$ , we have  $\varphi(a+) - \varphi(a-) = \delta_{S^1}(g(a+), g(a-))$ .

Property (i) was seen to be necessary for optimality in the proof of Lemma G1. Recall that, in addition to (i), equality in (G3) amounts to

$$(G6) \quad |\varphi(a+) - \varphi(a-)| = d_{S^1}(g(a+), g(a-)) \quad \forall a \in A.$$

If  $a \in A_1$ , then

$$|\varphi(a+) - \varphi(a-)| = \pi,$$

so that (ii) holds. Assume  $a \in A_2$ . Since  $e^{i\varphi(a+)} = g(a+)$  and  $e^{i\varphi(a-)} = g(a-)$ , then by (G6) we have

$$\varphi(a+) - \varphi(a-) = \arg \left( \frac{g(a+)}{g(a-)} \right),$$

which gives (iii). Conversely, it is easy to see that, if (i)–(iii) are fulfilled, then equality holds in (G3).

*Proof of Theorem 15.* We identify  $S^1 \setminus \{z\}$  with an interval  $I$ . Let  $A, A_1, A_2$  be defined as in the proof of Lemma G1. We claim that, for *each* choice of integers  $\varepsilon_a \in \{-1, 1\}$ ,  $a \in A_1$ , there is a canonical lifting  $\varphi$  of  $g$  on  $I$  such that

$$\varphi(a+) - \varphi(a-) = \varepsilon_a \pi \quad \forall a \in A_1.$$

This  $\varphi$  is obtained as in the proof of Lemma G1, as  $\varphi = \mu((0, x)) + C$ ,  $x \in I$ . One simply has to modify the definition of  $\mu$  by taking

$$\mu = -i\bar{g}g'd + \sum_{a \in A_1} \varepsilon_a \pi \delta_a + \sum_{a \in A_2} \delta_{S^1}(g(a+), g(a-)) \delta_a.$$

Moreover, the proof of Theorem 14 shows that, by this procedure, we obtain *all* canonical liftings. We claim that if  $\varphi$  is the canonical lifting corresponding to the choice  $\varepsilon_a$ ,  $a \in A_1$ , and  $\tilde{\varphi}$  the one corresponding to  $\tilde{\varepsilon}_a$ ,  $a \in A_1$ , then

$$\frac{\tilde{\varphi}(z-) - \tilde{\varphi}(z+)}{2\pi} = \frac{\varphi(z-) - \varphi(z+)}{2\pi} + \frac{1}{2} \sum_{a \in A_1} (\tilde{\varepsilon}_a - \varepsilon_a) \in \mathbb{Z}.$$

If we identify  $S^1 \setminus \{z\}$  with  $I = (0, \alpha)$ ,  $\alpha > 0$ , this amounts to proving that

$$\frac{\tilde{\varphi}(\alpha-) - \tilde{\varphi}(0+)}{2\pi} = \frac{\varphi(\alpha-) - \varphi(0+)}{2\pi} + \frac{1}{2} \sum_{a \in A_1} (\tilde{\varepsilon}_a - \varepsilon_a).$$

We have  $\tilde{\varphi}(\alpha-) - \tilde{\varphi}(0+) = \tilde{\mu}(I)$  (where  $\tilde{\mu}$  is the corresponding measure); a similar assertion holds for  $\varphi$ . Thus

$$\frac{\tilde{\varphi}(\alpha-) - \tilde{\varphi}(0)}{2\pi} - \frac{\varphi(\alpha-) - \varphi(0+)}{2\pi} = \frac{(\mu - \tilde{\mu})(I)}{2\pi} = \frac{1}{2} \sum_{a \in A_1} (\tilde{\varepsilon}_a - \varepsilon_a).$$

Let  $\varphi$  be the canonical lifting corresponding to the choice  $\varepsilon_a = -1$ ,  $a \in A_1$ . Then it is clear that, with  $d = \frac{\varphi(z-) - \varphi(z+)}{2\pi}$  and  $k = \text{card } A_1$ , we have

$$\text{Deg}_1 g = \{d, d + d, \dots, d + k\}.$$

We next prove that  $d$  is an integer. This follows easily from the fact that  $e^{i\varphi(z-)} = e^{i\varphi(z+)}$ , so that  $\varphi(z-) - \varphi(z+) \in 2\pi\mathbb{Z}$ .

It remains to establish that  $\text{Deg}_1 g$  does not depend on the choice of  $z$ . Let  $w$  be any other continuity point of  $g$ . Let  $\psi$  be a canonical lifting of  $g$  on  $S^1 \setminus \{w\}$ . Since  $g$  is continuous at  $w$ , there is some  $k \in \mathbb{Z}$  such that  $\psi(w-) = \psi(w+) + 2k\pi$ . We set

$$\varphi(\xi) = \begin{cases} \psi(\xi) & \text{if } \xi \in (w+, z-), \\ \psi(\xi) - 2k\pi & \text{if } \xi \in (z+, w-). \end{cases}$$

Clearly,  $\varphi \in BV$  and  $\varphi$  is continuous at  $w$ . It is obvious that

$$|\dot{\varphi}|_{\mathcal{M}(S^1 \setminus \{z\})} = |\dot{\psi}|_{\mathcal{M}(S^1 \setminus \{z, w\})} = |\dot{\psi}|_{\mathcal{M}(S^1 \setminus \{w\})}.$$

It follows that  $\varphi$  is a canonical lifting of  $g$  on  $S^1 \setminus \{z\}$ . Indeed, by Lemma G1 we have

$$E(g|_{S^1 \setminus \{z\}}) = E(g|_{S^1 \setminus \{w\}}) = |g|_{BV S^1}$$

if  $z, w$  are continuity points of  $g$ . Since

$$\begin{aligned} \varphi(z-) - \varphi(z+) &= \varphi(z-) - \varphi(w+) + \varphi(w+) - \varphi(w-) + \varphi(w-) - \varphi(z+) \\ &= \psi(w-) - \psi(w+), \end{aligned}$$

we find that the degrees obtained by cutting at  $z$  are among the ones obtained by cutting at  $w$ . By reversing the roles, we obtained that  $\text{Deg}_1 g$  is independent of  $z$ .

*Proof of Theorem 16.* Let  $z$  be a continuity point of  $g$  and let  $\varphi$  be a canonical lifting of  $g$  in  $S^1 \setminus \{z\}$ . Assume, e.g., that  $z = 1$ ; we identify  $S^1 \setminus \{1\}$  with  $(0, 2\pi)$ . Consider a sequence  $(\varphi_n) \subset C^\infty([0, 2\pi])$  such that

$$\int_0^{2\pi} |\dot{\varphi}_n| \rightarrow |\dot{\varphi}|_{\mathcal{M}((0, 2\pi))} \quad \text{and} \quad \varphi_n \rightarrow \varphi \quad \text{a.e.}$$



We may assume, in addition, that

$$\varphi_n(0) \rightarrow \varphi(0+) \quad \text{and} \quad \varphi_n(2\pi) \rightarrow \varphi(2\pi-).$$

(This is the case, e.g., if the functions  $\varphi_n$  are obtained from  $\varphi$  by mollification).

By replacing  $\varphi_n$  with  $\varphi_n + \alpha_n x + \beta_n$ , for some appropriate  $\alpha_n \rightarrow 0$  and  $\beta_n \rightarrow 0$ , we may further assume that

$$\varphi_n(0) = \varphi(0+) \quad \text{and} \quad \varphi_n(2\pi) = \varphi(2\pi-) \quad \forall n \geq 1.$$

Set  $g_n = e^{i\varphi_n}$ . Then, clearly,

$$g_n \in C^\infty(S^1 \setminus \{z\}) \cap C^0(S^1) \quad \text{and} \quad \deg g_n = \frac{\varphi(1-) - \varphi(1+)}{2\pi}.$$

By further mollifying  $g_n$ , we find a sequence  $(h_n) \subset C^\infty(S^1; S^1)$  such that

$$h_n \rightarrow g \quad \text{a.e.}, \quad \deg h_n = \frac{\varphi(1-) - \varphi(1+)}{2\pi} \quad \forall n \geq 1,$$

and

$$\int_{S^1} |\dot{h}_n| \rightarrow \int_{S^1 \setminus \{z\}} |\dot{\varphi}| = E(g|_{S^1 \setminus \{z\}}).$$

It follows that

$$\text{Deg}_2 g \supset \text{Deg}_1 g.$$

Conversely, let  $d \in \text{Deg}_2 g$  and let  $(g_n) \subset C^\infty(S^1; S^1)$  be such that

$$g_n \rightarrow g \quad \text{a.e.}, \quad \int_{S^1} |\dot{g}_n| \rightarrow |g|_{BV S^1} \quad \text{and} \quad \deg g_n = d \quad \forall n \geq 1.$$

Let  $z \in S^1$  be a continuity point of  $g$ . Write, in  $S^1 \setminus \{z\}$ ,  $g_n = e^{i\varphi_n}$ . Then

$$\int_{S^1} |\dot{\varphi}_n| \rightarrow |g|_{BV S^1}.$$

Up to some subsequence and after subtracting a suitable multiple of  $2\pi$ , we may assume that  $\varphi_n \rightarrow \varphi$  a.e., where  $\varphi \in BV$  is a lifting of  $g$ . Since  $|\dot{\varphi}|_{\mathcal{M}(S^1 \setminus \{z\})} \leq |g|_{BV S^1}$ , we find that  $\varphi$  has to be a canonical lifting of  $g$ . Let  $\varepsilon > 0$ . There is some  $\delta > 0$  such that, if  $I$  is the interval of size  $\delta$  centered at  $z$ , then we have  $|g|_{BV S^1(I)} < \varepsilon$ . We may further assume that  $g$  is continuous at the endpoints of  $I$ . Then

$$|g|_{BV S^1(S^1)} = |g|_{BV S^1(I)} + |g|_{BV S^1(S^1 \setminus I)}.$$

Arguing as above, we find that

$$\int_{S^1 \setminus I} |\dot{\varphi}_n| \rightarrow |g|_{BV S^1(S^1 \setminus I)} \quad \text{and} \quad \int_{I \setminus \{z\}} |\dot{\varphi}_n| \rightarrow |g|_{BV S^1(I)}.$$

In particular, for every  $n \geq 1$  sufficiently large,

$$|(\varphi_n(z-t) - \varphi_n(z+t)) - (\varphi_n(z-) - \varphi_n(z+))| < 2\varepsilon \quad \forall t \in (0, \delta).$$

We pick such  $t$  so that, in addition,

$$\varphi_n(z-t) \rightarrow \varphi(z-t) \quad \text{and} \quad \varphi_n(z+t) \rightarrow \varphi(z+t).$$

We then find

$$|\varphi(z-t) - \varphi(z+t) - 2\pi d| \leq 2\varepsilon.$$

Letting  $\varepsilon \rightarrow 0$  and  $\delta \rightarrow 0$ , we obtain

$$\varphi(z-) - \varphi(z+) = 2\pi d,$$

i.e.,  $d \in \text{Deg}_1 g$ .

We complete the proof of Theorem 16 by proving that  $g \mapsto \text{Deg } g$  is continuous in the multivalued sense. Since  $\text{Deg}$  is  $\mathbb{Z}$ -valued, this amounts to proving that, for each  $d \in \mathbb{Z}$ , the set

$$\{g \in BV(S^1; S^1) ; d \in \text{Deg } g\}$$

is open.

To this purpose, we start with the following

**Lemma G2.** *Let  $g \in BV(I; S^1)$ . Let  $\varphi \in BV(I; \mathbb{R})$  be a lifting of  $g$ . If  $\varphi$  is not a canonical lifting of  $g$ , then*

$$|\dot{\varphi}|_{\mathcal{M}(I)} \geq E(g) + \pi.$$

*Proof.* Let  $A, B$  be the set of jump points of  $g, \varphi$ , respectively. Recall that

$$\begin{aligned} |\dot{\varphi}|_{\mathcal{M}(I)} &\geq |\dot{g}d|_{\mathcal{M}(I)} + \sum_{a \in A} |\varphi(a+) - \varphi(a-)| + \sum_{a \in B \setminus A} [|\varphi(a+) - \varphi(a-)|] \\ &\geq E(g) + \sum_{a \in B \setminus A} |\varphi(a+) - \varphi(a-)|. \end{aligned}$$

If  $B \neq A$ , then

$$|\varphi(a+) - \varphi(a-)| \geq 2\pi \quad \forall a \in B \setminus A,$$

and the conclusion is clear. If  $B = A$ , then there is some  $a \in A$  such that

$$|\varphi(a+) - \varphi(a-)| > d_{S^1}(g(a+), g(a-)),$$

for otherwise  $\varphi$  would be a canonical lifting. For any such  $a$ , we have

$$|\varphi(a+) - \varphi(a-)| \equiv d_{S^1}(g(a+), g(a-)) \pmod{2\pi}.$$

Since  $d_{S^1}(g(a+), g(a-)) \leq \pi$ , we find that

$$|\varphi(a+) - \varphi(a-)| \geq d_{S^1}(g(a+), g(a-)) + \pi.$$

Since, for any  $b \in A \setminus \{a\}$ , we have

$$|\varphi(b+) - \varphi(b-)| \geq d_{S^1}(g(b+), g(b-)),$$

and the conclusion follows.

*Proof of Theorem 16 completed.* If  $g, h \in BV(S^1; S^1)$ , then

$$(G7) \quad |g\bar{h}|_{BV} = |g\bar{h} - 1|_{BV} = |g(\bar{h} - \bar{g})|_{BV} \leq |g|_{BV} \|h - g\|_{L^\infty} + |h - g|_{BV}.$$

Let  $g \in BV(S^1; S^1)$  and let  $d \in \text{Deg } g$ . In view of (G7), there is some  $\varepsilon > 0$  such that if

$$h \in BV(S^1; S^1) \quad \text{and} \quad \|g - h\|_{BV} < \varepsilon,$$

then

$$|g\bar{h}|_{BV} < \frac{1}{10}.$$

We claim that  $d \in \text{Deg } h$  for any such  $h$ . Indeed, let  $z$  be a continuity point for both  $g$  and  $h$ , and let  $\varphi$  be a canonical lifting of  $g$  in  $S^1 \setminus \{z\}$ . Set  $k = \bar{g}h$  and let  $\psi$  be a canonical lifting of  $k$ . Since  $|k|_{BV} < \frac{1}{10}$ , each jump point  $a$  of  $k$  is such that  $|k(a+) - k(a-)| < \frac{1}{10}$ . Thus

$$|\psi(a+) - \psi(a-)| \leq 2|k(a+) - k(a-)|$$

for any such  $a$ . It follows that

$$|\psi|_{BV} = |\dot{k}_d|_{\mathcal{M}(I)} + \sum_{\substack{\text{jump points} \\ \text{of } k}} |\psi(a+) - \psi(a-)| \leq 2|k|_{BV} < \frac{1}{5}.$$

Set  $\phi = \varphi + \psi$ . Then  $\phi$  is a lifting of  $h$  and

$$|(\phi(z-) - \phi(z+)) - (\varphi(z-) - \varphi(z+))| < \frac{2}{5},$$

so that

$$\phi(z-) - \phi(z+) = \varphi(z-) - \varphi(z+)$$

(since both quantities are multiple of  $2\pi$ ). In order to complete the proof of Theorem 16, it suffices to prove that  $\phi$  is a canonical lifting of  $h$ .

Indeed, on the one hand we have

$$E(h) \leq |\dot{\phi}|_{\mathcal{M}(S^1 \setminus \{z\})} \leq |\dot{\varphi}|_{\mathcal{M}(S^1 \setminus \{z\})} + \frac{1}{5},$$

so that  $E(h) \leq E(g) + \frac{1}{5}$ . By reversing the roles, we obtain on the other hand that  $E(g) \leq E(h) + \frac{1}{5}$ ; thus

$$E(h) \leq |\dot{\phi}|_{\mathcal{M}(S^1 \setminus \{z\})} \leq E(h) + \frac{2}{5}.$$

Lemma G2 implies that  $\phi$  is a canonical lifting of  $h$ .

*Proof of Theorem 17.* With the notation we already used, we have

$$\{g ; \text{Deg } g \text{ is single-valued}\} = \{g ; A_1(g) = \phi\} =: \mathcal{U}_1.$$

Thus, we have to prove that  $\mathcal{U}_1$  is dense in  $BV(S^1; S^1)$ .

Let  $g \in BV(S^1; S^1)$ ; then  $A_1(g)$  is finite. If  $A_1(g) = \phi$ , then  $g \in \mathcal{U}_1$ . Otherwise, we may assume, for simplicity, that  $A_1$  consists of a single point, say  $A_1 = \{1\}$ ; the general case can be treated along the same lines. We have  $|g(1-) - g(1+)| = 2$ . Without loss of generality, we may assume that  $g(1-) = -1$  and  $g(1+) = 1$ .

Given  $\varepsilon > 0$ , let  $h_\varepsilon : S^1 \rightarrow S^1$  be given by

$$h_\varepsilon(e^{i\theta}) = \begin{cases} e^{i\theta} & \text{if } 1 \leq \theta \leq 2\pi - 1, \\ e^{i(\varepsilon + (1-\varepsilon)\theta)} & \text{if } 0 \leq \theta \leq 1, \\ e^{i((2\pi-1)(2\pi+\varepsilon) - (2\pi-1+\varepsilon)\theta)} & \text{if } 2\pi - 1 \leq \theta < 2\pi. \end{cases}$$

It is immediate that  $h_\varepsilon(1+) = e^{i\varepsilon}$ ,  $h_\varepsilon(1-) = e^{-i\varepsilon}$ ,

$$|h_\varepsilon|_{BV} \rightarrow 0 \quad \text{and} \quad h_\varepsilon \rightarrow 1 \quad \text{uniformly.}$$

Thus,

$$gh_\varepsilon \rightarrow g \quad \text{in } BV \quad \text{as } \varepsilon \rightarrow 0.$$

On the other hand, since  $h_\varepsilon \in C^0(S^1 \setminus \{1\})$ , we have  $A_1(gh_\varepsilon) \setminus \{1\} = A_1(g) \setminus \{1\}$ . In particular,  $A_1(gh_\varepsilon) \subset \{1\}$ . Since, by construction,  $1 \notin A_1(gh_\varepsilon)$ , we have  $gh_\varepsilon \in \mathcal{U}_1$ . The proof of Theorem 17 is complete.

<sup>(1)</sup> LABORATOIRE J. -L. LIONS  
UNIVERSITÉ P. ET M. CURIE, B.C. 187  
4 PL. JUSSIEU  
75252 PARIS CEDEX 05  
*E-mail address:* brezis@ccr.jussieu.fr, ponce@ann.jussieu.fr

<sup>(2)</sup> RUTGERS UNIVERSITY  
DEPT. OF MATH., HILL CENTER, BUSCH CAMPUS  
110 FRELINGHUYSEN RD, PISCATAWAY, NJ 08854  
*E-mail address:* brezis@math.rutgers.edu, augponce@math.rutgers.edu

<sup>(3)</sup> DÉPARTEMENT DE MATHÉMATIQUES  
UNIVERSITÉ PARIS-SUD  
91405 ORSAY  
*E-mail address:* Petru.Mironescu@math.u-psud.fr