



HAL
open science

Lifting default for S^1 -valued maps

Petru Mironescu

► **To cite this version:**

Petru Mironescu. Lifting default for S^1 -valued maps. Comptes rendus de l'Académie des sciences. Série I, Mathématique, 2008, 346 (19-20), pp.1093-1044. hal-00747676

HAL Id: hal-00747676

<https://hal.science/hal-00747676>

Submitted on 31 Oct 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lifting default for \mathbb{S}^1 -valued maps

Petru Mironescu *

June 30th, 2008

Abstract

Let $\varphi \in C^\infty([0, 1]^N, \mathbb{R})$. When $0 < s < 1$, $p \geq 1$ and $1 \leq sp < N$, the $W^{s,p}$ -semi-norm $|\varphi|_{W^{s,p}}$ of φ is not controlled by $|g|_{W^{s,p}}$, where $g := e^{i\varphi}$ [3]. [This question is related to existence, for \mathbb{S}^1 -valued maps g , of a lifting φ as smooth as allowed by g .] In [4], the authors suggested that $|g|_{W^{s,p}}$ does control a weaker quantity, namely $|\varphi|_{W^{s,p}+W^{1,sp}}$. Existence of such control is due to J. Bourgain and H. Brezis [2] when $1 < p \leq 2$, $s = 1/p$ and to H.-M. Nguyen [10] when $N = 1$, $p > 1$ and $sp \geq 1$ or when $N \geq 2$, $p > 1$ and $sp > 1$. In this Note, we establish existence of control for all $s < 1$, $p \geq 1$ and N .

Résumé

Défaut de relèvement pour les applications à valeurs dans le cercle unité. Soit $\varphi \in C^\infty([0, 1]^N, \mathbb{R})$. Si $0 < s < 1$, $p \geq 1$ et $1 \leq sp < N$, alors la semi-norme $|\varphi|_{W^{s,p}}$ n'est pas contrôlée par $|g|_{W^{s,p}}$, où $g := e^{i\varphi}$ [3]. [Cette question est liée à l'existence, pour des g à valeurs dans \mathbb{S}^1 , de relèvements φ aussi réguliers que g le permet.] Dans [4], il est conjecturé que $|g|_{W^{s,p}}$ contrôle une quantité plus faible que $|\varphi|_{W^{s,p}}$, plus spécifiquement $|\varphi|_{W^{s,p}+W^{1,sp}}$. L'existence d'un tel contrôle est due à J. Bourgain and H. Brezis [2] pour $1 < p \leq 2$ et $s = 1/p$ et à H.-M. Nguyen [10] pour $N = 1$, $p > 1$ et $sp \geq 1$ ou pour $N \geq 2$, $p > 1$ et $sp > 1$. Dans cette Note, nous montrons l'existence d'un contrôle pour tout $s < 1$, $p \geq 1$ et N .

Version française abrégée

Soient $C = [0, 1]^N$, $0 < s < \infty$ et $1 \leq p < \infty$. $|\cdot|_{W^{s,p}}$ désigne une semi-norme standard sur l'espace de Sobolev $W^{s,p}(C)$; par exemple, pour $0 < s < 1$ nous considérons la semi-norme de Gagliardo,

$$|u|_{W^{s,p}} = \left(\iint_{C^2} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{1/p}.$$

Si $\varphi \in C^\infty(C, \mathbb{R})$ et $g := e^{i\varphi}$, alors $|\nabla\varphi| = |\nabla g|$. En particulier, $|g|_{W^{1,p}} = |\varphi|_{W^{1,p}}$. Plus généralement, $|g|_{W^{s,p}}$ contrôle $|\varphi|_{W^{s,p}}$ si $s \geq 1$, au sens où il existe une inégalité de la forme $|\varphi|_{W^{s,p}} \leq F(|g|_{W^{s,p}})$ avec F croissante [3]. Ceci n'est plus forcément vrai si $0 < s < 1$. Voici un exemple inspiré de [3] : si $N \geq 2$ et si $0 < s < 1$ et p sont tels que $1 < sp < N$, alors il existe une fonction $\psi \in W^{1,sp} \setminus W^{s,p}$. Si on considère $\varphi_\varepsilon := \psi * \rho_\varepsilon$, avec ρ noyau régularisant, alors $|\varphi_\varepsilon|_{W^{s,p}} \rightarrow \infty$ (car $\psi \notin W^{s,p}$), alors que $g_\varepsilon := e^{i\varphi_\varepsilon}$ reste bornée dans $W^{1,sp} \cap L^\infty$ (car $\psi \in W^{1,sp}$) et donc dans $W^{s,p}$, grâce à l'inclusion de Gagliardo-Nirenberg $W^{1,sp} \cap L^\infty \subset W^{s,p}$. Nonobstant la non inclusion $W^{1,1} \cap L^\infty \not\subset W^{s,p}$ si $sp = 1$, on peut adapter cet exemple au cas où $sp = 1$ et $N \geq 1$. Plus généralement, si $sp = 1$ ou $1 < sp < N$,

*Université de Lyon, Université Lyon1, CNRS, UMR 5208 Institut Camille Jordan, Bâtiment du Doyen Jean Braconnier, 43, blvd du 11 novembre 1918, F - 69200 Villeurbanne Cedex, France. Email address: mironescu@math.univ-lyon1.fr

alors on peut obtenir l'absence du contrôle à partir d'une fonction convenable $\psi \in W^{s,p} + W^{1,sp}$.

Dans le cas particulier $s = 1/2$ et $p = 2$, J. Bourgain et H. Brezis [2] ont montré que le contre-exemple ci-dessus est essentiellement le seul. Leur résultat est que toute fonction φ se décompose comme $\varphi = \varphi_1 + \varphi_2$, où $|\varphi_1|_{H^{1/2}}$ et $|\varphi_2|_{W^{1,1}}$ sont contrôlées par $|g|_{H^{1/2}}$. La preuve s'étend aux espaces $W^{1/p,p}$ avec $1 < p \leq 2$ et donne la décomposition (1). Ce résultat a motivé le problème suivant [4], [8] avec $0 < s < 1$ et $1 \leq p < \infty$:

$(D_{s,p})$ Tout $\varphi \in C^\infty(C, \mathbb{R})$ s'écrit $\varphi = \varphi_1 + \varphi_2$, où $|\varphi_1|_{W^{s,p}} \leq C|e^{i\varphi}|_{W^{s,p}}$ et $\|D\varphi_2\|_{L^{sp}} \leq C|e^{i\varphi}|_{W^{s,p}}^{1/s}$.

Récemment, H.-M. Nguyen [10] a résolu ce problème lorsque $p > 1$, $s = 1/p$ et $N = 1$; son argument s'applique aussi au cas $N \geq 2$, $p > 1$ et $sp > 1$.

Le but de cette Note est d'annoncer le

Théorème 1 *La décomposition $(D_{s,p})$ est valide pour tout $0 < s < 1$, $p \geq 1$ et N .*

Notons que la décomposition existe même si $sp < 1$.

Idée de la preuve. On étend g à une application $h : \mathbb{R}^N \rightarrow \mathbb{R}^2$, de sorte que h soit Lipschitz, constante en dehors d'un compact, $|h| \leq 3$ et $|h|_{W^{s,p}(\mathbb{R}^N)} \leq C|g|_{W^{s,p}(C)}$. On étend ensuite h à $\mathbb{R}^N \times \mathbb{R}_+$ par la formule $w(x, \varepsilon) = \Pi(h * \rho_\varepsilon(x))$. Ici, $\Pi \in C^\infty(\mathbb{R}^2; \mathbb{R}^2)$ est telle que $\Pi(z) = z/|z|$ si $|z| \geq 1/2$, tandis que $\rho \in C_0^\infty$ satisfait $\rho \geq 0$, $\int \rho = 1$, $\text{supp } \rho \subset B(0, 2) \setminus B(0, 1)$. Alors la conclusion du théorème est vérifiée par φ_j données par

$$\varphi_1(x) := - \int_0^\infty w(x, \varepsilon) \wedge \frac{\partial w}{\partial \varepsilon}(x, \varepsilon) d\varepsilon, \quad \varphi_2 := \varphi - \varphi_1.$$

L'inégalité $|\varphi_1|_{W^{s,p}} \leq C|h|_{W^{s,p}}$ découle d'estimations standard¹ pour les régularisées $h * \rho_\varepsilon$ d'une fonction $h \in W^{s,p}$. Elle implique $|\varphi_1|_{W^{s,p}} \leq C|g|_{W^{s,p}}$.

Pour estimer $D\varphi_2$, le point de départ est l'identité²

$$D\varphi_2(x) = -2 \int_0^\infty \frac{\partial}{\partial \varepsilon} w(x, \varepsilon) \wedge D_x w(x, \varepsilon) d\varepsilon.$$

En adaptant une idée de [4], cette identité permet d'obtenir l'estimation $\|D\varphi_2\|_{L^{sp}} \leq C|e^{i\varphi}|_{W^{s,p}}^{1/s}$. \square

Le lecteur trouvera les preuves détaillées dans [9]; entre autres, on y explique pourquoi notre décomposition est une sorte d'analogie continu de la décomposition trouvée dans [2].

1 Introduction

Let $C = [0, 1]^N$, $0 < s < \infty$ and $1 \leq p < \infty$. We will denote by $|\cdot|_{W^{s,p}}$ a standard semi-norm on the Sobolev space $W^{s,p}(C)$; e. g., when $0 < s < 1$, we take the Gagliardo semi-norm

$$|u|_{W^{s,p}} = \left(\iint_{C^2} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{1/p}.$$

Let $\varphi \in C^\infty(C, \mathbb{R})$ and set $g = e^{i\varphi}$. Since $|\nabla\varphi| = |\nabla g|$, we have $|g|_{W^{1,p}} = |\varphi|_{W^{1,p}}$. More generally, $|g|_{W^{s,p}}$ controls $|\varphi|_{W^{s,p}}$ when $s \geq 1$, i. e., there is some non decreasing F such that $|\varphi|_{W^{s,p}} \leq F(|g|_{W^{s,p}})$ [3]. This need not hold when $0 < s < 1$. Here is an example, essentially taken from [3]: let $N \geq 2$

1. L'une des estimations utilisées dans la preuve est $\int_{\mathbb{R}^N} \int_0^\infty \varepsilon^{p-sp-1} |D[h * \rho_\varepsilon](x)|^p d\varepsilon dx \leq C|h|_{W^{s,p}}^p$, bien connue par les spécialistes [11], II.12.

2. Évidente, du moins formellement.

and let $0 < s < 1$, $1 \leq p < \infty$ be such that $1 < sp < N$. By a Sobolev "non embedding", there is some $\psi \in W^{1,sp} \setminus W^{s,p}$. Let $\varphi_\varepsilon := \psi * \rho_\varepsilon$ and set $g_\varepsilon = e^{i\varphi_\varepsilon}$, where ρ is a mollifier. Then $|\varphi_\varepsilon|_{W^{s,p}} \rightarrow \infty$ (since $\psi \notin W^{s,p}$). On the other hand, $g_\varepsilon := e^{i\varphi_\varepsilon}$ is bounded in $W^{1,sp} \cap L^\infty$ (since $\psi \in W^{1,sp}$) and thus in $W^{s,p}$, by the Gagliardo-Nirenberg embedding $W^{1,sp} \cap L^\infty \subset W^{s,p}$. Despite the non embedding $W^{1,1} \cap L^\infty \not\subset W^{s,p}$ when $sp = 1$, one may easily adapt this example to the case $sp = 1$ and $N \geq 1$. More generally, when $sp = 1$ or $1 < sp < N$, one may prove lack of control with the help of an appropriate $\psi \in W^{s,p} + W^{1,sp}$.

In the special case $s = 1/2$ and $p = 2$, J. Bourgain and H. Brezis [2] proved that the above counter-example is essentially the only one. Their results asserts that each φ splits as $\varphi = \varphi_1 + \varphi_2$, where $|\varphi_1|_{H^{1/2}}$ and $|\varphi_2|_{W^{1,1}}$ are controlled by $|g|_{H^{1/2}}$. Their argument adapts steadily to the spaces $W^{1/p,p}$ where $1 < p \leq 2$ and yields

$$\text{If } 1 < p \leq 2, \text{ then } \varphi = \varphi_1 + \varphi_2 \quad \text{where } \varphi_j \in C^\infty(C), |\varphi_1|_{W^{1/p,p}} \leq C|g|_{W^{1/p,p}}, |\varphi_2|_{W^{1,1}} \leq C|g|_{W^{1/p,p}}^p. \quad (1)$$

This motivated the following open problem [4], [8]:

$$(D_{s,p}) \text{ Each } \varphi \in C^\infty(C, \mathbb{R}) \text{ splits as } \varphi = \varphi_1 + \varphi_2, \text{ where } |\varphi_1|_{W^{s,p}} \leq C|e^{i\varphi}|_{W^{s,p}} \text{ and } \|D\varphi_2\|_{L^{sp}} \leq C|e^{i\varphi}|_{W^{s,p}}^{1/s}.$$

[Here, $0 < s < 1$ and $1 \leq p < \infty$.] Very recently, H.-M. Nguyen [10] answered positively this problem when $p > 1$, $s = 1/p$ and $N = 1$; his argument adapts to the case where $N \geq 2$, $p > 1$ and $sp > 1$.

The main purpose of this Note is to announce the following

Theorem 1 *The decomposition $(D_{s,p})$ holds for each $0 < s < 1$, $p \geq 1$ and N .*

Unlike the proofs in [2], [10], our method applies to the case $sp < 1$.

2 Heuristics of the proof of Theorem 1

In order to explain the main idea, we consider, for simplicity, maps defined on \mathbb{R}^N which are constant at infinity (rather than maps defined on $[0, 1]^N$). Assume first that φ has small amplitude oscillations, say $|\varphi| \ll 1$. Then $|g - 1| \ll 1$ and $|\varphi|_{W^{s,p}} \sim |g|_{W^{s,p}}$. Thus, in this case, a convenient decomposition is $\varphi_1 = \varphi$ and $\varphi_2 = 0$. We next proceed as follows: we derive a formula for φ . This formula gives φ only when φ has small amplitude oscillations, but we may give it a meaning for each φ . In general (i. e., when φ may oscillate), we take this formula as the definition of φ_1 and simply let φ_2 be the phase excess, i. e., we set $\varphi_2 = \varphi - \varphi_1$.

In order to obtain a tractable formula for φ , we rely on the following remark: if $g = e^{i\varphi}$, then there is no formula giving φ in terms of g , but there is one for $D\varphi$, since $D\varphi = g \wedge Dg$. We consider a smooth extension $w : \mathbb{R}^N \times [0, +\infty[\rightarrow \mathbb{S}^1$ of g such that

$$\lim_{\varepsilon \rightarrow \infty} w(\cdot, \varepsilon) = \text{const.} \quad (2)$$

A natural choice is $w(x, \varepsilon) = \Pi(g * \rho_\varepsilon(x))$, where $\Pi(z) = z/|z|$ and ρ is a mollifier. This yields a smooth map whenever g (and thus $g * \rho_\varepsilon$) is close to 1. We may write $w = e^{i\psi}$, where $\psi(x, 0) = \varphi(x)$. Assuming that convergence in (2) is sufficiently fast, we then have $\psi(x, \infty) = C$ and thus

$$\varphi(x) = -\psi(x, \varepsilon) \Big|_{\varepsilon=0}^{\varepsilon=\infty} + C = C - \int_0^\infty w(x, \varepsilon) \wedge \frac{\partial w}{\partial \varepsilon}(x, \varepsilon) d\varepsilon. \quad (3)$$

As explained in the next section, (3) gives the right definition of φ_1 , provided we pick an appropriate ρ and we change slightly the definition of Π .

3 Sketch of the proof of Theorem 1

Let $\varphi \in C^\infty(C, \mathbb{R})$ and set $g = e^{i\varphi}$. We first extend g to a Lipschitz map $h : \mathbb{R}^N \rightarrow \mathbb{R}^2$ such that h is constant outside $[-1, 2]^N$, $|h| \leq 3$ and $|h|_{W^{s,p}(\mathbb{R}^N)} \leq C|g|_{W^{s,p}(C)}$. We next extend h to $\mathbb{R}^N \times \mathbb{R}_+$ through the formula $w(x, \varepsilon) = \Pi(h * \rho_\varepsilon(x))$. Here, $\Pi \in C^\infty(\mathbb{R}^2; \mathbb{R}^2)$ is such that $\Pi(z) = z/|z|$ if $|z| \geq 1/2$, while $\rho \in C_0^\infty$ satisfies $\rho \geq 0$, $\int \rho = 1$, $\text{supp } \rho \subset B(0, 2) \setminus B(0, 1)$. Then the conclusion of the theorem holds with φ_j given by

$$\varphi_1(x) := - \int_0^\infty w(x, \varepsilon) \wedge \frac{\partial w}{\partial \varepsilon}(x, \varepsilon) d\varepsilon, \quad \varphi_2 := \varphi - \varphi_1. \quad (4)$$

The inequality

$$|\varphi_1|_{W^{s,p}} \leq C|h|_{W^{s,p}} \quad (5)$$

follows from standard estimates¹ for regularizations $h * \rho_\varepsilon$ of maps $h \in W^{s,p}$. As a consequence of (5), we find that $|\varphi_1|_{W^{s,p}} \leq C|g|_{W^{s,p}}$.

[Estimate (5) is a cousin of the estimate

$$\left| x \mapsto \int_0^\infty u * \rho_\varepsilon(x) \frac{\partial}{\partial \varepsilon} [v * \rho_\varepsilon(x)] d\varepsilon \right|_{W^{s,p}} \leq C\|u\|_{L^\infty} \|v\|_{W^{s,p}}, \quad (6)$$

valid for any reasonable ρ and presumably well-known to experts. For special ρ 's, the "discrete" and much more popular analog of (6) is the "paraproduct inequality" [7]

$$\left| \sum_{j \leq k} u_j v_k \right|_{W^{s,p}} \leq C\|u\|_{L^\infty} \|v\|_{W^{s,p}}, \quad (7)$$

where $u = \sum u_j$, $v = \sum v_j$ are the Littlewood-Paley decompositions of u and v . Both (6) and (7)

are refinements of the standard inequality $\|uv\|_{W^{s,p}} \leq C(\|u\|_{W^{s,p}} \|v\|_{L^\infty} + \|v\|_{W^{s,p}} \|u\|_{L^\infty})$.

The starting point for estimating $D\varphi_2$ is the identity

$$D\varphi_2(x) = -2 \int_0^\infty \frac{\partial}{\partial \varepsilon} w(x, \varepsilon) \wedge D_x w(x, \varepsilon) d\varepsilon. \quad (8)$$

At least formally, this identity follows from

$$\begin{aligned} D\varphi_2(x) &= D\varphi(x) + \int_0^\infty D_x w(x, \varepsilon) \wedge \frac{\partial}{\partial \varepsilon} w(x, \varepsilon) d\varepsilon + \int_0^\infty w(x, \varepsilon) \wedge \frac{\partial}{\partial \varepsilon} D_x w(x, \varepsilon) d\varepsilon \\ &= w(x, \varepsilon) \wedge D_x w(x, \varepsilon)|_{\varepsilon=0} + \int_0^\infty D_x w(x, \varepsilon) \wedge \frac{\partial}{\partial \varepsilon} w(x, \varepsilon) d\varepsilon + w(x, \varepsilon) \wedge D_x w(x, \varepsilon) \Big|_{\varepsilon=0}^{\varepsilon=\infty} \\ &\quad - \int_0^\infty \frac{\partial}{\partial \varepsilon} w(x, \varepsilon) \wedge D_x w(x, \varepsilon) d\varepsilon = -2 \int_0^\infty \frac{\partial}{\partial \varepsilon} w(x, \varepsilon) \wedge D_x w(x, \varepsilon) d\varepsilon. \end{aligned}$$

Adapting an idea from [4], this identity implies the estimate $\|D\varphi_2\|_{L^{sp}} \leq C|e^{i\varphi}|_{W^{s,p}}^{1/s}$. \square

The interested reader will find the detailed proof in [9].

1. One of the key estimates used in the proof is $\int_{\mathbb{R}^N} \int_0^\infty \varepsilon^{p-sp-1} |D[h * \rho_\varepsilon](x)|^p d\varepsilon dx \leq C|h|_{W^{s,p}}^p$, well-known to experts [11], II.12.

We end this section by comparing our decomposition to the Bourgain-Brezis one. Assuming, for the sake of the simplicity, that φ and g are defined on the whole \mathbb{R}^N , the decomposition in [2] is (with $g = \sum g_j$ the Littlewood-Paley decomposition of g)

$$\varphi_1 := \sum_{j \leq k} g_j \wedge g_k, \quad \varphi_2 := \varphi - \varphi_1. \quad (9)$$

In the same way (6) is related to (7), one may interpret the formula defining φ_1 in (9) as a discrete analog of

$$x \mapsto - \int_0^\infty g * \rho_\varepsilon(x) \wedge \frac{\partial}{\partial \varepsilon} [g * \rho_\varepsilon(x)] d\varepsilon.$$

Thus our decomposition is a continuous version of the one in [2], with the additional sophistication that the regularizations of g are "almost projected" onto \mathbb{S}^1 (via Π).

4 Some applications

As a first application, we may achieve the description of $X^{s,p} = \overline{C^\infty(C; \mathbb{S}^1)}^{W^{s,p}}$, partly obtained in [3] and [5].

Theorem 2 *Let $0 < s < \infty$, $1 \leq p < \infty$. Then*

- a) ([3]) *When $sp < 1$ or $sp \geq N$, $X^{s,p} = W^{s,p}(C; \mathbb{S}^1)$;*
- b) ([5]) *When $s \geq 1$ and $sp \geq 2$, $X^{s,p} = W^{s,p}(C; \mathbb{S}^1)$;*
- c) ([5]) *When $s \geq 1$ and $1 \leq sp < 2$, $X^{s,p} = \{e^{i\varphi} ; \varphi \in W^{s,p} \cap W^{1,sp}(C, \mathbb{R})\}$;*
- d) *When $0 < s < 1$ and $1 < sp < N$, $X^{s,p} = \{e^{i\varphi} ; \varphi \in (W^{s,p} + W^{1,sp})(C, \mathbb{R})\}$.*
- e) *When $0 < s < 1$, $N \geq 2$ and $sp = 1$, $X^{s,p} = \{e^{i\varphi} ; \varphi \in (W^{s,p} + W^{1,1})(C, \mathbb{R})\} \cap W^{s,p}(C; \mathbb{S}^1)$.*

The lack of symmetry between statements d) and e) is explained by the fact that, when $sp > 1$ and $s < 1$, we have $\varphi \in W^{1,sp} \implies e^{i\varphi} \in W^{s,p}$; this implication fails when $sp = 1$ and $s < 1$. For the proof of d) and e), we send the reader to [9].

In a forthcoming joint paper with H. Brezis and H.-M. Nguyen [6], we investigate several consequences of Theorem 1. We end this Note by mentioning one of them.

Theorem 3 ([6]) *Assume that $N \geq 3$, $0 < s < 1$, $2 \leq sp < N$, $k \in \{2, 3, \dots\}$. Then, for each $u \in W^{s,p}(C; \mathbb{S}^1)$, there is some $v \in W^{s,p}(C; \mathbb{S}^1)$ such that $u = v^k$.*

This answers a question of F. Bethuel and D. Chiron [1]. [For the other values of s , p and N , the surjectivity of the map $v \mapsto v^k$ is clarified in [1]. The case considered in Theorem 3 is the only one left open in [1].]

Acknowledgements

The author thanks H.-M. Nguyen for sending him an early version of [10]. He warmly thanks P. Bousquet, H. Brezis and H.-M. Nguyen for useful discussions.

References

- [1] F. Bethuel, D. Chiron, Some questions related to the lifting problem in Sobolev spaces, in Perspectives in Nonlinear Partial Differential Equations, (H. Berestycki, M. Bertsch, F. Browder, L. Nirenberg, eds.), Contemporary Mathematics, Volume 446, 125–152.
- [2] J. Bourgain, H. Brezis, On the equation $\operatorname{div} Y = f$ and application to control of phases, J. Amer. Math. Soc. 16 (2003), 393–426.
- [3] J. Bourgain, H. Brezis, P. Mironescu, Lifting in Sobolev spaces, J. Anal. Math. 80 (2000), 37–86.
- [4] J. Bourgain, H. Brezis, P. Mironescu, Lifting, degree, and distributional Jacobian revisited, Commun. Pure Appl. Math. 58 (2005), 529–551.
- [5] H. Brezis, P. Mironescu, On some questions of topology for S^1 -valued fractional Sobolev spaces, Rev. R. Acad. Cien., Serie A Mat. 95 (2001), 121–143.
- [6] H. Brezis, P. Mironescu, H.-M. Nguyen, in preparation.
- [7] J.-Y. Chemin, Fluides parfaits incompressibles, Astérisque 230 (1995).
- [8] P. Mironescu, Sobolev maps on manifolds: degree, approximation, lifting, in Perspectives in Nonlinear Partial Differential Equations, In honor of Haïm Brezis (H. Berestycki, M. Bertsch, F. Browder, L. Nirenberg, L. A. Peletier, L. Véron eds.), Contemporary Mathematics, Amer. Math. Society 446 (2007), 413–436.
- [9] P. MIRONESCU, Lifting of S^1 -valued maps in sums of Sobolev spaces, to appear.
- [10] H.-M. Nguyen, Inequalities related to lifting and applications, to appear in C. R. Acad. Sci. Paris 346 (2008).
- [11] H. TRIEBEL, Theory of function spaces, Birkhäuser Verlag, 1983.