



HAL
open science

Continuous actions of the group of homeomorphisms of the circle on surfaces

Emmanuel Militon

► **To cite this version:**

Emmanuel Militon. Continuous actions of the group of homeomorphisms of the circle on surfaces. 2012. hal-00748512v1

HAL Id: hal-00748512

<https://hal.science/hal-00748512v1>

Preprint submitted on 5 Nov 2012 (v1), last revised 5 May 2014 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Continuous actions of the group of homeomorphisms of the circle on surfaces

E. Militon *

5 novembre 2012

Abstract

In this article we describe all the continuous group morphisms from the group of orientation preserving homeomorphisms of the circle to the group of homeomorphisms of the annulus or of the torus.

1 Introduction

For a compact manifold M , we denote by $\text{Homeo}(M)$ the group of homeomorphisms of M and by $\text{Homeo}_0(M)$ the connected component of the identity of this group (for the compact-open topology). By a theorem by Fischer (see [6] and [4]), the group $\text{Homeo}_0(M)$ is simple: the study of this group cannot be reduced to the study of other groups. One natural way to have a better understanding of this group is to try to compute its automorphism group. This task was completed by Whittaker who proved the following theorem.

Theorem. (*Whittaker [13]*) *Given two compact manifolds M and N and for any group isomorphism $\varphi : \text{Homeo}_0(M) \rightarrow \text{Homeo}_0(N)$, there exists a homeomorphism $h : M \rightarrow N$ such that the morphism φ is the map $f \mapsto h \circ f \circ h^{-1}$.*

This theorem was generalized by Filipkiewicz (voir [5]) to groups of diffeomorphisms by using a powerful theorem by Montgomery and Zippin which characterizes Lie groups among locally compact groups. The idea of the proof of Whittaker's theorem is the following: we see the manifold M algebraically by considering the subgroup G_x of the group of homeomorphism of M consisting of homeomorphisms which fix the point x in M . One proves then that, for any point x in M , there exists a unique point y_x in N such that $\varphi(G_x)$ is the group of homeomorphisms of N which fix the point y_x . Let us then define h by $h(x) = y_x$. We check then that h is a homeomorphism and that the homeomorphism h satisfies the conclusion of the theorem. However, Whittaker's proof uses crucially the fact that we have a group isomorphism and cannot be generalized easily a priori to the cases of group morphisms. Here is a conjecture that we can make in this case.

Conjecture 1.1. *For a compact manifold M , every group morphism $\text{Homeo}_0(M) \rightarrow \text{Homeo}_0(M)$ is either trivial or induced by a conjugacy by a homeomorphism.*

This conjecture is solved in [10] in the case of the circle. We can also be interested in morphisms from the group $\text{Homeo}_0(M)$ to the group of homeomorphisms of another manifold $\text{Homeo}_0(N)$. For instance, we can make the following conjecture (and generalizations of this one).

Conjecture 1.2. *Denote by $\mathbb{T}^2 - D$ the torus \mathbb{T}^2 with one open disc removed. Denote by $\text{Homeo}_0(\mathbb{T}^2 - D)$ the identity component of the group of homeomorphisms $\mathbb{T}^2 - D$ with support included in the interior of this surface. Every group morphism from $\text{Homeo}_0(\mathbb{T}^2 - D)$ to $\text{Homeo}_0(\mathbb{T}^2)$ is induced by an inclusion of $\mathbb{T}^2 - D$ in \mathbb{T}^2 .*

*supported by the Fondation Mathématique Jacques Hadamard

In this article, we investigate the case of continuous group morphisms from the group $\text{Homeo}_0(\mathbb{S}^1)$ of orientation-preserving homeomorphisms of the circle to the group $\text{Homeo}(S)$ of homeomorphisms of a compact orientable surface S . By simplicity of the group $\text{Homeo}_0(\mathbb{S}^1)$, such a morphism φ is either one-to-one or trivial. Moreover, as the quotient group $\text{Homeo}(S)/\text{Homeo}_0(S)$ is countable, any morphism $\text{Homeo}_0(\mathbb{S}^1) \rightarrow \text{Homeo}(S)/\text{Homeo}_0(S)$ is trivial. Hence, the image of the morphism φ is included in the group $\text{Homeo}_0(S)$. If the surface S is different from the sphere, the torus, the closed disc or the closed annulus, then any compact subgroup of the group $\text{Homeo}_0(S)$ is trivial by a theorem by Kerekjarto (see [3]). Therefore, the group $\text{Homeo}_0(S)$ does not contain a subgroup isomorphic to $\text{SO}(2)$ whereas the group $\text{Homeo}_0(\mathbb{S}^1)$ does and the morphism φ is necessarily trivial. We study in what follows the case where the surface S is the closed annulus or the torus. The case where the surface is the sphere or the closed disc remains open. However, by using the techniques of this article, it suffices to establish the existence of two global fixed point of any continuous action of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the sphere and of one global fixed point of any continuous action of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the disc to obtain theorems similar to those introduced below in the case of the torus and of the annulus.

In the following section, we state our classification theorem of continuous actions of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the torus and on the annulus. Any action will be obtained by gluing actions which preserve a foliation in circle and actions which are transitive on open annuli. In the third section, we describe the continuous actions of the group of compactly-supported homeomorphisms of the real line on the real line or on the circle: this description is useful for the classification theorem and interesting in itself. The fourth and fifth sections are devoted respectively to the proof of the classification theorem in the case of the closed annulus and in the case of the torus.

2 Description of the actions

All the actions of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the closed annulus and on the torus will be obtained by gluing elementary actions of this group on the closed annulus. In this section, we will first describe these elementary actions before describing some model actions to which any continuous action will be conjugate.

The easiest action of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the closed annulus $\mathbb{A} = [0, 1] \times \mathbb{S}^1$ is given by the preservation of a foliation in circle of this annulus:

$$\begin{aligned} p : \text{Homeo}_0(\mathbb{S}^1) &\rightarrow \text{Homeo}(\mathbb{A}) \\ f &\mapsto ((r, \theta) \mapsto (r, f(\theta))) \end{aligned} \cdot$$

The second elementary action we want to describe is a little more complex. Let $\pi : \mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} = \mathbb{S}^1$ be the projection. For a point θ on the circle $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$, we denote by $\tilde{\theta}$ a lift of θ , *i.e.* a point of the real line which projects on θ . For a homeomorphism f of the circle, we denote by \tilde{f} a lift of f , *i.e.* an element of the group $\text{Homeo}_{\mathbb{Z}}(\mathbb{R})$ of homeomorphisms of the real line which commute to integral translations such that $\pi \circ \tilde{f} = f \circ \pi$. A second elementary action is given by:

$$\begin{aligned} a_- : \text{Homeo}_0(\mathbb{S}^1) &\rightarrow \text{Homeo}(\mathbb{A}) \\ f &\mapsto ((r, \theta) \mapsto (\tilde{f}(\tilde{\theta}) - \tilde{f}(\tilde{\theta} - r), f(\theta))) \end{aligned} \cdot$$

Notice that the number $\tilde{f}(\tilde{\theta}) - \tilde{f}(\tilde{\theta} - r)$ does not depend on the lifts $\tilde{\theta}$ and \tilde{f} chosen and belongs to the interval $[0, 1]$. An action analogous to this last one is given by:

$$\begin{aligned} a_+ : \text{Homeo}_0(\mathbb{S}^1) &\rightarrow \text{Homeo}(\mathbb{A}) \\ f &\mapsto ((r, \theta) \mapsto (\tilde{f}(\tilde{\theta} + r) - \tilde{f}(\tilde{\theta}), f(\theta))) \end{aligned} \cdot$$

Notice that the actions a_+ and a_- are conjugate via the homeomorphism of \mathbb{A} given by $(r, \theta) \mapsto (1 - r, \theta)$, which is orientation-reversing, and via the orientation-preserving homeomorphism of \mathbb{A} given by $(r, \theta) \mapsto (r, \theta + r)$. We now describe another way to see the action a_- (and a_+). We see the torus \mathbb{T}^2 as the product $\mathbb{S}^1 \times \mathbb{S}^1$. Let

$$\begin{aligned} a_{\mathbb{T}^2} : \text{Homeo}_0(\mathbb{S}^1) &\rightarrow \text{Homeo}(\mathbb{T}^2) \\ f &\mapsto ((x, y) \mapsto (f(x), f(y))) \end{aligned} \cdot$$

It is easily checked that this defines a morphism. This action leaves the diagonal $\{(x, x), x \in \mathbb{S}^1\}$ invariant. The action obtained by cutting along the diagonal is conjugate to a_- . More precisely, let us define

$$h : \quad \mathbb{A} \rightarrow \mathbb{T}^2 \\ (r, \theta) \mapsto (\theta, \theta - r) .$$

Then, for any element f in the group $\text{Homeo}_0(\mathbb{S}^1)$:

$$h \circ a_-(f) = a_{\mathbb{T}^2}(f) \circ h.$$

Let G_{θ_0} be the group of homeomorphisms of the circle which fix a neighbourhood of the point θ_0 . For a point θ_0 of the circle, the image by the morphism a_- of the group G_{θ_0} leaves the sets $\{(r, \theta_0), r \in [0, 1]\}$ and $\{(r, \theta_0 + r), r \in [0, 1]\}$ globally invariant. On each of the connected components of the complementary of the union of these two sets with the boundary of the annulus, the action of G_{θ_0} is transitive (this is more easily seen by using the action $a_{\mathbb{T}^2}$).

Let us describe now the model actions on the closed annulus which are obtained by gluing the above actions. Take a non-empty compact subset $K \subset [0, 1]$ which contains the points 0 and 1 and a map $\lambda : [0, 1] - K \rightarrow \{-1, +1\}$ which is constant on each connected component of $[0, 1] - K$. Let us define now an action $\varphi_{K, \lambda}$ of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the closed annulus. To a homeomorphism f in $\text{Homeo}_0(\mathbb{S}^1)$, we associate a homeomorphism $\varphi_{K, \lambda}(f)$ which is defined as follows. If $r \in K$, we associate to a point $(r, \theta) \in \mathbb{A}$ the point $(r, f(\theta))$. If r belongs to a connected component (r_1, r_2) of the complementary of the compact set K and if $\lambda((r_1, r_2)) = \{-1\}$, we associate to the point $(r, \theta) \in \mathbb{A}$ the point

$$((r_2 - r_1)(\tilde{f}(\theta) - \tilde{f}(\theta - \frac{r - r_1}{r_2 - r_1})) + r_1, f(\theta)).$$

This action is continuous. This last map is obtained by conjugating the homeomorphism $a_-(f)$ with the map $(r, \theta) \mapsto (\zeta(r), \theta)$ where ζ is the unique linear increasing homeomorphism $[0, 1] \rightarrow [r_1, r_2]$. If r belongs to a connected component (r_1, r_2) of the complementary of the compact set K and if $\lambda((r_1, r_2)) = \{+1\}$, we associate to the point $(r, \theta) \in \mathbb{A}$ the point

$$((r_2 - r_1)(\tilde{f}(\theta + \frac{r - r_1}{r_2 - r_1}) - \tilde{f}(\theta)) + r_1, f(\theta)).$$

This last map is also obtained after renormalizing $a_+(f)$ on the interval (r_1, r_2) . This defines a continuous morphism $\varphi_{K, \lambda} : \text{Homeo}_0(\mathbb{S}^1) \rightarrow \text{Homeo}(\mathbb{A})$. To construct an action on the torus, it suffices to identify the point $(0, \theta)$ of the closed annulus with the point $(1, \theta)$. We denote by $\varphi_{K, \lambda}^{\mathbb{T}^2}$ the continuous action on the torus obtained this way. The main theorem of this article is the following:

Theorem 2.1. *Any non-trivial continuous action of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the closed annulus is conjugate to one of the actions $\varphi_{K, \lambda}$. Any non-trivial continuous action of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the torus is conjugate to one of the actions $\varphi_{K, \lambda}^{\mathbb{T}^2}$.*

In particular, any continuous action of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the torus admits an invariant circle. The following sections of this article are devoted to the proof of this theorem.

Notice that this theorem does not give directly a description of the conjugacy classes of such actions as two actions $\varphi_{K, \lambda}$ and $\varphi_{K', \lambda'}$ may be conjugate even though $K \neq K'$ or $\lambda \neq \lambda'$. Now, let $K \subset [0, 1]$ and $K' \subset [0, 1]$ be two compact sets which contain the finite set $\{0, 1\}$. Let $\lambda : [0, 1] - K \rightarrow \{-1, +1\}$ and $\lambda' : [0, 1] - K' \rightarrow \{-1, +1\}$ be two maps which are constant on each connected component of their domain of definition. The following theorem characterizes when the actions $\varphi_{K, \lambda}$ and $\varphi_{K', \lambda'}$ are conjugate.

Proposition 2.2. *The following statements are equivalent:*

- the actions $\varphi_{K, \lambda}$ and $\varphi_{K', \lambda'}$ are conjugate;
- either there exists an increasing homeomorphism $h : [0, 1] \rightarrow [0, 1]$ such that $h(K) = K'$ and such that $\lambda' \circ h = \lambda$ except on a finite number of connected component of $[0, 1] - K$ or there exists a decreasing homeomorphism $h : [0, 1] \rightarrow [0, 1]$ such that $h(K) = K'$ and such that $\lambda' \circ h = -\lambda$ except on a finite number of connected component of $[0, 1] - K$.

Proof. Let us begin by proving that the second statement implies the first one.

If there exists an increasing homeomorphism h which sends the compact subset $K \subset [0, 1]$ to the compact subset $K' \subset [0, 1]$, then the actions $\varphi_{K,\lambda}$ and $\varphi_{K',\lambda \circ h^{-1}}$ are conjugate. Indeed, denote by \hat{h} the homeomorphism $[0, 1] \rightarrow [0, 1]$ which coincides with h on K and which, on each connected component (r_1, r_2) of the complementary of K , is the unique increasing linear homeomorphism $(r_1, r_2) \rightarrow h((r_1, r_2))$. Then the actions $\varphi_{K,\lambda}$ and $\varphi_{K',\lambda \circ h}$ are conjugate via the homeomorphism

$$\begin{aligned} \mathbb{A} &\rightarrow \mathbb{A} \\ (r, \theta) &\mapsto (\hat{h}(r), \theta) \end{aligned} \cdot$$

Similarly, suppose that h is an orientation-reversing homeomorphism of the compact interval $[0, 1]$ which sends the compact subset $K \subset [0, 1]$ to the compact subset $K' \subset [0, 1]$. As above, we denote by \hat{h} the homeomorphism obtained from h by linearizing the homeomorphism on each connected component of the complementary of K . Then the actions $\varphi_{K,\lambda}$ and $\varphi_{K',-\lambda \circ h^{-1}}$ are conjugate via the map

$$\begin{aligned} \mathbb{A} &\rightarrow \mathbb{A} \\ (r, \theta) &\mapsto (\hat{h}(r), \theta) \end{aligned} \cdot$$

It suffices then to use the following lemma to complete the proof of the converse in the proposition.

Lemma 2.3. *Let $K \subset [0, 1]$ be a compact subset which contains the points 0 and 1. If $\lambda : [0, 1] - K \rightarrow \{-1, +1\}$ and $\lambda' : [0, 1] - K \rightarrow \{-1, +1\}$ are continuous maps which are equal except on one connected component (r_1, r_2) of the complementary of K where they differ, then the actions $\varphi_{K,\lambda}$ and $\varphi_{K,\lambda'}$ are conjugate.*

Proof. This lemma comes from the fact that the actions a_+ and a_- are conjugate via the homeomorphism of the annulus $(r, \theta) \mapsto (r, \theta + r)$. More precisely, suppose that $\lambda((r_1, r_2)) = \{+1\}$. Then the actions $\varphi_{K,\lambda}$ and $\varphi_{K,\lambda'}$ are conjugate via the homeomorphism h defined as follows. The homeomorphism h is equal to the identity on the Cartesian product of the complementary of (r_1, r_2) with the circle and is equal to $(r, \theta) \mapsto (r, \theta + \frac{r-r_1}{r_2-r_1})$ on the open set $(r_1, r_2) \times \mathbb{S}^1$. \square

Let us establish now the other implication. Suppose that there exists a homeomorphism g which conjugates the actions $\varphi_{K,\lambda}$ and $\varphi_{K',\lambda'}$. Now, for any angle α , if we denote by R_α the rotation of angle α :

$$g \circ \varphi_{K,\lambda}(R_\alpha) = \varphi_{K',\lambda'}(R_\alpha) \circ g.$$

Indeed, the homeomorphism $\varphi_{K,\lambda}(R_\alpha)$ is the rotation of angle α of the annulus. This means that, for any point (r, θ) of the closed annulus and any angle $\alpha \in \mathbb{S}^1$:

$$g((r, \theta + \alpha)) = g((r, \theta)) + (0, \alpha).$$

In particular, the homeomorphism g permutes the leaves of the foliation of the annulus by circle whose leaves are of the form $\{r\} \times \mathbb{S}^1$. Fix now a point θ_0 on the circle. The homeomorphism g sends the set $K \times \{\theta_0\}$ of fixed points of $\varphi_{K,\lambda}(G_{\theta_0})$ (the points which are fixed by every element of this group) to the set $K' \times \{\theta_0\}$ of fixed points of $\varphi_{K',\lambda'}(G_{\theta_0})$. From this and what is above, we deduce that for any $r \in K$ and any angle θ , $g(r, \theta) = (h(r), \theta)$, where $h : K \rightarrow K'$ is a homeomorphism. Moreover, if the homeomorphism g is orientation-preserving, then the homeomorphism h is increasing and, if the homeomorphism g is orientation-reversing, then the homeomorphism h is decreasing. We can extend the homeomorphism h to a homeomorphism of $[0, 1]$ which sends K to K' . Notice that, for a connected component (r_1, r_2) of the complementary of K in $[0, 1]$, the homeomorphism g sends the open set $(r_1, r_2) \times \mathbb{S}^1$ onto the open set $(r'_1, r'_2) \times \mathbb{S}^1$, where $(r'_1, r'_2) = h((r_1, r_2))$ is a connected component of the complementary of K' in $[0, 1]$.

It suffices now to establish that the condition on the maps λ and λ' is satisfied for the homeomorphism h . Suppose, to simplify the proof, that the homeomorphism g is orientation-preserving and hence the homeomorphism h is increasing: the case where the homeomorphism g is orientation reversing can be treated similarly. Suppose by contradiction that there exists a sequence $((r_{1,n}, r_{2,n}))_{n \in \mathbb{N}}$ of connected components of the open set $[0, 1] - K$ such that:

- $\lambda((r_{1,n}, r_{2,n})) = +1$;
- $\lambda'((r'_{1,n}, r'_{2,n})) = -1$, where $(r'_{1,n}, r'_{2,n}) = h((r_{1,n}, r_{2,n}))$;
- the sequence $(r_{1,n})_{n \in \mathbb{N}}$ is monotonous and converges to a real number r_∞ .

We will prove then that either the curve $\{(r, \theta_0), r \geq r_\infty\}$ (if the sequence $(r_{1,n})_{n \in \mathbb{N}}$ is decreasing) or the curve $\{(r, \theta_0), r \leq r_\infty\}$ (if the sequence $(r_{1,n})_{n \in \mathbb{N}}$ is increasing) is sent by the homeomorphism g to a curve which accumulates on $\{r_\infty\} \times \mathbb{S}^1$, which is not possible. The hypothesis $\lambda((r_{1,n}, r_{2,n})) = -1$ and $\lambda'((r'_{1,n}, r'_{2,n})) = +1$ for any n would lead to the same contradiction.

To achieve this, it suffices to prove that, for any positive integer n , the homeomorphism g sends the curve $\{(r, \theta_0), r_{1,n} < r < r_{2,n}\}$ onto the curve $\{(r, \theta_0 + \frac{r-r'_{1,n}}{r'_{2,n}-r'_{1,n}}), r'_{1,n} < r < r'_{2,n}\}$. Fix a positive integer n . Observe that the restriction of the group of homeomorphisms $\varphi_{K,\lambda}(G_{\theta_0})$ to $(r_{1,n}, r_{2,n}) \times \mathbb{S}^1$ has two invariant simple curves $\{(r, \theta_0), r_{1,n} < r < r_{2,n}\}$ and $\{(r, \theta_0 - \frac{r-r_{1,n}}{r_{2,n}-r_{1,n}}), r_{1,n} < r < r_{2,n}\}$ on which the action is transitive. The action of this group is transitive on the two connected components of the complementary of these two curves, which are open sets. Likewise, the restriction of the group of homeomorphisms $\varphi_{K',\lambda'}(G_{\theta_0})$ to $(r'_{1,n}, r'_{2,n})$ has two invariant simple curves $\{(r, \theta_0), r'_{1,n} < r < r'_{2,n}\}$ and $\{(r, \theta_0 + \frac{r-r'_{1,n}}{r'_{2,n}-r'_{1,n}}), r'_{1,n} < r < r'_{2,n}\}$ on which the action is transitive and the action of this group is transitive on the two connected components of the complementary of these two curves, which are open sets. Therefore, the homeomorphism g sends the curve $\{(r, \theta_0), r_{1,n} < r < r_{2,n}\}$ to one of the curves $\{(r, \theta_0), r'_{1,n} < r < r'_{2,n}\}$ or $\{(r, \theta_0 + \frac{r-r'_{1,n}}{r'_{2,n}-r'_{1,n}}), r'_{1,n} < r < r'_{2,n}\}$. Let us find now which of them is the image of the curve $\{(r, \theta_0), r_{1,n} < r < r_{2,n}\}$. We fix the orientation of the circle induced by the orientation of the real line \mathbb{R} and the covering map $\mathbb{R} \rightarrow \mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$. This orientation gives rise to an order on $\mathbb{S}^1 - \{\theta_0\}$. Take a homeomorphism f in G_{θ_0} different from the identity such that, for any $x \neq \theta_0$, $f(x) \geq x$. Then, for any $r \in (r_{1,n}, r_{2,n})$, $p_1 \circ \varphi_{K,\lambda}(f)(r, \theta_0) \geq r$, where $p_1 : \mathbb{A} = [0, 1] \times \mathbb{S}^1 \rightarrow [0, 1]$ is the projection, and the restriction of the homeomorphism $\varphi_{K,\lambda}(f)$ to the curve $\{(r, \theta_0), r_{1,n} < r < r_{2,n}\}$ is different from the identity. Likewise, for any $r \in (r'_{1,n}, r'_{2,n})$, $p_1 \circ \varphi_{K',\lambda'}(f)(r, \theta_0) \leq r$ and $p_1 \circ \varphi_{K',\lambda'}(f)(r, \theta_0 + r) \geq r$ and the restrictions of the homeomorphism $\varphi_{K',\lambda'}(f)$ to $\{(r, \theta_0), r'_{1,n} < r < r'_{2,n}\}$ and to $\{(r, \theta_0 + \frac{r-r'_{1,n}}{r'_{2,n}-r'_{1,n}}), r'_{1,n} < r < r'_{2,n}\}$ are different from the identity. Moreover, the map

$$\begin{aligned} (r_{1,n}, r_{2,n}) &\rightarrow (r'_{1,n}, r'_{2,n}) \\ r &\mapsto p_1 \circ g(r, \theta_0) \end{aligned}$$

is strictly increasing as the homeomorphism g was supposed to be orientation-preserving. This implies what we wanted to prove. \square

3 Continuous actions of $\text{Homeo}_c(\mathbb{R})$ on the line

Let us denote by $\text{Homeo}_c(\mathbb{R})$ the group of compactly supported homeomorphisms of the real line \mathbb{R} . The aim of this section is to give a proof of the following theorem.

Theorem 3.1. *Let $\psi : \text{Homeo}_c(\mathbb{R}) \rightarrow \text{Homeo}(\mathbb{R})$ be a continuous group morphism whose image has no fixed point. Then there exists a homeomorphism h of \mathbb{R} such that, for any compactly supported homeomorphism f of the real line:*

$$\psi(f) = h \circ f \circ h^{-1}.$$

Remarks: If ψ is chosen to be with values in $\text{Homeo}_c(\mathbb{R})$, then the conclusion of this theorem holds without continuity hypothesis.

This theorem also holds in the setting of diffeomorphisms, *i.e.* any continuous action of the group of compactly-supported C^r diffeomorphisms on the real line is topologically conjugate to the inclusion. Here also, one can avoid the continuity hypothesis by considering a compactly supported action. The proof in those cases is the same.

This theorem allows us to describe any continuous action of the group $\text{Homeo}_c(\mathbb{R})$ on the real line as it suffices to consider the action on each connected component of the complementary of the fixed point set of the action.

Proof. We fix a morphism ψ as in the statement of the theorem. During this proof, for a subset $A \subset \mathbb{R}$, we will denote by G_A the group of compactly supported homeomorphisms which pointwise fix a neighbourhood of A and by $F_A \subset \mathbb{R}$ the closed set of fixed points of $\psi(G_A)$ (the set of points which are fixed by every element of this group). Let us begin by giving a sketch of the proof of this theorem. We will first prove that, for a compact interval I with non-empty interior, the set F_I is compact and non-empty. As a consequence, for any real number x , the closed set F_x is non-empty. We will then prove that the sets F_x are single-point sets. Let $\{h(x)\} = F_x$. Then the map h is a homeomorphism and satisfies the relation required by the theorem. Let us detail this now.

Notice that, as the group $\text{Homeo}_c(\mathbb{R})$ is simple (see [6]), the image of ψ is included in the group of increasing homeomorphisms of \mathbb{R} (otherwise there would exist a group morphism $\text{Homeo}_c(\mathbb{R}) \rightarrow \mathbb{Z}/2\mathbb{Z}$) and the morphism ψ is a one-to-one map.

The following lemma is the only lemma in this proof where we need to use the continuity of the action. The lemmas which follow are proved without the use of continuity, even though the proof was sometimes simpler with the continuity hypothesis. The author was not able to prove this lemma without the continuity hypothesis but found that it sufficed to prove that there does not exist a non-trivial morphism from the group $\text{Homeo}_c(\mathbb{R})$ to the group of permutations of a countable set to prove this lemma without continuity assumption.

Lemma 3.2. *For a compact interval $I \subset \mathbb{R}$ with non-empty interior, the closed set F_I is non-empty.*

Proof. Fix a compact interval I with non-empty interior. Take a non-zero vector field $X : \mathbb{R} \rightarrow \mathbb{R}$ supported in I . The flow of this vector field defines a morphism

$$\begin{array}{ccc} \mathbb{R} & \rightarrow & \text{Homeo}_c(\mathbb{R}) \\ t & \mapsto & \varphi^t \end{array} .$$

Let us admit for the moment that the set F of fixed points of the subgroup $\{\psi(\varphi^t), t \in \mathbb{R}\}$ of $\text{Homeo}(\mathbb{R})$ is non-empty. Notice that this set is not \mathbb{R} as the morphism ψ is one-to-one. As each homeomorphism φ^t commutes with any element in G_I , we obtain that, for any element g in $\psi(G_I)$, $g(F) = F$. Moreover, as any element of the group G_I can be joined to the identity by a continuous path which is included in G_I and as the morphism ψ is continuous, any connected component C of F is invariant under $\psi(G_I)$. The upper bounds and the lower bounds of these components which lie in \mathbb{R} are then fixed points of the group $\psi(G_I)$. This proves the lemma.

It remains to prove that the set F is non-empty. Suppose by contradiction that it is empty. Then, for any real number x , the map

$$\begin{array}{ccc} \mathbb{R} & \rightarrow & \mathbb{R} \\ t & \mapsto & \psi(\varphi^t)(x) \end{array}$$

is a homeomorphism. Indeed, if it was not onto, the supremum or the infimum of the image would provide a fixed point for $(\psi(\varphi^t))_{t \in \mathbb{R}}$. If it was not one-to-one, there would exist $t_0 \neq 0$ such that $\psi(\varphi^{t_0})(x) = x$. Then, for any positive integer n , $\psi(\varphi^{t_0/2^n})(x) = x$ and, by continuity of ψ , for any real number t , $\psi(\varphi^t)(x) = x$ and the point x would be fixed by $(\psi(\varphi^t))_{t \in \mathbb{R}}$.

Fix a real number x_0 . Let $T_{x_0} : G_I \rightarrow \mathbb{R}$ be the map defined by

$$\psi(f)(x_0) = \psi(\varphi^{T_{x_0}(f)})(x_0).$$

The map T_{x_0} is a group morphism as, for any homeomorphisms f and g in G_I ,

$$\begin{aligned} \psi(\varphi^{T_{x_0}(fg)})(x_0) &= \psi(fg)(x_0) \\ &= \psi(f)\psi(\varphi^{T_{x_0}(g)})(x_0) \\ &= \psi(\varphi^{T_{x_0}(g)})\psi(f)(x_0) \\ &= \psi(\varphi^{T_{x_0}(g)+T_{x_0}(f)})(x_0). \end{aligned}$$

However, the group G_I , which is isomorphic to the group $\text{Homeo}_c(\mathbb{R}) \times \text{Homeo}_c(\mathbb{R})$, is a perfect group. Therefore, the morphism T_{x_0} is trivial. As the point x_0 is any point in \mathbb{R} , we deduce that the restriction of ψ to G_I is trivial, which is not possible as the morphism ψ is one-to-one. \square

Remark: if $\psi(\text{Homeo}_c(\mathbb{R})) \subset \text{Homeo}_c(\mathbb{R})$, this lemma can be proved without continuity hypothesis. Indeed, consider an element f in $\text{Homeo}_c(\mathbb{R})$ supported in I . One of the connected components of the set of fixed points of $\psi(f)$ is of the form $(-\infty, a]$ for some a in \mathbb{R} . This interval is necessarily invariant by the group $\psi(G_I)$ which commutes with f . Hence, the point a is a fixed point for the group $\psi(G_I)$.

During the proof, we will often use the following elementary result:

Lemma 3.3. *Let I and J be disjoint compact non-empty intervals. For any homeomorphism g in $\text{Homeo}_c(\mathbb{R})$, there exist homeomorphisms $g_1 \in G_I$, $g_2 \in G_J$ and $g_3 \in G_I$ such that:*

$$g = g_1 g_2 g_3.$$

Proof. Let g be a homeomorphism in $\text{Homeo}_c(\mathbb{R})$. Let h_1 be a homeomorphism in G_I which sends the interval $g(I)$ to an interval which is in the same connected component of $\mathbb{R} - J$ as the interval I . Let h_2 be a homeomorphism in G_J which is equal to $g^{-1} \circ h_1^{-1}$ on a neighbourhood of the interval $h_1 \circ g(I)$. Then, the homeomorphism $h_2 \circ h_1 \circ g$ belongs to G_I . It suffices then to take $g_1 = h_1^{-1}$, $g_2 = h_2^{-1}$ and $g_3 = h_2 \circ h_1 \circ g$ to conclude the proof of the lemma. \square

Before stating the next lemma, observe that, for compact intervals I with non-empty interiors, the sets F_I are pairwise homeomorphic by an increasing homeomorphism. Indeed, let I and J be two such intervals. Then there exists a homeomorphism λ in $\text{Homeo}_c(\mathbb{R})$ such that $\lambda(I) = J$. Then $\lambda G_I \lambda^{-1} = G_J$. Taking the image by ψ , we obtain that $\psi(\lambda)\psi(G_I)\psi(\lambda)^{-1} = \psi(G_J)$ and therefore $\psi(\lambda)(F_I) = F_J$.

Lemma 3.4. *For a compact interval $I \subset \mathbb{R}$ with non-empty interior, the closed set F_I is compact.*

Proof. Fix a compact interval I with non-empty interior. Suppose by contradiction that there exists a sequence $(a_k)_{k \in \mathbb{N}}$ of real numbers in F_I which tends to $+\infty$ (if we suppose that it tends to $-\infty$, we obtain of course an analogous contradiction). Let us choose a compact interval J which is disjoint from I . By the remark before the lemma, there exists a sequence $(b_k)_{k \in \mathbb{N}}$ of elements in F_J which tends to $+\infty$. Take positive integers n_1 , n_2 and n_3 such that $a_{n_1} < b_{n_2} < a_{n_3}$. Fix $x_0 < a_{n_1}$. We notice then that for any homeomorphisms $g_1 \in G_I$, $g_2 \in G_J$ and $g_3 \in G_I$, the following inequality is satisfied:

$$\psi(g_1)\psi(g_2)\psi(g_3)(x_0) < a_{n_3}.$$

Lemma 3.3 implies then that

$$\overline{\{\psi(g)(x_0), g \in \text{Homeo}_c(\mathbb{R})\}} \subset (-\infty, a_{n_3}).$$

The greatest element of the left-hand set is a fixed point of the image of ψ : this is not possible as this image was supposed to have no fixed point. \square

Lemma 3.5. *The closed sets F_x , where x is a point of the real line, are non-empty, compact, pairwise disjoint and have an empty interior.*

Proof. Notice that, if an interval J is included in an interval I , then $G_I \subset G_J$ and the same inclusion is true if we take the images of these subsets of $\text{Homeo}_c(\mathbb{R})$ by the morphism ψ . That is why $F_J \subset F_I$. Now, for any finite family of intervals $(I_n)_n$ whose intersection has non-empty interior, the intersection of the closed sets F_{I_n} which contains the non-empty closed set $F_{\bigcap_n I_n}$, is not empty. Fix now a point x in \mathbb{R} . Notice that

$$F_x = \bigcap_I F_I,$$

where the intersection is taken over the compact intervals I whose interior contains the point x . By compactness, this set is not empty and it is compact.

Take two points $x \neq y$ in \mathbb{R} et let us prove now that the sets F_x and F_y are disjoint. If they were not disjoint, the group generated by $\psi(G_x)$ and $\psi(G_y)$ would have a fixed point p . However, by lemma 3.3, the groups G_x and G_y generate the group $\text{Homeo}_c(\mathbb{R})$. Then, the point p would be a fixed point of the group $\psi(\text{Homeo}_c(\mathbb{R}))$, which is in contradiction with the hypothesis we made on the morphism ψ .

Let us prove now that the sets F_x have an empty interior. Of course, given two points x and y of the real line, if h is a homeomorphism in $\text{Homeo}_c(\mathbb{R})$ which sends the point x to the point y , then $\psi(h)(F_x) = F_y$. Therefore, the sets F_x are pairwise homeomorphic. If the sets F_x had a non-empty interior, there would exist uncountably many open intervals of the real line which are pairwise disjoint, which is false. \square

Lemma 3.6. *For every point x of the real line, every connected component of the complementary of F_x meets either each of the F_y , for $y > x$, or each of the F_y , for $y < x$.*

Proof. Let us fix a point x_0 in \mathbb{R} . Let (a_1, a_2) be a connected component of the complementary of F_{x_0} . It is possible that either $a_1 = -\infty$ or $a_2 = +\infty$.

Let us prove by contradiction that there exists $y_0 \neq x_0$ such that $F_{y_0} \cap (a_1, a_2) \neq \emptyset$. Suppose that, for any point $y \neq x_0$, $F_y \cap (a_1, a_2) = \emptyset$. For any couple of real numbers (z_1, z_2) , choose a homeomorphism h_{z_1, z_2} in $\text{Homeo}_c(\mathbb{R})$ such that $h_{z_1, z_2}(z_1) = z_2$. We claim that the open sets $\psi(h_{x_0, y})(a_1, a_2)$, for $y \in \mathbb{R}$, are pairwise disjoint, which is not possible as there would be uncountably many pairwise disjoint open intervals in \mathbb{R} . Indeed, if this was not the case, suppose that $\psi(h_{x_0, y_1})(a_1, a_2) \cap \psi(h_{x_0, y_2})(a_1, a_2) \neq \emptyset$ for $y_1 \neq y_2$. As the union of the closed sets F_y is invariant under the action ψ , then, for $i = 1, 2$, when a_i is finite, $\psi(h_{x_0, y_1})^{-1} \circ \psi(h_{x_0, y_2})(a_i) \notin (a_1, a_2)$, and $\psi(h_{x_0, y_2})^{-1} \circ \psi(h_{x_0, y_1})(a_i) \notin (a_1, a_2)$ so that $\psi(h_{x_0, y_1})(a_i) = \psi(h_{x_0, y_2})(a_i)$. But this last equality cannot hold as the point on the left-hand side belongs to F_{y_1} and the point on the right-hand side belongs to F_{y_2} and we observed that these two closed sets were disjoint.

In the case $y_0 > x_0$, let us prove now that, for any $y > x_0$, the closed set F_y meets the open interval (a_1, a_2) (it could be proved similarly that, if $y_0 < x_0$, then for any $y < x_0$, the closed set F_y meets the open interval (a_1, a_2)). Take $y > x_0$. There exists a homeomorphism h in G_{x_0} which sends y_0 to y . Then, as we saw before, the image of F_{y_0} under $\psi(h)$ is F_y . Moreover the homeomorphism $\psi(h)$ fixes the points x_1 and x_2 so that the closed set F_y meets the open interval (a_1, a_2) . \square

Lemma 3.7. *Each set F_x contains only one point.*

Proof. Suppose that there exists a real number x such that the set F_x contains two points $p_1 < p_2$. By lemma 3.6, there exists a real number y different from x such that the set F_y has a common point with the open interval (p_1, p_2) . Take a point $r < p_1$. Then, for any homeomorphisms g_1 in G_x , g_2 in G_y and g_3 in G_x ,

$$\psi(g_1) \circ \psi(g_2) \circ \psi(g_3)(r) < p_2.$$

By lemma 3.3, this implies that the following inclusion holds:

$$\{\psi(g)(r), g \in \text{Homeo}_c(\mathbb{R})\} \subset (-\infty, p_2].$$

The supremum of the left-hand set provides a fixed point for the action ψ : this is a contradiction. \square

Lemma 3.8. *Let us denote by $h(x)$ the only point in the set F_x . Then the map h is either strictly increasing or strictly decreasing.*

Proof. Notice that, as the sets F_x are pairwise disjoint, the map h is one-to-one. Take two real numbers $x_0 < y_0$. Then either $h(x_0) < h(y_0)$ or $h(y_0) < h(x_0)$. Take any two real numbers $x < y$. Then there exists a homeomorphism f in $\text{Homeo}_c(\mathbb{R})$ which sends the point x_0 to the point x and the point y_0 to the point y . Then the homeomorphism $\psi(f)$ sends the only point $h(x_0)$ in F_{x_0} to the only point $h(x)$ of F_x and sends the point $h(y_0)$ to the point $h(y)$. As the map $\psi(f)$ is strictly increasing, we obtain that the couple of points $(h(x), h(y))$ is in the same order as the couple of points $(h(x_0), h(y_0))$. Hence, the map h is either strictly increasing or strictly decreasing. \square

Lemma 3.9. *The sets F_x cover the real line. In other words, the map h is onto.*

Proof. Suppose by contradiction that the sets F_x do not cover the real line. Let C be a connected component of the complementary of the union of the F_x , where the point x varies over the real line. To

simplify the exposition of the proof, we suppose that the map h is increasing. Let us denote by x_0 the supremum of the set of points x such that $h(x)$ is lower than any point in the interval C . Then the point $h(x_0)$ is necessarily in the closure of C : otherwise, there would exist an interval in the complementary of the union of the F_x which strictly contains the interval C . We may suppose for instance that the point $h(x_0)$ is the supremum of the interval C . As in the proof of lemma 3.6, choose, for each couple (z_1, z_2) of real numbers, a homeomorphism h_{z_1, z_2} in $\text{Homeo}_c(\mathbb{R})$ which sends the point z_1 to the point z_2 . Then the sets $\psi(h_{x_0, x})(C)$, for x in \mathbb{R} , are pairwise disjoint: they are pairwise distinct as their suprema are pairwise distinct (the supremum of the set $\psi(h_{x_0, x})(C)$ is the point $h(x)$) and the fact that C does not contain any point of one of the F_x allows us to conclude (notice that the infima of those sets are accumulated by points in the union of the F_x , by definition of C). Then the set C has necessarily an empty interior as the topological space \mathbb{R} is second-countable. We conclude that $C = \{h(x_0)\} = F_{x_0}$ which is not possible. \square

By lemmas 3.8 and 3.9, the map h is a homeomorphism. Now take a homeomorphism f in $\text{Homeo}_c(\mathbb{R})$ and a point x in \mathbb{R} . Then the homeomorphism $\psi(f)$ sends the only point $h(x)$ in F_x to the only point $h(f(x))$ in $F_{f(x)}$. This implies that $\psi(f) \circ h = h \circ f$, which proves theorem 3.1. \square

Proposition 3.10. *Any action of the group $\text{Homeo}_c(\mathbb{R})$ on the circle has a fixed point.*

Hence, the description of the actions of the group $\text{Homeo}_c(\mathbb{R})$ on the circle is given by an action of the group $\text{Homeo}_c(\mathbb{R})$ on the real line which is homeomorphic to the circle minus one point.

Proof. As $H_b^2(\text{Homeo}_c(\mathbb{R}), \mathbb{Z}) = \{0\}$ (see [9]), the Euler class of this action necessarily vanishes so that this action admits a fixed point (see [7] about the Euler class of an action on the circle and its properties). \square

Finally, let us recall a result which is proved in [10].

Proposition 3.11. *Any non-trivial action of the group $\text{Homeo}_0(\mathbb{S}^1)$ on the circle is a conjugacy by a homeomorphism of the circle.*

4 Actions on the annulus

This section is dedicated to the proof of theorem 2.1 in the case of morphisms with values in the group of homeomorphisms of the annulus. Fix a continuous morphism $\varphi : \text{Homeo}_0(\mathbb{S}^1) \rightarrow \text{Homeo}(\mathbb{A})$.

First, as the group $\text{Homeo}_0(\mathbb{S}^1)$ is simple (see [6]), such a group morphism is either trivial or one-to-one. We assume in the rest of this section that this group morphism is one-to-one. The induced group morphism $\text{Homeo}_0(\mathbb{S}^1) \rightarrow \text{Homeo}(\mathbb{A})/\text{Homeo}_0(\mathbb{A})$ is not one-to-one as the source is uncountable and the target is countable. Therefore, it is trivial and the image of the morphism φ is included in the group $\text{Homeo}_0(\mathbb{A})$.

Now, the subgroup of rotations of the circle is continuously isomorphic to the topological group \mathbb{S}^1 . The image of this subgroup under the morphism φ is a compact subgroup of the group of homeomorphisms of the closed annulus which is continuously isomorphic to \mathbb{S}^1 . It is known that such a subgroup is conjugate to the rotation subgroup $\{(r, \theta) \mapsto (r, \theta + \alpha), \alpha \in \mathbb{S}^1\}$ of the group of homeomorphisms of the annulus (see [3]). After possibly conjugating φ , we may suppose from now on that, for any angle α , the morphism φ sends the rotation of angle α of the circle to the rotation of angle α of the annulus.

4.1 An invariant foliation

Fix a point θ_0 on the circle. Recall that the group G_{θ_0} is the group of homeomorphisms of the circle which fix a neighbourhood of the point θ_0 . Let us denote by $F_{\theta_0} \subset \mathbb{A}$ the closed subset of fixed points of $\varphi(G_{\theta_0})$ (*i.e.* points which are fixed under every homeomorphism in this group). The action φ induces

actions on the two boundary components of the annulus which are circles. The action of $\varphi(G_{\theta_0})$ on each of these boundary components admits a fixed point by proposition 3.11. Therefore, the set F_{θ_0} is non-empty. For any angle θ , let $\alpha = \theta - \theta_0$. Then $G_\theta = R_\alpha G_{\theta_0} R_\alpha^{-1}$, where R_α denotes the rotation of the circle of angle α . Therefore, $\varphi(G_\theta) = R_\alpha \varphi(G_{\theta_0}) R_\alpha^{-1}$, where R_α denotes here by abuse the rotation of angle α of the annulus, and $F_\theta = R_\alpha(F_{\theta_0})$. Let

$$B = \bigcup_{\theta \in \mathbb{S}^1} F_\theta = \bigcup_{\alpha \in \mathbb{S}^1} R_\alpha(F_{\theta_0}).$$

This set is of the form $B = K \times \mathbb{S}^1$, where K is the image of F_{θ_0} under the projection $\mathbb{A} = [0, 1] \times \mathbb{S}^1 \rightarrow [0, 1]$. The set K is compact and contains the points 0 and 1. Let \mathcal{F} be the foliation of the set B whose leaves are the circles of the form $\{r\} \times \mathbb{S}^1$, where the real number r belongs to the compact set K .

Lemma 4.1. *Each leaf of the foliation \mathcal{F} is preserved by the action φ .*

Proof. Fix an angle θ . Let us prove that, for any point x in F_θ the orbit of the point x under the action φ is contained in $\{R_\alpha(x), \alpha \in \mathbb{S}^1\} \subset \mathbb{A}$. First, for any orientation-preserving homeomorphism f of the circle which fixes the point θ , the homeomorphism $\varphi(f)$ pointwise fixes the set F_θ . Indeed, such a homeomorphism of the circle is the uniform limit of homeomorphisms which pointwise fix a neighbourhood of θ and the claim results from the continuity of the action φ . Any orientation-preserving homeomorphism of the circle g can be written $g = R_\beta f$, where f is a homeomorphism which fixes the point θ . Now, any point x in F_θ is fixed under $\varphi(f)$ and sent to a point in $\{R_\alpha(x), \alpha \in \mathbb{S}^1\} \subset \mathbb{A}$ under the rotation $R_\beta = \varphi(R_\beta)$, which proves the lemma. \square

Lemma 4.2. *Each closed set F_θ intersects each leaf of the foliation \mathcal{F} in exactly one point. Moreover, the map which, to any point (r, θ) of $K \times \mathbb{S}^1 \subset \mathbb{A}$, associates the only point of F_θ on the leaf $\{r\} \times \mathbb{S}^1$ is a homeomorphism of $K \times \mathbb{S}^1$. This homeomorphism conjugates the restriction of the action $\varphi_{K, \lambda}$ to $K \times \mathbb{S}^1$ to the restriction of the action φ to $K \times \mathbb{S}^1$ for any continuous map $\lambda : \mathbb{S}^1 - K \rightarrow \{-1, +1\}$. Moreover, this homeomorphism is of the form $(r, \theta) \mapsto (r, \eta(r) + \theta)$ where $\eta : K \rightarrow \mathbb{S}^1$ is a continuous function.*

Proof. By the lemma above, the action φ preserves each set of the form $\{r\} \times \mathbb{S}^1$ and induces on each of these set an action on a circle. By proposition 3.11, such actions are well-understood and the restriction of this action to a subgroup of the form G_θ admits exactly one fixed point (this action is non-trivial as the rotation subgroup acts non-trivially). This implies the first statement of the lemma.

Take a sequence $(r_n)_{n \in \mathbb{N}}$ of elements in K which converges to a real number r in K . Then, as the point $h(r_n, \theta_0)$ belongs to $\{r_n\} \times \mathbb{S}^1$ and to F_{θ_0} , the accumulation points of the sequence $(h(r_n, \theta_0))_n$ belong to $\{r\} \times \mathbb{S}^1$ and to F_{θ_0} : there is only one accumulation point which is the point $h(r, \theta_0)$. For any angle θ , we have $h(r, \theta) = R_{\theta - \theta_0} \circ h(r, \theta_0)$. This implies that the map h is continuous. This map is one-to-one: if $h(r, \theta) = h(r', \theta')$, this last point belongs to $\{r\} \times \mathbb{S}^1 = \{r'\} \times \mathbb{S}^1$ so that $r = r'$ and $R_{\theta - \theta_0} \circ h(r, \theta_0) = R_{\theta' - \theta_0} \circ h(r, \theta_0)$ so that $\theta = \theta'$. This map is onto by definition of the set $B = K \times \mathbb{S}^1$ which is the union of the F_θ . As this map is defined on a compact set, it is a homeomorphism.

Take any orientation-preserving homeomorphism f of the circle with $f(\theta) = \theta'$. Notice that $G_{\theta'} = f G_\theta f^{-1}$ and therefore $\varphi(f)(F_\theta) = F_{\theta'}$. So, for any $(r, \theta) \in K \times \mathbb{S}^1$, as the action φ preserves each leaf of the foliation \mathcal{F} , the homeomorphism $\varphi(f)$ sends the point $h(r, \theta)$, which is the unique point in the intersection $F_\theta \cap \{r\} \times \mathbb{S}^1$, to the unique point in the intersection $F_{\theta'} \cap \{r\} \times \mathbb{S}^1$, which is $h(r, \theta')$. This implies that

$$\varphi(f) \circ h(r, \theta) = h \circ \varphi_{K, \lambda}(f)(r, \theta).$$

Now, denote by $\eta(r)$ the second projection of $h(r, 0)$. As the set F_θ is the image by the rotation of angle θ of the set F_0 , we have $h(r, \theta) = h(r, 0) + (0, \theta)$ and $h(r, \theta) = (r, \eta(r) + \theta)$. \square

4.2 Action outside the foliation

In this section, we study the action φ on each connected component of $\mathbb{A} - K \times \mathbb{S}^1$. Let $A = [r_1, r_2] \times \mathbb{S}^1$ be the closure of such a component. By the last subsection, such a set is invariant under the action φ . This subsection is dedicated to the proof of the following proposition.

Proposition 4.3. *The restriction φ^A of the action φ to A is conjugate to a_+ (or equivalently to a_-) via an orientation preserving homeomorphism.*

Proof. Notice that, for any point θ of the circle, the action $\varphi_{G_\theta}^A$ admits no fixed point in the interior of A by definition of $K \times \mathbb{S}^1$: we will often use this fact.

Let us begin by sketching the proof of this proposition. We prove that, for any point θ of the circle, the morphism $\varphi_{G_\theta}^A$ can be lifted to a morphism $\tilde{\varphi}_\theta^A$ from the group G_θ to the group $\text{Homeo}_\mathbb{Z}([r_1, r_2] \times \mathbb{R})$ of homeomorphisms of the closed band which commute to the translations $(r, x) \mapsto (r, x + n)$, where n is an integer. Moreover, this group morphism can be chosen so that it has a bounded orbit. We will then find a continuum $\widetilde{G_\theta}$ with empty interior which touches both boundary components of the band and is invariant under the action $\tilde{\varphi}_\theta^A$. We prove then that the sets $G_\theta = \pi(\widetilde{G_\theta})$, where $\pi : [r_1, r_2] \times \mathbb{R} \rightarrow A$ is the projection, are pairwise disjoint and are simple paths which join the two boundary components of the annulus A . We see then that the group $\varphi(G_\theta \cap G_{\theta'})$, for $\theta \neq \theta'$, admits a unique fixed point $a(\theta, \theta')$ on G_θ . This last map a turns out to be continuous and allows us to build a conjugacy between φ^A and a_+ .

The following lemma is necessary to build the invariant sets G_θ :

For a homeomorphism g in $\text{Homeo}_0(A)$, we denote by \tilde{g} the lift of g to $\text{Homeo}_\mathbb{Z}([r_1, r_2] \times \mathbb{R})$ (this means that $\pi \circ \tilde{g} = g \circ \pi$) with $\tilde{g}([r_1, r_2] \times \{0\}) \in [r_1, r_2] \times [-1/2, 1/2]$. Let us denote by $D \subset \mathbb{R}^2$ the fundamental domain $[r_1, r_2] \times [-1/2, 1/2]$ for the action of \mathbb{Z} on $[r_1, r_2] \times \mathbb{R}$.

Lemma 4.4. *The map $\text{Homeo}_0(\mathbb{S}^1) \rightarrow \mathbb{R}_+$, which associates, to any homeomorphism f in $\text{Homeo}_0(\mathbb{S}^1)$, the diameter of the image under $\varphi(f)$ (or equivalently under any lift of $\varphi(f)$) of the fundamental domain D , is bounded.*

Remark: the continuity hypothesis on the morphism φ is not used in the proof of this lemma.

Proof. For a group G generated by a finite set S and for an element g in G , we denote by $l_S(g)$ the word length of g with respect to S , which is the minimal number of factors necessary to write g as a product of elements of $S \cup S^{-1}$, where S^{-1} is the set of inverses of elements in S . In order to prove the lemma, we need the following result which can be easily deduced from lemma 4.4 in [12], which is inspired from [1].

Lemma 4.5. *There exist constants $C > 0$ and $C' \in \mathbb{R}$ such that, for any sequence $(f_n)_{n \in \mathbb{N}}$ of elements of the group $\text{Homeo}_0(\mathbb{S}^1)$, there exists a finite set $S \subset \text{Homeo}_0(\mathbb{S}^1)$ such that:*

- for any integer n , the element f_n belongs to the group generated by S ;
- $\forall n \in \mathbb{N}$, $l_S(f_n) \leq C \log(n) + C'$.

Let us prove now lemma 4.4 by contradiction. Suppose that there exists a sequence $(f_n)_{n \in \mathbb{N}}$ of elements of the group $\text{Homeo}_0(\mathbb{S}^1)$ such that, for any integer n :

$$\text{diam}(\widetilde{\varphi(f_n)}(D)) \geq n.$$

We apply then the above lemma to this sequence to obtain a finite subset S included in $\text{Homeo}_0(\mathbb{S}^1)$ such that the conclusion of the lemma holds. Let $f_n = s_{1,n} s_{2,n} \dots s_{w_n,n}$, where $w_n \leq C \log(n) + C'$ and the $s_{i,j}$'s are elements of $S \cup S^{-1}$. We prove now that this implies that the diameter of $\widetilde{\varphi(f_n)}(D)$ grows at most at a logarithmic speed, which is a contradiction with the hypothesis we made on the sequence $(f_n)_n$.

Let us denote by M the maximum of the quantities $\|\widetilde{\varphi(s)}(x) - x\|$, where s varies over $S \cup S^{-1}$ and the point x varies over \mathbb{R}^2 (or equivalently over the compact set D) and where $\|\cdot\|$ denotes the euclidean norm. Take now two points x and y in D . Then for any integer n

$$\begin{aligned} & \left\| \widetilde{\varphi(s_{1,n})} \widetilde{\varphi(s_{2,n})} \dots \widetilde{\varphi(s_{w_n,n})}(x) - \widetilde{\varphi(s_{1,n})} \widetilde{\varphi(s_{2,n})} \dots \widetilde{\varphi(s_{w_n,n})}(y) \right\| \leq \\ & \left\| \widetilde{\varphi(s_{1,n})} \widetilde{\varphi(s_{2,n})} \dots \widetilde{\varphi(s_{w_n,n})}(x) - x \right\| + \left\| \widetilde{\varphi(s_{1,n})} \widetilde{\varphi(s_{2,n})} \dots \widetilde{\varphi(s_{w_n,n})}(y) - y \right\| + \|x - y\|. \end{aligned}$$

But for any point z in D , we have:

$$\begin{aligned} \left\| \widetilde{\varphi(s_{1,n})} \widetilde{\varphi(s_{2,n})} \dots \widetilde{\varphi(s_{w_n,n})}(z) - z \right\| &\leq \sum_{k=1}^{w_n-1} \left\| \widetilde{\varphi(s_{k,n})} \dots \widetilde{\varphi(s_{w_n,n})}(z) - \widetilde{\varphi(s_{k+1,n})} \dots \widetilde{\varphi(s_{w_n,n})}(z) \right\| \\ &\leq (w_n - 1)M. \end{aligned}$$

Hence $\text{diam}(\widetilde{\varphi(f_n)}(D)) \leq 2(w_n - 1)M + \text{diam}(D)$, which is in contradiction with the hypothesis we made on the sequence $(f_n)_n$. \square

Let θ_0 be a point of the circle.

Lemma 4.6. *There exists a group morphism $\tilde{\varphi}_{\theta_0}^A : G_{\theta_0} \rightarrow \text{Homeo}_{\mathbb{Z}}([r_1, r_2] \times \mathbb{R})$ such that:*

- *for any homeomorphism f in G_{θ_0} , $\Pi \circ \tilde{\varphi}_{\theta_0}^A(f) = \varphi^A(f)$, where $\Pi : \text{Homeo}_{\mathbb{Z}}([r_1, r_2] \times \mathbb{R}) \rightarrow \text{Homeo}_0(A)$ is the projection;*
- *the subset $\{\tilde{\varphi}_{\theta_0}^A(f)((r_1, 0)), f \in G_{\theta_0}\}$ of the band $[r_1, r_2] \times \mathbb{R}$ is bounded.*

Moreover, the morphism $\tilde{\varphi}_{\theta_0}^A$ is continuous.

Remark: The continuity hypothesis on the morphism φ is not necessary to prove the first part of this lemma.

Proof. As the topological space G_{θ_0} is contractible and the map $\Pi : \text{Homeo}_{\mathbb{Z}}([r_1, r_2] \times \mathbb{R}) \rightarrow \text{Homeo}_0(A)$ is a covering, there exists a (unique) continuous map $\eta : G_{\theta_0} \rightarrow \text{Homeo}_{\mathbb{Z}}([r_1, r_2] \times \mathbb{R})$ which lifts the map $\varphi|_{G_{\theta_0}}^A$ and sends the identity to the identity. Then the map

$$\begin{aligned} G_{\theta_0} \times G_{\theta_0} &\rightarrow \text{Homeo}_{\mathbb{Z}}([r_1, r_2] \times \mathbb{R}) \\ (f, g) &\mapsto \eta(fg)^{-1} \eta(f) \eta(g) \end{aligned}$$

is continuous and its image is contained in the discrete space of integral translations: it is constant and the map η is a group morphism. Two group morphisms which lift the group morphism $\varphi|_{G_{\theta_0}}^A$ differ by a morphism $G_{\theta_0} \rightarrow \mathbb{Z}$. However, as the group G_{θ_0} is simple (hence perfect), such a group morphism is trivial and $\eta = \tilde{\varphi}_{\theta_0}^A$. The action $\varphi|_{G_{\theta_0}}^A$ admits fixed points on the boundary of the annulus A . Hence, the action η admits fixed points on the boundary of $[r_1, r_2] \times \mathbb{R}$ and any orbit on one of these boundary components is bounded. \square

Let us denote

$$F = \overline{\bigcup_{f \in G_{\theta_0}} \tilde{\varphi}_{\theta_0}^A(f)([r_1, r_2] \times (-\infty, 0])}.$$

By the two above lemmas, there exists $M > 0$ such that $F \subset [r_1, r_2] \times (-\infty, M]$. Moreover, the closed set F is invariant under the action $\tilde{\varphi}_{\theta_0}^A$. Denote by U the connected component of the complementary of $F \cup [r_1, r_2] \times \mathbb{R}$ which contains the open subset $(r_1, r_2) \times (M, +\infty)$. By construction, the open set U is invariant under the action $\tilde{\varphi}_{\theta_0}^A$ (the interior of a fundamental domain far on the right must be sent in U by any homeomorphism in the image of $\tilde{\varphi}_{\theta_0}^A$, by the two above lemmas). Consider the topological space B which is the disjoint union of the band $[r_1, r_2] \times \mathbb{R}$ with a point $\{+\infty\}$ and for which a basis of neighbourhood of the point $+\infty$ are given by the sets of the form $[r_1, r_2] \times (A, +\infty) \cup \{+\infty\}$. Now, let us consider the prime end compactification of $U \subset B$ (see [9] for the background on prime end compactification). The space of prime ends of the simply connected open set U , on which there is a natural action ψ of the group G_{θ_0} induced by the action $\tilde{\varphi}_{\theta_0}^A$, is homeomorphic to a circle. By the following lemma, the action ψ is continuous.

Lemma 4.7. *Let W be a connected relatively compact open subset of the plane. Let us denote by $B(W)$ the space of prime ends of W . The map*

$$t : \text{Homeo}(\overline{W}) \rightarrow \text{Homeo}(B(W)),$$

which, to a homeomorphism f of \overline{W} , associates the induced homeomorphism on the space of prime ends of W , is continuous.

Proof. As the map t is a group morphism and as the space $B(W)$ is homeomorphic to a finite union of circles, it suffices to prove that, for any prime end ξ in $B(W)$, the map

$$\begin{aligned} \text{Homeo}(\overline{W}) &\rightarrow B(W) \\ f &\mapsto t(f)(\xi) \end{aligned}$$

is continuous at the identity. Let us fix such a prime end ξ . Let us denote by

$$V_1 \supset V_2 \supset \dots \supset V_n \supset \dots$$

a prime chain which defines the prime end ξ . If we denote by \tilde{V}_n the space of prime points of W which divide V_n , then the \tilde{V}_n 's are a basis of neighbourhoods of the point ξ (see section 3 in [9]). Let us fix a natural integer n and $p > n$. If the uniform distance between a homeomorphism f in $\text{Homeo}(\overline{W})$ and the identity is lower than the distance between the frontier $\text{Fr}_W(V_n)$ of V_n in W and the frontier $\text{Fr}_W(V_p)$ of V_p in W , then the set $f(\text{Fr}_W(V_n))$ does not meet $\text{Fr}_W(V_p)$. By lemma 4 in [9], if the homeomorphism f is sufficiently close to the identity, then $f(V_p) \subset V_n$ and $t(f)(\tilde{V}_p) \subset \tilde{V}_n$. This implies that, for f in such a neighbourhood, the point $t(f)(\xi)$ belongs to \tilde{V}_n , which is what we wanted to prove. \square

For a prime end ξ of U defined by a prime chain

$$V_1 \supset V_2 \supset \dots \supset V_n \supset \dots,$$

the *impression* of ξ is the intersection of the closures (in B) of the open sets V_n . This set does not depend on the prime chain representing ξ . The *principal set* of ξ is the set of points p in B such that there exists a prime chain

$$V_1 \supset V_2 \supset \dots \supset V_n \supset \dots$$

defining ξ such that the sequence of frontiers in U of V_n converges for the Hausdorff topology to the single-point set $\{p\}$. Notice that the subset of prime ends of U whose impression contains a point of $\{r_1, r_2\} \times \mathbb{R} \cup \{+\infty\}$ is closed. Denote by I a connected component of the complementary of this set (the set I is an open interval). By arc-connectedness of the group G_{θ_0} and continuity of the action ψ , the open interval I is invariant under the action ψ . Let ψ' be the restriction of the action ψ to the interval I .

Lemma 4.8. *The action ψ' has no fixed point.*

Proof. We prove this lemma by contradiction. More precisely, we prove that, if this action has a fixed point, then the action $\tilde{\varphi}_{\theta_0}^A$ has a fixed point in the interior of the band $[r_1, r_2] \times \mathbb{R}$, which is not the case. Suppose the action ψ' admits a fixed point. If the set of fixed points of the action ψ' has non-empty interior in I , as accessible prime ends are dense in I (see [9]), there exists an accessible prime end which is fixed under ψ . Therefore, the only point in the principal set of this prime end, which is contained in the interior of the band $[r_1, r_2] \times \mathbb{R}$, is fixed under the action $\tilde{\varphi}_{\theta_0}^A$.

Suppose now that this set of fixed points has an empty interior. Take a connected component of the complementary in I of the set of fixed points of ψ' and an endpoint e of this interval. Take an interval J of the circle whose interior contains the point θ_0 . Then, according to the previous section in which we describe the continuous actions of the group $\text{Homeo}_c(\mathbb{R})$ on the real line, the set of fixed points of $\psi(G_J)$ contains a closed interval \mathcal{J} with non-empty interior which contains the point e (not necessarily in its interior). As accessible prime ends are dense in P , this closed interval contains an accessible prime end. Then, as the principal set of this prime end is reduced to a point p , this point p is fixed by the group $\tilde{\varphi}_{\theta_0}^A(G_J)$. As a result, the group $\tilde{\varphi}_{\theta_0}^A(G_J)$ of homeomorphisms of the band $[r_1, r_2] \times \mathbb{R}$ admits a non-empty set H_J of fixed points which is included in the closure C_J of the union of the principal sets of prime ends in \mathcal{J} , which is compact. Moreover, this set is included in the interior of the band $[r_1, r_2] \times \mathbb{R}$ as it is contained in the union of the impression of prime ends in \mathcal{J} and by definition of the interval I . For any closed interval J' whose interior contains the point θ_0 and which is included in J , the set $H_{J'}$ of fixed points of $\tilde{\varphi}_{\theta_0}^A(G_{J'})$ which are included in C_J is non empty. Moreover, if the interval J'' is included in the interval J' , then $H_{J''} \subset H_{J'}$. By compactness, the intersection of those sets, which is contained in the set of fixed point of the action $\tilde{\varphi}_{\theta_0}^A$, is non-empty. \square

By the description of continuous actions of the group $\text{Homeo}_c(\mathbb{R})$ on the real line with no global fixed points (see section 3 of this article), the action ψ is transitive on the open interval I . Hence, all the prime ends in I are accessible, by density of accessible prime ends. Moreover, by this same description, for any point $\theta \neq \theta_0$ of the circle, there is a unique prime end $e_\theta \in I$ which is fixed under $\psi|_{G_{\theta_0} \cap G_\theta}$. Moreover, the interval I is the union of these prime ends. For any point $\theta \neq \theta_0$ of the circle, let us denote by \tilde{x}_θ the unique point in the principal set of the prime end e_θ . For any homeomorphism f in G_{θ_0} which sends the point $\theta \neq \theta_0$ to the point θ' , then $\tilde{\varphi}_{\theta_0}^A(f)(\tilde{x}_\theta) = \tilde{x}_{\theta'}$, as $G_{\theta'} = fG_\theta f^{-1}$. Let $\widetilde{L_{\theta_0}} = \{\tilde{x}_\theta, \theta \in \mathbb{S}^1 - \{\theta_0\}\}$ and $L_{\theta_0} = \pi(\widetilde{L_{\theta_0}})$.

Lemma 4.9. *The map*

$$\begin{aligned} \mathbb{S}^1 - \{\theta_0\} &\rightarrow (r_1, r_2) \times \mathbb{R} \\ \theta &\mapsto \tilde{x}_\theta \end{aligned}$$

is one-to-one and continuous. Moreover, the limit sets $\lim_{\theta \rightarrow \theta_0^+} \tilde{x}_\theta$ and $\lim_{\theta \rightarrow \theta_0^-} \tilde{x}_\theta$ each contain exactly one point of the boundary $\{r_1, r_2\} \times \mathbb{R}$ of the band.

Proof. Take $\theta' \neq \theta$ and suppose by contradiction that $\tilde{x}_\theta = \tilde{x}_{\theta'}$. Take $\theta'' \neq \theta$ in the same connected component of $\mathbb{S}^1 - \{\theta_0, \theta\}$ as θ' such that $\tilde{x}_{\theta''} \neq \tilde{x}_\theta$. Consider a homeomorphism f of the circle in $G_{\theta_0} \cap G_\theta$ which sends the point θ' to the point θ'' . Then the homeomorphism $\tilde{\varphi}_{\theta_0}^A(f)$ of the band fixes the point \tilde{x}_θ and sends the point $\tilde{x}_{\theta'}$ to the point $\tilde{x}_{\theta''}$, which is not possible. The map is one-to-one.

Now, let us prove that the considered map is continuous. Take a vector field of the circle which vanishes only on a small connected neighbourhood N of θ_0 . Let us denote by $(h^t)_{t \in \mathbb{R}}$ the one-parameter group generated by this vector field. Fix a point $\theta_1 \in \mathbb{S}^1 - N$. For any point $\theta \in \mathbb{S}^1 - N$, denote by $t(\theta)$ the unique time t such that $h^t(\theta_1) = \theta$. The map $\theta \mapsto t(\theta)$ is then a homeomorphism $\mathbb{S}^1 - N \rightarrow \mathbb{R}$. Now, the relation $\tilde{x}_\theta = \varphi(h^{t(\theta)})(\tilde{x}_{\theta_1})$ and the continuity of the action φ implies that the map $\theta \mapsto \tilde{x}_\theta$ is continuous, as the neighbourhood N can be taken arbitrarily small.

The following lemma implies that the intersection of each of the limit sets $\lim_{\theta \rightarrow \theta_0^+} \tilde{x}_\theta$ and $\lim_{\theta \rightarrow \theta_0^-} \tilde{x}_\theta$ with the boundary $\{r_1, r_2\} \times \mathbb{R}$ of the band is nonempty. Indeed, otherwise, these limit sets would provide a nonempty compact connected invariant set for the action $\tilde{\varphi}_{\theta_0}^A$.

Lemma 4.10. *The action $\tilde{\varphi}_{\theta_0}^A$ admits no nonempty compact connected invariant set included in $(r_1, r_2) \times \mathbb{R}$.*

Proof. Consider the unbounded component of the complementary of this set. The action $\tilde{\varphi}_{\theta_0}^A$ induces an action on the space of prime ends of this set, which is homeomorphic to a circle. By proposition 3.10, such an action has a fixed point. By an argument similar to the one in the proof of lemma 4.8, the action $\tilde{\varphi}_{\theta_0}^A$ admits then a fixed point in the interior of the band, which is excluded by hypothesis. \square

It remains to prove that the intersections $\lim_{\theta \rightarrow \theta_0^+} \tilde{x}_\theta \cap \{r_1, r_2\} \times \mathbb{R}$ and $\lim_{\theta \rightarrow \theta_0^-} \tilde{x}_\theta \cap \{r_1, r_2\} \times \mathbb{R}$ are reduced to a point. Suppose for instance that the intersection $\lim_{\theta \rightarrow \theta_0^+} \tilde{x}_\theta \cap \{r_1\} \times \mathbb{R}$ contains at least two points. Recall that, by proposition 3.11, we know that the restriction of the action φ to $\{r_1\} \times \mathbb{S}^1$ is a conjugacy by a homeomorphism $\mathbb{S}^1 \rightarrow \{r_1\} \times \mathbb{S}^1$. Therefore, the action $\varphi|_{G_{\theta_0}}$ fixes a point p and is transitive on $\{r_1\} \times \mathbb{S}^1 - \{p\}$. Moreover, as any orbit of the action $\tilde{\varphi}_{\theta_0}^A$ is bounded by lemmas 4.4 and 4.6, this last action pointwise fixes the set $\pi^{-1}(\{p\})$ and is transitive on each connected component of $\{r_1\} \times \mathbb{R} - \pi^{-1}(\{p\})$. Therefore, the intersection $\lim_{\theta \rightarrow \theta_0^+} \tilde{x}_\theta \cap \{r_1\} \times \mathbb{R}$, which is closed and invariant under the action $\tilde{\varphi}_{\theta_0}^A$, contains two distinct lifts \tilde{p} and \tilde{p}' of the point p . Take a sequence $(\tilde{x}_{\theta_n})_{n \in \mathbb{N}}$ of points in $\widetilde{L_{\theta_0}}$, where θ_n tends to θ_0^+ , which converges to the point \tilde{p} and a sequence $(\tilde{x}_{\theta'_n})_{n \in \mathbb{N}}$ of points in $\widetilde{L_{\theta_0}}$, where θ'_n tends to θ_0^+ , which converges to the point \tilde{p}' . Taking a subsequence if necessary, we may suppose that the sequences $(\theta_n)_n$ and $(\theta'_n)_n$ are monotonous and that, for any integer n , the angle θ'_n is between the angles θ_n and θ_{n+1} . Take a homeomorphism f in G_{θ_0} which, for any n , sends the point θ_n to the point θ'_n . Then, for any n , the homeomorphism $\tilde{\varphi}_{\theta_0}^A(f)$ sends the point \tilde{x}_{θ_n} to the point $\tilde{x}_{\theta'_n}$. By continuity, this homeomorphism sends the point \tilde{p} to the point \tilde{p}' . This is not possible as these points are fixed under the action $\tilde{\varphi}_{\theta_0}^A$. \square

Note that we do not know for the moment that the points $\lim_{\theta \rightarrow \theta_0^+} \tilde{x}_\theta$ and $\lim_{\theta \rightarrow \theta_0^-} \tilde{x}_\theta$ lie on different boundary components of A .

For any angle $\theta \neq \theta_0$, we define $L_\theta = R_{\theta - \theta_0}(L_{\theta_0})$. Notice that, for any angle θ , the set L_θ is invariant under the action $\varphi_{|G_\theta}^A$. Indeed $G_\theta = R_{\theta - \theta_0}G_{\theta_0}R_{\theta - \theta_0}^{-1}$ which implies that $\varphi^A(G_\theta) = R_{\theta - \theta_0}\varphi^A(G_{\theta_0})R_{\theta - \theta_0}^{-1}$ and, as the set L_{θ_0} is invariant under the action $\varphi_{|G_{\theta_0}}^A$, the claim follows. Moreover, for an orientation-preserving homeomorphism f of the circle which sends an angle θ to another angle θ' , the homeomorphism $\varphi(f)$ sends the set L_θ to the set $L_{\theta'}$. To prove this last fact, use that the homeomorphism f can be written as the composition of a homeomorphism which fixes the point θ with a rotation.

Lemma 4.11. *The sets L_θ are pairwise disjoint. Moreover, there exists a homeomorphism from the compact interval $[0, 1]$ to the closure of L_{θ_0} which sends the interval $(0, 1)$ to the set L_{θ_0} .*

Proof. By using rotations, we see that it suffices to prove that, for any angle $\theta \neq \theta_0$, $L_{\theta_0} \cap L_\theta = \emptyset$. Fix such an angle θ . Remember that the restriction of the action ψ' to $G_\theta \cap G_{\theta_0}$ has one fixed point e_θ and is transitive on each connected component of $I - \{e_\theta\}$. Denote by E_θ the subset of I consisting of prime ends whose only point in the principal set belongs to $\pi^{-1}(L_\theta)$. As this set is invariant under $\psi_{|G_\theta \cap G_{\theta_0}}$, it is either empty, or the one-point set $\{e_\theta\}$ or one of the connected components of $P - \{e_\theta\}$ or the closure of one of these components or P . In the last case, we would have $\widetilde{L_{\theta_0}} \subset \pi^{-1}(L_\theta)$ and $L_{\theta_0} \subset L_\theta$. Using homeomorphisms of the circle which fix the angle θ_0 and send the angle θ to another angle θ' , we see that, for any angle θ' , $L_{\theta_0} \subset L_{\theta'}$. This is not possible as the intersection of the closure of the set L_{θ_0} with the boundary of A is a two-point set which should be invariant under any rotation. The case where the set E_θ is a half-line (open or closed) leads to a similar contradiction by looking at one of the limit sets of L_{θ_0} . Hence, for any angle $\theta \neq \theta_0$, the intersection $L_{\theta_0} \cap L_\theta$ contains at most one point, the point x_θ .

This implies that any leaf of the form $\{r\} \times \mathbb{S}^1 \subset A$ contains at most two points of L_{θ_0} . We claim then that if one of these leaves contains two points of L_{θ_0} , then no leaf contains exactly one point of this set.

Take a point x_{θ_1} in L_{θ_0} which belongs to the leaf $\{r\} \times \mathbb{S}^1$. Suppose that there exists another point of the set L_{θ_0} on this leaf. This implies that the point x_{θ_1} belongs to L_{θ_1} . Using homeomorphisms in G_{θ_0} which send the point θ_1 to another point $\theta \neq \theta_0$ of the circle, we see that, for any angle $\theta \neq \theta_0$ the point x_θ belongs to L_θ . Hence, each circular leaf which meets the set L_{θ_0} contains exactly two points of this set.

If x is another point of L_{θ_0} on the leaf $\{r\} \times \mathbb{S}^1$, then there exists $\alpha \in \mathbb{S}^1 - \{0\}$ such that $R_\alpha(x) = x_\theta \in L_{\alpha + \theta_0}$. Therefore, $\alpha + \theta_0 = \theta$ and the point x is necessarily the point $R_{\theta_0 - \theta}(x_\theta) = x_{2\theta_0 - \theta}$. Hence, the map $\theta \mapsto p_1(x_\theta)$, where $p_1 : [r_1, r_2] \times \mathbb{R} \mapsto [r_1, r_2]$ is the projection, is strictly monotonous (as it is one-to-one) on $(\theta_0, \theta_0 + \frac{1}{2}]$ and on $[\theta_0 + \frac{1}{2}, \theta_0 + 1)$. Moreover, if this map was not globally one-to-one, the circular leaf which contains the point $x_{\theta_0 + \frac{1}{2}}$ would contain only one point, which is not possible. Hence, this map is strictly monotonous. This implies also that the sets L_θ , for $\theta \neq \theta_0$, are disjoint from L_{θ_0} . The monotonicity of the map $\theta \mapsto p_1(x_\theta)$, combined with lemma 4.9 implies the second part of the lemma. \square

Now, we can end the proof of the proposition. By theorem 3.1, for any $\theta \in \mathbb{S}^1$ and any $r \in (0, 1)$ the action $\varphi_{|G_\theta \cap G_{\theta+r}}^A$ admits a unique fixed point $a(r, \theta)$ in L_θ . Moreover, for any point θ of the circle, the map

$$\begin{aligned} (0, 1) &\rightarrow A \\ r &\mapsto a(r, \theta) \end{aligned}$$

is one-to-one, continuous and extends to a continuous map $[0, 1] \rightarrow A$ which allows us to define $a(0, \theta)$, which is the fixed point of $\varphi_{|G_\theta}^A$ on one boundary component of A , and $a(1, \theta)$, which is the fixed point of $\varphi_{|G_\theta}^A$ on the other boundary component of A . By the last lemma, the map a is one-to-one. Moreover, for every point α of the circle, and every point (r, θ) of the closed annulus $\mathbb{A} = [0, 1] \times \mathbb{S}^1$, $a(r, \theta + \alpha) = R_\alpha(a(r, \theta))$. Indeed, the point $R_\alpha(a(r, \theta))$ is the only fixed point of $\varphi_{|G_{\theta+\alpha} \cap G_{\theta+\alpha+r}}^A = R_\alpha \varphi_{|G_\theta \cap G_{\theta+r}}^A R_\alpha^{-1}$ in $L_{\theta+\alpha} = R_\alpha(L_\theta)$. This implies that the map a is continuous: it is a homeomorphism $\mathbb{A} \rightarrow A$. It remains to prove that it defines a conjugacy with a_+ . Take a homeomorphism f , and a point $(r, \theta) \in \mathbb{A}$. Notice that

$$fG_\theta \cap G_{\theta+r}f^{-1} = G_{f(\theta)} \cap G_{f(\theta+r)} = G_{f(\theta)} \cap G_{f(\theta) + (\tilde{f}(\theta+r) - \tilde{f}(\theta))}$$

and that $\varphi^A(f)(L_\theta) = L_{f(\theta)}$. Therefore, $\varphi^A(f)(a(r, \theta)) = a(a_+(f)(r, \theta))$. The action φ^A is conjugate to a_+ . As the actions a_+ and a_- are conjugate both by an orientation-preserving homeomorphism and an orientation-reversing homeomorphism, the proposition is proved. \square

4.3 Global conjugacy

This section is devoted to the end of the proof of theorem 2.1 in the case of actions on an annulus.

By lemma 4.2, our action φ , restricted to the union $K \times \mathbb{S}^1$ of the F_θ , where θ varies over the circle, is conjugate to the restriction of $\varphi_{K, \lambda}$ to $K \times \mathbb{S}^1$ by a homeomorphism of the form $(r, \theta) \mapsto (r, \eta(r) + \theta)$ for any λ . Conjugating φ by a homeomorphism of the annulus of the form $(r, \theta) \mapsto (r, \hat{\eta}(r) + \theta)$, where $\hat{\eta}$ is a continuous function which is equal to η on K , we may assume from now on that the action φ is equal to $\varphi_{K, \lambda}$ for any λ on $K \times \mathbb{S}^1$.

Moreover, for each connected component $(r_1^i, r_2^i) \times \mathbb{S}^1$ of the complementary of the closed set $K \times \mathbb{S}^1$, the restriction of this action to $[r_1^i, r_2^i] \times \mathbb{S}^1$ is conjugate to a_+ via an orientation-preserving homeomorphism $g_i : [0, 1] \times \mathbb{S}^1 \rightarrow [r_1^i, r_2^i] \times \mathbb{S}^1$, by proposition 4.3.

We now define a particular continuous function $\lambda_0 : [0, 1] - K \rightarrow \{-1, +1\}$ such that the action φ is conjugate to φ_{K, λ_0} . For any index i , let us denote by d_i^+ the diameter of the set $g_i(\{(r, 0), 0 \leq r \leq 1\})$ and by d_i^- the diameter of the set $g_i(\{(r, -r), 0 \leq r \leq 1\})$. If $d_i^+ \leq d_i^-$, we define λ_0 to be identically equal to $+1$ on the interval (r_1^i, r_2^i) . Otherwise, *i.e.* if $d_i^+ > d_i^-$, we define the map λ_0 to be identically equal to -1 on the interval (r_1^i, r_2^i) .

Now, let us define a conjugacy between the action φ and the action φ_{K, λ_0} . Notice that, by what is above, for any connected component $(r_1^i, r_2^i) \times \mathbb{S}^1$ of the complementary of $K \times \mathbb{S}^1$, there exists an orientation-preserving homeomorphism h_i of $[r_1^i, r_2^i] \times \mathbb{S}^1$ such that, for any orientation-preserving homeomorphism f of the circle,

$$h_i \circ \varphi_{K, \lambda_0}(f)|_{[r_1^i, r_2^i] \times \mathbb{S}^1} = \varphi(f)|_{[r_1^i, r_2^i] \times \mathbb{S}^1} \circ h_i.$$

Then, for any point $(r, \theta) \in [r_1^i, r_2^i] \times \mathbb{S}^1$ and any angle $\alpha \in \mathbb{S}^1$, $h_i(r, \theta + \alpha) = h_i(r, \theta) + (0, \alpha)$. (Recall that we supposed at the beginning of this section that the morphism φ sent the rotation of the circle of angle α on the rotation of the annulus of angle α .)

Now, let us denote by $h : \mathbb{A} \rightarrow \mathbb{A}$ the map which is equal to the identity on $K \times \mathbb{S}^1$ and which is equal to h_i on the connected component $(r_1^i, r_2^i) \times \mathbb{S}^1$ of the complementary of $K \times \mathbb{S}^1$. It is clear that the map h is a bijection and that, for a homeomorphism f in $\text{Homeo}_0(\mathbb{S}^1)$, the relation $h \circ \varphi_{K, \lambda_0}(f) = \varphi(f) \circ h$ holds. Moreover, the map h commutes with the rotations.

It remains to prove that the map h is continuous to end the proof. As the map h commutes with rotations, it suffices to prove that the map

$$\begin{aligned} \eta : [0, 1] &\rightarrow \mathbb{A} \\ r &\mapsto h(r, 0) \end{aligned}$$

is continuous. Notice first that, for a connected component (r_1^i, r_2^i) of the complementary of K , the limit $\lim_{r \rightarrow r_1^{i+}} \eta(r)$ is equal to $h_i(r_1, 0)$. This last point is the only point of F_0 on $\{r_1\} \times \mathbb{S}^1$ and is therefore equal to $\eta(r_1) = (r_1, 0)$. The map η is also continuous on the left at r_2^i . It suffices now to establish that, for any sequence $((r_1^n, r_2^n))_{n \in \mathbb{N}}$ of connected components of the complementary of K such that the sequence $(r_1^n)_{n \in \mathbb{N}}$ is monotonous and converges to a point r_∞ which belongs to the compact set K , for any sequence $(r_n)_{n \in \mathbb{N}}$ of real numbers such that, for any integer n , $r_n \in (r_1^n, r_2^n)$, the sequence $(\eta(r_n))_{n \in \mathbb{N}}$ converges to $\lambda(r_\infty) = (r_\infty, 0)$. Fix now such a sequence $((r_1^n, r_2^n))_{n \in \mathbb{N}}$.

Lemma 4.12. *For n sufficiently large, one of the curves $h([r_1^n, r_2^n] \times \{0\})$ and $h(\{(r, \theta - \frac{r-r_1^n}{r_2^n-r_1^n}, r_1^n \leq r \leq r_2^n)\})$ is homotopic with fixed extremities in the closed annulus $[r_1^n, r_2^n] \times \mathbb{S}^1$ to the curve $[r_1^n, r_2^n] \times \{0\}$. We denote by c_n this curve.*

Proof. Suppose by contradiction that there exists a strictly increasing map $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ such that, for any $n \in \mathbb{N}$, neither $h([r_1^{\sigma(n)}, r_2^{\sigma(n)}] \times \{0\})$ nor $h\left(\left(r, \theta - \frac{r - r_1^{\sigma(n)}}{r_2^{\sigma(n)} - r_1^{\sigma(n)}}, r_1^{\sigma(n)} \leq r \leq r_2^{\sigma(n)}\right)\right)$ are homotopic to $[r_1^n, r_2^n] \times \{0\}$. Then, one of the two curves $h([r_1^{\sigma(n)}, r_2^{\sigma(n)}] \times \{0\})$ and $h\left(\left(r, \theta - \frac{r - r_1^{\sigma(n)}}{r_2^{\sigma(n)} - r_1^{\sigma(n)}}, r_1^{\sigma(n)} \leq r \leq r_2^{\sigma(n)}\right)\right)$, which we denote by γ_n , admits a lift $\widetilde{\gamma}_n$ such that $\widetilde{\gamma}_n \cap \left\{r_1^{\sigma(n)}\right\} \times \mathbb{R} = \left\{(r_1^{\sigma(n)}, 0)\right\}$ and $\widetilde{\gamma}_n \cap \left\{r_2^{\sigma(n)}\right\} \times \mathbb{R} = \left\{(r_2^{\sigma(n)}, k)\right\}$, where k is an integer with $|k| \geq 2$. To simplify the notations, suppose that $k \geq 2$.

We now need an intermediate result. Let $(\tilde{x}_n)_{n \in \mathbb{N}}$ and $(\tilde{y}_n)_{n \in \mathbb{N}}$ be sequences of points of the band $[0, 1] \times \mathbb{R}$ converging respectively to the points \tilde{x}_∞ and \tilde{y}_∞ such that, for any n , the points \tilde{x}_n and \tilde{y}_n belong to the curve $\widetilde{\gamma}_n$ and are not endpoints of this curve. We suppose that $\tilde{x}_\infty \neq \tilde{y}_\infty$ and that the points \tilde{x}_∞ and \tilde{y}_∞ are not limit points of the endpoints of $\widetilde{\gamma}_n$. For any integer n , there exist unique real numbers r_n and $r'_n \in (0, 1)$ such that the point $x_n = \pi(\tilde{x}_n)$ (respectively $y_n = \pi(\tilde{y}_n)$) is the unique fixed point of the group $\varphi(G_0 \cap G_{r_n})$ (respectively $\varphi(G_0 \cap G_{r'_n})$) on the curve γ_n . We claim that, for any strictly increasing map $s : \mathbb{N} \rightarrow \mathbb{N}$ such that the sequences $(r_{s(n)})_{n \in \mathbb{N}}$ and $(r'_{s(n)})_{n \in \mathbb{N}}$ converge respectively to R and R' , then $R \neq R'$. Moreover, the real numbers R as R' are different from 0 or 1.

Let us begin by proving this claim. Suppose by contradiction that there exists a map s such that $R = R'$. We may suppose (by extracting a subsequence and by changing the roles of x_n and y_n if necessary) that, for any n , $r_{s(n)} < r'_{s(n)}$. Then the set of points in $\gamma_{s(n)}$ which are fixed by $\varphi|_{G_0 \cap G_r}$, with $r_{s(n)} \leq r \leq r'_{s(n)}$ defines a sequence of paths which converge to an interval included in $\{r_\infty\} \times \mathbb{S}^1$. This interval has non-empty interior and is necessarily pointwise fixed by $\varphi|_{G_0 \cap G_R}$: indeed, any homeomorphism f in $G_0 \cap G_R$ fixes the points between $r_{s(n)}$ and $r'_{s(n)}$ for n sufficiently large so that the homeomorphism $\varphi(f)$ fixes the projection of the set of points on $\widetilde{\gamma}_n$ between \tilde{x}_n and \tilde{y}_n , for n sufficiently large. Hence, it also fixes the projection of the interval between the points \tilde{x}_∞ and \tilde{y}_∞ on the line $\{r_\infty\} \times \mathbb{R}$. However, there exist no such interval with non-empty interior. Let us establish now for instance that $R \neq 1$. The set of points in $\gamma_{s(n)}$ which are fixed by $\varphi|_{G_0 \cap G_r}$, with $r_{s(n)} \leq r \leq 1$ is a path which admits a lift which joins the point $\tilde{x}_{s(n)}$ to either the point $(r_2^{\sigma(s(n))}, k)$ or the point $(r_1^{\sigma(s(n))}, k)$. Taking a subsequence if necessary, this sequence of sets converges to an interval with non-empty interior included in $\{r_\infty\} \times \mathbb{S}^1$. This interval is necessarily pointwise fixed by $\varphi|_{G_0 \cap G_R}$: this is a contradiction.

Let us come back to our proof. There exists a sequence $(\tilde{x}_n)_{n \in \mathbb{N}}$ of points of the band $[0, 1] \times \mathbb{R}$ converging to the point $(r_\infty, 1)$ such that, for any n , the point \tilde{x}_n belongs to the curve $\widetilde{\gamma}_n$. Let $x_n = \pi(\tilde{x}_n)$. For any integer n , there exists a unique real number $r_n \in (0, 1)$ such that the point x_n is the unique fixed point of the group $\varphi(G_0 \cap G_{r_n})$ on the curve γ_n . For any n , take a point \tilde{y}_n on the curve $\widetilde{\gamma}_n$ such that the sequence $(\tilde{y}_n)_n$ converges to a point \tilde{p} which is neither a point of the form (r_∞, k) , where k is an integer, nor a limit point of a sequence of endpoints of the curve $\widetilde{\gamma}_n$. As above, we denote by $r'_n \in (0, 1)$ the real number associated to the point $y_n = \pi(\tilde{y}_n)$. Consider a strictly increasing map $s : \mathbb{N} \rightarrow \mathbb{N}$ such that the sequences $(r_{s(n)})_{n \in \mathbb{N}}$ and $(r'_{s(n)})_{n \in \mathbb{N}}$ converge respectively to R and R' . Take a homeomorphism f in G_0 which sends the point R of the circle to the point R' of the circle. Let $z_n = \varphi(f)(x_n)$. Then, for any n , the real number associated to the point $z_{s(n)}$ is necessarily $f(r_{s(n)})$ and the sequence $(f(r_{s(n)}))_n$ converges to R' . By the claim above, the sequence $(z_{s(n)})_n$ has the same limit as the sequence $(y_{s(n)})_n$. By continuity, this homeomorphism of the annulus sends the point $(r_\infty, 0)$ to the point $p = \pi(\tilde{p})$. This is not possible as the point $(r_\infty, 0)$ of the annulus is fixed by the group $\varphi(G_0)$. \square

Lemma 4.13. *The diameter of the curve c_n tends to 0 as n tends to $+\infty$.*

Proof. This proof is similar to the proof of the above lemma. Suppose that the sequence of diameters of c_n does not converge to 0. Let us denote by \tilde{c}_n the lift of the curve c_n whose origin is the point $(r_1^{\sigma(n)}, 0)$. There exists a subsequence $\tilde{c}_{s(n)}$ of this sequence which converges (for the Hausdorff topology) to an interval with non-empty interior. As the projection of this interval on the annulus is invariant under the action $\varphi|_{G_0}$, this compact set projects onto the whole circle $\{r_\infty\} \times \mathbb{S}^1$. Therefore, there exists a sequence $(\tilde{x}_n)_{n \in \mathbb{N}}$, where the point \tilde{x}_n belongs to $\tilde{c}_{s(n)}$, which converges to a point of the form (r_∞, k) with $k \neq 0$ (which is not the limit of a sequence of endpoints of $\tilde{c}_{s(n)}$). There also exists a sequence $(\tilde{y}_n)_{n \in \mathbb{N}}$, where

the point \tilde{x}_n belongs to $\tilde{c}_{s(n)}$, which converges to a point \tilde{p} which is not of the form (r_∞, k) , where k is an integer. As in the proof of the above lemma, let us denote $r_n \in (0, 1)$ (respectively r'_n) the real number associated to the point $x_n = \pi(\tilde{x}_n)$ (respectively $y_n = \pi(\tilde{y}_n)$). Taking a subsequences if necessary, we may suppose that the sequences $(r_n)_n$ and $(r'_n)_n$ converge respectively to the real numbers R and R' . As in the proof of the above lemma, $R \neq R'$ and both these numbers are different from 0 and 1. Take then a homeomorphism f in G_0 which sends the point R to the point R' . Then the homeomorphism $\varphi(f)$ sends the point $(r_\infty, 0)$ to the point $p = \pi(\tilde{p}) \neq (r_\infty, 0)$, which is not possible as the point $(r_\infty, 0)$ is fixed by the group $\varphi(G_0)$. \square

Now, let us prove that $c_n = h([r_1^n, r_2^n] \times \{0\})$ for n sufficiently large, which proves the continuity of the map η by the above lemma. Let us denote by c'_n the curve among $h([r_1^n, r_2^n] \times \{0\})$ and $h(\{(r, \theta - \frac{r-r_1^n}{r_2^n-r_1^n}), r_1^n \leq r \leq r_2^n\})$ which is not equal to c_n . As the homotopy class of this curve c'_n is not the homotopy class of $[r_1^n, r_2^n] \times \{0\}$, the diameter of c'_n is bounded from below by $\frac{1}{2}$. Hence, for n sufficiently large (say for $n \geq N$), the diameter of c_n is smaller than the diameter of c'_n .

Fix now $n \geq N$. The orientation of the circle defines an order on $\mathbb{S}^1 - \{0\}$. Take a homeomorphism $f \neq Id$ in G_0 such that, for any $\theta \neq 0$, $f(\theta) \geq \theta$. Then the restriction of $\varphi(f)$ to $h([r_1^n, r_2^n] \times \{0\})$ and to $h(\{(r, \theta - \frac{r-r_1^n}{r_2^n-r_1^n}), r_1^n \leq r \leq r_2^n\})$ defines an orientation on both curves (the orientation such that an arc from a point x to its image by $\varphi(f)$ is positively oriented). If $d_n^+ \leq d_n^-$, the interior of the curve c_n is equal to the curve $g_n((0, 1) \times \{0\})$: this is the only simple curve in $(r_1^n, r_2^n) \times \mathbb{S}^1$ which is invariant under $\varphi|_{G_0}$ and which is oriented from the point $(r_1^n, 0)$ to the point $(r_2^n, 0)$. Moreover, the only simple curve in $(r_1^n, r_2^n) \times \mathbb{S}^1$ which is invariant under the action φ_{K, λ_0} restricted to G_0 which is oriented from the point $(r_1^n, 0)$ to the point $(r_2^n, 0)$ is the curve $(r_1^n, r_2^n) \times \{0\}$. Finally, as the homeomorphism h is continuous on $[r_1^n, r_2^n] \times \mathbb{S}^1$ and $h \circ \varphi_{K, \lambda_0} = \varphi \circ h$, we have $h([r_1^n, r_2^n] \times \{0\}) = c_n$. If $d_n^- < d_n^+$, the interior of the curve c_n is equal to the curve $g_n(\{(r, -r), 0 < r < 1\})$: this is the only simple curve in $(r_1^n, r_2^n) \times \mathbb{S}^1$ which is invariant under $\varphi|_{G_0}$ and which is oriented from the point $(r_2^n, 0)$ to the point $(r_1^n, 0)$. Moreover, the only simple curve in $(r_1^n, r_2^n) \times \mathbb{S}^1$ which is invariant under the action φ_{K, λ_0} restricted to G_0 which is oriented from the point $(r_2^n, 0)$ to the point $(r_1^n, 0)$ is the curve $(r_1^n, r_2^n) \times \{0\}$. Moreover, as the homeomorphism h is continuous on $[r_1^n, r_2^n] \times \mathbb{S}^1$ and $h \circ \varphi_{K, \lambda_0} = \varphi \circ h$, we have $h([r_1^n, r_2^n] \times \{0\}) = c_n$.

5 Continuous actions of the group of orientation-preserving homeomorphisms of the circle on the torus

Let $\varphi : \text{Homeo}_0(\mathbb{S}^1) \rightarrow \text{Homeo}_0(\mathbb{T}^2)$ be a one-to-one continuous group morphism (if such a group morphism is not one-to-one, it is trivial). The goal of this section is to prove the following theorem and its corollary:

Theorem 5.1. *For any point x of the circle, the image under the morphism φ of the group G_x admits a global fixed point.*

Corollary 5.2. *The action φ admits an invariant essential circle: the study of this action reduces to the study of an action on an annulus.*

Here, essential means non-separating so that the surface obtained by cutting along this curve is an annulus. Using the first part of theorem 2.1 which was proved in last section, this corollary implies directly the second part of this theorem. Let us first see why this theorem implies the corollary.

Proof of the corollary. First, the image under the morphism φ of the group \mathbb{S}^1 of rotations of the circle is conjugate to the subgroup of rotations of the torus of the form:

$$\begin{aligned} \mathbb{T}^2 = \mathbb{S}^1 \times \mathbb{S}^1 &\rightarrow \mathbb{T}^2 \\ (\theta_1, \theta_2) &\mapsto (\theta_1, \theta_2 + \alpha) \end{aligned} \text{ ,}$$

where $\alpha \in \mathbb{S}^1$. Therefore, after possibly conjugating, we may suppose that the morphism φ sends the rotation of the circle of angle α on the rotation $(\theta_1, \theta_2) \mapsto (\theta_1, \theta_2 + \alpha)$.

Fix a point x_0 on the circle. If p is a fixed point for the group $\varphi(G_{x_0})$, then the essential circle $\{\varphi(R_\alpha)(p), \alpha \in \mathbb{S}^1\}$ is invariant under the action φ . The proof of this last claim is similar to the proof of lemma 4.1. \square

Proof of theorem 5.1. Let us begin by giving a sketch of the proof of this theorem. First, we prove that the diameters of the images of the fundamental domain $[0, 1]^2$ under lifts of homeomorphisms in the image of φ are uniformly bounded. We then prove that the restriction of the morphism φ to a subgroup of the form G_x lifts to a group morphism $\widetilde{\varphi}_x : G_x \rightarrow \text{Homeo}_{\mathbb{Z}^2}(\mathbb{R}^2)$, where this last group is the group of homeomorphisms of \mathbb{R}^2 which commute to the integral translations. We prove that the action $\widetilde{\varphi}_x$ can be chosen so that it has a bounded orbit. Using these facts, we find a connected subset of \mathbb{R}^2 with empty interior which is invariant under the action of $\widetilde{\varphi}_x$. Using prime ends, we can prove that the action $\varphi|_{G_x}$ has a fixed point on the projection of this connected set.

For a homeomorphism g in $\text{Homeo}_0(\mathbb{T}^2)$, we denote by \tilde{g} the lift of g to $\text{Homeo}_{\mathbb{Z}^2}(\mathbb{R}^2)$ (this means that $\pi \circ \tilde{g} = g \circ \pi$ where $\pi : \mathbb{R}^2 \rightarrow \mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ is the projection) with $\tilde{g}(0) \in [-1/2, 1/2) \times [-1/2, 1/2)$. Let us denote by $D \subset \mathbb{R}^2$ the fundamental domain $[0, 1]^2$ for the action of \mathbb{Z}^2 on \mathbb{R}^2 .

Lemma 5.3. *The map $\text{Homeo}_0(\mathbb{S}^1) \rightarrow \mathbb{R}_+$ which associates, to any homeomorphism f in $\text{Homeo}_0(\mathbb{S}^1)$, the diameter of the image under $\varphi(f)$ (or equivalently under any lift of $\varphi(f)$) of the fundamental domain D is bounded.*

Proof. The proof of this lemma is almost identical to the proof of lemma 4.4. \square

Let x_0 be a point of the circle.

Lemma 5.4. *There exists a group morphism $\widetilde{\varphi}_{x_0} : G_{x_0} \rightarrow \text{Homeo}_{\mathbb{Z}^2}(\mathbb{R}^2)$ such that:*

- *for any homeomorphism f in G_{x_0} , $\Pi \circ \widetilde{\varphi}_{x_0}(f) = \varphi(f)$, where $\Pi : \text{Homeo}_{\mathbb{Z}^2}(\mathbb{R}^2) \rightarrow \text{Homeo}_0(\mathbb{T}^2)$ is the projection;*
- *the subset $\{\widetilde{\varphi}_{x_0}(f)(0), f \in G_{x_0}\}$ is bounded.*

Moreover, the morphism $\widetilde{\varphi}_{x_0}$ is continuous.

Proof. Let $G = \varphi(G_{x_0})$. Observe that the map

$$\begin{aligned} G \times G &\rightarrow \mathbb{Z}^2 \\ (f, g) &\mapsto \widetilde{fg}^{-1}(\tilde{f}(\tilde{g}(0))) \end{aligned}$$

defines a 2-cocycle on the group G (see [2] or [7] for more about the cohomology of groups). Moreover, by lemma 5.3, this cocycle is bounded. However, as the group G is isomorphic to the group $\text{Homeo}_c(\mathbb{R})$, the group $H_b^2(G, \mathbb{Z}^2)$ is trivial (see [8] and [11]). This implies that there exists a bounded map $b : G \rightarrow \mathbb{Z}^2$ such that:

$$\forall f, g \in G, \widetilde{fg}^{-1}(\tilde{f}(\tilde{g}(0))) = b(f) + b(g) - b(fg).$$

It suffices then to take for $\widetilde{\varphi}_{x_0}$ the composition of the morphism $\varphi|_{G_{x_0}}$ with the morphism

$$\begin{aligned} G &\rightarrow \text{Homeo}_{\mathbb{Z}^2}(\mathbb{R}^2) \\ f &\mapsto \tilde{f} + b(f) \end{aligned} .$$

For this action, the orbit of 0 is bounded by construction. It suffices now to prove that this action is continuous. As the topological space G_{x_0} is contractible and the map $\Pi : \text{Homeo}_{\mathbb{Z}^2}(\mathbb{R}^2) \rightarrow \text{Homeo}_0(\mathbb{T}^2)$ is a covering, there exists a (unique) continuous map $\eta : G_{x_0} \rightarrow \text{Homeo}_{\mathbb{Z}^2}(\mathbb{R}^2)$ which lifts the map $\varphi|_{G_{x_0}}$ and sends the identity to the identity. Then the map

$$\begin{aligned} G_{x_0} \times G_{x_0} &\rightarrow \text{Homeo}_{\mathbb{Z}^2}(\mathbb{R}^2) \\ (f, g) &\rightarrow \eta(fg)^{-1}\eta(f)\eta(g) \end{aligned}$$

is continuous and its image is contained in the discrete space of integral translations: it is constant and the map η is a group morphism. Two group morphisms which lift the group morphism $\varphi|_{G_{x_0}}$ differ by a group morphism $G_{x_0} \rightarrow \mathbb{Z}^2$. However, as the group G_{x_0} is simple (hence perfect), such a group morphism is trivial and $\eta = \widetilde{\varphi}_{x_0}$. \square

We can now end the proof of theorem 5.1. Let us denote by F the closure of the set

$$\bigcup_{f \in G_{x_0}} \widetilde{\varphi}_{x_0}(f)((-\infty, 0] \times \mathbb{R}).$$

By the two above lemmas, there exists $M > 0$ such that $F \subset (-\infty, M] \times \mathbb{R}$. Denote by U the connected component of the complementary of F which contains the open subset $(M, +\infty) \times \mathbb{R}$. Denote by U' the image of U under the projection $p_2 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}/\mathbb{Z}$ and by ψ_{x_0} the action on the annulus $\mathbb{R} \times \mathbb{R}/\mathbb{Z}$ defined by

$$\forall f \in \text{Homeo}_0(\mathbb{S}^1), \psi_{x_0}(f) \circ p_2 = p_2 \circ \widetilde{\varphi}_{x_0}(f).$$

Notice that $p_1 \circ \psi_{x_0} = \varphi|_{G_{x_0}} \circ p_1$, where $p_1 : \mathbb{R} \times \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z} = \mathbb{T}^2$.

By construction, the open set U' is invariant under the action ψ_{x_0} (a fundamental domain far on the right must be sent in U' by any homeomorphism in the image of ψ_{x_0} , by the two above lemmas). Now, this action can be extended to the set of prime ends of U' , giving a continuous action ψ of the group G_{x_0} (which is isomorphic to the group $\text{Homeo}_c(\mathbb{R})$) on the topological space of prime ends of U' , which is homeomorphic to \mathbb{S}^1 .

By proposition 3.10, this last action admits a fixed point. Moreover, for any closed interval I whose interior contains the point x_0 , the set of fixed points of this action contains an open interval and hence an accessible prime end. Therefore, the intersection of the set F_I of fixed points of the action $\psi_{x_0}|_{G_I}$ with $[0, M] \times \mathbb{R}/\mathbb{Z}$ is non-empty. Therefore, the set of fixed points of ψ_{x_0} , which is the intersection of the F_I 's, is non-empty. Theorem 5.1 is proved. \square

References

- [1] A. Avila, *Distortion elements in $\text{Diff}^\infty(\mathbb{R}/\mathbb{Z})$*
Available on arXiv:08082334
- [2] K. Brown, *Cohomology of groups*. Graduate Texts in Mathematics 87, Springer-Verlag, New-York, 1994.
- [3] C. Bonatti and B. Kolev, *Surface homeomorphisms with zero dimensional singular set*, Topology Appl. (1-3) 90 (1998), 69-95.
- [4] A. Bounemoura *Simplicité des groupes de transformations de surface*, Ensaos Matematicos, vol. 14 (2008), 1-143.
- [5] R.P. Filipkiewicz, *Isomorphisms between diffeomorphism groups*, Ergodic Th. and Dyn. Sys. 2 (1982), no 2, 159-171.
- [6] G.M. Fischer, *On the group of all homeomorphisms of a manifold*, Trans. of the Amer. Math. Soc. 97 (1960), 193-212.
- [7] E. Ghys, *Groups acting on the circle*, L'Enseignement Mathématique, vol.47 (2001), 329-407.
- [8] J.N. Mather, *The vanishing of the homology of certain groups of homeomorphisms*, Topology 10 (1971), 297-298.
- [9] J.N. Mather, *Topological proofs of some purely topological consequences of Caratheodory's theory of prime ends*, Selected studies: physics-astrophysics, mathematics, history of science, North-Holland, Amsterdam-New York (1982), 225-255.
- [10] S. Matsumoto, *Numerical invariants for semiconjugacy of homeomorphisms of the circle*, Proceedings of the American Mathematical Society, vol. 98 (1986), no 1, 163-168.
- [11] S. Matsumoto and S. Morita, *Bounded cohomology of certain groups of homeomorphisms*, Proceedings of the American Mathematical Society, vol. 94 (1985), 539-544.

- [12] E. Milton, *Distortion elements for surface homeomorphisms*, preprint.
Available on arXiv:1207.0291
- [13] J.V. Whittaker, *On isomorphic groups and homeomorphic spaces*, *Annals of Math.* 78 (1963), no 1, 74-91.