



HAL
open science

Dust rocket storms, gravity waves and their impact on the martian troposphere and thermosphere

A. Spiga, F. Gonzalez-Galindo, J. Faure, J.-B. Madeleine, F. Altieri, M.-A. Lopez-Valverde, F. Forget, Anni Määttänen, Franck Montmessin

► **To cite this version:**

A. Spiga, F. Gonzalez-Galindo, J. Faure, J.-B. Madeleine, F. Altieri, et al.. Dust rocket storms, gravity waves and their impact on the martian troposphere and thermosphere. EPSC 2012 (European Planetary Science Congress), Sep 2012, Madrid, Spain. pp.EPSC2012-272. <hal-00749778>

HAL Id: hal-00749778

<https://hal.science/hal-00749778v1>

Submitted on 27 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Dust rocket storms, gravity waves and their impact on the martian troposphere and thermosphere

A. Spiga (1), F. Gonzalez-Galindo (2), J. Faure (1), J.-B. Madeleine (1), F. Altieri (3), M.-A. Lopez-Valverde (2), F. Forget (1), A. Määttänen (4), F. Montmessin (4).
Contact: aymeric.spiga@upmc.fr. Website: <http://www.lmd.jussieu.fr/~aslmd>. (1) LMD, UPMC, Paris, France (2) IAA, CSIC, Granada, Spain (3) IAPS, Roma, Italy (4) LATMOS, CNRS, Guyancourt, France.

Background Recent studies have shed light on mesoscale phenomena in the Martian atmosphere, unresolved by global climate models [6]. A particular emphasis was put on near-surface circulations, less so on dynamical phenomena in the upper troposphere and mesosphere. This aspect is in need to be further investigated to better understand recent observations and, more generally, the martian climate.




Figure 1: Gravity waves, mesospheric cold pockets and CO₂ clouds. Results from idealized mesoscale simulations carried out in [7]. Vertical profiles of temperature in the vicinity of the domain center are shown : dashed line represents initial GCM profile, solid line represents predicted mesoscale profile after 3 simulated hours, orange “envelope” represents a larger set of predicted profiles ranging 2 – 4 simulated hours and 20 × 20 grid points around domain center. CO₂ condensation profile T_c is superimposed.

Gravity waves Many independent measurements have shown that extremely low temperatures (“cold pockets”) are found in the Martian mesosphere. Recent observational achievements also hint at such cold pockets by revealing mesospheric clouds formed through the condensation of CO₂ [e.g., 5]. Large-scale meteorological conditions are key factors to account

for the presence of those cold pockets in the Martian mesosphere [2]. Yet we show through multiscale meteorological modeling (Figure 1) that mesoscale gravity waves too could play a key role in the formation of mesospheric cold pockets propitious to CO₂ condensation [7]. Gravity waves are not detectable only through temperature disturbances. Wave patterns in the southern polar region of Mars were found in O₂ dayglow maps obtained with MEx/OMEGA. We show through mesoscale modeling (Figure 2) that the propagation of gravity waves in the Martian troposphere explains these patterns and that model predictions match spatial variability and dayglow fluctuations observed in the OMEGA maps [1].




Figure 2: Airglow and gravity waves. Estimated fluctuations of integrated airglow intensity in %, derived from predictions with mesoscale model by [6] for southern polar regions around spring equinox. See [1] for further details.

Dust rocket storms We use atmospheric mesoscale modeling with radiatively-active transported dust to predict the evolution of a local dust storm monitored by OMEGA onboard Mars Express [4]. In the afternoon, dust transport within storm is governed by deep convective motions (Figure 3). The vast majority of convective energy supply originates in the absorption of incoming sunlight by dust particles, mostly in the visible. What we propose to name a “dust rocket storm” subsequently forms and injects dust particles at high altitudes in the Martian troposphere (30 to 40 km above the surface). Combined to advection by horizontal winds, this contributes to form detached layers of dust reminiscent of those observed e.g. with instruments onboard Mars Reconnaissance Orbiter [3]. Such detached layers are stable over several days. We checked the robustness of our conclusions by performing sensitivity simulations without the radiative effect of transported dust, with various sizes and properties of dust disturbance and with lifting activated in a more realistic setting. Our conclusions (manuscript to be submitted to JGR planets) have strong implications for the Martian dust cycle and advocate for further studies of dust storms (including global dust storms) at finer scales than usually assumed in global climate models.

References

- [1] F. Altieri, A. Spiga, L. Zasova, G. Bellucci, and J. P. Bibring. Gravity waves mapped by the OMEGA/MEX instrument through O₂ dayglow at 1.27 microns: data analysis and atmospheric modeling. *Journal of Geophysical Research (Planets)*, 117(accepted):0–+, 2012.
- [2] F. González-Galindo, A. Määttänen, F. Forget, and A. Spiga. The martian mesosphere as revealed by co2 clouds observations and general circulation modeling. *Icarus*, 216:10–22, 2011.
- [3] N. G. Heavens, M. I. Richardson, A. Kleinböhl, D. M. Kass, D. J. McCleese, W. Abdou, J. L. Benson, J. T. Schofield, J. H. Shirley, and P. M. Wolkenberg. The vertical distribution of dust in the Martian atmosphere during northern spring and summer: Observations by the Mars Climate Sounder and analysis of zonal average vertical dust profiles. *Journal of Geophysical Research (Planets)*, 116(E15):4003, 2011.
- [4] A. Määttänen, T. Fouchet, O. Forni, R. Melchiorri, F. Forget, H. Savijarvi, J. P. Bibring, Y. Langevin, B. Gondet, V. Formisano, and M. Giuranna. A study of the properties of a local dust storm with Mars Express OMEGA and PFS data. *Icarus*, 201(2):504–516, 2009.




Figure 3: From dust rocket storms to detached layers of dust. *Longitude-altitude sections of “MCS-like” density-scaled dust optical depth predicted with our mesoscale model every two hours from local time 14 : 00 to 00 : 00. Initial state defined according to [4].*

- [5] F. Montmessin, B. Gondet, J.-P. Bibring, Y. Langevin, P. Drossart, F. Forget, and T. Fouchet. Hyperspectral imaging of convective CO₂ ice clouds in the equatorial mesosphere of Mars. *Journal of Geophysical Research (Planets)*, 112:11–+, 2007.
- [6] A. Spiga and F. Forget. A new model to simulate the Martian mesoscale and microscale atmospheric circulation: Validation and first results. *Journal of Geophysical Research (Planets)*, 114:E02009, 2009.
- [7] A. Spiga, F. González-Galindo, M.-Á. López-Valverde, and F. Forget. Gravity waves, cold pockets and CO₂ clouds in the Martian mesosphere. *Geophys. Res. Letters*, 39:2201, 2012.