



**HAL**  
open science

# Local null controllability of the three-dimensional Navier-Stokes system with a distributed control having two vanishing components

Jean-Michel Coron, Pierre Lissy

► **To cite this version:**

Jean-Michel Coron, Pierre Lissy. Local null controllability of the three-dimensional Navier-Stokes system with a distributed control having two vanishing components. 2012. hal-00750249v1

**HAL Id: hal-00750249**

**<https://hal.science/hal-00750249v1>**

Preprint submitted on 9 Nov 2012 (v1), last revised 29 Oct 2013 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Local null controllability of the three-dimensional Navier-Stokes system with a distributed control having two vanishing components

Jean-Michel Coron \*      Pierre Lissy \*

## Abstract

In this paper, we prove a local null controllability result for the three-dimensional Navier-Stokes equations on a (smooth) bounded domain of  $\mathbb{R}^3$  with null Dirichlet boundary conditions. The control is distributed into an (arbitrarily small) open subset and has two vanishing components. J.-L. Lions and E. Zuazua proved that the linearized system is not necessarily approximately controllable even if the control is distributed on the entire domain, hence it cannot be null controllable and the standard linearization method fails. We use the return method together with a new algebraic method inspired by the works of M. Gromov and previous results by M. Gueye.

Keywords: Navier-Stokes System Null controllability Return method.

MSC classification: MSC 35Q30 MSC 93B05 MSC 93C10

## 1 Introduction

### 1.1 Notations and statement of the theorem

Let  $T > 0$ , let  $\Omega$  be a nonempty bounded domain of  $\mathbb{R}^3$  of class  $C^\infty$  and let  $\omega$  be a nonempty open subset of  $\Omega$ . We define  $Q \subset \mathbb{R} \times \mathbb{R}^3$  by

$$Q := (0, T) \times \Omega = \{(t, x) \mid t \in (0, T) \text{ and } x \in \Omega\}.$$

The current point  $x \in \mathbb{R}^3$  is  $x = (x_1, x_2, x_3)$ . The  $i$ -th component of a vector (or a vector field)  $f$  is denoted  $f^i$ . The control is  $u = (u^1, u^2, u^3) \in L^2(Q)^3$ . We require that the support of  $u$  is included in  $\omega$ , which is our control domain. We impose that two components of  $u$  vanish, for example the first two:

$$u^1 = 0 \text{ and } u^2 = 0 \text{ in } Q, \tag{1.1}$$

---

<sup>1</sup>Work supported by the ERC advanced grant 266907 (CPDENL) of the 7th Research Framework Programme (FP7)

so that  $u$  will be written under the form  $(0, 0, 1_\omega v)$  with  $v \in L^2(Q)$  from now on, where  $1_\omega : \Omega \rightarrow \mathbb{R}$  is the characteristic function of  $\omega$ :

$$1_\omega = 1 \text{ in } \omega, \quad 1_\omega = 0 \text{ in } \Omega \setminus \omega.$$

Let us define

$$V := \{y \in H_0^1(\Omega)^3 \mid \nabla \cdot y = 0\}.$$

The space  $V$  is equipped with the  $H^1$ -norm. Let us denote by  $H$  the closure of  $V$  in  $L^2(\Omega)^3$ . The space  $H$  is equipped with the  $L^2$ -norm.

We are interested in the following Navier-Stokes control system:

$$\begin{cases} y_t - \Delta y + (y \cdot \nabla)y + \nabla p = (0, 0, 1_\omega v) & \text{in } Q, \\ \nabla \cdot y = 0 & \text{in } Q, \\ y = 0 & \text{on } [0, T] \times \partial\Omega. \end{cases} \quad (1.2)$$

From [27, Theorem 3.1, p. 282], we have the following existence result: For every  $y^0 \in H$ , there exist  $y \in L^2((0, T), V) \cap L^\infty((0, T), H)$  and  $p \in L^2(Q)$  satisfying

$$y(0, \cdot) = y^0 \text{ in } \Omega \quad (1.3)$$

such that (1.2) holds.

Our main result is the following theorem, which expresses the small-time local null-controllability of (1.2):

**Theorem 1.** *For every  $T > 0$  and for every  $r > 0$ , there exists  $\eta > 0$  such that, for every  $y^0 \in V$  verifying  $\|y^0\|_{H^1(\Omega)^3} \leq \eta$ , there exist  $v \in L^2(Q)$  and a solution  $(y, p) \in L^2((0, T), H^2(\Omega)^3 \cap V) \cap L^\infty((0, T), H^1(\Omega)^3 \cap V) \times L^2(Q)$  of (1.2)-(1.3) such that*

$$y(T, \cdot) = 0, \quad (1.4)$$

$$\|v\|_{L^2(Q)^3} \leq r, \quad (1.5)$$

$$\|y\|_{L^2((0, T), H^2(\Omega)^3) \cap L^\infty((0, T), H^1(\Omega)^3)} \leq r. \quad (1.6)$$

**Remark 1.** *Once a control  $v \in L^2(Q)$  is given, the corresponding solution  $(y, p)$  of (1.2), (1.3) and (1.6) given by Theorem 1 is unique (recall that for the Navier-Stokes system, the uniqueness of  $(y, p)$  means that  $y$  is unique and  $p$  is unique up to a constant depending on the time). This comes from the uniqueness result given in [27, Theorem 3.4, p. 297]: One has*

$$L^\infty((0, T), H^1(\Omega)^3) \subset L^8((0, T), H^1(\Omega)^3) \subset L^8((0, T), L^4(\Omega)^3)$$

thanks to a classical Sobolev embedding, and there is at most one solution  $(y, p)$  of (1.2) and (1.3) in the space

$$L^2((0, T), V) \cap L^\infty((0, T), H) \cap L^8((0, T), L^4(\Omega)^3) \times L^2(Q).$$

**Remark 2.** *One observes that in Theorem 1 the initial condition  $y^0$  is more regular than usual ( $y^0 \in H$ ). In fact, using the same arguments as in [15] and [16] (see also [6, Remark 1]), one can easily extend the previous theorem to small initial data in  $H \cap L^4(\Omega)^3$  with a solution  $(y, p) \in L^2((0, T), V) \cap L^\infty((0, T), H) \times L^2(Q)$ . In this case, Remark 1 is no longer true and there might possibly exist many solutions  $(y, p)$  verifying (1.2) and (1.3) once  $v$  is given.*

## 1.2 Some previous results

The controllability of the two or three-dimensional Navier-Stokes equations with a distributed control has been studied in numerous papers. In general, for Navier-Stokes equations, it is relevant to consider the approximate controllability, the null controllability or the controllability to the trajectories, the second one being a particular case of the third one.

In [21], a first result of local exact controllability to the trajectories was established under technical conditions:  $\Omega$  had to be homeomorphic to a ball, the control had to be supported in a nonempty open subset whose closure is included in  $\Omega$ , and the target trajectory had to be a stationary solution of the Navier-Stokes equation. Moreover, there were some technical regularity conditions for these stationary solutions. A similar result for the linearized Navier-Stokes equations was established but with the same strong conditions. Many of these hypotheses were removed in [22].

Then, it was proved in [15] the local exact controllability to the trajectories with regularity conditions that were weaker and more suitable for the study of the Navier-Stokes equations. In this article, the authors also proved some exact controllability results for linearized Navier-Stokes systems, with very weak regularity conditions. The same authors proved in [16] the local exact controllability with a control having one vanishing component, provided that  $\omega$  “touches” the boundary of the domain  $\Omega$  in some sense. Later on it was proved in [13] a local null controllability result on the Stokes system with a control having a vanishing component without the geometrical condition on  $\omega$ , but the authors were not able to extend it to the nonlinear Navier-Stokes system. A recent work ([6]) improved the previous one and proved the local null controllability of the Stokes system with an additional source member by means of a control having a vanishing component, which enabled the authors to prove the local null controllability of the Navier-Stokes system for a control having a vanishing component. In all these articles, the main points of the proof were to establish first the controllability of the linearized control system around the target trajectory thanks to Carleman estimates on the adjoint of the linearized equation, and then to use an inverse mapping theorem or a fixed-point theorem to deal with the nonlinear system.

The natural question is then: Can we remove another component of the control, which would provide an optimal result with respect to the number of controls? Reducing the number of components of the control is important for applications, and have already been studied many times for linear or parabolic systems of second order (that are quite similar to linearized Navier-Stokes systems), see for example [4], where a necessary and sufficient condition to control a system of coupled parabolic equations with constant coefficients and with less controls than equations is given, or [3, 25] for time-dependant coefficients. If the coefficients depend on the time and the space, there are no general results, notably if we consider two coupled parabolic systems where the coupling region and the control region do not intersect (a partial result under GCC is given in [1]). For a recent survey on the controllability of coupled linear parabolic equations, see [5].

### 1.3 The linear test

To obtain Theorem 1, the first natural idea is to linearize the system around 0, i.e. to consider the Stokes control system

$$\begin{cases} y_t - \Delta y + \nabla p = (0, 0, v1_\omega) & \text{in } Q, \\ \nabla \cdot y = 0 & \text{in } Q, \\ y(0, \cdot) = y^0 & \text{in } \Omega, \\ y = 0 & \text{on } [0, T] \times \partial\Omega. \end{cases} \quad (1.7)$$

It is well known (see for example [16], or [22]) that if this linear system were null controllable (with, in addition, an arbitrary source term in a suitable space), then applying an inverse mapping theorem (for example the one presented in [2]) in some relevant weighted spaces, we would obtain that (1.2) is locally null controllable around 0. However the linear control system (1.7) is in general not null controllable, and not even approximately controllable: In [24], it is proved that if we choose  $\Omega$  to be a cylinder with a circular generating set and with an axis parallel to  $e_3$  then (1.7) is not approximately controllable, even if we control on the entire cylinder. However, it is proved in [24] that the approximate controllability property holds generically with respect to the generating set of the cylinder.

Since linearizing around 0 is not relevant, we are going to use the return method, which consists in linearizing system (1.2) around a particular trajectory  $(\bar{y}, \bar{p}, \bar{u})$  (that we construct explicitly), verifying  $\bar{y}(0, \cdot) = \bar{y}(T, \cdot) = 0$ , proving that the linearized system (with a source term  $f$  verifying an exponential decrease condition at time  $t = T$ ) is null controllable, and then concluding by a usual inverse mapping argument that our system is locally null controllable. This method was introduced in [7] for a stabilization problem and first used in [8] for controllability issues for partial differential equations. The return method was already successfully used in [9, 11, 18] to obtain global controllability results for the Navier-Stokes equations and in [12] to prove the small-time local null controllability for the Navier-Stokes equations on the torus  $\mathbb{T}_2$  when the control has one vanishing component. For more explanations about the return method and other examples, see [10, Chapter 6].

## 2 Constructing a relevant trajectory

In this subsection, we construct explicit particular trajectories  $(\bar{y}, \bar{p}, \bar{u})$  going from 0 to 0 so that, as it will be shown in section 3, the linearized control system around them is null controllable.

Without loss of generality we may assume that  $0 \in \omega$ . Let  $g \in C^\infty(\mathbb{R}^3)$ ,

$$g : (t, w, x_3) \mapsto g(t, w, x_3),$$

and  $h \in C^\infty(\mathbb{R}^3)$ ,

$$h : (t, w, x_3) \mapsto h(t, w, x_3).$$

For  $(x_1, x_2, x_3) \in \mathbb{R}^3$ , let  $r := \sqrt{x_1^2 + x_2^2}$ . We define  $\bar{y} \in C^\infty(\mathbb{R}^4; \mathbb{R}^3)$  by

$$\bar{y}(t, x) := \begin{pmatrix} g(t, r^2, x_3)x_1 \\ g(t, r^2, x_3)x_2 \\ h(t, r^2, x_3) \end{pmatrix}, \forall t \in \mathbb{R}, \forall x = (x_1, x_2, x_3) \in \mathbb{R}^3. \quad (2.1)$$

Let  $r_1 > 0$  be small enough so that

$$\mathcal{C}_1 := \{(x_1, x_2, x_3) \in \mathbb{R}^3; r \leq r_1, |x_3| \leq r_1\} \subset \omega. \quad (2.2)$$

On the functions  $g$  and  $h$ , we also require that

$$\text{Supp}(g) \subset [T/4, T] \times (-\infty, r_1^2] \times [-r_1, r_1], \quad (2.3)$$

$$\text{Supp}(h) \subset [T/4, T] \times (-\infty, r_1^2] \times [-r_1, r_1]. \quad (2.4)$$

In (2.3), (2.4) and in the following,  $\text{Supp}(f)$  denotes the support of the function  $f$ . From (2.1), (2.2), (2.3) and (2.4), one obtains

$$\text{Supp}(\bar{y}) \subset [T/4, T] \times \mathcal{C}_1 \subset (0, T] \times \omega \subset (0, T] \times \Omega. \quad (2.5)$$

Let  $\tilde{p} \in C^\infty(\mathbb{R}^3)$  be defined by

$$\begin{aligned} \tilde{p}(t, w, x_3) := & \frac{1}{2} \int_{r_1^2}^\varepsilon (g_t - (4wg_{ww} + 8g_w + g_{33}) \\ & + 2wgg_w + |g|^2 + hg_3)(t, w', x_3) dw'. \end{aligned} \quad (2.6)$$

In (2.6), in order to lighten the notations, for  $l \in \mathbb{N}^*$ ,  $N \in \mathbb{N}^*$  and  $i = (i_1, \dots, i_N)$ , with  $i_k \in \{t, w, 3\}$  for every  $k \in \{1, \dots, N\}$ , and  $\varphi : D(\varphi) \subset \mathbb{R}^3 \rightarrow \mathbb{R}^l$ ,  $(t, w, x_3) \mapsto \varphi(t, w, x_3)$ ,  $\varphi_{i_1 i_2 \dots i_N}$  denotes the following derivative:

$$\varphi_{i_1 i_2 \dots i_N} := \frac{\partial^N \varphi}{\partial \xi_{i_1} \dots \partial \xi_{i_N}},$$

where

$$\xi_j = x_3 \text{ if } j = 3, \xi_j = t \text{ if } j = t \text{ and } \xi_j = w \text{ if } j = w.$$

We use similar notations throughout the paper. Let  $\bar{p} \in C^\infty(\mathbb{R}^4)$  be defined by

$$\bar{p}(t, x_1, x_2, x_3) := \tilde{p}(t, r^2, x_3). \quad (2.7)$$

From (2.2), (2.3), (2.4), (2.6) and (2.7), it follows that

$$\text{Supp}(\bar{p}) \subset [T/4, T] \times \mathcal{C}_1 \subset [T/4, T] \times \omega \subset (0, T] \times \Omega. \quad (2.8)$$

From (2.1), (2.6) and (2.7), one obtains

$$\bar{y}_t^1 - \Delta \bar{y}^1 + (\bar{y} \cdot \nabla) \bar{y}^1 + \bar{p}_1 = 0, \quad (2.9)$$

$$\bar{y}_t^2 - \Delta \bar{y}^2 + (\bar{y} \cdot \nabla) \bar{y}^2 + \bar{p}_2 = 0. \quad (2.10)$$

Let  $\bar{u} \in C^\infty(\mathbb{R}^4)^3$  be defined by

$$\bar{u} := (0, 0, \bar{y}_t^3 - \Delta \bar{y}^3 + (\bar{y} \cdot \nabla) \bar{y}^3 + \bar{p}_3). \quad (2.11)$$

From (2.11), one obtains (1.1). From (2.5), (2.8) and (2.11), we have

$$\text{Supp}(\bar{u}) \subset (0, T] \times \omega. \quad (2.12)$$

From (2.9), (2.10), (2.11) and (2.12), we have

$$\bar{y}_t - \Delta \bar{y} + (\bar{y} \cdot \nabla) \bar{y} + \nabla \bar{p} = 1_\omega \bar{u}. \quad (2.13)$$

Finally, in order to have

$$\text{div } \bar{y} = 0, \quad (2.14)$$

it suffices to impose

$$\frac{\partial h}{\partial x_3} = -2(g + wg_w). \quad (2.15)$$

Let  $\nu$  be a positive numerical constant which will be chosen later on. Let  $a \in C^\infty(\mathbb{R})$ ,  $b \in C^\infty(\mathbb{R})$  and  $c \in C^\infty(\mathbb{R})$  be such that

$$\text{Supp}(a) \subset [T/4, T] \text{ and } a(t) = e^{\frac{-\nu}{(T-t)^5}} \text{ in } [T/2, T], \quad (2.16)$$

$$\text{Supp}(b) \subset (-\infty, r_1^2) \text{ and } b(w) = w, \forall s \in (-\infty, r_1^2/4], \quad (2.17)$$

$$\text{Supp}(c) \subset (-r_1, r_1) \text{ and } c(x_3) = x_3^2 \text{ in } [-r_1/2, r_1/2]. \quad (2.18)$$

We choose

$$g(t, w, x_3) = \varepsilon a(t) b(w) c'(x_3) \quad (2.19)$$

and

$$h(t, w, x_3) = -2\varepsilon a(t) (b(w) + wb'(w)) c(x_3), \quad (2.20)$$

where  $\varepsilon > 0$  will be chosen later on. From (2.19) and (2.20), one obtains (2.15).

In the next section, we prove that, for every small enough  $T$ , for every small enough  $\varepsilon > 0$  and for a well-chosen  $\nu$ , the linearized control system around the trajectory  $(\bar{y}, \bar{p}, \bar{u})$  is controllable.

### 3 A controllability result on the linearized system

#### 3.1 Definitions and notations

The linearized control system around the trajectory  $(\bar{y}, \bar{p}, \bar{u})$  is the linear control system

$$\begin{cases} y_t^1 - \Delta y^1 + (\bar{y} \cdot \nabla) y^1 + (y \cdot \nabla) \bar{y}^1 + p_1 = f^1 & \text{in } Q, \\ y_t^2 - \Delta y^2 + (\bar{y} \cdot \nabla) y^2 + (y \cdot \nabla) \bar{y}^2 + p_2 = f^2 & \text{in } Q, \\ y_t^3 - \Delta y^3 + (\bar{y} \cdot \nabla) y^3 + (y \cdot \nabla) \bar{y}^3 + p_3 = 1_\omega v + f^3 & \text{in } Q, \\ \nabla \cdot y = 0 & \text{in } Q, \\ y = 0 & \text{on } [0, T] \times \partial\Omega, \end{cases} \quad (3.1)$$

where the state is  $y : Q \rightarrow \mathbb{R}^3$ ,  $f : Q \rightarrow \mathbb{R}^3$  is a source term (it will be specified later in which space exactly it shall be) and the control is  $v : Q \rightarrow \mathbb{R}$ . Let  $\omega_0$  be a nonempty open subset of

$$\mathcal{C}_2 := \left\{ (x_1, x_2, x_3); r < \frac{r_1}{2}, |x_3| < \frac{r_1}{2} \right\}, \quad (3.2)$$

which will be chosen more precisely in the next section. Let

$$Q_0 := (T/2, T) \times \omega_0.$$

The following figure represents the open subsets of  $\Omega$  we introduced up to now.

Let  $\mathcal{L} : C^\infty(Q_0)^5 \rightarrow C^\infty(Q_0)^4$  be defined by

$$\mathcal{L} \begin{pmatrix} y \\ p \\ v \end{pmatrix} := \begin{pmatrix} y_t^1 - \Delta y^1 + (\bar{y} \cdot \nabla) y^1 + (y \cdot \nabla) \bar{y}^1 + p_1 \\ y_t^2 - \Delta y^2 + (\bar{y} \cdot \nabla) y^2 + (y \cdot \nabla) \bar{y}^2 + p_2 \\ y_t^3 - \Delta y^3 + (\bar{y} \cdot \nabla) y^3 + (y \cdot \nabla) \bar{y}^3 + p_3 - v \\ \nabla \cdot y \end{pmatrix}, \quad (3.3)$$

for every  $y = (y^1, y^2, y^3) \in C^\infty(Q_0)^3$ , for every  $p \in C^\infty(Q_0)$  and for every  $v \in C^\infty(Q_0)$ . Let us denote by

$$\xi := (x_0, x_1, x_2, x_3) = (t, x_1, x_2, x_3)$$

the current point in  $Q_0$ . For  $\alpha = (\alpha_0, \alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^4$  and  $\varphi : Q_0 \rightarrow \mathbb{R}^k$ ,  $\partial^\alpha \varphi$ , denotes, as usual,

$$\partial_{t^{\alpha_0}}^{\alpha_0} \partial_{x_1^{\alpha_1}}^{\alpha_1} \partial_{x_2^{\alpha_2}}^{\alpha_2} \partial_{x_3^{\alpha_3}}^{\alpha_3} \varphi$$

(in this part, this notation will be more convenient than the one introduced in the previous section, which is more compact only for derivatives of small order). Let  $\mathcal{L}(\mathbb{R}^k; \mathbb{R}^l)$  be the set of linear maps from  $\mathbb{R}^k$  into  $\mathbb{R}^l$  and  $\mathcal{M}_{k,l}(\mathcal{R})$  be the set of matrices of size  $k \times l$  with value in the ring  $\mathcal{R}$ .

As usual, in the inequalities written in this article  $C$  denotes a constant (depending in general only on  $\omega$ ,  $\Omega$ ,  $T$ ) that may change from one line to another.

Let us give some other definitions.






Figure 1: The open subsets  $\mathcal{C}_1, \mathcal{C}_2, \omega_0, \omega$ .

**Definition 1.** A linear map  $\mathcal{M} : C^\infty(Q_0)^k \rightarrow C^\infty(Q_0)^l$  is called a linear partial differential operator of order  $m$  if, for every  $\alpha = (\alpha_0, \alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^4$  with  $|\alpha| := \alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 \leq m$ , there exists  $A_\alpha \in C^\infty(Q_0; \mathcal{L}(\mathbb{R}^k; \mathbb{R}^l))$  such that

$$(\mathcal{M}\varphi)(\xi) = \sum_{|\alpha| \leq m} A_\alpha(\xi) \partial^\alpha \varphi(\xi), \quad \forall \xi \in Q_0, \quad \forall \varphi \in C^\infty(Q_0)^k.$$

A linear map  $\mathcal{M} : C^\infty(Q_0)^k \rightarrow C^\infty(Q_0)^l$  is called a linear partial differential operator if there exists  $m \in \mathbb{N}$  such that  $\mathcal{M}$  is a linear partial differential operator of order  $m$ .

Let  $k$  be a positive integer and let  $\mathcal{B} := (\mathcal{B}^1, \mathcal{B}^2, \mathcal{B}^3) : C^\infty(Q_0)^k \rightarrow C^\infty(Q_0)^3$  be a linear partial differential operator. Let us consider the linear equation

$$\mathcal{L} \begin{pmatrix} y \\ p \\ v \end{pmatrix} = \begin{pmatrix} \mathcal{B}^1 f \\ \mathcal{B}^2 f \\ \mathcal{B}^3 f \\ 0 \end{pmatrix}, \quad (3.4)$$

where the data is  $f \in C^\infty(Q_0)^k$  and the unknown is  $(y, p, v) \in C^\infty(Q_0)^5$ . Following [19, Section 2.3.8, p. 148], we adopt the following definition.

**Definition 2.** *The linear equation (3.4) is algebraically solvable if there exists a linear partial differential operator  $\mathcal{M} : C^\infty(Q_0)^k \rightarrow C^\infty(Q_0)^5$  such that, for every  $f \in C^\infty(Q_0)^k$ ,  $\mathcal{M}f$  is a solution of (3.4), i.e. such that*

$$\mathcal{L} \circ \mathcal{M} = (\mathcal{B}, 0). \quad (3.5)$$

In the following, every function  $\varphi \in C^\infty(Q_0)^l$  with a compact support included in  $Q_0$  is extended by 0 in  $Q \setminus Q_0$  and we still denote this extension by  $\varphi$ .

The next proposition explains how the notion of algebraic solvability can be useful to reduce the number of controls as soon as a controllability result is already known for a large number of controls. In fact, the question of the null-controllability of (3.1) can be split into two distinct problems: One “algebraic” part (solving system(3.5)) and one “analytic” part (finding controls which are in the image of  $\mathcal{B}$ , the control acting possibly on all the equations and not only the third one). This proposition has a very general scope and could be formulated for more general control systems. It is inspired by techniques used in the control of ordinary differential equations (see, in particular, [10, Chapter 1, p. 13-15]).

**Proposition 1.** *Let us consider the linear control system*

$$\left\{ \begin{array}{ll} y_t^1 - \Delta y^1 + (\bar{y} \cdot \nabla) y^1 + (y \cdot \nabla) \bar{y}^1 + p_1 = \mathcal{B}^1 u + f^1 & \text{in } Q, \\ y_t^2 - \Delta y^2 + (\bar{y} \cdot \nabla) y^2 + (y \cdot \nabla) \bar{y}^2 + p_2 = \mathcal{B}^2 u + f^2 & \text{in } Q, \\ y_t^3 - \Delta y^3 + (\bar{y} \cdot \nabla) y^3 + (y \cdot \nabla) \bar{y}^3 + p_3 = \mathcal{B}^3 u + f^3 & \text{in } Q, \\ \nabla \cdot y = 0 & \text{in } Q, \\ y = 0 & \text{on } [0, T] \times \partial\Omega, \\ y(0, \cdot) = y^0 & \text{in } \Omega, \end{array} \right. \quad (3.6)$$

where the state is  $y : Q \rightarrow \mathbb{R}^3$ , the control is  $u \in C^\infty(Q)^k$ , which is required to have a support in  $Q_0$ , and  $f := (f^1, f^2, f^3) \in C^\infty(Q)$  is a source term. Let us assume that:

$\mathcal{A}_1$ . *The linear control system (3.6) is null controllable during the interval of time  $[0, T]$  in the sense that for every  $y^0 \in V$  and for every  $f \in C^\infty(Q)$  such that*

$$\text{there exists } \delta > 0 \text{ such that } f = 0 \text{ on } [T - \delta, T] \times \Omega, \quad (3.7)$$

*there exists  $u \in C^\infty(Q)^k$  with a compact support included in  $Q_0$  such that the solution  $(y, p)$  of (3.6) with initial condition  $y(0, \cdot) = y^0$  satisfies  $y(T, \cdot) = 0$ .*

$\mathcal{A}_2$ .

(3.4) is algebraically solvable.

Then, the linear control system (3.1) is null controllable during the interval of time  $[0, T]$ : For every  $y^0 \in V$  and for every  $f \in C^\infty(Q)$  satisfying (3.7), there exists  $v \in C^\infty(Q)$  with a compact support included in  $Q_0$  such that the the solution  $(y, p)$  of (3.1) satisfying the initial condition  $y(0, \cdot) = y^0$  satisfies  $y(T, \cdot) = 0$ .

**Proof of Proposition 1.** First of all, we use the null-controllability of (3.6) with controls in the image of  $\mathcal{B}$  (Assumption  $\mathcal{A}_1$ ): let  $y^0 \in V$  and let  $u^* \in C^\infty(Q)^k$  with support included in  $Q_0$  such that the solution  $(y^*, p^*)$  of the Cauchy problem (3.6) for the initial condition  $y^*(0, \cdot) = y^0$  satisfies  $y^*(T, \cdot) = 0$ . Let us remark that  $\mathcal{B}$  is a local operator, which implies that  $\mathcal{B}u^*_{|Q_0}$  still has a compact support in  $Q_0$ . Now we use the algebraic solvability of (3.4) (Assumption  $\mathcal{A}_2$ ): Let  $\mathcal{M}$  be as in Definition 2. For a map  $h \in C^\infty(Q)^k$  with a support included in  $Q_0$ , we denote by  $\mathcal{M}$  the map from  $Q$  into  $\mathbb{R}^5$  defined by

$$\mathcal{M}h = 0 \text{ in } Q \setminus Q_0, \mathcal{M}h = \mathcal{M}(h|_{Q_0}) \text{ in } Q_0.$$

We shall use this slight abuse of notation until the end of the paper. Note that, for every  $h \in C^\infty(Q)^k$  with a support included in  $Q_0$ ,  $\mathcal{M}h \in C^\infty(Q)^5$  and has a support included in  $Q_0$  (because  $\mathcal{M}$  is a local operator). Let us call

$$(\tilde{y}, \tilde{p}, v) := -\mathcal{M}u^*.$$

One observes that the support of  $(\tilde{y}, \tilde{p}, v)$  is still included in  $Q_0$ . In particular  $\tilde{y}(T, \cdot) = 0$ . Let

$$(y, p) := (y^* + \tilde{y}, p^* + \tilde{p}).$$

Then

$$\left\{ \begin{array}{ll} y_t^1 - \Delta y^1 + (\bar{y} \cdot \nabla) y^1 + (y \cdot \nabla) \bar{y}^1 + p_1 = f^1 & \text{in } Q, \\ y_t^2 - \Delta y^2 + (\bar{y} \cdot \nabla) y^2 + (y \cdot \nabla) \bar{y}^2 + p_2 = f^2 & \text{in } Q, \\ y_t^3 - \Delta y^3 + (\bar{y} \cdot \nabla) y^3 + (y \cdot \nabla) \bar{y}^3 + p_3 = 1_\omega v + f^3 & \text{in } Q, \\ \nabla \cdot y = 0 & \text{in } Q, \\ y = 0 & \text{on } [0, T] \times \partial\Omega, \\ y(0, \cdot) = y^0 & \text{in } \Omega, \\ y(T, \cdot) = 0 & \text{in } \Omega. \end{array} \right.$$

This shows that the linear control system (3.1) is indeed null controllable during the interval of time  $[0, T]$  and concludes the proof of Proposition 1.  $\blacksquare$

**Remark 3.** For the sake of simplicity, we have formulated Proposition 1 in a  $C^\infty$  setting. Let us assume that the control  $u$  coming from Assumption  $\mathcal{A}_1$  is not of class  $C^\infty$ , but is less regular (one observes that the regularities of  $y^*$ ,  $p^*$  and  $f$  do not matter for the proof of Proposition 1 since only  $u^*$  is differentiated by the linear partial differential operator  $\mathcal{M}$ ). For example, assume that  $u^* \in H_1$  where  $H_1$  is a functional space (for example a weighted Sobolev space), and assume

that  $\mathcal{M}$  can be extended on  $H_1$ ,  $\mathcal{M}u^*$  being then in another functional space  $H_2$  (for example another weighted Sobolev space of order less than  $H_1$  in order to take into account that  $\mathcal{M}$  is a linear partial differential operator). Then one easily verifies that Proposition 1 remains true as soon as every function of  $H_1$  (and its derivatives until the order at least the order of  $\mathcal{M}$ ) vanishes at time  $t = T$ , the first assumption  $\mathcal{A}_1$  being changed as the following: the linear control system (3.6) is null controllable during the interval of time  $[0, T]$ , i.e. for every  $y^0 \in V$  and for every  $f \in L^2(Q)$  satisfying (3.7) there exists  $u \in H_1$  with support included in  $Q_0$  such that the solution  $(y, p)$  of (3.6) satisfying the initial condition  $y(0, \cdot) = y^0$  satisfies  $y(T, \cdot) = 0$ . Similarly, we will need to relax property (3.7) by replacing it with a suitable decay rate near  $t = T$ . This will be detailed in Subsection 3.4.

It remains to deal, for a suitable choice of  $\mathcal{B}$ , with Assumption  $\mathcal{A}_2$  (we shall do it in Subsection 3.2) and with Assumption  $\mathcal{A}_1$  (we shall do it in Subsection 3.3).

### 3.2 Algebraic solvability of (3.4)

We choose  $k = 7$  and define  $\mathcal{B}$  by

$$\mathcal{B}(f^1, f^2, f^3, f^4, f^5, f^6, f^7) := \begin{pmatrix} f_1^1 + f_2^2 + f_3^3 \\ f_1^4 + f_2^5 + f_3^6 \\ f^7 \end{pmatrix}. \quad (3.8)$$

The main result of this subsection is the following proposition.

**Proposition 2.** *There exists  $\varepsilon^* > 0$ , there exists  $T^* > 0$  such that, for every  $\varepsilon \in (0, \varepsilon^*)$ , there exists a nonempty open subset  $\omega_0$  of  $\mathcal{C}_2$  such that Assumption  $\mathcal{A}_2$  holds for every  $T < T^*$ : There exists a linear partial differential operator  $\mathcal{M} : C^\infty(Q_0)^7 \rightarrow C^\infty(Q_0)^5$  such that (3.5) holds.*

#### 3.2.1 The adjoint problem

Let  $\mathcal{L}_0 : C^\infty(Q_0)^4 \rightarrow C^\infty(Q_0)^3$  be the linear partial differential operator defined by

$$\mathcal{L}_0 \begin{pmatrix} y \\ p \end{pmatrix} := \begin{pmatrix} y_t^1 - \Delta y^1 + (\bar{y} \cdot \nabla) y^1 + (y \cdot \nabla) \bar{y}^1 + p_1 \\ y_t^2 - \Delta y^2 + (\bar{y} \cdot \nabla) y^2 + (y \cdot \nabla) \bar{y}^2 + p_2 \\ \nabla \cdot y \end{pmatrix}, \quad (3.9)$$

for every  $y = (y^1, y^2, y^3) \in C^\infty(Q_0)^3$ , and every  $p \in C^\infty(Q_0)$ .

Let  $\mathcal{B}_0 : C^\infty(Q_0)^6 \rightarrow C^\infty(Q_0)^3$  be the linear partial differential operator defined by

$$\mathcal{B}_0(f^1, f^2, f^3, f^4, f^5, f^6) := \begin{pmatrix} f_1^1 + f_2^2 + f_3^3 \\ f_1^4 + f_2^5 + f_3^6 \\ 0 \end{pmatrix}. \quad (3.10)$$

Note that the third equation of (3.4) can be read as

$$v = y_t^3 - \Delta y^3 + (\bar{y} \cdot \nabla) y^3 + (y \cdot \nabla) \bar{y}^3 + p_3 - f^7.$$

Hence, one easily sees that Assumption  $\mathcal{A}_2$  is equivalent to the existence of a linear partial differential operator  $\mathcal{M}_0 : C^\infty(Q_0)^6 \rightarrow C^\infty(Q_0)^4$  such that

$$\mathcal{L}_0 \circ \mathcal{M}_0 = \mathcal{B}_0. \quad (3.11)$$

As in [19, p. 157], we study (3.11) by looking at the ‘‘adjoint equation’’. For every linear partial differential operator  $\mathcal{M} : C^\infty(Q_0)^k \rightarrow C^\infty(Q_0)^l$ ,  $\mathcal{M} = \sum_{|\alpha| \leq m} A_\alpha \partial^\alpha$ , we associate its (formal) adjoint

$$\mathcal{M}^* : C^\infty(Q_0)^l \rightarrow C^\infty(Q_0)^k$$

defined by

$$\mathcal{M}^* \psi := \sum_{|\alpha| \leq m} (-1)^{|\alpha|} \partial^\alpha (A_\alpha^{\text{tr}} \psi), \quad \forall \psi \in C^\infty(Q_0)^l, \quad (3.12)$$

where  $A_\alpha^{\text{tr}}(\xi)$  is the transpose of the matrix  $A_\alpha(\xi)$ . (Definition (3.12) makes sense since  $\sum_{|\alpha| \leq m} A_\alpha \partial^\alpha = 0$  implies that the  $A_\alpha$  are all equal to 0.) One has  $\mathcal{M}^{**} = \mathcal{M}$  and, if  $\mathcal{M} : C^\infty(Q_0)^k \rightarrow C^\infty(Q_0)^l$  and  $\mathcal{N} : C^\infty(Q_0)^l \rightarrow C^\infty(Q_0)^m$  are two linear partial differential operators, then  $(\mathcal{N} \circ \mathcal{M})^* = \mathcal{M}^* \circ \mathcal{N}^*$ .

Hence, (3.11) is equivalent to

$$\mathcal{M}_0^* \circ \mathcal{L}_0^* = \mathcal{B}_0^*. \quad (3.13)$$

Direct computations, together with (2.14), show that, for every  $z = (z^1, z^2) \in C^\infty(Q_0)^2$  and for every  $\pi \in C^\infty(Q_0)$ ,

$$\mathcal{L}_0^* \begin{pmatrix} z \\ \pi \end{pmatrix} = \begin{pmatrix} -z_t^1 - \Delta z^1 - (\bar{y} \cdot \nabla) z^1 + \bar{y}_1^1 z^1 + \bar{y}_1^2 z^2 - \pi_1 \\ -z_t^2 - \Delta z^2 - (\bar{y} \cdot \nabla) z^2 + \bar{y}_2^1 z^1 + \bar{y}_2^2 z^2 - \pi_2 \\ \bar{y}_3^1 z^1 + \bar{y}_3^2 z^2 - \pi_3 \\ -z_1^1 - z_2^2 \end{pmatrix}, \quad (3.14)$$

$$\mathcal{B}_0^* \begin{pmatrix} z \\ \pi \end{pmatrix} = (-z_1^1, -z_2^1, -z_3^1, -z_1^2, -z_2^2, -z_3^2). \quad (3.15)$$

Assumption  $\mathcal{A}_2$  is now equivalent to the following property: There exists a linear partial differential operator  $\mathcal{N} (= \mathcal{M}_0^*) : C^\infty(Q_0)^4 \rightarrow C^\infty(Q_0)^6$  such that for every  $\varphi = (\varphi^1, \varphi^2, \varphi^3, \varphi^4) \in C^\infty(Q_0)^4$ , if  $(z^1, z^2, \pi) \in C^\infty(Q_0)^3$  is a solution of

$$\begin{cases} -z_t^1 - \Delta z^1 - (\bar{y} \cdot \nabla) z^1 + \bar{y}_1^1 z^1 + \bar{y}_1^2 z^2 - \pi_1 = \varphi^1, \\ -z_t^2 - \Delta z^2 - (\bar{y} \cdot \nabla) z^2 + \bar{y}_2^1 z^1 + \bar{y}_2^2 z^2 - \pi_2 = \varphi^2, \\ \bar{y}_3^1 z^1 + \bar{y}_3^2 z^2 - \pi_3 = \varphi^3, \\ -z_1^1 - z_2^2 = \varphi^4, \end{cases} \quad (3.16)$$

then  $(z_1^1, z_2^1, z_3^1, z_1^2, z_2^2, z_3^2) = \mathcal{N} \varphi$ .

**Remark 4.** The most natural linear partial differential operator  $\mathcal{B}$  to try first would have been  $\mathcal{B} : C^\infty(Q_0)^3 \rightarrow C^\infty(Q_0)^3$  defined by

$$\mathcal{B}f := \begin{pmatrix} f^1 \\ f^2 \\ f^3 \end{pmatrix}, \forall f = (f^1, f^2, f^3) \in C^\infty(Q_0)^3. \quad (3.17)$$

Unfortunately, Proposition 2 does not hold with this  $\mathcal{B}$ . Indeed, in this case  $\mathcal{B}_0^* : C^\infty(Q_0)^3 \rightarrow C^\infty(Q_0)^2$  would be now (compare with (3.15)) such that, for every  $z = (z^1, z^2) \in C^\infty(Q_0)^2$  and for every  $\pi \in C^\infty(Q_0)$ ,

$$\mathcal{B}_0^*(z, \pi) = (z^1, z^2). \quad (3.18)$$

Let  $F_1 \in C^\infty(T/2, T)$  and let  $F_2 \in C^\infty(T/2, T)$ . We define  $z = (z^1, z^2) \in C^\infty(Q_0; \mathbb{R}^2)$  and  $\pi \in C^\infty(Q_0)$  by

$$z^1(t, x) := F_1(t), \quad z^2(t, x) := F_2(t),$$

and

$$\pi(t, x) := -F_1'(t)x_1 - F_2'(t)x_2 + F_1(t)\bar{y}^1 + F_2(t)\bar{y}^2.$$

Then  $\mathcal{L}_0^*(z, \pi) = 0$ . However, if  $(F_1, F_2) \neq (0, 0)$ , then  $\mathcal{B}_0^*(z, \pi) \neq 0$ . Hence, in this case, (3.13) does not hold whatever the linear partial differential operator  $\mathcal{M}_0$  is and whatever the trajectory  $(\bar{y}, \bar{p}, \bar{u})$  is.

### 3.2.2 Number of variables and equations

Let us give some algebraic results about the number of derivatives of a certain order.

**Definition 3.** Consider a scalar PDE with a smooth (enough) variable  $z$  depending on 4 variables  $x_0, x_1, x_2, x_3$ . We call equations of level  $n$  all the different equations we obtain by differentiating the PDE with respect to all the possible multi-integers of length  $n$ . The number of distinct equations of level  $n$  is denoted  $E(n)$ , and the number of distinct equations of a level less than or equal to  $n$  is denoted  $F(n)$ .

**Remark 5.** Clearly,  $E(n)$  is also the distinct number of derivatives of order  $n$  for (smooth enough) functions having 4 variables, and  $F(n)$  is also the distinct number of derivatives of an order less than  $n$  for (smooth enough) functions having 4 variables. Moreover, if we consider a scalar PDE with many variables  $z^1, \dots, z^k$  depending on  $x_0, x_1, x_2, x_3$  containing derivatives of  $z^1, \dots, z^k$  of order  $m$  at most, the maximum number of derivatives of  $z^1, \dots, z^k$  we may expect in the equations of a level less than or equal to  $n$  is  $kF(n+m)$ .

We want to compute  $E$  and  $F$  precisely. One has

$$E(n) = \frac{(n+1)(n+2)(n+3)}{6}, \quad (3.19)$$

$$F(n) = \frac{(n+1)(n+2)(n+3)(n+4)}{24}. \quad (3.20)$$

Indeed

$$(\alpha_0, \alpha_1, \alpha_2, \alpha_3) \mapsto \{\alpha_0 + 1, \alpha_0 + \alpha_1 + 2, \alpha_0 + \alpha_1 + \alpha_2 + 3, \alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 + 4\}$$

defines a bijection between the set of  $(\alpha_0, \alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^4$  such that  $\alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 \leq n$  and the set of subsets of  $\{1, 2, \dots, n+4\}$  having 4 elements. Hence,  $F(n)$  being the number of  $(\alpha_0, \alpha_1, \alpha_2, \alpha_3) \in \mathbb{N}^4$  such that  $\alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 \leq n$ , we have (3.20). In order to obtain (3.19), we just have to notice that

$$\begin{aligned} E(n) &= F(n) - F(n-1) \\ &= \frac{(n+1)(n+2)(n+3)}{6}. \end{aligned}$$

### 3.2.3 A related overdetermined system

Let us now study the equation (3.16), where the data is  $(\varphi^1, \varphi^2, \varphi^3, \varphi^4) \in C^\infty(Q_0)^4$  and the unknown is  $(z^1, z^2, \pi) \in C^\infty(Q_0)^3$ .

Let us explain the idea behind the reasoning we are going to develop in this subsection, which comes from [19, Section 2.3.8, p. 148]. Equation (3.16) is “analytically” overdetermined, since we have more equations (4) than unknowns (3). However, if we see (3.16) as a linear system of algebraic unknowns (the unknowns being  $z^1, z^2, \pi$  and their derivatives) the system is now “algebraically” underdetermined: We have 4 equations and 19 unknowns. But it is easy to obtain as many new equations as we want: It suffices to differentiate (3.16) enough times. Some new “algebraic unknowns” (the derivatives of  $z^1, z^2, \pi$ ) appear, but since the system was analytically overdetermined, one can hope that they are not too many new unknowns appearing. Notably, one can hope that, after differentiating a sufficient number of times, we obtain more equations than algebraic unknowns. We would then deduce Assumption  $\mathcal{A}_2$  by inverting in some sense this well-posed linear system (this will be explained in detail later).

We first eliminate  $\pi$  in our equation (3.16). To reach this goal, in (3.16), we apply  $\partial_3$  to the first and second lines, and use the third line. We obtain the following equations:

$$\left\{ \begin{array}{l} -2\bar{y}_3^1 z_1^1 - \bar{y}_3^2 z_2^1 + (\bar{y}_1^1 - \bar{y}_3^3) z_3^1 - \bar{y}^1 z_{13}^1 - \bar{y}^2 z_{23}^1 - \bar{y}^3 z_{33}^1 - z_{3t}^1 - \Delta z_3^1 \\ \quad - \bar{y}_3^2 z_1^2 + \bar{y}_1^2 z_3^2 = \varphi_3^1 - \varphi_1^3, \\ -\bar{y}_3^1 z_2^1 + \bar{y}_2^1 z_3^1 - \bar{y}_3^1 z_1^2 - \bar{y}^1 z_{13}^2 - 2\bar{y}_3^2 z_2^2 - \bar{y}^2 z_{23}^2 + (\bar{y}_2^2 - \bar{y}_3^3) z_3^2 \\ \quad - \bar{y}^3 z_{33}^2 - z_{3t}^2 - \Delta z_3^2 = \varphi_3^2 - \varphi_2^3, \\ \quad -z_1^1 - z_2^2 = \varphi^4. \end{array} \right. \quad (3.21)$$

The first and second equation of (3.21) contain derivatives of  $z^1$  and  $z^2$  up to order 3 and the third equation derivatives up to order 1. We would like to have the same derivatives of maximal order appearing in the three equations in order to be sure that the derivatives of maximal order appearing in the first and second equation might also appear in the third one. Hence we are going to differentiate the last equation 2 more times than the others. If we count the maximum

number of derivatives of  $z^1$  and  $z^2$  we create by differentiating  $n$  times the first and second equation and  $n + 2$  times the third one, we obtain

$$H(n) = 2F(n + 3) = \frac{(n + 4)(n + 5)(n + 6)(n + 7)}{12} \quad (3.22)$$

different derivatives. The number  $G(n)$  of equations we obtain is then

$$\begin{aligned} G(n) &= 2F(n) + F(n + 2) \\ &= \frac{(3 + n)(4 + n)(34 + 17n + 3n^2)}{24}. \end{aligned} \quad (3.23)$$

From (3.22) and (3.23), one sees that  $G(n) - H(n)$  is increasing with respect to  $n$  and that

$$G(18) - H(18) = -44 < 0 \text{ and } G(19) - H(19) = 460 > 0.$$

Hence, in order to have more equations than unknowns and as few equations as possible, we choose  $n = 19$ . We have  $G(19) = 30360$  equations and  $H(19) = 29900$  unknowns. We can see this system of 30360 partial differential equations as a linear system of the form

$$L_0(t, x)Z = \Phi,$$

where  $L_0 \in C^\infty(Q_0; \mathcal{M}_{30360 \times 29900}(\mathbb{R}))$ ,  $Z \in \mathbb{R}^{29900}$  ( $Z$  contains the derivatives of  $z^1$  and  $z^2$  up to the order 19) and  $\Phi \in \mathbb{R}^{30360}$  ( $\Phi$  contains the derivatives of  $\varphi$  up to the order 19). We order the 29900 lines of  $Z$  so that

$$\begin{cases} Z^1 = z_1^1, \\ Z^2 = z_2^1, \\ Z^3 = z_3^1, \\ Z^4 = z_1^2, \\ Z^5 = z_2^2, \\ Z^6 = z_3^2. \end{cases}$$

Assumption  $\mathcal{A}_2$  can then be written in the following form: Prove the existence of a nonempty open subset  $\omega_0$  of  $\mathcal{C}_2$  and of a map  $N \in C^\infty(Q_0; \mathcal{M}_{6 \times 30360}(\mathbb{R}))$  ( $N$  is the algebraic version of the linear partial differential operator  $\mathcal{N}$  introduced in Subsection 3.2, every linear partial differential operator can be alternatively considered as a matrix acting on the derivatives of the arguments of the operator) such that

$$N(t, x)L_0(t, x)Z = (Z^1, Z^2, Z^3, Z^4, Z^5, Z^6), \forall (t, x) \in Q_0, \forall Z \in \mathbb{R}^{29900}. \quad (3.24)$$

Since the size of the matrix  $L_0(t, x)$  is very large, it is impossible to find some  $N$  verifying System (3.24) by hand and we will have to do computations on a computer. Notably, it would be more convenient to make  $L_0$  be a sparse matrix in order to use relevant tools adapted to



the study of big sparse linear systems. This is the reason for our simple choices for  $a$ ,  $b$  and  $c$  given in (2.16), (2.17) and (2.18) (polynomials of small order do not create too many non zero coefficients in  $L_0$  when they are differentiated). Using (2.1), (2.16), (2.17), (2.18), (2.19) and (2.20), System (3.21) becomes simply, in  $Q_0$ ,

$$\left\{ \begin{array}{l} a(t)(-4x_1^3 - 4x_1x_2^2)\varepsilon z_1^1 + a(t)(-2x_1^2x_2 - 2x_2^3)\varepsilon z_2^1 \\ +a(t)(14x_1^2x_3 + 10x_2^2x_3)\varepsilon z_3^1 + a(t)(-2x_1^2x_2 - 2x_2^3)\varepsilon z_1^2 + 4a(t)x_1x_2x_3\varepsilon z_3^2 \\ +a(t)(-2x_1^3x_3 - 2x_1x_2^2x_3)\varepsilon z_{13}^1 + a(t)(-2x_1^2x_2x_3 - 2x_2^3x_3)\varepsilon z_{23}^1 \\ +a(t)(4x_1^2x_3^2 + 4x_2^2x_3^2)\varepsilon z_{33}^1 - z_{3t}^1 - z_{113}^1 - z_{223}^1 - z_{333}^1 = \varphi_3^1 - \varphi_1^1, \\ a(t)(-2x_1^3 - 2x_1x_2^2)\varepsilon z_1^2 + a(t)4x_1x_2x_3\varepsilon z_3^1 + a(t)(-2x_1^3 - 2x_1x_2^2)\varepsilon z_1^2 \\ +a(t)(-4x_1^2x_2 - 4x_2^3)\varepsilon z_2^2 + a(t)(10x_1^2x_3 + 14x_2^2x_3)\varepsilon z_3^2 \\ +a(t)(-2x_1^3x_3 - 2x_1x_2^2x_3)\varepsilon z_{13}^2 + a(t)(-2x_1^2x_2x_3 - 2x_2^3x_3)\varepsilon z_{23}^2 \\ +a(t)(4x_1^2x_3^2 + 4x_2^2x_3^2)z_{33}^2 - z_{3t}^2 - z_{113}^2 - z_{223}^2 - z_{333}^2 = \varphi_3^2 - \varphi_2^2, \\ -z_1^1 - z_2^2 = \varphi^4. \end{array} \right. \quad (3.25)$$

Let us consider the changing of variables

$$s := \varepsilon a(t)$$

and

$$e := \frac{1}{T-t}.$$

( $e$  appears when we differentiate  $t \mapsto a(t)$  on  $Q_0$ ). Let  $\mathbb{R}[E, S, X]$  be the set of polynomials in the variables  $e$ ,  $s$ ,  $x_1$ ,  $x_2$ ,  $x_3$ , with real coefficients. The  $30360 \times 29900$  entries of  $L_0$  can alternatively be seen as functions depending on  $(t, x_1, x_2, x_3, \varepsilon)$  or as elements of  $\mathbb{R}[E, S, X]$  and, from now on, we consider  $L_0$  as an element of  $\mathcal{M}_{30360 \times 29900}(\mathbb{R}[E, S, X])$ . As we will see after, it turns out that many of the entries of  $L_0$  are the 0 polynomial.

For a positive integer  $k$ , let us denote by  $\mathfrak{S}_k$  the set of permutations of  $\{1, \dots, k\}$ . To each  $\sigma \in \mathfrak{S}_k$ , we associate the matrix  $S_\sigma \in \mathcal{M}_{k,k}(\mathbb{R})$  defined by

$$\begin{aligned} S_{\sigma(i)i} &= 1, \forall i \in \{1, \dots, k\}, \\ S_{ji} &= 0, \forall i \in \{1, \dots, k\}, \forall j \in \{1, \dots, k\} \setminus \{\sigma(i)\}. \end{aligned} \quad (3.26)$$

For two positive integers  $k$  and  $l$ , let us denote by  $0_{k \times l}$  the null matrix of  $\mathcal{M}_{k \times l}(\mathbb{R})$  (which is included in  $\mathcal{M}_{k \times l}(\mathbb{R}[E, S, X])$ ). The following lemma is a key step for the proof of Proposition 2.

**Lemma 1.** *There exist*

$$\begin{aligned} \xi^0 &:= (e^0, s^0, x^0) \in \mathbb{R}^5, \\ \sigma &\in \mathfrak{S}_{29900}, \\ \tilde{\sigma} &\in \mathfrak{S}_{30360}, \end{aligned}$$

$$P \in \mathcal{M}_{7321 \times 7321}(\mathbb{R}[E, S, X]),$$

$$Q \in \mathcal{M}_{23039 \times 7321}(\mathbb{R}[E, S, X])$$

and

$$R \in \mathcal{M}_{23039 \times 22579}(\mathbb{R}[E, S, X])$$

such that

$$\sigma(i) = i, \forall i \in \{1, 2, 3, 4, 5, 6\}, \quad (3.27)$$

$$S_{\bar{\sigma}} L_0 S_{\sigma} = \begin{pmatrix} P & 0_{7321, 22579} \\ Q & R \end{pmatrix}, \quad (3.28)$$

$$\text{the rank of } P(\xi^0) \text{ is } 7321. \quad (3.29)$$

Let us assume for the moment that this lemma holds and end the proof of Proposition 2. A consequence of Lemma 1 is the following:

**Lemma 2.** *There exists a nonempty open subset  $\omega_0$  of  $\mathcal{C}_2$ ,  $T^* > 0$  and  $\varepsilon^* > 0$ , such that*

$$\det P\left(\frac{1}{T-t}, \varepsilon a(t), x\right) \neq 0, \forall T \in (0, T^*], \forall t \in [T/2, T), \forall \varepsilon \in (0, \varepsilon_0], \forall x \in \omega_0. \quad (3.30)$$

**Proof of Lemma 2.** Let us first point out that  $\det P \in \mathbb{R}[E, S, X]$  and, by (3.29), this polynomial is not the 0 polynomial. Hence there exist a nonnegative integer  $m$  and a polynomial  $\tilde{P} \in \mathbb{R}[E, S, X]$  such that

$$\det P(E, S, X) = S^m \tilde{P}(E, S, X), \quad (3.31)$$

$$\tilde{P}(E, 0, X) \in \mathbb{R}[E, X] \text{ is not the 0 polynomial.} \quad (3.32)$$

By (3.32), there exist  $\delta > 0$ ,  $C' > 0$  and a nonempty open subset  $\omega_0$  of  $\mathcal{C}_2$  such that

$$|\tilde{P}(e, 0, x)| \geq 2\delta, \forall e \in [C', +\infty), \forall x \in \omega_0. \quad (3.33)$$

By the mean value theorem, there exist a positive integer  $l$  and a positive real number  $C^*$  such that

$$|\tilde{P}(e, s, x) - \tilde{P}(e, 0, x)| \leq C^* |s| \left( |e|^l + |s|^l + 1 \right), \forall e \in \mathbb{R}, \forall s \in \mathbb{R}, \forall x \in \omega_0. \quad (3.34)$$

By (2.16), there exists  $\varepsilon^*$  such that

$$\varepsilon^* |a(t)| \left( (T-t)^{-l} + \varepsilon^{*l} |a(t)|^l + 1 \right) \leq \frac{\delta}{C^*}, \forall T \in (0, 2/C'], \forall t \in [T/2, T). \quad (3.35)$$

From (3.33), (3.34) and (3.35), we obtain that

$$\forall T \in (0, 2/C'], \forall t \in [T/2, T), \forall \varepsilon \in (0, \varepsilon^*], \forall x \in \omega_0,$$

$$|\tilde{P}((T-t)^{-1}, \varepsilon a(t), x)| \geq \delta,$$

which concludes the proof of Lemma 2. ■

Let us now go back to the proof of Proposition 2. For every positive integer  $l$ , we denote by  $\text{Id}_l$  the identity map of  $\mathbb{R}^l$ . By (3.30), there exists  $U \in C^\infty(Q_0; \mathcal{M}_{7321 \times 7321}(\mathbb{R}))$  such that

$$U(t, x)P(t, x, \varepsilon) = \text{Id}_{7321}, \forall x \in \omega_0. \quad (3.36)$$

Let  $\tilde{U} \in C^\infty(Q_0; \mathcal{M}_{7321 \times 30360}(\mathbb{R}))$  be defined by

$$\tilde{U}(t, x) := (U(t, x) \quad 0_{7321, 23039}), \forall x \in \omega_0. \quad (3.37)$$

From (3.28), (3.36) and (3.37), one has

$$\tilde{U}(t, x)S_{\bar{\sigma}}L_0(t, x) = (\text{Id}_{7321} \quad 0_{7321, 22579})S_{\bar{\sigma}}^{-1}, \forall x \in \omega_0. \quad (3.38)$$

Let  $K \in \mathcal{M}_{6, 7321}(\mathbb{R})$  be defined by

$$K := (\text{Id}_6 \quad 0_{6, 7315}). \quad (3.39)$$

From (3.27), (3.38) and (3.39), one has

$$K\tilde{U}(t, x)S_{\bar{\sigma}}L_0(t, x) = (\text{Id}_6 \quad 0_{6, 29894})S_{\bar{\sigma}}^{-1} = (\text{Id}_6 \quad 0_{6, 29894}), \forall x \in \omega_0,$$

which shows that (3.24) holds with  $N(t, x) := K\tilde{U}(t, x)S_{\bar{\sigma}}$ , and ends the proof of Proposition 2.  $\blacksquare$

To finish the proof of Proposition 2, it is now enough to prove Lemma 1.

**Proof of Lemma 1.** The fact that the dependence of  $\bar{y}$  and its derivatives in the time variable is quite complicated (it is both exponential and fractional) compared to the dependence in the space variable (which is polynomial) is problematic, because it is not very convenient to use for computations on a computer. In the previous proof we have seen  $\det P$  as a polynomial in  $s = \varepsilon a(t)$ ,  $e = \frac{1}{T-t}$  (which corresponds to terms appearing when we differentiate  $t \mapsto a(t)$ ) and  $x$ . Assume that we fix  $e = 0$ : This is equivalent to do as if the derivatives of  $a$  were all identically the null function, i.e. to do as if the function  $t \mapsto a(t)$  were replaced by a constant function, which is simpler than our original function  $a$ . We will then impose  $e^0 = 0$  for our computations. Let us set  $\xi^0 := (e^0, s^0, x^0)$  with  $e^0 = 0$ ,  $s^0 = 1$  and  $x^0 = (1.1, 1.2, 1.3)$ .

First of all, let us prove that a decomposition under the form of (3.28) exists at least at point  $\xi^0$ . We present in the Appendix A how we computed the matrix

$$L_0^0 := L_0(\xi^0) \in \mathcal{M}_{30360 \times 29900}(\mathbb{R})$$

thanks to a  $C^{++}$  program.

From now on we assume that we have matrix  $L_0^0$  at our disposal and we are going to explain how to exploit it to obtain Lemma 1.

We begin with reordering the columns so that the null columns of  $L_0^0$  are moved to the last columns. One verifies for example thanks to Matlab that there are exactly 140 such columns. There exist  $\bar{\sigma} \in \mathfrak{S}_{29900}$  and  $N^0 \in \mathcal{M}_{30360 \times 29760}(\mathbb{R})$  such that

$$L_0^0 S_{\bar{\sigma}} = \begin{pmatrix} N^0 & 0_{30360 \times 140} \end{pmatrix}. \quad (3.40)$$

One problem is that it could happen that some columns of  $L_0^0$  are equal to 0 but the corresponding columns of  $L_0$  are not identically null. However, we check that it is not the case (thanks to the evaluation function described in Appendix A).

Let us recall that our goal is to extract a well-chosen submatrix of  $L_0^0$  which is of maximal rank. A reasonable hope would have been that the matrix  $N^0$  (of size  $30360 \times 29760$ ) itself is of maximal rank 29760 (we would then have obtained something similar to Lemma 1 by choosing some squared extracted matrix of maximal rank  $P^0$  of  $N^0$ , which is always possible, the matrix  $P^0$  would then have been of size  $29760 \times 29760$  and the non-selected lines would be permuted to obtain matrices  $Q^0$  of size  $600 \times 29760$  and  $R^0$  of size  $600 \times 140$ ). However it turns out to be false, as we will see later on.

Since computing the rank of  $L_0^0$  on a computer is too long because of its size, we introduce the notion of structural rank.

**Definition 4.** Let  $A \in \mathcal{M}_{n,m}(\mathbb{R})$  and  $B \in \mathcal{M}_{n,m}(\mathbb{R})$ . We say that  $A$  and  $B$  are structurally equivalent if the following property is verified:

$$A_{ij} = 0 \Leftrightarrow B_{ij} = 0.$$

This is an equivalence relation on  $\mathcal{M}_{n,m}(\mathbb{R})$ , and we will call  $Cl(A)$  the equivalence class of  $A$ . The structural rank of a matrix  $A$  (denoted  $\mathbf{sprank}(A)$  in the following) is the maximal rank of the elements of  $Cl(A)$ . Equivalently, if we fill randomly the nonzero coefficients of  $A$  with probability 1 the rank of  $A$  is the structural rank.

One sees that the structural rank does not depend on the coefficients of the matrix but only on the distribution of the zeros in the matrix and is never less than the rank. The advantage of the structural rank is that it can be computed fast (in a couple of seconds in our case), especially on sparse matrices. It corresponds to the function  $\mathbf{sprank}$  in Matlab.

Computing the structural rank of  $N^0$  thanks to Matlab we find that

$$\mathbf{sprank}(N^0) = 28654 < 29760,$$

hence there is no hope that the rank of  $N^0$  is maximal.

To extract a submatrix of  $N^0$  which is of maximal rank, we can, for example, begin with extracting a submatrix of  $P^0$  which is of maximal structural rank, and verify that it is of maximal rank too. The right way to do this is to explore more carefully how the structural rank is computed. In fact the key point is the existence of a decomposition in block triangular form (which is related to the Dulmage-Mendelsohn decomposition for the bipartite graph associated to any matrix, see [14] and [26]) of a matrix.

**Proposition 3.** *Let  $A$  be a matrix. Then one can permute the columns and the lines of  $A$  to obtain a matrix of the following form:*

$$\begin{pmatrix} A_{11} & A_{12} & A_{13} & A_{14} \\ 0 & 0 & A_{23} & A_{24} \\ 0 & 0 & 0 & A_{34} \\ 0 & 0 & 0 & A_{44} \end{pmatrix} \quad (3.41)$$

where:

1.  $(A_{11}, A_{12})$  is the underdetermined part of the matrix, it always has more columns than rows.
2.  $(A_{13}, A_{14})$  is the overdetermined part of the matrix, it always has more rows than columns.
3.  $A_{12}, A_{23}, A_{34}$  are square matrices with nonzero diagonals (notably these matrices are of maximal structural rank)
4.  $A_{23}$  is the well-determined part of the matrix (if the matrix is square and non-singular, it is the entire matrix).

Moreover, one can permute rows and columns so that  $A_{23}$  is also block triangular. The decomposition obtained is called the block triangular form of matrix  $A$ . The structural rank of  $A$  is given by the sum of the structural ranks of  $A_{12}, A_{23}, A_{34}$ .

The block triangular form (3.41) (also called the coarse decomposition) of the matrix is in fact given by the `dmperm` function in Matlab, which also gives the permutation that makes the matrix be in the form of (3.41). It also gives the block form of the well-determined part (which is called the fine decomposition). One easily understands how to obtain a matrix of the form of (3.28) thanks to this decomposition: One can permute the blocks to obtain

$$\begin{pmatrix} A_{34} & 0 & 0 & 0 \\ A_{44} & 0 & 0 & 0 \\ A_{14} & A_{11} & A_{12} & A_{13} \\ A_{24} & 0 & 0 & A_{23} \end{pmatrix}, \quad (3.42)$$

from which we easily deduce decomposition (3.28).

To simplify the computations, we are not going to apply this block triangular decomposition directly to  $L_0^0$  but to a well-chosen submatrix  $\tilde{L}_0^0$ . First of all, we go back to  $L_0$  and select some equations and unknowns: There exist  $\sigma^0 \in \mathfrak{S}_{29900}$ ,  $\tilde{\sigma}^0 \in \mathfrak{S}_{30360}$ ,  $\tilde{Q}^0 \in \mathcal{M}_{16623 \times 14630}(\mathbb{R})$  and  $\tilde{R}^0 \in \mathcal{M}_{16623 \times 15270}(\mathbb{R})$  such that (see (3.42))

$$S_{\tilde{\sigma}^0} L^0 S_{\sigma^0} = \begin{pmatrix} \tilde{L}^0 & 0_{13737 \times 15270} \\ \tilde{Q}^0 & \tilde{R}^0 \end{pmatrix}, \quad (3.43)$$

where  $\tilde{L}^0$  corresponds to the equations we obtain by differentiating the two first equations of (3.21) 15 times and the last equation 17 times, so that  $\tilde{L}^0$  is of size  $(G(15), H(15)) = (13737, 14630)$  (here there are more unknowns than equations but we will see that this will not be a problem).

Let us call

$$\tilde{L}_0^0 := \tilde{L}^0(\xi^0),$$

and let us call

$$\begin{aligned}\tilde{Q}^0 &:= \tilde{Q}(\xi^0), \\ \tilde{R}^0 &:= \tilde{R}(\xi^0), \\ \tilde{S}^0 &:= \tilde{S}(\xi^0).\end{aligned}$$

One has

$$S_{\tilde{\sigma}^0} L_0^0 S_{\sigma^0} = \begin{pmatrix} \tilde{L}_0^0 & 0_{13737 \times 15270} \\ \tilde{Q}^0 & \tilde{R}^0 \end{pmatrix}.$$

Thanks to Matlab, we find the Dulmage-Mendelsohn decomposition of  $\tilde{L}_0^0$  and observe that there exists some permutations matrices  $\sigma^1$  and  $\tilde{\sigma}^1$  such that (see (3.42))

$$S_{\tilde{\sigma}^1} \tilde{L}_0^0 S_{\sigma^1} = \begin{pmatrix} \bar{L}_0^0 & 0_{9050 \times 5578} \\ \bar{Q}^0 & \bar{R}^0 \end{pmatrix},$$

with  $\bar{Q}^0 \in \mathcal{M}_{4687 \times 9050}(\mathbb{R})$ ,  $\bar{R}^0 \in \mathcal{M}_{4687 \times 5578}(\mathbb{R})$ , and  $\bar{L}_0^0$  is of maximal structural rank and square of size  $9050 \times 9050$  (it corresponds to the block  $A_{34}$  in the triangular block decomposition). Applying the Dulmage-Mendelsohn algorithm now on  $\bar{L}_0^0$ , we can write  $\bar{L}_0^0$  in an block upper triangular form with 352 diagonal blocks (this is the fine decomposition), the first 351 of them being of “small” size and the latter one being of size 7321. Let us call  $\bar{L}_{0(i,j)}^0$  (with  $(i,j) \in \llbracket 1; 352 \rrbracket^2$ ) the blocks of  $\bar{L}_0^0$ .

Using this decomposition, one can see (using Matlab) that  $\bar{L}_0^0$  is not of maximal rank. However, by computing the rank of the block  $\bar{L}_{0(352,352)}^0$  thanks to Matlab, one sees that it is of maximal rank 7321. Moreover, we verify that

$$\text{the columns corresponding to the unknowns } z_1^1, z_2^1, z_3^1, z_1^2, z_2^2, z_3^2 \text{ appear in this block,} \quad (3.44)$$

by looking carefully on Matlab where the columns corresponding to these unknowns have been moved under the action of the permutation matrices  $S_{\sigma^0}$  and  $S_{\sigma^1}$ . More precisely,  $z_i^1$  corresponds to the  $i$ -th column of  $\bar{L}_{0(352,352)}^0$  and  $z_i^2$  to the  $(3632 + i)$ -th column of  $\bar{L}_{0(352,352)}^0$ .

Let us call

$$P^0 := \bar{L}_{0(352,352)}^0.$$

There exist  $\sigma \in \mathfrak{S}_{29900}$ ,  $\tilde{\sigma} \in \mathfrak{S}_{30360}$ ,  $Q^0 \in \mathcal{M}_{23039 \times 7321}(\mathbb{R}[E, S, X])$  and  $R^0 \in \mathcal{M}_{23039 \times 22579}(\mathbb{R}[E, S, X])$  such that

$$S_{\tilde{\sigma}} \tilde{L}_0^0 S_{\sigma} = \begin{pmatrix} P^0 & 0_{7321 \times 22579} \\ Q^0 & R^0 \end{pmatrix}, \quad (3.45)$$

the rank of the first block  $P^0$  being maximal. The distribution of the nonzero coefficients of  $P^0$  is given in Figure 2.




Figure 2: Distribution of the nonzero coefficients of  $P^0$ .

Now, we come back to the matrix  $L_0$  and we consider the following matrices:

$$\tilde{M}^0 := S_{\tilde{\sigma}} \tilde{L}^0 S_{\sigma},$$

where  $S_{\tilde{\sigma}}$  and  $S_{\sigma}$  are introduced in (3.45).

Let us call  $\Theta$  the set of all coefficients of  $L_0$  (considered as a polynomial in the variables  $E, S, X$ ) that are not identically zero, and  $\Theta^0$  the set of all coefficients of  $L_0^0$  that are not equal to 0. Clearly  $\Theta^0 \subset \Theta$  (in fact thanks to Matlab one can see that  $\Theta^0$  is much smaller than  $\Theta$ ), moreover  $\Theta \setminus \Theta^0$  corresponds to the nonzero coefficients of the matrix that become identically null when we change  $a(t)$  into the function identically equal to 1 and apply it at point  $\xi^0$ . What could happen is that  $\tilde{M}^0$  is not of block triangular form as in (3.45) (the null block of the matrix  $\tilde{M}^0$  may contain some elements of  $\Theta \setminus \Theta^0$ ). However, since the only important thing is the location of the elements of  $\Theta \setminus \Theta^0$  and not their value, one can verify easily on Matlab that

the coefficients of  $\Theta \setminus \Theta^0$  do not influence the block form (3.45) (by looking where the elements of  $\Theta \setminus \Theta^0$  are moved under the action of  $S_{\bar{\sigma}}$  and  $S_{\sigma}$ ), i.e. the permutations  $S_{\bar{\sigma}}$  and  $S_{\sigma}$  also give a decomposition under the form of (3.45) for  $\tilde{L}^0$ : There exists

$$\tilde{P} \in \mathcal{M}_{7321 \times 7321}(\mathbb{R}[E, S, X]),$$

$$\tilde{Q} \in \mathcal{M}_{23039 \times 7321}(\mathbb{R}[E, S, X])$$

and

$$\tilde{R} \in \mathcal{M}_{23039 \times 22579}(\mathbb{R}[E, S, X])$$

such that

$$S_{\bar{\sigma}} \tilde{L}^0 S_{\sigma} = \begin{pmatrix} \tilde{P} & 0_{7321 \times 22579} \\ \tilde{Q} & \tilde{R} \end{pmatrix}, \quad (3.46)$$

with the relations  $P(\xi^0) = P^0, Q(\xi^0) = Q^0, R(\xi^0) = R^0$ .

Property (3.28) follows then directly from (3.43) and (3.46), (3.29) is a direct consequence of the above construction, and (3.27) can be easily deduced by permuting some lines and columns of  $S_{\bar{\sigma}}$  and  $S_{\sigma}$  (thanks to Property (3.44)). Finally, Lemma 1 holds. ■

Consequently Proposition 2 holds. Moreover, one observes that the linear partial differential operator  $\mathcal{M}_0$  that we have just created so that (3.11) holds is exactly  $\mathcal{P}^*$ , (where  $\mathcal{P}$  is the differential version of the matrix  $P$  seen as a partial differential operator acting on  $(z_1^1, z_2^1, z_3^1, z_1^2, z_2^2, z_3^2)$ ) and is of order 17 (because we have differentiated the equations of (3.16) 16 times, and (3.21) was obtained by differentiating System (3.16) 1 time). Hence, the corresponding operator  $\mathcal{M}$  in equality (3.5) is also of order 17.

This concludes the proof of Proposition 2. ■

### 3.3 Controllability of the linear control system (3.6)

In this subsection we prove some technical lemmas that imply the null controllability of (3.6) with controls which are derivatives of smooth enough functions having a small support. This is needed to ensure that the controls are in the image of  $\mathcal{B}$  (this is exactly Assumption  $\mathcal{A}_1$ ) and to take into account Remark 3.

The first lemma is the following one, (remind that  $\bar{y}$  is one of the trajectories constructed in Section 2). It consists in a Carleman estimate with curl observation.

**Lemma 3.** *Let  $\theta : \Omega \rightarrow [0, +\infty)$  be a lower semi-continuous function which is not identically 0 and let  $r \in (0, 1)$ . There exists  $C_1 > 0$  such that, for every  $K_1 > C_1$ , every  $\nu \geq K_1(1/r - 1)$ , there exists  $\varepsilon^0$  such that for every  $\varepsilon \in (0, \varepsilon^0)$ , there exists  $C > 0$  such that, for every  $g \in L^2((0, T) \times \Omega)^3$*



and for every solution  $z \in L^2((0, T), H) \cap L^\infty((0, T), V)$  of the adjoint of the linearized Navier-Stokes system

$$\begin{cases} -z_t - \Delta z - (\bar{y} \cdot \nabla^{tr})z - (z \cdot \nabla)\bar{y} + \nabla\pi = g & \text{in } Q, \\ \nabla \cdot z = 0 & \text{in } Q, \\ z = 0 & \text{on } [0, T] \times \partial\Omega, \end{cases} \quad (3.47)$$

one has

$$\begin{aligned} & \|e^{\frac{-\kappa_1}{2r(T-t)^5}} z\|_{L^2((T/2, T), H^1(\Omega)^3)}^2 + \|z(T/2, \cdot)\|_{L^2(\Omega)^3}^2 \\ & \leq C \left( \int_{(T/2, T) \times \Omega} \theta e^{-\frac{\kappa_1}{(T-t)^5}} |\nabla \wedge z|^2 + \int_{(T/2, T) \times \Omega} e^{-\frac{\kappa_1}{(T-t)^5}} |g|^2 \right). \end{aligned} \quad (3.48)$$

**Proof of Lemma 3.** In this proof, our system, which is initially defined on  $(0, T)$ , will only be considered on the interval of time  $(T/2, T)$ . In fact, in what follows (see, in particular, the proof of Proposition 4), we will not act on the system on the interval  $(0, T/2)$ , hence we only need a Carleman estimate on  $(T/2, T)$ . For our proof, we need to use the particular form of our  $\bar{y}$  in time, notably that (see (2.16)),

$$|\bar{y}(t, x)| + |\nabla \bar{y}(t, x)| \leq C \varepsilon e^{\frac{-\nu}{(T-t)^5}}, \quad \forall (t, x) \in (T/2, T) \times \Omega. \quad (3.49)$$

Without loss of generality, we may assume that there exists nonempty open subset  $\omega^*$  of  $\Omega$  such that  $\theta = 1_{\omega^*}$ . Let us now give some extra notations. Let  $\eta^0 \in C^2(\bar{\Omega})$  such that  $|\nabla \eta^0| > 0$  in  $\bar{\Omega} \setminus \omega^*$  and  $\eta^0 = 0$  on  $\partial\Omega$ . For the existence of  $\eta^0$ , see [17, Lemma 1.1, p. 4]. Let us call

$$\alpha(t, x) := \frac{e^{12\lambda\|\eta^0\|_\infty} - e^{\lambda(10\|\eta^0\|_\infty + \eta^0(x))}}{(t - T/2)^5 (T - t)^5}, \quad \xi(t, x) := \frac{e^{\lambda(10\|\eta^0\|_\infty + \eta^0(x))}}{(t - T/2)^5 (T - t)^5}$$

and

$$\alpha^*(t) := \max_{x \in \bar{\Omega}} \alpha(t, x).$$

We call

$$Q_{/2} := (T/2, T) \times \Omega.$$

Using (3.49) and [20, Proposition 3.1, p. 6] on the adjoint system (3.47) (where we see the first and zero order terms of this equation as a second member, because Proposition 3.1 of [20] concerns only the Stokes system), one has, for  $\lambda \geq C$  and  $s \geq C$ ,

$$\begin{aligned} & s^3 \lambda^4 \int_{Q_{/2}} e^{-2s\alpha - 2s\alpha^*} \xi^3 |\nabla \wedge z|^2 + s \lambda^2 \int_{Q_{/2}} e^{-2s\alpha - 2s\alpha^*} \xi |\nabla(\nabla \wedge z)|^2 \\ & \leq C \left( s^3 \lambda^4 \int_{(T/2, T) \times \omega^*} e^{-2s\alpha - 2s\alpha^*} \xi^3 |\nabla \wedge z|^2 + \varepsilon \int_{Q_{/2}} e^{-2s\alpha^*} e^{-\frac{2\nu}{(T-t)^5}} (|z|^2 \right. \\ & \quad \left. + |\nabla z|^2) + \int_{Q_{/2}} e^{-2s\alpha^*} |g|^2 \right). \end{aligned}$$

In fact, looking carefully at the proof of Proposition 3.1 of [20], one remarks (just by changing the weight  $\rho(t) := e^{-s\alpha^*}$  by  $\rho(t) := e^{-as\alpha^*}$  where  $a$  is a parameter that can be chosen as large as we wish), that the previous inequality can be improved in the following way, as soon as  $s$  is large enough, for every  $a > 0$  (the constant  $C$  depending on  $a$ ):

$$\begin{aligned} & s^3 \lambda^4 \int_{Q_{/2}} e^{-2s\alpha - 2as\alpha^*} \xi^3 |\nabla \wedge z|^2 + s\lambda^2 \int_{Q_{/2}} e^{-2s\alpha - 2as\alpha^*} \xi |\nabla(\nabla \wedge z)|^2 \\ & \leq C \left( s^3 \lambda^4 \int_{(T/2, T) \times \omega^*} e^{-2s\alpha - 2as\alpha^*} \xi^3 |\nabla \wedge z|^2 + \varepsilon \int_{Q_{/2}} e^{-2as\alpha^*} e^{-\frac{2\nu}{(T-t)^5}} (|z|^2 \right. \\ & \qquad \qquad \qquad \left. + |\nabla z|^2) + \int_{Q_{/2}} e^{-2as\alpha^*} |g|^2 \right). \end{aligned} \quad (3.50)$$

As usual, we now change our weights so that they do not vanish at time  $t = T/2$ . Let us call  $l : [T/2, T] \rightarrow \mathbb{R}$  defined by  $l(t) = T^2/16$  on  $[T/2, 3T/4]$  and  $l(t) = (t - T/2)(T - t)$  on  $[3T/4, T]$ . Let us define

$$\begin{aligned} \beta(t, x) & := \frac{e^{12\lambda\|\eta^0\|_\infty} - e^{\lambda(10\|\eta^0\|_\infty + \eta^0(x))}}{l^5(t)}, \quad \gamma(t, x) := \frac{e^{\lambda(10\|\eta^0\|_\infty + \eta^0(x))}}{l^5(t)}, \\ \beta^*(t) & := \max_{x \in \Omega} \beta(t, x), \quad \gamma^*(t) := \max_{x \in \Omega} \gamma(t, x). \end{aligned}$$

Clearly, the functions  $\alpha$  and  $\beta$  coincide on  $[3T/4, T]$ , as well as the functions  $\xi$  and  $\gamma$ . Using classical arguments (see for example the proof of Lemma 1 of [15]), one easily obtains (one can forget the weight  $s$  and  $\lambda$  in front of our terms from now on) the existence of  $C$  (depending now on  $s, \lambda$ , which are assumed to be large enough, and  $a$ ) such that

$$\begin{aligned} & \int_{Q_{/2}} e^{-2(1+a)s\beta^*} |\nabla \wedge z|^2 + \int_{Q_{/2}} e^{-2(1+a)s\beta^*} |\nabla(\nabla \wedge z)|^2 \\ & \leq C \left( \int_{(T/2, T) \times \omega^*} e^{-2as\beta^*} \gamma^*{}^3 |\nabla \wedge z|^2 + \varepsilon \int_{Q_{/2}} e^{-2as\beta^*} e^{-\frac{2\nu}{(T-t)^5}} (|z|^2 + |\nabla z|^2) \right. \\ & \qquad \qquad \qquad \left. + \int_{Q_{/2}} e^{-2as\beta^*} |g|^2 \right). \end{aligned} \quad (3.51)$$

One remarks that, since  $\nabla \cdot z = 0$  in  $Q$  and  $z = 0$  on  $(0, T) \times \partial\Omega$ , one has

$$C \|e^{-2(1+a)s\beta^*} \nabla \wedge z\|_{L^2(Q_{/2})^3}^2 \geq \|e^{-2(1+a)s\beta^*} \nabla z\|_{L^2(Q_{/2})^9}^2. \quad (3.52)$$

Using Poincaré's inequality, we also have

$$C \|e^{-2(1+a)s\beta^*} \nabla z\|_{L^2(Q_{/2})^9}^2 \geq \|e^{-2(1+a)s\beta^*} z\|_{L^2(Q_{/2})^3}^2. \quad (3.53)$$

Putting this into (3.51), one obtains

$$\begin{aligned} & \int_{Q_{/2}} e^{-2(1+a)s\beta^*} |\nabla z|^2 + \int_{Q_{/2}} e^{-2(1+a)s\beta^*} |z|^2 \\ & \leq C \left( \int_{(T/2, T) \times \omega^*} e^{-2as\beta^*} \gamma^{*3} |\nabla \wedge z|^2 + \varepsilon \int_{Q_{/2}} e^{-2as\beta^*} e^{-\frac{2\nu}{(T-t)^5}} (|z|^2 + |\nabla z|^2) \right. \\ & \quad \left. + \int_{Q_{/2}} e^{-2as\beta^*} |g|^2 \right). \end{aligned} \quad (3.54)$$

Let us define, for  $a > 1$ ,

$$K_0 := 2^6(1+a+\sqrt{a})s \frac{e^{12\lambda\|\eta^0\|_\infty} - e^{\lambda(10\|\eta^0\|_\infty + \min_{x \in \Omega} \eta^0(x))}}{T^5}, \quad (3.55)$$

$$K_1 := 2^6(a-\sqrt{a})s \frac{e^{12\lambda\|\eta^0\|_\infty} - e^{\lambda(10\|\eta^0\|_\infty + \min_{x \in \Omega} \eta^0(x))}}{T^5}. \quad (3.56)$$

One has the existence of  $C^* > 0$  (depending on  $a > 1$ ,  $\lambda \gg 1$  and  $s \gg 1$ ) such that

$$e^{-2as\beta^*(t)} \gamma^{*3}(t) \leq C^* e^{\frac{-K_1}{(T-t)^5}}, \quad \forall t \in (T/2, T). \quad (3.57)$$

Moreover, there exists  $\widehat{C} > 0$  (depending on  $a > 1$ ,  $\lambda \gg 1$  and  $s \gg 1$ ) such that

$$e^{-\frac{K_0}{(T-t)^5}} \leq \widehat{C} e^{-\frac{2(1+a)s\beta^*}{T^5(t)}}, \quad \forall t \in (T/2, T). \quad (3.58)$$

Fixing  $s, \lambda$  and making  $a \rightarrow +\infty$ , one easily sees that  $K_0/K_1 = \frac{1+a+\sqrt{a}}{a-\sqrt{a}} \rightarrow 1^+$  so that for every  $r \in (0, 1)$ , we have for  $a$  large enough,  $K_0 < K_1/r$ . For  $\varepsilon > 0$  small enough and for  $\nu$  large enough ( $\nu \geq K_0 - K_1$ ), one can absorb the undesired terms  $\varepsilon \int_{Q_{/2}} e^{-2as\beta^*} e^{-\frac{2\nu}{(T-t)^5}} (|z|^2 + |\nabla z|^2)$  from the right-hand side of (3.54). Then using some classical energy estimates together with (3.57) and (3.58), one obtains (3.48).  $\blacksquare$

From now on, we set

$$\rho_r(t) := e^{\frac{-K_1}{r(T-t)^5}}, \quad \rho_1(t) := e^{\frac{-K_1}{(T-t)^5}}.$$

Let us now derive from this Carleman inequality a result of null-controllability with controls which are derivatives of smooth functions. Let  $\widehat{1}_{\omega_0} : \mathbb{R}^3 \rightarrow [0, 1]$  be a function of class  $C^\infty$  which is not identically equal to 0 and having a support included in  $\omega_0$ , where  $\omega_0$  was introduced in Lemma 2. We apply Lemma 3 with  $\theta = \widehat{1}_{\omega_0}$ . One has the following proposition.

**Proposition 4.** *With the notations of Lemma 3, let  $f \in L^2(Q)^3$  be such that  $\rho_r^{-1/2}f \in L^2(Q)^3$  and let us consider the following linearized Navier-Stokes control system*

$$\begin{cases} y_t - \Delta y + (\bar{y} \cdot \nabla)y + (y \cdot \nabla)\bar{y} + \nabla p = f + \nabla \wedge ((\nabla \wedge u)\widehat{1}_{\omega_0}) & \text{in } Q, \\ \nabla \cdot y = 0 & \text{in } Q, \\ y = 0 & \text{on } [0, T] \times \partial\Omega, \end{cases} \quad (3.59)$$

where the control is  $u$ . Then, for every  $y_0 \in V$ , there exists a solution  $(y, p, u)$  of (3.59), such that  $y(0, \cdot) = y^0$  and for every  $\tilde{K}_1$  verifying  $0 < \tilde{K}_1 < K_1$ ,

$$e^{\frac{\tilde{K}_1(2-1/r)}{2(T-t)^5}} (\nabla \wedge u)\widehat{1}_{\omega_0} \in L^2((T/2, T), H^{53}(\Omega)^3) \cap H^{27}((T/2, T), H^{-1}(\Omega)^3), \quad (3.60)$$

$$e^{\frac{\tilde{K}_1}{2(T-t)^5}} y \in L^2((T/2, T), H^2(\Omega)^3) \cap L^\infty((T/2, T), H^1(\Omega)^3). \quad (3.61)$$

**Proof of Proposition 4.** For what follows, we only control on the interval of time  $(T/2, T)$ , i.e. we set  $u = 0$  on  $(0, T/2)$  and let the corresponding solution  $(y, p)$  of (3.59) on  $(0, T/2)$  evolve naturally until time  $T/2$ . Let  $y^{T/2} = y(T/2, \cdot)$ . We are now going to define  $(y, p, u)$  on  $(T/2, T)$ .

Let us call  $X_m := L^2((T/2, T), H^{2m}(\Omega)^3) \cap H^m((T/2, T), L^2(\Omega)^3)$ , which is an Hilbert space for the usual scalar product, and  $X'_m$  its dual with pivot space  $L^2(Q/2)^3$ .

Let  $\mathcal{L} : C^\infty(Q/2)^3 \rightarrow C^\infty(Q/2)^3$  be the linearized time-dependant Navier-Stokes operator

$$\mathcal{L}y = y_t - \Delta y + (\bar{y} \cdot \nabla)y + (y \cdot \nabla)\bar{y}$$

and  $\mathcal{L}^* : C^\infty(Q/2)^3 \rightarrow C^\infty(Q/2)^3$  be its adjoint. Let  $P_0$  be the set of  $(w, h)$  such that

$$w \in L^2((T/2, T), H^2(\Omega)^3) \cap H^1((T/2, T), L^2(\Omega)^3) \cap L^\infty((T/2, T), H^1(\Omega)^3),$$

$$h \in L^2((T/2, T), H^1(\Omega)) \text{ and } \int_{\Omega} h(t, x) dx = 0, \forall t \in (T/2, T),$$

$$\nabla \cdot w = 0 \text{ in } Q/2$$

$$w = 0 \text{ on } (T/2, T) \times \partial\Omega,$$

$$\mathcal{L}^*w + \nabla h \in C_0^\infty(Q/2)^3.$$

Let  $a$  be the following bilinear form defined on  $P_0$  (according to the right-hand side of (3.48)):

$$\begin{aligned} a((z, q), (w, h)) := & \langle \sqrt{\rho_1}(\mathcal{L}^*z + \nabla q), \sqrt{\rho_1}(\mathcal{L}^*w + \nabla h) \rangle_{X_{26}} \\ & + \int_{Q/2} \widehat{1}_{\omega_0} \rho_1 (\nabla \wedge z) \cdot (\nabla \wedge w). \end{aligned}$$

From the unique continuation property for Stokes-like systems induced by (3.48), we deduce that  $a$  is a scalar product on  $P_0$ . Let  $P$  be the completion of  $P_0$  for the norm associated to  $a$ . Then,  $(P, a)$  is a Hilbert space and  $a$  is continuous and coercive on  $P$ . Let us now consider the linear form  $l$  defined on  $P_0$  by

$$l(w, h) := \int_{Q/2} fw + \int_{\Omega} y^{T/2} w(T/2, \cdot).$$

Using the Cauchy-Schwarz inequality together with (3.48), one has, for every  $(w, h) \in P_0$ ,

$$\begin{aligned} |l(w, h)| &\leq \|\rho_r^{-1/2} f\|_{L^2(Q_{/2})^3} \|\sqrt{\rho_r} w\|_{L^2(Q_{/2})^3} + \|y^{T/2}\|_{L^2(\Omega)^3} \|w(T/2, \cdot)\|_{L^2(\Omega)^3} \\ &\leq C(\|\rho_r^{-1/2} f\|_{L^2(Q_{/2})^3} + \|y^{T/2}\|_{L^2(\Omega)^3}) \sqrt{a((w, h), (w, h))}. \end{aligned} \quad (3.62)$$

Hence  $l$  can be uniquely extended as a continuous linear map from  $P$  into  $\mathbb{R}$ . Then (3.62) holds for every  $(w, h) \in P$ . Applying the Lax-Milgram theorem, there exists a unique  $(\hat{z}, \hat{q}) \in P$  verifying

$$a((\hat{z}, \hat{q}), (w, h)) = l(w, h), \forall (w, h) \in P. \quad (3.63)$$

We then set

$$\begin{aligned} \hat{y} := \rho_1(\mathcal{L}^* \hat{z} + \nabla \hat{q}) + \sum_{k=1}^{52} (-1)^k (\tilde{\Delta}^k(\mathcal{L}^* \hat{z} + \nabla \hat{q})) \\ + \sqrt{\rho_1} \left( \sum_{k=1}^{26} (-1)^k \partial_t^{2k} (\sqrt{\rho_1}(\mathcal{L}^* \hat{z} + \nabla \hat{q})) \right) \end{aligned} \quad (3.64)$$

and

$$\hat{u} := \rho_1 \nabla \wedge (\hat{\mathbf{1}}_{\omega_0} \nabla \wedge \hat{z}),$$

where

$$\tilde{\Delta}^k f := \sum_{1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq 3} f_{i_1 i_2 i_2 \dots i_k i_k}.$$

Equation (3.63) implies that we have for every  $(w, h) \in P$ ,

$$\begin{aligned} \langle \sqrt{\rho_1}(\mathcal{L}^* \hat{z} + \nabla \hat{q}), \sqrt{\rho_1}(\mathcal{L}^* w + \nabla h) \rangle_{X_{26}} + \int_{Q_0} \hat{\mathbf{1}}_{\omega_0} \rho_1 \nabla \wedge \hat{z} \cdot \nabla \wedge w \\ = \int_{Q_{/2}} f w + \int_{\Omega} y^{T/2} w(T/2, \cdot) dx. \end{aligned} \quad (3.65)$$

From (3.65), one obtains that, for every  $g \in C_0^\infty(Q_{/2})^3$ ,

$$\int_{Q_{/2}} f w + \int_{Q_0} \hat{u} w + \int_{\Omega} y^{T/2} w(T/2, \cdot) dx = \langle \hat{y}, g \rangle_{X_{26'}, X_{26}},$$

as soon as  $(w, h)$  verifies

$$\begin{cases} \mathcal{L}^* w + \nabla h = g & \text{in } Q_{/2}, \\ \nabla \cdot w = 0 & \text{in } Q_{/2}, \\ w = 0 & \text{on } [T/2, T] \times \partial\Omega, \\ z(T, \cdot) = 0 & \text{in } \Omega, \end{cases} \quad (3.66)$$

that is to say there exists a pressure  $\widehat{p}$  such that  $(\widehat{y}, \widehat{p})$  is the solution by transposition of

$$\left\{ \begin{array}{ll} \widehat{y}_t - \Delta \widehat{y} + (\overline{y} \cdot \nabla) \widehat{y} + (\widehat{y} \cdot \nabla) \overline{y} + \nabla \widehat{p} = f + \widehat{u} & \text{in } Q_{/2}, \\ \nabla \cdot y = 0 & \text{in } Q_{/2}, \\ y = 0 & \text{on } [T/2, T] \times \partial\Omega, \\ y(T/2, \cdot) = y^{T/2} & \text{on } \Omega. \end{array} \right. \quad (3.67)$$

Moreover, the fact that  $a((\widehat{z}, \widehat{q}), (\widehat{z}, \widehat{q})) < +\infty$  easily implies (by coming back to the definition of the norm in  $X'_{26}$  and using equality (3.64)) that

$$(T-t)^{512} \rho_1^{-1/2} \widehat{y} \in X'_{26}.$$

Using regularity results about linearized Navier-Stokes systems and a bootstrap argument, one obtains that, for every  $\tilde{K}_1$  verifying  $0 < \tilde{K}_1 < K_1$ , we have, with  $\tilde{\rho}_1(t) := e^{-\frac{\tilde{K}_1}{(T-t)^5}}$ ,

$$\tilde{\rho}_1^{-1/2} \widehat{y} \in L^2((T/2, T), H^2(\Omega)^3) \cap L^\infty((T/2, T), H^1(\Omega)^3), \quad (3.68)$$

and  $(\widehat{y}, \widehat{p})$ , which was defined by transposition, is now the weak solution of (3.67).

Let us now focus on  $\widehat{u}$ . Thanks to inequality (3.48),  $a((\widehat{z}, \widehat{q}), (\widehat{z}, \widehat{q})) < +\infty$  implies that

$$\int_{Q_{/2}} \rho_r \widehat{z}^2 < \infty \quad (3.69)$$

If we set  $u := \rho_1 \widehat{z}$ , (3.69) is equivalent to

$$\rho_1^{-1} \rho_r^{1/2} u \in L^2(Q_{/2}).$$

Using regularity results for  $\mathcal{L}^*$  applied on  $\rho_1^{-1} \rho_r^{1/2} u$  and a bootstrap argument, one obtains

$$\tilde{\rho}_1^{-1} \rho_r^{1/2} u \in X_{27}. \quad (3.70)$$

Proposition 4 follows using (3.68) and (3.70). ■

### 3.4 Null-controllability of (3.1)

To finish, one can gather the results of Subsection 3.2 and Subsection 3.3 in order to apply Proposition 1 and obtain a controllability result on (3.1). However, we cannot work in the  $C^\infty$  setting of Proposition 1, so we need to take into account Remark 3 and to be careful concerning the spaces we are working with.

**Proposition 5.** *For every  $T > 0$  small enough, for every  $\alpha \in (0, 1)$ , there exists  $r_0 \in (0, 1)$  such that for every  $r \in (r_0, 1)$ , there exists  $C_1 > 0$  such that for every  $K_1 > C_1$ , for every  $f \in L^2(Q)$  be such that  $e^{\frac{K_1}{2r(T-t)^5}} f \in L^2(Q)^3$  and for every  $y^0 \in V$ , if*

$$\nu = \frac{1-r}{r} K_1, \quad (3.71)$$

there exists a solution  $(y, p, v)$  of the following linearized Navier-Stokes control system

$$\left\{ \begin{array}{ll} y_t - \Delta y + (\bar{y} \cdot \nabla) y + (y \cdot \nabla) \bar{y} + \nabla p = f + (0, 0, 1_{\omega_0} v) & \text{in } Q, \\ \nabla \cdot y = 0 & \text{in } Q, \\ y(0, \cdot) = y^0 \text{ in } \Omega, & \\ y(T, \cdot) = 0 \text{ in } \Omega, & \\ y = 0 & \text{on } [0, T] \times \partial\Omega, \end{array} \right. \quad (3.72)$$

such that

$$e^{\frac{\alpha K_1}{2(T-t)^5}} y \in L^2((0, T), H^2(\Omega)^3) \cap L^\infty((0, T), H^1(\Omega)^3), \quad (3.73)$$

$$e^{\frac{\alpha K_1}{2(T-t)^5}} v \in L^2(Q). \quad (3.74)$$

**Proof of Proposition 5.** We want to apply Proposition 1. First of all, we deal with Assumption  $\mathcal{A}_1$ . We apply Proposition 4: There exists a solution  $(y^*, p^*, u^*)$  of (3.59) such that  $y^*(0, \cdot) = y^0$  and, for every  $\tilde{K}_1$  verifying  $0 < \tilde{K}_1 < K_1$ ,

$$e^{\frac{\tilde{K}_1(2-1/r)}{2(T-t)^5}} \widehat{1}_{\omega_0}(\nabla \wedge u^*) \in L^2((T/2, T), H^{53}(\Omega)^3) \cap H^{27}((T/2, T), H^{-1}(\Omega)^3), \quad (3.75)$$

$$e^{\frac{\tilde{K}_1}{2(T-t)^5}} y^* \in L^2((T/2, T), H^2(\Omega)^3) \cap L^\infty((T/2, T), H^1(\Omega)^3). \quad (3.76)$$

Using well-known interpolation results (see for example [23, Section 13.2, p. 96]) and setting  $n := 27$ , we obtain that

$$e^{\frac{\tilde{K}_1(2-1/r)}{2(T-t)^5}} \widehat{1}_{\omega_0}(\nabla \wedge u^*) \in H^{(2n-1)/3}(Q_0) \subset H^{17}(Q_0).$$

Let us call  $w := \widehat{1}_{\omega_0}(\nabla \wedge u^*)$ , which is supported in  $Q_0$ . One observes that

$$\nabla \wedge w = \begin{pmatrix} w_3^2 - w_2^3 \\ w_3^1 - w_1^3 \\ w_2^1 - w_1^2 \end{pmatrix}.$$

Hence in view of equality (3.8) and setting

$$f^1 = 0, f^2 = -w^3, f^3 = w^2, f^4 = -w^3, f^5 = 0, f^6 = w^1, f^7 = w_2^1 - w_1^2,$$

one has  $\nabla \wedge w \in \text{Im}(\mathcal{B})$  and Assumption  $\mathcal{A}_1$  holds.

Now, we observe that Assumption  $\mathcal{A}_2$  follows from Proposition 2. Let  $(\tilde{y}, \tilde{p}, \tilde{v})$  be defined by

$$(\tilde{y}, \tilde{p}, \tilde{v}) := -\mathcal{M}w,$$

where  $\mathcal{M}$  is as in (the proof of) Proposition 2. It makes sense to apply  $\mathcal{M}$  to  $w$  because  $\mathcal{M}$  is a partial differential operator of order 17 and  $w \in H^{17}(Q_0)$ .

Using the fact that the operator  $\mathcal{M}$  is a partial differential operator of order 17 and that the coefficients of  $\mathcal{M}$  explodes at time  $t = T$  at rate at most  $e^{\frac{7321\nu}{(T-t)^5}}$ , as it follows from the construction of  $\tilde{y}$  given in Section 2 (see in particular (2.16)) and the construction of  $M$  given in the proof of Proposition 2, one has

$$e^{\frac{K_2}{2(T-t)^5}} v \in L^2(Q_0), \quad (3.77)$$

$$e^{\frac{K_2}{2(T-t)^5}} \tilde{y} \in L^2(Q_0), \quad (3.78)$$

for every  $K_2 < K_1(2 - 1/r) - 7321\nu$ . In order to obtain  $\tilde{y}(T, \cdot) = 0$ , it is enough to have an exponential decay for  $y$  at time  $T$ , i.e. to impose

$$K_1(2 - 1/r) - 7321\nu > 0,$$

which, with (3.71), is equivalent to

$$r > \frac{7322}{7323},$$

which can be insured since  $r$  can be arbitrarily chosen close to 1. Let  $\alpha \in (0, 1)$ . We take

$$r_0 := \frac{7322}{7323 - \alpha}. \quad (3.79)$$

Then if  $r \in (r_0, 1)$ , one has

$$\alpha K_1 < K_1 - 7321 K_1 \frac{1-r}{r}. \quad (3.80)$$

By (3.80), there exists  $K_2$  such that

$$\alpha K_1 < K_2 < K_1 - 7321 K_1 \frac{1-r}{r} = K_1 - 7321\nu. \quad (3.81)$$

Finally, one can apply Proposition 1 and we set

$$(y, p) := (y^* + \tilde{y}, p^* + \tilde{p}).$$

Thanks to (3.75), (3.76), (3.77) and (3.78), one has

$$\begin{aligned} e^{\frac{K_2}{2(T-t)^5}} v &\in L^2(Q_0), \\ e^{\frac{K_2}{2(T-t)^5}} y &\in L^2(Q_0). \end{aligned}$$

Then, using usual regularity results for the linearized Navier-Stokes operators on  $e^{\frac{K_2}{2(T-t)^5}} y$  now considered on the entire time interval  $(0, T)$  ( $y \in L^2((0, T/2) \times \Omega)$ , hence we also have  $e^{\frac{K_2}{2(T-t)^5}} y \in L^2((0, T/2) \times \Omega)$ ), we obtain

$$e^{\frac{\alpha K_1}{2(T-t)^5}} y \in L^2((0, T), H^2(\Omega)^3) \cap L^\infty((0, T), H^1(\Omega)^3),$$

as soon as  $y^0 \in V$ . The proof of Proposition 5 is completed.  $\blacksquare$



## 4 Proof of Theorem 1

To conclude, we are going to apply an inverse mapping theorem to go back to the nonlinear system, which is the following (see [2, Chapter 2, Section 2.3]):

**Proposition 6.** *Let  $E$  and  $F$  be two Banach spaces. Let  $e_0 \in E$  and  $\mathcal{A} : E \rightarrow F$  which is of class  $C^1$  in a neighborhood of  $e_0$ . Assume that  $d\mathcal{A}(e_0) \in \mathcal{L}_c(E, F)$  is a surjective operator. Then there exist  $\eta > 0$  and  $C > 0$  such that for every  $g \in F$  verifying  $\|g - \mathcal{A}(e_0)\| < \eta$ , there exists  $e \in E$  such that*

1.  $\mathcal{A}(e) = g$ ,
2.  $\|e - e_0\|_E \leq C\|g - \mathcal{A}(e_0)\|_F$ .

We are going to use the same techniques as in [12]. Let  $\alpha \in (0, 1)$ , and let us set consider a  $r \in (r_0, 1)$  where  $r_0$  verifies (3.79). We apply Proposition 6 with  $E$  and  $F$  defined in the following way. We let  $E$  be the space of the functions

$$(y, p, v) \in L^2(Q)^3 \times L^2(Q) \times L^2(Q)$$

such that

1.  $e^{\frac{\alpha K_1}{2(T-t)^5}} y \in L^\infty((T/2, T), H^1(\Omega))^3 \cap L^2((T/2, T), H^2(\Omega)^3)$ ,
2.  $\nabla \cdot y = 0$  in  $Q$ ,
3.  $\nabla p \in L^2(Q)$ ,
4.  $e^{\frac{\alpha K_1}{2(T-t)^5}} v \in L^2(Q)^3$  and  $v = 0$  on  $Q \setminus [T/2, T] \times \omega_0$ ,
5.  $e^{\frac{K_1}{2r(T-t)^5}} (y_t - \Delta y + (\bar{y} \cdot \nabla)y + (y \cdot \nabla)\bar{y} + \nabla p - (0, 0, v)) \in L^2(Q)^3$ ,
6.  $y(0, \cdot) \in V$ ,

equipped with the following norm which makes it a Banach space:

$$\begin{aligned} \|(y, p, v)\|_E &= \|e^{\frac{\alpha K_1}{2(T-t)^5}} y\|_{L^\infty((0, T), H^1(\Omega)^3) \cap L^2((0, T), H^2(\Omega)^3)} + \|p\|_{L^2(Q)} \\ &+ \|e^{\frac{\alpha K_1}{2(T-t)^5}} v\|_{L^2(Q)^3} + \|e^{\frac{K_1}{2r(T-t)^5}} (y_t - \Delta y + (\bar{y} \cdot \nabla)y + (y \cdot \nabla)\bar{y} + \nabla p - (0, 0, v))\|_{L^2(Q)^3} + \|y(0, \cdot)\|_{H^1(\Omega)^3}. \end{aligned}$$

We let  $F$  be the space of the functions  $(h, y^0) \in L^2(Q)^3 \times V$  such that

$$e^{\frac{\alpha K_1}{2(T-t)^5}} h \in L^2(Q)^3,$$

equipped with the following scalar product which makes it a Hilbert space:

$$((h, y^0)|(k, z^0)) = (e^{\frac{\alpha K_1}{2(T-t)^5}} h | e^{\frac{\alpha K_1}{2(T-t)^5}} k)_{L^2(Q)^3} + (y^0 | z^0)_{H^1(Q)^3}.$$

We define

$$\mathcal{A}(y, p, v) = (y_t - \Delta y + (y \cdot \nabla)y + (\bar{y} \cdot \nabla)y + (y \cdot \nabla)\bar{y} + \nabla p - (0, 0, v), y(0, \cdot)).$$

To apply the previous inverse mapping theorem, we first show the following lemma.

**Lemma 4.** *The map  $\mathcal{A}$  has its image included in  $F$  and is of class  $C^1$  on  $E$ .*

**Proof of Lemma 4.**

We see that  $\mathcal{A} = \mathcal{B} + \mathcal{C}$  with

$$\mathcal{B}(y, p, v) := (y_t - \Delta y + (\bar{y} \cdot \nabla)y + (y \cdot \nabla)\bar{y} + \nabla p - (0, 0, v), y(0, \cdot)).$$

and

$$\mathcal{C}(y, p, v) := ((y \cdot \nabla)y, 0).$$

Thanks to the construction of  $E$  and  $F$  we have  $\mathcal{B} : E \rightarrow F$  and  $\mathcal{B}$  is continuous, so, since  $\mathcal{B}$  is linear,  $\mathcal{B}$  is of class  $C^1$ . The map  $\mathcal{C}$  is a quadratic form, hence to prove that it maps  $E$  into  $F$  and is of class  $C^1$ , it is sufficient to prove that it is continuous, i.e. to prove that

$$\|e^{\frac{K_1}{2r(T-t)^5}}(y \cdot \nabla)y\|_{L^2(Q_0)^3} \leq C\|(y, p, v)\|_E^2. \quad (4.1)$$

We choose  $r$  and  $\alpha$  so that

$$\frac{K_1}{r} < 2\alpha K_1. \quad (4.2)$$

(One can take for example  $\alpha = 3/4$  and then  $r \in (0, 1)$  close enough to 1.) Let us call

$$\tilde{y}(t, x) := e^{\frac{K_1}{4r(T-t)^5}} y. \quad (4.3)$$

This definition of  $\tilde{y}$  and inequality (4.2) imply that

$$\|\tilde{y}\|_{L^\infty((0,T), H^1(\Omega)^3)} \leq C\|(y, p, v)\|_E,$$

which gives that

$$\|\nabla(\tilde{y})\|_{L^\infty((0,T), L^2(\Omega)^9)} \leq C\|(y, p, v)\|_E. \quad (4.4)$$

We also have

$$\|\tilde{y}\|_{L^2((0,T), H^2(\Omega)^3)} \leq C\|(y, p, v)\|_E. \quad (4.5)$$

A classical Sobolev inequality in dimension 3 together with (4.5) imply that

$$\|\tilde{y}\|_{L^2((0,T), L^\infty(\Omega)^3)} \leq C\|(y, p, v)\|_E. \quad (4.6)$$

Direct computations imply that

$$\|(\tilde{y} \cdot \nabla)\tilde{y}\|_{L^2((0,T), L^2(\Omega)^3)} \leq \|\nabla(\tilde{y})\|_{L^\infty((0,T), L^2(\Omega)^9)} \|\tilde{y}\|_{L^2((0,T), L^\infty(\Omega)^3)}. \quad (4.7)$$

From (4.4), (4.6) and (4.7), we obtain

$$\|(\tilde{y} \cdot \nabla)\tilde{y}\|_{L^2((0,T), L^2(\Omega)^3)} \leq C\|(y, p, v)\|_E^2,$$

which, together with (4.3), gives (4.1). ■

We now consider the element  $e_0 = (0, 0, 0)$  and we compute

$$d\mathcal{A}(e_0)(y, q, v) = y_t - \Delta y + (\bar{y} \cdot \nabla)y + (y \cdot \nabla)\bar{y} + \nabla p - (0, 0, v).$$

The application of Proposition 5 we gave in the previous subsection implies that this application is surjective. Hence, taking  $g = (0, y^0)$  and applying Proposition 6, Theorem 1 easily follows (notably because the trajectory  $\bar{y}$  can be chosen as small as we want since  $\varepsilon$  can be arbitrarily chosen small).

## A Appendix: Creation of the matrix $L_0$

In this appendix we explain how the matrix  $L_0$  at point  $\xi^0$  which represents all the differentiated equations of System (3.25) up to the order 19 was created. The program is written in  $C^{++}$ , using the library uBLAS which is well-adapted to the manipulation of sparse matrices. It is a parallel openMP algorithm, using 8 cores. We are not going to give all the technical details but just explain rapidly the spirit of the algorithm. To simplify, we will assume that the following “black boxes” (that had to be created) are at our disposal:

1. An evaluation function `ep` which evaluates a polynomial (represented by a vector) at  $\xi^0$ . This evaluation function can be created so that it can verify that  $\xi^0$  is not a root of the polynomial  $P(0, \cdot, \cdot)$ . (one just has to see if the evaluation is equal to 0 whereas the polynomial has nonzero coefficients).
2. A derivation function `deqex` which differentiates an equation of level  $m$  with respect to  $x_1, x_2, x_3$  or  $t$ .

A partial differential equation which is a derivative of order  $m$  of some of the equations of (3.25) will be represented in a matricial form in the following way: We know that there are at most  $F(n+3)$  derivatives appearing, and we observe that the coefficients are polynomials in  $(x_1, x_2, x_3)$  of an order less than 4 (it is a vector space of dimension 35). Hence the equation is represented by a matrix with  $F(n+3)$  lines and 35 columns, where on each line one can find the coefficient of the partial derivatives of  $z^1$  (or  $z^2$  appearing) corresponding to the number of this line, thanks to a natural bijection between  $\mathbb{N}^4$  and  $\mathbb{N}$ . Since we have 3 equations in (3.25) and 2 unknowns ( $z^1$  and  $z^2$ ), one can write the matrix  $M$  in the following way:

$$\begin{pmatrix} A_1 & B_1 \\ A_2 & B_2 \\ A_3 & B_3 \end{pmatrix}$$

For  $i = 1, 2, 3$ ,  $A_i$  represents the derivatives of  $z^1$  appearing in the derivatives of the  $i$ -th equation of (3.25) and  $B_i$  those of  $z^2$ . Hence, we can compute these  $A_i$  and  $B_i$  separately and then gather them to obtain  $L_0^0$ .

The algorithm is the following. We explain it for the first equation of (3.25) and for the unknown  $z^1$  (i.e. for  $A_1$ , but it is the same for the other matrices).

1. We create a matrix  $e$  that represents the equation. We use  $ep$  to fill the line of  $L_0^0$  corresponding to the equation in a .txt file under the form  $i\ j\ A_1(i, j)$ . We create a matrix  $h$  which is empty for the moment. In fact in  $e$  we will keep the equations of level  $m - 1$  and in  $h$  we will fill the equations of level  $m$ .
2. We create a “for” loop on  $m$  which will represent the level of equations we are creating. The integer  $m$  goes from 1 to 19 since we differentiate 19 times at most.
3. We create a second “for” loop in the interior of the first loop on a number  $n$  which represents one of the equations of level  $m$ . Thanks to the definition of the function  $F$  given in Subsection 3.2.2, we have  $F(m - 1) + 1 \leq n \leq F(m)$ . If  $m = 1$ , then  $n$  goes from  $F(0) + 1 = 2$  to  $F(1) = 5$  ( $n$  represents  $\partial_1, \partial_2, \partial_3$  or  $\partial_t$ ). If  $m = 2$ , then  $n$  goes from  $F(1) + 1 = 6$  to  $F(2) = 15$  ( $n$  represents  $\partial_{11}^2, \partial_{12}^2, \partial_{13}^2, \partial_{1t}^2, \partial_{22}^2, \partial_{23}^2, \partial_{2t}^2, \partial_{33}^2, \partial_{3t}^2$  or  $\partial_{tt}^2$ ), etc. This loop is parallelized on our 8 cores. In this loop, we want to create the  $n$ -th equation denoted  $E_n$ , which is of level  $m$ . Hence we take a suitable equation of level  $m - 1$  denoted  $E_r$  which is so that if we differentiate  $E_r$  with respect to 1, 2, 3 or  $t$ , we obtain  $E_n$ . For example, if we consider  $m = 2$  and we want to obtain the first equation of (3.25) differentiated two times with respect to 1, then we consider the equation  $E_r$  to be the first equation of (3.25) differentiated one time with respect to 1 and differentiated with respect to 1 to obtain  $E_n$ .
4. Once the loop on  $n$  is ended, we have in our matrix  $e$  all the equations of level  $m - 1$  and in  $h$  we have just created all the equations of level  $m$ . Now we just have to use our evaluation function  $ep$  on  $h$  to obtain the coefficients of the lines of  $A_1$  corresponding to the equations that are of level  $m$ , i.e. the equations numbered from  $F(m - 1) + 1$  to  $F(m)$ . We write these coefficients in our .txt file in the form  $i\ j\ A_1(i, j)$ .
5. We update now  $e$ , take  $e = h$ , we empty  $h$  and we can go to the following loop  $m + 1$ .

At the end we have created a file containing the coefficients of a sparse matrix  $A_1$  of size (8855, 14950). Using the same program with  $z^2$  and the two others equation we obtain five other files representing five matrices that we gather to obtain the matrix  $L_0(\xi^0) = L_0^0$ . Our matrix  $L_0^0$ , which represents all the equations, is of size (30360, 29900) and has 651128 nonzero coefficients. Only 0.0717% of the coefficients are different from 0, with an average of 21.45 nonzero coefficients on each row, which is logical since we are working with coefficients that are polynomials of small degree, so we do not create many terms on each line when we differentiate the equations. In Figure A, one can observe how the nonzero coefficients of  $L_0^0$  are distributed.

## B acknowledgements\*

The authors would like to thank Sergio Guerrero for fruitful discussions concerning Proposition 4.




Figure 3: Distribution of the nonzero coefficients of  $L_0^0$ .

## References

- [1] Fatiha Alabau-Boussouira and Matthieu Léautaud. Indirect controllability of locally coupled wave-type systems and applications. *J. Math. Pures Appl.*
- [2] Vladimir Mi. Alekseev, Vladimir Mi. Tikhomirov, and Sergej Vl. Fomin. *Optimal control*. Contemporary Soviet Mathematics. Consultants Bureau, New York, 1987. Translated from the Russian by V. M. Volosov.
- [3] Farid Ammar-Khodja, Assia Benabdallah, Cédric Dupaix, and Manuel González-Burgos. A generalization of the Kalman rank condition for time-dependent coupled linear parabolic systems. *Differ. Equ. Appl.*, 1(3):427–457, 2009.
- [4] Farid Ammar-Khodja, Assia Benabdallah, Manuel González-Burgos, and Luz de Teresa. The kalman condition for the boundary controllability of coupled parabolic systems. bounds on biorthogonal families to complex matrix exponentials. *Journal de Mathématiques Pures et Appliquées*, 96(6):555 – 590, 2011.
- [5] Farid Ammar-Khodja, Assia Benabdallah, Manuel González-Burgos, and Luz de Teresa. Recent results on the controllability of linear coupled parabolic problems: A survey. *Mathematical Control and Related Fields*, 1(3):267–306, 2011.

- [6] Nicolás Carreño and Sergio Guerrero. Local null controllability of the  $n$ -dimensional Navier-Stokes system with  $n-1$  scalar controls in an arbitrary control domain. *To appear in Journal of Mathematical Fluid Mechanics*, 2012.
- [7] Jean-Michel Coron. Global asymptotic stabilization for controllable systems without drift. *Math. Control Signals Systems*, 5(3):295–312, 1992.
- [8] Jean-Michel Coron. Contrôlabilité exacte frontière de l'équation d'Euler des fluides parfaits incompressibles bidimensionnels. *C. R. Acad. Sci. Paris Sér. I Math.*, 317(3):271–276, 1993.
- [9] Jean-Michel Coron. On the controllability of the 2-D incompressible Navier-Stokes equations with the Navier slip boundary conditions. *ESAIM Control Optim. Calc. Var.*, 1:35–75 (electronic), 1995/96.
- [10] Jean-Michel Coron. *Control and nonlinearity*, volume 136 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2007.
- [11] Jean-Michel Coron and Andrei V. Fursikov. Global exact controllability of the 2D Navier-Stokes equations on a manifold without boundary. *Russian J. Math. Phys.*, 4(4):429–448, 1996.
- [12] Jean-Michel Coron and Sergio Guerrero. Local null controllability of the two-dimensional Navier-Stokes system in the torus with a control force having a vanishing component. *J. Math. Pures Appl. (9)*, 92(5):528–545, 2009.
- [13] Jean-Michel Coron and Sergio Guerrero. Null controllability of the  $N$ -dimensional Stokes system with  $N-1$  scalar controls. *J. Differential Equations*, 246(7):2908–2921, 2009.
- [14] Andrew Ll. Dulmage and Nathan Sa. Mendelsohn. Coverings of bipartite graphs. *Canad. J. Math.*, 10:517–534, 1958.
- [15] Enrique Fernández-Cara, Sergio Guerrero, Oleg Yu. Imanuvilov, and Jean-Pierre Puel. Local exact controllability of the Navier-Stokes system. *J. Math. Pures Appl. (9)*, 83(12):1501–1542, 2004.
- [16] Enrique Fernández-Cara, Sergio Guerrero, Oleg Yu. Imanuvilov, and Jean-Pierre Puel. Some controllability results for the  $N$ -dimensional Navier-Stokes and Boussinesq systems with  $N-1$  scalar controls. *SIAM J. Control Optim.*, 45(1):146–173 (electronic), 2006.
- [17] Andrei Vl. Fursikov and Oleg Yu. Imanuvilov. *Controllability of evolution equations*, volume 34 of *Lecture Notes Series*. Seoul National University Research Institute of Mathematics Global Analysis Research Center, Seoul, 1996.
- [18] Andrei Vl. Fursikov and Oleg Yu. Imanuvilov. Exact controllability of the Navier-Stokes and Boussinesq equations. *Russian Math. Surveys*, 54:565–618, 1999.

- [19] Mikhael Gromov. *Partial differential relations*, volume 9 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1986.
- [20] Mamadou Gueye. Insensitizing controls for the Navier-Stokes equations. *To appear in Annales de l'IHP*, 2012.
- [21] Oleg Yu. Imanuvilov. On exact controllability for the Navier-Stokes equations. *ESAIM Control Optim. Calc. Var.*, 3:97–131 (electronic), 1998.
- [22] Oleg Yu. Imanuvilov. Remarks on exact controllability for the Navier-Stokes equations. *ESAIM Control Optim. Calc. Var.*, 6:39–72 (electronic), 2001.
- [23] Jacques-Louis Lions and Enrico Magenes. *Problèmes aux limites non homogènes et applications. Vol. 1*. Travaux et Recherches Mathématiques, No. 17. Dunod, Paris, 1968.
- [24] Jacques-Louis Lions and Enrique Zuazua. A generic uniqueness result for the Stokes system and its control theoretical consequences. In *Partial differential equations and applications*, volume 177 of *Lecture Notes in Pure and Appl. Math.*, pages 221–235. Dekker, New York, 1996.
- [25] Karine Mauffrey. On the null controllability of a  $3 \times 3$  parabolic system with non constant coefficients by one or two control forces. *J. Math. Pures Appl.*, in press.
- [26] Alex Pothén and Chin-Ju Fan. Computing the block triangular form of a sparse matrix. *ACM Trans. Math. Software*, 16(4):303–324, 1990.
- [27] Roger Temam. *Navier-Stokes equations*, volume 2 of *Studies in Mathematics and its Applications*. North-Holland Publishing Co., Amsterdam, revised edition, 1979. Theory and numerical analysis, With an appendix by F. Thomasset.