



HAL
open science

About Pillai and Fermat-Catalan conjectures

Jamel Ghannouchi

► **To cite this version:**

| Jamel Ghannouchi. About Pillai and Fermat-Catalan conjectures. 2014. ⟨hal-00751215v12⟩

HAL Id: hal-00751215

<https://hal.science/hal-00751215v12>

Preprint submitted on 13 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

About Pillai and Fermat-Catalan conjectures

Jamel Ghanouchi

Ecole Supérieure des Sciences et Techniques de Tunis

jamel.ghanouchi@topnet.tn

Abstract

(MSC=11D04) We begin with Fermat-Catalan equation $Z^c = X^a + Y^b > 2Y^b$ and see if it has solutions for $q + 1 = 2p$ or $q + 2 = 2p$ or $q = 2p$.

(Keywords : Diophantine equations, Fermat-Catalan equation ; Approach)

An attempt of resolution of Fermat-Catalan and Pillai equations

Let Fermat-Catalan equation

$$Y^p = X^q + Z^a > 2Z^a$$

The truth is Fermat-Catalan consists in three equations : the one higher and $Y^p = Z^a - X^q$ and $Y^p = X^q - Z^a$, we will treat only the first as the others have the same solutions by the same reasoning. Then let $Z^a = 1_a 1_a$. And Pillai one

$$Y^p = X^q + a = X^q + 1_a 1_a$$

Thus

$$Y^p = X^q + 1_a 1_a$$

We suppose firstly that

$$c = \frac{X^p - 1_a}{Y^{\frac{p}{2}}} > 0, \quad c' = \frac{7_a - X^p}{Y^{\frac{p}{2}}} < 0$$

Hence

$$Y^{\frac{p}{2}} = \frac{6_a}{c + c'}$$

$$X^p = cY^{\frac{p}{2}} + 1_a = \frac{6_a c}{c + c'} + 1_a = \frac{7_a c + 1_a c'}{c + c'}$$

And

$$X^q = Y^p - 1_a 1_a = \frac{36_a 1_a - 1_a 1_a (c + c')^2}{(c + c')^2}$$

As $X^p > 7_a$ then $c' < 0$ and $c > c'$ but $Y^{\frac{p}{2}} \geq 3$ then $0 < c + c' \leq 2_a$ Two cases $c^2 > 1$ et $0 < c^2 \leq 1$

$c^2 > 1$ We have

$$X^q - X^{2p} = \frac{36_a 1_a - 1_a 1_a (c + c')^2 - (7_a c + 1_a c')^2}{(c + c')^2}$$

$$< \frac{36_a 1_a - 36_a 1_a c^2}{(c + c')^2} < 0$$

Or

$$q + 1 \leq 2p$$

But

$$c' Y^{\frac{p}{2}} = 7_a - X^p < -Y^{\frac{p}{2}}$$

Then $c' < -1$. But, let u, v verifying

$$\frac{7uc + vc'}{c + c'} = 4$$

Or

$$(7u - 4)c = (4 - v)c' > 0$$

(The reasoning is the same if it is negative). As $c > 0$ and $c' < 0$ it means $7u - 4 > 0$ and $4 - v < 0$. Then

$$(7u - 4)(X^p - 1_a) = (4 - v)(7_a - X^p)$$

Or

$$(7u - v)X^p = 24_a + 7_a(u - v)$$

Hence

$$X^p = \frac{7_a(u - v) + 24_a}{7u - v}$$

But

$$c + c' = \frac{4 - v}{7u - 4}c' + c' = \frac{7u - v}{7u - 4}c'$$

Then

$$Y^{\frac{p}{2}} = \frac{6_a}{c + c'} = \frac{6_a(7u - 4)}{(7u - v)c'}$$

And

$$\frac{X^p}{Y^{\frac{p}{2}}} = \frac{(7_a(u - v) + 24_a)c'}{6_a(7u - 4)} = \frac{(7(u - v) + 24)c'}{6(7u - 4)}$$

But as $X^p > 28_a$ (else X, p can be calculated and Pillai equation has a finite number of solutions!) Thus

$$(9c' + 7c)Y^{\frac{p}{2}} = 9(7_a - X^p) + 7(X^p - 1_a) = 56_a - 2X^p < 0 \Rightarrow 9c' + 7c < 0$$

And

$$(4 - v)(9c' + 7c) > 0$$

Thus

$$9(7u - 4)c = 9(4 - v)c' > 7(v - 4)c$$

It means

$$7(7u - 4) > 7v - 14u - 20 > 7(4 - v)c' + 2(4 - 7u)$$

But if $X^p < X^{\frac{q+1}{2}} + 7 < X^{\frac{q+2}{2}}$ then $X^{q+1} \leq X^{2p} < X^{q+2}$ else

$$\begin{aligned} X^q(X - 49) > X^q > 49a1_a &\Rightarrow (X^p - 7_a)^2 - 49a1_a \geq X^{q+1} - 49a1_a > 49X^q \\ \Rightarrow (X^p - 7_a)^2 &\geq 49(X^q + 1_a1_a) = 49Y^p \Rightarrow X^p - 7_a \geq 7Y^{\frac{p}{2}} \end{aligned}$$

And

$$\begin{aligned} X^p - 7_a &> 7Y^{\frac{p}{2}} > Y^{\frac{p}{2}} \left(\frac{7(v-4) + 2(4-7u)}{7u-4} \right) \\ &> Y^{\frac{p}{2}} \left(\frac{7(4-v)c' + 2(4-7u)}{7u-4} \right) \end{aligned}$$

Then

$$-7_a + X^p + 2Y^{\frac{p}{2}} > \left(\frac{7(4-v)c'}{7u-4} \right) Y^{\frac{p}{2}}$$

Or

$$\begin{aligned} &\frac{-7_a + X^p + 2Y^{\frac{p}{2}}}{6c'} \\ &= \left(\frac{2}{6c'} - \frac{1}{6} \right) Y^{\frac{p}{2}} < \frac{7(4-v)}{6(7u-4)} Y^{\frac{p}{2}} \end{aligned}$$

Or

$$\left(\frac{2}{6c'} - \frac{1}{6} \right) c' > \frac{7(4-v)c'}{6(7u-4)}$$

And

$$\begin{aligned} &\frac{c'}{6} + \frac{7(4-v)c'}{6(7u-4)} \\ &= \frac{(7(u-v) + 24)c'}{6(7u-4)} = \frac{X^p}{Y^{\frac{p}{2}}} < \frac{2}{6} < 1 \end{aligned}$$

Thus

$$X^{q+1} \leq X^{2p} < Y^p = X^q + 1_a1_a < X^{q+1}$$

It is impossible! Thus $q + 1 = 2p$ or $q + 2 = 2p$

$$0 < c^2 \leq 1$$

$$\begin{aligned} 2X^q - X^{2p} &= \frac{72a1_a - 2a1_a(c+c')^2 - (7ac + 1ac')^2}{(c+c')^2} \\ &> 1_a1_a \frac{64 - 49c^2}{(c+c')^2} > 0 \end{aligned}$$

Or

$$X^q \geq X^{2p-1}$$

Or

$$q + 1 \geq 2p$$

We have

$$c'Y^{\frac{p}{2}} = 7a - X^p > -Y^{\frac{p}{2}}$$

Thus $c' > -1$. But, let u, v verifying

$$\frac{7uc + vc'}{c + c'} = 4$$

Or

$$(7u - 4)c = (4 - v)c' > 0$$

As $c > 0$ and $c' < 0$ it means $7u - 4 > 0$ and $4 - v < 0$. Then $7u - 4 > 0$ and $4 - v < 0$ and

$$(7u - 4)(X^p - 1_a) = (4 - v)(7a - X^p)$$

Or

$$(7u - v)X^p = 24a + 7a(u - v)$$

Hence

$$X^p = 1_a \frac{7(u - v) + 24}{7u - v}$$

But

$$c + c' = \frac{4 - v}{7u - 4}c' + c' = \frac{7u - v}{7u - 4}c'$$

Then

$$Y^{\frac{p}{2}} = \frac{6a}{c + c'} = \frac{6a(7u - 4)}{(7u - v)c'}$$

And

$$\frac{X^p}{Y^{\frac{p}{2}}} = \frac{(7(u - v) + 24)c'}{6(7u - 4)}$$

But

$$(3c' + 7c)Y^{\frac{p}{2}} = 3(7a - X^p) + 7(X^p - 1_a) = 4X^p + 14a > 0 \Rightarrow 3c' + 7c > 0$$

And

$$(4 - v)(3c' + 7c) < 0$$

Or

$$3(7u - 4)c = 3(4 - v)c' < 7(v - 4)c$$

It means

$$7u - 4 < 7v - 14u - 20 < 7(4 - v)c' + 2(4 - 7u)$$

But

$$X^q + 1_a 1_a = Y^p \geq X^{2p} + 1_a 1_a > (X^p - 7a)^2$$

Thus

$$\begin{aligned} X^p - 7a &< Y^{\frac{p}{2}} < Y^{\frac{p}{2}} \left(\frac{7(v - 4) + 2(4 - 7u)}{7u - 4} \right) \\ &< Y^{\frac{p}{2}} \left(\frac{7(4 - v)c' + 2(4 - 7u)}{7u - 4} \right) \end{aligned}$$

Then

$$-7a + X^p + 2Y^{\frac{p}{2}} < \left(\frac{7(4 - v)c'}{7u - 4} \right) Y^{\frac{p}{2}}$$

Or

$$\begin{aligned}\frac{-7a + X^p + 2Y^{\frac{p}{2}}}{6c'} &= \left(\frac{2}{6c'} - \frac{1}{6}\right)Y^{\frac{p}{2}} \\ &> \frac{7(4-v)}{6(7u-4)}Y^{\frac{p}{2}}\end{aligned}$$

Or

$$\left(\frac{2}{6c'} - \frac{1}{6}\right)c' < \frac{7(4-v)c'}{6(7u-4)}$$

And

$$\begin{aligned}\frac{c'}{6} + \frac{7(4-v)c'}{6(7u-4)} \\ = \frac{(7(u-v) + 24)c'}{6(7u-4)} = \frac{X^p}{Y^{\frac{p}{2}}} > \frac{2}{6}\end{aligned}$$

Thus

$$3X^{2p+1} > 36X^{2p} > 4Y^p > 3X^q$$

It means that finally $q = 2p$ or $q + 1 = 2p$ or $q + 2 = 2p$. We suppose now that

$$c = \frac{X^p - 1_a}{Y^{\frac{p}{2}}} < 0, \quad c' = \frac{7a - X^p}{Y^{\frac{p}{2}}} > 0$$

Hence

$$Y^{\frac{p}{2}} = \frac{6a}{c + c'} > 3$$

Thus

$$0 < c + c' < 2a$$

And

$$X^p = cY^{\frac{p}{2}} + 1_a = \frac{6ac}{c + c'} + 1_a = \frac{7ac + 1_ac'}{c + c'}$$

And

$$X^q = Y^p - 1_a 1_a = \frac{36a 1_a - 1_a 1_a (c + c')^2}{(c + c')^2}$$

As $X^p < 1_a < 7a$ then $c < 0$ and $c' > -c > c$ but $Y^{\frac{p}{2}} \geq 3$ then $0 < c + c' \leq 2a$ Two cases $c'^2 > 1$ et $0 < c'^2 \leq 1$

$c'^2 > 1$ We have

$$c^2 Y^p = (X^p - 1_a)^2 > (X^p - 7a)^2 = c'^2 Y^p > Y^p = X^q + 1_a 1_a > (X^{\frac{q}{2}} - 1_a)^2$$

Or

$$q + 1 \leq 2p$$

But if $7a < X^p + Y^{\frac{p}{2}}$ then $7a < 3X^p < 21a$ and X, p can be calculated and Pillai equation has a finite number of solutions, else

$$c'Y^{\frac{p}{2}} = 7a - X^p > Y^{\frac{p}{2}}$$

Then $-c' < -1$. But, let u, v verifying

$$\frac{7uc + vc'}{c + c'} = 4$$

Or

$$(7u - 4)c = (4 - v)c' > 0$$

As $c < 0$ and $c' > 0$ it means $7u - 4 < 0$ and $4 - v > 0$. Then

$$(7u - 4)(X^p - 1_a) = (4 - v)(7_a - X^p)$$

Or

$$(7u - v)X^p = 24_a + 7_a(u - v)$$

Hence

$$X^p = \frac{7_a(u - v) + 24_a}{7u - v}$$

But

$$c + c' = \frac{4 - v}{7u - 4}c' + c' = \frac{7u - v}{7u - 4}c'$$

Then

$$Y^{\frac{p}{2}} = \frac{6_a}{c + c'} = \frac{6_a(7u - 4)}{(7u - v)c'}$$

And

$$\frac{X^p}{Y^{\frac{p}{2}}} = \frac{(7_a(u - v) + 24_a)c'}{6_a(7u - 4)} = \frac{(7(u - v) + 24)c'}{6(7u - 4)}$$

But

$$(9c' + 7c)Y^{\frac{p}{2}} = 9(7_a - X^p) + 7(X^p - 1_a) = 56_a - 2X^p > 0 \Rightarrow 9c' + 7c > 0$$

And

$$(4 - v)(9c' + 7c) > 0$$

Thus

$$9(7u - 4)c = 9(4 - v)c' > 7(v - 4)c$$

It means

$$7(7u - 4) > 7v - 14u - 20 > 7(4 - v)c' + 2(4 - 7u)$$

But if $X^p < X^{\frac{q+1}{2}} + 7 < X^{\frac{q+2}{2}}$ then $X^{q+1} \leq X^{2p} < X^{q+2}$ else

$$\begin{aligned} X^q(X - 49) > X^q > 49_a 1_a &\Rightarrow (X^p - 7_a)^2 - 49_a 1_a \geq X^{q+1} - 49_a 1_a > 49X^q \\ \Rightarrow (X^p - 7_a)^2 &\geq 49(X^q + 1_a 1_a) = 49Y^p \Rightarrow X^p - 7_a \geq 7Y^{\frac{p}{2}} \end{aligned}$$

And

$$\begin{aligned} X^p - 7_a &> 7Y^{\frac{p}{2}} > Y^{\frac{p}{2}} \left(\frac{7(v - 4) + 2(4 - 7u)}{7u - 4} \right) \\ &> Y^{\frac{p}{2}} \left(\frac{7(4 - v)c' + 2(4 - 7u)}{7u - 4} \right) \end{aligned}$$

Then

$$-7_a + X^p + 2Y^{\frac{p}{2}} > \left(\frac{7(4 - v)c'}{7u - 4} \right) Y^{\frac{p}{2}}$$

Or

$$\frac{-7a + X^p + 2Y^{\frac{p}{2}}}{6c'} \\ = \left(\frac{2}{6c'} - \frac{1}{6}\right)Y^{\frac{p}{2}} < \frac{7(4-v)}{6(7u-4)}Y^{\frac{p}{2}}$$

Or

$$\left(\frac{2}{6c'} - \frac{1}{6}\right)c' > \frac{7(4-v)c'}{6(7u-4)}$$

And

$$\frac{c'}{6} + \frac{7(4-v)c'}{6(7u-4)} \\ = \frac{(7(u-v) + 24)c'}{6(7u-4)} = \frac{X^p}{Y^{\frac{p}{2}}} < \frac{2}{6} < 1$$

Thus

$$X^{q+1} \leq X^{2p} < Y^p = X^q + 1_a 1_a < X^{q+1}$$

It is impossible! Thus $q+1 = 2p$ or $q+2 = 2p$

$$0 < c'^2 \leq 1$$

$$(X^p - 7a)^2 < Y^p = X^q + 1_a 1_a$$

Or

$$X^q \geq X^{2p-1}$$

Or

$$q+1 \geq 2p$$

We have

$$c'Y^{\frac{p}{2}} = 7a - X^p > -Y^{\frac{p}{2}}$$

Thus $c' > -1$. But, let u, v verifying

$$\frac{7uc + vc'}{c + c'} = 4$$

Or

$$(7u-4)c = (4-v)c' > 0$$

As $c < 0$ and $c' > 0$ it means $7u-4 < 0$ and $4-v > 0$. And

$$(7u-4)(X^p - 1_a) = (4-v)(7a - X^p)$$

Or

$$(7u-v)X^p = 24a + 7a(u-v)$$

Hence

$$X^p = 1_a \frac{7(u-v) + 24}{7u-v}$$

But

$$c + c' = \frac{4-v}{7u-4}c' + c' = \frac{7u-v}{7u-4}c'$$

Then

$$Y^{\frac{p}{2}} = \frac{6a}{c + c'} = \frac{6a(7u - 4)}{(7u - v)c'}$$

And

$$\frac{X^p}{Y^{\frac{p}{2}}} = \frac{(7(u - v) + 24)c'}{6(7u - 4)}$$

But

$$(3c' + 7c)Y^{\frac{p}{2}} = 3(7a - X^p) + 7(X^p - 1a) = 4X^p + 14a > 0 \Rightarrow 3c' + 7c > 0$$

And

$$(4 - v)(3c' + 7c) > 0$$

Or

$$3(7u - 4)c = 3(4 - v)c' > 7(v - 4)c$$

It means $3(7u - 4) < 7(v - 4)$ and

$$7u - 4 < 7v - 14u - 20 < 7(4 - v)c' + 2(4 - 7u)$$

But

$$X^q + 1a1a = Y^p \geq X^{2p} + 1a1a > (X^p - 7a)^2$$

Thus

$$\begin{aligned} X^p - 7a &< Y^{\frac{p}{2}} < Y^{\frac{p}{2}} \left(\frac{7(v - 4) + 2(4 - 7u)}{7u - 4} \right) \\ &< Y^{\frac{p}{2}} \left(\frac{7(4 - v)c' + 2(4 - 7u)}{7u - 4} \right) \end{aligned}$$

Then

$$-7a + X^p + 2Y^{\frac{p}{2}} < \left(\frac{7(4 - v)c'}{7u - 4} \right) Y^{\frac{p}{2}}$$

Or

$$\begin{aligned} \frac{-7a + X^p + 2Y^{\frac{p}{2}}}{6c'} &= \left(\frac{2}{6c'} - \frac{1}{6} \right) Y^{\frac{p}{2}} \\ &> \frac{7(4 - v)}{6(7u - 4)} Y^{\frac{p}{2}} \end{aligned}$$

Or

$$\left(\frac{2}{6c'} - \frac{1}{6} \right) c' < \frac{7(4 - v)c'}{6(7u - 4)}$$

And

$$\begin{aligned} &\frac{c'}{6} + \frac{7(4 - v)c'}{6(7u - 4)} \\ &= \frac{(7(u - v) + 24)c'}{6(7u - 4)} = \frac{X^p}{Y^{\frac{p}{2}}} > \frac{2}{6} \end{aligned}$$

Thus

$$3X^{2p+1} > 36X^{2p} > 4Y^p > 3X^q$$

It means that finally $q = 2p$ or $q + 1 = 2p$ or $q + 2 = 2p$. We suppose finally

$$c = \frac{X^p - 1a}{Y^{\frac{p}{2}}} > 0, \quad c' = \frac{7a - X^p}{Y^{\frac{p}{2}}} > 0$$

Hence

$$Y^{\frac{p}{2}} = \frac{6a}{c+c'}$$

$$X^p = cY^{\frac{p}{2}} + 1_a = \frac{6ac}{c+c'} + 1_a = \frac{7ac + 1_ac'}{c+c'}$$

And

$$X^q = Y^p - 1_a 1_a = \frac{36a1_a - 1_a 1_a (c+c')^2}{(c+c')^2}$$

As $1_a < X^p < 7a$ then X, p can be calculated and Pillai equation has a finite number of solutions!

main result and relation with Fermat and Catalan equations It means that finally $q = 2p$ or $q + 1 = 2p$ or $q + 2 = 2p$. We see that for Catalan equation (as q and $2p$ do not have the same parity) $q + 1 = 2p$ (for this equation, we have $c > 0$ and $c' < 0$) and for Fermat equation there are only two solutions $q + 2 = 2p = 2n = n + 2 = 4$ or $q + 1 = 2p = n + 1 = 2n = 2$. We will prove now that there is only one value of p for the solution, it is $p = 2$. If the exponent of X is even, we must do as it follows

$$Y^p = X^{2p-2u} + X + 1_a 1_a - X = (X+1)X(X^{2p-2u-2} - X^{2p-2u-3} + \dots - X+1) + 1_a 1_a - X = uX(X+1) + 1_a 1_a - X$$

We have also

$$Y^p = X^{2p-2u} - X + 1_a 1_a + X = (X-1)X(X^{2p-2u-2} + X^{2p-2u-3} + \dots + X+1) + 1_a 1_a + X = vX(X-1) + 1_a 1_a + X$$

Else if the exponent of X is odd and it will be the case that we will treat :

$$Y^p = X^{2p-1} + 1 + 1_a 1_a - 1 = (X+1)(X^{2p-2} - X^{2p-3} + \dots - X+1) + 1_a 1_a - 1 = u(X+1) + 1_a 1_a - 1$$

We have also

$$Y^p = X^{2p-2u} - 1 + 1_a 1_a + 1 = (X-1)(X^{2p-2} + X^{2p-3} + \dots + X+1) + 1_a 1_a + 1 = v(X-1) + 1_a 1_a + 1$$

Hence in all cases

$$2 = u(X+1) - v(X-1) = X+1 - (X-1)$$

And

$$2Y^p - 2_a 1_a = uX(X+1) + vX(X-1)$$

Thus

$$(u-1)(X+1) = (v-1)(X-1)$$

Thus

$$X = \frac{u+v-2}{v-u}$$

And

$$2Y^p - 2_a 1_a = (u+v)X + u - v = \frac{(u+v)(u+v-2)}{v-u} + u - v$$

$$= \frac{u^2 + v^2 + 2uv - 2u - 2v - u^2 - v^2 + 2uv}{v-u} = \frac{2u(v-1) + 2v(u-1)}{v-u}$$

Hence

$$Y^p = \frac{u(v-1) + v(u-1) + 1_a 1_a (v-u)}{v-u} = X^{2p-1} + 1_a 1_a$$

If we suppose p even

$$Y^p - Y = v'Y(Y-1) = X^{2p-1} + 1_a 1_a - Y$$

$$Y^p + Y = u'Y(Y+1) = X^{2p-1} + 1_a 1_a + Y$$

$$2Y = u'Y(Y+1) - v'Y(Y-1) = Y(Y+1) - Y(Y-1)$$

And

$$2X^{2p'-1} + 2_a 1_a = u'Y(Y+1) + v'Y(Y-1)$$

Or

$$(u'-1)(Y+1) = (v'-1)(Y-1)$$

Thus

$$Y = \frac{u' + v' - 2}{v' - u'}$$

And

$$2X^{2p-1} + 2_a 1_a = Y((u'+v')Y + (u'-v')) = \frac{u' + v' - 2}{v' - u'} \left(\frac{2u'(v'-1) + 2v'(u'-1)}{v' - u'} \right)$$

Or

$$Y^p = X^{2p-1} + 1_a 1_a = \frac{u' + v' - 2}{v' - u'} \left(\frac{u'(v'-1) + v'(u'-1)}{v' - u'} \right)$$

We have for equations with four unknowns

$$\begin{aligned} Y^p &= \left(\frac{u' + v' - 2}{v' - u'} \right)^p = X^{2p-1} + 1_a 1_a = \left(\frac{u + v - 2}{v - u} \right)^{2p-1} + 1_a 1_a \\ &= \frac{u' + v' - 2}{v' - u'} \left(\frac{u'(v'-1) + v'(u'-1)}{v' - u'} \right) = \frac{u(v-1) + v(u-1) + 1_a 1_a (v-u)}{v-u} \end{aligned}$$

The solutions of those equations are $u' = v' = 2$. Let us prove it : we have also

$$Y^p = Y^p - Y + Y = Y(Y-1)v' + Y = X^{2p-1} + 1_a 1_a = u(X+1) + 1_a 1_a - 1 \quad (1)$$

$$Y^p = Y^p - Y + Y = Y(Y-1)v' + Y = X^{2p-1} - 1 + 1_a 1_a + 1 = v(X-1) + 1_a 1_a + 1 \quad (2)$$

$$Y^p = Y^p + Y - Y = Y(Y+1)u' - Y = X^{2p-1} + 1 + 1_a 1_a - 1 = u(X+1) + 1_a 1_a - 1 \quad (3)$$

$$Y^p = Y^p + Y - Y = Y(Y+1)u' - Y = X^{2p-1} - 1 + 1_a 1_a + 1 = v(X-1) + 1_a 1_a + 1 \quad (4)$$

Hence, with

$$X = \frac{u + v - 2}{v - u}, \quad Y = \frac{u' + v' - 2}{v' - u'}$$

We have

$$(1) \quad v' \frac{u' + v' - 2}{v' - u'} \frac{2u' - 2}{v' - u'} + \frac{u' + v' - 2}{v' - u'} = u \frac{2v - 2}{v - u} + 1_a 1_a - 1$$

$$(2) \quad v' \frac{u' + v' - 2}{v' - u'} \frac{2u' - 2}{v' - u'} + \frac{u' + v' - 2}{v' - u'} = v \frac{2u - 2}{v - u} + 1_a 1_a + 1$$

$$(3) \quad u' \frac{u' + v' - 2}{v' - u'} \frac{2v' - 2}{v' - u'} - \frac{u' + v' - 2}{v' - u'} = u \frac{2v - 2}{v - u} + 1_a 1_a - 1$$

$$(4) \quad u' \frac{u' + v' - 2}{v' - u'} \frac{2v' - 2}{v' - u'} - \frac{u' + v' - 2}{v' - u'} = v \frac{2u - 2}{v - u} + 1_a 1_a + 1$$

Or

$$(1) \quad (v - u)(u' + v' - 2)(2v'(u' - 1) + v' - u') = (2u(v - 1) + (1_a 1_a - 1)(v - u))(v' - u')^2$$

$$(2) \quad (v - u)(u' + v' - 2)(2v'(u' - 1) + v' - u') = 2v(u - 1)(v' - u')^2 + (1_a 1_a + 1)(v - u)(v' - u')^2$$

$$(3) \quad (v - u)(u' + v' - 2)(2u'(v' - 1) - (v' - u')) = (2u(v - 1) + (1_a 1_a - 1)(v - u))(v' - u')^2$$

$$(4) \quad (v - u)(u' + v' - 2)(2u'(v' - 1) - (v' - u')) = 2v(u - 1)(v' - u')^2 + (1_a 1_a + 1)(v - u)(v' - u')^2$$

But

$$(1), (3) \quad (v - u)(u' + v' - 2)(2u'v' - u' - v') = (2uv - 2u)(v' - u')^2 + (1_a 1_a - 1)(v - u)(v' - u')^2$$

$$(2), (4) \quad (v - u)(u' + v' - 2)(2u'v' - u' - v') = (2vu - 2v)(v' - u')^2 + (1_a 1_a + 1)(v - u)(v' - u')^2$$

Hence if we suppose $u' - v' \neq 0$

$$\begin{aligned} & (v - u)((u' - 1)(Y - 1) + (v' - 1)(Y - 1))(u'(v' - 1)(Y - 1)) + v'(u' - 1)(Y - 1) \\ &= (v - u)((u' - 1)(Y - 1) + (u' - 1)(Y + 1))(u'(u' - 1)(Y + 1) + v'(u' - 1)(Y - 1)) \\ & \quad = 2(v - u)(u' - 1)^2 Y(u'(Y + 1) + v'(Y - 1)) \\ & \quad = 4(v - u)(u' - 1)^2 Y(X^{2p-1} + 1) \\ &= (2vu - 2u)(v' - u')^2(Y - 1)(Y - 1) + (1_a 1_a - 1)(v - u)(v' - u')^2(Y - 1)(Y + 1) \\ &= (2vu - 2u + (1_a 1_a - 1)(v - u))(v' - 1 - (u' - 1))(Y - 1)(v' - 1 - (u' - 1))(Y - 1) \\ &= (2vu - 2u + (1_a 1_a - 1)(v - u))((u' - 1)(Y + 1) - (u' - 1)(Y - 1))((u' - 1)(Y + 1) - (u' - 1)(Y - 1)) \\ & \quad = (8u(v - 1) + 4(1_a 1_a - 1)(v - u))(u' - 1)^2 \\ & \quad (u' - 1)^2(2u(v - 1) + (1_a 1_a - 1)(v - u) - (v - u)Y(X^{2p-1} + 1_a 1_a)) = 0 \end{aligned}$$

If the expression in the right of the parenthesis is equal to zero

$$\begin{aligned} (v - 1)(2u + 1_a 1_a - 1 - Y(X^{2p-1} + 1_a 1_a)) &= -(u - 1)Y(X^{2p-1} + 1_a 1_a - 1_a 1_a + 1) \\ &= \frac{(u - 1)(X + 1)}{X - 1}(2u + 1_a 1_a - 1 - Y(X^{2p-1} + 1_a 1_a)) \end{aligned}$$

Thus

$$\begin{aligned} (u - 1)((X + 1)(2u + 1_a 1_a - 1 - Y(X^{2p-1} + 1_a 1_a)) + (X - 1)Y(X^{2p-1} + 1_a 1_a - 1_a 1_a)) &= 0 \\ = (u - 1)(2Y^p + (1_a 1_a - 1)(X - 1) - 2Y(X^{2p-1} + 1_a 1_a) + (1 - 1_a 1_a)(X - 1)) &= 0 \\ = (u - 1)Y^p(2 - 2Y) &= 0 \end{aligned}$$

Consequently

$$u - 1 = 0 \Rightarrow (u - 1)(X + 1) = (v - 1)(X - 1) = 0 \Rightarrow v = 1$$

Impossible (because Y^p is not equal to $X + 1$)! Thus

$$v' = u'$$

$$(1), (3) \quad (v - u)(u' + v' - 2) = \frac{2u(v - 1)(v' - u')^2}{2v'u' - v' - u'}$$

$$(2), (4) \quad (v - u)(u' + v' - 2) = \frac{(2uv + 2v - 4u)(v' - u')^2}{u'v' - u' - v'} = \frac{(2uv - 2u)(v' - u')^2}{u'v' - u' - v'}$$

If $v - u = 0$ then

$$(v - u)X = u + v - 2 = 2u - 2 = 2v - 2 = 0$$

And it is impossible, because Y^p is different of $X + 1$. Thus $v' - u' = 0$, it means

$$(v' - u')Y = u' + v' - 2 = 2u' - 2 = 2v' - 2 = 0$$

Thus

$$v'Y(Y - 1) + Y = u(X + 1) = Y^2 = Y^p$$

And $p = 2$. And if p is odd

$$Y^p - 1 = v'(Y - 1)$$

$$Y^p + 1 = u'(Y + 1)$$

$$2 = u'(Y + 1) - v'(Y - 1) = (Y + 1) - (Y - 1)$$

$$2Y^p = 2X^{2p-1} + 2_a1_a = (u' + v')Y + u' - v'$$

And

$$(u' - 1)(Y + 1) = (v' - 1)(Y - 1)$$

Thus

$$Y = \frac{u' + v' - 2}{v' - u'}$$

$$X^{2p-1} + 1_a1_a = Y^p = \frac{u'(v' - 1) + v'(u' - 1)}{v' - u'}$$

We have four equations with four unknowns

$$\begin{aligned} Y^p &= \left(\frac{u' + v' - 2}{v' - u'}\right)^p = X^{2p-1} + 1_a1_a = \left(\frac{u + v - 2}{v - u}\right)^{2p-1} + 1_a1_a \\ &= \frac{u'(v' - 1) + v'(u' - 1)}{v' - u'} = \frac{u(v - 1) + v(u - 1) + 1_a1_a(v - u)}{v - u} \end{aligned}$$

The solutions are $u' = v' = 1$. Let us prove it : we have also

$$Y^p = Y^p - 1 + 1 = (Y - 1)v' + 1 = X^{2p-1} + 1_a1_a = u(X + 1) + 1_a1_a - 1 \quad (1)$$

$$Y^p = Y^p - 1 + 1 = (Y - 1)v' + 1 = X^{2p-1} - 1 + 1_a1_a + 1 = v(X - 1) + 1_a1_a + 1 \quad (2)$$

$$Y^p = Y^p + 1 - 1 = (Y + 1)u' - 1 = X^{2p-1} + 1 + 1_a1_a - 1 = u(X + 1) + 1_a1_a - 1 \quad (3)$$

$$Y^p = Y^p + 1 - 1 = (Y + 1)u' - 1 = X^{2p-1} - 1 + 1_a1_a + 1 + 2 = v(X - 1) + 1_a1_a + 1 \quad (4)$$

Hence, with

$$X = \frac{u + v - 2}{v - u}, \quad Y = \frac{u' + v' - 2}{v' - u'}$$

We have

$$(1) \quad v' \frac{2u' - 2}{v' - u'} + 1 = u \frac{2v - 2}{v - u} + 1_a 1_a - 1$$

$$(2) \quad v' \frac{2u' - 2}{v' - u'} + 1 = v \frac{2u - 2}{v - u} + 1_a 1_a + 1$$

$$(3) \quad u' \frac{2v' - 2}{v' - u'} - 1 = u \frac{2v - 2}{v - u} + 1_a 1_a - 1$$

$$(4) \quad u' \frac{2v' - 2}{v' - u'} - 1 = v \frac{2u - 2}{v - u} + 1_a 1_a + 1$$

Or

$$(1) \quad (v - u)(2v'(u' - 1) + v' - u') = 2u(v - 1)(v' - u') + (1_a 1_a - 1)(v - u)(v' - u')$$

$$(2) \quad (v - u)(2v'(u' - 1) + v' - u') = 2v(u - 1)(v' - u') + (1_a 1_a + 1)(v - u)(v' - u')$$

$$(3) \quad (v - u)(2u'(v' - 2) - (v' - u')) = 2u(v - 1)(v' - u') + (1_a 1_a - 1)(v - u)(v' - u')$$

$$(4) \quad (v - u)(2u'(v' - 1) - (v' - u')) = 2v(u - 1)(v' - u') + (1_a 1_a + 1)(v - u)(v' - u')$$

But

$$(1), (3) \quad (v - u)(2u'v' - u' - v') = 2u(v - 1)(v' - u') + (1_a 1_a - 1)(v - u)(v' - u')$$

$$(2), (4) \quad (v - u)(2u'v' - u' - v') = (2vu - 2v)(v' - u') + (1_a 1_a + 1)(v - u)(v' - u')$$

Hence if we suppose $u' - v' \neq 0$

$$\begin{aligned} & (v - u)(u'(v' - 1)(Y - 1) + v'(u' - 1)(Y - 1)) \\ &= (v - u)(u'(u' - 1)(Y + 1) + v'(u' - 1)(Y - 1)) \\ &= (v - u)(u' - 1)(2X^{2p-1} + 2_a 1_a - 2) \\ &= (2vu - 2u + (1_a 1_a - 1))(v' - u')(Y - 1) = (2u(v - 1) + (1_a 1_a - 1)(v - u))(v' - 1 - (u' - 1))(Y - 1) \\ &= (2u(v - 1) + (1_a 1_a - 1)(v - u))((u' - 1)(Y + 1) - (u' - 1)(Y - 1)) \\ &= (4u(v - 1) + 2(1_a 1_a - 1)(v - u))(u' - 1) \end{aligned}$$

Thus

$$(u' - 1)(2u(v - 1) + (1_a 1_a - 1)(v - u) - (v - u)(X^{2p-1} + 1_a 1_a)) = 0$$

If

$$\begin{aligned} (v - 1)(X^{2p'-1} + 1 - 2u) &= (u - 1)(X^{2p'-1} + 1) \\ &= \frac{(u - 1)(X + 1)}{X - 1}(X^{2p'-1} + 1 - 2u) \end{aligned}$$

And

$$\begin{aligned} (u - 1)((X + 1)(X^{2p'-1} + 1 - 2u) - (X - 1)(X^{2p'-1} + 1)) &= 0 \\ &= (u - 1)(2X^{2p'-1} + 3 - 2u) = 0 \end{aligned}$$

Consequently

$$u - 1 = 0 \Rightarrow (u - 1)(X + 1) = (v - 1)(X - 1) = 0 \Rightarrow v = 1$$

Impossible (because Y^p is not equal to $X + 1$)! Thus

$$v' = u'$$

And

$$u' + v' = 2 = 2u' = 2v'$$

Thus

$$v'(Y - 1) + 1 = u(X + 1) = Y = Y^p$$

It is impossible! Thus $p = 2$. We deduce that Fermat-Catalan equation implies six equations-solutions which are

$$(I) \quad Y^2 = X^3 + Z^a$$

$$(II) \quad Y^2 = X^4 + Z^a$$

$$(III) \quad Y^2 + X^3 = Z^a$$

$$(IV) \quad Y^2 + X^4 = Z^a$$

$$(V) \quad Y^2 + Z^a = X^3$$

$$(VI) \quad Y^2 + Z^a = X^4$$

Known solutions We have

$$1 + 2^3 = 3^2$$

$$7^2 + 3^5 = 3^4$$

$$13^2 + 7^3 = 2^9$$

$$71^2 = 17^3 + 2^7$$

$$3^5 + 11^4 = 122^2$$

$$1549304^2 + 33^8 = 15613^3$$

$$2213459^2 + 1413^3 = 65^7$$

$$15312283^2 = 113^7 - 9262^3$$

$$21062938^2 = 76271^3 + 17^7$$

$$30042907^2 = 96222^3 + 43^8$$

Conclusion

Fermat-Catalan equation $Z^c = X^a + Y^b$ does not have solutions for both $a > 2$, $b > 2$ and $c > 2$. In fact, at least one of the exponents must be equal to 2. We have shown a way to discuss the solutions.