



**HAL**  
open science

## Prey selection by the common dolphin: fulfilling high energy requirements with high quality food

Jérôme Spitz, Emeline Mourocq, Jean-Pierre Léauté, Jean-Claude Quéro,  
Vincent Ridoux

### ► To cite this version:

Jérôme Spitz, Emeline Mourocq, Jean-Pierre Léauté, Jean-Claude Quéro, Vincent Ridoux. Prey selection by the common dolphin: fulfilling high energy requirements with high quality food. *Journal of Experimental Marine Biology and Ecology*, 2010, 390 (2), pp.73-77. hal-00751689

**HAL Id: hal-00751689**

**<https://hal.science/hal-00751689v1>**

Submitted on 14 Nov 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 Prey selection by the common dolphin: fulfilling high energy 2 requirements with high quality food

3  
4 Jérôme Spitz<sup>a,b\*</sup>, Emeline Mourocq<sup>c</sup>, Jean-Pierre Leauté<sup>d</sup>, Jean-Claude Quéro<sup>e</sup>, Vincent  
5 Ridoux<sup>a,c</sup>

6  
7 <sup>a</sup>Littoral, Environnement et Sociétés, UMR 6250, Université de La Rochelle / CNRS, 2 rue Olympe de Gouges,  
8 17032 La Rochelle, Cedex, France

9 <sup>b</sup>Parc zoologique de La Flèche, Le Tertre Rouge, 72200 La Flèche, France

10 <sup>c</sup>Centre de Recherche sur les Mammifères Marins, Fédération de Recherche en Environnement et  
11 Développement Durable, Université de La Rochelle, 17071 La Rochelle, France

12 <sup>d</sup>Ifremer, Station RH de La Rochelle, Place du Séminaire, BP 7, 17137 L'Houmeau, France.

13 <sup>e</sup>Muséum d'Histoire Naturelle de La Rochelle, 28 rue Albert 1<sup>er</sup>, 17000 La Rochelle

14  
15 \*Corresponding author. Tel.: +33 5 46507658; fax.: +33 5 46449945; *E-mail address*: [jspitz@univ-lr.fr](mailto:jspitz@univ-lr.fr) (J.  
16 Spitz)

## 17 18 ABSTRACT

19 Which characteristics define the prey species constituting the diet of a given predator?  
20 Answering this question would help predict a predator's diet and improve our understanding  
21 of how an ecosystem functions. The aim of this study was to test if the diet of common  
22 dolphins, *Delphinus delphis*, in the oceanic Bay of Biscay reflected prey availability or a  
23 selection shaped by prey energy densities (ED). To do this, the community of potential prey  
24 species, described both in terms of relative abundance and energy densities, was compared to  
25 the common dolphin diet in this area. This analysis of a predator's diet and its prey field  
26 revealed that the common dolphin selected its diet on the basis of prey energy densities  
27 (significant values of Chesson's index for  $ED > 5 \text{kJ.g}^{-1}$ ). High-energy prey were positively  
28 selected in the diet [*e.g.* *Notoscopelus kroeyeri*,  $ED = 7.9 \text{kJ.g}^{-1}$ , 9% of relative abundance in

29 the environment (%Ne); 62% of relative abundance in the diet (%Nd)] and low-energy prey  
30 disregarded (*Xenodermichthys copei*, ED=2.1kJ.g<sup>-1</sup>, 20%Ne, 0%Nd). These results supported  
31 the hypothesis that common dolphins selected high energy density prey species to meet their  
32 energetically expensive life style and disregard prey organisms of poor energy content even  
33 when abundant in the environment.

34

35 *Keyword:* field of prey; energy density; active prey selection; trophic *cul-de-sac*

36

### 37 **1. Introduction**

38 Beyond the simple description of the diet, a major ecological issue is also to understand its  
39 ecological significance: why does the observed array of prey species compose the diet of a  
40 given predator? What are the prey key traits for this predator? Answering this would help  
41 predict a predator's diet and its variations according to the availability of prey functional  
42 traits, improve our understanding of ecosystem functioning and anticipate the consequences  
43 of possible changes.

44 The optimal foraging theory (OFT) is a classical paradigm used to explain or even  
45 predict the diet of a forager (Emlen, 1966; MacArthur and Pianka, 1966; Charnov, 1976). In  
46 initial studies placed in the OFT framework, some predictions depended on unrealistic  
47 assumptions and OFT was a much debated theory (Zach and Smith, 1981; Stephens and Krebs,  
48 1986; Pierce and Ollason, 1987). However, numerous studies provided results which matched  
49 quantitatively or qualitatively OFT predictions (Sih and Christensen, 2001). Today, three  
50 major robust and fundamental predictions are commonly assumed (Stephens and Krebs, 1986;  
51 Sih and Christensen, 2001): (1) predators should prefer prey that yield more energy compared  
52 to foraging costs, (2) as abundance of higher profitable prey species increases in the  
53 environment, lower profitable prey should be dropped out from the diet and predators should

54 become more specialised, and (3) foragers should obey a quantitative threshold rule for when  
55 specific prey types should be included or excluded from an optimal diet. Thus, predators have  
56 to develop an adapted foraging strategy related with their specific energy requirements.

57 From killer whales eating marine mammals to herbivorous dugong, marine mammals  
58 exhibit a broad range of biological models and have developed diverse feeding strategies  
59 (Berta and Sumich, 1999). But, some species, particularly small cetaceans and pinnipeds, are  
60 often described as opportunistic feeders: predators which consume their prey without  
61 selection, *i.e.* proportionately to their availability in the environment. Contrastingly, active  
62 prey choice by marine mammals has rarely been actually tested. The difficulty to describe the  
63 prey field exposed to a predator species (species diversity and abundance, distributions,  
64 energy contents...) is an important limitation in such investigations (*e.g.* Santos and Pierce,  
65 2003).

66 The common dolphin *Delphinus delphis* is the most abundant delphinid in offshore warm-  
67 temperate waters in the Atlantic (Perrin, 2002). This small delphinid is likely an energetically  
68 expensive biological model because it is fairly small among cetaceans and a very active  
69 swimmer, both of these characteristics being associated with high metabolic costs per unit  
70 body mass (Berta and Sumich, 1999; Costa and Williams, 1999). In the Bay of Biscay,  
71 common dolphin feed preferentially on small schooling pelagic fish: scads *Trachurus* spp.,  
72 pilchard *Sardina pilchardus*, anchovy *Engraulis encrasicolus* and mackerels *Scomber*  
73 *scombrus*. Between seasons or years, the specific composition of its diets can change  
74 significantly (Meynier et al., 2008), but, if the proximate composition of forage species was  
75 considered (Spitz et al., 2010), dolphins seemed always to switch from a fat and high-quality  
76 prey species to another high-quality prey species. Low-quality prey would be neglected;  
77 indeed several of them, such as gadids, are very abundant in the area (Poulard and Blanchard,  
78 2005), but never constitute a measurable share of the common dolphin diet (Meynier et al.,

79 2008). Hence, prey selection by common dolphins based on prey energy densities was  
80 suggested to occur in this neritic habitat. Recently and for the first time in a fully oceanic  
81 area, the diet of common dolphins off the Bay of Biscay was shown to be largely constituted  
82 of myctophids (Pusineri et al., 2007). By comparison with other predators studied in the same  
83 area and collected in the same condition, *i.e.* living in the same prey field, it appeared that  
84 similarly-sized pelagic predators such as blue shark, *Prionace glauca*, or swordfish, *Xiphias*  
85 *gladius*, ate larger and probably leaner prey types (Pusineri et al., 2008).

86 The present study aimed at testing if, within the fish prey field available to top  
87 predators, common dolphins would select the most profitable food sources, *i.e.* high energy  
88 density prey, to fulfil their high energy requirements. To investigate this issue, we compared  
89 the composition of the community of potential dolphin prey species off the Bay of Biscay,  
90 both in terms of relative abundance and energy density with the diet of common dolphins in  
91 order to test if the diet reflected prey availability or a selection shaped by prey energy  
92 densities.

93

## 94 **2. Materials and methods**

95

### 96 *2.1. Composition of the epi- and mesopelagic fish community*

97

98 The epi- to mesopelagic oceanic fish community off the Bay of Biscay was  
99 investigated during EVHOE (*EValuation des ressources Halieutiques de l'Ouest Européen*)  
100 research cruises onboard the R/V *Thalassa* in October 2002, 2003 and 2008. Fourteen hauls  
101 were performed with a 25 m vertical opening pelagic trawl. The mesh size decreases  
102 gradually from 76 mm to 44 mm in the bag. The trawl-haul duration was one hour at 4 kn.  
103 The hauls were carried out during the night at various depths supposedly accessible to

104 dolphins from sub-surface to 500 m deep off the continental slope from 43.91–46.60°N and  
105 2.40–5.11°W (Table 1). Fish were counted and identified following published guides (*e.g.*  
106 Quéro et al., 2003).

107 The occurrence of a given species was the number of haul in which this species was  
108 observed. The relative abundance was given by the number of individuals of the same species  
109 found throughout the complete haul series. These two indices can be expressed by their  
110 percentage frequency as percentage of occurrence (%O) and percentage by number (%N),  
111 respectively.

$$112 \quad \%O_i = n_i / N * 100$$

113 where  $n_i$  is the number of hauls where species  $i$  was found and  $N$  the total number of hauls;

$$114 \quad \%N_i = x_i / X * 100$$

115 where  $x_i$  is the number of individuals belonging to species  $i$  and  $X$  the total number of fish  
116 caught.

117 Confidence intervals around relative abundance were generated by bootstrap  
118 simulations. The bootstrapping routine was written by using the *R* software (Ihaka and  
119 Gentleman, 1996). Random samples were drawn with replacement and the procedure was  
120 repeated 1000 times.

121

## 122 *2.2. Diet of common dolphin and energy content of prey species: origin of data*

123

124 The dietary composition of common dolphin used in the present work comes from a  
125 previous analysis of stomach contents carried out from dolphins incidentally caught in tuna  
126 driftnet fisheries during the summers 1992-1993 off the Bay of Biscay (Pusineri et al. 2007;  
127 summary in Table 2). Briefly described, stomach contents from 63 common dolphins were  
128 analyzed by prey occurrence, number and mass, following standard methods (*e.g.* Ridoux,

129 1994, Spitz et al., 2006). The diet was dominated by myctophid fish, mostly *Notoscopelus*  
130 *kroeyeri*.

131 Quality of forage species from the Bay of Biscay was recently explored by analysis of  
132 proximate composition and energy content for a wide range of species including mesopelagic  
133 fish (Spitz et al., 2010; Table 3).

134

### 135 2.3. Comparison between diet and prey availability

136

137 The overlap between hauls and stomach contents was assessed by calculating the  
138 Pianka index of overlap on relative abundance data of fish species in the water *versus* in the  
139 diet (Pianka, 1973):

$$140 \quad O = \frac{\sum p_{iA} p_{iB}}{\sqrt{\sum p_{iA}^2 \sum p_{iB}^2}}$$

141

142 where  $p_{iA}$  is the percentage by number of species  $i$  in hauls and  $p_{iB}$  is the percentage by  
143 number of the species  $i$  in the diet of the common dolphin. This index varies from 0 (no  
144 overlap) to 1 (complete overlap).

145

146 Prey selectivity was tested by using the Chesson's index (Chesson, 1978):

$$147 \quad \alpha_{i(1 \rightarrow m)} = \frac{p_i r_i}{\sum p_i r_i}$$

148 where  $\alpha_i$  is the selectivity for prey type  $i$ ;  $r_i$  is the percentage by number of species  $i$  in the diet  
149 of the common dolphin;  $p_i$  percentage by number of species  $i$  in hauls and  $m$  is the total  
150 number of species found in hauls and in stomach contents. Values of  $\alpha_i$  close to  $1/m$  represent

151 feeding at random whereas values greater (*versus* smaller) than 1/m correspond to positive  
152 (*versus* negative) selection of prey *i*.

153 Pearson's correlation tests between species abundance in the hauls *versus* in the diet  
154 was performed to examine if the diet reflected prey availability (percentage by number >1%  
155 either in diet or in hauls). Similar tests were carried out between Chesson's index of  
156 selectivity and energetic density to investigate if prey choice was associated to prey quality.  
157 All correlations were computed by using the *R* software (Ihaka and Gentleman, 1996).

158

### 159 **3. Results**

160

#### 161 *3.1. Composition of the epi- and mesopelagic fish community*

162

163 From the fourteen epi- to mesopelagic trawl hauls, 3258 fish were captured. The  
164 pelagic fish community was composed of 26 species from 12 families. Three families  
165 accounted for up to 88% of total number of fish caught (Table 4). The alepocephalid  
166 *Xenodermichtys copei* was the most abundant species with a total contribution of 22.2%N,  
167 95% non-parametric confidence interval being [12.9–32]. Myctophid fish were represented by  
168 at least eight species and amounted to 44.5%N; among them *Benthoosema glaciale* (17.2%N  
169 [2.2–40]) and *Notoscopelus kroeyeri* (8.9%N [5.1–13.8]) were the most prevalent. The  
170 sternoptychids were the third dominant family with three species identified: *Argyropelecus*  
171 *olfersii* (11.3%N [4.6–21.8]), *A. hemigymnus* (3.4%N [0.6–8.2]) and *Maurolicus muelleri*  
172 (6.6%N [0.1–18.1]). Lastly, the paralepid *Arctozenus risso* accounted for 8.1%N [3.7–14.2].  
173 Abundances of all other species were negligible.

174

#### 175 *3.2. Comparison between diet and prey availability*

176



177 A first approach to the comparison between common dolphin diet and prey availability  
178 was by using the Pianka index of overlap which revealed a fairly low degree of overlap (<0.4)  
179 between the two datasets. Similarly, no correlation (Pearson test,  $P>0.05$ ) existed between  
180 relative abundances of each prey species observed in hauls *versus* in stomach contents (Figure  
181 1). Actually, the Chesson's index of selectivity suggested a positive selection by which  
182 common dolphin would specifically target *M. punctatum*, *M. muelleri* and even more strongly  
183 *B. glaciale* and *N. kroeyeri* (Table 5). All other species appeared to be negatively selected by  
184 the predator. Finally, a significant correlation (Pearson test,  $P<0.005$ ) was found between the  
185 values of Chesson's index of selectivity and fish energy content (Figure 2), with all species  
186 poorer than  $5 \text{ kJ.g}^{-1}$  being negatively selected and the other being increasingly selected as  
187 their energy content increased.

188

## 189 **4. Discussion**

190

### 191 *4.1. General comments*

192

193 Prey choice or selection by marine predators is difficult to measure, especially in fully  
194 oceanic area. The degree of selectivity or opportunism of such predators was often inferred  
195 from dietary analyses only. For instance, predators showing a broad diversity in their diet or  
196 feeding upon reportedly abundant prey species were classified as opportunistic feeders (*e.g.*  
197 Bearzi et al., 2009). In combining results of stomach content analysis and description of prey  
198 field, the present study revealed a strongly significant relationship between prey selection  
199 indices and prey energy densities. Thus, common dolphins selected high energy density prey  
200 species to face their energetically expensive life style and disregarded prey organisms poorer  
201 than  $5 \text{ kJ.g}^{-1}$  even when abundant in the environment.

202 Both stomach content analysis and trawling are subject to their own selectivity and  
203 biases which could affect our perception of dolphin diets and fish abundance. The  
204 representativeness of diet described by stomach content analysis was often subject to the  
205 difficulty of controlling the sampling design and to the differential digestion of ingested prey  
206 (*e.g.* Tollit et al., 1997; Pierce et al., 2007). Especially for protected species such as marine  
207 mammals, the dietary results were often limited by the size, the spatio-temporal coverage, or  
208 for instance, the age/sex/reproductive status composition of the individuals providing the  
209 stomach content sample set. In spite of these limitations, however, stomach content analysis is  
210 still the best and most widely used approach to investigate marine top predator diets and  
211 provide the most detailed information on prey composition. On the other hand, the  
212 representativeness of a fish community described from trawling survey is equally affected by  
213 sampling design (*i.e.* number of trawls, spatio-temporal coverage, immersion/depth/daytime  
214 stratification,...) and the differential escape capabilities of target species for a given trawl  
215 (Wardle, 1993). In the present work, the low number of hauls increased these limitations.  
216 However, oceanic fish communities are often poorly described around the world, in particular  
217 because the cost of fish survey is generally higher in oceanic habitats than in coastal ones.  
218 The present study provides original data on the eastern North Atlantic mesopelagic fish  
219 community off the Bay of Biscay. Despite the fairly limited number of trawl hauls on which  
220 the description of the mesopelagic fish community was based, the results obtained were  
221 consistent with previous data in the same region (Quéro, 1969; Quéro et al., 2002) or in  
222 adjacent areas (Roe et al., 1984; Fock et al., 2004). Myctophids with *N. kroeyeri* and *B.*  
223 *glaciale*, alepocephalids with *X. copei*, sternoptichids with *A. olfersii*, *A. hemigymnus* and *M.*  
224 *muelleri* and paralepids with *A. risso* were the main species amongst a highly diversified fish  
225 community. Given the sources of uncertainty and biases both in diet and fish community  
226 descriptions, the composition values used in the present work should be considered as

227 revealing general patterns. Hence, the strong significant relationship observed between prey  
228 selection indices and prey energy content reveals a general pattern that supports the  
229 hypothesis of a quality-based prey selection by the common dolphin.

230

#### 231 *4.2. Considerations on foraging strategies*

232

233 This active selection based on prey quality suggests that, within a given prey field,  
234 predators would adapt the quality of their diets according to their specific energy  
235 requirements. Hence, in the oceanic Bay of Biscay the diets of the top predator community  
236 studied sympatrically revealed a large diversity of potential prey species (Pusineri et al.,  
237 2008). Among this field of available prey, observed differences in the quality of forage  
238 species (Spitz et al., 2010) could reflect differences in predator-specific metabolic needs and  
239 associated foraging strategies. Indeed, in addition to being the main prey of the common  
240 dolphin, *N. kroeyeri*, a high-quality prey, was also the main fish prey of the striped dolphins  
241 *Stenella coeruleoalba*, the other homoeothermic top predator (Ringelstein et al., 2006). Thus,  
242 predators with high energy requirement would be constrained to feeding on the most  
243 profitable food sources constituted of small, gregarious and high energy density prey types.  
244 Whereas in the diet of large ectothermic predators, high-quality prey were either rare (<5  
245 %M) in the swordfish *Xiphias gladius* (Chancollon et al., 2006) or absent in the blue shark  
246 *Prionace glauca* (Pusineri et al., 2008). These latter predators with lower energy requirements  
247 would be better fitted to exploit lower quality food sources made of larger and leaner prey  
248 species.

249

#### 250 *4.3. Ecosystem implications*

251

252 Beyond a standard description of food habits, to attempt explaining the diet  
253 composition of marine top predators increases the knowledge of prey-predator functional  
254 relationship. Indeed, prey selection by top predators could shape the structure of communities,  
255 but the challenge is to identify the key functional traits which influence food web structure  
256 and ecosystem functioning (Lazzaro et al., 2009). Here, the lowest quality, albeit most  
257 abundant, mesopelagic fish species, such as *X. copei*, were neglected, or negatively selected,  
258 by all top-predators studied so far in the area (Pusineri et al., 2008) and should therefore be  
259 considered as trophic *cul-de-sacs*. In contrast, the highest quality prey (e.g. *N. kroeyeri* and *M.*  
260 *muelleri*) are positively selected by predators with energetically expensive life styles and  
261 should be considered as trophic highways to these predators. Predation being a major  
262 selective pressure for organisms of intermediate trophic level, forage species developed many  
263 adaptations allowing predation risk to be reduced, such as toxicity, camouflage or specific  
264 behavioral characteristics (Caro, 2005). Thus, prey with traits which reduce predation risk  
265 would be favored within a trophic level (Preisser et al., 2007). In this evolutionary context,  
266 one may interpret low energy density and associated poor nutritional quality as a successful  
267 adaptation for mesopelagic fish that would limit predation risk and could explain their high  
268 abundance within this fish community.

269

## 270 **Acknowledgments**

271 We are particularly grateful to J.C. Mahé, R. Bellail and J.C. Poulard (Ifremer) and the whole  
272 staff of the R/V *Thalassa* for their help in the mesopelagic fish community investigations.  
273 CRMM is funded by *Ministère de l'Ecologie et du Développement Durable* and by  
274 *Communauté d'Agglomération de la Ville de La Rochelle*. The PhD thesis of J. Spitz is  
275 supported by the *Agence Nationale de la Recherche Technique* with a CIFRE grant.

276

## 277 **References**

278

279 Bearzi, G., Fortuna, C.M., Reeves, R.R., 2009. Ecology and conservation of common  
280 bottlenose dolphins *Tursiops truncatus* in the Mediterranean Sea. *Mammal Rev.* 39, 92-  
281 123.

282 Berta, A., Sumich, J.L., 1999. *Marine Mammals: Evolutionary Biology*, San Diego:  
283 Academic Press.

284 Caro, T., 2005. *Antipredator Defenses in Birds and Mammals*, Chicago: University of  
285 Chicago Press.

286 Chancollon, O., Pusineri, C., Ridoux, V., 2006. Food and feeding ecology of the North  
287 Atlantic swordfish, *Xiphias gladius*, off the Bay of Biscay. *ICES J. Mar. Sci.* 63, 1075-  
288 1085.

289 Charnov, E.L., 1976. Optimal foraging: attack strategy of a mantid. *Am. Nat.* 110, 141–151.

290 Chesson, J., 1978. Measuring performance in selective predation. *Ecology* 59, 211-215.

291 Costa, D.P., Williams, T.M., 1999. Marine mammal energetic. In *Biology of Marine*  
292 *Mammals* (eds Reynolds, J.E., Rommel, S.A.), pp 176-217. Washington: Smithsonian  
293 Institution Press.

294 Emlen, J.M., 1966. The role of time and energy in food preference. *Am. Nat.* 100, 611-617.

295 Fock, H.O., Pusch, C., Ehrich, S. 2004. Structure of deep-sea pelagic fish assemblages in  
296 relation to the Mid-Atlantic Ridge (45°–50°N). *Deep-Sea Res.* 51, 953-978.

297 Ihaka, R., Gentleman, R., 1996. R: a language for data analysis and graphics. *J. Comput.*  
298 *Graph. Stat.* 5, 299-314.

299 Lazzaro, X., Lacroix G., Gauzens, B., Gignoux J., Legendre, S., 2009. Predator foraging  
300 behaviour drives food-web topological structure. *J. Anim. Ecol.* 78, 1307–1317.

301 MacArthur, R.H., Pianka, E.R., 1966. On the optimal use of a patchy environment. *Am. Nat.*  
302 100, 603-609.

303 Meynier, L., Pusineri, C., Spitz, J., Santos, M.B., Pierce, G.J., Ridoux, V., 2008. Diet and  
304 feeding ecology of the common dolphin, *Delphinus delphis*, in the Bay of Biscay and its  
305 intraspecific variations. *Mar. Ecol. Prog. Ser.* 354, 277-287.

306 Perrin, W.F., 2002. Common dolphin. In *Encyclopedia of Marine Mammals* (eds Perrin,  
307 W.F., Würsig, B., Thewissen, J.G.M.), pp 245-248. London: Academic Press.

308 Pianka, E.R., 1973. The structure of lizard communities. *Ann. Rev. Ecol. Syst.* 4, 53-74.

309 Pierce, G.J., Ollason, J.G., 1987. Eight reasons why optimal foraging theory is a complete  
310 waste of time. *Oikos* 49, 111–118.

311 Pierce, G.J., Santos, M.B., Cerviño, S. 2007. Assessing sources of variation underlying  
312 estimates of cetacean diet composition: A simulation study on analysis of harbour  
313 porpoise diet in Scottish (UK) waters. *J. Mar. Biol. Assoc. UK* 87, 213-221.

314 Poulard, J.C., Blanchard, F., 2005. The impact of climate change on the fish community  
315 structure of the eastern continental shelf of the Bay of Biscay. *ICES J. Mar. Sci.* 62,  
316 1436-1443.

317 Preisser, E.L., Orrock, J.L., Schmitz, O.J., 2007. Predator hunting mode and habitat domain  
318 alter nonconsumptive effects in predator–prey interactions. *Ecology* 88, 2744–2751.

319 Pusineri, C., Chancollon, O., Ringelstein, J., Ridoux, V., 2008. Feeding niche segregation  
320 among the Northeast Atlantic community of oceanic top predators. *Mar. Ecol. Prog. Ser.*  
321 361, 21-34.

322 Pusineri, C., Magnin, V., Meynier, L., Spitz, J., Hassani, S., Ridoux, V., 2007. Food and  
323 feeding ecology of the common dolphin (*Delphinus delphis*) in the oceanic northeast  
324 Atlantic and comparison with its diet in neritic areas. *Mar. Mam. Sci.* 23, 30-47.

325 Quéro, J.C., 1969. Liste des poissons capturés au cours des pêches pélagiques profondes de la  
326 *Thalassa* dans l'Atlantique nord-est. *CIEM, CM 1969/L: 25*, pp 1-5.

327 Quéro, J.C., Porché, P., Vayne, J.J., 2003. Guide des poissons de l'Atlantique européen, Paris:  
328 Delachaux et Niestlé.

329 Quéro, J.C., Spitz, J., Vayne, J.J., 2002. Observations ichtyologiques effectuées en 2001. Ann.  
330 Soc. Sci. nat. Char. Mar. 9, 179-186.

331 Ridoux, V., 1994. The diets and dietary segregation of seabirds at the subantarctic Crozet  
332 Islands. Mar. Ornithol. 22, 1-192.

333 Ringelstein, J., Pusineri, C., Hassani, S., Meynier, L., Nicolas, R., Ridoux, V., 2006. Food and  
334 feeding ecology of the striped dolphin, *Stenella coeruleoalba*, in the oceanic waters of  
335 the north-east Atlantic. J. Mar. Biol. Assoc. UK 86, 909-918.

336 Roe, H.S.J., Angel, M.V., Badcock, P., Domanski, P.T., Pugh, P.R., Thurston, M.H., 1984.  
337 The diel migration and distributions within a mesopelagic community in the North East  
338 Atlantic. Progr. Oceanogr. 13, 245-511.

339 Santos, M.B., Pierce, G.J., 2003. The diet of harbour porpoise (*Phocoena phocoena*) in the  
340 northeast Atlantic. Oceanogr. Mar. Biol. 41, 355-390.

341 Sih, A., Christensen, B., 2001. Optimal diet theory: when does it work, and when and why  
342 does it fail? Anim. Behav. 61, 379-390.

343 Spitz, J., Mourocq, E., Shoen, V., Ridoux, V., 2010. Proximate composition and energy  
344 content of forage species from the Bay of Biscay: high or low quality food? ICES J.  
345 Mar. Sci., 67: xxx-xxx.

346 Spitz, J., Rousseau, Y., Ridoux, V., 2006. Diet overlap between harbour porpoise and  
347 bottlenose dolphin: an argument in favour of interference competition for food? Estuar.  
348 Coast. Shelf. S. 70, 259-270.

349 Stephens, D.W., Krebs, J.R., 1986. Foraging Theory. Princeton: Princeton University Press.

- 350 Tollit, D.J., Steward, M., Thompson, P.M., Pierce, G.J., Santos, M.B., Hughes, S. 1997.  
351 Species and size differences in the digestion of otoliths and beaks; implications for  
352 estimates of pinniped diet composition. *Can. J. Fish. Aquat. Sci.* 54, 105–119.
- 353 Wardle, C.S., 1993. Fish behaviour and fishing gear. In *Behaviour of Teleost Fishes* (ed  
354 Pitcher TJ), pp 609–643. London: Chapman & Hall.
- 355 Zach, R., Smith, J.M.N., 1981. Optimal foraging in wild birds? In *Foraging Behavior:*  
356 *Ecological, Ethological and Psychological* (eds Kamil, A.C., Sargent, T.D.), pp 95–107.  
357 New York: Garland STPM Press.



358 Figure 1. Correlation between relative abundance in the environment for major fish species  
359 and their relative abundance in the diet of common dolphin.

360

361 Figure 2. Correlation between energetic value for major fish species and their value of  
362 Chesson's index.

363

364 Table 1. Trawling station data for the RV *Thalassa*

365

366 Table 2. Composition of the diet of common dolphin off the Bay of Biscay in percentage by  
367 number (Pusineri et al., 2007)

368


369 Table 3. Energy content of main mesopelagic fish species off the Bay of Biscay (Spitz et al.,  
370 2010)

371

372 Table 4. Composition of fish community in the 500 meters depth surface layer off the Bay of  
373 Biscay


374

375 Table 5. Values of Chesson's index ( $\alpha_i$ )



377

378 Figure 1



380

381 Figure 2

382

383 Table 1

384

Date	Bottom depth (m)	Catch depth (m)	Start of the haul	Latitude °N	Longitude °W
15/10/2002	1500	500	20:13	44.34	2.46
16/10/2002	1500	200	19:04	44.58	2.40
19/10/2002	2000	20	19:56	44.90	2.57
20/10/2003	700	200	20:13	45.78	3.76
22/10/2003	600	500	19:40	46.60	4.95
24/10/2003	3700	500	20:10	45.30	3.69
25/10/2003	1500	200	20:36	43.92	2.81
26/10/2003	1500	500	19:51	43.91	2.80
02/11/2003	3700	200	20:07	46.27	5.11
22/10/2008	1200	500	19:47	44.33	2.29
23/10/2008	800	200	20:02	43.75	2.24
25/10/2008	3300	20	21:06	44.79	2.74
26/10/2008	1500	500	20:05	44.99	2.67
28/10/2008	700	500	19:35	45.28	3.33

385

Family	Species	%N
<b>Sternoptichidae</b>		
	<i>Maurolicus muelleri</i>	9.4
<b>Platyroctidae</b>		
	Platyroctidae unidentified	0.1
<b>Stomiidae</b>		
	<i>Stomias boa ferox</i>	0.1
<b>Chauliodontidae</b>		
	<i>Chauliodus sloani</i>	0.1
<b>Chiasmodontidae</b>		
	Chiasmodontidae unidentified	0.1
<b>Bathylagidae</b>		
	All Bathylagidae species	0.2
<b>Paralepididae</b>		
	<i>Arctozenus risso</i>	1.1
	<i>Paralepis coregonoides</i>	0.7
	<i>Macroparalepis affinis</i>	0.1
<b>Myctophidae</b>		
	<i>Notoscopelus kroeyeri</i>	62.4
	<i>Myctophum punctatum</i>	5.3
	<i>Benthoosema glaciale</i>	11.3
	<i>Symbolophorus veranyi</i>	0.7
	<i>Lampanyctus</i> spp.	2.3
	<i>Ceratoscopelus maderensis</i>	0.4
	<i>Lobianchia gemellarii</i>	0.3
	<i>Electrona risso</i>	0.1
	Myctophidae unidentified	0.5
<b>Nomeidae</b>		
	<i>Cubiceps gracilis</i>	0.2
<b>Others fish species</b>		0.6
<b>Cephalopod species</b>		4.0

<b>Family</b>	<b>Species</b>	<b>Gross energy (kJ/g)</b>
<b>Serrivomeridae</b>	<i>Serrivomer beanii</i>	2.1
<b>Alepocephalidae</b>	<i>Xenodermichtys copei</i>	2.2
<b>Platyroctidae</b>	<i>Normichthys operosa</i>	2.6
<b>Sternoptychidae</b>	<i>Argyropelecus olfersii</i>	3.5
	<i>Maurolicus muelleri</i>	4.2
<b>Stomiidae</b>	<i>Stomias boa ferox</i>	2.8
<b>Paralepididae</b>	<i>Arctozenus risso</i>	4.3
<b>Myctophidae</b>	<i>Lampanyctus crocodilus</i>	4.1
	<i>Notoscopelus kroeyeri</i>	7.9
	<i>Benthoosema glaciale</i>	5.9

Family	Species	%O	N	%N	95% IC
<b>Serrivomeridae</b>	<i>Serrivomer beanii</i>	21.4	11	0.3	[0-1.1]
<b>Bathylagidae</b>	<i>Dolicholagus longirostris</i>	7.1	1	0.0	[0-0.1]
<b>Alepocephalidae</b>	<i>Xenodermichtys copei</i>	85.7	724	22.2	[12.9-32]
<b>Platyroctidae</b>	<i>Holthyrnia macrops</i>	7.1	1	0.0	[0-0.1]
<b>Sternoptychidae</b>	<i>Argyrolepecus hemigymnus</i>	57.1	110	3.4	[0.6-8.2]
	<i>Argyrolepecus olfersii</i>	78.6	369	11.3	[4.6-21.8]
	<i>Maurolicus muelleri</i>	28.6	214	6.6	[0.1-18.1]
<b>Stomiidae</b>	<i>Melanostomias bartonbeani</i>	7.1	1	0.0	[0-0.1]
	<i>Rhadinesthes decimus</i>	7.1	1	0.0	[0-0.1]
	<i>Stomias boa feroc</i>	71.4	69	2.1	[0.9-3.8]
<b>Gonostomatidae</b>	<i>Cyclothone microdon</i>	14.3	15	0.5	[0-1.4]
<b>Chiasmodontidae</b>	<i>Pseudoscopelus pierbartus</i>	7.1	1	0.0	[0-0.1]
<b>Paralepididae</b>	<i>Arctozenus risso</i>	71.4	264	8.1	[3.7-14.2]
	<i>Lestidiops affinis</i>	7.1	3	0.1	[0-0.3]
	<i>Macroparalepis affinis</i>	57.1	16	0.5	[0.2-0.8]
	<i>Paralepis coregonoides</i>	7.1	1	0.0	[0-0.1]
<b>Myctophidae</b>	<i>Benthoosema glaciale</i>	85.7	561	17.2	[2.2-40]
	<i>Ceratoscopelus maderensis</i>	78.6	146	4.5	[1.9-7.5]
	<i>Electrona risso</i>	7.1	2	0.1	[0-0.2]
	<i>Lobianchia gemellarii</i>	7.1	1	0.0	[0-0.1]
	<i>Myctophum punctatum</i>	78.6	327	10.0	[2.4-25.5]
	<i>Notoscopelus kroeyeri</i>	85.7	290	8.9	[5.1-13.8]
	<i>Lampanyctus spp.</i>	57.1	124	3.8	[0.2-10.4]
	<i>Diaphus mollis</i>	7.1	1	0.0	[0-0.1]
<b>Merlucciidae</b>	<i>Cynogadus brachycolus</i>	7.1	1	0.0	[0-0.1]
<b>Syngnathidae</b>	<i>Entelurus aequoerus</i>	14.3	4	0.1	[0-0.4]

<b>Species</b>	<b><math>\alpha</math></b>	<b>Selection</b>
<i>Serrivomer beanii</i>	0.000	Negative
<i>Xenodermichtys copei</i>	0.000	Negative
<i>Argyropelecus hemigymnus</i>	0.000	Negative
<i>Argyropelecus olfersii</i>	0.000	Negative
<i>Maurollicus muelleri</i>	0.070	<b>Positive</b>
<i>Stomias boa ferox</i>	0.000	Negative
<i>Cyclothone microdon</i>	0.000	Negative
<i>Chauliodus sloani</i>	0.000	Negative
<i>Arctozenus risso</i>	0.010	Negative
<i>Macroparalepis affinis</i>	0.000	Negative
<i>Paralepis coregonoides</i>	0.000	Negative
<i>Benthoosema glaciale</i>	0.220	<b>Positive</b>
<i>Ceratoscopelus maderensis</i>	0.002	Negative
<i>Lampanyctus</i> spp.	0.000	Negative
<i>Lobianchia gemellarii</i>	0.000	Negative
<i>Myctophum punctatum</i>	0.060	<b>Positive</b>
<i>Notoscopelus kroeyeri</i>	0.628	<b>Positive</b>
<i>Electrona risso</i>	0.000	Negative
<i>Symbolophorus veranyi</i>	0.000	Negative
<i>Cubiceps gracilis</i>	0.000	Negative
<i>Scomberesox saurus</i>	0.000	Negative
Bathylagidae	0.000	Negative
Chiasmodontidae	0.000	Negative
Platyroctidae	0.000	Negative
Others Stomidae	0.000	Negative
Syngnathidae	0.000	Negative