



HAL
open science

Large-time asymptotics for an uncorrelated stochastic volatility model

Martin Forde

► **To cite this version:**

Martin Forde. Large-time asymptotics for an uncorrelated stochastic volatility model. *Statistics and Probability Letters*, 2011, 81 (8), pp.1230. 10.1016/j.spl.2011.03.021 . hal-00753944

HAL Id: hal-00753944

<https://hal.science/hal-00753944>

Submitted on 20 Nov 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Large-time asymptotics for an uncorrelated stochastic volatility model

Martin Forde


PII: S0167-7152(11)00104-0
DOI: 10.1016/j.spl.2011.03.021
Reference: STAPRO 5956

To appear in: *Statistics and Probability Letters*

Received date: 23 December 2010

Revised date: 8 March 2011

Accepted date: 9 March 2011



Please cite this article as: Forde, M., Large-time asymptotics for an uncorrelated stochastic volatility model. *Statistics and Probability Letters* (2011), doi:10.1016/j.spl.2011.03.021

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Large-time asymptotics for an uncorrelated stochastic volatility model

Martin Forde

Department of Mathematical Sciences, Dublin City University, Glasnevin, Dublin 9, Ireland^{1, 2}

Abstract

We derive a large-time large deviation principle for the log stock price under an uncorrelated stochastic volatility model. For this we use a Donsker-Varadhan-type large deviation principle for the occupation measure of the Ornstein-Uhlenbeck process, combined with a simple application of the contraction principle and exponential tightness.

Key words: Stochastic volatility, Donsker-Varadhan large deviations for occupation measures, Large time implied volatility asymptotics.

1. The uncorrelated Ornstein-Uhlenbeck model with σ bounded

We work on a model $(\Omega, \mathcal{F}, \mathbb{P})$ with a filtration $(\mathcal{F}_t)_{t \geq 0}$ supporting two Brownian motions which satisfies the usual conditions.

Set $f(y) = \sigma^2(y)$, and assume that $0 < f_{\min} \leq f \leq f_{\max} < \infty$. We consider an uncorrelated stochastic volatility model for a log stock price process $X_t = \log S_t$ defined by the following stochastic differential equations

$$\begin{cases} dX_t = -\frac{1}{2}\sigma^2(Y_t)^2 dt + \sigma(Y_t)dW_t^1, \\ dY_t = -\alpha Y_t dt + dW_t^2, \end{cases} \quad (1)$$

for $\alpha > 0$, $X_0 = x_0, Y_0 = y_0$, where W_1, W_2 are two independent standard Brownian motions and Y is an Ornstein-Uhlenbeck process. We set $S_0 = 1$ (i.e. $x_0 = 0$) without loss of generality, because $X_t - x_0$ is independent of x_0 as the SDEs have no dependence on x .

¹work supported by SFI Grant for the Edgeworth centre for Financial Mathematics.

²We thank one anonymous referee for feedback on the article.

1.1. Large deviations for the occupation measure of the OU process

For each $t > 0$ and $A \in \mathcal{B}(\mathbb{R})$, let

$$\mu_t(A) = \frac{1}{t} \int_0^t 1_A(Y_s) ds \quad (2)$$

denote the proportion of time up to t that the sample path of Y spends in A . For each $t > 0$ and ω , $\mu_t(\omega, \cdot)$ is a probability measure on \mathbb{R} . Let $\mathcal{P}(\mathbb{R})$ denote the space of probability measures on \mathbb{R} . Then $\mu_t(A)$ satisfies a large-time large deviation principle in the topology of weak convergence, with convex, lower semicontinuous rate function given by

$$I_B(\mu) = \frac{1}{2} \int_{-\infty}^{\infty} |\partial_y \sqrt{\frac{d\mu}{d\mu_\infty}}(y)|^2 \mu_\infty(dy) \quad (3)$$

for $\mu \in \mathcal{P}(\mathbb{R})$, where $\mu_\infty(y) = \left(\frac{\alpha}{\pi}\right)^{\frac{1}{2}} e^{-\alpha y^2}$ is the unique stationary distribution for Y , i.e. $N(0, 1/2\alpha)$ (see Donsker and Varadhan (1975a, 1975b, 1975c, 1976), Stroock (1984) and pages 367-8 in Feng and Kurtz (2006)). If μ is not absolutely continuous with respect to μ_∞ , then $I_B(\mu) = \infty$.

Remark 1.1. Clearly $I_B(\mu)$ attains its minimum value of zero at $\mu = \mu_\infty$. Moreover, any measure which makes the rate function zero is a stationary distribution, and Y has a unique stationary distribution, so μ_∞ is the unique minimizer of $I_B(\mu)$.

We note that $F : \mathcal{P}(\mathbb{R}) \mapsto [f_{\min}, f_{\max}]$ given by

$$F(\mu) = \langle f, \mu \rangle = \int_{-\infty}^{\infty} f(y) \mu(dy) \quad (4)$$

is a bounded, continuous functional.

1.2. The Prokhorov metric

Given two measures μ and ν in $\mathcal{P}(\mathbb{R})$, the Prokhorov metric is defined by $d(\mu, \nu) = \inf\{\delta > 0 : \mu(C) \leq \nu(C^\delta) + \delta, \nu(C) \leq \mu(C^\delta) + \delta \text{ for all closed } C \subset \mathbb{R}\}$. $\mathcal{P}(\mathbb{R})$ then becomes a compact metric space (note that $d(\mu, \nu) \leq 1$), so the rate function $I_B(\mu)$ is *good*. Convergence of measures in the Prokhorov metric is equivalent to weak convergence of measures.

1.3. Contraction principle

By the contraction principle, the quantity

$$A_t = \frac{1}{t} \int_0^t f(Y_s) ds = \int_{-\infty}^{\infty} f(y) \mu_t(dy)$$

also satisfies the LDP, with good lower semicontinuous rate function given by

$$I_f(a) = \inf_{\mu \in \mathcal{P}(\mathbb{R}): \langle f, \mu \rangle = a} I_B(\mu) \quad , \quad a \in [f_{\min}, f_{\max}]. \quad (5)$$

Remark 1.2. $I_B(\cdot)$ is non-negative and $I_B(\mu_\infty) = 0$, so

$$I_f(\bar{\sigma}^2) = 0,$$

where

$$\bar{\sigma}^2 = \langle f, \mu_\infty \rangle = \int_{-\infty}^{\infty} \sigma^2(y) \mu_\infty(y) dy. \quad (6)$$

Moreover, μ_∞ is the unique minimizer of I_B , so $\bar{\sigma}^2$ is the unique minimizer of I_f .

2. A joint large deviation principle for $(X_t/t, A_t)$

Recall the definition of $A_t = \frac{1}{t} \int_0^t f(Y_s) ds$. The following proposition establishes a joint LDP for X_t/t and A_t in the large-time limit.

Proposition 2.1. $(X_t/t, A_t)$ satisfies a joint LDP as $t \rightarrow \infty$ with good rate function $I(x, a) = \frac{(x + \frac{1}{2}a)^2}{2a} + I_f(a)$.

Proof. See Appendix. □

From this we obtain the following proposition:

Proposition 2.2. (X_t/t) satisfies the LDP as $t \rightarrow \infty$ with a good rate function given by

$$I(x) = \inf_{a \in [f_{\min}, f_{\max}]} \left[\frac{(x + \frac{1}{2}a)^2}{2a} + I_f(a) \right] \leq \frac{(x + \frac{1}{2}\bar{\sigma}^2)^2}{2\bar{\sigma}^2}$$

and I attains its minimum value of zero at $x = -\frac{1}{2}\bar{\sigma}^2$.

Proof. The LDP with a good rate function just follows from the contraction principle. Setting $a = \bar{\sigma}^2$ defined in 6 and using that $I_f(\bar{\sigma}^2) = 0$, we see that $I(-\frac{1}{2}\bar{\sigma}^2) = 0$. \square

Remark 2.1 For non-zero correlation and/or unbounded σ , the approach outlined here will not work. However, we can transform the problem to a *small-noise, fast mean-reverting regime*, which is the same scaling used in the recent paper by Feng et al. (2010), aside from the fact that the drift of the log Stock price process is not small in this case. This problem then falls into the class of homogenization and averaging problems for nonlinear HJB type equations, where the fast volatility variable lives on a non-compact space. The Feng et al. argument based on viscosity solutions can be easily adapted to the large-time regime, using Bryc's lemma combined with exponential tightness to prove a large deviation principle. The leading order term is the unique viscosity solution to a HJB equation where the Hamiltonian is given in terms of the limiting log mgf for the integrated variance; this will be dealt with in a sequel article. For the well known SABR model with $\beta = 1$, we can derive large-time asymptotics for the correlated case using the Willard mixing formula, see Forde (2010).

References

- [1] Dembo, A. and Zeitouni, O., 1998. Large deviations techniques and applications. Jones and Bartlet publishers, Boston.
- [2] Donsker, M.D. and Varadhan, S.R.S., 1975. On a variational formula for the principal eigenvalue for operators with maximum principle., Proc. Nat. Acad. Sci. USA. 72 No.3, pp. 780-783.
- [3] Donsker, M.D. and Varadhan, S.R.S., 1975. Asymptotic evaluation of Markov process expectations for large time, I. Comm. Pure Appl. Math. 28, pp.1-47.
- [4] Donsker, M.D. and Varadhan, S.R.S., 1975. Asymptotic evaluation of Markov process expectations for large time, I. Comm. Pure Appl. Math. 28, pp.279-301.

- [5] Donsker, M.D. and Varadhan, S.R.S., 1976. Asymptotic evaluation of Markov process expectations for large time. III. *Comm. Pure Appl. Math*, 29, pp.389-461.
- [6] Feng, J. and Kurtz, T.G., 2006. *Large Deviations for Stochastic Processes*. American Mathematical Society.
- [7] Feng, J., Fouque, J.P. and Kumar, R., 2010. Small-time asymptotics for fast mean-reverting stochastic volatility models. Forthcoming in *Annals of Applied Probability*.
- [8] Forde, M., 2010. The Large-maturity smile for the SABR model with non-zero correlation. Submitted.
- [9] Forde, M. and Jacquier, A., 2009. The Large-maturity smile for the Heston model. Forthcoming in *Finance and Stochastics*.
- [10] Stroock, D.W., 1984. *An introduction to the theory of large deviations*. Springer-Verlag, Berlin.

A. Proof of Proposition 2.1

Let $Z_t = X_t/t$. We first note that $(Z_t, A_t) \stackrel{d}{=} (W_{tA_t} - \frac{1}{2}A_t, A_t)$. We first assume $x + \frac{1}{2}a > 0$. Now choose δ so that $0 < \delta < x + \frac{1}{2}a$. Then

$$\begin{aligned} \mathbb{P}(|Z_t - x| < \frac{\delta}{\sqrt{2}}, |A_t - a| < \frac{\delta}{\sqrt{2}}) &\leq \mathbb{P}(\|(Z_t, A_t) - (x, a)\| < \delta) \\ &\leq \mathbb{P}(|Z_t - x| < \delta, |A_t - a| < \delta). \end{aligned}$$

From the Gärtner-Ellis theorem, we can verify that $\frac{W_{at}^1 - \frac{1}{2}at}{t}$ satisfies the LDP as $t \rightarrow \infty$ with rate $\frac{(x + \frac{1}{2}a)^2}{2a}$. Then $\forall \epsilon > 0$, conditioning on A_t and using the LDPs for A_t and $\frac{W_{at}^1 - \frac{1}{2}at}{t}$, we see that there exists a $t = t^*(\epsilon, \delta)$ such that

$$\mathbb{P}(|Z_t - x| < \delta, |A_t - a| < \delta) \leq e^{-t[-\epsilon + \inf_{y \in B_\delta(x)} \frac{(y + \frac{1}{2}(a - \delta))^2}{2(a + \delta)}]} e^{-t[-\epsilon + \inf_{a_1 \in \bar{B}_a(\delta)} I_f(a_1)]}.$$

Then

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(|Z_t - x| < \delta, |A_t - a| < \delta) \leq - \inf_{y \in B_\delta(x)} \frac{(y + \frac{1}{2}(a - \delta))^2}{2(a + \delta)} - \inf_{a_1 \in \bar{B}_a(\delta)} I_f(a_1),$$

and by the lower semicontinuity of $I_f(a)$ we have $\lim_{\delta \rightarrow 0} \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(|Z_t - x| < \delta, |A_t - a| < \delta) \leq -[\frac{(x + \frac{1}{2}a)^2}{2a} + I_f(a)]$. Using a similar argument for the lower bound, we replace the limsup here by a genuine limit, so (Z_t, A_t) satisfies the weak LDP with rate function $\frac{(x + \frac{1}{2}a)^2}{2a} + I_f(a)$. $I_f(a)$ is good, so (A_t) is exponentially tight; hence for all $R > 0, a > 0$, there exists a compact $K_a \subset \mathbb{R}$ such that $\limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}((Z_t, A_t) \in [-R, R] \times K_a^c) \leq \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(A_t \in K_a^c) \leq -a$, so (Z_t, A_t) is exponentially tight, hence (Z_t, A_t) satisfies the full LDP and the rate function is good. We proceed similarly for $x + \frac{1}{2}a \leq 0$.