



**HAL**  
open science

## Multiyear surveillance of Influenza A virus in wild birds in Portugal

Ana Margarida Henriques, Teresa Fagulha, Sílvia C Barros, Fernanda Ramos,  
Margarida Duarte, Tiago Luis, Miguel Fevereiro

► **To cite this version:**

Ana Margarida Henriques, Teresa Fagulha, Sílvia C Barros, Fernanda Ramos, Margarida Duarte, et al.. Multiyear surveillance of Influenza A virus in wild birds in Portugal. *Avian Pathology*, 2011, 40 (06), pp.597-602. 10.1080/03079457.2011.618943 . hal-00755787

**HAL Id: hal-00755787**

**<https://hal.science/hal-00755787>**

Submitted on 22 Nov 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Multiyear surveillance of Influenza A virus in wild birds in Portugal**

Journal:	<i>Avian Pathology</i>
Manuscript ID:	CAVP-2011-0062.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	04-Jul-2011
Complete List of Authors:	Henriques, Ana; Laboratorio Nacional de Investigacao Veterinaria (LNIV), Virologia Fagulha, Teresa; Laboratorio Nacional de Investigacao Veterinaria (LNIV), Virologia Barros, Sílvia; Laboratorio Nacional de Investigacao Veterinaria (LNIV), Virologia Ramos, Fernanda; Laboratorio Nacional de Investigacao Veterinaria (LNIV), Virologia Duarte, Margarida; Laboratorio Nacional de Investigacao Veterinaria (LNIV), Virologia Luis, Tiago; Laboratorio Nacional de Investigacao Veterinaria (LNIV), Virologia Fevereiro, Miguel; Laboratorio Nacional de Investigacao Veterinaria (LNIV), Virologia
Keywords:	Influenza A virus, Avian Influenza, AIV, wild birds

SCHOLARONE™  
Manuscripts

Cavp-2011-0062.R1

Multiyear surveillance of Influenza A virus in wild birds in Portugal

**Ana M. Henriques, Teresa Fagulha, Sílvia C. Barros, Fernanda Ramos, Margarida Duarte, Tiago Luís and Miguel Fevereiro\***

*Laboratório Nacional de Investigação Veterinária, Departamento de Virologia, Lisboa, Portugal*

Short Title: Avian influenza in wild birds in Portugal

\*Corresponding Author

Tel.: +351217115288

Fax: +351217115387

E-mail: [miguel.fevereiro@Iniv-min.agricultura.pt](mailto:miguel.fevereiro@Iniv-min.agricultura.pt)

*Received: 4 July 2011*

## Abstract

This report presents the results of a multiyear (2005-2009) study of avian influenza virus (AIV) occurrence in wild birds in Portugal. A total of 5691 samples from wild birds belonging to 13 different Orders were examined. Ninety three samples tested positive for AIV by matrix RT-PCR, giving a total prevalence of 1.63 %. Twenty one viruses were successfully cultured in embryonated chicken eggs, which represent a rate of viral infectivity of 22.6 % in the samples. Nine subtypes of hemagglutinin (H1, H3-H7, H9-H11) and eight subtypes of neuraminidase (N1-N4, N6-N9) were identified in 20 different combinations. The most prevalent subtypes of hemagglutinin detected were H5, H1 and H4 while for neuraminidase, subtypes N2 and N6 were the most common. The subtype combinations H4N6 and H1N1 were predominant (15.1 %). All H5 and H7 viruses detected in this study were low pathogenic for poultry as determined by the sequence of amino acids at the cleavage site of hemagglutinin. The full-length nucleotide sequences of five H5, one H7 and five N3 genes were phylogenetically analyzed. The Bayesian analysis revealed that all but one of the strains analyzed were closely related to isolates detected in the same period in North and Central European countries. Three H5N3 isolates, all from 2007, formed a separate cluster in both H5 and N3 phylogenetic trees. This study provides evidence that various subtypes of avian influenza virus, including subtypes H5 and H7, circulate in Portugal which may pose a risk to industrial poultry.

## Introduction

The influenza A virus, and most of its hemagglutinin (HA) and neuraminidase (NA) subtype combinations, is commonly present in wild bird species of wetlands and aquatic environments, with most infections thought to be inapparent (Swayne & Suarez, 2000; Webster *et al.*, 1992). Clinically, AI viruses can be classified into two main pathotypes (Capua & Alexander, 2004); the highly pathogenic avian influenza (HPAI) viruses that cause a systemic disease in which mortality may approach 100 % in fowl; and low pathogenic avian influenza (LPAI) viruses that cause a much milder disease in susceptible poultry. HPAI is caused only by certain strains of H5 and H7 viruses.

The segmented nature of AIV genome enables viral evolution by genetic reassortment. A recent study indicates that genetic sublineages do not persist, but frequently reassort with other viruses (Munster *et al.*, 2007). Gaining information on the actual frequency of reassortment in the wild bird population and the impact of these events on LPAI virus evolution is of relevant interest.

Avian influenza viruses (AIV) have diversified into two separate avian lineages, North American and European, probably as a consequence of long-term ecological and geographical separation of the hosts (Krauss *et al.*, 2004; Olsen *et al.*, 2006). Several multiyear studies have been made in countries of Northern Europe in wild birds (Fouchier *et al.*, 2003; Munster *et al.*, 2007; Wallensten *et al.*, 2007) but to our knowledge no such study has been conducted in Portugal so far. Situated in the east Atlantic migratory flyway the country is visited by many species that migrate annually between Africa and the Iberian Peninsula, mainly from March to July, which corresponds with spring migration and breeding season for birds. On the other hand, wintering migratory birds, especially waterfowl, come predominantly from the northern regions of Europe. These two migratory populations mingle with resident free living birds which may favor the introduction of new AIV in the country. In the frame of avian

influenza preparedness, a surveillance program was carried out in wild birds aiming to assess the risk of AIV introduction, either of low (LPAI) or high pathogenicity (HPAI) to domestic poultry. This study reports the epidemiological and virological findings of AIV occurrence in wild birds in Portugal from January 2005 to December 2009.

## Materials and Methods

**Collection of samples.** The majority of samples originated from migratory or resident wild birds and was collected in bird sanctuaries located in the vicinity of poultry farms. The birds were caught throughout the year. A total of 5691 samples consisting of swabs collected from the oropharynx and cloaca of wild birds belonging to thirteen different orders were analyzed during the five years study (Table 1). The swabs were transported dried and kept at 5°C to 8°C until tested.

**AIV screening.** The content of the oropharynx and cloaca swabs was eluted and pooled into vials containing 2 ml of cold PBS with antibiotics (2000 u/ml penicillin, 2 mg/ml streptomycin and 50 µg/ml gentamycin) and submitted to RNA extraction on a nucleic acid extraction workstation BioSprint 96, following the manufacturer's protocol (Qiagen). The samples were firstly screened for AIV by matrix (M) gene rRT-PCR using the method described by Spackman *et al.* (Spackman *et al.*, 2002) and Qiagen One-step rRT-PCR kit in 25 µl final volume. A positive control was included in each assay.

**Virus subtyping.** Samples matrix rRT-PCR positives were further screened for subtypes H5 and H7, using rRT-PCR protocols recommended by the EU Community Reference Laboratory (Slomka, Pavlidis *et al.*, 2007; Slomka *et al.*, 2009). To determine all other AIV subtypes, the RNA obtained from swabs or tissues was used to partially amplify the HA and NA genes by conventional RT-PCR with universal primers described by Hoffmann *et al.* (Hoffmann *et al.*, 2001) and primers designed in our laboratory (NA-F: CAC CAT GGA TCC AAA TCA GAA GAT; N7-F: CAC CAT GGA TCC AAA TC AGA AAC TAT TTG; NeurUniR: CCD AWY CTD AYN SCR TTG TCY TT), respectively. The RT-PCR products were analyzed by 2 % (w/v) agarose gel electrophoresis and the amplicons were purified after being excised from the gel using Qiagen QIAquick gel extraction kit as described by the manufacturer. Nucleotide sequencing was carried out with the BigDye DNA sequencing kit (Applied Biosystems (AB)) in the Applied Biosystems 3130 genetic analyzer.

**Viruses Pathotyping.** The RNA obtained from swabs or tissues was used to accomplish the pathotyping of H5 and H7 viruses. The genomic region encoding the HA protein cleavage site was amplified with primer pairs recommended by the EU Community Reference Laboratory, Kha1/Kha3 for H5 and GK7.3/ GK7.4 for H7 (Slomka, Coward *et al.*, 2007) and the nucleotide sequences of the purified PCR products were determined as described above.

**Virus culture.** Virus isolation in specific-pathogen-free 11 day-old embryonated chicken eggs was also attempted with all matrix rRT-PCR positive samples. Samples tested negative after first-passage, were passaged two more times to confirm the absence of AIV. Haemagglutinating positive allantoic fluids were tested against

reference sera to H5, H7 and Newcastle disease virus (as described in chapter V of the EU Diagnostic manual in accordance with EU Directive 2005/94) and for the M, H5 and H7 genes by rRT-PCR as described above.

**Phylogenetic analysis.** The complete nucleotide sequences of five H5, one H7 and five N3 genes were determined and phylogenetically analyzed. The viruses included in the study were selected after blast analysis and comprised not only European strains showing high homology to sequenced viruses but also viruses from other parts of the world. Only complete sequences were used in the analysis. The phylogenetic trees were obtained with a Bayesian inference of phylogeny throughout the MrBayes v3.1.2 software, using the GTR model (nst=6) with gamma-shaped rate variation with a proportion of invariable sites (rates=invgamma). The analysis used ngen=106, nchains=4 and samplefreq=10.

**Accession numbers.** The GenBank accession numbers HM849026 to HM849030 and HM849003 represent the H5 and H7 genes of 28003/H5N3, 28006/H5N3, 29497-10/H5N2, 37369/H5N3, 4228/H5N3, and 13771/H7N3, respectively. The N3 genes of 13771/H7N3, 28003/H5N3, 28006/H5N3, 37369/H5N3 and 4228/H5N3 were given respectively accession numbers HM849005 and HM849031 to HM849034.

## Results and Discussion

**AIV detection and virus isolation.** Total prevalence of influenza A virus in the birds sampled during the five-year period was 1.63 %. The large proportion (88.2 %) of the



93 AIV-matrix gene rRT-PCR positive samples was from mallards, and only eleven samples came from other species of ducks, and from seagulls, storks, partridges and flamingos. Active monitoring reached 70 % of the sampling and 76.3 % of the rRT-PCR positive samples. These data demonstrate the importance of active monitoring in the success of avian influenza surveillance. The main places of the sampling as well as the districts where the positive birds were found are shown in Figure 1. Figure 2 represents the temporal distribution of sampling and rRT-PCR positive samples.

Twenty one viruses were successfully cultured in embryonated chicken eggs, which represent a rate of viral infectivity (no viruses isolated / no rRT-PCR positive samples) of 22.6 % in the samples (Table 2). The swabs were transported dried which may have affected the success of AIV isolation. However, the low rate of virus isolation recorded in present study is consistent with that found in other studies (Dusek *et al.*, 2009; Munster *et al.*, 2007). Usually, rRT-PCR is more sensitive than virus isolation, enabling detection of genome fragments of non-infective viruses.

**Virus subtyping.** In this study, nine subtypes of HA (H1, H3-H7, H9-H11) and eight subtypes of NA (N1-N4, N6-N9) were identified in 20 different combinations (Figure 3). The most prevalent subtypes of HA detected were H5 (23.7 %), H1 and H4 (21.5 %). The high prevalence of H5 subtype is unusual and different to that found in other studies in Europe and North America. However, this fact can be explained by the increased sampling of a population of mallards semi-resident in a natural reserve, after the detection of H5N2. As in other studies (Krauss *et al.*, 2004; Suss *et al.*, 1994; Wallensten *et al.*, 2007), H4 and H1 also dominated followed by H9 and H6, while subtypes H2, H8, H12-H16 were never detected in this study. Concerning NA the most prevalent subtype was N2 (32.3 %), followed by N6 (17.2 %). With exception of N5, all

other known subtypes of NA were found. The subtype combinations H4N6 and H1N1 (15.1 %) were the most common, similar in rates previously reported in Sweden and Germany (Suss *et al.*, 1994; Wallensten *et al.*, 2007). Our study also shows that H5N3 and H5N2 combinations were frequently detected in Portugal, with prevalence of 10.8 % and 7.5 %, respectively.

**Viruses Pathotyping.** The twenty two H5 and four H7 AIV identified in this study were found to be of low pathogenicity for poultry since they lack basic amino acids residues at the cleavage site of HA. The pathogenicity of four isolates was also assessed by plaque forming assays in MDCK cells in the presence and absence of trypsin. The isolates tested, one H7N3 and three H5N3, required trypsin for plaque formation, thus corroborating the molecular prediction of low pathogenicity.

**Nucleotide sequence analysis.** The molecular analysis of non-structural 1 (NS1) proteins and of the stalk region of neuraminidase of these strains revealed full-length proteins, typical of viruses from wild birds (Matrosovich *et al.*, 1999). In addition, all viruses studied possess Asp92 and the ESEV amino acid motif at the C-terminal of NS1 protein, typical of avian influenza viruses (Dundon *et al.*, 2006; Soubies *et al.*, 2010). Analysis of matrix 2 protein (M2) of fully sequenced isolates showed that none of the isolates were resistant to amantidine.

**Phylogenetic analysis.** Phylogenetic analysis was performed with the complete nucleotide sequence of H5 gene of five AIV detected during this study (Figure 4). The analysis revealed that the H5 isolates from this study were closely related to LPAI strains from the North and Central European countries and more distantly related to

African, Asian and American viruses. Three strains detected in 2007 (28003/H5N3, 28006/H5N3 and 37369/H5N3) were more closely related to each other and to German isolate A/mallard/Bavaria/35/2006(H5) with 99 % nucleotide identity. The other two H5 viruses (4228/H5N3 and 29497-10/H5N2) showed a close relationship with isolates from Sweden and Switzerland (Figure 4A). The identity of the nucleotide sequences of the H5 viruses reported here ranged between 93 % and 99 %. The 2006 H7 isolate (13771/H7N3) was also analyzed, showing greatest nucleotide identity (98% to 99%) with isolates from Hungary, Netherlands, Sweden and United Kingdom detected between 2005 and 2007 (data not shown).

Five neuraminidases of subtype N3 chosen at random were fully sequenced and analyzed phylogenetically. Four of them belonged to H5 strains from ducks and one to a H7 virus isolated in a tufted duck. The nucleotide variation within the N3 sequences of these five viruses ranged between 94 % and 99 %. Interspersed clusters of European and Asian viruses were observed. American and Oceanian viruses formed a distinct group, as expected. With the exception of 4228/H5N3, all other strains from this study clustered with European viruses. Three H5N3 viruses (28003/H5N3, 28006/H5N3 and 37369/H5N3) detected in 2007 and with different geographic origins, formed a separate branch with a French isolate of 2002. Isolate 13771/H7N3 branched with Swiss and Dutch isolates (Figure 4B).

It was interesting to note that H5N3 strains 28003, 28006 and 37369, all from 2007 but with different geographic origins, clustered together and formed a distinct sub-cluster in both H5 and N3 dendograms. Also of note was strain 4228/H5N3 that may be a reassortant virus since H5 and N3 genes clustered with European and Asian viruses respectively. The North American and Eurasia virus lineages were clearly evidenced both in H5 and N3 analysis.

The results of this study indicate the active circulation of influenza A viruses in wild birds in Portugal. Amongst all wild birds sampled, mallards harbored the majority of AIV obtained during this multiyear surveillance. The greater susceptibility of mallards to avian influenza infections has also been reported in other studies (Olsen *et al.*, 2006; Webster *et al.*, 1992) being this species considered the main target for sampling in AI surveillance programs. The isolation of H5 and H7 subtypes and the geographical vicinity between wild bird sanctuaries and poultry farms represents a risk for industrial poultry of the country.

This study also provides the genotypic analysis and a detailed phylogenetic data of several strains collected during a five-year period. Compiling sequence data and the identification of new viruses and their relationships will contribute to a better understanding of the evolution of the viruses and also for the establishment of an updated plan for molecular diagnosis of influenza A viruses.

### **Acknowledgments**

We are grateful to Dr. David Rodrigues for the submission of numerous samples from mallards. We thank Arminda Batistae Ana Almeida for excellent technical support. We are thankful to Portuguese Fundação para a Ciência e a Tecnologia (FCT) for Contrato-Programa “Ciência 2008” to A. M. Henriques. Financial assistance for this study was partially obtained from FCT; project CVT/67189/2006. The authors acknowledge technical support received from FLU-LAB-NET - “Development and enhancement of Laboratory Networks for Avian Influenza” (Contract no SSPE-044453).

## References

- Capua, I. & Alexander, D.J. (2004). Avian influenza: recent developments. *Avian Pathology*, 33, 393-404.
- Dundon, W.G., Milani, A., Cattoli, G. & Capua, I. (2006). Progressive truncation of the Non-Structural 1 gene of H7N1 avian influenza viruses following extensive circulation in poultry. *Virus Research*, 119, 171-176.
- Dusek, R.J., Bortner, J.B., DeLiberto, T.J., Hoskins, J., Franson, J.C., Bales, B.D., *et al.* (2009). Surveillance for high pathogenicity avian influenza virus in wild birds in the Pacific Flyway of the United States, 2006-2007. *Avian Diseases*, 53, 222-230.
- Fouchier, R.A., Olsen, B., Bestebroer, T.M., Herfst, S., van der Kemp, L., Rimmelzwaan, G.F., *et al.* (2003). Influenza A virus surveillance in wild birds in Northern Europe in 1999 and 2000. *Avian Diseases*, 47, 857-860.
- Hoffmann, E., Stech, J., Guan, Y., Webster, R.G. & Perez, D.R. (2001). Universal primer set for the full-length amplification of all influenza A viruses. *Archives of Virology*, 146, 2275-2289.
- Krauss, S., Walker, D., Pryor, S.P., Niles, L., Chenghong, L., Hinshaw, V.S., *et al.* (2004). Influenza A viruses of migrating wild aquatic birds in North America. *Vector Borne Zoonotic Diseases*, 4, 177-189.
- Matrosovich, M., Zhou, N., Kawaoka, Y. & Webster, R. (1999). The surface glycoproteins of H5 influenza viruses isolated from humans, chickens, and wild aquatic birds have distinguishable properties. *Journal of Virology*, 73, 1146-1155.

- Munster, V.J., Baas, C., Lexmond, P., Waldenstrom, J., Wallensten, A., Fransson, T., *et al.* (2007). Spatial, temporal, and species variation in prevalence of influenza A viruses in wild migratory birds. *PLoS Pathogens*, 3, e61.
- Olsen, B., Munster, V.J., Wallensten, A., Waldenstrom, J., Osterhaus, A.D. & Fouchier, R.A. (2006). Global patterns of influenza a virus in wild birds. *Science*, 312, 384-388.
- Slomka, M.J., Coward, V.J., Banks, J., Londt, B.Z., Brown, I.H., Voermans, J., *et al.* (2007). Identification of sensitive and specific avian influenza polymerase chain reaction methods through blind ring trials organized in the European Union. *Avian Dis*, 51, 227-234.
- Slomka, M.J., Pavlidis, T., Banks, J., Shell, W., McNally, A., Essen, S., *et al.* (2007). Validated H5 Eurasian real-time reverse transcriptase-polymerase chain reaction and its application in H5N1 outbreaks in 2005-2006. *Avian Dis*, 51, 373-377.
- Slomka, M.J., Pavlidis, T., Coward, V.J., Voermans, J., Koch, G., Hanna, A., *et al.* (2009). Validated RealTime reverse transcriptase PCR methods for the diagnosis and pathotyping of Eurasian H7 avian influenza viruses. *Influenza Other Respi Viruses*, 3, 151-164.
- Soubies, S.M., Volmer, C., Croville, G., Loupiau, J., Peralta, B., Costes, P., *et al.* (2010). Species-specific contribution of the four C-terminal amino acids of influenza A virus NS1 protein to virulence. *Journal of Virology*, 84, 6733-6747.
- Spackman, E., Senne, D.A., Myers, T.J., Bulaga, L.L., Garber, L.P., Perdue, M.L., *et al.* (2002). Development of a real-time reverse transcriptase PCR assay for

type A influenza virus and the avian H5 and H7 hemagglutinin subtypes.

*Journal of Clinical Microbiology*, 40, 3256-3260.

Suss, J., Schafer, J., Sinnecker, H. & Webster, R.G. (1994). Influenza virus subtypes in aquatic birds of eastern Germany. *Archives of Virology*, 135(1-2), 101-114.

Swayne, D.E. & Suarez, D.L. (2000). Highly pathogenic avian influenza. *Revue Scientifique et Technique* 19, 463-482.

Wallensten, A., Munster, V.J., Latorre-Margalef, N., Brytting, M., Elmberg, J., Fouchier, R.A., *et al.* (2007). Surveillance of influenza A virus in migratory waterfowl in northern Europe. *Emerging Infectious Diseases*, 13, 404-411.

Webster, R.G., Bean, W.J., Gorman, O.T., Chambers, T.M. & Kawaoka, Y. (1992). Evolution and ecology of influenza A viruses. *Microbiological Reviews*, 56, 152-179.

**Table 1.** *Total of wild birds screened for influenza A virus (2005-2009)*

Bird Order	2005	2006	2007	2008	2009	Total
Accipitriformes	54	83	14	18	33	202
Anseriformes	361	1479	637	787	297	3561
Charadriiformes	6	58	70	12	181	327
Ciconiiformes	70	208	37	17	14	346
Falconiformes	27	28	3	6	12	76
Galiformes	6	0	0	1	0	7
Gruiformes	6	57	0	12	30	105
Passeriformes	32	170	7	66	11	286
Pelecaniformes	26	135	26	4	3	194
Phoenicopteriformes	23	159	3	2	1	188
Podicipediformes	2	6	1	0	0	9
Procellariiformes	5	17	35	90	109	256
Strigiformes	29	63	10	8	24	134
Total	647	2463	843	1023	715	5691



**Table 2.** AIV prevalence and subtypes found in wild birds (2005-2009)

Common name (Scientific name)	Tested	No of positives (%)	Viable virus (%)	Subtype (no of positives)
Mallard ( <i>Anas platyrhynchos</i> )	1495	82 (5.5)	20 (24.4)	H1N1 (14), H1N2 (3), H1N6 (1), H1N? (1), H3N2 (1), H3N3 (1), H3N8 (1), H4N2 (5), H4N6 (14), H4N? (1), H5N2 (6), H5N3 (9), H5N6 (1), H5N? (3), H6N2 (1), H6N7 (1), H6N? (1); H7N3 (1); H9N2 (6), H9N4 (1), H9N? (1); H10N2 (2), H11N9 (5), H?N2 (2)
Common teal ( <i>A. crecca</i> )	204	3 (1.5)	0	H1N2 (1), H7N7 (1), H9N2 (1)
Tufted duck ( <i>Aythya fuligula</i> )	24	1(4.2)	1 (100)	H7N3 (1)
Seagull ( <i>Larus sp.</i> )	522	3 (0.6)	0	H6N2 (1), H6N3 (1), H6N? (1)
Flamingo ( <i>Phoenicopterus sp.</i> )	194	1 (0.5)	0	H5N2 (1)
Partridge ( <i>Perdix perdix</i> )	227	2 (0.9)	0	H5N3 (1); H5N? (1)
Stork ( <i>Ciconia ciconia</i> )	178	1 (0.6)	0	H7 N? (1)


## Figure legends

**Figure 1.** *Map of Portugal presenting the distribution of the sampling by district. The number of positive samples, with percentage in brackets, is indicated.*


**Figure 2.** *Temporal distribution of samples. The bars represent the sampling distribution. The number of positive birds, with percentage in brackets, is indicated.*

**Figure 3.** *The HA and NA subtypes of AIV found in wild birds in Portugal during 2006-2009. The numbers above the bars in the graphs represent the percentage of positive samples.*

**Figure 4.** *Bayesian analysis of the complete nucleotide sequences of H5 (A) and N3 (B) genes of influenza A virus detected in Portugal in wild ducks during 2006-2009. The phylogenetic trees were obtained with a Bayesian inference of phylogeny throughout the MrBayes v3.1.2 software, using the GTR model ( $nst=6$ ) with gamma-shaped rate variation with a proportion of invariable sites ( $rates=invgamma$ ). The analysis used  $ngen=10^6$ ,  $nchains=4$  and  $samplefreq=10$ . The numbers included on each boot strap represent the Bayesian posterior probability. Viruses reported in this study are presented in bold.*



**Figure 1:** Map of Portugal presenting the distribution of the sampling by district. The number of positive samples, with percentage in brackets, is indicated.



**Figure 2:** Temporal distribution of samples. The bars represent the sampling distribution. The number of positive birds, with percentage in brackets, is indicated.





Figure 3: The HA and NA subtypes of AIV found in wild birds in Portugal during 2006-2009. The numbers above the bars in the graphs represent the percentage of positive samples.



**Figure 4:** Bayesian analysis of the complete nucleotide sequences of H5 (A) and N3 (B) genes of influenza A virus detected in Portugal in wild ducks during 2006-2009. The phylogenetic trees were obtained with a Bayesian inference of phylogeny throughout the MrBayes v3.1.2 software, using the GTR model (nst=6) with gamma-shaped rate variation with a proportion of invariable sites (rates=invgamma). The analysis used ngen=10<sup>6</sup>, nchains=4 and samplefreq=10. The numbers included on each boot strap represent the Bayesian posterior probability.