



## Estimates for some Weighted Bergman Projections

Philippe Charpentier, Yves Dupain, Modi Mounkaila,

► **To cite this version:**

Philippe Charpentier, Yves Dupain, Modi Mounkaila,. Estimates for some Weighted Bergman Projections. 2012. hal-00761375v1

**HAL Id: hal-00761375**

**<https://hal.science/hal-00761375v1>**

Preprint submitted on 5 Dec 2012 (v1), last revised 23 May 2013 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ESTIMATES FOR SOME WEIGHTED BERGMAN PROJECTIONS

P. CHARPENTIER, Y. DUPAIN & M. MOUNKAILA

ABSTRACT. In this paper we investigate the regularity properties of weighted Bergman projections for smoothly bounded pseudo-convex domains of finite type in  $\mathbb{C}^n$ . The main result is obtained for weights comparable to a non negative rational power of the distance to the boundary: we prove Sobolev- $L^p$  and Lipschitz estimates for domains in  $\mathbb{C}^2$  and for convex domains.

## INTRODUCTION

Let  $\Omega$  be a bounded open set in  $\mathbb{C}^n$ . Let  $\omega$  be a non negative measurable function on  $\Omega$  and  $\lambda$  be the Lebesgue measure on  $\mathbb{C}^n$ . Then  $\omega$  is called an *admissible weight* (or simply a *weight*) for  $\Omega$  if the set  $A^2(\Omega, \omega d\lambda)$  of square integrable holomorphic functions with respect to the measure  $\omega d\lambda$  is a closed subspace of the Hilbert space  $L^2(\Omega, \omega d\lambda)$  (see [PW90]). Then, if  $\omega$  is a weight on  $\Omega$ , the weighted Bergman projection  $P_\omega^\Omega$ , i.e. the orthogonal projection of  $L^2(\Omega, \omega d\lambda)$  onto  $A^2(\Omega, \omega d\lambda)$ , is well defined.

The aim of this paper is to investigate Lipschitz and Sobolev  $L^p$  regularity of  $P_\omega^\Omega$  when  $\Omega$  is pseudo-convex.

A special class of weight has been introduced by L. Hörmander in [Hör65] for bounded pseudo-convex domain in  $\mathbb{C}^n$ . Let  $\varphi$  be a pluri-subharmonic function defined in  $\Omega$ . Hörmander's theorem solves the so-called  $\bar{\partial}_\varphi$ -Neumann problem associated to the weight  $e^{-\varphi}$  proving the existence of the Neumann operator  $\mathcal{N}_\varphi$  inverting the complex laplacian  $\square_\varphi$ . Then the Bergman projection  $P_{e^{-\varphi}}^\Omega$  is closely related to  $\mathcal{N}_\varphi$  by the formula  $P_{e^{-\varphi}}^\Omega = \text{Id} - \bar{\partial}^* \mathcal{N}_\varphi \bar{\partial}$  and regularity results for  $\mathcal{N}_\varphi$  can give regularity results for  $P_{e^{-\varphi}}^\Omega$ .

For  $\varphi = 0$  many work have been done in this direction in various function spaces. In particular, for the Bergman projection a lot of results have been obtained in [NRSW89, McN94, MS94, Cho96, Cho02, CD06b, CD08], and the theorems obtained in this paper are in the same spirit.

For  $(L^2)$  Sobolev regularity there is a very large bibliography essentially based on J. J. Koh's work (see [Str10] for a good general presentation). For other spaces, a lot of sharp results where obtained by several authors, but there are still basic open problems (see for example [CD08, CD06b, CD06a, CNS92, BC00, McN94, MS94, MS97] and references therein). If  $\varphi$  is  $\mathcal{C}^\infty$  on  $\bar{\Omega}$  it is not difficult to see that the operators associated to the  $\bar{\partial}_\varphi$ -Neumann problem have exactly the same regularity than the corresponding operators associated to the unweighted  $\bar{\partial}$ -Neumann problem.

For general (admissible) weights  $\omega$ , very few results where obtained for  $P_\omega^\Omega$ .

In the case of the unit ball of  $\mathbb{C}^n$ , for weights equal to a power greater than  $-1$  of the distance to the boundary, the kernels of these operators can be written explicitly (see [Cha80, HP84b, HP84a]) and then it is possible to obtain very sharp estimates. Generalizations of these results to strictly pseudo-convex domains have also been done by several authors (see [LR86, LR87, LR88, Cum90]).

For smoothly bounded pseudo-convex domains of finite type in  $\mathbb{C}^2$ , A. Bonami & S. Grellier ([BG95]) proved Lipschitz and Sobolev  $L^p$  estimates for the weighted Bergman projection when the weight is an entire power of a defining function of the domain. In [CL97], D. C. Chang & B. Q. Li generalize these results to "decoupled" domains in  $\mathbb{C}^n$ .

In general, even in dimension 1,  $L^p$  estimates for weighted Bergman projections can be true only for  $p = 2$ , and, in general are not easy to obtain as shown in [Zey11b, Zey11a, Zeya, Zeyb].

In this paper we will obtain sharp Lipschitz and Sobolev  $L^p$  estimates for weighted Bergman projection for a quite general class of weights in two cases: pseudo-convex domains of finite type in  $\mathbb{C}^2$  and convex domains of finite type in  $\mathbb{C}^n$ . In both cases, the main results will be obtain when the weight is "equivalent" to a rational power of the distance to the boundary. In particular this generalizes the results obtained by A. Bonami & S. Grellier and D. C. Chang & B. Q. Li. We will also study the Sobolev ( $L^2$ ) regularity of weighted  $\bar{\partial}$ -Neumann problem for general pseudo-convex domains of finite type in  $\mathbb{C}^n$ .

For the estimates of a weighted Bergman projection, a useful point is a result of Z. Pasternak-Winiarski ([PW90], see section 1) which implies that the weighted Bergman projections associated to two "equivalent" weights have the same regularity properties. Thus, to prove some estimate for the Bergman projections associated to the weights of a class of equivalent weights it suffices to prove it for one weight of the class.

The paper is organized as follows. In the first Section we present the main result on weighted Bergman projections and explain how Pasternak-Winiarski's result is used.

---

2000 *Mathematics Subject Classification.* 32F17, 32T25, 32T40.

*Key words and phrases.* pseudo-convex, finite type, Levi form locally diagonalizable, convex, extremal basis, geometric separation, weighted Bergman projection,  $\bar{\partial}_\varphi$ -Neumann problem.

All other Sections of the paper concern the construction of special weights for which we can prove some estimates for the Bergman projection or the  $\bar{\partial}$ -Neumann problem. In particular, for those weights we will obtain estimates not stated in the main results.

## 1. MAIN RESULTS AND METHODS

For simplicity, we only state here the main result concerning the Bergman projection for weights equivalent to a rational power of the distance to the boundary. Detailed results for other operators, other weights and for the Bergman kernel will be given in the next Sections.

If  $k$  is a positive function on an open set  $\Omega$  in  $\mathbb{C}^n$  whose inverse is locally integrable, we denote by  $P_k = P_k^\Omega$  the orthogonal projection of  $L^2(kd\lambda)$  onto the (closed [PW90]) subspace of holomorphic functions (i.e. the Bergman projection associated to the weight  $k$ ).

**Theorem 1.1.** *Let  $\Omega$  be a smoothly bounded pseudo-convex domain of finite type  $M$  satisfying one of the two following conditions:*

- $\Omega$  is a domain in  $\mathbb{C}^2$ ;
- $\Omega$  is convex.

*Let  $\omega$  be a positive function defined on  $\Omega$  for which there exists a non negative rational number  $r$  such that  $\omega = k\delta^r$ ,  $\delta$  being the distance to  $\partial\Omega$ . Let us denote by  $P_\omega^\Omega$  the weighted Bergman projection of  $\Omega$  associated to the weight  $\omega d\lambda$ ,  $d\lambda$  denoting the Lebesgue measure.*

- (1) *Let  $s \in \mathbb{N}$ . Assume that  $k$  is  $s$ -time differentiable,  $k^{(s)}$  is measurable and, for  $0 \leq i \leq s$ , there exist positive constants  $c_0 > 0$  and  $C_i > 0$ ,  $0 \leq i \leq s$ , such that  $c_0 \leq k \leq C_0$  and  $|k^{(i)}| \leq C_i$ ,  $1 \leq i \leq s$  on  $\Omega$ . Then, for  $p \in ]1, +\infty[$  and  $-1 < \beta < p(r+1) - 1$ ,  $P_\omega^\Omega$  maps continuously the Sobolev space  $L_s^p(\delta^\beta)$ ,  $s \in \mathbb{N}$ , into itself.*
- (2) *Assume that  $k$  is of class  $\mathcal{C}^1$  and that there exist constants  $c_0$  and  $C_i$ ,  $i = 0, 1$ , such that  $c_0 \leq k \leq C_0$  and  $|k'| \leq C_1$  on  $\Omega$ . Then, for  $\alpha < 1/M$ ,  $P_\omega^\Omega$  maps continuously the Lipschitz space  $\Lambda_\alpha$  into itself.*

The main step of the proof of this theorem together with other results (for other weights and all Lipschitz spaces) will be given in Sections 4.1, 4.2 and 4.3. The first step consists in showing that it suffices to prove the estimates for one special weight: this fact is based on a theorem of Z. Pasternak-Winiarski, which, for convenience, we state as follows:

**Theorem 1.2** ([PW90]). *Let  $\Omega$  be a bounded open set in  $\mathbb{C}^n$ . Let  $\omega$  be an admissible weight. Let  $h$  be a positive function on  $\Omega$  such that  $0 < c \leq h \leq C < 1/2$ . Then, denoting by  $M_h$  the operator  $g \mapsto hg$ ,*

$$P_{(1+h)\omega}^\Omega = P_\omega^\Omega + \sum_{k=1}^{\infty} (-1)^{k-1} \left[ P_\omega^\Omega \circ M_h \right]^k \circ (\text{Id} - P_\omega^\Omega),$$

where  $[...]^k$  is the power for the composition operator, the series converging in  $\mathcal{L}(L^2(\omega d\lambda))$  norm.

**Corollary.** *Assume that the conclusion of (1) or (2) of Theorem 1.1 is true for one admissible weight  $\omega_0$ . Then the same is also true for all weights  $\omega = k\omega_0$ , where the function  $k$  satisfies the properties described in Theorem 1.1.*

*Proof.* This is almost immediate. The hypothesis made on the weights implies that, for any  $\varepsilon > 0$ , there exists an integer  $N$  such that  $k = (1+h)^N$  with  $\|h\|_{\infty, s} \leq \varepsilon$ , where  $\|h\|_{\infty, s} = \max_{0 \leq i \leq s} \|h^{(i)}\|_\infty$ . Thus it is enough to show that if the Theorem is true for the weight  $\omega$  it is also true for the weight  $\omega' = \omega(1+h)$ .

Let us first consider the  $L_s^p(\delta^\beta)$  continuity and denote by  $\|\cdot\|_{p,s,\beta}$  the norm in  $L_s^p(\delta^\beta)$ . By hypothesis, there exists a constant  $K_{p,s,\beta}$  such that  $\|P_\omega^\Omega g\|_{p,s,\beta} \leq K_{p,s,\beta} \|g\|_{p,s,\beta}$ . Let  $f \in L_s^p(\delta^\beta)$ ; then  $f_1 = (\text{Id} - P_\omega^\Omega)f \in L_s^p(\delta^\beta)$  and  $\|f_1\|_{p,s,\beta} \leq (K_{p,s,\beta} + 1) \|f\|_{p,s,\beta}$ . For  $\varepsilon$  small enough and  $g \in L_s^p(\delta^\beta)$ , we have  $\|hg\|_{p,s,\beta} \leq \frac{1}{K_{p,s,\beta} + 1} \|g\|_{p,s,\beta}$ . Then writing

$$P_{(1+h)\omega}^\Omega f = P_\omega^\Omega f + \sum_{l=1}^{\infty} N_l(f),$$

where  $N_1(f) = P_\omega^\Omega(hf_1)$  and  $N_l(f) = P_\omega^\Omega(hN_{l-1}(f))$ ,  $l \geq 1$ , we get  $\|N_l(f)\|_{p,s,\beta} \leq \frac{1}{(K_{p,s,\beta} + 1)^{l-2}} \|f\|_{p,s,\beta}$ ,  $l \geq 2$ , proving the convergence of the series in  $L_s^p(\delta^\beta)$ .

The proof for the Lipschitz norm  $\|\cdot\|_\alpha$  is similar because

$$\|hf\|_\alpha \leq \left( \|h\|_{\infty, 1} \max(1, \text{diam}(\Omega)) + 2 \|h\|_\infty \right) \|f\|_\alpha.$$

□

In the next Sections, we construct special weights on a pseudo-convex domain of finite type  $\Omega$  for which we can prove estimates for  $P_\omega^\Omega$ , or for operators associated to a weighted  $\bar{\partial}$ -Neumann problem, using known results for unweighted Bergman projections and  $\bar{\partial}$ -Neumann problem. The method we use is a generalization of the well known construction of Forelli-Rudin. Such method has been used by many authors for the same kind of investigations ([Lig89, Zeyb]). To try to obtain weights which are not only powers of the defining function of  $\Omega$ , we consider a more general situation investigating the properties of a domain

$\tilde{\Omega}$  defined in  $\mathbb{C}^{n+m}$  by an equation of the form  $\rho(z) + h(w) < 0$  where  $\rho$  is a defining function of  $\Omega$  and  $h$  a positive function. The discussion of these properties is done in section 2.

*Remark.* The restriction  $r \in \mathbb{Q}_*^+$  in Theorem 1.1 is due to the method and we don't know if the theorem can be extended to the natural scale of powers which is  $]-1, +\infty[$ .

## 2. A HARTOGS DOMAIN $\tilde{\Omega}$ IN $\mathbb{C}^{n+m}$ BASED ON A DOMAIN $\Omega$ IN $\mathbb{C}^n$

For a given smoothly bounded pseudo-convex domain  $\Omega$  in  $\mathbb{C}^n$  with a defining function  $\rho$ , we consider a smooth non negative function  $h$  defined in  $\mathbb{C}^m$  such that  $h(w) = 0 \Leftrightarrow w = 0$  and  $\lim_{|w| \rightarrow +\infty} h(w) = +\infty$  and we denote by  $\tilde{\Omega}$  the smooth bounded domain

$$(2.1) \quad \tilde{\Omega} = \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^m, \text{ s. t. } \rho(z) + h(w) < 0\}.$$

Then, for particular functions  $h$ , there are, very simple relations between the standard Bergman kernel of  $\tilde{\Omega}$  and a weighted Bergman kernel of  $\Omega$ , and between the unweighted  $\bar{\partial}$ -Neumann problem on  $\tilde{\Omega}$  and a weighted  $\bar{\partial}_\varphi$ -Neumann problem on  $\Omega$  (see section 3). The aim of this part is to study the needed conditions on  $\rho$  and  $h$  to be able to obtain enough properties on  $\tilde{\Omega}$  so that we can get sharp estimates on the Bergman projection (or on the  $\bar{\partial}$ -Neumann problem of  $\tilde{\Omega}$ ) or sufficiently precise information on the Bergman kernel of  $\tilde{\Omega}$  on  $\{w = 0\}$ .

More precisely, in this section we discuss the following questions: suppose  $\nabla h(w) \neq 0$  if  $w \neq 0$ ; under which conditions on  $\rho$  and  $h$  the domain  $\tilde{\Omega}$  is:

- pseudo-convex;
- pseudo-convex of finite type if  $\Omega$  is of finite type.
- completely geometrically separated, in the sense of [CD08] if  $\Omega$  is so.

To get these properties, quite strong conditions have to be imposed to  $\rho$  and  $h$ , and, to simplify the reading of the paper, we state now these different conditions.

### 2.1. A special defining function for $\Omega$

As we will see in section 2.3, even with the function  $h(w) = |w|^2$ ,  $w \in \mathbb{C}$ ,  $\tilde{\Omega}$  is not always pseudo-convex (Remark 2.2). Thus, we have to use a special defining function  $\rho = \rho_s$  of  $\Omega$ :

A celebrated theorem of K. Diederich & J. E. Forneaess ([DF77a, Theorem 1]) proves that for any smooth bounded pseudo-convex domain  $\Omega$  there exists  $s_\Omega \in ]0, 1]$  such that, for  $s \in ]0, s_\Omega[$  there exists a smooth defining function  $\rho$  of  $\Omega$  such that the function  $-(-\rho)^s$  is strictly pluri-subharmonic in  $\Omega$ . Then

*In all this paper, for  $s \in ]0, s_\Omega[$ , we will denote by  $\rho_s$  a defining function such that*

$$(2.2) \quad -(-\rho_s)^s \text{ is strictly pluri-subharmonic in } \Omega.$$

Of course such a function  $\rho_s$  is not unique.

*Remark 2.1.*

- (1) The pluri-subharmonicity of  $-(-\rho_s)^s$  means

$$i\partial\bar{\partial}\rho_s \geq i\frac{1-s}{\rho_s}\partial\rho_s \wedge \bar{\partial}\rho_s.$$

Thus  $i\partial\bar{\partial}\rho_s \geq \frac{i}{\rho_s}\partial\rho_s \wedge \bar{\partial}\rho_s$ , and, as this means that  $-\log(-\rho_s)$  is pluri-subharmonic in  $\Omega$ , the  $\bar{\partial}_{-r\log(-\rho_s)}$ -Neumann problem is well defined for  $r \geq 0$ .

- (2) If  $U$  is an open set in  $\mathbb{C}^n$  and  $\rho$  is a  $\mathcal{C}^\infty$  function on  $U$  whose gradient does not vanish, then, if  $z^0 \in U$  with  $\rho(z^0) = 0$  and if  $i\partial\bar{\partial}\rho \geq i\frac{\mu}{\rho}\partial\rho \wedge \bar{\partial}\rho$ ,  $\mu \in \mathbb{R}$ , on the open set  $\{z \in U \text{ such that } \rho(z) < 0\}$  then the hypersurface

$$\Xi = \{z \in U \text{ such that } \rho(z) = 0\}$$

is pseudo-convex since the restriction of  $i\partial\bar{\partial}\rho$  to the complex tangent space of  $\Xi$  is non negative.

### 2.2. Hypothesis on the function $h$

Depending on the properties we want to have for  $\tilde{\Omega}$ , several conditions will be imposed to  $h$ . We define now five conditions that we will refer in the results stated in the following Sections.

Let  $\Omega$  be a bounded smooth pseudo-convex domain in  $\mathbb{C}^n$  and let  $h$  be a smooth real function on  $\mathbb{C}^m$ .

**Condition I:**

$\rho = \rho_s$  is a defining function of  $\Omega$  satisfying (2.2) as stated in section 2.1.  $h$  is non negative,  $h(w) = 0 \Leftrightarrow w = 0$  (and thus  $\nabla h(0) = 0$ ),  $\nabla h(w) \neq 0$  if  $w \neq 0$ ,  $\lim_{|w| \rightarrow +\infty} h(w) = +\infty$  and there exists  $s' \in [0, s[$  such that  $h^{s'}$  is pluri-subharmonic (i.e.  $i\partial\bar{\partial}h \geq i\frac{1-s'}{h}\partial h \wedge \bar{\partial}h$  which implies that  $h$  is strictly pluri-subharmonic at every point  $w$  such that  $\frac{\partial h}{\partial w_i}(w) \neq 0$  for all  $i$ ).

**Condition II:**

$h(w) = \sum_{i=1}^p h_i(w_i)$ ,  $w = (w_1, \dots, w_p)$ ,  $w_i \in \mathbb{C}^{m_i}$ , the functions  $h_i$  being non negative smooth pluri-subharmonic on  $\mathbb{C}^{m_i}$ , and satisfying:

- (a) for every  $i$ ,  $h_i(w_i) = 0 \Leftrightarrow w_i = 0$ ,  $\nabla h_i(w_i) \neq 0$  if  $w_i \neq 0$ ,  $\lim_{|w| \rightarrow +\infty} h_i(w) = +\infty$ ;
- (b)  $\log(h_i)$  is pluri-subharmonic (i.e.  $i\partial\bar{\partial}h_i \geq i\frac{1}{h_i}\partial h_i \wedge \bar{\partial}h_i$ ) and  $h_i$  is strictly pluri-subharmonic outside the origin;
- (c)  $h_i$  is of finite type  $2q_i = \text{typ}_0(h_i)$  at the origin (in the sense introduced at the beginning of section 2.4).

**Condition III:**

$h$  satisfies Condition II with  $m_i = 1$  for every  $i$  (thus the first part of the second condition of Condition II means  $\Delta h_i \geq \frac{1}{h_i}|h_i'|^2$ ), and, for each  $i$  there exists a function  $\alpha_i$ ,  $\mathcal{C}^\infty$  in a neighborhood of the origin in  $\mathbb{C}$ ,  $\lim_{w_i \rightarrow 0} \alpha_i(w_i) = 0$ , such that  $\frac{\partial h_i}{\partial w_i} = \alpha_i \frac{\partial^2 h_i}{\partial w_i \partial \bar{w}_i}$  in that neighborhood.

**Condition IV:**

$\Omega$  is of finite type,  $h$  satisfies Condition III, and, for each  $i$ ,  $h_i(w_i) = k_i(|w_i|)$ ,  $k_i(t) \asymp t^{2q_i}$  (where  $f \asymp g$  means that there exist two constants  $c > 0$  and  $C > 0$  such that  $cf \leq g \leq Cf$ ) and  $2q_i$  strictly larger than the type of  $\Omega$ .

**Condition V:**

$\Omega$  is of finite type,  $h$  satisfies Condition II with  $m_i = 1$  for every  $i$ , and, for each  $i$ ,  $\left| \frac{\partial^2 h_i}{\partial w_i^2} \right| \leq \Delta h_i$ ,  $h_i(w_i) = k_i(|w_i|)$  and  $k_i(t) \asymp t^{2q_i}$  and  $q_i$  strictly larger than the type of  $\Omega$ .

Condition I is used in section 2.3 to get the pseudo-convexity of  $\tilde{\Omega}$ . Condition II is used in section 2.4 to ensure that  $\tilde{\Omega}$  is of finite type. In section 2.5 we use Condition III to obtain that  $\tilde{\Omega}$  have a Levi form locally diagonalizable (and thus is ‘‘completely geometrically separated’’ ([CD08])) when  $\Omega$  is a finite type domain in  $\mathbb{C}^2$ . Finally, Conditions IV and V are used in Sections section 4.1 and section 4.2 to get pointwise estimates of a weighted Bergman kernel in these two cases.

**Example 2.1.** Let  $m_i \in \mathbb{N}_*$ ,  $1 \leq i \leq p$ ,  $m = \sum_i m_i$ ,  $q_i \in \mathbb{N}_*$ ,  $1 \leq i \leq p$ . Then the function

$$h : w = (w_1, \dots, w_p) \in \prod_i \mathbb{C}^{m_i} = \mathbb{C}^m \mapsto \sum_i |w_i|^{2q_i}$$

satisfies Conditions I (whatever is  $s$ ) and II, and all other conditions if  $m_i = 1$  for every  $i$  and the  $q_i$  large enough.

*Proof.* Let us denote  $h_i(w_i) = |w_i|^{2q_i}$ ,  $1 \leq i \leq p$ . Then  $\bar{\partial}h = \sum_i \bar{\partial}h_i$ ,  $\partial\bar{\partial}h = \sum_i \partial\bar{\partial}h_i$  and, a simple calculus and Cauchy-Schwarz inequality gives, for  $1 \leq i \leq p$ ,

$$\begin{aligned} i\partial\bar{\partial}h_i &= iq_i^2 |w_i|^{2q_i-4} \left( \sum_j \bar{w}_i^j dw_i^j \right) \wedge \left( \sum_j w_i^j d\bar{w}_i^j \right) + \\ &\quad + iq_i |w_i|^{2q_i-2} \left( \sum_j dw_i^j \wedge d\bar{w}_i^j \right) - iq_i |w_i|^{2q_i-4} \left( \sum_j \bar{w}_i^j dw_i^j \right) \wedge \left( \sum_j w_i^j d\bar{w}_i^j \right) \\ &\geq i \frac{1}{h_i} \partial h_i \wedge \bar{\partial} h_i, \end{aligned}$$

Then, by Cauchy-Schwarz inequality,

$$ih \langle \partial\bar{\partial}h; t, \bar{t} \rangle \geq \left( \sum_i h_i \right) \left( \sum \frac{i \langle \partial h_i \wedge \bar{\partial} h_i; t_i, \bar{t}_i \rangle}{h_i} \right) \geq \left( \sum \langle \partial h_i \wedge \bar{\partial} h_i; t_i, \bar{t}_i \rangle^{1/2} \right)^2 \geq i \langle \partial h \wedge \bar{\partial} h; t, \bar{t} \rangle$$

and Condition I is satisfied for any  $s' \geq 0$ . □

**2.3. Pseudo-convexity of  $\tilde{\Omega}$** 

In general, even for very simple function  $h$ , the domain  $\tilde{\Omega}$  is not pseudo-convex. For example, if  $\Omega$  is the unit ball it is very easy to write a defining function  $\rho$  of  $\Omega$  such that the domain  $\{(z, w) \in \mathbb{C}^n \times \mathbb{C}, \text{ s. t. } \rho(z) + |w|^2 < 0\}$  is not pseudo-convex (see Remark 2.2).

If  $\Omega$  admits a smooth defining function which is pluri-subharmonic in  $\bar{\Omega}$ , it suffices to take  $h$  pluri-subharmonic. But this is not the general case (c.f. [DF77b]), so, we have to choose a convenient defining function for  $\Omega$ :

**Proposition 2.1.** *Let  $Z \in \mathbb{C}^n$  and let  $U$  (resp.  $V$ ) be an open neighborhood of  $Z$  (resp. of the origin in  $\mathbb{C}^m$ ). Let  $\rho : U \rightarrow \mathbb{R}$  (resp.  $h : V \rightarrow \mathbb{R}_+$ ) be a smooth function such that  $\rho(Z) = 0$  and  $\nabla \rho$  does not vanishes in  $U$  (resp.  $\nabla h(w) \neq 0$  if and only if  $w \neq 0$  and  $h(w) \neq 0$  if  $w \neq 0$ ). Assume that there exists  $s \in ]0, 1[$  and  $s' \in [0, s[$  such that  $-(\rho)^s$  is strictly pluri-subharmonic in the open set  $G = \{z \in U \text{ s. t. } \rho(z) < 0\}$  and  $h^{s'}$  is pluri-subharmonic in  $V$ . Then:*

- (1) *The hypersurface  $\partial\tilde{G} = \{(z, w) \in U \times V \text{ s. t. } r(z, w) = \rho(z) + h(w) = 0\}$  is pseudo-convex (in the sense that the restriction of  $i\partial\bar{\partial}r$  to the complex tangent space of  $\partial\tilde{G}$  is non negative).*
- (2) *Moreover,  $\partial\tilde{G}$  is strictly pseudo-convex at a point  $(z^1, w^1)$  if  $h$  is strictly pluri-subharmonic at  $w^1 \neq 0$ .*

*Proof.* Note that, since the gradient of  $r$  does not vanishes on  $U \times V$ ,  $\partial\tilde{G}$  is an hypersurface if it is not empty. Moreover,  $s'$  being  $\leq 1$ ,  $h$  is pluri-subharmonic on  $V$ . The hypothesis made on  $\rho$  is

$$(2.3) \quad i\partial\bar{\partial}\rho \geq i\frac{1-s}{\rho}\partial\rho \wedge \bar{\partial}\rho + \varepsilon i\partial\bar{\partial}|z|^2,$$

where  $\varepsilon$  is a non negative function, strictly positive in  $G$ .

Let  $(z^1, w^1)$  be a point of the boundary of  $\tilde{G}$  and  $t = (t_z, t_w) \in \mathbb{C}^n \times \mathbb{C}^m$  be a vector of the complex tangent space of  $\partial\tilde{G}$  at  $(z^1, w^1)$ .

If  $w^1 = 0$ , as the hypothesis on  $h$  imply  $\nabla h(0) = 0$ ,  $t_z$  is tangent to  $\rho$  at  $z^0$ , and the pseudo-convexity of  $\partial\tilde{G}$  (Remark 2.1) and the pluri-subharmonicity of  $h$  imply  $\langle i\partial\bar{\partial}r; t, \bar{t} \rangle \geq 0$  showing that  $\partial\tilde{G}$  is pseudo-convex at  $(z^1, 0)$ .

Suppose now  $w^1 \neq 0$ . Note that,  $t$  being tangent to  $r$ , we have  $\sum_i \frac{\partial\rho}{\partial z_i}(z^1) t_z^i = -\sum_j \frac{\partial h}{\partial w_j}(w^1) t_w^j$ , and  $\rho(z^1) = -h(w^1)$ . Then (2.3) gives

$$\langle i\partial\bar{\partial}\rho(z^1); t_z, \bar{t}_z \rangle \geq \frac{1-s}{-h(w^1)} \left| \sum_j \frac{\partial h}{\partial w_j}(w^1) t_w^j \right|^2 + \varepsilon |t_z|^2.$$

Thus

$$\langle i\partial\bar{\partial}r(z^1, w^1); t, \bar{t} \rangle \geq \frac{1-s}{-h(w^1)} \left| \sum_j \frac{\partial h}{\partial w_j}(w^1) t_w^j \right|^2 + \langle i\partial\bar{\partial}h(w^1); t_w, \bar{t}_w \rangle + \varepsilon |t_z|^2.$$

The conclusion comes then from the fact that the hypothesis made on  $h$  is

$$i\partial\bar{\partial}h \geq \max \left\{ i\frac{1-s'}{h}\partial h \wedge \bar{\partial}h, \varepsilon_1 i\partial\bar{\partial}|w|^2 \right\},$$

with  $0 \leq s' < s$  and  $\varepsilon_1$  a non negative function, which is positive at  $w^1$  if  $h$  is strictly pluri-subharmonic at that point:

$s' \leq s$  gives immediately (1), and, if  $h$  is strictly pluri-subharmonic at  $w^1$ , then  $\varepsilon(z^1) > 0$ ,  $\varepsilon_1(w^1) > 0$  and

$$\langle i\partial\bar{\partial}r(z^1, w^1); t, \bar{t} \rangle \geq \varepsilon_2 |t|^2,$$

for  $\varepsilon_2 > 0$  small. □

**Proposition 2.2.** *Let  $\Omega$  be a bounded pseudo-convex domain, with smooth boundary. Let  $h$  be a smooth non negative function defined in  $\mathbb{C}^m$ . Let  $\rho = \rho_s$  be a smooth defining function of  $\Omega$  satisfying (2.2) as stated in section 2.1.*

(1) *Then, if  $h$  satisfies Condition 1 of section 2.2,  $\tilde{\Omega}$  is pseudo-convex.*

(2) *Moreover, for  $(z^0, w^0) \in \partial\tilde{\Omega}$ ,  $w^0 \neq 0$  if, in addition,  $h$  is strictly pluri-subharmonic at  $w^0$ ,  $\tilde{\Omega}$  is strictly pseudo-convex at  $(z^0, w^0)$ .*

*Proof.* Let  $(z^0, w^0) \in \partial\tilde{\Omega}$ . If  $\partial\rho(z^0) \neq 0$ ,  $\tilde{\Omega}$  is pseudo-convex at  $(z^0, w^0)$  by Proposition 2.1. If  $\partial\rho(z^0) = 0$ , then  $z^0 \in \Omega$  and (2.3) shows that  $\rho$  is strictly pluri-subharmonic in a neighborhood of  $z^0$ . Thus  $r$  is pluri-subharmonic in a neighborhood of  $(z^0, w^0)$  and strictly pluri-subharmonic if  $h$  is strictly pluri-subharmonic at  $w^0$ . □

*Remark 2.2.* When  $m = 1$  and  $h(w) = |w|^2$ , the hypersurface  $\{\rho(z) + h(w) = 0\}$  is pseudo-convex at a point  $(z^1, w^1)$ ,  $w^1 \neq 0$ , if and only if

$$i\partial\bar{\partial}\rho(z^1) \geq \frac{i}{\rho(z^1)} (\partial\rho \wedge \bar{\partial}\rho)(z^1).$$

For example, if  $\rho$  is the signed distance to the boundary of  $\Omega$ , by Oka's theorem,  $\{\rho(z) + |w|^2 < 0\}$  is pseudo-convex.

**Example 2.2.** With the function  $h$  given in Example 2.1, (1) of Proposition 2.2 applies for any pseudo-convex domain  $\Omega$ . Note also that,  $\tilde{h}_1(w) = h(w) + |w|^{2q}$ ,  $q \in \mathbb{N}_*$  satisfies the condition stated in (2) of the Proposition.

*Proof.* As noted in the proof of Example 2.1,  $i\partial\bar{\partial}(|w|^{2q}) \geq \frac{i}{|w|^{2q}} \partial(|w|^{2q}) \wedge \bar{\partial}(|w|^{2q})$ . □

*Remark 2.3.*  $\Omega$  being pseudo-convex in  $\mathbb{C}^n$ , E. Ligocka considered, in [Lig89], the domains

$$\tilde{\Omega}_{N,k} = \left\{ (z, w) \in \mathbb{C}^n \times \mathbb{C}^m \text{ s. t. } \rho(z) + |w|^{2Nk} < 0 \right\},$$

where  $k$  is a sufficiently large integer such that the defining function  $\rho$  of  $\Omega$  satisfies that  $-(-\rho)^{1/k}$  is strictly plurisubharmonic in  $\Omega$  and  $N$  a positive integer. She showed that if  $\Omega$  is "weakly regular", so is  $\tilde{\Omega}_{N,k}$ .

## 2.4. Type finiteness of $\tilde{\Omega}$

We now investigate the question concerning the type of the domain  $\tilde{\Omega}$  defined by equation (2.1) with a defining function  $\rho$  of  $\Omega$  which does not necessarily satisfy equation (2.2). Recall first that, if  $g$  is a function defined in a neighborhood of the origin, the order of cancellation of  $g$  at 0, denoted here by  $\text{ord}_0(g)$ , is the minimal length of the derivatives of  $g$  at 0 which are not equal to 0. Then,  $h$  being a smooth function defined on  $\mathbb{C}^m$ , we call the *type* of  $h$  at the origin the supremum of  $\frac{\text{ord}_0(h \circ \varphi)}{\text{ord}_0(\varphi)}$ , taken over all non zero holomorphic function  $\varphi$  from the unit disc of the complex plane into  $\mathbb{C}^m$  such that  $\varphi(0) = 0$ . If this supremum is finite, we will say that  $h$  is of *finite type* at the origin and we will denote this supremum by  $\text{typ}_0(h)$ .

**Proposition 2.3.** *Let  $z_0 \in \mathbb{C}^n$  and  $U$  (resp.  $V$ ) be an open neighborhood of  $z_0$  (resp. of the origin in  $\mathbb{C}^m$ ). Let  $\rho : U \rightarrow \mathbb{R}$  (resp.  $h : V \rightarrow \mathbb{R}_+$ ) be a smooth function such that  $\rho(z_0) = 0$  and  $\nabla \rho$  does not vanishes in  $U$  (resp.  $\nabla h(w) \neq 0$  if  $w \neq 0$  and  $h(w) \neq 0$  if and only if  $w \neq 0$ ). Assume that the hypersurface  $\partial G = \{z \in U \text{ s. t. } \rho(z) = 0\}$  is pseudo-convex (in the sense that the restriction of  $i\partial\bar{\partial}\rho$  to the complex tangent space to  $\partial G$  is non negative) and of finite type  $\tau$  at the point  $z_0$  and that  $h$  is of finite type  $\text{typ}_0(h)$  at the origin.*

*Then the boundary of  $\tilde{G} = \{(z, w) \in U \times V \text{ s. t. } r(z, w) = \rho(z) + h(w) < 0\}$  is of finite type  $\max(\tau, \text{typ}_0(h))$  at the point  $(z_0, 0)$ .*

*Proof.* To simplify the notations, we can assume  $z_0 = 0$ . Let  $\Phi = (\varphi, \psi)$  be a non constant holomorphic function from the unit disc of the complex plane into  $U \times V$  such that  $\Phi(0) = 0$ . We want to estimate  $\frac{\text{ord}_0(r \circ \Phi)}{\text{ord}_0(\Phi)}$ , and thus we can assume that the first derivative of  $r \circ \Phi$  vanishes at the origin. As the gradient of  $h$  vanishes at the origin, this implies that the first derivative of  $\rho \circ \varphi$  vanishes at the origin which means that the gradient of  $\varphi$  at 0 is tangent to  $\rho$  at 0. Clearly, we can assume that the order of cancellation of  $h \circ \psi$  at the origin is not infinite, and then, by Lemma 8.1 of [CD08] and the hypothesis made on  $h$ , there exists an integer  $k \geq 1$  such that all the derivatives of order  $< 2k$  of  $h \circ \psi$  at the origin vanish and  $\Delta^k(h \circ \psi)(0) > 0$ .

As  $\text{ord}_0(\Phi)$  is the minimum of  $\text{ord}_0(\varphi)$  and  $\text{ord}_0(\psi)$ , to prove the Proposition it suffices to prove that

$$\text{ord}_0(r \circ \Phi) \leq \min\{\text{ord}_0(\rho \circ \varphi), \text{ord}_0(h \circ \psi)\} = \min\{\text{ord}_0(\rho \circ \varphi), 2k\}$$

(note that this not a ‘‘trivial’’ consequence of the fact that  $\rho$  and  $h$  are decoupled).

Note first that, if  $\text{ord}_0(\rho \circ \varphi) < 2k$  or  $\text{ord}_0(\rho \circ \varphi) > 2k$ , the inequality is obvious. Thus we can assume  $\text{ord}_0(\rho \circ \varphi) = 2k$ , and we have to prove that  $\text{ord}_0(r \circ \Phi) = 2k$ .

Suppose it is not the case: it then follows that all the derivatives of  $r \circ \Phi$  of order  $\leq 2k$  vanish at 0. Consider  $\Delta^k(\rho \circ \varphi)(0)$ . As  $\varphi$  is holomorphic, we have

$$(2.4) \quad \Delta^k(\rho \circ \varphi)(0) = \Delta^{k-1} \left( \left\langle \partial\bar{\partial}\rho(\varphi); \varphi', \overline{\varphi'} \right\rangle \right) (0).$$

As all the derivatives of  $\rho \circ \varphi$  of order less or equal than  $2k - 1$  vanish at the origin, we have  $\rho \circ \varphi(\zeta) = O(\zeta^{2k})$ . For  $\zeta$  small, let  $\xi = \xi(\zeta)$  be the projection of  $\varphi(\zeta)$  on  $\{z \in U \text{ s. t. } \rho(z) = 0\}$  so that  $\varphi(\zeta) - \xi = O(\zeta^{2k})$ . By the hypothesis  $\text{ord}_0(\rho \circ \varphi) = 2k$ , all the derivatives of  $\langle \partial\bar{\partial}\rho(\varphi); \varphi' \rangle$  of order less or equal than  $2k - 2$  vanish at the origin, which implies that there exists a vector  $T = T(\zeta)$  tangent to  $\rho$  at the point  $\xi$  such that  $\varphi'(\zeta) = T + O(\zeta^{2k-1})$ . Then, the hypersurface  $\partial G$  being pseudo-convex (by hypothesis), we have

$$\left\langle i\partial\bar{\partial}\rho(\varphi(\zeta)); \varphi'(\zeta), \overline{\varphi'(\zeta)} \right\rangle = \left\langle i\partial\bar{\partial}\rho(\xi); T, \overline{T} \right\rangle + O(\zeta^{2k-1}) \geq -C|\zeta|^{2k-1}.$$

Applying Lemma 8.1 of [CD08] to the positive function

$$\left\langle i\partial\bar{\partial}\rho(\varphi(\zeta)); \varphi'(\zeta), \overline{\varphi'(\zeta)} \right\rangle + C|\zeta|^{2k-1},$$

by (2.4), we get  $\Delta^k(\rho \circ \varphi)(0) \geq 0$  which is impossible since  $0 = \Delta^k(r \circ \Phi)(0) = \Delta^k(\rho \circ \varphi)(0) + \Delta^k(h \circ \psi)(0) > 0$ .  $\square$

Recall that in Proposition 2.3 we do not assume that  $\rho$  necessarily satisfies (2.2).

Applying this Proposition to a general pseudo-convex domain, we get:

**Corollary 1.** *Assume that  $\rho = \rho_s$  where  $\rho_s$  satisfy (2.2) as stated in section 2.1.*

- (1) *Assume that  $h$  satisfies Condition I of section 2.2. Let  $z^0$  be a boundary point of  $\Omega$ . If  $\partial\Omega$  is of finite type  $\tau$  at  $z^0$  and if  $h$  is of finite type  $\text{typ}_0(h)$  at the origin, then  $\partial\tilde{\Omega}$  is of finite type  $\max(\tau, \text{typ}_0(h))$  at the point  $(z^0, 0)$ . Moreover, if  $h$  is strictly pluri-subharmonic in  $\mathbb{C}^n \setminus \{0\}$  then  $\partial\tilde{\Omega}$  is strictly pseudo-convex at every boundary point  $(z^0, w^0)$  such that  $w^0 \neq 0$ .*
- (2) *Assume that  $h$  satisfy Condition II of section 2.2. Then  $\tilde{\Omega}$  is pseudo-convex and, at every point  $(z^0, w^0) \in \partial\tilde{\Omega}$ ,  $w^0 \neq 0$ ,  $\partial\tilde{\Omega}$  is of finite type  $\max_i \text{ s. t. } w_i^0 = 0 \{ \text{typ}_0(h_i) \}$  if there exists some  $i$  such that  $w_i^0 = 0$  and strictly pseudo-convex if  $w_i^0 \neq 0$  for all  $i$ .*

*Proof.* The first part of (1) is a special case of Proposition 2.3, and the second part is stated in Proposition 2.2.

Let us now prove (2). The pseudo-convexity of  $\tilde{\Omega}$  follows the results of section 2.3. If  $w_i^0 \neq 0$  for all  $i$ , as before,  $\partial\tilde{\Omega}$  is strictly pseudo-convex at  $(z^0, w^0)$ . Then, assume that there exists some  $i$  such that  $w_i^0 = 0$ . Without loss of generality, we can suppose that  $w_{k+1}^0 = \dots = w_p^0 = 0$ ,  $k < p$ , and  $w_l^0 \neq 0$  for  $1 \leq l \leq k$ . Let us denote  $w = (w', w'')$ , with  $w' = (w_1, \dots, w_k)$ ,  $w'' = (w_{k+1}, \dots, w_p)$ ,

and  $\rho_1(z, w') = \rho(z) + \sum_{i=1}^k h_i(w_i)$ . In a neighborhood of  $(z^0, w'^0)$ ,  $\nabla \rho_1$  does not vanish,  $\{\rho_1 < 0\}$  is strictly pseudo-convex, and we can apply Proposition 2.3 to the domain  $\rho_1(z, w') + \sum_{i=k+1}^p h_i(w_i) < 0$  and the function  $h_1(w')$  at the point  $(z^0, w'^0, 0)$ .  $\square$

When  $\Omega$  admits a pluri-subharmonic defining function which is of finite type everywhere, the Proposition gives:

**Corollary 2.** *Assume  $\Omega$  admits a defining function  $\rho$  pluri-subharmonic in a neighborhood of  $\bar{\Omega}$  and of finite type in  $\bar{\Omega}$ . Assume that  $h$  satisfies Condition II of section 2.2. Then the domain  $\tilde{\Omega}$ , defined with  $\rho$ , is pseudo-convex of finite type. More precisely, at every point  $(z^1, w^1) \in \partial\tilde{\Omega}$  the type of  $\partial\tilde{\Omega}$  is bounded by  $\max\{\text{typ}_{z^1}(\partial\Omega), \text{typ}_0(h_i), 1 \leq i \leq p\}$  if  $w^1 = 0$  and by*

$$\max\left\{2\text{typ}_{z^1}(\rho), \max_{i \text{ such that } w_i \neq 0} \{\text{typ}_0(h_i)\}\right\}$$

otherwise.

*Proof.* Let us first consider the case  $p = 1$ . By Proposition 2.3,  $\tilde{\Omega}$  is of finite type  $\max\{\text{typ}_{z^1}(\partial\Omega), \text{typ}_0(h)\}$  at every point  $(z^1, 0) \in \partial\tilde{\Omega}$ , and we have to study the finiteness at points  $(z^1, w^1) \in \partial\tilde{\Omega}$  such that  $w^1 \neq 0$ . Let  $\Phi = (\varphi, \psi)$  be a non constant holomorphic function from the unit disc of the complex plane into a neighborhood of  $(z^1, w^1)$  such that  $\Phi(0) = (z^1, w^1)$ . We have to estimate  $\tau_\Phi = \frac{\text{ord}_0(r \circ \Phi)}{\text{ord}_0(\Phi - \Phi(0))}$ , and we can, of course, assume that  $\text{ord}_0(r \circ \Phi) \geq 2$ .

Let  $k$  be a positive integer and let us assume that all the derivatives of order  $\leq 2k$  of  $r \circ \Phi$  vanish at the origin. Then

$$\Delta(r \circ \Phi)(0) = \langle \partial\bar{\partial}\rho; \varphi', \bar{\varphi}' \rangle(0) + \langle \partial\bar{\partial}h; \psi', \bar{\psi}' \rangle(0) = 0,$$

and the hypothesis on  $\rho$  and  $h$  (pluri-subharmonicity of  $\rho$  and strict pluri-subharmonicity of  $h$ ) imply  $\langle \partial\bar{\partial}\rho; \varphi', \bar{\varphi}' \rangle(0) = \langle \partial\bar{\partial}h; \psi', \bar{\psi}' \rangle(0) = 0$  and the last equality implies  $\psi'(0) = 0$ . Moreover Lemma 8.1 of [CD08] implies that, for  $1 \leq j \leq k-1$ ,  $\Delta^j(\langle \partial\bar{\partial}h; \psi', \bar{\psi}' \rangle)(0) = 0$ , and, by induction, a simple calculus shows that this implies  $\psi^{(j+1)}(0) = 0$ ,  $1 \leq j \leq k-1$ . Then all the derivatives of order  $\leq k$  of  $h \circ \psi$  vanish at the origin, and, thus, the same is true for the derivatives of  $\rho \circ \varphi$ .

This implies first that the order of cancellation of  $r \circ \Phi$  cannot be infinite at 0. Assume it is  $l$ , and let  $k$  be a positive integer such that  $l = 2k + 1$  or  $l = 2k + 2$ . In both cases, the order of cancellation of  $\rho \circ \varphi - \rho(z^1)$ ,  $h \circ \psi - h(w^1)$  and  $\psi - \psi(0)$  are  $\geq k + 1$  and we have  $\frac{\text{ord}_0(r \circ \Phi)}{\text{ord}_0(\Phi - \Phi(0))} \leq \frac{2\text{ord}_0(\rho \circ \varphi - \rho(z^1))}{\text{ord}_0(\Phi - \Phi(0))}$  which implies  $\tau_\Phi \leq 2$  if  $\text{ord}_0(\varphi - \varphi(0)) \geq k + 1$  and  $\tau_\Phi \leq 2\text{typ}_{z^1}(\rho)$  if not.

The case  $p \geq 2$  follows easily. If, for all  $i$ ,  $1 \leq i \leq p$ ,  $w_i^1 \neq 0$  then  $h$  is strictly pluri-subharmonic at  $w^1$  and the previous proof applies. Otherwise, to simplify notations, we can assume that  $w_i^1 \neq 0$  for  $1 \leq i \leq r < p$  and  $w_i^1 = 0$  for  $r + 1 \leq i \leq p$ . Denoting  $u = (w_1, \dots, w_r)$ ,  $v = (w_{r+1}, \dots, w_p)$ ,  $\rho_1(z, u) = \rho(z) + \sum_{i=1}^r h_i(w_i)$  and  $h_1(v) = \sum_{i=r+1}^p h_i(w_i)$ , the previous case shows that  $\rho_1$  is pluri-subharmonic and the type of  $\rho$  at  $(z^1, u^1)$  is bounded by  $2\text{typ}_{z^1}(\rho)$ . The conclusion is obtained applying Proposition 2.3.  $\square$

## 2.5. Geometric separation

If the domain  $\Omega$  is completely geometrically separated at a boundary point  $z_0$  (see [CD08] for definition), we do not know, in general, if  $\tilde{\Omega}$  has the same property at the point  $(z_0, 0)$ . We can only prove the weaker following result (for which we will not give a proof because we don't have any application):

**Proposition 2.4.** *Assume that  $\Omega$  is of finite type at  $z_0 \in \partial\Omega$  and that  $h(w) = \sum |w_i|^{2q_i}$ ,  $w_i \in \mathbb{C}$ . Then for all Diederich-Fornaess defining function  $\rho$  of  $\Omega$  of the form  $\rho = \sigma e^{-L|z|^2}$  (see [DF77a]) with  $L$  large enough (depending only on  $\Omega$ ), we have:*

*if there exist a neighborhood  $V$  of  $z_0$ ,  $K > 0$  and a finite dimensional vector space  $E_0$  of complex tangent vector fields to  $\rho$  in  $V$  such that, at any point of  $V \cap \Omega$  and for any  $\delta > 0$  there exists a  $(K, \delta)$ -extremal basis for  $\rho$  whose elements belong to  $E_0$ , then  $\tilde{\Omega}$  is geometrically separated at  $(z_0, 0)$ .*

Note that the hypothesis in this Proposition is stronger than the simple fact that  $\partial\Omega$  is geometrically separated at  $z_0$ : the existence of extremal basis is assumed not only on the points of  $\partial\Omega \cap V$  but on all  $\Omega \cap V$  (condition which depends not only on  $\Omega$  but also on the choice of  $\rho$ ). Unfortunately, if we add the hypothesis that all the level set of  $\rho$  are ‘‘completely geometrically separated’’ in  $\Omega \cap V$  we were not able to prove, in general, that  $\tilde{\Omega}$  has the same property at  $(z_0, 0)$ . The only general result we have is when  $\Omega$  is in  $\mathbb{C}^2$  (see Remark 2.4):

**Theorem 2.1.** *Assume  $\Omega$  is pseudo-convex of finite type in  $\mathbb{C}^2$ . Assume that  $\rho = \rho_s$  with  $\rho_s$  satisfying (2.2) as stated in section 2.1 and that  $h$  satisfies Condition III of section 2.2. Then the domain*

$$\tilde{\Omega} = \left\{ (z, w) = (z, w_1, \dots, w_m) \in \mathbb{C}^2 \times \mathbb{C}^m \text{ s. t. } r(z, w) = \rho(z) + \sum_{i=1}^m h_i(w_i) < 0 \right\}$$

*is pseudo-convex of finite type and has a Levi form which is locally diagonalizable at every point of its boundary. In particular  $\tilde{\Omega}$  is completely geometrically separated (c.f. [CD08]).*

*Proof.* Let  $(z^0, w^0)$  be a boundary point of  $\tilde{\Omega}$ . If  $w_i^0 \neq 0$  for all  $i$ , by (2) of the Corollary of Proposition 2.3,  $\partial\tilde{\Omega}$  is strictly pseudo-convex at  $(z^0, w^0)$ . Thus we have only two cases to consider:



- (1)  $w^0 = 0$ ;
- (2) there exist  $i$  and  $j$  such that  $w_i^0 = 0$  and  $w_j^0 \neq 0$ .

Let us consider the first case. Denote by  $L$  (resp.  $N$ ) the vector field complex tangent (resp. normal) to  $\rho$  in a neighborhood of  $z^0$ . We assume that  $N$  is chosen so that  $N\rho \equiv 1$  in that neighborhood. Without changing the notation, we will consider these vector fields defined in a neighborhood of  $(z^0, 0)$  so that  $L$  is tangent to  $r$  and  $Nr \equiv 1$  in this neighborhood. Let us define  $m$  vector fields,  $Z_i$ , complex tangent to  $r$ , in a neighborhood of  $(z^0, 0)$  by

$$Z_i = \frac{\partial}{\partial w_i} - \frac{\partial h_i}{\partial w_i} N,$$

and then  $m$  new vector fields,  $W_i$ , also complex tangent to  $r$ , by

$$\begin{aligned} W_1 &= Z_1, \\ W_{k+1} &= Z_{k+1} - \sum_{j=1}^k a_{k+1}^j W_j \text{ for } k \geq 2. \end{aligned}$$

We now show, by induction over  $k$ , that it is possible to choose the coefficients  $a_k^j$  so that the coefficient of the Levi form of  $r$ ,  $[W_k, \overline{W}_k](\partial r)$ , vanishes identically on the neighborhood of  $(z^0, 0)$ . To simplify notations, in this proof, the character  $*$  will denote a  $\mathcal{C}^\infty$  function in a neighborhood of the origin.

Suppose that the vector fields  $W_i$ ,  $1 \leq i \leq k$ , has been constructed with coefficients  $a_i^j$ ,  $2 \leq i \leq k$ ,  $1 \leq j \leq i-1$  satisfying the two following properties:

- (1)  $a_i^j = * \frac{\partial h_i}{\partial w_i}$ ,
- (2)  $[W_i, \overline{W}_i](\partial r) = \alpha_i \frac{\partial^2 h_i}{\partial w_i \partial \overline{w}_i}$ , where  $\alpha_i$  is a  $\mathcal{C}^\infty$  real function in a neighborhood of the origin of modulus greater than  $1/2$ ,

and let us prove that  $W_{k+1}$  can be constructed, the coefficients  $a_{k+1}^j$ ,  $1 \leq j \leq k$  satisfying the above conditions. Note that the hypothesis made on  $h_i$  imply first that (2) follows (1) because (1) implies

$$[W_i, \overline{W}_i](\partial r) = \frac{\partial^2 h_i}{\partial w_i \partial \overline{w}_i} + \left| \frac{\partial h_i}{\partial w_i} \right|^2 [N, \overline{N}](\partial \rho) + * \frac{\partial h_i}{\partial w_i} + * \frac{\partial h_i}{\partial \overline{w}_i}.$$

Note also that  $W_1$  satisfy trivially (2).

Thus we construct  $W_{k+1}$  with coefficients  $a_{k+1}^j$  satisfying (1).

For  $j \leq k$ , by induction, we have

$$[W_{k+1}, \overline{W}_j](\partial r) = -\frac{\partial h_j}{\partial \overline{w}_j} [N, \overline{W}_j](\partial r) - a_{k+1}^j [W_j, \overline{W}_j](\partial r)$$

with

$$\begin{aligned} [N, \overline{W}_j](\partial r) &= -\frac{\partial h_j}{\partial \overline{w}_j} [N, \overline{N}](\partial \rho) - \sum_{1 \leq l < j} \overline{a}_l^j [N, \overline{W}_l](\partial r) \\ &= * \frac{\partial h_j}{\partial \overline{w}_j}, \end{aligned}$$

if  $2 \leq j \leq k$ , and

$$[N, \overline{W}_1](\partial r) = * \frac{\partial h_1}{\partial \overline{w}_1}.$$

This shows, by Condition III of  $h$ , that the  $a_{k+1}^j$  can be defined satisfying (1) and such that  $[W_{k+1}, \overline{W}_j](\partial r) \equiv 0$ .

To finish the proof of the first case, we modify the vector field  $L$  replacing it by  $L_1 = L - \sum_{k=1}^m b_k W_k$  choosing the  $b_k$  so that the basis  $(L_1, W_1, \dots, W_m)$  diagonalizes the Levi form of  $r$  in a neighborhood of  $(z^0, 0)$  which means, now,  $[L_1, \overline{W}_i](\partial r) \equiv 0$  in that neighborhood:

$$\begin{aligned} [L_1, \overline{W}_i](\partial r) &= [L, \overline{W}_i](\partial r) - b_i [W_i, \overline{W}_i](\partial r) \\ &= \begin{cases} -\frac{\partial h_i}{\partial \overline{w}_i} [L, \overline{N}](\partial r) - \sum_{1 \leq l < i} \overline{a}_l^i [L, \overline{W}_l](\partial r) - b_i [W_i, \overline{W}_i](\partial r) & \text{if } i \geq 2 \\ -\frac{\partial h_1}{\partial \overline{w}_1} [L, \overline{N}](\partial r) - b_i [W_i, \overline{W}_i](\partial r) & \text{if } i = 1 \end{cases} \\ &= * \frac{\partial h_i}{\partial \overline{w}_i} - b_i [W_i, \overline{W}_i](\partial r), \end{aligned}$$

and, by (2),  $b_i$  can be chosen  $\mathcal{C}^\infty$  in a neighborhood of  $(z^0, 0)$ .

Let us now consider the second case (2). To simplify the notations, we assume that  $w_i^0 \neq 0$  for  $1 \leq i \leq m_0 < m$  and  $w_i^0 = 0$  for  $m_0 + 1 \leq i \leq m$ . We denote  $w = (w', w'')$ , with  $w' = (w_1, \dots, w_{m_0})$ ,  $w'' = (w_{m_0+1}, \dots, w_m)$ ,  $\rho^1(z, w') = \rho(z) + \sum_{i=1}^{m_0} h_i(w_i)$  and  $h^1(w'') = \sum_{i=m_0+1}^m h_i(w_i)$ .

By Proposition 2.1, in a neighborhood of  $(z^0, w^0)$ , the hypersurface  $\{\rho^1 = 0\}$  is strictly pseudo-convex. Then, reducing eventually the neighborhood, there exists a basis of vector fields  $(L_1, \dots, L_{m_0-1})$  complex tangent to  $\rho^1$  which diagonalizes the

Levi form of  $\rho^1$  in that neighborhood. Let us denote by  $N$  the complex normal vector field to  $\rho^1$  in that neighborhood such that  $N\rho^1 \equiv 1$  (note that, reducing the neighborhood if necessary, we can assume that the gradient of  $\rho^1$  does not vanishes in the neighborhood). We now consider the following  $m - m_0$  vector fields (which are complex tangent to  $\rho^1$  in the neighborhood)

$$\begin{aligned} W_1 &= \frac{\partial}{\partial w_{m_0+1}} - \frac{\partial h_{m_0+1}}{\partial w_{m_0+1}} N - \sum_{i=1}^{m_0-1} a_1^i L_i, \\ W_j &= \frac{\partial}{\partial w_{m_0+j}} - \frac{\partial h_{m_0+j}}{\partial w_{m_0+j}} N - \sum_{i=1}^{m_0-1} a_j^i L_i - \sum_{l=1}^{j-1} b_j^l W_l, \text{ for } j \geq 2. \end{aligned}$$

To finish the proof of the Theorem, we show that it is possible to choose the coefficients  $a_j^i$  and  $b_j^l$   $\mathcal{C}^\infty$  in a neighborhood of  $(z^0, w^0)$  so that the basis of vector field  $(L_1, \dots, L_{m_0-1}, W_1, \dots, W_{m-m_0})$  diagonalizes the Levi form of  $r$  in that neighborhood. This is done using an induction argument similar to the one used in the first case: assume that the vector fields  $W_j$ ,  $1 \leq j \leq k$  has been constructed and that their coefficients satisfy

- (1)  $a_j^i$  and  $b_j^l$  are  $\ast \frac{\partial h_{m_0+j}}{\partial w_{m_0+j}}$ , where  $\ast$  is a  $\mathcal{C}^\infty$  function in a neighborhood of  $(z^0, w^0)$ ,
- (2)  $[W_j, \overline{W_j}](\partial r) = \alpha \frac{\partial^2 h_{m_0+j}}{\partial w_{m_0+j} \partial \overline{w_{m_0+j}}}$ , where  $\alpha$  is a  $\mathcal{C}^\infty$  real function in a neighborhood of  $(z^0, w^0)$  greater, in modulus, than  $1/2$ .

As for the first case, note that (2) follows (1). Then, for  $1 \leq j \leq m_0 - 1$ ,

$$[W_{k+1}, \overline{L_j}](\partial r) = -\frac{\partial h_{m_0+k+1}}{\partial w_{m_0+k+1}} [N, \overline{L_j}](\partial r) - a_{k+1}^j [L_j, \overline{L_j}](\partial r),$$

and, for  $j \leq k$ ,

$$[W_{k+1}, \overline{W_j}](\partial r) = -\frac{\partial h_{m_0+k+1}}{\partial w_{m_0+k+1}} [N, \overline{W_j}](\partial r) - b_{k+1}^j [W_j, \overline{W_j}](\partial r),$$

and the results follow, noting that  $[L_j, \overline{L_j}](\partial r)$  is bounded from below by a strictly positive constant in a neighborhood of  $(z^0, w^0)$ , and, as in the first case, that  $[N, \overline{W_j}](\partial r) = \ast \frac{\partial h_{m_0+j}}{\partial w_{m_0+j}}$ , with  $\ast$  is a  $\mathcal{C}^\infty$  function in a neighborhood of  $(z^0, w^0)$ .  $\square$

**Example.** If  $h(w) = \sum h_i(w_i)$ ,  $w_i \in \mathbb{C}$ , each function  $h_i$  being a positive radial analytic function vanishing at the origin, then the hypothesis of the Theorem are verified.

The proof of the second case, shows that Theorem 2.1 is also valid if  $\Omega$  is a smooth strictly pseudo-convex domain in  $\mathbb{C}^n$  (of any dimension). Moreover, applying first the method of the second case and then the one of the first case, this is also true when the rank of the Levi form of  $\rho$  is  $\geq n - 2$ . Thus:

**Theorem 2.2.** Assume that  $\rho = \rho_s$  with  $\rho_s$  satisfying (2.2) as stated in section 2.1 and that  $h$  satisfy Condition III of section 2.2. If the rank of the Levi form of  $\rho$  is  $\geq n - 2$ , then  $\tilde{\Omega}$  is locally diagonalizable at every point of its boundary.

*Remark 2.4.* If  $\Omega$  is a smooth bounded convex domain of finite type in  $\mathbb{C}^n$  we do not know if it is always possible to choose a defining function  $\rho$  and a function  $h$  so that  $\tilde{\Omega}$  is ‘‘completely geometrically separated’’ at any boundary point (we will see in section 4.2 that this is possible near  $\{w = 0\}$ ).

### 3. RELATIONS BETWEEN OPERATORS RELATED TO $\tilde{\Omega}$ AND TO $\Omega$

Assume that  $\Omega$  is a smooth bounded pseudo-convex domain of finite type in  $\mathbb{C}^n$ , that  $\rho = \rho_s$  is a defining function of  $\Omega$  where  $\rho_s$  satisfies (2.2) as stated in section 2.1, and that  $h$  satisfies, at least, Condition II of section 2.2. Thus, by the Corollary of Proposition 2.3, the domain

$$\tilde{\Omega} = \{(z, w) \in \mathbb{C}^{n+m} \text{ s. t. } \rho(z) + h(w) < 0\},$$

is a smooth bounded pseudoconvex domain of  $\mathbb{C}^{n+m}$  of finite type.

In this Section, taking into account the properties of the  $\bar{\partial}$ -Neumann problem for  $\tilde{\Omega}$ , we derive properties for Solutions of the  $\bar{\partial}$ -equation and for the Bergman projections related to the weight

$$(3.1) \quad \omega(z) = \int_{\{h(w) < -\rho(z)\}} d\lambda(w).$$

Suppose  $f = \sum_{i=1}^n f_i d\bar{z}_i$  is a  $(0, 1)$ -form on  $\Omega$ . Consider it as a  $(0, 1)$ -form  $\tilde{f}$  in  $\tilde{\Omega}$ . If  $f$  is  $\bar{\partial}$ -closed, so is  $\tilde{f}$ , and if  $\tilde{u}$  is a solution of  $\bar{\partial}\tilde{u} = \tilde{f}$  in  $\tilde{\Omega}$ , then  $\tilde{u}$  is holomorphic in the variable  $w$  and the function  $u$  defined by  $u(z) = \tilde{u}(z, 0)$  is a solution of the equation  $\bar{\partial}u = f$  in  $\Omega$ . Moreover, for all  $\alpha \in \mathbb{N}^n$ , denoting  $D_z^\alpha = \frac{\partial^{|\alpha|}}{\partial z_1^{\alpha_1} \dots \partial z_n^{\alpha_n}}$ , we have  $D_z^\alpha u(z) = D_z^\alpha \tilde{u}(z, 0)$ ,  $w \mapsto D_z^\alpha \tilde{u}(z, w)$  is holomorphic, for any integer  $p \geq 0$ ,  $w \mapsto |D_z^\alpha \tilde{u}(z, w)|^p$  is pluri-subharmonic, and, by the mean value property,

$$D_z^\alpha u(z) = (\omega(z))^{-1} \int_{\{h(w) < -\rho(z)\}} D_z^\alpha \tilde{u}(z, w) d\lambda(w),$$

and

$$\int_{\Omega} |D_z^\alpha u(z)|^p \omega(z) d\lambda(z) \leq \int_{\tilde{\Omega}} |D_z^\alpha \tilde{u}(z, w)|^p d\lambda(z, w).$$

Thus:

**Lemma 3.1.** *With the conditions and notations stated above, for any  $p \in [1, +\infty]$  and any integer  $t \geq 0$ , denote by  $L_\omega^{p,t}(\Omega)$  the Sobolev space of functions  $g$  (resp. of  $(0, 1)$ -forms  $g = \sum_{i=1}^n g_i d\bar{z}_i$ ) such that, for all  $\alpha \in \mathbb{N}^n$ ,  $|\alpha| \leq t$ ,  $D_z^\alpha g$  belongs to the weighted  $L^p$  space  $L_\omega^p(\Omega) = L^p(\Omega, \omega(z)d\lambda(z))$  (resp. to the weighted space  $L_{(0,1),\omega}^{p,t}(\Omega)$  of  $(0, 1)$ -forms  $g$  on  $\Omega$  whose coefficients  $g_i$  belong to  $L_\omega^{p,t}(\Omega)$ ) equipped with the norm  $\|g\|_{L_\omega^{p,t}(\Omega)} = \sum_{|\alpha| \leq t} \|D_z^\alpha g\|_{L_\omega^p(\Omega)}$  (resp.  $\|g\|_{L_{(0,1),\omega}^{p,t}(\Omega)} = \sum_{i=1}^n \sum_{|\alpha| \leq t} \|D_z^\alpha g_i\|_{L_\omega^p(\Omega)}$ ). Then*

- (1)  $f$  is in  $L_{(0,1),\omega}^{p,t}(\Omega)$  if and only if  $\tilde{f}$  belongs to  $L_{(0,1)}^{p,t}(\tilde{\Omega})$  and, in this case,  $\|f\|_{L_{(0,1),\omega}^{p,t}(\Omega)} = \|\tilde{f}\|_{L_{(0,1)}^{p,t}(\tilde{\Omega})}$ ;
- (2) If  $\tilde{u}$  belongs to  $L^{p,t}(\tilde{\Omega})$  then  $u$  belongs to  $L_\omega^{p,t}(\Omega)$  and  $\|u\|_{L_\omega^{p,t}(\Omega)} \leq \|\tilde{u}\|_{L^{p,t}(\tilde{\Omega})}$ .

Similarly, a function  $u$  belongs to  $L_\omega^2(\Omega)$  if and only if the function  $\tilde{u}$ , defined on  $\tilde{\Omega}$  by  $\tilde{u}(z, w) = u(z)$ , belongs to  $L^2(\tilde{\Omega})$ . So, if  $P_\omega^{\tilde{\Omega}}$  denotes the Bergman projection of  $\tilde{\Omega}$  and  $P_\omega^\Omega$  the Bergman projection of  $\Omega$  with the weight  $\omega$ , if  $h(w) = \sum_{i=1}^p h_i(|w_i|)$ ,  $w_i \in \mathbb{C}^{m_i}$ , by the mean value property we have  $P_\omega^\Omega(u)(z) = P_\omega^{\tilde{\Omega}}(\tilde{u})(z, 0)$ . Then:

**Lemma 3.2.** *With the above notations,*

- (1) We have  $\|u\|_{L_\omega^{p,t}(\Omega)} = \|\tilde{u}\|_{L^{p,t}(\tilde{\Omega})}$ ;
- (2) If  $h(w) = \sum_{i=1}^p h_i(|w_i|)$ ,  $w_i \in \mathbb{C}^{m_i}$ , we have  $P_\omega^\Omega(u)(z) = P_\omega^{\tilde{\Omega}}(\tilde{u})(z, 0)$  and  $\|P_\omega^\Omega(u)\|_{L_\omega^{p,t}(\Omega)} \leq \|P_\omega^{\tilde{\Omega}}(\tilde{u})\|_{L^{p,t}(\tilde{\Omega})}$ ;
- (3) If  $h(w) = \sum_{i=1}^p h_i(|w_i|)$ ,  $w_i \in \mathbb{C}^{m_i}$ , for any  $\bar{\partial}$ -closed  $(0, 1)$ -form  $f \in L_\omega^2(\Omega)$ , denoting  $\tilde{f}(z, w) = f(z)$ ,  $\bar{\partial}^* \mathcal{N}_\Omega^{\tilde{f}}(z, 0)$  is the solution of the equation  $\bar{\partial} u = f$  orthogonal to holomorphic functions in  $L_\omega^2(\Omega)$ ;
- (4) If  $h(w) = \sum_{i=1}^p h_i(|w_i|)$ ,  $w_i \in \mathbb{C}^{m_i}$ , and  $K_{B,\omega}^\Omega$  (resp.  $K_B^\Omega$ ) denotes the Bergman kernel of  $\Omega$  associated to the measure  $\omega(z)d\lambda(z)$  (resp. of  $\tilde{\Omega}$  associated to the Lebesgue measure), we have  $K_{B,\omega}^\Omega(z, \zeta) = K_B^\Omega((z, 0), (\zeta, 0))$ .

Now, we will derive from these Lemmas some simple weighted estimates on  $\Omega$  when the corresponding unweighted estimates are known on  $\tilde{\Omega}$ .

### 3.1. Sobolev estimates for general pseudo-convex domain

As  $\tilde{\Omega}$  is of finite type, by the fundamental result of D. Catlin ([Cat87]) the  $\bar{\partial}$ -Neumann problem of  $\tilde{\Omega}$  satisfies a subelliptic estimate. Then, all the associated operators map continuously the  $L^2$  Sobolev spaces of  $\tilde{\Omega}$  into themselves.

To respect traditional notations, let us denote, for  $t \in \mathbb{N}$ ,  $W_\omega^t(\Omega) = L_\omega^{2,t}(\Omega)$  and  $W_{(0,1),\omega}^t(\Omega) = L_{(0,1),\omega}^{2,t}(\Omega)$ .

Lemmas 3.2 and 3.1 imply thus:

**Theorem 3.1.** *Let  $\Omega$  be a smooth bounded pseudo-convex domain of finite type in  $\mathbb{C}^n$ . Let  $\rho = \rho_s$  be a defining function of  $\Omega$  where  $\rho_s$  satisfies (2.2) as stated in section 2.1. Let  $h$  be a smooth function on  $\mathbb{C}^m$  satisfying Condition II of section 2.2 the functions  $h_i$  being radial. Then,  $\omega$  being the weight defined by (3.1):*

- (1) For any integer  $t$ , if  $f$  is a  $\bar{\partial}$ -closed  $(0, 1)$ -form in  $W_{(0,1),\omega}^t(\Omega)$ , then the solution to the equation  $\bar{\partial} u = f$  orthogonal to holomorphic functions in  $L_\omega^2(\Omega)$  satisfies  $\|u\|_{W_\omega^t(\Omega)} \leq C \|f\|_{W_{(0,1),\omega}^t(\Omega)}$ , the constant  $C$  depending on  $\rho$ ,  $h$  and  $t$ ;
- (2) For any integer  $t$ , the weighted Bergman projection  $P_\omega^\Omega$  maps continuously the Sobolev space  $W_\omega^t(\Omega)$  into itself.

**Corollary.** *Let  $\Omega$  and  $\rho$  be as in the Theorem. Let  $r \geq 0$  be a rational number and  $\varphi_r$  be the pluri-subharmonic function  $\varphi_r = -r \log(-\rho)$  (c.f. Remark 2.1). Let us denote by  $\mathcal{N}_{\varphi_r}$  the  $\bar{\partial}$ -Neumann operator for the weight  $e^{-\varphi_r}$  acting on  $(0, 1)$ -forms and by  $\bar{\partial}_{\varphi_r}^* \mathcal{N}_{\varphi_r}^{(0,1)}$  the restriction to the space of  $\bar{\partial}$ -closed forms in  $L_{(0,1),(-\rho)^r}^2(\Omega)$  of the operator  $\bar{\partial}_{\varphi_r}^* \mathcal{N}_{\varphi_r}$  giving the  $L_{(-\rho)^r}^2$  minimal solution of the  $\bar{\partial}$ -equation. Let us denote by  $\mathcal{B}_{\varphi_r}$  the Bergman projection of  $L_{\varphi_r}^2(\Omega)$ . Then:*

- (1) For all  $t \geq 0$ ,  $\bar{\partial}_{\varphi_r}^* \mathcal{N}_{\varphi_r}^{(0,1)}$  maps continuously the subspace of  $\bar{\partial}$ -closed forms of  $W_{(0,1),(-\rho)^r}^t(\Omega)$  into  $W_{(-\rho)^r}^t(\Omega)$ .
- (2) For all real number  $t$ ,  $\mathcal{B}_{\varphi_r}$  maps continuously  $W_{(-\rho)^r}^t(\Omega)$  into itself.

*Remark 3.1.*

- (1) In the Corollary, the function  $h$  is equal to  $\sum |w_i|^{2q_i}$ ,  $w_i \in \mathbb{C}$ , the integers  $q_i$  being chosen so that  $r = \sum \frac{1}{q_i}$ .  $\tilde{\Omega}$  being of finite type, there is a gain in the Sobolev scale for the estimates of the  $\bar{\partial}$ -Neumann problem on  $\tilde{\Omega}$ . This implies a similar gain for  $\bar{\partial}_{\varphi_r}^* \mathcal{N}_{\varphi_r}^{(0,1)}$ . But this gain is the inverse on the type of  $\tilde{\Omega}$  which is given in Corollary 1 of Proposition 2.3 and, then, can be very small depending on  $r$ .
- (2) If  $\Omega$  is a smooth bounded pseudo-convex domain in  $\mathbb{C}^n$  admitting a defining function  $\rho$  which is pluri-subharmonic in  $\bar{\Omega}$  then:

- (a) If  $h$  is a positive pluri-subharmonic function satisfying  $\nabla h(w) \neq 0$  if  $w \neq 0$ ,  $\lim_{|w| \rightarrow +\infty} h(w) = +\infty$  and  $h(w) = \sum_{i=1}^p h_i(|w_i|)$ ,  $w = (w_1, \dots, w_p)$ ,  $w_i \in \mathbb{C}^{m_i}$ , then  $\tilde{\Omega}$  admits a pluri-subharmonic defining function and, applying a theorem of H. Boas & E. Straube ([BS91]), we get that, for all real number  $t$ , the weighted Bergman projection  $P_{\omega}^{\Omega}$  maps continuously the Sobolev spaces  $W_{\omega}^t(\Omega)$  into themselves.
- (b) Moreover, if  $\rho$  is of finite type in  $\tilde{\Omega}$  then Corollary 2 of Proposition 2.3 shows that the results of the Theorem and the Corollary are also valid using  $\rho$  to define  $\tilde{\Omega}$  and thus for other weights  $\omega$ .
- (3) By the Corollary of Theorem 1.2, in the previous Theorem and it's Corollary, the estimates concerning the minimal solution of the  $\bar{\partial}$ -equation and the Bergman kernel are also valid for any weight of the form  $k\omega$  where  $k$  satisfies  $0 < c \leq k \leq C$ ,  $|k^{(i)}| \leq C_i$ ,  $i = 1, 2$ . In particular, for the Corollary, these weights are exactly the weights equivalent (in that sense) to a rational power of the distance to the boundary of  $\Omega$ .

### 3.2. Lipschitz estimates for domains in $\mathbb{C}^2$

Here we obtain estimates on weighted Bergman projections of  $\Omega$  using only properties of the Bergman projection of the domain  $\tilde{\Omega}$ . For this we need  $\tilde{\Omega}$  to be “completely geometrically separated” and we assume:

$\Omega$  is a domain in  $\mathbb{C}^2$ , or the rank of the Levi form of  $\partial\Omega$  is  $\geq n - 2$ ,  $\rho = \rho_s$  with  $\rho_s$  satisfying (2.2) as stated in section 2.1 and  $h$  satisfies Condition III of section 2.2. We denote by  $\omega = \omega_{\rho, h}$  the associated weight (equation (3.1)).

Let  $M$  be the type of  $\Omega$ . By [CD06b, CD06a], we know that  $P_{\tilde{\Omega}}$  maps continuously the Lipschitz space  $\Lambda_{\alpha}(\tilde{\Omega})$ ,  $\alpha \geq 0$ , into itself and that the space of holomorphic functions in  $\Lambda_{\alpha}(\Omega)$  is continuously embedded in the anisotropic Lipschitz space  $\Gamma_{\alpha}(\Omega)$  for  $\alpha < 1/M$ . Then Lemma 3.2 and the Corollary of Theorem 1.2 give immediately:

**Theorem 3.2.** *In the conditions stated above for  $\Omega$ ,  $\rho$  and  $h$ , let  $\omega_1$  be a weight equivalent to  $\omega$  (i.e.  $\omega_1 = k\omega_{\rho, h}$ ,  $\omega_{\rho, h}$  given by equation (3.1), the positive function  $k$  satisfying the conditions stated in (2) of Theorem 1.1). Then the weighted Bergman projection  $P_{\omega_1}^{\Omega}$  maps continuously the Lipschitz space  $\Lambda_{\alpha}(\Omega)$  into itself for all  $\alpha \geq 0$  and into the anisotropic Lipschitz space  $\Gamma_{\alpha}(\Omega)$  for  $\alpha < 1/M$ .*

*Remark.*

- (1) In the next Section, using pointwise estimates of the kernel of  $P_{\omega}^{\Omega}$  we will extend the Lipschitz estimate for  $P_{\omega}^{\Omega}$  to convex domains of finite type in  $\mathbb{C}^n$  but for a smaller class of weights  $\omega$ .
- (2) In the conditions of the preceding Theorem, assuming that the weight  $\omega$  is equivalent to a positive rational power of the distance to the boundary ( $\omega = k\delta^q$ ,  $\delta$  being the distance to  $\partial\Omega$ ,  $q \in \mathbb{Q}_*^+$  and  $k$  satisfying the conditions stated in (1) of Theorem 1.1) using that the Bergman projection  $P_{\tilde{\Omega}}$  of  $\tilde{\Omega}$  maps continuously the Sobolev spaces  $L_s^p$  ( $1 < p < +\infty$ ,  $s \in \mathbb{N}$ ) into themselves ([CD06b, CD06a]), we get immediately that the weighted Bergman projections  $P_{\omega}^{\Omega}$  maps continuously the (weighted) Sobolev spaces  $L_s^p(\delta^q d\lambda)$  into themselves. Using pointwise estimates of the kernel of  $P_{\tilde{\Omega}}$  ([CD06b]) these estimates can be extended to the Sobolev spaces  $L_s^p(\delta^t d\lambda)$  for  $-1 < t - q < p - 1$ .

In the next Section, establishing pointwise estimates of the kernel of  $P_{\omega}^{\Omega}$  we will get the (better) results stated in Theorem 1.1.

## 4. SHARP ESTIMATES OF THE WEIGHTED BERGMAN KERNEL AND PROOF OF THEOREM 1.1

The aim of this Section is to establish precise pointwise estimates of the kernel of the weighted Bergman projection  $P_{\omega}^{\Omega}$  in terms of the geometry of  $\Omega$  (from which we will deduce Theorem 1.1) using pointwise estimates of the kernel of  $P_{\tilde{\Omega}}$  and Lemma 3.2. Hence we need, at least, that the domain  $\tilde{\Omega}$  is “completely geometrically separated” near the set  $\{(z, 0) \in \tilde{\Omega}\}$  and to have a precise comparison of the geometries of  $\tilde{\Omega}$  and  $\Omega$ . We are able to do this in the two following cases:

- $\Omega$  is a finite type domain in  $\mathbb{C}^2$ ,  $\rho = \rho_s$  satisfies (2.2) and  $h$  satisfies Condition IV of section 2.2
- $\Omega$  is a convex domain of finite type in  $\mathbb{C}^n$ ,  $\rho = g^4 e^{-1/g} - 1$  where  $g$  is the gauge function of  $\Omega$ , so that  $\rho$  is convex and of finite type in a neighborhood of  $\partial\Omega$ , and  $h$  satisfies Condition V of section 2.2

(the weight  $\omega$  being given by (3.1)).

Note that, for the convex case, we are not able, in general, to get complete geometric separation of  $\tilde{\Omega}$  near  $\partial\Omega \times \{0\}$  if we use a Diederich-Fornaess defining function (see Proposition 2.4). This property being indispensable in this Section we need to use another defining function defined using the gauge: doing this, we loose the property of finite type everywhere on  $\partial\tilde{\Omega}$  and so the global properties of the Bergman projection  $P_{\tilde{\Omega}}$  but local estimates of the kernel of  $P_{\tilde{\Omega}}$  will suffice for our purpose.

### 4.1. The case of finite type domains in $\mathbb{C}^2$

#### 4.1.1. Precise comparison between the geometries of $\Omega$ and $\tilde{\Omega}$

We assume that the defining function  $\rho$  of  $\Omega$  is  $\rho = \rho_s$  where  $\rho_s$  satisfies (2.2) as stated in section 2.1 and that the function  $h$  satisfies Condition IV of section 2.2 (so that  $\tilde{\Omega}$  is of finite type and has a Levi form locally diagonalizable at every point of  $\partial\tilde{\Omega}$ ).

We use the notations of the proof of Theorem 2.1 for  $L$ ,  $N$ ,  $L_1$  and  $W_k$ , and let us denote by  $\tilde{N}$  the complex normal to the defining function  $\tilde{\rho}$  of  $\tilde{\Omega}$  (i.e.  $\tilde{N}(\tilde{\rho}) \equiv 1$  in a neighborhood of the boundary of  $\tilde{\Omega}$ ). Moreover, for the geometries, using the

notation “ $F^\Omega(L, z, \delta)$ ” introduced in Section 2 of [CD08], we denote  $\tilde{F}_1(\tilde{\zeta}, \delta) = \delta^{-2}$ ,  $\tilde{F}_2(\tilde{\zeta}, \delta) = F^{\tilde{\Omega}}(L_1, \tilde{\zeta}, \delta)$ ,  $\tilde{F}_i(\tilde{\zeta}, \delta) = F^{\tilde{\Omega}}(W_{i-2}, \tilde{\zeta}, \delta)$  and  $F_L(\zeta, \delta) = F^\Omega(L, \zeta, \delta)$ .

Let us first compare the weights  $\tilde{F}_i(\tilde{\zeta}, \delta)$  and  $F_L(\zeta, \delta)$  constructed with the extremal basis defined in section 2.5.

**Lemma 4.1.** *We have:*

- (1)  $L_1 = L - \sum * W_k = L - \sum * \frac{\partial}{\partial w_i} - \left( \sum * \frac{\partial h_i}{\partial w_i} \right) N$ , where  $*$  are  $\mathcal{C}^\infty$  functions;
- (2)  $N = \beta \tilde{N} + \sum \left( * \frac{\partial h_i}{\partial w_i} + \sum_{j>i} * \left| \frac{\partial h_j}{\partial w_j} \right|^2 \right) W_i$ , where  $\beta$  and  $*$  are  $\mathcal{C}^\infty$  functions,  $\beta \simeq 1$  for  $|w|$  small.

*Proof.* Part (1) is a trivial consequence of the definitions of the vector fields. Let us give some indications for part (2). We have

$$\tilde{N} = \frac{\bar{\nabla} \rho + \bar{\nabla} h}{|\bar{\nabla} \rho|^2 + |\bar{\nabla} h|^2} \text{ and } N = \frac{\bar{\nabla} \rho}{|\bar{\nabla} \rho|^2}.$$

Thus

$$\begin{aligned} N &= \frac{|\bar{\nabla} \rho|^2 + |\bar{\nabla} h|^2}{|\bar{\nabla} \rho|^2} \tilde{N} + * \bar{\nabla} h \cdot \frac{\partial}{\partial w} \\ &= \beta \tilde{N} + * \sum \frac{\partial h_i}{\partial w_i} \left( Z_i + \frac{\partial h_i}{\partial w_i} N \right), \end{aligned}$$

and

$$\left( 1 - \sum * \left| \frac{\partial h_i}{\partial w_i} \right|^2 \right) N = \beta \tilde{N} + \sum \left( * \frac{\partial h_i}{\partial w_i} + \sum_{j>i} * \left| \frac{\partial h_j}{\partial w_j} \right|^2 \right) W_i.$$

□

Now we apply this Lemma to estimate the weights  $\tilde{F}_i(\tilde{\zeta}, \delta)$  for  $\tilde{\zeta} = (\zeta, 0) \in \partial \tilde{\Omega}$ .

Denoting  $c_{11} = [L, \bar{L}](\partial \rho)$ , we have

$$\begin{aligned} \tilde{c}_{11} &= [L_1, \bar{L}_1](\partial r) \\ &= \left[ L - \sum * \frac{\partial}{\partial w_i} - \left( \sum * \frac{\partial h_i}{\partial w_i} \right) N, \bar{L} - \sum * \frac{\partial}{\partial w_i} - \left( \sum * \frac{\partial h_i}{\partial w_i} \right) \bar{N} \right](\partial r) \\ &= c_{11} + \sum * \frac{\partial h_i}{\partial w_i} + \sum * \frac{\partial h_i}{\partial w_i}. \end{aligned}$$

Then the order of cancellation of the functions  $h_i$  being greater than the type of  $\Omega$ , it is obvious that  $\tilde{F}_1(\tilde{\zeta}, \delta) \simeq F_1(\zeta, \delta)$ .

Furthermore, in 2, we saw that  $[W_k, \bar{W}_k](\partial r) = \alpha_k \frac{\partial^2 h_k}{\partial w_k \partial \bar{w}_k}$ ,  $\alpha_k \in \mathcal{C}^\infty$ ,  $|\alpha_k| \geq 1/2$ , then, as  $W_k = \frac{\partial}{\partial w_k} + * N + \sum_{i<k} * \frac{\partial}{\partial w_i}$  and

$$\tilde{F}_{k+2} = \sum_{\mathcal{L} \in \mathcal{L}_k} \left( \frac{\mathcal{L}([W_k, \bar{W}_k](\partial r))}{\delta} \right)^{2/|\mathcal{L}|+2},$$

it follows clearly that

$$\tilde{F}_{k+2}(\tilde{\zeta}, \delta) \simeq \left( \frac{1}{\delta} \right)^{1/q_k} \lesssim \tilde{F}_1(\tilde{\zeta}, \delta).$$

Finally, we compare the pseudo-distances in  $\Omega$  and  $\tilde{\Omega}$ .

Let  $\gamma$  and  $\tilde{\gamma}$  be the respective pseudo-distances in  $\partial \Omega$  and  $\partial \tilde{\Omega}$  defined by the exponential map of tangent vectors fields associated to extremal basis (see [CD08] and [CD06b, p. 75 and 100]). Then

**Lemma 4.2.** *With the above notations,  $\gamma(z, \zeta) \simeq \tilde{\gamma}(\tilde{z}, \tilde{\zeta})$ , for  $z$  and  $\zeta$  in  $\partial \Omega$ .*

*Proof.* We use the notations introduced in [CD06b]: for  $\Omega$ ,  $\mathcal{Y}_1 = \Re N$ ,  $\mathcal{Y}_2 = \Im N$ ,  $\mathcal{Y}_3 = \Re L_1$ ,  $\mathcal{Y}_4 = \Im L_1$  and, for  $\tilde{\Omega}$ ,  $\tilde{\mathcal{Y}}_1 = \Re \tilde{N}$ ,  $\tilde{\mathcal{Y}}_2 = \Im \tilde{N}$ ,  $\tilde{\mathcal{Y}}_3 = \Re \tilde{L}_1$ ,  $\tilde{\mathcal{Y}}_4 = \Im \tilde{L}_1$  and  $\tilde{\mathcal{Y}}_{2k+3} = \Re W_k$ ,  $\tilde{\mathcal{Y}}_{2k+4} = \Im W_k$ . Then, following [CD06b, p. 100], there exists  $\varphi : [0, 1] \rightarrow \mathbb{C}^2$ , piecewise  $\mathcal{C}^1$ , such that  $\varphi(0) = z$ ,  $\varphi(1) = \zeta$ ,  $\varphi'(t) = \sum a_i(t) \mathcal{Y}_i(\varphi(t))$  a.e. with  $|a_i(t)| \lesssim G_i(z, \gamma(z, \zeta))^{-1/2}$ , where  $G_1 = G_2 = 1/\delta^2$  and  $G_3 = G_4 = F_1(z, \delta)$ . Now, let  $\tilde{\varphi}(t) = (\varphi(t), 0)$  be the same curve considered in  $\mathbb{C}^{2+m}$ . Then Lemma 4.1 and  $G_1(z, \delta) = \tilde{G}_1(\tilde{z}, \delta) = \frac{1}{\delta^2} \gtrsim F_1(z, \delta) \simeq \tilde{F}_1(\tilde{z}, \delta) \gtrsim \tilde{F}_k(\tilde{z}, \delta) = \tilde{G}_{2k+1}(\tilde{z}, \delta) = \tilde{G}_{2k+2}(\tilde{z}, \delta)$ ,  $k \geq 2$  show that  $\tilde{\varphi}$  satisfies  $\tilde{\varphi}'(t) = \sum b_i(t) \tilde{\mathcal{Y}}_i(\tilde{\varphi}(t))$  with  $|b_i(t)| \lesssim \tilde{G}_i(\tilde{z}, \gamma(z, \zeta))^{-1/2}$ . Thus

$$\tilde{\gamma}(\tilde{z}, \tilde{\zeta}) \lesssim \gamma(z, \zeta).$$

To show the converse inequality, consider a curve  $\tilde{\varphi} : [0, 1] \rightarrow \mathbb{C}^{2+m}$  such that  $\tilde{\varphi}(0) = \tilde{z}$ ,  $\tilde{\varphi}(1) = \tilde{\zeta}$ ,  $\tilde{\varphi}'(t) = \sum b_i(t) \tilde{\mathcal{Y}}_i(\tilde{\varphi}(t))$  a.e. with  $|b_i(t)| \lesssim \tilde{G}_i(\tilde{z}, \tilde{\gamma}(\tilde{z}, \tilde{\zeta}))^{-1/2}$ .

First, we consider the component  $\tilde{\varphi}_{2+i}(t)$  of the curve  $\tilde{\varphi}$ . Let us decompose  $\tilde{\varphi}'$  on the basis  $L, N, \frac{\partial}{\partial w_i}$  and their conjugates. Then the coefficients  $\tilde{w}_i(\tilde{\varphi}'(t))$  of  $\tilde{\varphi}'(t)$  in the directions  $\frac{\partial}{\partial w_i}$  or  $\frac{\partial}{\partial \bar{w}_i}$  is, in modulus, bounded from above by

$$F_1\left(z, \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)\right)^{-1/2} + \tilde{F}_i\left(\tilde{z}, \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)\right)^{-1/2} + \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right) \lesssim \tilde{F}_i\left(\tilde{z}, \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)\right)^{-1/2} \lesssim \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)^{1/q_i},$$

and thus, since  $\tilde{\varphi}_{2+i}(0) = 0$ ,

$$|\tilde{\varphi}_{2+i}(t)| \lesssim \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)^{1/q_i}.$$

Now let us denote  $\varphi(t) = (\tilde{\varphi}_1(t), \tilde{\varphi}_2(t))$  the projection of  $\tilde{\varphi}$  onto  $\mathbb{C}^2$ , and let us write  $\varphi'(t) = \sum_{i=1}^4 c_i(t) \mathcal{D}_i(\varphi(t))$ . We have to estimate the contribution of the coefficients  $b_j$  to the coefficient  $c_i$ .

Suppose  $j > 4$ . The contribution to  $c_3$  and  $c_4$  is null and to  $c_1$  and  $c_2$  is bounded by  $\frac{\partial h}{\partial w_j} \tilde{F}_{j+2}\left(\tilde{z}, \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)\right)^{-1/2}$  (with a evident correspondence  $j' \leftarrow j$ ). As  $\left|\frac{\partial h_{j'}}{\partial w_{j'}}\right| \lesssim |w_{j'}|^{2q_{j'}-1} \lesssim \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)^{\frac{2q_{j'}-1}{2q_{j'}}}$ , this contribution is bounded by

$$\tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)^{\frac{2q_{j'}-1}{2q_{j'}}} \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)^{\frac{1}{2q_{j'}}} = \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right).$$

If  $j = 4$  or  $3$ , the contribution to  $c_4$  and  $c_3$  is bounded by  $\tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)$ , and the contribution to  $c_1$  and  $c_2$  is bounded by

$$\sum \left| \frac{\partial h_i}{\partial w_i} \right| \tilde{F}_1\left(\tilde{z}, \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)\right)^{-1/2} \lesssim \tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right).$$

When  $j = 1$  or  $2$ , the contribution is bounded by  $\tilde{\gamma}\left(\tilde{z}, \tilde{\zeta}\right)$ .

This proves the Lemma.  $\square$

#### 4.1.2. Pointwise estimate of the Bergman kernel

**Theorem 4.1.** *Assume that  $\Omega$  is pseudo-convex of finite type in  $\mathbb{C}^2$  and that the hypothesis on  $\rho, h$ , stated at the beginning of the section are satisfied. Let  $L$  be the complex tangent vector field to  $\rho$  defined by  $L = \frac{\partial \rho}{\partial z_2} \frac{\partial}{\partial z_1} - \frac{\partial \rho}{\partial z_1} \frac{\partial}{\partial z_2}$  and  $N$  be the normal one such that  $N\rho \equiv 1$  in a small neighborhood  $U$  of  $\partial\Omega$ . Let  $\mathcal{L}$  be a list of vector fields belonging to  $\{L, \bar{L}, N, \bar{N}\}$ .*

*Let us denote by  $K_\omega^\Omega$  the Bergman kernel of  $L_\omega^2(\Omega)$  for the weight  $\omega$ . Then for sufficiently close points  $p_1$  and  $p_2$  in  $U$ , we have the following estimate:*

$$\begin{aligned} \left| \mathcal{L} K_\omega^\Omega(p_1, p_2) \right| &\leq C_{|\mathcal{L}|} \left( \frac{1}{\delta(p_1, p_2)^2} \right)^{1+l_N/2} F_L^{1+l_L/2}(p_1, \delta(p_1, p_2)) \prod_{j=1}^m \left( \frac{1}{\delta(p_1, p_2)} \right)^{1/q_j} \\ &\simeq C_{|\mathcal{L}|} \frac{\left( \frac{1}{\delta(p_1, p_2)^2} \right)^{l_N/2} F_L^{l_L/2}(p_1, \delta(p_1, p_2))}{\text{Vol}_\omega(B(p_1, \delta(p_1, p_2)))}, \end{aligned}$$

where  $F_L$  is the weight associated to  $L$ ,  $l_L$  (resp.  $l_N$ ) denotes the number of times  $L$  or  $\bar{L}$  (resp.  $N$  or  $\bar{N}$ ) appears in the list  $\mathcal{L}$ ,  $\delta(p_1, p_2) = |\rho(p_1)| + |\rho(p_2)| + \delta_\Omega(p_1, p_2)$ ,  $\delta_\Omega(p_1, p_2) = \gamma(\pi(p_1), \pi(p_2))$ ,  $\gamma$  being the pseudo-distance on  $\partial\Omega$ , and  $B(p_1, \delta(p_1, p_2))$  the associated pseudo-ball, of the geometry describe in [CD06b] and  $\text{Vol}_\omega$  denotes the volume with respect to the measure  $\omega(z)d\lambda(z)$ .

*Proof.* We use (3) of Lemma 3.2 and the sharp estimates on  $K_B^\Omega$  deduced from [CD06b].

Suppose  $|\mathcal{L}| = 0$ . We have  $\left| K_B^\Omega(\tilde{z}, \tilde{\zeta}) \right| \lesssim \text{Vol}\left(\tilde{B}\left(\tilde{z}, \tilde{\delta}\left(\tilde{z}, \tilde{\zeta}\right)\right)\right)^{-1}$ , by [CD06b, Main Theorem on the Bergman kernel, part II, p. 77], with, by Lemma 4.2,

$$\tilde{\delta}\left(\tilde{z}, \tilde{\zeta}\right) = |r(\tilde{z})| + |r(\tilde{\zeta})| + \delta_{\tilde{\Omega}}\left(\tilde{z}, \tilde{\zeta}\right) \simeq |\rho(z)| + |\rho(\zeta)| + \delta_\Omega(z, \zeta) =: \delta,$$

and, by [CD06b, Section 3],  $\left| K_B^\Omega(\tilde{z}, \tilde{\zeta}) \right| \lesssim \text{Vol}\left(\tilde{B}\left(\pi(\tilde{z}), \tilde{\delta}\left(\tilde{z}, \tilde{\zeta}\right)\right)\right)^{-1}$ .

The estimates of the functions  $F_i$  and  $\tilde{F}_i$  in the proofs of the Lemmas show that

$$\text{Vol}\left(\tilde{B}\left(\pi(\tilde{z}), \delta\right)\right) \simeq \delta^2 \tilde{F}_1\left(\pi(\tilde{z}), \delta\right) \prod_{k=1}^m \delta^{1/q_k} \simeq \delta^{2+\sum_k 1/q_k} F_1(z, \delta).$$

To finish the proof, we have to estimate  $\text{Vol}(B(z, \delta)) = \int_{B(z, \delta)} \omega(\xi) dV(\xi)$ :

$$\begin{aligned} \omega(\xi) &= \text{Vol}\{w \text{ such that } h(w) < -\rho(\xi)\} \\ &\simeq \prod \text{Vol}\{w_i \text{ such that } h_i(w_i) < \rho(\xi)\} \simeq \prod (-\rho(\xi))^{\Sigma 1/q_i}. \end{aligned}$$

Then, using the fact that  $\xi \in B(z, \delta)$  and  $|\rho(z)| \lesssim \delta$  imply  $|\rho(\xi)| \lesssim \delta$  (see [CD06b, CD08]), we obtain

$$\text{Vol}(B(z, \delta)) \simeq \delta^{2+\Sigma 1/q_i} F_1(z, \delta)^{-1},$$

which finishes the proof when  $\|\mathcal{L}\| = 0$ .

When  $|\mathcal{L}| \geq 1$ , the proof is done similarly using Lemma 4.1, 4.2 and the inequalities on the derivatives of  $K_B^{\tilde{\Omega}}$  given in [CD06b].  $\square$

*Remark.* This proof easily generalizes in higher dimensions  $n$  when the Levi form of  $\rho$  has a rank  $\geq n - 2$ .

## 4.2. The case of convex domains of finite type in $\mathbb{C}^n$

Now we assume that the function  $h$  satisfies Condition V of section 2.2 (for example  $h(w) = \sum |w_i|^{2q_i}$ ,  $w_i \in \mathbb{C}$ ).

### 4.2.1. Choice of the defining function and geometry of $\tilde{\Omega}$

Because of Remark 2.4, we have to choose a special defining function to obtain useful properties on  $\tilde{\Omega}$ .

Let  $g$  be the gauge function for  $\Omega$ . Then  $\rho = g^4 e^{-1/g} - 1$  is a smooth convex defining function for  $\Omega$  which is of finite type in a neighborhood of  $\partial\Omega$ . Thus the domain  $\tilde{\Omega} = \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^m \text{ such that } \rho(z) + h(w) < 0\}$  is smooth, convex and of finite type in a neighborhood of  $\partial\Omega \times \{0\}$ .

To get a useful estimate of the Bergman kernel of  $\tilde{\Omega}$ , we need a precise comparison between the geometries of  $\partial\Omega$  and of  $\partial\tilde{\Omega}$  near the points of  $\partial\Omega \times \{0\}$ .

Let  $P_0 \in \partial\Omega$ ,  $\tilde{P}_0 = (P_0, 0) \in \partial\tilde{\Omega}$  and  $\delta > 0$  sufficiently small. We now investigate extremal basis at  $P_0$  and  $\tilde{P}_0$  (in the sense of [McN94, Hef04]).

**Lemma 4.3.** *Let  $(z_1^\delta, \dots, z_n^\delta)$  be a  $\delta$ -extremal coordinate system at  $P_0$ . Then it is possible to choose a  $\delta$ -extremal coordinate system at  $\tilde{P}_0$ ,  $(\tilde{z}_1^\delta, \dots, \tilde{z}_{m+n}^\delta)$ , such that, for  $1 \leq i \leq n$ ,  $\tilde{z}_i^\delta = (z_i^\delta, 0)$ .*

*Proof.* Let

$$\tilde{H}_\delta = \left\{ \tilde{P} \in \mathbb{C}^{n+m} \text{ such that } r(\tilde{P}) = \delta \right\}.$$

Let  $\tilde{P}_1 \in \tilde{H}_\delta$  such that  $|\tilde{P}_1 - \tilde{P}_0|$  is the euclidean distance  $d(\tilde{P}_0, \tilde{H}_\delta)$  from  $\tilde{P}_0$  to  $\tilde{H}_\delta$ . Let  $Q_1$  be the projection of  $\tilde{P}_1$  to  $\mathbb{C}^n$  so that  $\tilde{P}_1 = (Q_1, w)$ . We have

$$\delta - \rho(Q_1) = h(w) \asymp \sum |w_i|^{2q_i},$$

and, by the condition on the  $q_i$  (recall that  $h$  satisfies Condition V and then  $q_i > \text{typ}(\Omega) = \tau$ ) we obtain

$$|w|^2 \gtrsim (\delta - \rho(Q_1))^{2/2\tau+2}.$$

On the other hand, the geometric properties of  $\Omega$  show that there exists

$$Q_2 \in H_\delta = \{P \in \mathbb{C}^n \text{ such that } \rho(P) = \delta\}$$

such that the distance  $d(Q_2, H_\delta)$  from  $Q_2$  to  $H_\delta$  is less than  $C(\delta - \rho(Q_1))^{1/\tau}$  (with a constant  $C$  independent of  $\delta$ ), and, by the definition of  $\tilde{P}_1$ ,

$$|P_1 - Q_1|^2 + c(\delta - \rho(Q_1))^{2/2\tau+2} \leq \left( |P_1 - Q_1| + C(\delta - \rho(Q_1))^{1/\tau} \right)^2$$

which implies, for  $\delta$  small enough (depending on  $c, C$  and  $\tau$ , i.e. on  $\Omega$ ),  $\rho(Q_1) = \delta$  and we can choose  $\tilde{z}_1^\delta = (z_1^\delta, 0)$ .

Define now  $\tilde{H}_{2,\delta}$  as the intersection of  $\tilde{H}_\delta$  with the affine complex space orthogonal to  $\tilde{z}_1^\delta$  passing through  $\tilde{P}_0$ . Let  $\tilde{P}_2$  be such that  $|\tilde{P}_2 - \tilde{P}_0|$  is the euclidean distance from  $\tilde{P}_0$  to  $\tilde{H}_{2,\delta}$ . Let  $w_0 \in \partial\Omega$  and  $U$  be a small neighborhood of  $w_0$ . Arguing as before, it is easy to show that  $\tilde{P}_2 \in \partial\Omega \times \{0\}$  and we can choose  $\tilde{z}_2^\delta = (z_2^\delta, 0)$ . The proof is finished by induction.  $\square$

**Corollary.** *Let  $L_1, \dots, L_n$  be the  $\delta$ -extremal basis of vector fields associated to the  $\delta$ -extremal coordinate system at the point  $P_0 \in \partial\Omega \cap U$  defined in Lemma 4.3 (see [CD08]). Then the basis  $\tilde{L}_1, \dots, \tilde{L}_{n+m}$  defined by:*

- (1) for  $1 \leq i \leq n$ ,  $L_i = \tilde{L}_i$ ,
- (2) for  $1 \leq j \leq m$ ,  $\tilde{L}_{n+j} = \frac{\partial}{\partial w_j} - \beta_{n+j} \frac{\partial}{\partial Z_1}$ , where  $Z_1$  is the complex normal to  $\partial\Omega$  at the point  $w_0$ ,  $\beta_{n+j}$  being so that  $\tilde{L}_{n+j}$  is tangent to  $\partial\tilde{\Omega}$ .

is  $\delta$ -extremal at  $\tilde{P}_0$ .

*Proof.* For the point (1). note that, for  $i \geq 2$ ,  $L_i = \frac{\partial}{\partial z_i} - \beta_i \frac{\partial}{\partial Z_1}$  (see [CD08, Section 7.1]) (recall that  $L_1 = \tilde{L}_1 = N$ ). For (2), without loss of generality, we can assume  $q_{j+1} \geq q_j$ ,  $1 \leq j \leq m - 1$ , and the result is trivial if  $h_i(w_i) = |w_i|^{2q_i}$  and “easy” to prove in the general case using [CD08].  $\square$

Let  $F_i$  and  $\tilde{F}_i$  be the weights defined with the vector fields  $L_i$  and  $\tilde{L}_i$ . Then

**Lemma 4.4.** *For  $1 \leq i \leq n$ ,  $F_i(z, \delta) = \tilde{F}_i(\tilde{z}, \delta)$  and, for  $1 \leq j \leq m$ ,  $\tilde{F}_{n+j}(\tilde{z}, \delta) \simeq \left(\frac{1}{\delta}\right)^{1/q_j}$ , for  $z \in U$  and  $\tilde{z} = (z, 0)$ .*

*Proof.* The first part is a trivial consequence of the preceding Corollary and the second is proved, as in the case of dimension 2, noting that  $\beta_{n+j} = -\frac{\partial h_j(w_j)}{\partial w_j} / \frac{\partial \rho}{\partial Z_1}$ , with  $\frac{\partial \rho}{\partial Z_1} \in \mathcal{C}^\infty$  and close to 1 for  $\delta$  small.  $\square$

### 4.2.2. Pointwise estimate of the Bergman kernel

**Theorem 4.2.** Assume  $\Omega$  is convex of finite type in  $\mathbb{C}^n$  and that the hypothesis on  $\rho$ ,  $h$ , and  $\omega$  stated at the beginning of the section are satisfied. Let  $w_0$  be a boundary point of  $\Omega$  and  $U$  a small neighborhood of  $w_0$ . Let  $N$  be the complex normal to  $\partial\Omega$  (i.e.  $N\rho \equiv 1$  in a neighborhood  $U$  of  $\partial\Omega$ ). Let  $p_1$  and  $p_2$  be two points in  $U$  and  $\delta_\Omega(p_1, p_2)$  as in Theorem 4.1. Let  $\{L_2, \dots, L_n\}$  be a  $\delta(p_1, p_2)$ -extremal basis associated to  $\rho$  at the point  $p_1$  (with  $\delta(p_1, p_2) = |\rho(p_1)| + |\rho(p_2)| + \delta_\Omega(p_1, p_2)$ ). Let us denote  $L_1 = N$ . Let  $\mathcal{L}$  be a list of vector fields belonging to  $\{L_1, \bar{L}_1, \dots, L_n, \bar{L}_n, N, \bar{N}\}$ . Let  $K_\omega^\Omega$  be the Bergman kernel of  $L_\omega^2(\Omega)$  for the weight  $\omega$ . Then

$$\begin{aligned} \left| \mathcal{L} K_\omega^\Omega(p_1, p_2) \right| &\leq C_{|\mathcal{L}|} \left( \frac{1}{\delta(p_1, p_2)^2} \right)^{1+l_N/2} \mathcal{F}^{1+\mathcal{L}/2}(p_1, \delta(p_1, p_2)) \prod_{j=1}^m \left( \frac{1}{\delta(p_1, p_2)} \right)^{1/q_j} \\ &\simeq C_{|\mathcal{L}|} \frac{\left( \frac{1}{\delta(p_1, p_2)^2} \right)^{l_N/2} \mathcal{F}^{\mathcal{L}/2}(p_1, \delta(p_1, p_2))}{\text{Vol}_\omega(B(p_1, \delta(p_1, p_2)))}, \end{aligned}$$

where  $l_N$  denotes the number of times  $N$  or  $\bar{N}$  appears in the list  $\mathcal{L}$ ,  $\mathcal{F}^{1+\mathcal{L}/2}(p_1, \delta(p_1, p_2)) = \prod_{i=2}^n \mathcal{F}_{L_i}^{1+l_i/2}$ ,  $l_i$  being the number of times  $L_i$  or  $\bar{L}_i$  appears in the list  $\mathcal{L}$  and  $\text{Vol}_\omega$  denotes the volume with respect to the measure  $\omega(z)d\lambda(z)$ .

*Proof.* The construction made before shows that the estimate is immediate because, the exponential map being a local diffeomorphism ([CD06b, p. 75]), the fact that  $L_i = \tilde{L}_i$ ,  $1 \leq i \leq n$  (Corollary of Lemma 4.3), implies  $\delta_\Omega(p_1, p_2) = \delta_{\tilde{\Omega}}(p_1, p_2)$ .  $\square$

### 4.3. Proof of Theorem 1.1

In the two cases we consider here,  $\partial\tilde{\Omega}$  is of finite type at every point of the form  $(z, 0)$ . Then, by Catlin's Theorem ([Cat87]), the results of [KN65] show that the Neumann operator of  $\tilde{\Omega}$  is pseudolocal at these points, and, the method introduced by N. Kerzman in [Ker72] proves that the restriction of the Bergman kernel of  $\tilde{\Omega}$  to  $(\bar{\Omega} \times \{0\})^2$  is  $\mathcal{C}^\infty$  outside the diagonal of  $(\partial\Omega \times \{0\})^2$ . Thus, the identity  $K_\omega^\Omega(p_1 p_2) = K^{\tilde{\Omega}}((p_1, 0), (p_2, 0))$  implies that the estimates of Theorems 4.1 and 4.2 are valid everywhere.

These estimates, the hypothesis on  $h$  (i.e.  $h(w) \asymp \sum |w_i|^{2q_i}$ ), an immediate generalization of Proposition 2.1 of [BCG96] and a standard application of Hölder inequality imply that  $P_\omega^\Omega$  maps continuously  $L^p(\Omega, (-\rho)^\alpha d\lambda)$  into itself for  $-1 < \alpha < p \left(1 + \sum \frac{1}{q_i}\right) - 1$ .

The Lipschitz estimate is also standard.

Now, choosing the special function  $h(w) = \sum |w_i|^{2q_i}$ ,  $w_i \in \mathbb{C}$ , the weight  $\omega$  is equal to  $C(-\rho)^{\sum 1/q_i}$ , and Theorem 1.1 follows the Corollary of Theorem 1.2.

*Remark.* Note that same method gives trivially Theorem 1.1 for pseudo-convex decoupled domains of finite type in  $\mathbb{C}^n$ .

## REFERENCES

- [BC00] Bo Berndtsson and Philippe Charpentier, *A Sobolev mapping property of the Bergman kernel*, Math. Z. **235** (2000), no. 1, 1–10.
- [BCG96] A. Bonami, D. Chang, and S. Grellier, *Commutation properties and Lipschitz estimates for the Bergman and Szegő projections*, Math. Z. **223** (1996), 275–302.
- [BG95] A. Bonami and S. Grellier, *Weighted Bergman projections in domains of finite type in  $\mathbb{C}^2$* , Contemp. Math. **189** (1995), 65–80.
- [BS91] Harold P. Boas and Emil J. Straube, *Sobolev estimates for the  $\bar{\partial}$ -Neumann operator on domains in  $\mathbb{C}^n$  admitting a defining function that is plurisubharmonic on the boundary*, Math. Z. **206** (1991), no. 1, 81–88.
- [Cat87] David Catlin, *Subelliptic estimates for the  $\bar{\partial}$ -Neumann problem on pseudoconvex domains*, Ann. of Math. **126** (1987), no. 1, 131–191.
- [CD06a] P. Charpentier and Y. Dupain, *Estimates for the Bergman and Szegő projections for pseudo-convex domains of finite type with locally diagonalizable Levi form*, Publications Mathématiques **50** (2006), no. 2, 413–446.
- [CD06b] ———, *Geometry of Pseudo-convex Domains of Finite Type with Locally Diagonalizable Levi Form and Bergman Kernel*, Jour. Math. Pures et Appl. **85** (2006), 71–118.
- [CD08] ———, *Extremal Basis, Geometrically Separated Domains and Applications*, <http://fr.arxiv.org/abs/0810.1884> (2008).
- [Cha80] P. Charpentier, *Solutions minimales de l'équation  $\bar{\partial}u = f$  dans la boule et le polydisque*, Ann. Inst. Fourier **30** (1980), 121–153.
- [Cho96] S. Cho, *Estimates of the Bergman kernel function on certain pseudoconvex domains in  $\mathbb{C}^n$* , Math Z. **222** (1996), 329–339.
- [Cho02] ———, *Estimates of the Bergman kernel function on pseudoconvex domains with comparable Levi form*, J. Korean Math. Soc. **39** (2002), no. 3, 425–437.
- [CL97] Chang D. C. and Li B. Q., *Sobolev and Lipschitz estimates for weighted Bergman projections*, Nagoya Math. J. **147** (1997), 147–178.
- [CNS92] D. C. Chang, A. Nagel, and E. Stein, *Estimates for the  $\bar{\partial}$ -Neumann problem for pseudoconvex domains in  $\mathbb{C}^2$  of finite type*, Acta Math. **169** (1992), 153–228.
- [Cum90] A. Cumenge, *Estimations limites pour la solution canonique de l'équation  $\bar{\partial}u = f$* , Math. Ann. **286** (1990), no. 4, 639–654.
- [DF77a] K. Diederich and J. E. Fornaess, *Pseudoconvex Domains : Bounded Strictly Plurisubharmonic Exhaustion Functions*, Inventiones math. **39** (1977), 129–141.
- [DF77b] ———, *Pseudoconvex domains: An example with nontrivial nebenhuelle*, Math. Ann. **225** (1977), 275–292.
- [Hef04] T. Hefer, *Extremal bases and Hölder Estimates for  $\bar{\partial}$  on Convex Domains of Finite Type*, Michigan Math. J. **52** (2004), 573–602.
- [Hör65] L. Hörmander,  *$L^2$  estimates and existence theorems for the  $\bar{\partial}$ -operator*, Acta Math. **113** (1965), 89–152.
- [HP84a] R. Harvey and J. Polking, *The  $\bar{\partial}$ -Neumann kernel in the ball in  $\mathbb{C}^n$* , Proc. Sympos. Pure Math. **41** (1984), 117–136.
- [HP84b] ———, *The  $\bar{\partial}$ -Neumann solution to the inhomogeneous Cauchy-Riemann equation in the ball in  $\mathbb{C}^n$* , Trans. Amer. Math. Soc. **281** (1984), no. 2, 587–613.
- [Ker72] N. Kerzman, *The Bergman kernel function. Differentiability at the boundary*, Math. Ann. **195** (1972), 149–158.



- [KN65] J. J. Kohn and L. Nirenberg, *Non coercive boundary value problems*, Comm. Pure Appl. Math. **18** (1965), 443–492.
- [Lig89] E. Ligočka, *On the Forelli-Rudin construction and weighted Bergman projections*, Studia Math. **94** (1989), no. 3, 257–272.
- [LR86] I. Lieb and M. Range, *Integral representations and estimates in the theory of the  $\bar{\partial}$ -Neumann problem.*, Ann. of Math. **123** (1986), no. 2, 265–301.
- [LR87] ———, *The kernel of the  $\bar{\partial}$ -Neumann operator on strictly pseudoconvex domains*, Math. Ann. **278** (1987), no. 1–4, 151–173.
- [LR88] ———, *The  $\bar{\partial}$ -Neumann kernel for codimension-one forms on strictly pseudoconvex domains*, Math. Notes, Princeton Univ. Press **38** (1988), 473–782.
- [McN94] J. McNeal, *Estimates on Bergman Kernels of Convex Domains*, Advances in Math (1994), 108–139.
- [MS94] J. D. McNeal and E. M. Stein, *Mapping properties of the Bergman projection on convex domains of finite type*, Duke Math. J. **73** (1994), no. 1, 177–199.
- [MS97] ———, *The Szegő projection on convex domains*, Math. Z. **224** (1997), no. 4, 519–553.
- [NRSW89] A. Nagel, J. P. Rosay, E. M. Stein, and S. Wainger, *Estimates for the Bergman and Szegő kernels in  $\mathbb{C}^2$* , Annals of Math. **129** (1989), 113–147.
- [PW90] Z. Pasternak-Winiarski, *On the Dependence of the Reproducing Kernel on the Weight of Integration*, Jour. Func. Anal. **94** (1990), 110–134.
- [Str10] E. Straube, *Lectures on the  $L^2$  Sobolev theory on the -Neumann problem*, European Mathematical Society, 2010, 2010.
- [Zeya] Y. Zeytuncu,  *$L^p$  regularity of some weighted Bergman projections on the unit disc*, Turk. J. Math. To appear.
- [Zeyb] ———,  *$L^p$  Regularity of weighted Bergman Projections*, Tran. of Amer. Math. Soc. To appear.
- [Zey11a] ———, *Sobolev regularity of weighted Bergman projections on the unit disc*, Complex Variables and Elliptic Equations (2011).
- [Zey11b] ———, *Weighted Bergman projections and kernels:  $L^p$  regularity and zeros*, Proc. Amer. Math. Soc. **139** (2011), 2105–2112.

P. CHARPENTIER & Y. DUPAIN, UNIVERSITÉ BORDEAUX I, INSTITUT DE MATHÉMATIQUES DE BORDEAUX, 351, COURS DE LA LIBÉRATION, 33405, TALENCE, FRANCE

M. MOUNKAILA, UNIVERSITÉ ABDOU MOUMOUNI, FACULTÉ DES SCIENCES, B.P. 10662, NIAMEY, NIGER

*E-mail address:* P. Charpentier: philippe.charpentier@math.u-bordeaux1.fr

*E-mail address:* Y. Dupain: yves.dupain@math.u-bordeaux1.fr

*E-mail address:* M. Mounkaila: modi.mounkaila@yahoo.fr