HAL open science

Optimal estimate of the spectral gap for the degenerate Goldstein-Taylor model

Etienne Bernard, Francesco Salvarani

▶ To cite this version:

Etienne Bernard, Francesco Salvarani. Optimal estimate of the spectral gap for the degenerate Goldstein-Taylor model. 2012. hal-00766211v1

HAL Id: hal-00766211

https://hal.science/hal-00766211v1

Preprint submitted on 18 Dec 2012 (v1), last revised 3 May 2013 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

OPTIMAL ESTIMATE OF THE SPECTRAL GAP FOR THE DEGENERATE GOLDSTEIN-TAYLOR MODEL

ÉTIENNE BERNARD AND FRANCESCO SALVARANI

Abstract. In this paper we study the decay to the equilibrium state for the solution of a generalized version of the Goldstain-Taylor system, posed in the one-dimensional torus $\mathbb{T} = \mathbb{R}/\mathbb{Z}$, by allowing that the non-negative cross section σ can vanish in a subregion $X := \{x \in \mathbb{T} \,|\, \sigma(x) = 0\}$ of the domain with meas $(X) \geq 0$ with respect to the Lebesgue measure.

We prove that the solution converges in time, with respect to the strong L^2-topology, to its unique equilibrium with an exponential rate whenever meas $(\mathbb{T} \setminus X) \geq 0$ and we give an optimal estimate of the spectral gap.

1. Introduction

The investigation about explicit rates of approach to equilibrium in large time, for kinetic equations, is an active field of research and many results have been obtained, both in the linear and in the non linear case.

An important concept, in this context, is *hypocoercivity*. This property appears in many evolution equations which have a conservative part and a dissipative one. Even if the conservative part alone does not induce relaxation and the dissipative one is not sufficient to induce convergence to equilibrium, sometimes the combination of the two parts leads to relaxation. When this situation occurs, the equation is said to be *hypocoercive*.

For kinetic equations, the conservative term is usually the free transport operator, which mixes the space and the velocity variables, whereas the dissipative part is a collision operator, whose null space does not depend on the space variable.

Furthermore, the key ingredient of many proofs is based upon the independence of the null space of the dissipative operator on the space variable. This allows indeed a local control of the dissipative properties of the equation, and hence the solution is locally "attracted" everywhere toward its local equilibrium (see, for instance, [9, 5]).

The situation is, however, quite different in the degenerate case, i.e. when the collision operator can vanish in the spatial domain of the problem (even if the degeneracy happens only at isolated points). In the region of degeneracy, indeed, the null space of the collision operator becomes trivial.

This problem has been studied for the first time by Desvillettes and the second author in [4]. In this article, they proved, under very stringent hypotheses on the degeneracy of the cross section, that the solution of the linear Boltzmann equation and the solution of a reduced two-velocity model, namely the generalized Goldstein-Taylor system, converge in time to their equilibrium with (at least) polynomial speed. They also conjectured that some explicit rate should still exist even in more general degenerate situations.

The deep reason of this conjecture is based on the fact that the hypocoercivity properties, which are locally lost in the regions of degeneracy, can be recovered

Date: December 17, 2012.

Key words and phrases. Goldstein-Taylor model, Spectral gap, Degenerate cross section.

at a global level, through the action of the mixing term, here the free transport operator.

However, at least in more than one space dimension, the decay rate to equilibrium cannot be – in general – exponential, as shown by a counterexample of the authors in [2].

Indeed, we proved that the linear Boltzmann equation on the torus $\mathbb{T}^d$, $d \geq 2$, with velocities on the sphere $\mathbb{S}^{d-1}$, has, for a wide class of cross sections in L^∞, a L^2-distance to equilibrium that cannot decay faster than $t^{-1/2}$.

The exponential rate can be recovered only by assuming an additional hypothesis on the geometrical structure of the cross section.

This additional requirement has been introduced by the authors in [3] and has been called the *geometrical condition*, as a reminiscence of the Bardos-Lebeau-Rauch condition that guarantees the exponential stabilization of the damped wave equation [1]. We proved that the geometrical condition is necessary and sufficient to recover the exponential decay in time to equilibrium. However, our proof in [3] is not constructive, since it is based on a compactness argument.

Hence, the problem of finding an explicit exponential rate for the linear Boltzmann equation, with cross sections satisfying the geometrical condition, is still an open problem. Likewise, finding the best convergence rate to equilibrium in the general case is still open.

The aim of this article is to go beyond the actual state-of-the-art and give a quantitative study of the spectral gap in the degenerate case. Here we will restrict ourselves to consider the simplest possible degenerate kinetic equation.

We prove that the convergence to equilibrium for the generalized Goldstein-Taylor system proposed in [4] is exponential in time and, moreover, we can exactly characterize the convergence rate. The geometrical condition of [3] is automatically fulfilled here, since the spatial domain is one-dimensional. Consequently, our result agrees with the general theory.

The proof is based on the equivalent formulation of the problem in terms of the damped telegrapher's equation, by following a trick by Kac [7], and on a careful estimate of the time decay of the energy for the non-homogeneous telegrapher's equation, proved by Rauch and Taylor in [11].

The question of obtaining similar results when the velocities are not discrete has not been treated here, as it requires different techniques. This fact is a consequence of the non-closure of the moment equations when the number of admissible velocities of the system is infinite.

The paper is organized as follows: in Section 2 we state precisely our problem and then, in Section 3, we prove our main theorem about the long-time behavior of the solution of the problem.

2. The problem and its basic properties

We consider a simplified one-dimensional version of the linear Boltzmann equation, in which the velocity set is discrete and reduced only to two equal and opposite velocities.

It is a variant of the well-known Goldstein-Taylor model [6, 12]. It describes the behavior of a gas composed of two kinds of particles moving parallel to the x-axis with constant speeds, of equal modulus $c = 1$, the first one in the positive x-direction with density $u(x,t)$, the other one in the negative x-direction with density $v(x,t)$, that experience switches in velocity distributed under a Poisson law, weighted by the cross section σ, which takes into account possible anisotropy

effects. The corresponding system of equations is:

$$\begin{cases} \dfrac{\partial u}{\partial t} + \dfrac{\partial u}{\partial x} = \sigma(x)(v-u) \\ \\ \dfrac{\partial v}{\partial t} - \dfrac{\partial v}{\partial x} = \sigma(x)(u-v), \end{cases} \tag{2.1}$$

where $u := u(t,x)$, $v := v(t,x)$, $x \in \mathbb{T} = \mathbb{R}/\mathbb{Z}$, $t \geq 0$.

Such set of equations will satisfy the initial conditions

$$u(0,x) = u^{\text{in}}(x), \qquad v(0,x) = v^{\text{in}}(x). \tag{2.2}$$

We will suppose henceforth that $(u^{\text{in}}, v^{\text{in}}) \in H^1(\mathbb{T}) \times H^1(\mathbb{T})$, and that

$$\sigma \in L^\infty(\mathbb{T}), \text{ with } \sigma \geq 0 \text{ a.e. and } \int_{\mathbb{T}} \sigma(x)\,\mathrm{d}x > 0.$$

We now state our main result:

Theorem 2.1. *Let $(u^{\text{in}}, v^{\text{in}}) \in H^1(\mathbb{T}) \times H^1(\mathbb{T})$, and $\sigma \in L^\infty(\mathbb{T})$ be non-negative functions. Suppose moreover that*

$$\int_{\mathbb{T}} \sigma(x)\,\mathrm{d}x > 0.$$

Denote also

$$u_\infty := \frac{1}{2}\int_{\mathbb{T}} (u^{\text{in}} + v^{\text{in}})\,\mathrm{d}x.$$

Then, there exists an explicit positive constant A_* depending on $\|u^{\text{in}}\|_{L^2(\mathbb{T})}$, $\|v^{\text{in}}\|_{L^2(\mathbb{T})}$ and σ such that

$$\|u - u_\infty\|^2_{L^2(\mathbb{T})} + \|v - u_\infty\|^2_{L^2(\mathbb{T})} \leq A_* \exp\left(-\|\sigma\|_{L^1(\mathbb{T})} t\right).$$

Moreover, the decay rate is optimal.

3. A Lyapunov functional of the system

We underline that the results presented below need different regularity requirements.

For this reason, we will write explicitly the hypotheses on the regularity of the initial conditions and of the cross section in the statement of the corresponding theorems.

3.1. A priori estimates.

In order to make the paper self-consistent, we summarize here some basic properties of the generalized Goldstein-Taylor system (2.1)-(2.2). The first result concerns the well-posedness of the problem.

Proposition 3.1. *Consider the Goldstein-Taylor model* (2.1)-(2.2) *with non negative initial data $(u^{\text{in}}, v^{\text{in}}) \in L^1(\mathbb{T}) \times L^1(\mathbb{T})$ and with cross section $\sigma \in L^\infty(\mathbb{T})$ such that $\sigma \geq 0$ a.e.. Then there exists a unique (generalized) non negative solution (u,v) to this system in $C(\mathbb{R}^+; L^1(\mathbb{T}) \times L^1(\mathbb{T}))$.*

Proof. The result is immediate since the collision term can be treated as a bounded perturbation of the transport operator [10]. □

We then give some a-priori estimates of the Cauchy problem (2.1)-(2.2). The following lemma holds:

Lemma 3.2. *Let (u, v) be the solution of the generalized Goldstein-Taylor system* (2.1)-(2.2)*, with non-negative initial data $(u^{\text{in}}, v^{\text{in}}) \in L^\infty(\mathbb{T}) \times L^\infty(\mathbb{T})$ and with non-negative cross section $\sigma \in L^\infty(\mathbb{T})$. Then, for any smooth convex function $\varphi(r)$, $r \geq 0$, we have*

$$\int_{\mathbb{T}} [\varphi(u(t,x)) + \varphi(v(t,x))] \, \mathrm{d}x \leq \int_{\mathbb{T}} \left[\varphi(u^{\text{in}}(x)) + \varphi(v^{\text{in}}(x))\right] \, \mathrm{d}x. \tag{3.1}$$

In particular, the conservation of mass and the maximum principle hold, i.e.

$$\|u(t,\,\cdot\,) + v(t,\,\cdot\,)\|_{L^1(\mathbb{T})} = \|u^{\text{in}} + v^{\text{in}}\|_{L^1(\mathbb{T})}$$

and

$$\max\{\|u(t,\,\cdot\,)\|_{L^\infty(\mathbb{T})}\,,\, \|v(t,\,\cdot\,)\|_{L^\infty(\mathbb{T})}\} \leq \|u^{\text{in}} + v^{\text{in}}\|_{L^\infty(\mathbb{T})}.$$

Proof. We apply the strategy of [8]. Let $\varphi(r)$, $r \geq 0$, a smooth convex function. We multiply the first equation of (2.1) by $\varphi'(u)$ and the second one by $\varphi'(v)$. We integrate on $\mathbb{T}$ and, by summing the two equations, we obtain

$$\frac{d}{dt} \int_{\mathbb{T}} \varphi(u) \, \mathrm{d}x = - \int_{\mathbb{T}} \sigma(x)(u - v) \left[\varphi'(u) - \varphi'(v)\right] \, \mathrm{d}x.$$

Since $\varphi \in C^2(\Omega)$ and is a convex function, then φ' is monotone. Hence

$$\sigma(x)(u - v)[\varphi'(u) - \varphi'(v)] \geq 0$$

because σ is non-negative and then, for all $t > 0$,

$$\int_{\mathbb{T}} [\varphi(u(t,x)) + \varphi(v(t,x))] \, \mathrm{d}x \leq \int_{\mathbb{T}} \left[\varphi(u^{\text{in}}(x)) + \varphi(v^{\text{in}}(x))\right] \, \mathrm{d}x.$$

By choosing $\varphi(r) = r$, we obtain the conservation of mass:

$$\int_{\mathbb{T}} [u(t,x) + v(t,x)] \, \mathrm{d}x = \int_{\mathbb{T}} \left[u^{\text{in}}(x) + v^{\text{in}}(x)\right] \, \mathrm{d}x. \tag{3.2}$$

By taking $\varphi(r) = r^p$ for all $p \geq 1$, we obtain the boundedness of any L^p-norm

$$\int_{\mathbb{T}} [u(t,x)^p + v(t,x)^p] \, \mathrm{d}x \leq \int_{\mathbb{T}} \left[u^{\text{in}}(x)^p + v^{\text{in}}(x)^p\right] \, \mathrm{d}x.$$

Moreover, when $p \to +\infty$, the lemma implies also the L^∞ bound:

$$0 < u(t,x), v(t,x) \leq \|u^{\text{in}} + v^{\text{in}}\|_{L^\infty(\mathbb{T})}. \tag{3.3}$$

□

We give here a result, whose proof is similar to Proposition 3.1 of [4], showing that the H^1-regularity of the solution of the problem (2.1)-(2.2) is preserved in time.

Proposition 3.3. *Let $(u^{\text{in}}, v^{\text{in}}) \in H^1(\mathbb{T}) \times H^1(\mathbb{T})$. Then, there exists a constant γ (depending explicitly on $||u^{\text{in}}||_{H^1}$, $||v^{\text{in}}||_{H^1}$ and $||\sigma||_{L^\infty}$) such that the solution (u, v) of system* (2.1)-(2.2) *satisfies the bound*

$$\sup_{t \geq 0} \int_{\mathbb{T}} \left((\partial_t u)^2 + (\partial_x u)^2 \right) \mathrm{d}x \leq \gamma,$$

$$\sup_{t \geq 0} \int_{\mathbb{T}} \left((\partial_t v)^2 + (\partial_x v)^2 \right) \mathrm{d}x \leq \gamma.$$

Proof of Proposition 3.3: We differentiate the equations of system (2.1) with respect to the variable t, and multiply the result by $2\partial u/\partial t$ and $2\partial v/\partial t$ respectively. After integrating with respect to $x \in \mathbb{T}$, we end up with

$$\frac{d}{dt}\int_{\mathbb{T}}\left[\left(\frac{\partial u}{\partial t}\right)^2+\left(\frac{\partial v}{\partial t}\right)^2\right]dx=-2\int_{\mathbb{T}}\sigma(x)\left[\frac{\partial u}{\partial t}-\frac{\partial v}{\partial t}\right]^2\mathrm{d}x\leq 0. \tag{3.4}$$

We finally observe that

$$\partial_x u=-\partial_t u+\sigma(v-u),\qquad \partial_x v=\partial_t v+\sigma(v-u).$$

Hence, using the bound above and the fact that $\sigma \in L^\infty$, we can conclude the proof of Proposition 3.3.

3.2. **A damped wave equation.** We define the *macroscopic density*

$$\rho(t,x):=u(t,x)+v(t,v),\qquad t>0, x\in\mathbb{T},$$

and the *flux*

$$j(t,x):=u(t,x)-v(t,v),\qquad t>0, x\in\mathbb{T}.$$

From (2.1), it is easy to show that (ρ, j) verifies

$$\begin{cases}\partial_t\rho+\partial_x j=0,\\ \\ \partial_t j+\partial_x\rho=-2\sigma j,\end{cases}\qquad (t,x)\in\mathbb{R}^+\times\mathbb{T} \tag{3.5}$$

with initial conditions $\rho(0,x)=u^{\mathrm{in}}(x)+v^{\mathrm{in}}(x)$ and $j(0,x)=u^{\mathrm{in}}(x)-v^{\mathrm{in}}(x)$.

By using a trick that comes back to Kac, it can be shown that j is the solution of a damped waves equation.

We have indeed the following result [7].

Proposition 3.4. *Let $j=u-v$ be the flux corresponding to the Goldstein-Taylor system* (2.1)-(2.2)*. Then j is the solution of the damped telegrapher's equation*

$$\begin{cases}\partial^2_{tt}j-\partial^2_{xx}j+2\sigma\partial_t j=0,\\ \\ j(0,x)=u^{\mathrm{in}}(x)-v^{\mathrm{in}}(x),\\ \\ \partial_t j(0,x)=2\sigma(x)[v^{\mathrm{in}}(x)-u^{\mathrm{in}}(x)]-\partial_x u^{\mathrm{in}}(x)-\partial_x v^{\mathrm{in}}(x),\end{cases}\qquad (t,x)\in\mathbb{R}^+\times\mathbb{T}. \tag{3.6}$$

In [11], the exponential decay in time to zero of the energy

$$E(t):=\|\partial_t j(t,\,\cdot\,)\|_{L^2(\mathbb{T})}+\|\partial_x j(t,\,\cdot\,)\|_{L^2(\mathbb{T})} \tag{3.7}$$

of the damped telegrapher's equation (3.6) has been proved, and explicit estimates have been provided on the decay rate. More precisely, the following result holds:

Proposition 3.5. *Let j be the solution of the damped telegrapher equation* (3.6)*, posed in the periodic torus $\mathbb{T}$.*

Then, there exist two explicit positive constants c and α such that

$$E(t)=\int_{\mathbb{T}}[(j_t)^2+(j_x)^2]\,\mathrm{d}x\leq ce^{-\alpha t}.$$

In particular,

$$\alpha=\int_{\mathbb{T}}\sigma(x)\,\mathrm{d}x.$$

Remark 3.6. *The original formulation of the previous proposition has been written in terms of a Fourier expansion (see* [11]*, Section 2), but some straightforward computations show that we can deduce our formulation from the original one.*

We now prove the main result of this article, namely the exponential decay in time of the solution (u, v) of the generalized Goldstein-Taylor system (2.1)-(2.2) to the stationary state (u_∞, u_∞).

The proof is based on the fact that the quantity

$$H(t) := \int_{\mathbb{T}} [(u - u_\infty)^2 + (v - u_\infty)^2] \, \mathrm{d}x$$

is a Lyapunov functional of the system and is controlled by the energy defined in (3.7). A quantitative estimate on the long-time decay of H will give directly the asymptotic behavior of the L^2 distance between the solution of (2.1)-(2.2) and the stationary state.

The proof will be performed in several steps. The first one concerns a bound on the weighted L^2-norm of the flux.

Lemma 3.7. *There exists a constant $B \geq 0$ such that*

$$\left\| j\sqrt{\sigma} \right\|^2_{L^2(\mathbb{T})} \leq BE(t).$$

Moreover,

$$B = \left(\|u^{\mathrm{in}}\|_{L^2(\mathbb{T})} + \|v^{\mathrm{in}}\|_{L^2(\mathbb{T})} \right).$$

Proof. We recall that

$$\partial_t j + \partial_x \rho = -2\sigma j.$$

Multiplying both sides of the equality above by j and integrating in $x \in \mathbb{T}$, we obtain immediately that

$$\int_{\mathbb{T}} j\partial_t j \, \mathrm{d}x - \int_{\mathbb{T}} \rho \partial_x j \, \mathrm{d}x = -2 \int_{\mathbb{T}} \sigma j^2 \, \mathrm{d}x.$$

The previous equation implies that

$$\int_{\mathbb{T}} 2\sigma j^2 \, \mathrm{d}x \leq \left| \int_{\mathbb{T}} j\partial_t j \, \mathrm{d}x \right| + \left| \int_{\mathbb{T}} \rho \partial_x j \, \mathrm{d}x \right|.$$

By Cauchy-Schwarz inequality, the inequality above leads to

$$2 \int_{\mathbb{T}} \sigma j^2 \, \mathrm{d}x \leq \|\partial_t j\|_{L^2(\mathbb{T})} \|j\|_{L^2(\mathbb{T})} + \|\rho\|_{L^2(\mathbb{T})} \|\partial_x j\|_{L^2(\mathbb{T})}$$

or

$$2 \int_{\mathbb{T}} \sigma j^2 \, \mathrm{d}x \leq \left(\|j\|_{L^2(\mathbb{T})} + \|\rho\|_{L^2(\mathbb{T})} \right) E(t).$$

By Lemma 3.2,

$$\int_{\mathbb{T}} \sigma j^2 \, \mathrm{d}x \leq BE(t)$$

with

$$B := \left(\left\|u^{\mathrm{in}}\right\|_{L^2(\mathbb{T})} + \left\|v^{\mathrm{in}}\right\|_{L^2(\mathbb{T})} \right). \tag{3.8}$$

This ends the proof of the lemma. □

The next result gives a bound of the full L^2-norm of the flux j, which will be evaluated in terms of the energy E.

Lemma 3.8. *There exists a constant $C > 0$ such that*

$$\|j\|^2_{L^2(\mathbb{T})} \leq CE(t).$$

Moreover we have

$$C = \frac{2}{\|\sqrt{\sigma}\|^2_{L^1(\mathbb{T})}} \left[B + \frac{1}{\pi^2} \left(\|\sqrt{\sigma}\|^2_{L^1(\mathbb{T})} + \|\sigma\|_{L^1(\mathbb{T})} \right) \right].$$

Proof. First, we introduce the notation

$$\beta := \int_{\mathbb{T}} \sqrt{\sigma}(x)dx.$$

Notice that $\sqrt{\sigma}/\beta$ is a unit measure on $\mathbb{T}$. Keeping that in mind, we have

$$\begin{aligned} \|j\|^2_{L^2(\mathbb{T})} &= \int_{\mathbb{T}} (j)^2 \, \mathrm{d}x \\ &= \int_{\mathbb{T}} \left(j - \frac{1}{\beta} \int_{\mathbb{T}} j\sqrt{\sigma} \, \mathrm{d}x + \frac{1}{\beta} \int_{\mathbb{T}} j\sqrt{\sigma} \, \mathrm{d}x \right)^2 \mathrm{d}x \\ &\leq 2 \int_{\mathbb{T}} \left(j - \frac{1}{\beta} \int_{\mathbb{T}} j\sqrt{\sigma} \, \mathrm{d}x \right)^2 \mathrm{d}x + \frac{2}{\beta^2} \left(\int_{\mathbb{T}} j\sqrt{\sigma} \, \mathrm{d}x \right)^2 . \end{aligned} \tag{3.9}$$

By Jensen inequality and Lemma 3.7, we have

$$\begin{aligned} \frac{2}{\beta^2} \left(\int_{\mathbb{T}} j\sqrt{\sigma} \, \mathrm{d}x \right)^2 &\leq \frac{2}{\beta^2} \int_{\mathbb{T}} j^2 \sigma \, \mathrm{d}x \\ &= \frac{2}{\beta^2} \left\| j\sqrt{\sigma} \right\|^2_{L^2(\mathbb{T})} \\ &\leq \frac{2B}{\beta^2} E(t) \end{aligned} \tag{3.10}$$

where B is defined in (3.8).

Since $\sqrt{\sigma}/\beta$ is a unit measure on $\mathbb{T}$, the first term in the right-hand of the inequality gives

$$\begin{aligned} \left(j - \frac{1}{\beta} \int_{\mathbb{T}} j\sqrt{\sigma} \, \mathrm{d}x \right)^2 &= \left(j - \int_{\mathbb{T}} j(x) \, \mathrm{d}x - \frac{1}{\beta} \int_{\mathbb{T}} \sqrt{\sigma(x)} \left(j - \int_{\mathbb{T}} j(x) \, \mathrm{d}x \right) \mathrm{d}x \right)^2 \\ &\leq 2 \left(j - \int_{\mathbb{T}} j(x) \, \mathrm{d}x \right)^2 + 2 \frac{1}{\beta^2} \left(\int_{\mathbb{T}} \sqrt{\sigma(x)} \left(j - \int_{\mathbb{T}} j(x) \, \mathrm{d}x \right) \mathrm{d}x \right)^2 . \end{aligned}$$

Thus, by Cauchy-Schwartz inequality and Poincaré-Wirtinger inequality,

$$2 \int_{\mathbb{T}} \left(j - \frac{1}{\beta} \int_{\mathbb{T}} j\sqrt{\sigma} \, \mathrm{d}x \right)^2 \mathrm{d}x \leq \frac{2}{\pi^2} \left(1 + \frac{1}{\beta^2} \|\sigma\|_{L^1(\mathbb{T})} \right) E(t). \tag{3.11}$$

Hence we obtain the result by applying inequalities (3.11) and (3.10) in inequality (3.9). □

Our last preliminary result is the control of the L^2-norm of the spatial derivative of the macroscopic density ρ in terms of the energy E.

Lemma 3.9. *There exists a constant $K > 0$ such that*

$$\|\partial_x \rho\|^2_{L^2(\mathbb{T})} \leq K E(t).$$

with

$$K := 1 + 2 \|\sigma\|^2_{L^\infty(\mathbb{T})} C,$$

where C is defined in Lemma 3.8.

Proof. We recall that

$$\partial_t j + \partial_x \rho = -2\sigma j.$$

Therefore we have:

$$|\partial_x \rho|^2 \leq |\partial_t j|^2 + 2 \|\sigma\|^2_{L^\infty(\mathbb{T})} |j|^2 .$$

Integrating both sides of the inequality above, we obtain

$$\|\partial_x \rho\|^2_{L^2(\mathbb{T})} \leq \|\partial_t j\|^2_{L^2(\mathbb{T})} + 2 \|\sigma\|^2_{L^\infty(\mathbb{T})} \|j\|^2_{L^2(\mathbb{T})}$$

thus, by Lemma 3.8, there exists $K > 0$ such that

$$\|\partial_x \rho\|^2_{L^2(\mathbb{T})} \leq KE(t).$$

□

4. Proof of the main result

We are now ready to prove Theorem 2.1.

Proof. Assume that $u^{\text{in}}, v^{\text{in}} \in H^1(\mathbb{T})$. Since u_∞ is constant and nonnegative, a direct computation shows that

$$H(t) = \frac{1}{2}\left(\|\rho - 2u_\infty\|^2_{L^2} + \|j\|^2_{L^2}\right). \tag{4.1}$$

By the conservation of the mass, given in Lemma 3.2, for each $t > 0$ we have

$$\int_{\mathbb{T}} \rho(t,x)\,\mathrm{d}x = 2u_\infty.$$

Poincaré-Wirtinger's inequality hence assures that

$$\|\rho - 2u_\infty\|^2_{L^2} \leq \|\partial_x \rho\|^2_{L^2}.$$

Using the inequality above in equality (4.1) implies

$$H(t) \leq \frac{1}{2}\left(\|\partial_x \rho\|^2_{L^2} + \|j\|^2_{L^2}\right).$$

Consequently by Lemma 3.8 and Lemma 3.9 there exists a constant $A > 0$ such that

$$H(t) \leq AE(t).$$

Thee constant A can be explicitly computed. Indeed

$$A := C + K,$$

where C and K are defined in Lemma 3.8 and Lemma 3.9. Proposition 3.5 allows to conclude that the inequality in Theorem 2.1 holds for every solution with u^{in}, $v^{\text{in}} \in H^1(\mathbb{T})$.

Finally, since the cross section σ is non negative by hypotheses, and Proposition 3.1 guarantees the non negativity of u and v, then Duhamel's formula implies the pair of inequalities

$$u(t,x) \geq u^{\text{in}}(x-t)\exp\left(-\int_0^t \sigma(x-s)\,\mathrm{d}s\right), \qquad \text{for all } (t,x) \in \mathbb{R}^+ \times \mathbb{T}$$

$$v(t,x) \geq v^{\text{in}}(x+t)\exp\left(-\int_0^t \sigma(x+s)\,\mathrm{d}s\right), \qquad \text{for all } (t,x) \in \mathbb{R}^+ \times \mathbb{T}.$$

This show the optimality of the decay rate, and hence the complete characterization of the spectral gap.

□

5. Conclusion

In this paper, we completely characterize the spectral gap of the degenerate Goldstein-Taylor system, with general degenerate cross section, in terms of the geometrical properties of the cross section itself.

Because of the peculiar situation of the one-dimensional Euclidean space, all uniformly bounded cross sections that differ from the null function (in L^1 sense) satisfy the geometrical condition, and hence generate hypocoercive effects in agreement with our general result in [3].

Our result is optimal, and provides the definitive answer to the estimation of the spectral gap of the Goldstein-Taylor model.

Part of the proof depends heavily on the fact that the set of admissible velocities is discrete.

In the case of a continuous set of velocities, a similar result needs new techniques of proof, since the set of the moment equations is no longer closed.

Acknowledgements. This paper has been partially supported by the Italian national institute of higher mathematics (INDAM), GNFM project "Study of complex kinetic systems: theoretical analysis and numerical simulation".

References

[1] Claude Bardos, Gilles Lebeau, and Jeffrey Rauch. Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary. *SIAM J. Control Optim.*, 30(5):1024–1065, 1992.

[2] Étienne Bernard and Francesco Salvarani. On the convergence to equilibrium for degenerate transport problems. *Arch. Rational Mech. Anal.*, 2012. In press.

[3] Étienne Bernard and Francesco Salvarani. On the exponential decay to equilibrium of the degenerate linear boltzmann equation, 2012.

[4] Laurent Desvillettes and Francesco Salvarani. Asymptotic behavior of degenerate linear transport equations. *Bull. Sci. Math.*, 133(8):848–858, 2009.

[5] Jean Dolbeault, Clément Mouhot, and Christian Schmeiser. Hypocoercivity for kinetic equations with linear relaxation terms. *C. R. Math. Acad. Sci. Paris*, 347(9-10):511–516, 2009.

[6] S. Goldstein. On diffusion by discontinuous movements, and on the telegraph equation. *Quart. J. Mech. Appl. Math.*, 4:129–156, 1951.

[7] Mark Kac. A stochastic model related to the telegrapher's equation. *Rocky Mountain J. Math.*, 4:497–509, 1974. Reprinting of an article published in 1956, Papers arising from a Conference on Stochastic Differential Equations (Univ. Alberta, Edmonton, Alta., 1972).

[8] Pierre Louis Lions and Giuseppe Toscani. Diffusive limit for finite velocity Boltzmann kinetic models. *Rev. Mat. Iberoamericana*, 13(3):473–513, 1997.

[9] Clément Mouhot and Lukas Neumann. Quantitative perturbative study of convergence to equilibrium for collisional kinetic models in the torus. *Nonlinearity*, 19(4):969–998, 2006.

[10] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1983.

[11] Jeffrey Rauch and Michael Taylor. Decay of solutions to nondissipative hyperbolic systems on compact manifolds. *Comm. Pure Appl. Math.*, 28(4):501–523, 1975.

[12] G. I. Taylor. Diffusion by Continuous Movements. *Proc. London Math. Soc.*, S2-20(1):196.

(É. B.) Institut Géographique National, Laboratoire de Recherche en Géodésie, Université Paris Diderot, Bâtiment Lamarck A, 5, rue Thomas Mann, Case courrier 7071, 75205 Paris Cedex 13, France

E-mail address: `etienne.bernard@ign.fr`

(F. S.) Dipartimento di Matematica F. Casorati, Università degli Studi di Pavia, Via Ferrata 1, 27100 Pavia, Italy

E-mail address: `francesco.salvarani@unipv.it`