



HAL
open science

Some properties of generalized Fisher information in the context of nonextensive thermostatistics

Jean-François Bercher

► **To cite this version:**

Jean-François Bercher. Some properties of generalized Fisher information in the context of nonextensive thermostatistics. 2012. hal-00766699v1

HAL Id: hal-00766699

<https://hal.science/hal-00766699v1>

Preprint submitted on 18 Dec 2012 (v1), last revised 4 Jun 2013 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Some properties of generalized Fisher information in the context of nonextensive thermostatics

J.-F. Bercher

*Université Paris-Est, LIGM, UMR CNRS 8049, ESIEE-Paris
5 bd Descartes, 77454 Marne la Vallée Cedex 2, France
tel: 33-1-45-92-65-15 fax: 33-1-45-92-66-99*

Abstract

We present two extended forms of Fisher information that fit well in the nonextensive thermostatics context. We show that there exists an interplay between these generalized Fisher information, the generalized q -Gaussian distributions and the q -entropies. The minimum of the generalized Fisher information among distributions with a fixed moment, or with a fixed q -entropy is attained, in both cases, by a generalized q -Gaussian distribution. This complements the fact that the q -Gaussians maximize the q -entropies subject to a moment constraint, and yields new variational characterizations of the generalized q -Gaussians. We show that the generalized Fisher information naturally pop up in the expression of the time derivative of the q -entropies, for distributions satisfying a certain nonlinear heat equation. This result includes as a particular case the classical de Bruijn identity. We then study further properties of the generalized Fisher information and of their minimization. We show that, though non additive, the generalized Fisher information of a combined system is upper bounded. In the case of mixing, we show that the generalized Fisher information is convex for $q \geq 1$. Finally, we show that the minimization of the generalized Fisher information subject to moment constraints satisfies a Legendre structure analog to the Legendre structure of thermodynamics.

Keywords: Generalized Fisher information, Generalized Rényi and Tsallis entropies, Generalized q -gaussian distributions, Information theoretic inequalities

PACS: 02.50.-r, 05.90.+m, 89.70.+c

Email address: jf.bercher@esiee.fr (J.-F. Bercher)

1. Introduction and preliminary definitions

Information measures are important both for the foundation of information sciences and for practical applications of information processing. They are of outmost importance in several areas of physics, beginning of course with statistical physics. Entropies and Fisher information have been identified as useful and versatile tools for characterizing complex systems, see e.g. [1, 2, 3]. Classical information theoretic inequalities, as described for instance in [4, 5], interrelate information measures. These inequalities have proved to be useful for communication theoretic problems and engineering applications. They are also connected to uncertainty relations in physics, and to functional inequalities in mathematics. The aim of the paper is to present and study two generalized Fisher information measure that fit well in the context of nonextensive thermostatics, and to describe the interplay between information measures and generalized q -Gaussians, thus providing new characterization of these generalized q -Gaussians. We first recall the context, our notations and main definitions.

Generalized entropies

The Boltzmann entropy, linked to the Shannon entropy of information theory, is at the heart of thermodynamics. For energy constraints, it is well-known that the maximum entropy distribution is the Gaussian distribution. For the analysis of complex systems, generalized entropies, which reduces to the standard one as a particular case, have been proposed. In particular, the thermodynamics derived from Tsallis entropy, the nonextensive thermodynamics, has received a high attention and there is a wide variety of applications where experiments, numerical results and analytical derivations fairly agree with the new formalisms [6]. Some physical applications of the generalized entropies, – including statistics of cosmic rays, defect turbulence, optical lattices, systems with long-range interactions, superstatistical systems, etc., can be found in the recent review [7] and references therein. Let us recall here that if x is a vector of $\Omega \subseteq \mathbb{R}^n$, and $f(x)$ a probability density defined with respect to the Lebesgue measure, then following [8], the information generating function is the quantity $M_q[f] = \int_{\Omega} f(x)^q dx$, for $q \geq 0$. The Tsallis entropy is given by $S_q[f] = \frac{1}{1-q} (M_q[f] - 1)$, while the Rényi entropy is given by $H_q[f] = \frac{1}{1-q} \log M_q[f]$. For our purposes, it is not always necessary to distinguish between the two entropies that we will call collectively q -entropies. Finally, we will also call “entropy power” of order q ,

or q -entropy power the quantity

$$N_q[f] = M_q[f]^{\frac{1}{1-q}} = \exp(H_q[f]) = \left(\int_{\Omega} f(x)^q dx \right)^{\frac{1}{1-q}}, \quad (1)$$

for $q \neq 1$. For $q = 1$, we let $N_q[f] = \exp(H_1[f])$, where $H_1[f]$ is the Boltzmann-Shannon entropy.

Generalized q -Gaussian distributions

In the context of nonextensive thermostatics, the role devoted to the standard Gaussian distribution is extended to generalized q -Gaussian distributions, which include the standard Gaussian as a special case. These generalized q -Gaussian distributions form a versatile family that can describe problems with compact support as well as problems with heavy tailed distributions. They are also analytical solutions of actual physical problems, see e.g. [9, 10] [11], and are sometimes known as Barenblatt-Pattle functions, following their identification by [12, 13]. We shall also mention that the Generalized q -Gaussian distributions appear in other fields, namely as the solution of non-linear diffusion equations, or as the distributions that saturate some sharp inequalities in functional analysis [14, 15, 16, 17].

Definition 1. Let x be a random vector of \mathbb{R}^n , and let $|x|$ denote its Euclidean norm. For $\alpha \in (0, \infty)$, γ a real positive parameter and $q > (n - \alpha)/n$, the generalized Gaussian with scale parameter γ has the radially symmetric probability density

$$G_{\gamma}(x) = \begin{cases} \frac{1}{Z(\gamma)} (1 - (q - 1) \gamma |x|^{\alpha})_+^{\frac{1}{q-1}} & \text{for } q \neq 1 \\ \frac{1}{Z(\gamma)} \exp(-\gamma |x|^{\alpha}) & \text{if } q = 1 \end{cases} \quad (2)$$

where we use the notation $(x)_+ = \max\{x, 0\}$, and where $Z(\gamma)$ is the partition function such that $G_{\gamma}(x)$ integrates to one. Its expression is given in the Appendix, section 5, eq. (78).

In the sequel, we will denote G the generalized Gaussian obtained with $\gamma = 1$. For $q > 1$, the density has a compact support, while for $q \leq 1$ it is defined on the whole \mathbb{R}^n and behaves as a power distribution for $|x| \rightarrow \infty$. A shorthand notation for the expression of the generalized q -Gaussian density is

$$G_{\gamma}(x) = \frac{1}{Z(\gamma)} \exp_{q^*}(-\gamma |x|^{\alpha}), \quad (3)$$

with $q^* = 2 - q$, and where the so-called q -exponential function, and its inverse the q -logarithm, are given by

$$\exp_q(x) := (1 + (1 - q)x)_+^{\frac{1}{1-q}} \text{ for } q \neq 1 \text{ and } \exp_{q=1}(x) := \exp(x), \quad (4)$$

$$\ln_q(x) := \frac{x^{1-q} - 1}{1 - q} \text{ for } q \neq 1 \text{ and } \ln_{q=1}(x) := \ln(x). \quad (5)$$

The general expressions of the main information measures attached to the generalized Gaussians are derived in section 5.

It is well known that the Gaussian distribution is a central distribution with respect to classical information measures and inequalities. In particular, the Gaussian distribution is both a maximum entropy and a minimum Fisher information distribution over all distributions with the same variance. We will see that the same kind of result holds for the family of generalized q -Gaussians, for Rényi or Tsallis entropy and suitable extensions of the Fisher information.

Generalized (β, q) -Fisher information

In the context of nonextensive thermostatics, several authors have introduced and studied generalized versions of the Fisher information. Among these contributions we note the series of papers [18, 19, 20, 21], and the proposals of Naudts [22, 23] and Furuichi [24, 25], which are related to the generalized Fisher information that are used here – see [26] for a discussion of these links.

In our recent work [26, 27], we have thrown a bridge between concepts in estimation theory and tools of nonextensive thermostatics. Using the notion of escort distribution, we have introduced a generalized Fisher information and established an extended version of the Cramér-Rao inequality for parameter estimation. In the case of a location parameter, it reduces to an extended version of the standard Cramér-Rao inequality, which is saturated by the generalized q -Gaussians. A closely related generalized Fisher information has been originally introduced by Lutwak *et al.* [28] in information theory and extended to the multidimensional in [29] and independently in [30]. These two extensions of the Fisher information measure, the (β, q) -Fisher information $\phi_{\beta, q}[f]$ and $I_{\beta, q}[f]$, depend on an entropic index q and on a parameter β . Both information reduce to the standard Fisher information in the case $q = 1$, $\beta = 2$. They are defined as follows.

Definition 2. Let $f(x)$ be a probability density function defined over a subset Ω of \mathbb{R}^n . Let $|x|$ denote the Euclidean norm of x and ∇f the gradient

operator. If $f(x)$ is continuously differentiable over Ω , then for $q \geq 0$, $\beta > 1$, the generalized (β, q) -Fisher information is defined by

$$\phi_{\beta,q}[f] = \int_{\Omega} f(x)^{\beta(q-1)+1} \left(\frac{|\nabla f(x)|}{f(x)} \right)^{\beta} dx \quad (6)$$

$$= E \left[f(x)^{\beta(q-1)} |\nabla \ln f(x)|^{\beta} \right] = E \left[|\nabla \ln_{q^*} f(x)|^{\beta} \right] \quad (7)$$

with $q^* = 2 - q$.

The second form of generalized Fisher information appears in the derivation and formulation of a general Cramér-Rao inequality in estimation problems [26, 27].

Definition 3. With the same notations and assumptions as above, the second generalized (β, q) -Fisher information is

$$I_{\beta,q}[f] = \left(\frac{q}{M_q[f]} \right)^{\beta} \phi_{\beta,q}[f] = \left(\frac{q}{\int_{\Omega} f(x)^q dx} \right)^{\beta} \int_{\Omega} f(x)^{\beta(q-1)+1} \left(\frac{|\nabla f(x)|}{f(x)} \right)^{\beta} dx \quad (8)$$

$$= \left(\frac{q}{M_q[f]} \right)^{\beta} E \left[|\nabla \ln_{q^*} f(x)|^{\beta} \right]. \quad (9)$$

Structure and contributions of the paper

As already mentioned, the goal of the paper is to introduce two generalized Fisher information in the context of nonextensive thermostatics, to document their main properties and study the interplay between generalized information measures and generalized q -Gaussians. Though the paper reviews some of our previous results, most of the results presented here are new. The paper is organized into two main parts. In the first part, we present several variational characterizations of the generalized q -Gaussian distributions, where the generalized Fisher information play a fundamental role. The new findings include (i) the second generalized Stam inequality (11) which lower bounds the product of the entropy power and of the extended Fisher information, (ii) the definition of several information functionals minimized by the generalized q -Gaussians, (iii) the derivation of the extended de Bruijn inequality (36, 37) and of the de Bruijn entropy power inequality (38) which intimately links the q -entropies to the generalized Fisher information through a doubly nonlinear diffusion equation. In the second part, we establish several properties of the generalized Fisher information which might be useful for thermodynamics considerations. The discussion includes the additivity and mixing properties of the generalized Fisher information, as well as an original derivation of the preservation of the Legendre structure.

2. Variational characterizations of generalized q -Gaussians and generalized Fisher information

As already pointed out, the q -Gaussians maximize the Tsallis or Rényi entropy among all probability distributions with a given variance or covariance [31]. This means that the q -Gaussians are the canonical distributions of the generalized thermostatics based on Tsallis or Rényi entropy. The maximum entropy formalism directly induces a variational characterization of the generalized q -Gaussians as the extremal functions of the Lagrangian

$$L[f] = H_q[f] + \mu m_\alpha[f],$$

where μ is a Lagrange multiplier and $m_\alpha[f]$ denotes the moment of order α of the probability distribution f . A remarkable point is that the generalized q -Gaussians are also solutions of other variational problems as well, and that these problems involve the generalized Fisher information we defined above. Such characterizations are described now.

2.1. The minimum of generalized Fisher information among distributions with a given q -entropy is attained for generalized q -Gaussians

Another characterization of the generalized q -Gaussian is indeed the fact that they minimize the extended Fisher information among all distributions with a given q -entropy. This result is a consequence of a generalized Stam inequality which lower bounds a product of the entropy power and of the generalized Fisher information. This inequality is stated in the Proposition below and has been established in [28] in the monodimensional case, in [29, 30] in the multidimensional case, and in [27] for arbitrary norms. The statement of the case (b) is new.

Proposition 1. *[Generalized Stam inequality] For $n \geq 1$, β and α Hölder conjugates of each other, $\alpha > 1$, and $q > \max\{(n-1)/n, n/(n+\alpha)\}$, then for any probability density on \mathbb{R}^n , supposed continuously differentiable, the following generalized Stam inequalities holds*

$$(a) \quad \phi_{\beta,q}[f]^{\frac{1}{\beta}} N_q[f]^{\frac{\lambda}{n}} \geq \phi_{\beta,q}[G]^{\frac{1}{\beta}} N_q[G]^{\frac{\lambda}{n}}, \quad (10)$$

$$(b) \quad I_{\beta,q}[f]^{\frac{1}{\beta}} N_q[f]^{\frac{1}{n}} \geq I_{\beta,q}[G]^{\frac{1}{\beta}} N_q[G]^{\frac{1}{n}}. \quad (11)$$

with $\lambda = n(q-1)+1 > 0$ and with equality if and only if f is any generalized q -Gaussian (3).

In the previous expressions, $\phi_{\beta,q}[G]$, $I_{\beta,q}[G]$ and $N_q[G]$ are the values taken by the Fisher information and the entropy power when the probability density f is the generalized q -Gaussian G . The exact expressions of these quantities are given in the Appendix, section 5.

Proof. The inequality (a) is proved in [29] and [30] using a general sharp Gagliardo-Nirenberg inequality due to Cordero et al. [16]. It is quite easy to get the same kind of generalized Stam inequality for the second generalized Fisher information. Indeed, for the inequality (b), it suffices to replace $\phi_{\beta,q}[f]^{\frac{1}{\beta}}$ in (10) by $I_{\beta,q}[f]^{\frac{1}{\beta}} M_q[f]/q$, to use the fact that $M_q[f] = N_q[f]^{1-q}$ and to simplify the exponent to obtain (11). \square

The generalized Stam inequalities imply that the generalized q -Gaussian minimize the extended Fisher information within the set of probability distributions with a fixed q -entropy power: G_γ is the solution of

$$\inf_f \{ \phi_{\beta,q}[f] \text{ or } I_{\beta,q}[f] : f \in \mathcal{P}, M_q[f] = M_q[G_\gamma] \}, \quad (12)$$

where \mathcal{P} is the probability simplex. In turn, this leads to the following variational description.

Proposition 2. *The generalized q -Gaussian G minimize the following two information functionals*

$$J_1[f] = \frac{1}{\beta\lambda} \phi_{\beta,q}[f] N_q[G] + \frac{1}{n} \phi_{\beta,q}[G] N_q[f] \quad (13)$$

$$J_2[f] = \frac{1}{\beta} I_{\beta,q}[f] N_q[G] + \frac{1}{n} I_{\beta,q}[G] N_q[f] \quad (14)$$

where the values $\phi_{\beta,q}[G]$, $I_{\beta,q}[G]$ and $N_q[G]$ are given in the Appendix, section 5.

Another statement of this result could be in terms of an energy functional. Let $f = u^k$, with $k = \beta / (\beta(q-1) + 1)$. With this notation, the generalized Fisher information reduces to

$$\phi_{\beta,q}[f] = |k|^\beta \int_{\Omega} |\nabla u(x)|^\beta dx, \quad (15)$$

which is the β -Dirichlet energy. Additionally, $M_q[f] = \int_{\Omega} u(x)^{kq} dx$ is a measure of informational energy of order kq in the sense of [32, 33]. Then, (13) becomes

$$J_1[f] = \frac{1}{\beta\lambda} N_q[G] \int_{\Omega} |\nabla u(x)|^{\beta} dx + \frac{1}{n} \phi_{\beta,q}[G] \left(\int_{\Omega} u(x)^{kq} dx \right)^{\frac{1}{1-q}} \quad (16)$$

a weighted sum of these two energies.

Proof. Consider the first generalized Stam inequality. As already indicated, a direct consequence of this inequality is that the generalized q -Gaussian solves problem (12). The Lagrangian corresponding to this problem is

$$J(f) = \phi_{\beta,q}[f] + \mu N_q[f]. \quad (17)$$

This Lagrangian is minimum for a q -Gaussian G_{γ} with parameter γ . This parameter γ is a function of the Lagrange parameter μ . Suppose that μ is chosen such that the optimum distribution is the generalized q -Gaussian with $\gamma = 1$. In such conditions, we have $J(G_{\gamma}) \geq J(G)$, $\forall \gamma$. On the other hand, we have the following scaling identities:

$$\begin{cases} M_q[G_{\gamma}] = \gamma^{\frac{n}{\alpha}(q-1)} M_q[G], \\ \phi_{\beta,q}[G_{\gamma}] = \gamma^{\frac{\beta\lambda}{\alpha}} \phi_{\beta,q}[G], \\ I_{\beta,q}[G_{\gamma}] = \gamma^{\frac{\beta}{\alpha}} I_{\beta,q}[G]. \end{cases} \quad (18)$$

Thus, the inequality $J(G_{\gamma}) \geq J(G)$ means that

$$J(G_{\gamma}) = \gamma^{\frac{\beta\lambda}{\alpha}} \phi_{\beta,q}[G] + \mu \gamma^{-\frac{n}{\alpha}} N_q[G] \quad (19)$$

is minimum in γ for $\gamma = 1$. The derivative with respect to γ then vanishes at $\gamma = 1$, and this yields the value

$$\mu = \frac{\beta\lambda}{n} \frac{\phi_{\beta,q}[G]}{N_q[G]}. \quad (20)$$

Then, it only remains to set $J_1[f] = \frac{N_q[G]}{\beta\lambda} J[f]$ to obtain (13). The second information functional (14) is established in the same way. \square

2.2. The minimum of generalized Fisher information among distributions with a given moment is attained for generalized q -Gaussians

In addition to be the maximum entropy distribution in the set of all distributions with a given variance, it is well-known that the Gaussian distribution is also a minimum Fisher information distribution over all distributions

with the same variance. This can be seen as a consequence of the Cramér-Rao inequality for a location parameter. The same kind of characterization holds in the multidimensional case for the generalized q -Gaussian and the generalized Fisher information.

In [28, 29, 30], a Cramér-Rao inequality for generalized q -Gaussian involving the Fisher information $\phi_{\beta,q}[f]$ has been established. The Fisher information $I_{\beta,q}[f]$ has been introduced in the context of parameter estimation in [26, 27] and a related Cramér-Rao inequality proved. These inequalities mean that among all distributions with a given moment, the generalized q -Gaussians are the minimizers of extended versions of the Fisher information, just as the standard Gaussian minimizes Fisher information over all distributions with a given variance. We begin by recalling the generalized Cramér-Rao inequalities involving the generalized Fisher information measures and saturated by the generalized q -Gaussians. Then, we show that the generalized q -Gaussians can also be characterized as the minimizers of two new variational problems.

Proposition 3. *For $n \geq 1$, β and α Hölder conjugates of each other, $\alpha > 1$, $q > \max\{(n-1)/n, n/(n+\alpha)\}$ then for any probability density f on $\Omega \subseteq \mathbb{R}^n$, supposed continuously differentiable and such that the involved information measures are finite,*

$$m_\alpha[f]^\frac{1}{\alpha} \phi_{\beta,q}[f]^\frac{1}{\beta\lambda} \geq m_\alpha[G]^\frac{1}{\alpha} \phi_{\beta,q}[G]^\frac{1}{\beta\lambda} \quad (21)$$

$$m_\alpha[f]^\frac{1}{\alpha} I_{\beta,q}[f]^\frac{1}{\beta} \geq m_\alpha[G]^\frac{1}{\alpha} I_{\beta,q}[G]^\frac{1}{\beta} = n \quad (22)$$

with $\lambda = n(q-1)+1$, where the general Fisher information are defined in (6) and (8) and where the equality holds iff f is a generalized Gaussian $g = G_\gamma$.

Proof. The proof of (21) can be found in [29, 30] in the multidimensional case. As far as (22) is concerned, the proof is given in [26, 27] as a particular case of the more general Cramér-Rao inequality for parameter estimation. The fact that the lower bound $m_\alpha[G]^\frac{1}{\alpha} I_{\beta,q}[G]^\frac{1}{\beta}$ is exactly equal to n is a direct consequence of the proof of (22). This is also checked here in Appendix 5 where we derive the information measures for the generalized q -Gaussian G . \square

The Cramér-Rao inequalities (21)-(22) imply that G_γ realizes the minimum of the generalized Fisher information in the set of all probability densities with a given moment of order α :

$$\inf_f \{ \phi_{\beta,q}[f] \text{ or } I_{\beta,q}[f] : f \in \mathcal{P}, m_\alpha[f] = m_\alpha[G_\gamma] \}, \quad (23)$$

where \mathcal{P} is the probability simplex. As above and as a direct consequence, we obtain a variational characterization of the generalized q -Gaussians:

Proposition 4. *The generalized q -Gaussian G minimize the following two information functionals*

$$J_3[f] = \frac{1}{\beta\lambda} \phi_{\beta,q}[f] m_\alpha[G] + \frac{1}{\alpha} \phi_{\beta,q}[G] m_\alpha[f] \quad (24)$$

$$J_4[f] = \frac{1}{\beta} I_{\beta,q}[f] m_\alpha[G] + \frac{1}{\alpha} I_{\beta,q}[G] m_\alpha[f] \quad (25)$$

Proof. The proof proceeds just as the proof of Proposition 2. Let us consider the first Cramér-Rao inequality (21) and the problem (23). The corresponding Lagrangian is

$$J(f) = \phi_{\beta,q}[f] + \mu m_\alpha[f]. \quad (26)$$

Assume that μ is chosen such that the optimum distribution is the generalized q -Gaussian with $\gamma = 1$. In such conditions, we have $J(G_\gamma) \geq J(G)$, $\forall \gamma$. By the scaling identities (18) and the additional fact that $m_\alpha[G_\gamma] = \gamma^{-1} m_\alpha[G]$, we get that

$$J(G_\gamma) = \gamma^{\frac{\beta\lambda}{\alpha}} \phi_{\beta,q}[G] + \mu \gamma^{-1} m_\alpha[G]. \quad (27)$$

Since this functional is minimum for $\gamma = 1$, its derivative vanishes at $\gamma = 1$, which gives

$$\mu = \frac{\beta\lambda}{\alpha} \frac{\phi_{\beta,q}[G]}{m_\alpha[G]} \quad (28)$$

and in turn the inequality (24). The second information functional (25) is established in the same way. \square

We shall note that up to the change of function $u(x)^k = f(x)$, the Euler-Lagrange equation associated with the two functionals (24) and (25) is a β -Laplace equation of the form

$$\Delta_\beta u(x) - \frac{1}{\alpha k^{\beta-1}} \frac{\phi_{\beta,q}[G]}{m_\alpha[G]} |x|^\alpha u(x)^{k-1} = 0, \quad (29)$$

where we used the β -Laplacian operator $\Delta_\beta u := \operatorname{div}(|\nabla u|^{\beta-2} \nabla u)$. In the monodimensional case and if $\beta = 2$, we obtain an instance of the generalized Emden-Fowler equation $u''(x) + h(x) u(x)^\gamma = 0$, where $h(x)$ is a given function. This kind of equations arise in studies of gaseous dynamics in astrophysics, in certain problems in fluid mechanics and pseudoplastic flow, see [34], as well as in some reaction-diffusion processes.

2.3. *The derivative of q -entropies give the generalized Fisher information – An extended de Bruijn identity*

A fundamental connection between the Boltzmann-Shannon entropy, Fisher information, and the Gaussian distribution is given by the de Bruijn identity [35]. We show here that this important connection can be extended to the q -entropies, the generalized Fisher information and the generalized q -Gaussians.

The de Bruijn identity states that if $Y_t = X + \sqrt{2t}Z$ where Z is a standard Gaussian vector and X a random vector of \mathbb{R}^n , independent of Z , then

$$\frac{d}{dt}H[f_{Y_t}] = I_{2,1}[f_{Y_t}] = \phi_{2,1}[f_{Y_t}], \quad (30)$$

where f_{Y_t} denotes the density of $Y_t = X + \sqrt{2t}Z$. In thermodynamics, this can be seen as a consequence of the second law for an isolated system out of equilibrium. The standard proof of the de Bruijn identity uses the fact that Y_t satisfies the well-known heat equation [36]

$$\frac{\partial f}{\partial t} = \Delta f, \quad (31)$$

where Δ denotes the Laplace operator.

It is possible to consider more general versions of the heat equation. In particular, it is known that the porous medium equation, or the fast diffusion equation, admit q -Gaussians as solutions. This connects the q -Gaussian distributions to differential equations of physics, and has been noticed in the nonextensive context, see e.g. [10, 37]. The porous medium equation and fast diffusion equation correspond to the differential equation

$$\frac{\partial f}{\partial t} = \Delta f^m, \quad (32)$$

with $m > 1$ for the porous medium equation and < 1 for the fast diffusion. These two equations have been exhaustively studied and characterized by J. L. Vázquez, e.g. in the two books [38, 39]. These equations appear in a large number of physical situations, including fluid mechanics, nonlinear heat transfer or diffusion. Other applications have been reported in mathematical biology, lubrication, boundary layer theory, etc, see the series of applications presented in [38, chapters 2 and 21] and references therein.

The porous medium equation (32) can be generalized to a wider form, the *doubly nonlinear equation*, which involves a p -Laplacian operator $\Delta_p f :=$

$\operatorname{div}(|\nabla f|^{p-2}\nabla f)$, and the power m of the porous medium or fast diffusion equation. This doubly nonlinear equation takes the form

$$\frac{\partial}{\partial t}f = \Delta_{\beta}f^m = \operatorname{div}\left(|\nabla f^m|^{\beta-2}\nabla f^m\right), \quad (33)$$

where we use $p = \beta$ for convenience and coherence with the other notations in the paper. The β -Laplacian operator typically appears for the problem of minimization of a Dirichlet energy like $\int |\nabla f|^{\beta}dx$ which leads to the Euler-Lagrange equation $\Delta_{\beta}f = 0$. A discussion on the p -Laplace equation can be found in [40].

As we can see, the doubly nonlinear equation includes the standard heat equation ($\beta = 2, m = 1$), the β -Laplace equation ($\beta \neq 2, m = 1$), the porous medium equation ($\beta = 2, m > 1$) and the fast diffusion equation ($\beta = 2, m < 1$). It can be shown, see [39, page 192], that for $m(\beta-1) + (\beta/n) - 1 > 0$, (33) has a unique self-similar solution, whose initial value is the Dirac mass at the origin. This fundamental solution is given by

$$f(x, t) = \frac{1}{t^{\frac{n}{\delta}}}B\left(\frac{x}{t^{\frac{1}{\delta}}}\right), \quad \text{with } B(x) = \begin{cases} (C - k|x|^{\alpha})_{+}^{\frac{\beta-1}{m(\beta-1)-1}} & \text{for } m \neq \frac{1}{\beta-1} \\ \frac{1}{\sigma} \exp\left(-\frac{|\beta-1|}{\beta\alpha}|x|^{\alpha}\right) & \text{for } m = \frac{1}{\beta-1} \end{cases} \quad (34)$$

with

$$\delta = n(\beta-1)m + \beta - n > 0, \quad k = \frac{m(\beta-1) - 1}{\beta} \left(\frac{1}{\delta}\right)^{\frac{1}{\beta-1}} \quad \text{and } \alpha = \frac{\beta}{\beta-1}. \quad (35)$$

The constants C and σ are uniquely determined by mass conservation, e.g. $\int f(x, t)dx = 1$. Of course, we observe that the function $B(x)$ above is analog to the generalized q -Gaussian (3). In the literature, $B(x)$ is called the Barenblatt profile following its identification as a solution for heat release from a point source [12, 13].

The doubly nonlinear diffusion equation also enables to derive a nice extension of the de Bruijn identity (30). A remarkable point is that the extended Fisher information naturally pops up in this identity, generalizing the role of the standard Fisher information in the classical de Bruijn identity. This is stated in the following Proposition.

Proposition 5. *[Extended de Bruijn identity] Let $f(x, t)$ a probability distributions satisfying the doubly nonlinear equation (33). For $q = m + 1 - \frac{\alpha}{\beta}$,*

$\lambda = n(q - 1) + 1$, and $\delta = \beta\lambda - n(q - 1)$, we have

$$\frac{d}{dt} H_q[f] = \frac{q m^{\beta-1}}{M_q[f]} \phi_{\beta,q}[f] = \left(\frac{m}{q}\right)^{\beta-1} M_q[f]^{\beta-1} I_{\beta,q}[f]. \quad (36)$$

$$\frac{d}{dt} S_q[f] = q m^{\beta-1} \phi_{\beta,q}[f] = \left(\frac{m}{q}\right)^{\beta-1} M_q[f]^\beta I_{\beta,q}[f]. \quad (37)$$

where $M_q[f] = \int f^q$ and where $H_q[f] = \frac{1}{1-q} \log M_q[f]$ is the Rényi entropy and $S_q[f] = \frac{1}{1-q} (M_q[f] - 1)$ the Tsallis entropy. In terms of the entropy power $N_q[f] = M_q[f]^{\frac{1}{1-q}}$ we also have

$$\frac{d}{dt} N_q[f]^{\frac{\delta}{n}} = q \frac{\delta}{n} m^{\beta-1} N_q[f]^{\frac{\beta\lambda}{n}} \phi_{\beta,q}[f] = \frac{\delta}{n} \left(\frac{m}{q}\right)^{\beta-1} N_q[f]^{\frac{\beta}{n}} I_{\beta,q}[f], \quad (38)$$

and

$$\frac{d}{dt} N_q[f]^{\frac{\delta}{n}} \geq \frac{d}{dt} N_q[G]^{\frac{\delta}{n}}. \quad (39)$$

Of course, the classical de Bruijn identity

$$\frac{d}{dt} H[f] = \phi_{2,1}[f] = I_{2,1}[f]. \quad (40)$$

is recovered from the extended de Bruijn identity (36), for $\alpha = \beta = 2$, and $q = m = 1$. We recognize in the right side of (38) the very same products of entropy power and Fisher information that appear in the the generalized Stam inequalities (10) and (11). By these inequalities, we obtain (39) which indicates that the derivative of the entropy power in minimum for the generalized q -Gaussian, with equality if and only if $f = G_\gamma$, $\forall \gamma$. Furthermore, it is known that the solutions of the doubly nonlinear equation converge to the Barenblatt profile [39]. Thus, we see that the minimum of the derivative of the entropy power is always attained asymptotically. In a very recent and nice paper [41], Savaré and Toscani have shown that in the case $\beta = 2$, $m = q$, the entropy power $N_q[f]^{\frac{\delta}{n}}$ is a concave function of t . A consequence of this result is that the derivative (38) of entropy power is non increasing in time, with a minimum for $f = G$, reached at least asymptotically. It would be very interesting to get the same kind of result in the present setting. For now, let us proceed with the proof of extended de Bruijn identity.

Proof. Considering the Rényi entropy

$$H_q[f] = \frac{1}{1-q} \log \int_{\Omega} f(x,t)^q dx, \quad (41)$$

the derivative with respect to t is

$$(1 - q) \frac{\partial}{\partial t} H_q[f] = \frac{1}{\int_{\Omega} f(x, t)^q dx} \int_{\Omega} \frac{\partial}{\partial t} f(x, t)^q dx = \frac{q}{\int_{\Omega} f(x, t)^q dx} \int_{\Omega} f(x, t)^{q-1} \frac{\partial}{\partial t} f(x, t) dx \quad (42)$$

$$= \frac{q}{\int_{\Omega} f(x, t)^q dx} \int_{\Omega} f(x, t)^{q-1} \Delta_{\beta} f(x, t)^m dx \quad (43)$$

$$= \frac{q}{\int_{\Omega} f(x, t)^q dx} \int_{\Omega} f(x, t)^{q-1} \nabla \cdot [|\nabla f^m|^{\beta-2} \nabla f^m] dx \quad (44)$$

where we used the fact that $f(x, t)$ satisfies the doubly nonlinear heat equation (33). Let ψ be a scalar function and A a vector valued function defined on a subset U of \mathbb{R}^n . By the divergence theorem and the product rule, we have

$$\int_U \nabla \cdot (\psi A) dV = \int_{\partial U} \psi A \cdot n dS = \int_U \psi \nabla \cdot A dV + \int_U A \cdot \nabla \psi dV. \quad (45)$$

Thus, considering

$$\int_{\Omega} f(x, t)^{q-1} \Delta_{\beta} f(x, t)^m dx = \int_{\Omega} f(x, t)^{q-1} \nabla \cdot [|\nabla f^m|^{\beta-2} \nabla f^m] dx, \quad (46)$$

we get, with $\psi = f(x, t)^{q-1}$ and $A = |\nabla f^m|^{\beta-2} \nabla f^m$,

$$\begin{aligned} \int_{\Omega} f(x, t)^{q-1} \nabla \cdot [|\nabla f^m|^{\beta-2} \nabla f^m] dx &= \int_{\partial \Omega} f(x, t)^{q-1} |\nabla f^m|^{\beta-2} \nabla f^m \cdot n dx \\ &\quad - \int_{\Omega} (q-1) f(x, t)^{q-2} \nabla f \cdot [|\nabla f^m|^{\beta-2} \nabla f^m] dx. \end{aligned} \quad (47)$$

Assuming that both $f(x, t)$ and $|\nabla f|$ tend to zero on the boundary $\partial \Omega$ of the domain, it remains

$$\int_{\Omega} f(x, t)^{q-1} \nabla \cdot [|\nabla f^m|^{\beta-2} \nabla f^m] dx = -(q-1) \int_{\Omega} f(x, t)^{q-2} \nabla f \cdot [|\nabla f^m|^{\beta-2} \nabla f^m] dx, \quad (48)$$

$$= -(q-1) m^{\beta-1} \int_{\Omega} f(x, t)^{\beta m + q - m - 1} \left(\frac{|\nabla f|}{f} \right)^{\beta} dx, \quad (49)$$

where we used twice the fact that $\nabla f^m = m f^{m-1} \nabla f$. Finally, if we choose $q = m + 1 - \frac{\alpha}{\beta}$, the exponent in the last equation reduces to $\beta(q-1) + 1$,

and

$$\frac{d}{dt} H_q[f] = \frac{q m^{\beta-1}}{\int f(x)^q dx} \int_{\Omega} f(x)^{\beta(q-1)+1} \left(\frac{|\nabla f|}{f} \right)^{\beta} dx \quad (50)$$

$$= \frac{q m^{\beta-1}}{M_q[f]} \phi_{\beta,q}[f]. \quad (51)$$

Using the relationship (8) between $I_{\beta,q}[f]$ and $\phi_{\beta,q}[f]$, this identity becomes

$$\frac{d}{dt} H_q[f] = \left(\frac{m}{q} \right)^{\beta-1} M_q[f]^{\beta-1} I_{\beta,q}[f]. \quad (52)$$

Of course, the second inequality (37) follows immediately. If we consider the entropy power $N_q = M_q^{\frac{1}{1-q}}$ instead of the entropy, we obtain along similar steps the following relation:

$$\frac{\partial}{\partial t} N_q[f]^{\frac{\delta}{n}} = q \frac{\delta}{n} m^{\beta-1} N_q[f]^{\frac{\beta\lambda}{n}} \phi_{\beta,q}[f] = \frac{\delta}{n} \left(\frac{m}{q} \right)^{\beta-1} N_q[f]^{\frac{\beta}{n}} I_{\beta,q}[f]. \quad (53)$$

with $\lambda = n(q-1) + 1$ and $\delta = n(\beta-1)(q-1) + \beta = \beta\lambda - n(q-1)$. □

3. Thermodynamics considerations

Given an information measure or a generalized entropy, it is possible to investigate whether it is possible to construct a thermodynamics based on this measure. Several of such constructions have been proposed, namely the nonextensive thermostatics based on the Tsallis entropy, see the books [42, 6], the construction of thermodynamics based on the Rényi entropy, with connections with the multifractals [43, 44, 45, 46], or based on the Sharma-Mittal entropy [47]. Generalized entropies obtained through Beck and Cohen's superstatistics have also been considered [48, 7, 49]. Finally, following the pioneering work of Frieden, thermodynamics based on the Fisher information have been proposed [50, 51, 52, 53, 54].

Though it is not the objective of the present paper to discuss and compare the pros and cons of these different thermodynamics, these prior works naturally lead us to examine the status of some basic properties of conventional thermodynamics in the case of the generalized Fisher information. In this spirit, we consider the additivity property of the information, the mixing property, and finally the Legendre structure of thermodynamics.

3.1. Additivity

In classical thermodynamics, additivity is the principle that if a global system is constituted of two independent components, then the total entropy is the sum of entropies of each subsystem. However, the use of a different form of entropy or complexity measure can lead to reconsider this property. It is well known, for instance, that the Tsallis entropy is non additive. Indeed, if we use a measure of information, it is understandable that the information given by a whole system can be different of the sum of the information given by its components ; the difference representing an information gain attached to the combining of the elementary systems. In the case of Fisher information, the information is additive for independent systems. However, this property is not preserved for the generalized Fisher information as soon as $q \neq 1$ or $\beta \neq 2$. Nevertheless, we still have an interesting inequality which bounds the Fisher information of a combined system. Indeed, if X and Y are two independent random vectors with densities $f_X(x)$ and $f_Y(y)$, then the generalized Fisher information $\phi_{\beta,q}$ attached to the joint distribution $f_{X,Y}(x,y) = f_X(x)f_Y(y)$ is

$$\phi_{\beta,q}[f_X f_Y] = \int (f_X(x)f_Y(y))^{\beta(q-2)+1} |\nabla f_X(x) f_Y(y) + f_X(x)\nabla f_Y(y)|^\beta dx dy. \quad (54)$$

The Minkovski inequality indicates that the triangle inequality holds in L_p spaces, e.g. for a weightened L_p norm

$$\left(\int w(x)|f(x) + g(x)|^p dx \right)^{\frac{1}{p}} \leq \left(\int w(x)|f(x)|^p dx \right)^{\frac{1}{p}} + \left(\int w(x)|g(x)|^p dx \right)^{\frac{1}{p}}, \quad (55)$$

with $w(x) > 0$ and $p > 1$. Applying this to the generalized Fisher information (54), we immediately obtain

$$\phi_{\beta,q}[f_X f_Y]^{\frac{1}{\beta}} \leq M_{\beta(q-1)+1}[f_Y]^{\frac{1}{\beta}} \phi_{\beta,q}[f_X]^{\frac{1}{\beta}} + M_{\beta(q-1)+1}[f_X]^{\frac{1}{\beta}} \phi_{\beta,q}[f_Y]^{\frac{1}{\beta}}, \quad (56)$$

where $M_q[f]$ is the information generating function. By the general power mean inequality, we know that $M_a[f]^{\frac{1}{a}} \leq M_b[f]^{\frac{1}{b}}$ for $a < b$. Consequently, when $q < 1$, $M_{\beta(q-1)+1} < M_1 = 1$, and the inequality (56) reduces to

$$\phi_{\beta,q}[f_X f_Y]^{\frac{1}{\beta}} \leq \phi_{\beta,q}[f_X]^{\frac{1}{\beta}} + \phi_{\beta,q}[f_Y]^{\frac{1}{\beta}}. \quad (57)$$

3.2. Mixing property

It is well known that the Boltzmann entropy increases when two or more different substances are mixed. A simple interpretation is that the uncertainty on (or the complexity of) the mixed system increases and consequently

its the entropy. This mixing property directly implies that the entropy is a concave function. If we substitute to the classical entropy a different form which measures differently the amount of information in a system, then the mixing property, though thermodynamically appealing, is not necessarily preserved. This is what happens in the case of our generalized Fisher information. It is known that the standard Fisher information is a convex function [55, 50]. It has also be shown that the Fisher information with $q = 1$ and any exponent $\beta > 1$ is also a convex function [56]. We show here that this is also true for any $q \geq 1$. Consider the integrand in the definition of the Fisher information, and denote $x = |\nabla f|$ and $y = f$. Then, let us define

$$h(x, y) = x^\beta y^{(b-\beta)}, \quad (58)$$

with $b = \beta(q-1) + 1$. This function will be jointly convex, with respect to its two arguments, if the associated Hessian is non negative definite. Accordingly, the integral of h will be a convex functional of f . Hence, we only have to determine the conditions for the non negativity of the Hessian.

By direct computation, we readily obtain that the Hessian matrix for the function $h(x, y)$ is

$$H = \begin{bmatrix} \frac{\partial^2 h}{\partial x^2} & \frac{\partial^2 h}{\partial x \partial y} \\ \frac{\partial^2 h}{\partial y \partial x} & \frac{\partial^2 h}{\partial y^2} \end{bmatrix} = x^{\beta-2} y^{b-\beta-2} \begin{bmatrix} y^2 \beta (\beta - 1) & x y \beta (b - \beta) \\ x y \beta (b - \beta) & x^2 (b - \beta) (b - \beta - 1) \end{bmatrix}. \quad (59)$$

The determinant of the matrix is

$$D = -x^{2\beta-2} y^{2(b-\beta)-2} \beta (b - \beta) (b - 1), \quad (60)$$

and its trace is

$$T = x^{\beta-2} y^{b-\beta-2} (\beta y^2 (\beta - 1) + x^2 (b - \beta) (b - \beta - 1)). \quad (61)$$

Since we know that the determinant and trace of a matrix are respectively the product and sum of the eigenvalues, we see that the two eigenvalues are non negative if both the determinant and trace are non negative, for any x and y . This entails here the conditions

$$\begin{cases} \beta (b - \beta) (b - 1) & \leq 0, \\ \beta (\beta - 1) & \geq 0, \\ (b - \beta) (b - \beta - 1) & \geq 0. \end{cases}$$

Actually, the first two conditions are sufficient since they imply the third one. With $\beta > 0$, we finally obtain $\beta \geq a \geq 1$, which, given that $b = \beta(q - 1) + 1$, leads to

$$2 - \frac{1}{\beta} \geq q \geq 1. \quad (62)$$

Though the generalized Fisher information $\phi_{\beta,q}[f]$ is not a convex function of the probability density function for $q < 1$, we should notice that the ‘thermodynamic’ Fisher information $\phi_{\beta,q}(m_\alpha)$ and $I_{\beta,q}(m_\alpha)$ considered as functions of an observable $m_\alpha = E[|X|^\alpha]$, are convex in m_α for any $q > \max\{(n - 1)/n, n/(n + \alpha)\}$. This is actually an immediate consequence of the Cramér-Rao inequalities (21) and (22). Indeed, any generalized q -Gaussian G_γ with parameter γ reaches the Cramér-Rao bound and we have, for instance for the first Fisher information :

$$m_\alpha [G_\gamma]^\frac{1}{\alpha} \phi_{\beta,q} [G_\gamma]^\frac{1}{\beta\lambda} = m_\alpha [G]^\frac{1}{\alpha} \phi_{\beta,q} [G]^\frac{1}{\beta\lambda}. \quad (63)$$

Choosing γ such that $m_\alpha = m_\alpha [G_\gamma]$, we obtain that the ‘thermodynamic’ Fisher information $\phi_{\beta,q}(m_\alpha)$ is

$$\phi_{\beta,q}(m_\alpha) = K m_\alpha^{-\frac{\beta\lambda}{\alpha}}, \quad (64)$$

with $K = m_\alpha [G]^\frac{\beta\lambda}{\alpha} \phi_{\beta,q} [G]$. Hence, since α , β and λ are positive, $\phi_{\beta,q}(m_\alpha)$ is a convex function of m_α .

3.3. Reciprocity relations and Legendre thermodynamics relationships

In the formulation of standard thermodynamics, the Legendre structure is at heart. We show here that the standard reciprocity relations and Legendre structure of thermodynamics are still valid for generalized Fisher information. The fact that the Legendre structure of thermodynamics is preserved for general maximum-entropy-like problems and is just a consequence of Jaynes’ maximum entropy principle have already be shown in [57], and re-derived in the case of the Fisher information in [50, 54]. We extend this to the case of the generalized Fisher information. Actually, we provide here a simple and self-contained derivation of the fact that the thermodynamics-like Legendre structure is a general and direct consequence of a minimization principle.

We look for the more general distribution, in the Fisher sense, which is compatible with a prior information given as a mean value of some observable. This distribution is selected as the distributions with minimum Fisher

information, and the related variational problem is stated as

$$I_{\beta,q}(m) = \inf_f \{I_{\beta,q}[f] : f \in \mathcal{P}, E[A_i(X)] = m_i, i = 1..M - 1 \text{ and } E[1] = m_0 = 1\}. \quad (65)$$

The problem consists in finding a distribution with minimum Fisher information on the set of all probability distributions with a series of fixed moments $E[A_i(X)] = m_i, i = 0..M - 1$. The value of the information attached to the optimum distribution is denoted $I_{\beta,q}(m)$ – the use of the square brackets and parenthesis distinguishes between the functions of the state and the functions of the observable.

Let λ_i denote the Lagrange multipliers associated with the M constraints, and let us define the dual function $D(\lambda)$ by

$$D(\lambda) := \inf_f \left\{ I_{\beta,q}[f] - \sum_{i=0}^M \lambda_i \left(\int_{\Omega} A_i(x) f(x) dx - m_i \right) \right\}. \quad (66)$$

It is well known that the maximum of the dual function is always less than or equal to the value of the primal problem: $\sup_{\lambda} D(\lambda) \leq I_{\beta,q}(m)$. Clearly, the dual function can be expressed in term of the function $I_{\beta,q}(\lambda)$ defined by

$$I_{\beta,q}(\lambda) := \sup_f \left\{ \sum_{i=0}^M \lambda_i \int_{\Omega} A_i(x) f(x) dx - I_{\beta,q}[f] \right\}, \quad (67)$$

and that gives

$$D(\lambda) = \sum_{i=0}^M \lambda_i m_i - I_{\beta,q}(\lambda). \quad (68)$$

If we denote by \bar{A} the vector of moments $\bar{A}_i = \int_X A_i(x) f(x) dx$, then among all distributions with the same moments \bar{A} , there exists a distribution with minimum Fisher information, say $I_{\beta,q}(\bar{A})$, and this distribution realizes the supremum in (67) for a given moment. Then it remains to explore all the possible means \bar{A} and select the one which maximizes the right hand side of (67).

In other words, if $S_{\bar{A}}$ is the set of distributions with given moments $S_{\bar{A}} = \{f : \bar{A}_i = \int_{\Omega} A_i(x) f(x) dx, i = 0..M - 1\}$, then

$$I_{\beta,q}(\lambda) = \sup_{\bar{A}} \sup_{f \in S_{\bar{A}}} \left\{ \sum_{i=0}^M \lambda_i \int_{\Omega} A_i(x) f(x) dx - I_{\beta,q}[f] \right\}, \quad (69)$$

$$= \sup_{\bar{A}} \left\{ \sum_{i=0}^M \lambda_i \bar{A}_i - I_{\beta,q}(\bar{A}) \right\} \quad (70)$$

where $I_{\beta,q}(\bar{A})$ denotes the Fisher information of the distribution which realizes the supremum in (67) for a given moment. These simple relations directly imply the so-called reciprocity relations. Indeed, the maximum in (70) is attained for a value \bar{A}_λ such that

$$\nabla_{\bar{A}} I_{\beta,q}(\bar{A}_\lambda) = \lambda. \quad (71)$$

On the other hand, (70) then gives

$$I_{\beta,q}(\lambda) = \sum_{i=0}^M \lambda_i \bar{A}_{i\lambda} - I_{\beta,q}(\bar{A}_\lambda). \quad (72)$$

Differentiating $I_{\beta,q}(\bar{A}_\lambda)$ with respect to λ and taking into account (71), we obtain

$$\nabla_\lambda I_{\beta,q}(\bar{A}_\lambda) = (\nabla_\lambda \otimes \bar{A}_\lambda) \cdot \nabla_{\bar{A}} I_{\beta,q}(\bar{A}_\lambda) = (\nabla_\lambda \otimes \bar{A}_\lambda) \cdot \lambda. \quad (73)$$

This result is nothing but a general form of the Euler theorem. Now, the differentiation $I_{\beta,q}(\lambda)$ in (72) with respect to λ gives

$$\nabla_\lambda I_{\beta,q}(\lambda) = \bar{A}_\lambda + (\nabla_\lambda \otimes \bar{A}_\lambda) \cdot \lambda - (\nabla_\lambda \otimes \bar{A}_\lambda) \cdot \nabla_{\bar{A}} I_{\beta,q}(\bar{A}_\lambda) \quad (74)$$

which, taking into account the Euler formula (73) yields the simple relation

$$\nabla_\lambda I_{\beta,q}(\lambda) = \bar{A}_\lambda. \quad (75)$$

In fact, the relation (70) expresses the fact that $I_{\beta,q}(\lambda)$ is the Legendre transform of $I_{\beta,q}(\bar{A})$, and the relationships (71) and (75) simply express that λ and \bar{A} are the dual variables linked by the induced Legendre structure. Let us nevertheless emphasize here that $I_{\beta,q}(\bar{A})$ is not necessarily the Legendre transform of $I_{\beta,q}(\lambda)$; this will only be true if $I_{\beta,q}(\bar{A})$ is a convex function of \bar{A} .

Finally, observe that when we select the value of the Lagrange parameter λ which maximizes the dual function, we obtain

$$\nabla_\lambda I_{\beta,q}(\lambda) = m. \quad (76)$$

This gives the way to compute a value of λ , and therefore the associated probability distribution, such that the moment constraints $E[A_i] = m_i$, $i = 1..M - 1$ are satisfied.

4. Conclusion

In this paper, we have presented two extended forms of Fisher information that fit well in the nonextensive thermostatics context. Indeed, these Fisher information satisfy several extended properties that interrelate generalized q -Gaussians, the generalized Fisher information and the q -entropies. Among these properties, we have stressed that the generalized q -Gaussians are the solution of variational problems such as the minimization of the generalized Fisher information among distributions with a given q -entropy, or such as the minimization of the generalized Fisher information among distributions with a fixed moment. This complements the known fact that the generalized q -Gaussians maximize the q -entropies subject to a moment constraint. These results recover, as a particular case, known characterizations of the standard Gaussian. We have also introduced several information functionals minimized by the generalized q -Gaussian. In information theory, the important de Bruijn identity links the Fisher information and the derivative of the entropy. We have shown here that this identity can be extended to generalized versions of entropy and Fisher information. More precisely, we have shown that for all distributions satisfying a nonlinear heat equation, then the generalized Fisher information naturally pop up in the expression of the derivative of the entropy. In a second step, we have examined further properties of the generalized Fisher information and of their minimization. In particular, we have considered the combination of two independent systems and shown that, though non additive, the generalized Fisher information of the combined system is upper bounded. In the case of mixing, we have shown that the generalized Fisher information is convex for $q \geq 1$. Finally, we have shown that the minimization of the generalized Fisher information subject to moment constraints leads to a Legendre structure analog to the Legendre structure of thermodynamics.

Further work should examine whether a principle of minimum generalized Fisher information, together with the underlying Legendre structure, could lead to interesting predictions. It would also be interesting to look for the uncertainty relationships that could be derived from the generalized Cramér-Rao and Stam inequalities. Whatever the outcome, it is clear that the two generalized Fisher information appear as useful companions to the q -entropies, as demonstrated eg. by de Bruijn identity. The interplay between the extended q -entropies and generalized (β, q) -Fisher information can certainly be of value for the analysis and characterization of complex systems, e.g. following [20, 58, 59, 2].

5. Appendix - Information measures of generalized q -Gaussians

Let $g_\gamma(x) = (1 - s\gamma|x|^\alpha)_+^{\frac{\nu}{s}}$, and the corresponding generalized q -Gaussian is obtained as $G_\gamma(x) = g_\gamma(x) / \int g_\gamma(x) dx$, with $\nu = 1$ and $s = q - 1$. In order to express the different information measures of generalized q -Gaussians, we use a closed-form expression given in the next proposition.

Proposition 6. *Let $\alpha, p > 0$, then consider the following quantity:*

$$\mu_{p,\nu} = \int |x|^p (1 - s\gamma|x|^\alpha)_+^{\frac{\nu}{s}} dx$$

By direct calculation in polar coordinates, one gets the formula

$$\begin{aligned} \mu_{p,\nu} = & \frac{2}{\alpha} (\gamma)^{-\frac{p+n}{\alpha}} n \omega_n \times \\ & \begin{cases} (-s)^{-\frac{p+n}{\alpha}} B\left(\frac{p+n}{\alpha}, -\frac{\nu}{s} - \frac{p+n}{\alpha}\right) & \text{for } -\frac{\nu\alpha}{(p+n)} < s < 0 \\ s^{-\frac{p+n}{\alpha}} B\left(\frac{p+n}{\alpha}, \frac{\nu}{s} + 1\right) & \text{for } s > 0 \\ (\nu)^{-\frac{p+n}{\alpha}} \Gamma\left(\frac{p+n}{\alpha}\right) & \text{if } s = 0 \end{cases} \end{aligned} \quad (77)$$

where ω_n is the volume of the n -dimensional ball.

From this expression, we immediately identify that the partition function of a generalized q -Gaussian G_γ with scale factor γ , we have $Z(\gamma) = (\gamma)^{-\frac{n}{\alpha}} \mu_{0,1}$, that is

$$Z(\gamma) = \frac{2}{\alpha} (\gamma)^{-\frac{n}{\alpha}} n \omega_n \times \begin{cases} (1-q)^{-\frac{n}{\alpha}} B\left(\frac{n}{\alpha}, -\frac{1}{q-1} - \frac{n}{\alpha}\right) & \text{for } 1 - \frac{\alpha}{n} < q < 1 \\ (q-1)^{-\frac{n}{\alpha}} B\left(\frac{n}{\alpha}, \frac{1}{q-1} + 1\right) & \text{for } q > 1 \\ \Gamma\left(\frac{n}{\alpha}\right) & \text{if } q = 1. \end{cases} \quad (78)$$

Similarly, we obtain the information generating function

$$M_q[G_\gamma] = \int G_\gamma(x)^q dx = \frac{\int g_\gamma(x)^q dx}{\left(\int g_\gamma(x) dx\right)^q} = \frac{\mu_{0,q}}{(\mu_{0,1})^q}, \quad (79)$$

$$= \frac{\alpha q}{n(q-1) + \alpha q} \times \frac{1}{(\mu_{0,1})^{q-1}} \quad (80)$$

where the second line is obtained by simplification using the properties of the Beta functions, namely $B(x, y+1) = \frac{y}{x+y} B(x, y)$, and $B(x, y-1) = \frac{x+y-1}{y-1} B(x, y)$.

The information generating function, and thus the associated Rényi and Tsallis entropies are finite for $q > \max \{n/(n + \alpha), 1 - \alpha/n\}$.

Likewise, the moment of order p is given by

$$m_p[G_\gamma] = \frac{\int |x|^p g(x) dx}{\int g(x) dx} = \frac{\mu_{p,1}}{\mu_{0,1}}. \quad (81)$$

If $p = \alpha$, by the properties of the Beta functions, the expressions for the moment of order p simplifies into

$$m_\alpha[G_\gamma] = \frac{\mu_{\alpha,1}}{\mu_{0,1}} = \frac{1}{\gamma(\alpha q + q - 1)} \text{ for } q > n/(n + \alpha) \quad (82)$$

Let $b = \beta(q - 1) + 1$. The generalized Fisher information of the generalized Gaussian has the expression

$$\phi_{\beta,q}[G_\gamma] = \frac{(\alpha\gamma)^\beta}{(\mu_{0,1})^b} \int |x|^\alpha (1 - (q - 1)|x|^\alpha)_+^{\frac{b}{(q-1)} - \beta} dx \quad (83)$$

which reduces easily to

$$\phi_{\beta,q}[G_\gamma] = (\alpha\gamma)^\beta \frac{\mu_{\alpha,1}}{\mu_{0,1}} \times \frac{1}{(\mu_{0,1})^{\beta(q-1)}}. \quad (84)$$

The second generalized Fisher information is given by

$$I_{\beta,q}[G_\gamma] = \frac{\phi_{\beta,q}[G_\gamma]}{M_q[G_\gamma]^\beta} = (\alpha\gamma)^\beta \frac{\mu_{\alpha,1}}{\mu_{0,1}} \times \left(\frac{\mu_{0,1}}{\mu_{0,q}}\right)^\beta.$$

Let us finally evaluate the lower bound in the Cramér-Rao inequality (22). We have

$$I_{\beta,q}[G_\gamma]^{\frac{1}{\beta}} m_\alpha[G_\gamma]^{\frac{1}{\alpha}} = \alpha\gamma \left(\frac{\mu_{0,1}}{\mu_{0,q}}\right) \left(\frac{\mu_{\alpha,1}}{\mu_{0,1}}\right)^{\frac{1}{\beta}} \left(\frac{\mu_{\alpha,1}}{\mu_{0,1}}\right)^{\frac{1}{\alpha}} = \alpha\gamma \left(\frac{\mu_{\alpha,1}}{\mu_{0,q}}\right),$$

since $\alpha^{-1} + \beta^{-1} = 1$. By properties of the Beta functions, we obtain that the ratio simplifies to $\mu_{\alpha,1}/\mu_{0,q} = n/(\gamma\alpha)$. Therefore, we get the equality $I_{\beta,q}[G_\gamma]^{\frac{1}{\beta}} m_\alpha[G_\gamma]^{\frac{1}{\alpha}} = I_{\beta,q}[G] m_\alpha[G]^{\frac{1}{\alpha}} = n$.

6. References

- [1] J. Dehesa, P. Sánchez-Moreno, R. J. Yáñez, Cramér-Rao information plane of orthogonal hypergeometric polynomials, *Journal of Computational and Applied Mathematics* 186 (2) (2006) 523–541.

- [2] J. S. Dehesa, S. López-Rosa, B. Olmos, R. J. Yáñez, Fisher information of d -dimensional hydrogenic systems in position and momentum spaces, *Journal of Mathematical Physics* 47 (5) (2006) 052104.
- [3] J. Dehesa, A. Plastino, P. Sánchez-Moreno, C. Vignat, Generalized Cramér-Rao relations for non-relativistic quantum systems, *Applied Mathematics Letters* 25 (11) (2012) 1689–1694.
- [4] T. M. Cover, J. A. Thomas, *Elements of Information Theory*, 2nd Edition, 2nd Edition, Wiley-Interscience, 2006.
- [5] A. Dembo, T. Cover, J. Thomas, Information theoretic inequalities, *IEEE Transactions on Information Theory* 37 (6) (1991) 1501–1518.
- [6] C. Tsallis, *Introduction to Nonextensive Statistical Mechanics*, illustrated edition Edition, Springer, 2009.
- [7] C. Beck, Generalised information and entropy measures in physics, *Contemporary Physics* 50 (4) (2009) 495.
- [8] S. Golomb, The information generating function of a probability distribution, *IEEE Transactions on Information Theory* 12 (1) (1966) 75–77.
- [9] E. Lutz, Anomalous diffusion and Tsallis statistics in an optical lattice, *Physical Review A* 67 (5) (2003) 051402.
- [10] V. Schwämmle, F. D. Nobre, C. Tsallis, q -Gaussians in the porous-medium equation: stability and time evolution, *The European Physical Journal B-Condensed Matter and Complex Systems* 66 (4) (2008) 537–546.
- [11] C. Vignat, A. Plastino, Why is the detection of q -Gaussian behavior such a common occurrence?, *Physica A* 388 (5) (2009) 601–608.
- [12] G. I. Barenblatt, On some unsteady motions of a liquid and a gas in a porous medium, *Prikladnaja Matematika i Mechanika* 16 (1952) 67–78.
- [13] R. E. Pattle, Diffusion from an instantaneous point source with concentration dependent coefficient, *Quart. J. Mech. Appl. Math.* 12 (1959) 407–409.
- [14] M. Del Pino, J. Dolbeault, Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions, *Journal de Mathématiques Pures et Appliquées* 81 (9) (2002) 847–875.

- [15] M. Del Pino, J. Dolbeault, The optimal euclidean L_p -Sobolev logarithmic inequality, *Journal of Functional Analysis* 197 (1) (2003) 151–161.
- [16] D. Cordero-Erausquin, B. Nazaret, C. Villani, A mass-transportation approach to sharp Sobolev and Gagliardo-Nirenberg inequalities, *Advances in Mathematics* 182 (2) (2004) 307–332.
- [17] M. Agueh, Sharp Gagliardo-Nirenberg inequalities via p -Laplacian type equations, *Nonlinear Differential Equations and Applications NoDEA* 15 (2008) 457–472.
- [18] L. Chimento, F. Pennini, A. Plastino, Naudts-like duality and the extreme Fisher information principle, *Physical Review E* 62 (5 Pt B) (2000) 7462–7465.
- [19] M. Casas, L. Chimento, F. Pennini, A. Plastino, A. R. Plastino, Fisher information in a Tsallis non-extensive environment, *Chaos, Solitons, & Fractals* 13 (3) (2002) 451–459.
- [20] F. Pennini, A. R. Plastino, A. Plastino, Rényi entropies and Fisher informations as measures of nonextensivity in a Tsallis setting, *Physica A* 258 (3-4) (1998) 446–457.
- [21] F. Pennini, A. Plastino, G. Ferri, Semiclassical information from deformed and escort information measures, *Physica A* 383 (2) (2007) 782–796.
- [22] J. Naudts, Generalised exponential families and associated entropy functions, *Entropy* 10 (3) (2008) 131–149.
- [23] J. Naudts, The q -exponential family in statistical physics, *Central European Journal of Physics* 7 (3) (2009) 405–413.
- [24] S. Furuichi, On the maximum entropy principle and the minimization of the Fisher information in Tsallis statistics, *Journal of Mathematical Physics* 50 (1) (2009) 013303–12.
- [25] S. Furuichi, On generalized Fisher informations and Cramér-Rao type inequalities, *Journal of Physics: Conference Series* 201 (2010) 012016.
- [26] J.-F. Bercher, On generalized Cramér-Rao inequalities, generalized Fisher information and characterizations of generalized q -Gaussian distributions, *Journal of Physics A: Mathematical and Theoretical* 45 (25) (2012) 255303.

- [27] J.-F. Bercher, On generalized Cramér-Rao inequalities and characterizations of generalized q -Gaussian distributions: the multidimensional case, Submitted, available at arXiv:1211.2008.
- [28] E. Lutwak, D. Yang, G. Zhang, Cramér-Rao and moment-entropy inequalities for Rényi entropy and generalized Fisher information, *IEEE Transactions on Information Theory* 51 (2) (2005) 473–478.
- [29] E. Lutwak, S. Lv, D. Yang, G. Zhang, Extensions of Fisher information and Stam’s inequality, *IEEE Transactions on Information Theory* 58 (3) (2012) 1319 –1327.
- [30] J.-F. Bercher, On a (β, q) -generalized Fisher information and inequalities involving q -Gaussian distributions, *Journal of Mathematical Physics* 53 (6) (2012) 063303.
- [31] O. Johnson, C. Vignat, Some results concerning maximum Rényi entropy distributions (2007) 339–351.
- [32] L. Pardo, Order- α weighted information energy, *Information sciences* 40 (2) (1986) 155–164.
- [33] O. Onicescu, Energie informationelle, *Comptes Rendus de la Académie des Sciences Paris, Ser. A* 263 (1966) 841.
- [34] A. Nachman, A. Callegari, A nonlinear singular boundary value problem in the theory of pseudoplastic fluids, *SIAM Journal on Applied Mathematics* 38 (2) (1980) 275–281.
- [35] A. Stam, Some inequalities satisfied by the quantities of information of Fisher and Shannon, *Information and Control* 2 (2) (1959) 101–112.
- [36] D. V. Widder, *The Heat Equation*, Academic Press, 1975.
- [37] A. Ohara, Geometric study for the Legendre duality of generalized entropies and its application to the porous medium equation, *The European Physical Journal B* 70 (1) (2009) 15–28.
- [38] J. L. Vázquez, *The Porous Medium Equation: Mathematical Theory*, 1st Edition, Oxford University Press, USA, 2006.
- [39] J. L. Vázquez, *Smoothing and Decay Estimates for Nonlinear Diffusion Equations: Equations of Porous Medium Type*, Oxford University Press, USA, 2006.

- [40] P. Lindqvist, Notes on the p-Laplace equation, Vol. 102, University of Jyväskylä, 2006.
- [41] G. Savaré, G. Toscani, The concavity of Rényi entropy power, arXiv:1208.1035.
- [42] J. Naudts, Generalised Thermostatistics, 1st Edition, Springer, 2011.
- [43] A. G. Bashkirov, Rényi entropy as a statistical entropy for complex systems, Theoretical and Mathematical Physics 149 (2) (2006) 1559–1573.
- [44] P. Jizba, T. Arimitsu, The world according to Rényi: thermodynamics of multifractal systems, Annals of Physics 312 (1) (2004) 17–59.
- [45] E. Lenzi, R. Mendes, L. da Silva, Statistical mechanics based on Rényi entropy, Physica A: Statistical Mechanics and its Applications 280 (3-4) (2000) 337–345.
- [46] A. Parvan, T. Biró, Rényi statistics in equilibrium statistical mechanics, Physics Letters A 374 (19-20) (2010) 1951–1957.
- [47] T. Frank, A. Plastino, Generalized thermostatistics based on the Sharma-Mittal entropy and escort mean values, The European Physical Journal B - Condensed Matter and Complex Systems 30 (4) (2002) 543–549.
- [48] C. Beck, E. G. D. Cohen, Superstatistics, Physica A 322 (2003) 267–275.
- [49] C. Tsallis, A. M. C. Souza, Constructing a statistical mechanics for Beck-Cohen superstatistics, Physical Review E 67 (2) (2003) 026106.
- [50] B. R. Frieden, A. Plastino, A. R. Plastino, B. H. Soffer, Fisher-based thermodynamics: Its Legendre transform and concavity properties, Physical Review E 60 (1) (1999) 48–53.
- [51] B. R. Frieden, P. M. Binder, Physics from Fisher information: A unification, American Journal of Physics 68 (11) (2000) 1064.
- [52] B. R. Frieden, A. Plastino, A. R. Plastino, B. H. Soffer, Non-equilibrium thermodynamics and Fisher information: An illustrative example, Physics Letters A 304 (3-4) (2002) 73–78.
- [53] B. R. Frieden, Science from Fisher Information: A Unification, Cambridge University Press, 2004.

- [54] A. Plastino, A. Plastino, M. Casas, Fisher variational principle and thermodynamics, in: S. Sieniutycz, H. Farkas (Eds.), *Variational and Extremum Principles in Macroscopic Systems*, Elsevier, Oxford, 2005, pp. 379–394.
- [55] M. Cohen, The Fisher information and convexity, *IEEE Transactions on Information Theory* 14 (4) (1968) 591–592.
- [56] D. Boekee, An extension of the Fisher information measure, in: I. Csiszár, P. Elias (Eds.), *Topics in Information Theory*, Vol. 16, János Bolyai Mathematical Society and North-Holland, Keszthely, Hungary, 1977, pp. 113–123.
- [57] A. Plastino, On the universality of thermodynamics' Legendre transform structure, *Physics Letters A* 226 (5) (1997) 257–263.
- [58] C. Vignat, J. F. Bercher, Analysis of signals in the Fisher-Shannon information plane, *Physics Letters A* 312 (1-2) (2003) 27–33.
- [59] E. Romera, J. S. Dehesa, The Fisher-Shannon information plane, an electron correlation tool, *The Journal of Chemical Physics* 120 (19) (2004) 8906–12.