



HAL
open science

The mod 2 cohomology rings of SL_2 of the imaginary quadratic integers

Ethan Berkove, Alexander D. Rahm

► **To cite this version:**

Ethan Berkove, Alexander D. Rahm. The mod 2 cohomology rings of SL_2 of the imaginary quadratic integers. 2014. hal-00769261v3

HAL Id: hal-00769261

<https://hal.science/hal-00769261v3>

Preprint submitted on 8 Oct 2014 (v3), last revised 20 Aug 2016 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE MOD 2 COHOMOLOGY RINGS OF SL_2 OF THE IMAGINARY QUADRATIC INTEGERS

ETHAN BERKOVE AND ALEXANDER D. RAHM
(WITH AN APPENDIX BY AUREL PAGE)

ABSTRACT. We establish general dimension formulae for the second page of the equivariant spectral sequence of the action of the SL_2 groups over imaginary quadratic integers on their associated symmetric space. On the way, we extend the torsion subcomplex reduction technique to cases where the kernel of the group action is nontrivial. Using the equivariant and Lyndon–Hochschild–Serre spectral sequences, we investigate the second page differentials and show how to obtain the mod 2 cohomology rings of our groups from this information.

1. INTRODUCTION

The objects of study in this paper are the groups SL_2 over the ring \mathcal{O}_{-m} of integers in the imaginary quadratic number field $\mathbb{Q}(\sqrt{-m})$, with m a square-free positive integer. These groups, as well as their central quotients $PSL_2(\mathcal{O}_{-m})$, are known as Bianchi groups. The determination of the (co)homology of Bianchi groups, motivated in [19], has a long history of case-by-case computations (see [16] for a list of references). This changed with the recently introduced technique of torsion subcomplex reduction, which provided general formulae for the Farrell cohomology of the $PSL_2(\mathcal{O}_{-m})$ groups for any m [14]. In this article, we extend the subcomplex reduction technique to the case where the kernel of a group action is nontrivial in order to obtain the mod 2 cohomology rings of the $SL_2(\mathcal{O}_{-m})$ groups. For this purpose, we study a variant of the long exact sequence in Borel cohomology. The reason why we consider only \mathbb{F}_2 -coefficients is that for any prime $\ell > 2$, the Lyndon–Hochschild–Serre spectral sequence with \mathbb{F}_ℓ -coefficients associated to the central extension

$$1 \rightarrow \{\pm 1\} \rightarrow SL_2(\mathcal{O}_{-m}) \rightarrow PSL_2(\mathcal{O}_{-m}) \rightarrow 1$$

is concentrated in the horizontal axis, yielding $H^*(SL_2(\mathcal{O}_{-m}); \mathbb{F}_\ell) = H^*(PSL_2(\mathcal{O}_{-m}); \mathbb{F}_\ell)$. Furthermore, Bianchi groups contain only 2- and 3-torsion; results for $H^*(PSL_2(\mathcal{O}_{-m}); \mathbb{F}_3)$ can be found in [14].

Philosophically, the cohomology of the Bianchi groups can be thought of as coming from two sources. Since the Bianchi groups act on a 2-dimensional retract of hyperbolic 3-space X , the Bianchi groups have virtual (co)homological dimension 2. Consequently, in dimensions 2 and below, many of the cohomology classes of the Bianchi groups come from the topology of the quotient space and are detected with rational coefficients [18]; calculations exclusively for rational coefficients have been carried out in [22]. Above the virtual cohomological dimension, all cohomology classes are torsion and originate from finite subgroups. In particular, one can use the equivariant spectral sequence to determine the (co)homology of Bianchi groups. Results of the second author have made this precise. The article [13] introduced the ℓ -torsion subcomplex, and contains a proof that this subcomplex determines the homology of $PSL_2(\mathcal{O}_{-m})$ above dimension 2; in addition, that the homology is a direct sum of generic modules associated to the homeomorphism types of connected components of the subcomplex.

In this work, we will show how to extend the torsion subcomplex results from the projective special linear group to the linear group. The difficulty that we need to overcome is that the kernel of the action of $SL_2(\mathcal{O}_{-m})$ on X is nontrivial. We solve this problem by using a technique reminiscent of the long exact sequence in relative Borel cohomology associated to a pair

Date: October 8, 2014.

2000 Mathematics Subject Classification. 11F75, Cohomology of arithmetic groups.

The second author is funded by the Irish Research Council.

of complexes. This is the content of Section 5, where we describe the E_2 page of the equivariant spectral sequence in terms of components of the 2-torsion subcomplex. We also need to determine whether any loops in the subcomplexes are homologous in the quotient $\mathrm{SL}_2(\mathcal{O}_{-m}) \backslash X$. This interaction is tracked by the variable c in our calculations. The last piece of the puzzle in the determination of the mod 2 cohomology of the SL_2 Bianchi groups is the rank of the second page differential. We can say a lot about this rank by individually analyzing component types of the reduced 2-torsion subcomplex using the cohomology ring structure. This is the subject of Section 7, where we work with both the equivariant and Lyndon–Hochschild–Serre spectral sequences.

We conclude with some sample calculations. For instance, Example (ι) in Section 8 considers the case when $m \equiv 3 \pmod{8}$, $\mathbb{Q}(\sqrt{-m})$ has precisely one finite ramification place over \mathbb{Q} , and the ideal class number of the totally real number field $\mathbb{Q}(\sqrt{m})$ is 1. These three conditions are equivalent to the quotient of the 2-torsion subcomplex having the shape $\bullet \rightarrow \bullet$, as worked out in [14]. Under these assumptions, our cohomology ring has the following dimensions:

$$\dim_{\mathbb{F}_2} \mathrm{H}^q(\mathrm{SL}_2(\mathcal{O}_{-m}); \mathbb{F}_2) = \begin{cases} \beta^1 + \beta^2, & q = 4k + 5, \\ \beta^1 + \beta^2 + 2, & q = 4k + 4, \\ \beta^1 + \beta^2 + 3, & q = 4k + 3, \\ \beta^1 + \beta^2 + 1, & q = 4k + 2, \\ \beta^1, & q = 1, \end{cases}$$

where $\beta^q := \dim_{\mathbb{F}_2} \mathrm{H}^q(\mathrm{SL}_2(\mathcal{O}_{-m}) \backslash X; \mathbb{F}_2)$. Let $\beta_1 := \dim_{\mathbb{Q}} \mathrm{H}_1(\mathrm{SL}_2(\mathcal{O}_{-m}) \backslash X; \mathbb{Q})$. For all absolute values of the discriminant less than 296, numerical calculations yield $\beta^2 + 1 = \beta^1 = \beta_1$. In this range, the numbers m subject to the above dimension formula and β_1 are given as follows (the Betti numbers are taken from [16]).

m	11	19	43	59	67	83	107	131	139	163	179	211	227	251	283
β_1	1	1	2	4	3	5	6	8	7	7	10	10	12	14	13

We add a few remarks about the approach taken in this paper. The finite subgroups we consider have no subgroups isomorphic to $\mathbb{Z}/2 \oplus \mathbb{Z}/2$, so their cohomology is periodic of period 4. Furthermore, the restriction map on cohomology from any finite subgroup to the central $\mathbb{Z}/2$ subgroup is onto in dimensions divisible by 4. We expect something similar for the Bianchi groups we consider, since by results of Quillen [11] their mod-2 cohomology is F -isomorphic to the cohomology of the maximal abelian subgroups, here the center $\mathbb{Z}/2$. We show this in Proposition 12, where we identify a class α in dimension 4 which provides the periodicity. We thank the referee for pointing out that this class α is the second Chern class of the natural representation of $\mathrm{SL}_2(\mathbb{C})$.

Under these observations, one might expect that a reasonable alternative approach to calculating cohomology would come from using the Lyndon–Hochschild–Serre spectral sequence associated to the short exact sequence

$$1 \rightarrow \{\pm 1\} \rightarrow \mathrm{SL}_2(\mathcal{O}_{-m}) \rightarrow \mathrm{PSL}_2(\mathcal{O}_{-m}) \rightarrow 1$$

to determine cohomology up to degree 5. Unfortunately, although one can often easily determine d_2 and d_3 for this spectral sequence, calculations in an unpublished manuscript by the first author yield that in cases with discriminant as low as $m = 3$ and 11, there is a non-trivial internal d_4 differential whose existence can only be determined by knowing $\mathrm{H}^*(\mathrm{SL}_2(\mathcal{O}_{-m}))$ beforehand. For this reason, an approach relying exclusively on this spectral sequence does not appear feasible.

This paper grew out of an attempt to take advantage of much of what is already known about the $\mathrm{PSL}_2(\mathcal{O}_{-m})$ Bianchi groups. Results in [13], for example, show that information about $\mathrm{H}^*(\mathrm{PSL}_2(\mathcal{O}_{-m}))$ above the cohomological dimension resides in “geometric” data. Specifically, for a large range of values of m , the geometric data resides in four types of subcomplexes in the quotient $\mathrm{SL}_2(\mathcal{O}_{-m}) \backslash X$ whose multiplicity we are going to denote by o , ι , θ , and ρ . Other calculations exist for the rational homology of $\mathrm{SL}_2(\mathcal{O}_{-m}) \backslash X$ ([22], for example) which provide values of the Betti numbers β_1 and β_2 for the quotient space. Our work shows that these data are almost sufficient for a full answer, and give very tight bounds on the cohomology of

$SL_2(\mathcal{O}_{-m})$ in all dimensions. We note that prior machine calculations for $H^*(SL_2(\mathcal{O}_{-m}))$ have been performed for many values of m , but the time to complete a calculation usually increases with m . Consequently, the results in this paper provide a constructive argument which can be used to complement, corroborate, and extend existing results.

Acknowledgments. The first author thanks the De Brún Centre for its hospitality and for funding a stay at NUI Galway devoted to the present work. We are grateful for helpful comments by Matthias Wendt on the core of our long exact sequence analysis, and for valuable assistance by Graham Ellis and Tuan Anh Bui on importing our cell complexes into HAP [6] for the example calculations. Of special note, Tuan Anh Bui's resolutions for the cusp stabilizer groups have been of great help. We would even more like to thank the anonymous referee, who provided an approach which streamlined a number of our original arguments, and whose careful and thoughtful comments helped to greatly improve the overall quality of this paper.

2. SPECTRAL SEQUENCES AND CENTRAL EXTENSIONS

The calculations in this paper involve two spectral sequences. We use the Lyndon–Hochschild–Serre spectral sequence to determine cohomology of groups via group extensions. We also use the equivariant spectral sequence since the Bianchi groups act cellularly on low-dimensional contractible complexes. We introduce both spectral sequences briefly in this section; more details can be found in [1], [4], and [9].

For the development of the Lyndon–Hochschild–Serre spectral sequence we follow the approach in [1]. Given a short exact sequence of groups

$$(1) \quad 1 \rightarrow H \rightarrow \Gamma \xrightarrow{\pi} \Gamma/H \rightarrow 1,$$

there is an associated fibration of classifying spaces. One can then apply the Leray–Serre spectral sequence ([9], Chapters 5 and 6). This spectral sequence has $E_2^{i,j} \cong H^i(\Gamma/H; H^j(H; M))$ for untwisted coefficients M and converges to $H^{i+j}(\Gamma; M)$. We note that the Leray–Serre spectral sequence can also be developed more algebraically, as in [4], VII.5.

In the short exact sequence of groups in Equation 1, when the normal subgroup H is central in Γ , it is possible to say more. This is a central extension, and in such cases Γ/H acts trivially on the homotopy fiber B_H . Then with field coefficients M , the E_2 -term of the resulting spectral sequence has the form

$$E_2^{i,j} \cong H^i(\Gamma/H; M) \otimes H^j(H; M).$$

For the remainder of this article, unless specified otherwise, we only consider cohomology with (necessarily trivial) \mathbb{F}_2 -coefficients. We omit these coefficients from the notation.

Lemma IV.1.12 in [1] identifies a unique cohomology class k , called the k -invariant, which generates the kernel of $B_\pi^* : H^2(\Gamma/H) \rightarrow H^2(\Gamma)$. The k -invariant is also the cohomology class associated to the extension. The Leray–Serre spectral sequence, of which the Lyndon–Hochschild–Serre spectral sequence is just one example, has many useful properties. It is compatible with cup products, for example.

For the equivariant spectral sequence, we follow the development in [4, Chapter VII]. Let the group Γ act cellularly on a CW-space X in such a way that the stabilizer of any cell fixes that cell pointwise. Consider the *equivariant cohomology groups* $H^*(\Gamma, C^\bullet(X))$ with coefficients in the cellular cochain complex $C^\bullet(X)$; in our setting, our cochains will take on values in \mathbb{F}_2 . One can define these cohomology groups by taking a projective resolution, F , of $\mathbb{F}_2[\Gamma]$ over \mathbb{F}_2 , then setting $H^*(\Gamma, C^\bullet(X)) = H^*(\text{Hom}_{\mathbb{F}_2[\Gamma]}(F, C^\bullet(X)))$. When X is a contractible space, the equivariant cohomology groups can be identified with the cohomology of Γ , as

$$H^*(\Gamma, C^\bullet(X)) \cong H^*(\Gamma, C^\bullet(\text{point})) \cong H^*(\Gamma).$$

Using the horizontal and vertical filtrations of the double complex $\text{Hom}_\Gamma(F, C^\bullet(X))$ we get a spectral sequence with

$$E_1^{i,j} \cong H^j(\Gamma, C^i(X))$$

converging to $H^{i+j}(\Gamma)$. Let $X^{(i)}$ be a set of representatives of i -cells in X and let Γ_σ be the stabilizer in Γ of a cell σ . Then

$$C^i(X) = \prod_{\sigma \in X^{(i)}} \mathbb{F}_2 \cong \prod_{\sigma \in \Gamma \backslash X^{(i)}} \text{Coind}_{\Gamma_\sigma}^{\Gamma} \mathbb{F}_2.$$

Once we apply Shapiro's Lemma and sum over the representatives of i -cells in $\Gamma \backslash X_i$, the spectral sequence takes the form $E_1^{i,j} \cong \prod_{\sigma \in \Gamma \backslash X^{(i)}} H^j(\Gamma_\sigma)$.

The equivariant spectral sequence has a number of additional desirable properties that we will use in the sequel.

- (1) There is a product on the spectral sequence, $E_r^{pq} \otimes E_r^{st} \rightarrow E_r^{p+s, q+t}$, which is compatible with the standard cup product on $H^*(\Gamma)$ [4, VII.5]
- (2) On the E_2 -page, the products in $E_2^{0,*}$, the vertical edge, are compatible with the products in $\prod_{\sigma \in \Gamma \backslash X^{(0)}} H^q(\Gamma_\sigma)$. [4, X.4.5.vi].
- (3) The differential d_1 is a difference of restriction maps (cohomology analog of [4, VII.8])

$$\prod_{\sigma \in \Gamma \backslash X^{(i)}} H^j(\Gamma_\sigma) \xrightarrow{d_1^{i,j}} \prod_{\tau \in \Gamma \backslash X^{(i+1)}} H^j(\Gamma_\tau).$$

Remark 1. In both the equivariant and Lyndon–Hochschild–Serre spectral sequences, the differentials are derivations for $r \geq 2$. Working mod 2, this means that given cohomology classes $u, v \in E_r$, the differential satisfies the Leibniz rule,

$$d_r(uv) = d_r(u)v + ud_r(v).$$

A proof that the differential is a derivation for the Leray Serre spectral sequence can be found in [9] (Section 1.4 and Theorem 2.14). For the equivariant spectral sequence, we lay out a sketch following Brown [4, X.4]. Recall that $H^*(\Gamma, C^\bullet(X)) = H^*(\text{Hom}_\Gamma(F, C^\bullet(X)))$, and note that $F \otimes F$ is also a projective resolution of $\mathbb{F}_2\Gamma$ over \mathbb{F}_2 . Therefore, there is a diagonal approximation $\Delta : F \rightarrow F \otimes F$. Consider the composition

$$\begin{aligned} \text{Hom}_\Gamma(F, C^\bullet(X)) \otimes \text{Hom}_\Gamma(F, C^\bullet(X)) &\longrightarrow \text{Hom}_\Gamma(F \otimes F, C^\bullet(X) \otimes C^\bullet(X)) \\ (2) \quad &\xrightarrow{\Delta} \text{Hom}_\Gamma(F, C^\bullet(X) \otimes C^\bullet(X)) \\ &\xrightarrow{\cup} \text{Hom}_\Gamma(F, C^\bullet(X)) \end{aligned}$$

where \cup is the cochain cup product on $C^\bullet(X)$. The composition respects both the horizontal and vertical filtrations of the double complex $\text{Hom}_\Gamma(F, C^\bullet(X))$. This immediately implies Property 1 for the equivariant spectral sequence. And since d_r is induced from the product on the double complex $\text{Hom}_\Gamma(F, C^\bullet(X))$ which satisfies the Leibniz rule, the differential d_r satisfies the Leibniz rule for $r \geq 2$ as well. We note that in the equivariant spectral sequence, in light of the composition given above, it may be difficult to calculate the products in the derivation. For further details on this construction we direct the reader to Brown [4, X.4]. Having the differential be a derivation is helpful. In particular, it implies two useful results: 1) when $d_r(u) = 0$, then $d_r(uv) = ud_r(v)$; and 2) when u is a class on the E_r page, then $d_r(u^2) = 2d_r(u) = 0$.

Finally, for both spectral sequences, one can define Steenrod operations which are compatible with differentials as well as the Steenrod squares in the abutment. A complete account can be found in [21]. However, we will only need the following facts:

- (1) There are well-defined operations, $Sq^k : E_r^{p,q} \rightarrow E_r^{p, q+k}$ when $0 \leq k \leq q$ [21, Theorem 2.15].
- (2) When $k \leq q - 1$, $d_2 Sq^k u = Sq^k d_2 u$, i.e., the operations commute with the differential [21, Theorem 2.17].
- (3) Sq^k commutes with the appropriate differentials to give the Kudo transgression theorem. [21, Corollary 2.18].

3. THE NON-CENTRAL TORSION SUBCOMPLEX

In this section we recall the ℓ -torsion subcomplexes theory of [14] and compare it with the *non-central* ℓ -torsion subcomplexes we are going to study. We require any discrete group Γ under our study to be provided with what we will call a *polytopal Γ -cell complex*, that is, a finite-dimensional simplicial complex X with cellular Γ -action such that each cell stabilizer fixes its cell point-wise. In practice, we relax the simplicial condition to a polyhedral one, merging finitely many simplices to a suitable polytope. We could obtain the simplicial complex back as a triangulation.

Definition 2. Let ℓ be a prime number. The *ℓ -torsion subcomplex* of a polytopal Γ -cell complex X consists of all the cells of X whose stabilizers in Γ contain elements of order ℓ .

We further require that the fixed point set X^G be acyclic for every nontrivial finite ℓ -subgroup G of Γ . Then Brown's proposition X.(7.2) [4] specializes as follows.

Proposition 3. *There is an isomorphism between the ℓ -primary parts of the Farrell cohomology of Γ and the Γ -equivariant Farrell cohomology of the ℓ -torsion subcomplex.*

For a given Bianchi group, the non-central torsion subcomplex can be quite large. It turns out to be useful to reduce this subcomplex, and we identify two conditions under which we can do this in a way that Proposition 3 still holds.

Condition A. In the ℓ -torsion subcomplex, let σ be a cell of dimension $n - 1$ which lies in the boundary of precisely two n -cells representing different orbits, τ_1 and τ_2 . Assume further that no higher-dimensional cells of the ℓ -torsion subcomplex touch σ ; and that the n -cell stabilizers admit an isomorphism $\Gamma_{\tau_1} \cong \Gamma_{\tau_2}$.

Condition B. The inclusion of the cell stabilizer Γ_σ into Γ_{τ_1} and Γ_{τ_2} induces isomorphisms on mod ℓ cohomology.

When both conditions are satisfied in the ℓ -torsion subcomplex, we merge the cells τ_1 and τ_2 along σ and do so for their entire orbits. The effect of this merging is to decrease the size of the ℓ -torsion subcomplex without changing its Γ -equivariant Farrell cohomology. This process can often be repeated: by a "terminal vertex," we will denote a vertex with no adjacent higher-dimensional cells and precisely one adjacent edge in the quotient space, and by "cutting off" the latter edge, we will mean that we remove the edge together with the terminal vertex from our cell complex.

Definition 4. The *reduced ℓ -torsion subcomplex* associated to a polytopal Γ -cell complex X is the cell complex obtained by recursively merging orbit-wise all the pairs of cells satisfying conditions A and B, and cutting off edges that admit a terminal vertex when condition B is satisfied.

The following theorem, stating that Proposition 3 still holds after reducing, is proved in [14].

Theorem 5. *There is an isomorphism between the ℓ -primary parts of the Farrell cohomology of Γ and the Γ -equivariant Farrell cohomology of the reduced ℓ -torsion subcomplex.*

In the case of a trivial kernel of the action on the polytopal Γ -cell complex, this allows one to establish general formulae for the Farrell cohomology of Γ [14]. In contrast, our action of $SL_2(\mathcal{O}_{-m})$ on hyperbolic 3-space has the 2-torsion group $\{\pm 1\}$ in the kernel; since every cell stabilizer contains 2-torsion, the 2-torsion subcomplex does not ease our calculation in any way. We can remedy this situation by considering the following object, on whose cells we impose a supplementary property.

Definition 6. The *non-central ℓ -torsion subcomplex* of a polytopal Γ -cell complex X consists of all the cells of X whose stabilizers in Γ contain elements of order ℓ that are not in the center of Γ .

We note that this definition yields a correspondence between the *non-central* ℓ -torsion subcomplex for a group acting by conjugation and the ℓ -torsion subcomplex for its central quotient

group. We use this correspondence in order to identify the *non-central* ℓ -torsion subcomplex for the action of $\mathrm{SL}_2(\mathcal{O}_{-m})$ on hyperbolic 3-space as the ℓ -torsion subcomplex of $\mathrm{PSL}_2(\mathcal{O}_{-m})$. However, incorporating the non-central condition for $\mathrm{SL}_2(\mathcal{O}_{-m})$ introduces significant technical obstacles, and we have to analyze a variant of the long exact sequence in Borel cohomology in Section 5 in order to overcome them.

We recall the following information from [14] on the ℓ -torsion subcomplex of $\mathrm{PSL}_2(\mathcal{O}_{-m})$. Let Γ be a finite index subgroup in $\mathrm{PSL}_2(\mathcal{O}_{-m})$. Then any element of Γ fixing a point inside hyperbolic 3-space \mathcal{H} acts as a rotation of finite order. By Felix Klein's work, we know conversely that any torsion element α is elliptic and hence fixes some geodesic line. We call this line *the rotation axis of α* . Every torsion element acts as the stabilizer of a line conjugate to one passing through the Bianchi fundamental polyhedron. We obtain the *refined cellular complex* from the action of Γ on \mathcal{H} as described in [12], namely we subdivide \mathcal{H} until the stabilizer in Γ of any cell σ fixes σ point-wise. We achieve this by computing Bianchi's fundamental polyhedron for the action of Γ , taking as a preliminary set of 2-cells its facets lying on the Euclidean hemispheres and vertical planes of the upper-half space model for \mathcal{H} , and then subdividing along the rotation axes of the elements of Γ . It is well-known [18] that if γ is an element of finite order n in a Bianchi group, then n must be 1, 2, 3, 4 or 6, because γ has eigenvalues ρ and $\bar{\rho}$, with ρ a primitive n -th root of unity, and the trace of γ is $\rho + \bar{\rho} \in \mathcal{O}_{-m} \cap \mathbb{R} = \mathbb{Z}$. For ℓ being one of the two occurring prime numbers 2 and 3, the orbit space of this sub-complex is a graph, because the cells of dimension greater than 1 are trivially stabilized in the refined cellular complex. We can see that this graph is finite either from the finiteness of the Bianchi fundamental polyhedron, or from studying conjugacy classes of finite subgroups as in [8].

As in [17], we make use of a 2-dimensional deformation retract X of the refined cellular complex, equivariant with respect to a Bianchi group $\Gamma := \mathrm{SL}_2(\mathcal{O}_{-m})$. This retract has a cell structure in which each cell stabilizer fixes its cell pointwise. Since X is a deformation retract of \mathcal{H} and hence acyclic, $H_\Gamma^*(X) \cong H_\Gamma^*(\mathcal{H}) \cong H^*(\Gamma)$.

Observation 7. From a case-by-case calculation of roughly 140 Bianchi groups with discriminant absolute value less than 1556, there are four observed non-central reduced 2-torsion subcomplex quotient types. There may exist more types for the Bianchi groups, but of the four known types, only three occur more than twice in our sample (see Appendix A and a few more examples in Figure 3 of [12]).

Joint work in progress of Grant Lakeland with the authors indicates the existence of more component types for congruence subgroups in the Bianchi groups. However, it seems out of reach of current software and hardware to make a substantial enough extension of our sample through direct numerical computation to be able to say more about existence of types for the full Bianchi groups.

A connected component of the 2-torsion subcomplex can be considered as a tree with action of the subgroup G of the Bianchi group that sends the tree to itself. We obtain G as the *groupe fondamentale du graphe de groupes* ([20], not to be confused with the fundamental group of the underlying graph) of a connected component of the quotient of the non-central reduced 2-torsion subcomplex with attached stabilizer groups and monomorphisms from the edge stabilizers into the vertex stabilizers. Such G in this paper can be built iteratively using amalgamated products and HNN extensions. We collect the connected component types that have appeared so far in numerical computations [12] in Table 1. The homotopy type of the quotient space is given in the right-most column. All edges have stabilizer type $\mathbb{Z}/4$, and the stabilizer types of the vertices are given. We note that the degree of each vertex in the quotient space is the same as the number of distinct conjugacy classes of $\mathbb{Z}/4$ in the vertex stabilizer. This information determines the HNN extensions up to ordering of the conjugacy classes. We also observe that all stabilizers which contain a copy of \mathbf{Q}_8 are associated to vertices. This observation will be used in Section 5.

TABLE 1. Connected components of non-central reduced 2-torsion subcomplex quotients

	Type of group G	Quotient of tree acted on by G
(o)	HNN extension $\mathbb{Z}/4 *_{\mathbb{Z}/4}$	$\circlearrowright \mathbb{Z}/4$
(ι)	Amalgamated product $\mathbf{Te} *_{\mathbb{Z}/4} \mathbf{Te}$	$\mathbf{Te} \leftrightarrow \mathbf{Te}$
(θ)	Double HNN extension $((\mathbf{Q}_8 *_{\mathbb{Z}/4} \mathbf{Q}_8) *_{\mathbb{Z}/4}) *_{\mathbb{Z}/4}$	$\mathbf{Q}_8 \Theta \mathbf{Q}_8$
(ρ)	Iterated construction $(\mathbf{Q}_8 *_{\mathbb{Z}/4}) *_{\mathbb{Z}/4} \mathbf{Te}$	$\mathbf{Q}_8 \circlearrowright \mathbf{Te}$

4. MAPS INDUCED BY FINITE SUBGROUPS IN THE BIANCHI GROUPS

In this section, we classify the possible cell stabilizers of the $SL_2(\mathcal{O}_{-m})$ action and determine the restriction maps between subgroups. We will use this information to determine the cohomology of components of the reduced 2-torsion subcomplexes in Section 6

Since the action of $SL_2(\mathcal{O}_{-m})$ on X is properly discontinuous, cell stabilizers are finite subgroups of $SL_2(\mathcal{O}_{-m})$. The enumeration of the finite subgroups of $SL_2(\mathbb{C})$ is a classical result, and the list of finite subgroups which appear in the Bianchi groups $PSL_2(\mathcal{O}_{-m})$ is also well known [7]: They are the cyclic groups 1, $\mathbb{Z}/2$ and $\mathbb{Z}/3$, the Klein four group \mathbf{D}_2 , the symmetric group on three letters \mathbf{S}_3 , and the alternating group on four letters \mathbf{A}_4 . The preimages of these groups in $SL_2(\mathcal{O}_{-m})$ are the cyclic groups $\mathbb{Z}/2$, $\mathbb{Z}/4$, $\mathbb{Z}/6$, the quaternion group \mathbf{Q}_8 of order 8, the dicyclic group \mathbf{Di} of order 12, and the binary tetrahedral group \mathbf{Te} of order 24.

We start by recalling some mod 2 cohomology rings. In what follows, the symbol \mathbb{F}_2 represents the field of two elements, and all the cohomology rings and groups in which we omit the coefficients are meant to be taken with (obviously trivial) \mathbb{F}_2 -coefficients. We write reduced cohomology classes [in square brackets] and nilpotent cohomology classes (in parentheses). The index of a class specifies its degree.

Theorem 8. [1] *The cohomology rings for finite subgroups of Bianchi groups are given below, where a subscript denotes the degree of the generator.*

$$\begin{aligned}
 H^*(\mathbb{Z}/2) &\cong H^*(\mathbf{S}_3) \cong \mathbb{F}_2[x_1] \\
 H^*(\mathbb{Z}/3) &\cong H^*(1) \cong \mathbb{F}_2 \\
 H^*(\mathbf{D}_2) &\cong \mathbb{F}_2[x_1, y_1] \\
 H^*(\mathbf{A}_4) &\cong \mathbb{F}_2[u_2, v_3, w_3] / (u_2^3 + v_3^2 + w_3^2 + v_3 w_3)
 \end{aligned}$$

□

We use the Lyndon–Hochschild–Serre spectral sequence to determine the cohomology rings of the finite subgroups of $SL_2(\mathcal{O}_{-m})$. In particular, for each finite subgroup G of $SL_2(\mathcal{O}_{-m})$, there is a corresponding finite subgroup Q of $PSL_2(\mathcal{O}_{-m})$ which fits into the central extension

$$1 \rightarrow \mathbb{Z}/2 \rightarrow G \rightarrow Q \rightarrow 1$$

where $\mathbb{Z}/2$ is the subgroup of $SL_2(\mathcal{O}_{-m})$ containing $\{\pm I\}$. We note that all finite subgroups of $SL_2(\mathbb{C})$ also sit inside $SU(2)$ and act freely on it. Since $SU(2)$ can be identified with the 3-sphere, the cohomology rings for the finite subgroups of $SL_2(\mathcal{O}_{-m})$ are periodic of period dividing 4 [4]. In particular, the cohomology rings can all be expressed as a tensor product where one term is a polynomial ring on one generator.

Theorem 9. *The following are the mod 2 cohomology rings of the finite subgroups of $\mathrm{SL}_2(\mathcal{O}_{-m})$.*

$$\begin{aligned} \mathrm{H}^*(\mathbb{Z}/4) &\cong \mathrm{H}^*(\mathbf{Di}) \cong \mathbb{F}_2[e_2](b_1) \\ \mathrm{H}^*(\mathbb{Z}/2) &\cong \mathrm{H}^*(\mathbb{Z}/6) \cong \mathbb{F}_2[e_1] \\ \mathrm{H}^*(\mathbf{Q}_8) &\cong \mathbb{F}_2[e_4](x_1, y_1, x_2, y_2, x_3) \\ \mathrm{H}^*(\mathbf{Te}) &\cong \mathbb{F}_2[e_4](b_3) \end{aligned}$$

Proof. The cohomology results for cyclic group results are straightforward, and the calculation of the other cohomology rings (and a classification for all periodic groups) are contained in [1]. However, we briefly review the derivations here as the descriptions will be useful when we determine restriction maps to subgroups.

The dicyclic group \mathbf{Di} is a semidirect product and fits into the short exact sequence

$$1 \rightarrow \mathbb{Z}/3 \rightarrow \mathbb{Z}/3 \rtimes \mathbb{Z}/4 \rightarrow \mathbb{Z}/4 \rightarrow 1.$$

Since $\mathrm{H}^*(\mathbb{Z}/3)$ has trivial mod 2 homology, in the Lyndon–Hochschild–Serre spectral sequence the only nontrivial cohomology occurs on the horizontal axis and is isomorphic to $\mathrm{H}^*(\mathbb{Z}/4)$.

For $\mathbb{Z}/4$, \mathbf{Q}_8 and \mathbf{Te} , we again use the Lyndon–Hochschild–Serre spectral sequence for central extensions. In the spectral sequence of the extension, let $\mathbb{F}_2[e_1]$ be the cohomology ring corresponding to the central subgroup $\mathbb{Z}/2$. On the E_2 page of the spectral sequence, the image of $d_2(e_1)$ can be identified with the k -invariant. In addition, the Kudo transgression theorem describes the images of $d_{2^n+1}((e_1)^{2^n})$.

We start with the calculation of $\mathrm{H}^*(\mathbb{Z}/4)$. Let $\mathrm{H}^*(\mathbb{Z}/2) \cong \mathbb{F}_2[x_1]$, and consider the spectral sequence associated to the central extension. In this case, the k -invariant $d_2(e_1) = x_1^2$, so $d_2(e_1 x_i^k) = x_i^{k+2}$ and $d_2(e_1^2) = 0$. As the result, the spectral sequence collapses at the E_2 page with classes in even rows and in columns 0 and 1 only. The class e_1^2 represents the polynomial class in $\mathrm{H}^2(\mathbb{Z}/4)$, and the class in $E_2^{0,1}$ is the one-dimensional exterior class.

We next calculate $\mathrm{H}^*(\mathbf{Q}_8)$. The extension is central, so $E_1^{p,q} \cong \mathrm{H}^p(\mathbf{D}_2) \otimes \mathrm{H}^q(\mathbb{Z}/2)$, and by [1, Proposition IV.2.10], the k -invariant $d_2(e_1) = x_1^2 + x_1 y_1 + y_1^2 \in \mathrm{H}^*(\mathbf{D}_2)$. Also, $d_2(e_1^2) = 0$, which completely determines the differentials on the E_2 page. Therefore, the only classes which survive to the E_3 page lie in even rows. Next, using the Kudo transgression theorem,

$$d_3(e_1^2) = d_3(Sq^1(e_1)) = Sq^1(x_1^2 + x_1 y_1 + y_1^2) = x_1^2 y_1 + x_1 y_1^2.$$

Through careful accounting, one can show that the classes that remain in E_3 are in columns 0 through 3, and that the spectral sequence has non-zero terms in rows congruent to 0 mod 4. Consequently, all higher differentials vanish and $E_3 \cong E_\infty$. The only classes left correspond to 1, $x_1, y_1, x_1^2, y_1^2, x_1^2 y_1 = x_1 y_1^2$ and the product of these classes with powers of e_1^4 , which represents a four-dimensional polynomial class. We find that as a ring, $\mathrm{H}^*(\mathbf{Q}_8) \cong \mathbb{F}_2[e_4](x_1, y_1) / \langle R \rangle$, where R is the ideal generated by $x_1^2 + x_1 y_1 + y_1^2$ and $x_1^2 y_1 + x_1 y_1^2$. The result follows.

Finally, we consider the calculation of $\mathrm{H}^*(\mathbf{Te})$, which proceeds in a similar manner as the case for $\mathrm{H}^*(\mathbf{Q}_8)$, except now the quotient group is \mathbf{A}_4 and we have to consider

$$\mathrm{H}^*(\mathbf{A}_4) \cong \mathbb{F}_2[u_2, v_3, w_3] / (u_2^3 + v_3^2 + w_3^2 + v_3 w_3).$$

Briefly, $d_2(e_1) = u_2$, and $d_3(e_1^2) = v_3 = Sq^1(u_2)$ [3]. Once again, after a careful check, the spectral sequence collapses at the E_3 page and the only classes left correspond to 1, the four-dimensional class represented by e_1^4 , $w_3 \in E_3^{3,0}$, and the product of w_3 with powers of e_1^4 . \square

Given the following commuting diagram of groups,

$$\begin{array}{ccccc} \mathbb{Z}/2 & \longrightarrow & G_1 & \longrightarrow & Q_1 \\ & & \parallel & & \downarrow i_2 \\ & & \parallel & & \downarrow i_1 \\ \mathbb{Z}/2 & \longrightarrow & G_2 & \longrightarrow & Q_2 \end{array}$$

if one knows the induced map $i_1^* : \mathrm{H}^*(Q_2) \rightarrow \mathrm{H}^*(Q_1)$, it is often possible to calculate the effect of $i_2^* : \mathrm{H}^*(G_2) \rightarrow \mathrm{H}^*(G_1)$. The commutative diagram gives rise to two Lyndon–Hochschild–Serre spectral sequences, one for each extension, and the two are compatible via the isomorphism on the fiber. Specifically, the maps between the spectral sequences are given by i_1^* along the

x -axis and the identity along the y -axis. Once the spectral sequences converge, for the cases we consider in this work the result gives the effect of the restriction map $res_{G_1}^{G_2} = i_2^*$ for classes on the edges.

Theorem 10. *The following are the nontrivial restriction maps on cohomology generators for finite subgroups of $SL_2(\mathcal{O}_{-m})$:*

$$\begin{aligned} res_{\mathbb{Z}/2}^{\mathbb{Z}/4}(e_2) &= e_1^2 \\ res_{\mathbb{Z}/6}^{\mathbf{Di}}(e_2) &= res_{\mathbb{Z}/2}^{\mathbf{Di}}(e_2) = e_1^2 \\ res_{\mathbb{Z}/4}^{\mathbf{Q}_8}(e_4) &= e_2^2, res_{\mathbb{Z}/4}^{\mathbf{Q}_8}(x_1) = (b_1) \\ res_{\mathbb{Z}/2}^{\mathbf{Q}_8}(e_4) &= e_1^4 \\ res_{\mathbb{Z}/6}^{\mathbf{Te}}(e_4) &= res_{\mathbb{Z}/2}^{\mathbf{Te}}(e_4) = e_1^4 \\ res_{\mathbb{Z}/4}^{\mathbf{Te}}(e_4) &= e_2^2 \end{aligned}$$

In addition, $res_{\mathbb{Z}/2}^{\mathbb{Z}/6}$ and $res_{\mathbb{Z}/4}^{\mathbf{Di}}$ are isomorphisms.

Proof. The isomorphism cases follow as $\mathbb{Z}/6 \cong \mathbb{Z}/2 \oplus \mathbb{Z}/3$ and $H^*(S_3) \cong H^*(\mathbb{Z}/2)$. For the other restriction maps, we can prove the results concerning the polynomial classes using a Lyndon–Hochschild–Serre spectral sequence calculation in cohomology associated to the central extension

$$1 \rightarrow \mathbb{Z}/2 \rightarrow G \rightarrow Q \rightarrow 1.$$

In every calculation, the polynomial class of $H^*(G)$ arises from the 0 column of the spectral sequence, so it corresponds to some power of e_1^k . The choice of k invariant of the extension determines when d_{2^n+1} vanishes on the vertical axis; at this point, $(e_1)^{2^n}$ survives to E_∞ and represents the polynomial class in the abutment. This observation determines the restriction maps on polynomial classes since the map between spectral sequences on the fiber (the y -axis) is the identity.

For $res_{\mathbb{Z}/4}^{\mathbf{Q}_8}(x_1)$, write $H^*(\mathbf{D}_2) = \mathbb{F}_2[x_1, y_1]$ and $H^*(\mathbb{Z}/2) = \mathbb{F}_2[e_1]$. Using the commutative diagram above with $Q_1 \cong \mathbb{Z}/2$ and $Q_2 \cong \mathbf{D}_2$, without loss of generality we have $i_1^*(x_1^j) = res_{\mathbb{Z}/2}^{\mathbf{D}_2}(x_1^j) = e_1^j$ and $i_1^*(x_1^j y_1^k) = res_{\mathbb{Z}/2}^{\mathbf{D}_2}(x_1^j y_1^k) = 0$ when $k > 0$. After the Lyndon–Hochschild–Serre spectral sequence is applied, $res_{\mathbb{Z}/4}^{\mathbf{Q}_8}(x_1)$ surjects onto the stated class in $H^1(\mathbb{Z}/4)$. We note that the other nilpotent classes of $H^*(\mathbf{Q}_8)$ in dimensions 2 and 3 must restrict to 0, since from Theorem 9, the E_2 page for the calculation of $H^*(\mathbb{Z}/4)$ is zero in the second column and to the right.

Finally, we need to show that $res_{\mathbb{Z}/4}^{\mathbf{Te}}(b_3) = 0$. We recall from the calculation of $H^*(\mathbf{Te})$ in Theorem 9 that the exterior class b_3 is in $E_\infty^{3,0}$. However, in the calculation of $H^*(\mathbb{Z}/4)$, $E_\infty^{i,j} = 0$ when $i > 1$. Therefore, the restriction map must send b_3 to 0. \square

Remark 11. We note that the restriction map is natural with respect to cup products, so the results of Theorem 10 can be used to determine the effect of the restriction map on other cohomology classes. In particular, the only nontrivial restriction map on classes with nilpotent components is $res_{\mathbb{Z}/4}^{\mathbf{Q}_8}(e_4^i x_1) = e_2^{2i} b_1$. In addition, in the cohomology of the groups we consider, only polynomial classes restrict nontrivially to polynomial classes.

We noted in the discussion after Theorem 8 that the cohomology of the finite subgroups of $SL_2(\mathcal{O}_{-m})$ are all periodic of period dividing 4. Above the virtual cohomological dimension, which is 2 for the Bianchi groups, the same period can be observed for the SL_2 Bianchi groups and their subgroups. The proof of this is based on ideas of [2] and [5].

Proposition 12. *Any (not necessarily finite) subgroup Γ in $SL_2(\mathcal{O}_{-m})$ has periodic cohomology above the virtual cohomological dimension. The periodicity can be realized by cup product with a 4-dimensional class α .*

Proof. The Bianchi groups act properly on X , which is homeomorphic to \mathbb{R}^3 . In addition, from the discussion after Theorem 8, we know that the finite subgroups in the Bianchi groups act

freely on $SU(2) \subset SL_2(\mathbb{C})$. Therefore the groups $SL_2(\mathcal{O}_{-m})$ and their subgroups act freely and properly on $\mathbb{R}^3 \times \mathbb{S}^3$.

We essentially follow the argument in [5] and set $Y := \mathbb{R}^3 \times \mathbb{S}^3$ with the Γ -action described above. There is a fibration

$$Y \rightarrow E\Gamma \times_{\Gamma} Y \rightarrow B\Gamma,$$

where $B\Gamma$ is the classifying space for Γ and $E\Gamma$ is its universal cover. Since Y has the homotopy type of \mathbb{S}^3 , this is a spherical fibration which is orientable as we are working with \mathbb{F}_2 -coefficients. The result follows by applying the Gysin exact sequence.

The class α is the image under d_4 of the generator of $H^3(\mathbb{S}^3)$. As mentioned in the introduction, the referee has pointed out that this class is the second Chern class of the natural representation of $SL_2(\mathbb{C})$. \square

5. CALCULATION OF THE E_2 PAGE

In this section, we describe the E_2 page of the equivariant spectral sequence for Bianchi groups in a fairly general way. Some of our analyses will hinge on distinguishing types of cohomology classes. Given a finite group G , we note that the nilpotent classes in $H^*(G)$ form an ideal. We will denote this ideal by $H_{\text{nil}}^*(G)$. We next define the reduced submodule $H_{\text{red}}^*(G)$ as a set of representatives for the quotient by the ideal of nilpotent classes, so

$$0 \rightarrow H_{\text{nil}}^*(G) \rightarrow H^*(G) \rightarrow H_{\text{red}}^*(G) \rightarrow 0$$

is exact.

Denote by X the cell complex described in Section 3 before Observation 7, and by X_s the non-central 2-torsion subcomplex of X with respect to Γ . Further, denote by X'_s the subcomplex of X_s consisting of cells whose stabilizer group contains a copy of \mathbf{Q}_8 , i.e. the stabilizer being of type either \mathbf{Q}_8 or \mathbf{Te} . We note that if a cell of X is not in X'_s , then by Theorem 9 the cohomology of that cell's stabilizer is isomorphic to $H^*(\mathbb{Z}/4)$ if the cell is in X_s , respectively to $H^*(\mathbb{Z}/2)$ if the cell is not in X_s . We also note that X'_s is a 0-dimensional subcomplex of X .

Note 13. In the calculations that follow, we require that for any cell $\tau \in X_s$, the inclusion of the center $Z(\Gamma) \hookrightarrow \Gamma_{\tau}$ into the cell stabilizer of τ induces a monomorphism on the reduced parts of the cohomology rings, $H_{\text{red}}^*(\Gamma_{\tau}) \hookrightarrow H_{\text{red}}^*(Z(\Gamma))$. Furthermore, for any cell $\sigma \in X$ not in X_s , we want that the inclusion $Z(\Gamma) \hookrightarrow \Gamma_{\sigma}$ induces an isomorphism $H_{\text{red}}^*(\Gamma_{\sigma}) \rightarrow H_{\text{red}}^*(Z(\Gamma))$. By Theorem 10, the action of the Bianchi groups on the cell complex X obtained from hyperbolic space satisfies these conditions.

In the following equivariant spectral sequence material, we will assume that we are working with Γ -equivariant cohomology, i.e., $E_2^{*,q}(Y)$ stands for the $E_2^{*,q}$ -term of the equivariant spectral sequence associated to the action of Γ on Y , unless specified otherwise.

For the next theorem, we will show how to write each row in the equivariant spectral sequence converging to $H_{\Gamma}^*(X)$ as the direct sum of terms from X_s and X'_s with terms from a relative spectral sequence, which is defined as follows. Given a cellular subcomplex $X' \subseteq X$ there is a short exact sequence of cochain complexes

$$0 \rightarrow C^*(X, X') \rightarrow C^*(X) \rightarrow C^*(X') \rightarrow 0.$$

Let F be a free resolution for Γ . Applying $\text{Hom}_{\mathbb{F}_2}(F, -)$ and then cohomology yields a long exact sequence

$$(3) \quad \dots \rightarrow E_1^{*,q}(X, X') \rightarrow E_1^{*,q}(X) \rightarrow E_1^{*,q}(X') \xrightarrow{\delta} E_1^{*+1,q}(X, X') \rightarrow \dots$$

Theorem 14. *In the equivariant spectral sequence with \mathbb{F}_2 -coefficients converging to $H_{\Gamma}^*(X)$, the E_2 page is given by the following rows, k running through $\mathbb{N} \cup \{0\}$:*

$$\begin{aligned} E_2^{*,4k+3}(X) &\cong E_2^{*,3}(X_s) \oplus E_2^{*,3}(X, X_s), \\ E_2^{0,4k+2}(X) &\cong H_{\Gamma}^2(X'_s) \oplus E_2^{0,2}(X, X'_s), & E_2^{p,4k+2}(X) &\cong E_2^{p,2}(X, X'_s) \text{ for } p \geq 1, \\ E_2^{*,4k+1}(X) &\cong E_2^{*,1}(X_s) \oplus E_2^{*,1}(X, X_s), \\ E_2^{*,4k}(X) &\cong H^*(\Gamma \backslash X). \end{aligned}$$

not in X_s has mod-2 cohomology isomorphic to $H^*(\mathbb{Z}/2)$. The results of Theorem 10 then imply that δ is the zero map, hence the long exact sequence splits into short exact sequences. The result is an isomorphism of chain complexes

$$E_2^{*,q}(X) \cong E_2^{*,q}(X_s) \oplus E_2^{*,q}(X, X_s).$$

□

Theorem 14 describes the rows of the E_2 page in terms of rows of E_2 pages of subcomplexes and relative complexes. The cohomology of the former terms will be calculated in Section 6. We calculate the latter terms next, first establishing some notation.

Notation 15. Here and in what follows, $\beta_q(\Gamma \backslash X_s) := \dim_{\mathbb{Q}} H_q(\Gamma \backslash X_s; \mathbb{Q})$. Denote by $\chi(\Gamma \backslash X_s) = \beta_0(\Gamma \backslash X_s) - \beta_1(\Gamma \backslash X_s)$ the Euler characteristic of the orbit space $\Gamma \backslash X_s$ of the non-central 2-torsion subcomplex.

For \mathcal{O} not the Gaussian or Eisensteinian integers, the Euler characteristic of the hyperbolic orbit space $\Gamma \backslash X$ vanishes because its boundary consists of disjoint 2-tori [19, p. 513]. Consequently, the Betti numbers β_1 and β_2 of the hyperbolic orbit space $\Gamma \backslash X$ satisfy $1 - \beta_1 + \beta_2 = 0$, so we can replace β_2 by $\beta_1 - 1$ when it is convenient.

Notation 16. For $q \in \{1, 2\}$, denote the dimension $\dim_{\mathbb{F}_2} H^q(\Gamma \backslash X; \mathbb{F}_2)$ by β^q .

The Universal Coefficient Theorem yields

$$H^q(\Gamma \backslash X; \mathbb{F}_2) \cong \text{Hom}(H_q(\Gamma \backslash X; \mathbb{Z}), \mathbb{F}_2) \oplus \text{Ext}(H_{q-1}(\Gamma \backslash X; \mathbb{Z}), \mathbb{F}_2).$$

As X is 2-dimensional, the group $H_2(\Gamma \backslash X; \mathbb{Z})$ contains no torsion, so we obtain that the dimension β^2 equals the Betti number β_2 plus the number N of 2-torsion summands of $H_1(\Gamma \backslash X; \mathbb{Z})$. As $H_0(\Gamma \backslash X; \mathbb{Z})$ contains no torsion, we obtain that $\beta^1 = \beta_1 + N$. The number N vanishes for all absolute values of the discriminant less than 296 and has been determined in [16] on a database which includes all the Bianchi groups of ideal class numbers 1, 2, 3 and 5, most of the cases of ideal class number 4, as well as all of the cases of discriminant absolute value bounded by 500.

Notation 17. Denote by c the co-rank (i.e., the rank of the cokernel) of the map $H^1(\Gamma \backslash X; \mathbb{F}_2) \rightarrow H^1(\Gamma \backslash X_s; \mathbb{F}_2)$ induced by the inclusion $X_s \subset X$.

Remark 18. By the definition of the co-rank c , we have a lower bound $0 \leq c$. As X_s is 1-dimensional, it cannot have torsion in its integral homology, so $\beta_1(\Gamma \backslash X_s) = \beta^1(\Gamma \backslash X_s)$; and hence the co-rank c is obviously identically 0 whenever $\beta_1(\Gamma \backslash X_s) = 0$. When $\beta_1(\Gamma \backslash X_s) > 0$, we can expect an upper bound

$$c \leq \beta_1(\Gamma \backslash X_s) - 1.$$

We arrive at this expectation by a geometric study of $\Gamma \backslash X_s$ based on [12, 14]. Using the description of $\Gamma \backslash X_s$ of those two articles, the argument for the upper bound can be outlined as follows. The subcomplex X_s consists of rotation axes that are geodesic lines L through hyperbolic 3-space \mathcal{H} , with both open ends of L being in the boundary of \mathcal{H} . If an element γ of finite order in Γ sends L to a line with non-empty intersection with L , then L is sent to itself either identically, or by inverting its open ends. In the latter case, $\Gamma \backslash X_s$ has an endpoint or bifurcation point given by an orbit containing the intersection point of L with the fixed line of γ . Hence L can only be quotiented into a loop by a translation of L itself, which has to be carried out by an element of infinite order in Γ . This implies that there can be no facets of the Bianchi fundamental polyhedron filling in such a loop. Thus, every loop in $\Gamma \backslash X_s$ comes from a loop in $\Gamma \backslash X$, and the map

$$H^1(\Gamma \backslash X; \mathbb{F}_2) \rightarrow H^1(\Gamma \backslash X_s; \mathbb{F}_2)$$

has at least rank one. This produces the expected upper bound for the co-rank c . Summarizing, c can be non-zero only for non-central 2-torsion subcomplex quotients containing at least two distinct loops. An example for the latter case is given in Example $(\theta \ o \ o)$ in Section 8.

Notation 19. Let v denote the number of conjugacy classes of subgroups of quaternionic type \mathbf{Q}_8 in $\text{SL}_2(\mathcal{O}_{-m})$, whether or not they are contained in a binary tetrahedral group \mathbf{Te} , and define

$$\text{sign}(v) := \begin{cases} 0, & v = 0, \\ 1, & v > 0. \end{cases}$$

There is a formula for v in terms of the prime divisors of the discriminant of the ring \mathcal{O}_{-m} of integers [8]. Recall from [12, 14] that the endpoints of $\Gamma \backslash X_s$ are precisely the orbits of vertices with stabilizer group \mathbf{Te} , and that the bifurcation points of $\Gamma \backslash X_s$ are precisely the orbits of vertices with stabilizer group \mathbf{Q}_8 . Hence we can think of v as the number of necessary vertices (endpoints or bifurcation points) of $\Gamma \backslash X_s$.

Proposition 20. *For q odd, $E_2^{p,q}(X, X_s) \cong H^p(\Gamma \backslash X, \Gamma \backslash X_s)$. In particular,*

$$\dim_{\mathbb{F}_2} E_2^{p,q}(X, X_s) = \begin{cases} 0 & p = 0 \\ \beta^1(\Gamma \backslash X) + c + \chi(\Gamma \backslash X_s) - 1 & p = 1 \\ \beta^2(\Gamma \backslash X) + c & p = 2 \end{cases}$$

Proof. We start with the split short exact sequence of chain complexes from Sequence (4),

$$(5) \quad 0 \rightarrow E_1^{*,q}(X, X_s) \rightarrow E_1^{*,q}(X) \rightarrow E_1^{*,q}(X_s) \rightarrow 0.$$

Every cell stabilizer in (X, X_s) has cohomology isomorphic to $H^*(\mathbb{Z}/2)$, so

$$E_1^{p,q}(X, X_s) \cong \bigoplus_{\sigma \in \Gamma \backslash (X, X_s)^{(p)}} H^q(\mathbb{Z}/2).$$

In addition, we have the decomposition

$$E_1^{p,q}(X) \cong \bigoplus_{\sigma \in \Gamma \backslash (X, X_s)^{(p)}} H^q(\Gamma_\sigma) \oplus \bigoplus_{\tau \in \Gamma \backslash X_s^{(p)}} H^q(\Gamma_\tau).$$

In Sequence (5), there is a one-to-one correspondence between cohomology classes associated to cells in $X_s^{(*)}$ in the middle term and cohomology classes associated to cells in the rightmost term. Therefore, for fixed odd q we can artificially replace the stabilizers associated to cells in $X_s^{(p)}$ by $\mathbb{Z}/2$ to get a new short exact sequence with the same leftmost term as in Sequence (5):

$$0 \rightarrow \bigoplus_{\sigma \in \Gamma \backslash (X, X_s)^{(p)}} H^q(\mathbb{Z}/2) \rightarrow \bigoplus_{\sigma \in \Gamma \backslash X^{(p)}} H^q(\mathbb{Z}/2) \rightarrow \bigoplus_{\sigma \in \Gamma \backslash X_s^{(p)}} H^q(\mathbb{Z}/2) \rightarrow 0.$$

The differential on the leftmost chain complex is $d_1^{p,q}$, and the next two differentials are induced by the standard coboundary operator with trivial action on the $H^*(\mathbb{Z}/2)$ terms. As X is a 2-dimensional cell complex and X_s is 1-dimensional, upon applying the snake lemma we get the following long exact sequence concentrated in eight terms:

$$\begin{array}{ccccccc} \hookrightarrow & E_2^{2,q}(X, X_s) & \longrightarrow & H^2(\Gamma \backslash X) \otimes_{\mathbb{F}_2} H^q(\mathbb{Z}/2) & \longrightarrow & 0 & \longrightarrow \dots \\ \hookrightarrow & E_2^{1,q}(X, X_s) & \longrightarrow & H^1(\Gamma \backslash X) \otimes_{\mathbb{F}_2} H^q(\mathbb{Z}/2) & \longrightarrow & H^1(\Gamma \backslash X_s) \otimes_{\mathbb{F}_2} H^q(\mathbb{Z}/2) & \longrightarrow \dots \\ 0 & \longrightarrow & E_2^{0,q}(X, X_s) & \longrightarrow & H^0(\Gamma \backslash X) \otimes_{\mathbb{F}_2} H^q(\mathbb{Z}/2) & \longrightarrow & H^0(\Gamma \backslash X_s) \otimes_{\mathbb{F}_2} H^q(\mathbb{Z}/2) \end{array}$$

As any action on $H^q(\mathbb{Z}/2) \cong \mathbb{F}_2$ is trivial, tensoring with $H^q(\mathbb{Z}/2)$ is an isomorphism, and we can drop the factors $\otimes_{\mathbb{F}_2} H^q(\mathbb{Z}/2)$ in the above long exact sequence. Then we can identify the above sequence with the long exact sequence associated to the relative cohomology of the pair $(\Gamma \backslash X, \Gamma \backslash X_s)$,

$$\begin{array}{ccccccc} \hookrightarrow & H^2(\Gamma \backslash X, \Gamma \backslash X_s) & \longrightarrow & H^2(\Gamma \backslash X) & \longrightarrow & 0 & \longrightarrow \dots \\ \hookrightarrow & H^1(\Gamma \backslash X, \Gamma \backslash X_s) & \longrightarrow & H^1(\Gamma \backslash X) & \longrightarrow & H^1(\Gamma \backslash X_s) & \longrightarrow \dots \\ 0 & \longrightarrow & H^0(\Gamma \backslash X, \Gamma \backslash X_s) & \longrightarrow & H^0(\Gamma \backslash X) & \longrightarrow & H^0(\Gamma \backslash X_s) \end{array}$$

This implies our first claim,

$$E_2^{p,q}(X, X_s) \cong H^p(\Gamma \backslash X, \Gamma \backslash X_s).$$

Since $X_s \subset X$ with X connected, we immediately see that $H^0(\Gamma \backslash X)$ maps isomorphically to a 1-dimensional \mathbb{F}_2 -subspace in $H^0(\Gamma \backslash X_s)$, yielding $H^0(\Gamma \backslash X, \Gamma \backslash X_s) = 0$. Therefore, the map from $H^0(\Gamma \backslash X_s)$ to $H^1(\Gamma \backslash X, \Gamma \backslash X_s)$ has rank $\beta^0(\Gamma \backslash X_s) - 1$. In addition, using the co-rank c from Notation 17, we have to complement $H^1(\Gamma \backslash X, \Gamma \backslash X_s)$ by an \mathbb{F}_2 -subspace of dimension $\beta^1(\Gamma \backslash X) - (\beta^1(\Gamma \backslash X_s) - c)$. This yields the claimed formula $\chi(\Gamma \backslash X_s) - 1 + \beta^1(\Gamma \backslash X) + c$ for the dimension of $H^1(\Gamma \backslash X, \Gamma \backslash X_s)$. The remaining terms of the long exact sequence produce

$$H^2(\Gamma \backslash X, \Gamma \backslash X_s) \cong H^2(\Gamma \backslash X) \oplus (\mathbb{F}_2)^c.$$

□

Proposition 21. *In the second row of the equivariant spectral sequence,*

$$\dim_{\mathbb{F}_2} E_2^{p,2}(X, X'_s) = \begin{cases} 1 - \text{sign}(v) & p = 0 \\ \beta^1(\Gamma \backslash X) + v - \text{sign}(v) & p = 1 \\ \beta^2(\Gamma \backslash X) & p = 2 \end{cases}$$

Proof. We claim that there is a short exact sequence of chain complexes,

$$(6) \quad 0 \rightarrow E_1^{*,2}(X, X'_s) \rightarrow E_1^{*,0}(X) \rightarrow (\mathbb{F}_2)^v \rightarrow 0,$$

where the term $(\mathbb{F}_2)^v$ is concentrated in dimension 0. To see this, we note that from Theorem 9, the cohomology rings of stabilizers of cells in (X, X'_s) are either isomorphic to $H^*(\mathbb{Z}/4)$ or $H^*(\mathbb{Z}/2)$. Both cohomology rings have period evenly dividing 2. Consequently, $E_1^{*,2}(X, X'_s)$ is isomorphic to $E_1^{*,0}(X, X'_s)$. Then Sequence (6) arises by composing this isomorphism with the short exact sequence of cell complexes defining $E_1^{*,0}(X, X'_s)$,

$$0 \rightarrow E_1^{*,0}(X, X'_s) \rightarrow E_1^{*,0}(X) \rightarrow E_1^{*,0}(X'_s) \rightarrow 0,$$

since $E_1^{0,0}(X'_s) \cong (\mathbb{F}_2)^v$ and $E_1^{p,0}(X'_s) = 0$ for $p \geq 1$. In expanded view, Sequence (6) has the following form.

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ & & d_1^{2,q} & & d_1^{2,q} & & d_1^{2,q} \\ 0 & \longrightarrow & E_1^{2,2}(X, X'_s) & \longrightarrow & E_1^{2,0}(X) & \longrightarrow & 0 \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ & & d_1^{1,q} & & d_1^{1,q} & & d_1^{1,q} \\ 0 & \longrightarrow & E_1^{1,2}(X, X'_s) & \longrightarrow & E_1^{1,0}(X) & \longrightarrow & 0 \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ & & d_1^{0,q} & & d_1^{0,q} & & d_1^{0,q} \\ 0 & \longrightarrow & E_1^{0,2}(X, X'_s) & \longrightarrow & E_1^{0,0}(X) & \longrightarrow & (\mathbb{F}_2)^v \longrightarrow 0 \\ & & \uparrow & & \uparrow & & \uparrow \\ & & 0 & & 0 & & 0 \end{array}$$

In the resulting long exact sequence obtained from the Snake Lemma, the terms corresponding to the top row immediately imply that $\dim_{\mathbb{F}_2} E_2^{p,2}(X, X'_s) = \beta^2(\Gamma \backslash X)$, since from Theorem 14, $E_2^{*,0}(X) \cong H^*(\Gamma \backslash X)$. The bottom two rows generate a five-term exact sequence

$$0 \rightarrow E_2^{0,2}(X, X'_s) \rightarrow H^0(\Gamma \backslash X) \rightarrow (\mathbb{F}_2)^v \rightarrow E_2^{1,2}(X, X'_s) \rightarrow H^1(\Gamma \backslash X) \rightarrow 0.$$

Now $H^0(\Gamma \backslash X) \cong \mathbb{F}_2$, and $E_2^{0,2}(X, X'_s)$ is isomorphic to the kernel of the map $H^0(\Gamma \backslash X) \rightarrow (\mathbb{F}_2)^v$, which is non-trivial precisely when $v > 0$. Consequently, $\dim_{\mathbb{F}_2} E_2^{0,2}(X, X'_s) = 1 - \text{sign}(v)$. Then there is a short exact sequence

$$0 \rightarrow (\mathbb{F}_2)^{v - \text{sign}(v)} \rightarrow E_2^{1,2}(X, X'_s) \rightarrow H^1(\Gamma \backslash X) \rightarrow 0,$$

which implies $\dim_{\mathbb{F}_2} E_2^{1,2}(X, X'_s) \cong \beta^1(\Gamma \backslash X) + v - \text{sign}(v)$. \square

Corollary 22. *The E_2 page of the equivariant spectral sequence with \mathbb{F}_2 -coefficients associated to the action of Γ on X is concentrated in the columns $n \in \{0, 1, 2\}$ and has the following form.*

$q = 4k + 3$	$E_2^{0,3}(X_s)$	$E_2^{1,3}(X_s) \oplus (\mathbb{F}_2)^{a_1}$	$(\mathbb{F}_2)^{a_2}$
$q = 4k + 2$	$H^2_\Gamma(X'_s) \oplus (\mathbb{F}_2)^{1 - \text{sign}(v)}$	$(\mathbb{F}_2)^{a_3}$	$H^2(\Gamma \backslash X)$
$q = 4k + 1$	$E_2^{0,1}(X_s)$	$E_2^{1,1}(X_s) \oplus (\mathbb{F}_2)^{a_1}$	$(\mathbb{F}_2)^{a_2}$
$q = 4k$	\mathbb{F}_2	$H^1(\Gamma \backslash X)$	$H^2(\Gamma \backslash X)$
$k \in \mathbb{N} \cup \{0\}$	$n = 0$	$n = 1$	$n = 2$

where

$$\begin{aligned} a_1 &= \chi(\Gamma \backslash X_s) - 1 + \beta^1(\Gamma \backslash X) + c \\ a_2 &= \beta^2(\Gamma \backslash X) + c \\ a_3 &= \beta^1(\Gamma \backslash X) + v - \text{sign}(v) \end{aligned}$$

Proof. Theorem 14 describes the structure of the E_2 page of the spectral sequence by row as cohomology of the quotient space ($q = 4k$) and direct sums of cohomology groups of subcomplexes and relative complexes (other values of q). Propositions 20 and 21 provide a description of the cohomology of the relative complexes. \square

6. CALCULATION OF THE SPECTRAL SEQUENCE ON THE SUBCOMPLEX

In Corollary 22, we have expressed the E_2 page of the equivariant spectral sequence converging to $H^*(SL_2(\mathcal{O}_{-m}))$ in terms of invariants of the quotient space and the E_2 page associated to the non-central 2-torsion subcomplex X_s . In this section, we will investigate that latter E_2 page, $E_2^{p,q}(X_s)$. Recall that the only 2-torsion elements which stabilize cells outside of the non-central 2-torsion subcomplex are in the $\mathbb{Z}/2$ center. This allows us to relate back to and use results about the 2-torsion subcomplex for $PSL_2(\mathcal{O}_{-m})$. What we have to do in order to establish this relation is to show that $E_2^{p,q}(X_s)$ splits as a direct sum indexed by the connected components of X_s ; then further that the reduction of subcomplex components is in complete agreement with the analogous reduction for $PSL_2(\mathcal{O}_{-m})$. We do this in the following lemma.

Lemma 23. *The terms $E_2^{p,q}(X_s)$ split into direct summands each with support on one connected component of the quotient of the reduced non-central 2-torsion subcomplex (again denoted X_s).*

Proof. An argument in Section 6 of [14] explains why the mod 2 cohomology of $PSL_2(\mathcal{O}_{-m})$ splits into a direct sum above the virtual cohomological dimension. The argument references the calculations in [3], and notes that classes that arise in one component of the non-central 2-torsion subcomplex do not restrict to subgroups in others. This implies that products between classes that come from distinct components multiply trivially in cohomology.

We need to extend this result from the projective special linear group to the special linear group. In other words, we need to make sure that the result is compatible with the central extension of $PSL_2(\mathcal{O}_{-m})$ by $\{\pm 1\} \cong \mathbb{Z}/2$. Summarizing results in [12] where the 2-torsion subcomplex is developed, we note that all edge fusions, which happen during the reduction of the subcomplex for $PSL_2(\mathcal{O}_{-m})$, remove a vertex with stabilizer $\mathbb{Z}/2$ or S_3 . The adjacent edges which are fused both have stabilizer $\mathbb{Z}/2$. Since $H^*(S_3) \cong H^*(\mathbb{Z}/2)$ and since this is an

isomorphism of rings, the mod 2 cohomology of the component corresponding to the 2-torsion subcomplex is ring-isomorphic to the mod 2 cohomology coming from the original component.

Now the central $\{\pm 1\}$ group acts trivially on the retracted cell complex, so as noted in the discussion after Definition 6, the *non-central* 2-torsion subcomplex for $\mathrm{SL}_2(\mathcal{O}_{-m})$ is identical to the 2-torsion subcomplex for $\mathrm{PSL}_2(\mathcal{O}_{-m})$. The stabilizers for $\mathrm{SL}_2(\mathcal{O}_{-m})$ are extensions of the stabilizers in $\mathrm{PSL}_2(\mathcal{O}_{-m})$ by $\{\pm 1\}$. In particular, $\mathbb{Z}/2$ is extended to $\mathbb{Z}/4$ and S_3 is extended to \mathbf{Di} . Since there is a ring isomorphism $H^*(\mathbf{Di}) \cong H^*(\mathbb{Z}/4)$, we conclude that at vertices where there was an edge fusion in the 2-torsion subcomplex in the $\mathrm{PSL}_2(\mathcal{O}_{-m})$ case, there will also be an edge fusion in the non-central 2-torsion subcomplex in the $\mathrm{SL}_2(\mathcal{O}_{-m})$ case. Therefore, the non-central 2-torsion subcomplex can be reduced as in the $\mathrm{PSL}_2(\mathcal{O}_{-m})$ case presented in [12]. This yields a splitting for all of the terms $E_2^{p,q}(X_s)$. \square

The main task in the remainder of this section will be to use the cohomology and restriction information gathered in Section 4 in order to determine the equivariant cohomology supported on the individual connected components of the quotient of the reduced non-central 2-torsion subcomplex. The key observation for this task is that the individual connected components of the quotient of the reduced non-central 2-torsion subcomplex correspond to groups which can be described as amalgamated products and HNN extensions. We recall the definition of an HNN extension.

Definition 24. Let $\varphi : G_2 \rightarrow G_1$ be an injection of G_2 into G_1 . An *HNN extension* of G_1 is a group with presentation

$$G_1 *_{G_2} = \langle t, G_1 \mid t^{-1}gt = \varphi(g), g \in G_2 \rangle .$$

The element t is often referred to as the *free letter*.

We note that the notation $G_1 *_{G_2}$ is not completely descriptive, since there may be many possible injective maps φ .

Via Bass–Serre theory, it is known that amalgamated products and HNN extensions both act on trees. In the amalgamated product $G = G_1 *_H G_2$, there is an action where the fundamental domain is given by two vertices with stabilizers G_1 and G_2 , connected by an edge with stabilizer H . In the HNN extension $G = G_1 *_{G_2}$, the fundamental domain is a single vertex with a loop where the vertex stabilizer is G_1 and the edge stabilizer is G_2 .

We can calculate the cohomology of HNN extensions and amalgamated products using the next result.

Theorem 25. *Let σ_0 and σ_1 be 0-cells and 1-cells in a fundamental domain, and let G_v and G_e be the vertex and edge stabilizers respectively. Then there is a long exact sequence in cohomology*

$$(7) \quad \cdots \xrightarrow{\alpha} \bigoplus_{e \in \sigma_1} H^{i-1}(G_e) \xrightarrow{\delta} H^i(G) \xrightarrow{\beta} \bigoplus_{v \in \sigma_0} H^i(G_v) \xrightarrow{\alpha} \bigoplus_{e \in \sigma_1} H^i(G_e) \xrightarrow{\delta} \cdots$$

The direct sum is over one edge and two vertices if G is an amalgamated product (resulting in a Mayer–Vietoris sequence), and over one edge and one vertex if G is an HNN extension.

The following is another result from Bass–Serre theory.

Proposition 26. *In the long exact sequence of Theorem 25, the map β is the restriction map. In an amalgamated product, $\alpha = \mathrm{res}_H^{G_1} - \mathrm{res}_H^{G_2}$. In an HNN extension, $\alpha = \mathrm{res}_{G_2}^{G_1} - \varphi^*$, where φ^* is the map induced by conjugating G_2 by the free letter.*

In fact, when the equivariant spectral sequence only has two non-zero columns, the E_1 page degenerates into a Wang sequence which is precisely the long exact sequence (7) in Theorem 25. Furthermore, the map α is the d_1 differential. We note that although we are primarily interested in an additive calculation of the cohomology, the long exact sequence can also be used to determine ring information, as $\ker(\alpha) = \mathrm{im}(\beta)$ and this is a ring isomorphism.

We will use the equivariant spectral sequence to carry out our cohomology calculations. We recall that Remark 11 states that concerning the occurring restriction maps on cohomology, only reduced classes restrict nontrivially to reduced classes.

(o) **Lemma 27.** Let $G \cong \mathbb{Z}/4 *_{\mathbb{Z}/4}$. Then $\dim_{\mathbb{F}_2} H^q(G) = \begin{cases} 1, & q = 0; \\ 2, & q \geq 1. \end{cases}$

Proof. The quotient of the tree acted on by G has shape \circlearrowleft , with vertex and edge stabilizers both isomorphic to $\mathbb{Z}/4$. Using the restriction maps from Theorem 10 we set up the E_1 and E_2 pages of the equivariant spectral sequence for this HNN extension below.

$$\begin{array}{c|ccc} & E_1 \text{ page} & & E_2 \text{ page} \\ 4 & \mathbb{F}_2 & \longrightarrow & \mathbb{F}_2 \\ 3 & \mathbb{F}_2 & \longrightarrow & \mathbb{F}_2 \\ 2 & \mathbb{F}_2 & \longrightarrow & \mathbb{F}_2 \\ 1 & \mathbb{F}_2 & \longrightarrow & \mathbb{F}_2 \\ 0 & \mathbb{F}_2 & \longrightarrow & \mathbb{F}_2 \\ \hline & 0 & & 1 \end{array} \xrightarrow{H_*(d_1)} \begin{array}{c|ccc} & E_2 \text{ page} & & E_2 \text{ page} \\ 4 & \mathbb{F}_2 & & \mathbb{F}_2 \\ 3 & \mathbb{F}_2 & & \mathbb{F}_2 \\ 2 & \mathbb{F}_2 & & \mathbb{F}_2 \\ 1 & \mathbb{F}_2 & & \mathbb{F}_2 \\ 0 & \mathbb{F}_2 & & \mathbb{F}_2 \\ \hline & 0 & & 1 \end{array}$$

The arrows in the spectral sequence are the d_1 differentials given by $d_1 = res_{\mathbb{Z}/4}^{\mathbb{Z}/4} - \varphi^*$. Since both maps $res_{\mathbb{Z}/4}^{\mathbb{Z}/4}$ and φ^* have identical effects on $H^*(\mathbb{Z}/4)$, we obtain $d_1 = 0$. So the E_1 page of the spectral sequence is the same as the E_∞ page. \square

(l) **Lemma 28.** Let $G \cong \mathbf{Te} *_{\mathbb{Z}/4} \mathbf{Te}$. Then $\dim_{\mathbb{F}_2} H^q(G) = \begin{cases} 1, & q = 0; \\ 2, & q = 4k, k \geq 1; \\ 0, & q = 4k + 1; \\ 1, & q = 4k + 2; \\ 3, & q = 4k + 3. \end{cases}$

Proof. The quotient of the tree acted on by G has shape $\bullet \rightarrow \bullet$. Both vertex stabilizers are isomorphic to \mathbf{Te} , and the edge stabilizer is isomorphic to $\mathbb{Z}/4$. Theorems 9 and 10 allow us to build the E_1 and E_2 pages of the equivariant spectral sequence for this amalgamated product. Both pages are below.

$$\begin{array}{c|ccc} & E_1 \text{ page} & & E_2 \text{ page} \\ 6 & 0 & \longrightarrow & \mathbb{F}_2 \\ 5 & 0 & \longrightarrow & \mathbb{F}_2 \\ 4 & (\mathbb{F}_2)^2 & \longrightarrow & \mathbb{F}_2 \\ 3 & (\mathbb{F}_2)^2 & \longrightarrow & \mathbb{F}_2 \\ 2 & 0 & \longrightarrow & \mathbb{F}_2 \\ 1 & 0 & \longrightarrow & \mathbb{F}_2 \\ 0 & (\mathbb{F}_2)^2 & \longrightarrow & \mathbb{F}_2 \\ \hline & 0 & & 1 \end{array} \xrightarrow{H_*(d_1)} \begin{array}{c|ccc} & E_2 \text{ page} & & E_2 \text{ page} \\ 6 & & & \mathbb{F}_2 \\ 5 & & & \mathbb{F}_2 \\ 4 & \mathbb{F}_2 & & \mathbb{F}_2 \\ 3 & (\mathbb{F}_2)^2 & & \mathbb{F}_2 \\ 2 & & & \mathbb{F}_2 \\ 1 & & & \mathbb{F}_2 \\ 0 & \mathbb{F}_2 & & \mathbb{F}_2 \\ \hline & 0 & & 1 \end{array}$$

On the E_1 page, the only non-zero restriction maps are the ones on reduced classes in rows $4k$. \square

We consider the group $G \cong ((\mathbf{Q}_8 *_{\mathbb{Z}/4} \mathbf{Q}_8) *_{\mathbb{Z}/4}) *_{\mathbb{Z}/4}$, constructed as follows. The copies of \mathbf{Q}_8 are amalgamated over a copy of $\mathbb{Z}/4$. The group G is formed from $\mathbf{Q}_8 *_{\mathbb{Z}/4} \mathbf{Q}_8$ via an iterated HNN extension. The first HNN extension takes a second $\mathbb{Z}/4$ subgroup, non-conjugate to the first, in one copy of \mathbf{Q}_8 to a non-conjugate $\mathbb{Z}/4$ subgroup in the other. The second HNN extension is defined similarly using the third non-conjugate $\mathbb{Z}/4$ subgroups.

(\theta) **Lemma 29.** Let $G \cong ((\mathbf{Q}_8 *_{\mathbb{Z}/4} \mathbf{Q}_8) *_{\mathbb{Z}/4}) *_{\mathbb{Z}/4}$ with HNN extensions as specified above. Then

$$\dim_{\mathbb{F}_2} H^q(G) = \begin{cases} 1, & q = 0; \\ 4, & q = 4k, k \geq 1; \\ 4, & q = 4k + 1; \\ 5, & q = 4k + 2; \\ 5, & q = 4k + 3. \end{cases}$$

Proof. The quotient of the tree acted on by G has shape Θ . The two vertex stabilizers are isomorphic to \mathbf{Q}_8 , and the three non-conjugate edge stabilizers are all isomorphic to $\mathbb{Z}/4$.

The E_1 page of the spectral sequence has two copies of $H^*(\mathbf{Q}_8)$ in column 0 and three copies of $H^*(\mathbb{Z}/4)$ in column 1. Our main concern is to determine the action of the d_1 differential from $E_1^{0,1}$ to $E_1^{1,1}$. By Theorem 10, we know that in dimension 1, a non-zero restriction $res_{\mathbb{Z}/4}^{\mathbf{Q}_8}$ can be traced back to a non-zero restriction $res_{\mathbb{Z}/2}^{\mathbf{D}_2}$. In cohomology, let b_{11}, b_{12} , and b_{13} denote the exterior classes in $\oplus_3 H^1(\mathbb{Z}/4)$ corresponding to the three edge stabilizers, and let $x_{11}, y_{11}, x_{12}, y_{12}$ be the nilpotent classes in $H^1(\mathbf{Q}_8) \oplus H^1(\mathbf{Q}_8)$ corresponding to the two vertex stabilizers. Further, assume that x_{11} and x_{12} both restrict isomorphically to b_{11} via the correspondence from Theorem 10, and that y_{11} and y_{12} both restrict isomorphically to b_{12} . In \mathbf{D}_2 , the product of any two non-zero elements is the final non-zero element. Therefore, in cohomology, x_{11}, y_{11}, x_{12} and y_{12} all restrict to b_{13} . That is,

$$\begin{aligned} d_1^{0,1}(x_{11}) &= b_{11} + b_{13} \\ d_1^{0,1}(y_{11}) &= b_{12} + b_{13} \\ d_1^{0,1}(x_{12}) &= b_{11} + b_{13} \\ d_1^{0,1}(y_{12}) &= b_{12} + b_{13}. \end{aligned}$$

So $d_1^{0,1}$ has rank 2. The only other non-zero differential is between polynomial classes in rows $4k$. Let e_{41} and e_{42} be the polynomial generators of $H^4(\mathbf{Q}_8) \oplus H^4(\mathbf{Q}_8)$, and let $e_{21}^2, e_{22}^2, e_{23}^2$ be the squares of the two-dimensional polynomial generators in $\oplus_3 H^4(\mathbb{Z}/4)$. Based on the geometry of the quotient of the tree acted on by G , we have

$$d_1(e_{41}) = d_1(e_{42}) = e_{21}^2 + e_{22}^2 + e_{23}^2.$$

This implies that both the kernel and image of d_1 are 1-dimensional. We can now completely determine the E_1 and E_2 pages of the equivariant spectral sequence.

E_1 page				E_2 page		
6	$(\mathbb{F}_2)^4 \longrightarrow$	$(\mathbb{F}_2)^3$	$H_*(d_1)$ \implies	6	$(\mathbb{F}_2)^4$	$(\mathbb{F}_2)^3$
5	$(\mathbb{F}_2)^4 \longrightarrow$	$(\mathbb{F}_2)^3$		5	$(\mathbb{F}_2)^2$	\mathbb{F}_2
4	$(\mathbb{F}_2)^2 \longrightarrow$	$(\mathbb{F}_2)^3$		4	\mathbb{F}_2	$(\mathbb{F}_2)^2$
3	$(\mathbb{F}_2)^2 \longrightarrow$	$(\mathbb{F}_2)^3$		3	$(\mathbb{F}_2)^2$	$(\mathbb{F}_2)^3$
2	$(\mathbb{F}_2)^4 \longrightarrow$	$(\mathbb{F}_2)^3$		2	$(\mathbb{F}_2)^4$	$(\mathbb{F}_2)^3$
1	$(\mathbb{F}_2)^4 \longrightarrow$	$(\mathbb{F}_2)^3$		1	$(\mathbb{F}_2)^2$	\mathbb{F}_2
0	$(\mathbb{F}_2)^2 \longrightarrow$	$(\mathbb{F}_2)^3$		0	\mathbb{F}_2	$(\mathbb{F}_2)^2$
		1		0	1	

□

(ρ) **Lemma 30.** Let $G \cong (\mathbf{Q}_8 *_{\mathbb{Z}/4}) *_{\mathbb{Z}/4} \mathbf{Te}$, where the HNN extension identifies two non-conjugate copies of $\mathbb{Z}/4$ in \mathbf{Q}_8 ; and the third conjugacy class of $\mathbb{Z}/4$ in \mathbf{Q}_8 is the amalgamated subgroup with \mathbf{Te} . Then

$$\dim_{\mathbb{F}_2} H^q(G) = \begin{cases} 1 & q = 0 \\ 3, & q = 4k, \geq 1; \\ 2, & q = 4k + 1; \\ 3, & q = 4k + 2; \\ 4, & q = 4k + 3. \end{cases}$$

Proof. The quotient of the tree acted on by G has shape $\Theta \bullet$. The vertex stabilizer incident to the loop is isomorphic to \mathbf{Q}_8 . The other vertex stabilizer is isomorphic to \mathbf{Te} , and the two edge stabilizers are isomorphic to $\mathbb{Z}/4$. This is the most complicated case, since the fundamental domain for G consists of two vertices and two edges, and the restriction maps arising from the edges have to be taken into account.

We follow the approach used for a similar mod 2 calculation in Lemma 3.1 of [3]. We start with an analysis of the component of $d_1 : H^*(\mathbf{Q}_8) \rightarrow H^*(\mathbb{Z}/4)$, where the copy of $\mathbb{Z}/4$ is the stabilizer of the loop. Referring back to the long exact sequence given in Equation 7, this component of d_1

is the difference $res_{\mathbb{Z}/4}^{\mathbf{Q}_8} - \varphi^*$, where the restriction map is induced by subgroup injection on one side of the loop, and by the twisting from the HNN extension on the other one. We tweak the notation for classes from Theorem 9 by denoting by $b_{11} \in H^1(\mathbb{Z}/4)$ the class in the loop stabilizer and $b_{12} \in H^1(\mathbb{Z}/4)$ the class from the edge stabilizer. (We use the same convention for reduced classes.) By the mod 2 calculation in [3], which carefully tracks the effect of the twisting map on cohomology, one can show that the generators x_1 and y_1 mentioned in Theorem 9 have images

$$res_{\mathbb{Z}/4}^{\mathbf{Q}_8}(x_1) = b_{11}, \text{ but } res_{\mathbb{Z}/4}^{\mathbf{Q}_8}(y_1) = 0.$$

That is, the restriction only detects the subgroup on one side of the HNN extension. The map on cohomology induced by φ , on the other hand, detects the subgroup on both sides:

$$\varphi^*(x_1) = \varphi^*(y_1) = b_{11}.$$

Consequently, the component of d_1 mapping to the loop sends x_1 to 0 and y_1 to b_{11} . In addition, as $res_{\mathbb{Z}/4}^{\mathbf{Q}_8}(e_4) = \varphi^*(e_4) = e_{21}^2$, the same component of d_1 sends e_4 to 0. The map $res_{\mathbb{Z}/4}^{\mathbf{Q}_8}$ to the unlooped edge with $\mathbb{Z}/4$ stabilizer is given by Theorem 10, although we need to determine whether the nilpotent class with a nontrivial restriction is x_1 , y_1 , or $x_1 + y_1$. In fact, a basis for $H^*(\mathbf{Q}_8)$ can be chosen so that $res_{\mathbb{Z}/4}^{\mathbf{Q}_8}$ sends x_1 to 0 and y_1 to b_{12} . Summarizing,

$$\begin{aligned} res_{\mathbb{Z}/4 \oplus \mathbb{Z}/4}^{\mathbf{Q}_8}(y_1) &= b_{11} + b_{12} \\ res_{\mathbb{Z}/4 \oplus \mathbb{Z}/4}^{\mathbf{Q}_8}(e_4) &= e_{22}^2 \end{aligned}$$

and all other classes are sent to 0. The rest of d_1 is given by the restriction map $res_{\mathbb{Z}/4}^{\mathbf{Te}}$, and this map is nontrivial only on the reduced class. We have now completely determined the d_1 differential, so we can write down the E_1 and E_2 pages.

E_1 page				E_2 page		
6	$(\mathbb{F}_2)^2$	\longrightarrow	$(\mathbb{F}_2)^2$		$(\mathbb{F}_2)^2$	$(\mathbb{F}_2)^2$
5	$(\mathbb{F}_2)^2$	\longrightarrow	$(\mathbb{F}_2)^2$		\mathbb{F}_2	\mathbb{F}_2
4	$(\mathbb{F}_2)^2$	\longrightarrow	$(\mathbb{F}_2)^2$		\mathbb{F}_2	\mathbb{F}_2
3	$(\mathbb{F}_2)^2$	\longrightarrow	$(\mathbb{F}_2)^2$	$\xrightarrow{H_*(d_1)}$	$(\mathbb{F}_2)^2$	$(\mathbb{F}_2)^2$
2	$(\mathbb{F}_2)^2$	\longrightarrow	$(\mathbb{F}_2)^2$		$(\mathbb{F}_2)^2$	$(\mathbb{F}_2)^2$
1	$(\mathbb{F}_2)^2$	\longrightarrow	$(\mathbb{F}_2)^2$		\mathbb{F}_2	\mathbb{F}_2
0	$(\mathbb{F}_2)^2$	\longrightarrow	$(\mathbb{F}_2)^2$		\mathbb{F}_2	\mathbb{F}_2
	0		1		0	1

□

Note 31. As the sub-subcomplex X'_s is 0-dimensional, we can read off the equivariant cohomology $H_q^\Gamma(X'_s)$ as the $E_1^{0,q}$ -terms of its spectral sequence. This furthermore splits as a direct sum over the connected components of X_s , and we write $\bullet \rightarrow \bullet$, \ominus' , respectively $\bigcirc \rightarrow \bullet$ for subsets of X'_s with orbit space constituting the vertices of a connected component $\bullet \rightarrow \bullet$, \ominus , respectively $\bigcirc \rightarrow \bullet$ of $\Gamma \backslash X_s$. Then using Theorem 9, we obtain on the known types of connected components

$$H_2^\Gamma(\bullet \rightarrow \bullet) = 0, H_2^\Gamma(\ominus') \cong (\mathbb{F}_2)^4, H_2^\Gamma(\bigcirc \rightarrow \bullet) \cong (\mathbb{F}_2)^2.$$

The set \bigcirc' is empty, but we can abuse notation and also write $H_2^\Gamma(\bigcirc') = 0$.

7. DECOMPOSITION OF THE SECOND PAGE DIFFERENTIAL

Looking back at the four groups associated to the non-central 2-torsion subcomplex quotient components shown in Table 1, we see that the groups associated to the quotient types $\bullet \rightarrow \bullet$, \ominus , and $\bigcirc \rightarrow \bullet$ have periodic cohomology of period 4 starting above degree $d = 0$, where α is the unique polynomial class in degree 4 which is detected on the fiber of the extension. In contrast, for the quotient type \bigcirc , the periodic cohomology has period 2, where the periodicity generator is again the unique polynomial class in dimension 2.

In the equivariant spectral sequence for the action of $SL_2(\mathcal{O}_{-m})$ on X , which we will denote by E3SL, we have completely determined the d_1 differentials. The final goal is to determine as much

as possible about the d_2 differentials. By periodicity, once we know what happens in the first four rows, we know what happens in the entire spectral sequence. We use a couple of techniques. One is an analysis of Steenrod operations in the E3SL. The target of the d_2 differential is the second column of the E3SL. All 2-cell stabilizers are of type $\mathbb{Z}/2$, and the Steenrod algebra on $H^*(\mathbb{Z}/2)$ is generated by the operation $Sq^1(x_1) = x_1^2$. We will also compare the E3SL with the Lyndon–Hochschild–Serre spectral sequence associated to the extension

$$1 \rightarrow \mathbb{Z}/2 \rightarrow \mathrm{SL}_2(\mathcal{O}_{-m}) \rightarrow \mathrm{PSL}_2(\mathcal{O}_{-m}) \rightarrow 1,$$

and which we will denote by LH3S.

Lemma 32. *All classes in $E_2^{0,2q}$ of the E3SL are d_2 -cocycles.*

Proof. From Theorem 9, we see that Sq^1 is trivial in even degrees. The only classes that might need to be checked are in $H^2(\mathbf{Q}_8)$, and the result follows since these classes arise from squares on the horizontal edge of the LH3S associated to the extension

$$1 \rightarrow \mathbb{Z}/2 \rightarrow \mathbf{Q}_8 \rightarrow \mathbf{D}_2 \rightarrow 1.$$

A non-zero target of the d_2 differential would be an odd-dimensional class in $H^*(\mathbb{Z}/2)$, which has a non-trivial Sq^1 . However, since $Sq^1 d_2 = d_2 Sq^1$ this is impossible. \square

As our cell complex X is 2-dimensional, we obtain the following corollary from this lemma.

Corollary 33. *The d_2 differential can only be nontrivial on nilpotent classes on the vertical edge of the equivariant spectral sequence associated to $\mathrm{SL}_2(\mathcal{O}_{-m})$.*

We note that the $d_2^{0,q}$ differentials are block-diagonalizable with matrix blocks supported each by one connected component of the quotient of the non-central 2-torsion subcomplex. Working one component type at a time, we will show that in the equivariant spectral sequence, for components of type \circlearrowright , $d_2^{0,1}$ is trivial if and only if $d_2^{0,3}$ is. We will also show that the d_2 differential is trivial on components of type \blackrightarrow and \ominus . At the time of the writing of this paper we have been unable to determine d_2 's behavior on components of type \circlearrowright . However, this component type only appears in the database summarized in Appendix A for $m = 2$, and then only with multiplicity 1.

(o) **Lemma 34.** The d_2 differential is nontrivial on cohomology on components of type \circlearrowright in degrees $q \equiv 1 \pmod{4}$ if and only if it is nontrivial on these components in degrees $q \equiv 3 \pmod{4}$.

Proof. The vertex stabilizers for components of type \circlearrowright are $\mathbb{Z}/4$, and $Sq^2(H^3(\mathbb{Z}/4)) \neq 0$. Similarly, classes in $E_2^{2,2}$ are generated by $H^2(\mathbb{Z}/2)$, and Sq^2 of these classes are also non-zero. The result follows by applying periodicity, and noting that $d_2 Sq^2 = Sq^2 d_2$. \square

(l) **Lemma 35.** The d_2^{E3SL} differential is trivial on connected components of type \blackrightarrow .

Proof. We thank the referee of this paper for outlining the following proof. We start with the long exact sequence in Borel cohomology:

$$\dots \xrightarrow{\delta} H_\Gamma^*(X, X_s) \rightarrow H_\Gamma^*(X) \rightarrow H_\Gamma^*(X_s) \xrightarrow{\delta} H_\Gamma^{*+1}(X, X_s) \rightarrow \dots$$

We wish to show that the coboundary operator δ is 0 in dimension 3 when restricted to components of type \blackrightarrow . This will imply that these cohomology classes in $H_\Gamma^3(X_s)$ correspond to classes in $H_\Gamma^3(X)$. At the level of spectral sequences, the class in $H_\Gamma^3(X_s)$ sits in $E_2^{0,3}$ in the E3SL. If $\delta = 0$, then this class is a d_2 -cocycle.

We note that as X_s is one-dimensional and X is path-connected, the Γ -equivariant spectral sequences for (X, X_s) , respectively for X_s , collapse at the E_2 page, as the E_2 -terms are concentrated in the two columns $p = 1$ and $p = 2$, respectively $p = 0$ and $p = 1$. In addition, by Theorem 20, $E_2^{p,q}(X, X_s)$ is isomorphic to $H^q(\mathbb{Z}/2) \otimes H^p(\Gamma \backslash X, \Gamma \backslash X_s)$, with a trivial Steenrod action on the second tensor factor. We also note that the connecting homomorphism δ commutes with Steenrod operations, and that the filtration of the target is compatible with the Steenrod action.

There are (potentially trivial) contributions to $H_\Gamma^4(X, X_s)$ from both columns of the spectral sequence. We begin our analysis of δ by considering the long exact sequence

$$\dots \xrightarrow{\delta'} E_2^{p,q}(X, X_s) \rightarrow E_2^{p,q}(X) \rightarrow E_2^{p,q}(X_s) \xrightarrow{\delta'} E_2^{p+1,q}(X, X_s) \rightarrow \dots$$

which comes from the relative pair (X, X_s) . When $p = 0$, the kernel of δ' is isomorphic to $E_2^{0,q}(X)$, and the d_2 -differential in the Γ -equivariant spectral sequence for X determines a map from this kernel to $E_2^{2,q-1}(X)$. The differential can also be identified with the restriction of δ to the kernel of δ' . Let ι denote a $\bullet \rightarrow$ component from X_s . Then from the proof of Theorem 14, we know that δ' is the zero map when q is odd, in particular, from $E_2^{0,3}(\iota)$ to $E_2^{1,3}(X, \iota)$.

Therefore, any non-zero element in $\text{im } \delta$ must lie in $H_{\text{red}}^2(\mathbb{Z}/2) \otimes H^2(\Gamma \backslash X, \Gamma \backslash X_s)$. However, as $H_\Gamma^5(\iota) = 0$, we see that Sq^2 is trivial on $H_\Gamma^3(\iota)$. On the other hand, Sq^2 is non-trivial on $H_{\text{red}}^2(\mathbb{Z}/2) \otimes H^2(\Gamma \backslash X, \Gamma \backslash X_s)$. We conclude that δ vanishes on all of $H_\Gamma^3(\iota)$, and these classes can be identified with classes in $H_\Gamma^3(X)$. That is, d_2 is the 0-map on connected components of quotient type $\bullet \rightarrow$. □

For the Θ case, we will directly show that d_2^{E3SL} vanishes on $E_2^{0,1}$, which requires a comparison of computations from the E3SL and LH3S. We start by recalling the following result:

(θ) **Lemma 36.** In $\text{PSL}_2(\mathcal{O}_{-m})$, the group associated to Θ is

$$\Gamma_0 = ((\mathbf{D}_2 *_{\mathbb{Z}/2} \mathbf{D}_2) *_{\mathbb{Z}/2}) *_{\mathbb{Z}/2}.$$

Its mod 2 cohomology ring is $H^*(\Gamma_0) \cong \mathbb{F}_2[l_1, m_1, m_3][\sigma_1, \sigma_2, \tau_1]/m_3(m_3 + l_1^2 m_1 + l_1 m_1^2)$, where all products with nilpotent classes are trivial.

Proof. We provide a brief sketch of this argument, which is similar to the proof of Lemma 29. In fact, the restriction maps in degree 1 cohomology are analogous, and these maps determine all of d_1 . One finds that, as in the case of Lemma 29, d_1 has rank 1 in row 0 of the spectral sequence and rank 2 in row 1. In rows 3 and higher, d_1 has rank three and is surjective. The first few rows of the $E_2 = E_\infty$ page of the spectral sequence are as follows, with $E_2^{0,q} \cong (\mathbb{F}_2)^{2q-1}$:

4	$(\mathbb{F}_2)^7$	0
3	$(\mathbb{F}_2)^5$	0
2	$(\mathbb{F}_2)^3$	0
1	$(\mathbb{F}_2)^2$	\mathbb{F}_2
0	\mathbb{F}_2	$(\mathbb{F}_2)^2$
	0	1

The ring relations are not obvious from this table. However, the Bianchi group $\text{PSL}_2(\mathcal{O}_{-10})$ contains exactly one Θ component in its reduced 2-torsion subcomplex, and its ring structure is calculated in [3, Theorem 4.6]. The vertical edge of spectral sequence is generated by two 1-dimensional classes and a 3-dimensional class subject to the relation in the lemma. The three nilpotent classes in the lemma lie in column 1. □

(θ) **Lemma 37.** The differential d_2^{E3SL} vanishes on $E_2^{0,1}$ on connected components of type Θ .

Proof. We start with the Lyndon–Hochschild–Serre spectral sequence associated to the extension

$$1 \rightarrow \mathbb{Z}/2 \longrightarrow ((\mathbf{Q}_8 *_{\mathbb{Z}/4} \mathbf{Q}_8) *_{\mathbb{Z}/4}) *_{\mathbb{Z}/4} \longrightarrow ((\mathbf{D}_2 *_{\mathbb{Z}/2} \mathbf{D}_2) *_{\mathbb{Z}/2}) *_{\mathbb{Z}/2} \rightarrow 1$$

with the following low-dimensional rows of its E_2 page.

3	z_1^3	four classes	four classes	five classes
2	z_1^2	$l_1 z_1^2, m_1 z_1^2, \sigma_1 z_1^2, \tau_1 z_1^2$	four classes	five classes
1	z_1	$l_1 z_1, m_1 z_1, \sigma_1 z_1, \tau_1 z_1$	$l_1^2 z_1, l_1 m_1 z_1, m_1^2 z_1, \tau_2 z_1$	five classes
0		$l_1, m_1, \sigma_1, \tau_1$	$l_1^2, l_1 m_1, m_1^2, \tau_2$	$l_1^3, l_1^2 m_1, l_1 m_1^2, m_1^3, m_3$
	0	1	2	3

An argument similar to the one used for the calculation of $H^*(\mathbf{Q}_8)$ from $H^*(\mathbf{D}_2)$ in Theorem 9 determines $d_2^{0,1}$:

$$d_2(z_1) = l_1^2 + l_1 m_1 + m_1^2.$$

As d_2 is a derivation, $d_2(z_1^2) = 0$. In addition, for any class $c \in H^*((\mathbf{D}_2 *_{\mathbb{Z}/2} \mathbf{D}_2) *_{\mathbb{Z}/2} *_{\mathbb{Z}/2})$ on the horizontal edge of the spectral sequence, $d_2(cz_1) = cd_2(z_1)$. These observations completely determine the d_2 differential. In dimensions ranging from 0 to 3, the resulting E_3 page is concentrated in the following terms.

2	z_1^2	$\sigma_1 z_1^2, \tau_1 z_1^2$		
1		$\sigma_1 z_1, \tau_1 z_1$	$\tau_2 z_1$	
0		$l_1, m_1, \sigma_1, \tau_1$	three classes	three classes
	0	1	2	3

On the E_3 page, we calculate

$$d_3(z_1^2) = d_3(Sq^1(z_1)) = Sq^1(d_2(z_1)) = l_1^2 m_1 + l_1 m_1^2.$$

This latter class is non-zero in E_3 ; passing to the E_4 page, the number of resulting cohomology generators exactly matches the results in Lemma 29 up through dimension 3.

Assume that there is a Bianchi group $SL_2(\mathcal{O}_{-m})$ so that d_2^{E3SL} is nontrivial on $E_2^{0,1}$ on a \ominus component. The classes supported on the \ominus component in $E_2^{0,1}$ of the E3SL correspond to the classes l_1 and m_1 in the calculation of $H^*((\mathbf{Q}_8 *_{\mathbb{Z}/4} \mathbf{Q}_8) *_{\mathbb{Z}/4} *_{\mathbb{Z}/4})$, so without loss of generality $d_2^{E3SL}(l_1) \neq 0$ and the class l_1 does not survive to the E_3 page. Then the class l_1 will be missing from the horizontal edge of the LH3S in the calculation of $H^*(SL_2(\mathcal{O}_{-m}); \mathbb{F}_2)$. We claim this cannot happen. By [14, Theorem 2], in dimensions 3 and higher, $H^*(PSL_2(\mathcal{O}_{-m}); \mathbb{F}_2)$ is isomorphic to the direct sum of the cohomology rings associated to the components of its 2-torsion subcomplex. Therefore, regardless of the d_2 differential in the equivariant spectral sequence for $PSL_2(\mathcal{O}_{-m})$, there will be a 3-dimensional class $l_1(l_1^2 + l_1 m_1 + m_1^2) \in H^3(PSL_2(\mathcal{O}_{-m}))$ in the horizontal edge of the LH3S. In addition, since we accounted for all classes up to dimension 3 from the cohomology of the \ominus component above, the class $l_1(l_1^2 + l_1 m_1 + m_1^2)$, which is $d_2(l_1 z_1)$, cannot survive to E_∞ . The only possibility is that there is a class $b \in H^1(PSL_2(\mathcal{O}_{-m}))$ so that

$$d_2(bz_1) = b(l_1^2 + l_1 m_1 + m_1^2) = l_1(l_1^2 + l_1 m_1 + m_1^2).$$

The class b must be a geometric class coming from $SL_2(\mathcal{O}_{-m}) \backslash X$. As such, b is nilpotent, and $d_2(bz_1) = bd_2(z_1)$ is also nilpotent. On the other hand, $d_2(l_1 z_1)$ is not. We conclude that there can be no class b with the required properties; and hence l_1 survives to E_∞ . \square

(\theta) Lemma 38. The d_2 differential of the E3SL is trivial on $E_2^{0,3}$ on components of type \ominus .

Proof. We determine part of the ring structure of the cohomology of the subgroup associated to \ominus in $SL_2(\mathcal{O}_{-m})$. Using the notation introduced in Lemma 29, let $H^1(\mathbf{Q}_8) \oplus H^1(\mathbf{Q}_8)$ be generated by $x = x_{11}, y = y_{11}$ and $s = x_{12}, t = y_{12}$. Then in the 0-column, the two 1-dimensional classes in \ominus can be represented by $a = (y, t)$ and $b = (x, s)$. Continuing in the 0-column, the four 2-dimensional classes can be represented by $c = (0, t^2)$, $a^2 + c = (y^2, 0)$, $d = (0, s^2)$, $b^2 + d = (x^2, 0)$; and the two 3-dimensional classes by $b_{31} = b(a^2 + c) = a(b^2 + d)$ and $b_{32} = ad = bc$; these relations follow from the ring structure detailed in Theorem 9. The ring relations also imply that $a^2 b = ab^2$ and $ac = bd = 0$, because $x^3 = y^3 = s^3 = t^3 = 0$.

For our purposes, all 3-dimensional classes in the 0-column are products of classes of degree 1 and 2. By Lemmas 32 and 37, d_2 vanishes on these smaller degree classes, so $d_2^{0,3}$ also vanishes as d_2 is a derivation. \square

8. EXAMPLE CALCULATIONS

In the examples computed in this section, we use Corollary 22 and the lemmata of Section 6 to determine the E_2 page of the equivariant spectral sequence converging to the cohomology of $SL_2(\mathcal{O}_{-m})$. Since the equivariant spectral sequence collapses at the E_3 page, once we understand the d_2 differential we can read off $H^*(SL_2(\mathcal{O}_{-m}); \mathbb{F}_2)$. On components of types \blackrightarrow and \ominus , results

in Section 7 shows that this d_2 differential vanishes. On components of type \bigcirc , Lemma 34 shows that we need only to know the rank of $d_2^{0,1}$, for which Appendix A provides numerical results. When we cannot completely determine the rank of d_2 , or when the value of the co-rank c is unclear (see Notation 17), we still can use Bianchi.gp [15] and HAP [6] to compute $H^*(SL_2(\mathcal{O}_{-m}); \mathbb{F}_2)$ in order to resolve this indeterminacy.

Example. (ι). Let $\Gamma \backslash X_s = \bullet \rightarrow \bullet$. Then $v = 2$, $\chi(\Gamma \backslash X_s) = 1$ and $c = 0$. By Lemma 35, the differential $d_2^{E_{3SL}}$ vanishes. Then applying Corollary 22, Note 31 and the proof of Lemma 28, we obtain the following dimensions for $H^*(\Gamma)$.

$$\dim_{\mathbb{F}_2} H^q(SL_2(\mathcal{O}_{-m}); \mathbb{F}_2) = \begin{cases} \beta^1(\Gamma \backslash X) + \beta^2(\Gamma \backslash X), & q = 4k + 5, \\ \beta^1(\Gamma \backslash X) + \beta^2(\Gamma \backslash X) + 2, & q = 4k + 4, \\ \beta^1(\Gamma \backslash X) + \beta^2(\Gamma \backslash X) + 3, & q = 4k + 3, \\ \beta^1(\Gamma \backslash X) + \beta^2(\Gamma \backslash X) + 1, & q = 4k + 2, \\ \beta^1(\Gamma \backslash X), & q = 1. \end{cases}$$

The type $\Gamma \backslash X_s = \bullet \rightarrow \bullet$ occurs for instance for the cases $\Gamma = SL_2(\mathcal{O}_{-m})$ with $m \in \{11, 19, 43, 67, 163\}$, where $\beta^2 = \beta_2 = \beta_1 - 1$ and $\beta^1 = \beta_1$ is given as follows.

m	11	19	43	67	163
β_1	1	1	2	3	7.

In these cases, the results for $\dim_{\mathbb{F}_2} H^q(SL_2(\mathcal{O}_{-m}); \mathbb{F}_2)$ have been checked numerically using Bianchi.gp and HAP.

Example. (θ). Let $\Gamma \backslash X_s = \bigcirc$. Then $v = 2$ and $\chi(\Gamma \backslash X_s) = -1$. By Lemma 38, the differential $d_2^{E_{3SL}}$ vanishes. Applying Corollary 22, Note 31 and the proof of Lemma 29, we obtain the following dimensions for $H^*(\Gamma)$.

$$\dim_{\mathbb{F}_2} H^q(SL_2(\mathcal{O}_{-m}); \mathbb{F}_2) = \begin{cases} c + \beta^1(\Gamma \backslash X) + \beta^2(\Gamma \backslash X) + 2, & q = 4k + 5, \\ c + \beta^1(\Gamma \backslash X) + \beta^2(\Gamma \backslash X) + 2, & q = 4k + 4, \\ c + \beta^1(\Gamma \backslash X) + \beta^2(\Gamma \backslash X) + 3, & q = 4k + 3, \\ c + \beta^1(\Gamma \backslash X) + \beta^2(\Gamma \backslash X) + 3, & q = 4k + 2, \\ \beta^1(\Gamma \backslash X) + 2, & q = 1. \end{cases}$$

$$\text{Instances are } \Gamma = SL_2(\mathcal{O}_{-m}) \text{ with } m = \begin{cases} 5. & \text{Then } \beta^1 = \beta_1 = 2 \text{ and } \beta^2 = \beta_2 = 1. \\ 10 \text{ or } 13. & \text{Then } \beta^1 = \beta_1 = 3 \text{ and } \beta^2 = \beta_2 = 2. \\ 58. & \text{Then } \beta^1 = \beta_1 = 12 \text{ and } \beta^2 = \beta_2 = 11. \end{cases}$$

The authors' numerical calculation produces compatible results for $\dim_{\mathbb{F}_2} H^q(SL_2(\mathcal{O}_{-m}); \mathbb{F}_2)$ at a vanishing co-rank c in all these four cases.

Example. (o). For the case $\Gamma \backslash X_s = \bigcirc$, there is just one loop in $\Gamma \backslash X_s$. By Remark 18, this loop does not become contractible in $\Gamma \backslash X$, so the co-rank c vanishes. Also v and $\chi(\Gamma \backslash X_s)$ are zero. Corollary 22 and the proof of Lemma 27 yield a “stationary” (i.e., period 1) E_2 -page:

$$E_2^{n,q}(X) \cong H^n(\Gamma \backslash X).$$

Let $\Gamma = SL_2(\mathcal{O}_{-m})$ with $m = 7$. Making use of the fact that in this case the quotient space $\Gamma \backslash X$ is homotopy equivalent to a Möbius strip [18], $\beta^1 = \beta_1 = 1$, $\beta^2 = \beta_2 = 0$ and hence $d_2 = 0$. This allows us to conclude that $H^q(SL_2(\mathcal{O}_{-7}); \mathbb{F}_2) \cong (\mathbb{F}_2)^2$ for all $q \geq 1$.

Let $\Gamma = SL_2(\mathcal{O}_{-m})$ with $m = 15$. Then $\beta^1 = \beta_1 = 2$ and $\beta^2 = \beta_2 = 1$. The numerical computation yields

$$\dim_{\mathbb{F}_2} H^q(SL_2(\mathcal{O}_{-15}); \mathbb{F}_2) = \begin{cases} 4, & q \geq 2, \\ 3, & q = 1. \end{cases}$$

We again infer that the d_2 -differential vanishes.

Let $\Gamma = \mathrm{SL}_2(\mathcal{O}_{-m})$ with $m = \begin{cases} 35. \text{ Then } \beta^1 = \beta_1 = 3 \text{ and } \beta^2 = \beta_2 = 2. \\ 91. \text{ Then } \beta^1 = \beta_1 = 5 \text{ and } \beta^2 = \beta_2 = 4. \\ 115. \text{ Then } \beta^1 = \beta_1 = 7 \text{ and } \beta^2 = \beta_2 = 6. \end{cases}$ The numerical

computation yields

$$\dim_{\mathbb{F}_2} H^q(\mathrm{SL}_2(\mathcal{O}_{-35}); \mathbb{F}_2) = \begin{cases} 5, & q \geq 2, \\ 3, & q = 1, \end{cases} \quad \dim_{\mathbb{F}_2} H^q(\mathrm{SL}_2(\mathcal{O}_{-91}); \mathbb{F}_2) = \begin{cases} 9, & q \geq 2, \\ 5, & q = 1, \end{cases}$$

$$\text{and } \dim_{\mathbb{F}_2} H^q(\mathrm{SL}_2(\mathcal{O}_{-115}); \mathbb{F}_2) = \begin{cases} 13, & q \geq 2, \\ 7, & q = 1. \end{cases}$$

So we infer that the d_2 -differential has full rank on the column $n = 0$ both in rows $q = 4k + 1$ and $q = 4k + 3$, while it is zero on classes in even rows.

Example. (ρ). As there is just one loop in $\Gamma \backslash X_s := \mathcal{O} \bullet$, which by Remark 18 does not become contractible in $\Gamma \backslash X$, the co-rank c vanishes. We observe that $v = 2$ and $\chi(\Gamma \backslash X_s) = 0$. Applying Corollary 22, Note 31 and the proof of Lemma 30, we obtain the following dimensions for the $E_2^{n,q}$ -page.

$q = 4k + 3$	2	$\beta^1(\Gamma \backslash X) + 1$	$\beta^2(\Gamma \backslash X)$
$q = 4k + 2$	2	$\beta^1(\Gamma \backslash X) + 1$	$\beta^2(\Gamma \backslash X)$
$q = 4k + 1$	1	$\beta^1(\Gamma \backslash X)$	$\beta^2(\Gamma \backslash X)$
$q = 4k$	1	$\beta^1(\Gamma \backslash X)$	$\beta^2(\Gamma \backslash X)$
$k \in \mathbb{N} \cup \{0\}$	$n = 0$	$n = 1$	$n = 2$

For $\Gamma = \mathrm{SL}_2(\mathcal{O}_{-m})$ with $m = 2$, the orbit space $\Gamma \backslash X$ is homotopy equivalent to a cylinder [18], so $\beta^1(\Gamma \backslash X) = 1$, $\beta^2(\Gamma \backslash X) = 0$.

This allows us to conclude that

$$\dim_{\mathbb{F}_2} H^q(\mathrm{SL}_2(\mathcal{O}_{-2}); \mathbb{F}_2) = \begin{cases} 2, & q = 4k + 5, \\ 3, & q = 4k + 4, \\ 4, & q = 4k + 3, \\ 3, & q = 4k + 2, \\ 2, & q = 1. \end{cases}$$

The examples of Euclidean rings \mathcal{O}_{-2} , \mathcal{O}_{-7} and \mathcal{O}_{-11} have been checked in HAP [6] with the cellular complex imported from Bianchi.gp [15]; and the example \mathcal{O}_{-2} additionally by a paper-and-pencil calculation by the first author using classical methods. We also observe that these three Euclidean examples are compatible with the homology with Steinberg coefficients calculated in [18] up to a minor typo in the case \mathcal{O}_{-2} .

Example. ($o \iota$). Let $\Gamma \backslash X_s = \mathcal{O} \bullet \bullet$. Then $c = 0$, $v = 2$ and $\chi(\Gamma \backslash X_s) = 1$. Using our calculations in the above examples, we obtain the following dimensions for the $E_2^{n,q}$ -page.

$q = 4k + 3$	3	$\beta^1(\Gamma \backslash X) + 2$	$\beta^2(\Gamma \backslash X)$
$q = 4k + 2$	0	$\beta^1(\Gamma \backslash X) + 1$	$\beta^2(\Gamma \backslash X)$
$q = 4k + 1$	1	$\beta^1(\Gamma \backslash X) + 2$	$\beta^2(\Gamma \backslash X)$
$q = 4k$	1	$\beta^1(\Gamma \backslash X)$	$\beta^2(\Gamma \backslash X)$
$k \in \mathbb{N} \cup \{0\}$	$n = 0$	$n = 1$	$n = 2$


This case is realized for $\Gamma = \mathrm{SL}_2(\mathcal{O}_{-m})$ with $m = \begin{cases} 6. & \text{Then } \beta^1 = \beta_1 = 2 \text{ and } \beta^2 = \beta_2 = 1. \\ 22. & \text{Then } \beta^1 = \beta_1 = 5 \text{ and } \beta^2 = \beta_2 = 4. \end{cases}$

With our numerical calculations, we obtain

$$\dim_{\mathbb{F}_2} H^q(\mathrm{SL}_2(\mathcal{O}_{-6}); \mathbb{F}_2) = \begin{cases} 4, & q = 4k + 5, \\ 6, & q = 4k + 4, \\ 7, & q = 4k + 3, \\ 5, & q = 4k + 2, \\ 3, & q = 1, \end{cases} \quad \dim_{\mathbb{F}_2} H^q(\mathrm{SL}_2(\mathcal{O}_{-22}); \mathbb{F}_2) = \begin{cases} 10, & q = 4k + 5, \\ 12, & q = 4k + 4, \\ 13, & q = 4k + 3, \\ 11, & q = 4k + 2, \\ 6, & q = 1, \end{cases}$$

so we conclude that $d_2 = 0$.

$E_{p,q}^2$ -page of this spectral sequence is concentrated as follows in the three columns $p \in \{0, 1, 2\}$,



where the ‘‘Farrell supplement’’ is the cokernel of the map

$$\bigoplus_{\sigma \in \Gamma \backslash X^{(0)}} H_1(\Gamma_\sigma; \mathbb{Z}) \xleftarrow{d_{1,1}^1} \bigoplus_{\sigma \in \Gamma \backslash X^{(1)}} H_1(\Gamma_\sigma; \mathbb{Z})$$

induced by inclusion of cell stabilizers. The Farrell supplement and $H_1(\Gamma \backslash X; \mathbb{Z})$ have been computed on a database of Bianchi groups [16].

For the cases in this database for which the origin of $d_{2,0}^2$ is nontrivial and its target contains 2-torsion, Aurel Page has computed the abelianization $\Gamma^{ab} \cong H_1(\Gamma; \mathbb{Z})$, i.e. the commutator factor subgroup, of $\Gamma = \mathrm{PSL}_2(\mathcal{O}_{-m})$. As the above spectral sequence converges to the group homology of Γ with integer coefficients, we obtain a short exact sequence

$$0 \rightarrow \text{Farrell supplement} / \text{image } d_{2,0}^2 \rightarrow \mathrm{PSL}_2(\mathcal{O}_{-m})^{ab} \rightarrow H_1(\Gamma \backslash X; \mathbb{Z}) \rightarrow 1;$$

and we deduce from it the image of $d_{2,0}^2$ in Appendix A.

Observation 39. Within the scope of the database in Appendix A, the dimension over \mathbb{F}_2 of the 2-primary part of the image of the $d_{2,0}^2$ -differential of the equivariant spectral sequence associated to $\mathrm{PSL}_2(\mathcal{O}_{-m})$ is at most the number of connected components of type \circ in the non-central 2-torsion subcomplex quotient. It is clear from $H_1(\mathcal{A}_4; \mathbb{Z}) \cong \mathbb{Z}/3$ that the target of $d_{2,0}^2$ has no 2-torsion on the connected components of type $\bullet \rightarrow$, and the same property follows for connected components of type $\circ \rightarrow$ by a lemma of [12] specifying the matrix block induced in the $d_{1,1}^1$ -differential by the inclusions into \mathcal{D}_2 . The connected components on which our observation is backed only by the numerical results are the ones of type \ominus .

APPENDIX A. NUMERICAL RESULTS

The program [10] computes a presentation of the Bianchi groups. Aurel Page has carried this out and calculated the commutator factor groups (abelianizations) $\mathrm{SL}_2(\mathcal{O}_{-m})^{Ab}$ and $\mathrm{PSL}_2(\mathcal{O}_{-m})^{Ab}$ in the cases that we need in order to deduce the image of $d_{2,0}^2$ in the way described in Section 9. The outcome of this procedure is included in the below tables. We denote by Δ the discriminant of \mathcal{O}_{-m} , i.e.

$$\Delta = \begin{cases} -m, & m \equiv 3 \pmod{4}, \\ -4m, & \text{otherwise.} \end{cases}$$

and by β_1 the first Betti number of both $\mathrm{PSL}_2(\mathcal{O}_{-m})$ and $\mathrm{SL}_2(\mathcal{O}_{-m})$. This Betti number has been computed with the two independent programs [10] and [15] with identical results. We provide the remaining torsion parts $\mathrm{SL}_2(\mathcal{O}_{-m})_{\mathrm{tors}}^{Ab}$ and $\mathrm{PSL}_2(\mathcal{O}_{-m})_{\mathrm{tors}}^{Ab}$ obtained with [10]. We insert the quotient of the reduced non-central 2-torsion subcomplex from [14] and [15]:

Let o denote the number of connected components of homeomorphism type \circ , let ι the one for $\bullet \rightarrow$, θ for \ominus and ρ for $\circ \rightarrow$. By Corollary 22 and Lemmata 27 through 30, we deduce the rank of $d_{2,0}^2$ for the equivariant spectral sequence with \mathbb{F}_2 -coefficients associated to the action of $\mathrm{SL}_2(\mathcal{O}_{-m})$ on X as the difference

$$\mathrm{rank}_{\mathbb{F}_2} d_{2,0}^2 = o + 2\theta + \rho - \mathrm{rank}_{\mathbb{F}_2} \left(\mathrm{SL}_2(\mathcal{O}_{-m})_{\mathrm{tors}}^{Ab} \right).$$

$-\Delta$	m	o	ι	θ	ρ	β_1	$PSL_2(\mathcal{O}_{-m})_{\text{tors}}^{\text{Ab}}$	$\text{image } d_{2,0}^2(\text{PSL})$	$SL_2(\mathcal{O}_{-m})_{\text{tors}}^{\text{Ab}}$	$\text{rank}_{\mathbb{F}_2} d_2^{0,1}(\text{SL})$
35	35	1	0	0	0	3	$\mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/3$	1
40	10	0	0	1	0	3	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2$	0
47	47	1	0	0	0	5	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
52	13	0	0	1	0	3	$(\mathbb{Z}/2)^2$	0	$(\mathbb{Z}/2)^2$	0
55	55	1	0	0	0	5	$\mathbb{Z}/2$	$\mathbb{Z}/3$	$\mathbb{Z}/4$	0
56	14	1	1	0	0	5	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0
68	17	0	1	1	0	5	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0
79	79	3	0	0	0	6	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/4$	1
84	21	3	0	0	0	7	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
87	87	1	0	0	0	8	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
88	22	1	1	0	0	5	$\mathbb{Z}/2$	$\mathbb{Z}/3$	$\mathbb{Z}/2$	0
91	91	1	0	0	0	5	0	$\mathbb{Z}/2$	0	1
95	95	1	0	0	0	9	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
103	103	1	0	0	0	7	$\mathbb{Z}/2$	0	$\mathbb{Z}/4$	0
104	26	0	0	1	0	8	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0
111	111	1	0	0	0	10	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
115	115	1	0	0	0	7	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	1
116	29	0	0	1	0	9	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0
119	119	2	0	0	0	11	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	1
120	30	3	0	0	0	10	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	2
127	127	1	0	0	0	8	$\mathbb{Z}/2$	0	$\mathbb{Z}/4$	0
132	33	3	2	0	0	10	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
136	34	2	0	0	2	8	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2$	2
143	143	1	0	0	0	12	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/3$	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
148	37	2	0	1	0	8	$(\mathbb{Z}/2)^3$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^3$	1
151	151	1	0	0	0	10	$\mathbb{Z}/2$	0	$\mathbb{Z}/4$	0
152	38	1	1	0	0	10	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0
155	155	1	0	0	0	10	$\mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/3$	1
159	159	1	0	0	0	14	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
164	41	0	1	1	0	12	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0
167	167	1	0	0	0	13	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
168	42	3	0	0	0	13	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	2
183	183	1	0	0	0	14	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
184	46	2	0	0	0	11	$\mathbb{Z}/2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	1
191	191	1	0	0	0	15	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
195	195	2	0	0	0	15	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/3$	2
199	199	1	0	0	0	13	$\mathbb{Z}/2$	0	$\mathbb{Z}/4$	0
203	203	1	0	0	0	12	$\mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/3$	1
212	53	0	0	1	0	14	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0
215	215	1	0	0	0	18	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
219	219	1	2	0	0	13	$\mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/3$	1
223	223	3	0	0	0	15	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/4$	2
228	57	3	2	0	0	16	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
231	231	2	0	0	0	21	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	1
232	58	0	0	1	0	12	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2$	0
235	235	3	0	0	0	13	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	2
239	239	1	0	0	0	18	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
244	61	0	0	1	0	15	$(\mathbb{Z}/2)^2$	0	$(\mathbb{Z}/2)^2$	0
247	247	1	0	0	0	14	$\mathbb{Z}/2$	0	$\mathbb{Z}/4$	0
248	62	2	0	0	0	16	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	1
255	255	2	0	0	0	23	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	1
259	259	1	0	0	0	14	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	1
260	65	1	0	2	0	20	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	1
263	263	1	0	0	0	18	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
264	66	2	2	0	0	20	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	1
271	271	1	0	0	0	17	$\mathbb{Z}/2$	0	$\mathbb{Z}/4$	0
276	69	3	0	0	0	23	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
280	70	3	0	0	0	19	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/2$	2
287	287	2	0	0	0	21	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	1
291	291	1	2	0	0	17	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0
292	73	0	1	1	0	16	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2$	0
295	295	1	0	0	0	19	$\mathbb{Z}/2$	$\mathbb{Z}/3$	$\mathbb{Z}/4$	0
296	74	0	0	1	0	19	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	0
299	299	1	0	0	0	18	$(\mathbb{Z}/3)^2$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	$(\mathbb{Z}/3)^2$	1
303	303	1	0	0	0	22	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
308	77	3	0	0	0	23	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
311	311	1	0	0	0	23	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0

$-\Delta$	m	o	ι	θ	ρ	β_1	$\mathrm{PSL}_2(\mathcal{O}_{-m})_{\mathrm{tors}}^{\mathrm{Ab}}$	$\mathrm{image} d_{2,0}^2(\mathrm{PSL})$	$\mathrm{SL}_2(\mathcal{O}_{-m})_{\mathrm{tors}}^{\mathrm{Ab}}$	$\mathrm{rk}d_2^{0,1}$
312	78	3	0	0	0	22	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
319	319	1	0	0	0	20	$\mathbb{Z}/2$	$\mathbb{Z}/3$	$\mathbb{Z}/4$	0
323	323	1	2	0	0	16	$\mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/3$	1
327	327	1	0	0	0	24	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
328	82	1	1	1	0	17	$(\mathbb{Z}/2)^3$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^3$	0
335	335	1	0	0	0	26	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
340	85	0	0	2	0	23	$(\mathbb{Z}/2)^4$	$(\mathbb{Z}/3)^2$	$(\mathbb{Z}/2)^4$	0
344	86	1	1	0	0	21	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/3$	0	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/3$	0
355	355	1	0	0	0	20	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	1
356	89	0	1	1	0	24	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0
359	359	3	0	0	0	25	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3$	1
367	367	1	0	0	0	20	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/3$	$\mathbb{Z}/4 \oplus (\mathbb{Z}/3)^2$	0
371	371	1	0	0	0	22	$\mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/3$	1
372	93	3	0	0	0	27	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
376	94	2	0	0	0	22	$\mathbb{Z}/2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	1
383	383	1	0	0	0	25	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
388	97	0	1	1	0	21	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2$	0
391	391	2	0	0	0	25	$\mathbb{Z}/2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/4$	1
395	395	1	0	0	0	24	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
399	399	4	0	0	0	33	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3$	2
403	403	1	0	0	0	19	0	$\mathbb{Z}/2$	0	1
404	101	2	0	1	0	28	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/3$	1
407	407	1	0	0	0	29	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/3$	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
408	102	1	2	0	0	27	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^4$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	0
415	415	1	0	0	0	28	$\mathbb{Z}/2$	$\mathbb{Z}/3$	$\mathbb{Z}/4$	0
420	105	8	0	0	0	41	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^6 \oplus (\mathbb{Z}/3)^3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	6
424	106	0	0	1	0	23	$(\mathbb{Z}/2)^4$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^4$	0
427	427	3	0	0	0	21	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/2$	2
431	431	1	0	0	0	29	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
435	435	2	0	0	0	31	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^5$	$\mathbb{Z}/3$	2
436	109	0	0	1	0	25	$(\mathbb{Z}/2)^4$	0	$(\mathbb{Z}/2)^4$	0
439	439	5	0	0	0	26	$(\mathbb{Z}/2)^3$	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/4$	2
440	110	3	0	0	0	32	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
443	443	1	1	0	0	21	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0
447	447	1	0	0	0	32	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
452	113	0	1	1	0	27	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	0	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	0
455	455	2	0	0	0	39	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
456	114	2	2	0	0	32	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^3$	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/3$	1
463	463	1	0	0	0	23	$\mathbb{Z}/2$	0	$\mathbb{Z}/4$	0
471	471	1	0	0	0	34	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/3$	0
472	118	1	1	0	0	25	$(\mathbb{Z}/2)^3$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^3$	0
479	479	1	0	0	0	33	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^3$	0	$\mathbb{Z}/4 \oplus (\mathbb{Z}/3)^3$	0
483	483	2	0	0	0	33	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/3$	2
487	487	1	0	0	0	24	$\mathbb{Z}/2 \oplus (\mathbb{Z}/13)^2$	0	$\mathbb{Z}/4 \oplus (\mathbb{Z}/13)^2$	0
488	122	0	0	1	0	28	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^4 \oplus \mathbb{Z}/3$	0
499	499	2	1	0	0	22	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^2$	0	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^2$	0
520	130	0	0	2	0	32	$(\mathbb{Z}/2)^6$	$(\mathbb{Z}/3)^2$	$(\mathbb{Z}/2)^6$	0
532	133	3	0	0	0	33	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	$(\mathbb{Z}/2)^2$	1
555	555	2	0	0	0	39	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/3$	2
568	142	6	0	0	0	29	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/4$	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^3 \oplus \mathbb{Z}/4$	4
595	595	2	0	0	0	37	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^3$	$\mathbb{Z}/3$	2
667	667	1	0	0	0	32	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	1
696	174	3	0	0	0	50	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	2
715	715	2	0	0	0	43	0	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^2$	0	2
723	723	1	2	0	0	41	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$	1
760	190	3	0	0	0	46	$(\mathbb{Z}/2)^4$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	$(\mathbb{Z}/2)^4$	1
763	763	1	0	0	0	38	0	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	0	1
795	795	2	0	0	0	55	$\mathbb{Z}/3$	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/3$	2
955	955	1	0	0	0	50	$(\mathbb{Z}/2)^4 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/2)^4 \oplus (\mathbb{Z}/3)^2$	1
1003	1003	1	2	0	0	48	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	$\mathbb{Z}/3$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2$	0
1027	1027	1	0	0	0	48	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^3$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^3$	1
1227	1227	1	2	0	0	69	$\mathbb{Z}/4 \oplus \mathbb{Z}/8 \oplus \mathbb{Z}/3$	0	$\mathbb{Z}/4 \oplus \mathbb{Z}/8 \oplus \mathbb{Z}/3$	1
1243	1243	1	2	0	0	58	$(\mathbb{Z}/3)^4$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/3)^4$	1
1387	1387	1	2	0	0	62	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2 \oplus (\mathbb{Z}/167)^2$	$\mathbb{Z}/3$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/3)^2 \oplus (\mathbb{Z}/167)^2$	0
1411	1411	1	2	0	0	64	$\mathbb{Z}/2 \oplus (\mathbb{Z}/16)^2 \oplus (\mathbb{Z}/43)^2$	$\mathbb{Z}/3$	$\mathbb{Z}/2 \oplus (\mathbb{Z}/16)^2 \oplus (\mathbb{Z}/43)^2$	0
1507	1507	1	2	0	0	70	$(\mathbb{Z}/3)^2 \oplus (\mathbb{Z}/5)^4$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/3)^2 \oplus (\mathbb{Z}/5)^4$	1
1555	1555	1	0	0	0	80	$(\mathbb{Z}/4)^8 \oplus (\mathbb{Z}/11)^2$	$\mathbb{Z}/2 \oplus \mathbb{Z}/3$	$(\mathbb{Z}/4)^8 \oplus (\mathbb{Z}/11)^2$	1

REFERENCES

- [1] Alejandro Adem and R. James Milgram, *Cohomology of finite groups*, 2nd ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 309, Springer-Verlag, Berlin, 2004. MR2035696 (2004k:20109)
- [2] Alejandro Adem and Jeff H. Smith, *Periodic complexes and group actions*, Ann. of Math. (2) **154** (2001), no. 2, 407–435, DOI 10.2307/3062102. MR1865976 (2002i:57031)
- [3] Ethan Berkove, *The mod 2 cohomology of the Bianchi groups*, Trans. Amer. Math. Soc. **352** (2000), no. 10, 4585–4602. MR1675241 (2001b:11043)
- [4] Kenneth S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York, 1994. Corrected reprint of the 1982 original. MR1324339 (96a:20072)
- [5] Francis X. Connolly and Stratos Prassidis, *Groups which act freely on $\mathbf{R}^m \times S^{n-1}$* , Topology **28** (1989), no. 2, 133–148, DOI 10.1016/0040-9383(89)90016-5. MR1003578 (90h:57052)
- [6] Graham Ellis, *Homological algebra programming*, Computational group theory and the theory of groups, Contemp. Math., vol. 470, Amer. Math. Soc., Providence, RI, 2008, pp. 63–74, DOI 10.1090/conm/470/09186. MR2478414 (2009k:20001)
- [7] Felix Klein, *Ueber binäre Formen mit linearen Transformationen in sich selbst*, Math. Ann. **9** (1875), no. 2, 183–208. MR1509857
- [8] Norbert Krämer, *Imaginärquadratische Einbettung von Maximalordnungen rationaler Quaternionenalgebren, und die nichtzyklischen endlichen Untergruppen der Bianchi-Gruppen*, preprint, 2013
<http://hal.archives-ouvertes.fr/hal-00720823/en/> (German).
- [9] John McCleary, *User's Guide to Spectral Sequences, Second Edition*, Cambridge Studies in Advanced Mathematics, vol. 58, Cambridge University Press, Cambridge, UK, 2001.
- [10] Aurel Page, *Computing arithmetic Kleinian groups*, accepted for publication in Mathematics of Computation, <http://hal.archives-ouvertes.fr/hal-00703043>, 2014.
- [11] Daniel Quillen, *The spectrum of an equivariant cohomology ring: I, II*, Ann. of Math. **94** (1971), no. 3, 549–572 and 573–602.
- [12] Alexander D. Rahm, *The homological torsion of PSL_2 of the imaginary quadratic integers*, Trans. Amer. Math. Soc. **365** (2013), no. 3, 1603–1635. MR3003276
- [13] ———, *Homology and K-theory of the Bianchi groups*, C. R. Math. Acad. Sci. Paris **349** (2011), no. 11-12, 615–619. MR2817377 (2012e:20116)
- [14] ———, *Accessing the cohomology of discrete groups above their virtual cohomological dimension*, Journal of Algebra **404** (2014), no. C, 152–175. DOI: 10.1016/j.jalgebra.2014.01.025, <http://hal.archives-ouvertes.fr/hal-00618167>
- [15] ———, *Bianchi.gp*, Open source program (GNU general public license), validated by the CNRS: <http://www.projet-plume.org/fiche/bianchigp> Part of the Pari/GP Development Center scripts library, 2010.
- [16] ———, *Higher torsion in the Abelianization of the full Bianchi groups*, LMS J. Comput. Math. **16** (2013), 344–365. MR3109616
- [17] Alexander D. Rahm and Mathias Fuchs, *The integral homology of PSL_2 of imaginary quadratic integers with nontrivial class group*, J. Pure Appl. Algebra **215** (2011), no. 6, 1443–1472, DOI 10.1016/j.jpaa.2010.09.005. Zbl 1268.11072
- [18] Joachim Schwermer and Karen Vogtmann, *The integral homology of SL_2 and PSL_2 of Euclidean imaginary quadratic integers*, Comment. Math. Helv. **58** (1983), no. 4, 573–598, DOI 10.1007/BF02564653. MR728453 (86d:11046)
- [19] Jean-Pierre Serre, *Le problème des groupes de congruence pour SL_2* , Ann. of Math. (2) **92** (1970), 489–527 (French). MR0272790 (42 #7671)
- [20] ———, *Arbres, amalgames, SL_2* , Société Mathématique de France, Paris, 1977 (French). Avec un sommaire anglais; Rédigé avec la collaboration de Hyman Bass; Astérisque, No. 46. Note that there exists an English translation (MR1954121), *Trees*. MR0476875 (57 #16426)
- [21] William M. Singer, *Steenrod Squares in Spectral Sequences*, Mathematical Surveys and Monographs, vol. 129, American Mathematical Society (AMS), Providence, RI, 2006. Zbl 1124.55005
- [22] Karen Vogtmann, *Rational homology of Bianchi groups*, Math. Ann. **272** (1985), no. 3, 399–419, DOI 10.1007/BF01455567. MR799670 (87a:22025)

MATHEMATICS DEPARTMENT, LAFAYETTE COLLEGE, QUAD DRIVE, EASTON, PA 18042, USA

E-mail address: berkovee@lafayette.edu

URL: <http://sites.lafayette.edu/berkovee/>

NATIONAL UNIVERSITY OF IRELAND, DEPARTMENT OF MATHEMATICS, UNIVERSITY ROAD, GALWAY, IRELAND

E-mail address: Alexander.Rahm@nuigalway.ie

URL: <http://www.maths.nuigalway.ie/~rahm/>