

# Global Changes in Ocean Carbon: Variability and Vulnerability

Nicolas Metzl, Bronte Tilbrook, Dorothee C. E. Bakker, Corinne Le Quéré, Scott C. Doney, Richard A. Feely, Maria Hood, Roger Dargaville

## ▶ To cite this version:

Nicolas Metzl, Bronte Tilbrook, Dorothee C. E. Bakker, Corinne Le Quéré, Scott C. Doney, et al.. Global Changes in Ocean Carbon: Variability and Vulnerability: Surface Ocean CO2 Variability and Vulnerability Workshop, Paris, France, 11-14 April 2007. Eos, Transactions American Geophysical Union, 2007, 88 (28), pp.287. 10.1029/2007EO280005 . hal-00770693

## HAL Id: hal-00770693 https://hal.science/hal-00770693

Submitted on 20 Aug 2021  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Global Changes in Ocean Carbon: Variability and Vulnerability

Nicolas Metzl, Bronte Tilbrook, Dorothee C. E. Bakker, Corinne Le Quéré, Scott C. Doney, Richard A. Feely, Maria Hood, Roger Dargaville

## ► To cite this version:

Nicolas Metzl, Bronte Tilbrook, Dorothee C. E. Bakker, Corinne Le Quéré, Scott C. Doney, et al.. Global Changes in Ocean Carbon: Variability and Vulnerability. Eos, Transactions American Geophysical Union, American Geophysical Union (AGU), 2007, 88, pp.287-287. 10.1029/2007EO280005 . hal-00770693

## HAL Id: hal-00770693 https://hal.archives-ouvertes.fr/hal-00770693

Submitted on 20 Aug 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## AmeriFlux Network Aids Global Synthesis

#### PAGE 286

The section again reference processing the section of the secti

The Southern of the South South South States of the South South South South South Southern South South

New of the state o

- part is support to constant instantial information plant in a set of plantic information of the set of the set of the set of the set of the plantic is a set of the set of the set of the plantic is a set of the set of the set of the plantic is a set of the set of the set of the mathematical is a set of the set of the

Indexest in the new birth part theory part theory of the state space of the state of the state of the state space of the state of the state of the state space of the state of the state of the state space of the state of the Chick Control - Provide Application (Control - Control - Contro

(b) Constrained a constraint to the barrow of series and the constraint of the provident of the order barrow of the series of the barrow of sector of the series of the barrow of the sector of the barrow of the series of the formation probability of the constraint of the barrow of the sector of the sector of the sector of the formation barrow of the sector of the sector of the formation barrow of the sector of the sector of the formation barrow of the sector of the sector of the formation.

(a) A set of the se

ha luga ay Agen. Yanaya

(c) and A they had the expression in a contract of a second se

# MEETINGS

### Global Changes in Ocean Carbon: Variability and Vulnerability

# Surface Ocean CO<sub>2</sub> Variability and Vulnerability Workshop, Paris, France, 11–14 April 2007

#### **PAGE 287**

The oceans have taken up approximately half of the anthropogenic  $CO_2$  emissions. This uptake reduces climate change but also lowers ocean pH, with the potential to disrupt ecosystems. Climate change affects ocean biology and physics and could lead to reduced efficiency of the carbon sinks, a process that atmospheric data and ocean models indicate is already occurring in the

Southern Ocean. Attempts to set a baseline stabilization target for the atmospheric  $CO_2$  concentration will ultimately depend on our understanding and prediction of oceanic  $CO_2$  sinks. While we are now close to monitoring oceanic  $CO_2$  uptake on decadal and regional scales, meaningful predictions of its future behavior are difficult. There is a critical and urgent need to better understand the ocean processes regulating  $CO_2$  uptake

and to identify research and observational priorities for the future.

On 11-14 April 2007, over 100 scientists from 20 countries gathered at the United Nations Educational, Scientific, and Cultural Organization (UNESCO), in Paris, to review current knowledge on the magnitude, variability, and processes governing ocean sources and sinks of carbon. The discussions covered observations, process-based models, and atmospheric and oceanic inversions. This workshop was cosponsored by the International Ocean Carbon Coordination Project (IOCCP), the Surface Ocean Lower Atmosphere Study (SOLAS), the Integrated Marine Biogeochemistry and Ecosystem Research (IMBER), and the Global Carbon Project (GCP) programs.

New results were presented for a global airsea  $CO_2$  flux climatology and detection of decadal variations in oceanic partial pressures of  $CO_2$  ( $pCO_2$ ). Observations conducted over more than 20 years show the long-term  $pCO_2$  increase of surface waters is

#### Eos, Vol. 88, No. 28, 10 July 2007

generally close to the atmospheric  $CO_2$ increase, indicating relatively constant sinks. In recent years, significant decadal changes in  $pCO_2$  have been observed in some parts of the ocean, e.g., in the North Atlantic and the equatorial Pacific; however, for many regions there are still no routine observations. Quantification of the decadal changes of the air-sea  $CO_2$  fluxes has been improved using atmospheric data, especially for the vulnerable Southern Ocean, where oceanic data are sparse. Presentations, posters, working group reports, and maps and tables of the ocean carbon observation network are available on the meeting Web page: http:// www.ioc.unesco.org/ioccp/pCO2\_2007.htm.

A major outcome of the workshop was the widespread recognition and strong support for sustained funding and further development of the global observing system for surface  $\rho CO_2$ . The workshop resulted in actions for developing joint synthesis papers, establishing a standard and well-documented global surface  $\rho CO_2$  data set, and producing a regular atlas of surface ocean  $\rho CO_2$ . Regional synthesis groups were formed to analyze the underlying causes for variability and vulnerability in the system and to develop plans for a sustained observing system.

—NICOLAS METZL, Institute Pierre Simon Laplace, Paris; BRONTE TILBROOK, CSIRO Marine and Atmospheric Research, Hobart, Tasmania, Australia; DOROTHEE BAKKER, University of East Anglia (UEA), Norwich, UK; CORINNE LE QUÉRÉ, UEA and British Antarctic Survey, Cambridge, UK; SCOTT DONEY, Woods Hole Oceanographic Institute, Woods Hole, Mass.; RICHARD FEELY, NOAA Pacific Marine Environmental Laboratory, Seattle, Wash.; MARIA HOOD and ROGER DARGAVILLE, United Nations Educational, Scientific, and Cultural Organization (UNESCO), Paris.

PAGE 287