



HAL
open science

The stochastic Weiss conjecture for bounded analytic semigroups

Jamil Abreu, Bernhard Hermann Haak, Jan van Neerven

► **To cite this version:**

Jamil Abreu, Bernhard Hermann Haak, Jan van Neerven. The stochastic Weiss conjecture for bounded analytic semigroups. *Journal London Mathematical Society*, 2013, 88 (1), pp.181-201. 10.1112/jlms/jdt003 . hal-00771883

HAL Id: hal-00771883

<https://hal.science/hal-00771883>

Submitted on 2 Nov 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

THE STOCHASTIC WEISS CONJECTURE FOR BOUNDED ANALYTIC SEMIGROUPS

JAMIL ABREU, BERNHARD HAAK, AND JAN VAN NEERVEN

ABSTRACT. Suppose $-A$ admits a bounded H^∞ -calculus of angle less than $\pi/2$ on a Banach space E which has Pisier's property (α) , let B be a bounded linear operator from a Hilbert space H into the extrapolation space E_{-1} of E with respect to A , and let W_H denote an H -cylindrical Brownian motion. Let $\gamma(H, E)$ denote the space of all γ -radonifying operators from H to E . We prove that the following assertions are equivalent:

- (a) the stochastic Cauchy problem $dU(t) = AU(t) dt + B dW_H(t)$ admits an invariant measure on E ;
- (b) $(-A)^{-1/2} B \in \gamma(H, E)$;
- (c) the Gaussian sum $\sum_{n \in \mathbb{Z}} \gamma_n 2^{n/2} R(2^n, A) B$ converges in $\gamma(H, E)$ in probability.

This solves the stochastic Weiss conjecture of [8].

1. INTRODUCTION

Let A be the generator of a strongly continuous bounded analytic semigroup $S = (S(t))_{t \geq 0}$ on a Banach space E , let F be another Banach space, and let $C : D(A) \rightarrow F$ be a bounded operator. If there exists a constant $M \geq 0$ such that

$$\int_0^\infty \|CS(t)x\|_F^2 dt \leq M^2 \|x\|_E^2, \quad \forall x \in D(A),$$

an easy Laplace transform argument shows that

$$\sup_{\lambda > 0} \lambda^{1/2} \|CR(\lambda, A)\|_{\mathcal{L}(E, F)} \leq M/\sqrt{2}.$$

Here, as usual, $R(\lambda, A) = (\lambda - A)^{-1}$ denotes the resolvent of A at λ .

The celebrated *Weiss conjecture* in linear systems theory is the assertion that the converse also holds. It was solved affirmatively for normal operators A acting on a Hilbert space by Weiss [29], for generators of analytic Hilbert space contraction semigroups with $F = \mathbb{C}$ by Jacob and Partington [10], and subsequently for operators admitting a bounded H^∞ -calculus of angle $< \pi/2$ acting on an L^p -space, $1 < p < \infty$, by Le Merdy [18, 19]. Counterexamples to the general statement were found by Jacob, Partington and Pott [11], Zwart, Jacob, and Staffans [30], and Jacob and Zwart [12].

Whereas the Weiss conjecture is concerned with *observation operators*, in the context of stochastic evolution equations it is natural to consider a 'dual' version of the conjecture in terms of *control operators*. To be more precise, we consider the

Date: December 10, 2012.

2000 Mathematics Subject Classification. Primary: 93B28, Secondary: 35R15, 46B09, 47B10, 47D06.

Key words and phrases. Stochastic Weiss conjecture, stochastic evolution equations, invariant measures, H^∞ -calculus, Rademacher interpolation.

The first named author is supported by FAPESP project 2007/08220-9. The second named author is partially supported by the ANR project ANR-09-BLAN-0058-01. The third named author is supported by VICI subsidy 639.033.604 of the Netherlands Organisation for Scientific Research (NWO).

following situation. Let $W_H = (W_H(t))_{t \in [0, T]}$ be a cylindrical Brownian motion in a Hilbert space H and let $B \in \mathcal{L}(H, E_{-1})$ be a bounded linear operator. Here, E_{-1} denotes the extrapolation space of E with respect to A (see Subsection 2.5). The *stochastic Weiss conjecture*, proposed recently in [8], is the assertion that, under suitable assumptions on the linear operator A , the existence of an invariant measure for the linear stochastic Cauchy problem

$$(SCP)_{(A, B)} \quad \begin{cases} dU(t) = AU(t) dt + B dW_H(t), & t \in [0, T], \\ U(0) = 0, \end{cases}$$

is equivalent to an appropriate condition on the operator-valued function $\lambda \mapsto \lambda^{1/2} R(\lambda, A)B$. This conjecture is justified by the observation (cf. Proposition 2.5 below) that an invariant measure exists if and only if $t \mapsto S(t)B$ defines an element of the space $\gamma(L^2(\mathbb{R}_+; H), E)$ (see Subsection 2.3 for the definition of this space).

In the paper just cited, an affirmative solution was given in the case where A and B are simultaneously diagonalisable. The aim of this article is to prove the stochastic Weiss conjecture for the class of operators admitting a bounded H^∞ -calculus of angle $< \pi/2$. Denoting by $S(E)$ the class of all sectorial operators $-A$ on E of angle $< \pi/2$ that are injective and have dense range, our main result reads as follows.

Theorem 1.1. *Let E have property (α) and assume that $-A \in S(E)$ admits a bounded H^∞ -calculus of angle $< \pi/2$ on E . Let $B : H \rightarrow E_{-1}$ be a bounded operator. Then the following assertions are equivalent:*

- (a) $(SCP)_{(A, B)}$ admits an invariant measure on E ;
- (b) $(-A)^{-1/2} B \in \gamma(H, E)$;
- (c) $\lambda \mapsto \lambda^{1/2} R(\lambda, A)B$ defines an element in $\gamma(L^2(\mathbb{R}_+, \frac{d\lambda}{\lambda}; H), E)$;
- (d) for all $\lambda > 0$ we have $R(\lambda, A)B \in \gamma(H, E)$ and the Gaussian sum

$$\sum_{n \in \mathbb{Z}} \gamma_n 2^{n/2} R(2^n, A)B$$

converges in $\gamma(H, E)$ in probability (equivalently, in $L^p(\Omega; \gamma(H, E))$ for some (all) $1 \leq p < \infty$).

Since B maps into the extrapolation space E_{-1} , some care has to be taken in giving a rigorous interpretations of these assertions. The details will be explained below.

In the special case when E is a Hilbert space and H is a separable Hilbert space with orthonormal basis $(h_k)_{k \geq 1}$, condition (a) is equivalent to

$$\sum_{k=1}^{\infty} \int_0^{\infty} \|S(t)Bh_k\|^2 dt < \infty, \quad (1.1)$$

and condition (d) reduces to

$$\sum_{k=1}^{\infty} \sum_{n \in \mathbb{Z}} 2^n \|R(2^n, A)Bh_k\|^2 < \infty. \quad (1.2)$$

Compared to the Weiss conjecture, we see that a uniform boundedness condition on $\lambda^{1/2} R(\lambda, A)B$ gets replaced by a (dyadic) square summability condition along $(h_k)_{k \geq 1}$ in (1.2); this is consistent with the square summability condition along $(h_k)_{k \geq 1}$ in (1.1).

All spaces are real. When we use spectral arguments, we turn to the complexifications without further notice.

2. PRELIMINARIES

In this section we collect some notations and results that will be used in the proof of Theorem 1.1.

2.1. Property (α) . A *Rademacher sequence* is a sequence of independent random variables taking the values ± 1 with probability $1/2$. Let $(r'_j)_{j=1}^\infty$ and $(r''_k)_{k=1}^\infty$ be Rademacher sequences on probability spaces (Ω', \mathbb{P}') and (Ω'', \mathbb{P}'') , and let $(r_{jk})_{j,k=1}^\infty$ be a doubly indexed Rademacher sequence on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. It is important to observe that the sequence $(r'_j r''_k)_{j,k=1}^\infty$ is not a Rademacher sequence. By standard randomisation techniques one proves (see, e.g., [26]):

Proposition 2.1. *For a Banach space E the following assertions are equivalent:*

- (1) *there exists a constant $C \geq 0$ such that for all finite sequences $(a_{jk})_{j,k=1}^n$ in \mathbb{R} and $(x_{jk})_{j,k=1}^n$ in E we have*

$$\mathbb{E}' \mathbb{E}'' \left\| \sum_{j,k=1}^n a_{jk} r'_j r''_k x_{jk} \right\|^2 \leq C^2 \left(\max_{1 \leq j,k \leq n} |a_{jk}| \right)^2 \mathbb{E}' \mathbb{E}'' \left\| \sum_{j,k=1}^n r'_j r''_k x_{jk} \right\|^2;$$

- (2) *there exists a constant $C \geq 0$ such that for all finite sequences $(x_{jk})_{j,k=1}^n$ in E we have*

$$\frac{1}{C^2} \mathbb{E} \left\| \sum_{j,k=1}^n r_{jk} x_{jk} \right\|^2 \leq \mathbb{E}' \mathbb{E}'' \left\| \sum_{j,k=1}^n r'_j r''_k x_{jk} \right\|^2 \leq C^2 \mathbb{E} \left\| \sum_{j,k=1}^n r_{jk} x_{jk} \right\|^2.$$

A Banach space E is said to have *property (α)* if it satisfies the above equivalent conditions. Examples of spaces having this property are Hilbert spaces and the spaces $L^p(\mu)$ with $1 \leq p < \infty$. Property (α) was introduced by Pisier [27], who proved that a Banach lattice has property (α) if and only if it has finite cotype. In particular, the space c_0 fails property (α) .

2.2. γ -Boundedness. A family $\mathcal{T} \subseteq \mathcal{L}(E, F)$ is called *γ -bounded* if there exists a constant $C \geq 0$ such that for all finite sequences $(T_n)_{n=1}^N$ in \mathcal{T} and $(x_n)_{n=1}^N$ in E we have

$$\mathbb{E} \left\| \sum_{n=1}^N \gamma_n T_n x_n \right\|^2 \leq C^2 \mathbb{E} \left\| \sum_{n=1}^N \gamma_n x_n \right\|^2.$$

Here, and in the rest of the paper, $(\gamma_n)_{n \geq 1}$ denotes a sequence of independent standard Gaussian random variables on a probability space (Ω, \mathbb{P}) . The least admissible constant C in the above inequality is called the *γ -bound* of \mathcal{T} .

By letting $N=1$ it is seen that γ -bounded families are uniformly bounded. For Hilbert spaces E and F , the notions of uniform boundedness and γ -boundedness are equivalent. For detailed expositions on γ -boundedness and the closely related notion of R -boundedness, as well as for references to the extensive literature we refer the reader to [2, 5, 17, 28].

2.3. γ -Radonifying operators. Let \mathcal{H} be a Hilbert space and E a Banach space. For a finite rank operator $T : \mathcal{H} \rightarrow E$ of the form

$$T = \sum_{n=1}^N h_n \otimes x_n,$$

where $(h_n)_{n=1}^N$ is an orthonormal sequence in \mathcal{H} and $(x_n)_{n=1}^N$ is a sequence in E , we define

$$\|T\|_{\gamma(\mathcal{H}, E)} := \left\| \sum_{n=1}^N \gamma_n x_n \right\|_{L^2(\Omega; E)}. \quad (2.1)$$

The Banach space $\gamma(\mathcal{H}, E)$ is defined as the completion of the linear space of finite rank operators with respect to this norm. The identity mapping on the finite rank operators extends to a contractive embedding of $\gamma(\mathcal{H}, E)$ into $\mathcal{L}(\mathcal{H}, E)$. This allows us to view elements of $\gamma(\mathcal{H}, E)$ as operators from \mathcal{H} to E . It may be shown (see [22, Theorem 3.20, (3.1)]) that any $T \in \gamma(\mathcal{H}, E)$ induces an orthogonal decomposition $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$ such that \mathcal{H}_0 is separable and $T|_{\mathcal{H}_1} \equiv 0$; in this situation, for any orthonormal basis $(h_n)_{n \geq 1}$ of \mathcal{H}_0 the Gaussian sum $\sum_{n \geq 1} \gamma_n T h_n$ converges in $L^2(\Omega; E)$ and

$$\|T\|_{\gamma(\mathcal{H}, E)}^2 = \mathbb{E} \left\| \sum_{n \geq 1} \gamma_n T h_n \right\|^2.$$

The distribution of the random variable $\sum_{n \geq 1} \gamma_n T h_n$ is a centred Gaussian Radon measure μ on E whose covariance is given by

$$\int_E \langle x, x^* \rangle \langle x, y^* \rangle d\mu(x) = [T^* x^*, T^* y^*]_{\mathcal{H}}. \quad (2.2)$$

The following γ -Fatou lemma holds (see [15] and [22, Proposition 3.18]). Suppose $(T_n)_{n=1}^\infty$ is a bounded sequence in $\gamma(\mathcal{H}, E)$ and $T \in \mathcal{L}(\mathcal{H}, E)$ is an operator such that

$$\lim_{n \rightarrow \infty} \langle T_n h, x^* \rangle = \langle T h, x^* \rangle, \quad \forall h \in \mathcal{H}, x^* \in E^*.$$

Then, if E does not contain a closed subspace isomorphic to c_0 , we have $T \in \gamma(\mathcal{H}, E)$ and

$$\|T\|_{\gamma(\mathcal{H}, E)} \leq \liminf_{n \rightarrow \infty} \|T_n\|_{\gamma(\mathcal{H}, E)}. \quad (2.3)$$

The *Kalton–Weis extension theorem* [15, Proposition 4.4] (see also [22, Corollary 6.3]) asserts that if $T : H_1 \rightarrow H_2$ is a bounded linear operator, then the tensor extension $T : H_1 \otimes E \rightarrow H_2 \otimes E$,

$$T(h \otimes x) := T h \otimes x$$

extends to a bounded operator (with the same norm) from $\gamma(H_1, E)$ to $\gamma(H_2, E)$.

The *Kalton–Weis multiplier theorem* [15, Proposition 4.11] (see [22, Theorems 4.3 and 5.2] for the formulation given here) asserts that if (X, μ) is a σ -finite measure space, E and F are Banach spaces with F not containing a closed subspace isomorphic to c_0 , and if $M : X \rightarrow \mathcal{L}(E, F)$ is strongly measurable with respect to the strong operator topology and has γ -bounded range, then the mapping

$$(\mathbf{1}_B \otimes h) \otimes x \mapsto (\mathbf{1}_B \otimes h) \otimes Mx$$

has a unique extension to a bounded linear operator from $\gamma(L^2(X, \mu; H), E)$ into $\gamma(L^2(X, \mu; H), F)$ (with norm less than or equal to the γ -bound of the range of M).

Below we shall use (see [26]) that a Banach space E has property (α) if and only if, whenever \mathcal{H}_0 and \mathcal{H}_1 are nonzero Hilbert spaces, the mapping $(h_0 \otimes h_1) \otimes x \mapsto h_0 \otimes (h_1 \otimes x)$ extends to an isomorphism of Banach spaces

$$\gamma(\mathcal{H}_0 \widehat{\otimes} \mathcal{H}_1, E) \simeq \gamma(\mathcal{H}_0, \gamma(\mathcal{H}_1, E)).$$

Here, $\mathcal{H}_0 \widehat{\otimes} \mathcal{H}_1$ denotes the Hilbert space completion of the algebraic tensor product $\mathcal{H}_0 \otimes \mathcal{H}_1$. We will be particularly interested in the case $\mathcal{H}_0 = L^2(\mathbb{R}_+, \frac{dt}{t})$, in which case the above isomorphism then takes the form

$$\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), E) \simeq \gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), \gamma(H, E)).$$

2.4. Stochastic integration. Let H be a Hilbert space and let (Ω, \mathbb{P}) be a probability space. A *cylindrical Brownian motion in H* is a mapping $W_H : L^2(\mathbb{R}_+; H) \rightarrow L^2(\Omega)$ such that $W_H f$ is a centred Gaussian random variable for all $f \in L^2(\mathbb{R}_+; H)$ and

$$\mathbb{E}(W_H f \cdot W_H g) = [f, g]_{L^2(\mathbb{R}_+; H)}$$

for all $f, g \in L^2(\mathbb{R}_+; H)$. Such a mapping is linear and bounded.

A function $\Phi : \mathbb{R}_+ \rightarrow \mathcal{L}(H, E)$ is said to be *stochastically integrable* with respect to W_H if it is scalarly square integrable, i.e., for all $x^* \in E^*$ the function $\Phi^* x^* : t \mapsto \Phi^*(t)x^*$ belongs to $L^2(\mathbb{R}_+; H)$, and for all Borel sets $B \subseteq \mathbb{R}_+$ there exists a random variable $X_B \in L^2(\Omega; E)$ such that

$$\int_B \Phi^* x^* dW_H := W_H(\mathbf{1}_B \Phi^* x^*) = \langle X_B, x^* \rangle, \quad \forall x^* \in E^*.$$

In that case we define

$$\int_B \Phi dW_H := X_B.$$

The following result was proved in [24].

Proposition 2.2. *A scalarly square integrable function $\Phi : \mathbb{R}_+ \rightarrow \mathcal{L}(H, E)$ is stochastically integrable with respect to W_H if and only if there exists an operator $R \in \gamma(L^2(\mathbb{R}_+; H), E)$ such that $R^* x^* = \Phi^* x^*$ in $L^2(\mathbb{R}_+; H)$ for all $x^* \in E^*$. In this situation one has*

$$\mathbb{E} \left\| \int_{\mathbb{R}_+} \Phi dW_H \right\|^2 = \|R\|_{\gamma(L^2(\mathbb{R}_+; H), E)}^2.$$

Suppose W_H is a given cylindrical Brownian motion in H . To any $B \in \gamma(H, E)$ it is possible to associate an E -valued Brownian motion $(W_H^B(t))_{t \geq 0}$ by defining

$$W_H^B(t) = \int_{\mathbb{R}_+} \mathbf{1}_{(0,t)} \otimes B dW_H, \quad t \geq 0.$$

Here we identify $\mathbf{1}_{(0,t)} \otimes B$ with an element in $\gamma(L^2(\mathbb{R}_+; H), E)$ of norm $\sqrt{t} \|B\|_{\gamma(H, E)}$ in the natural way, i.e., by the action

$$f \mapsto \int_{\mathbb{R}_+} \mathbf{1}_{(0,t)}(s) B f(s) ds, \quad f \in L^2(\mathbb{R}_+; H).$$

We may then define the stochastic integral of a simple function $\Psi : \mathbb{R}_+ \rightarrow \mathcal{L}(E)$ with respect to W_H^B by

$$\int_{\mathbb{R}_+} \Psi dW_H^B := \int_{\mathbb{R}_+} \Psi \circ B dW_H. \quad (2.4)$$

With these notations we have the following result [26, Theorem 1.1].

Proposition 2.3. *Let H be a Hilbert space and E a Banach space with property (α) . Let $B \in \gamma(H, E)$ be a given operator and let μ denote the centred Gaussian Radon measure on E associated with B as in (2.2). Let w be a real-valued Brownian motion. For an operator-valued function $\Psi : (0, T) \rightarrow \mathcal{L}(E)$ the following assertions are equivalent:*

- (a) Ψ is stochastically integrable with respect to W_H^B ;
- (b) Ψx is stochastically integrable with respect to w for μ -almost all $x \in E$.

In this situation we have

$$\mathbb{E} \left\| \int_0^T \Psi dW_H^B \right\|^2 \approx \int_E \mathbb{E} \left\| \int_0^T \Psi x dw \right\|^2 d\mu(x) \quad (2.5)$$

with proportionality constants depending on E only.

For later use we point out that, by using Proposition 2.2 and (2.4), the equivalence (2.5) may be reformulated as

$$\|\Psi \circ B\|_{\gamma(L^2(\mathbb{R}_+; H), E)}^2 \approx \int_E \|\Psi x\|_{\gamma(L^2(\mathbb{R}_+; E))}^2 d\mu(x).$$

2.5. Existence, uniqueness and invariant measures. Let A be the generator of a strongly continuous semigroup $S = (S(t))_{t \geq 0}$ on a Banach space E . We define $E_{-1} := (E \times E)/\mathcal{G}(A)$, where $\mathcal{G}(A) = \{(x, Ax) : x \in D(A)\}$ is the graph of A . The mapping

$$i_{-1} : x \mapsto (0, x) + \mathcal{G}(A)$$

defines a dense embedding i_{-1} from E into E_{-1} . We shall always identify E with its image $i_{-1}(E)$ in E_{-1} .

The operator A extends to a bounded operator A_{-1} from E into E_{-1} by defining

$$A_{-1}x := (-x, 0) + \mathcal{G}(A).$$

To see that this indeed gives an extension of A , note that for $x \in D(A)$ we have

$$i_{-1}Ax = (0, Ax) + \mathcal{G}(A) = (-x, 0) + \mathcal{G}(A) = A_{-1}x.$$

It is easy to see that the operator A_{-1} , which is densely defined and closed as a linear operator in E_{-1} with domain $D(A_{-1}) = E$, generates a strongly continuous semigroup $S_{-1} = (S_{-1}(t))_{t \geq 0}$ on E_{-1} which satisfies $S_{-1}(t)i_{-1}x = i_{-1}S(t)x$ for all $x \in E$ and $t \geq 0$.

For a bounded operator $B : H \rightarrow E_{-1}$ we are interested in E -valued solutions to the stochastic evolution equation (SCP) $_{(A, B)}$. To formulate this problem rigorously, we first consider the problem (SCP) $_{(A_{-1}, B)}$ in E_{-1} :

$$(SCP)_{(A_{-1}, B)} \quad \begin{cases} dU_{-1}(t) = A_{-1}U_{-1}(t) dt + B dW_H(t), & t \in [0, T], \\ U_{-1}(0) = 0. \end{cases}$$

Here, as always, W_H is a cylindrical Brownian motion in H , and we adopt the standard notation $W_H(t)h := W_H(\mathbf{1}_{(0, t)} \otimes h)$.

An E -valued process $U = (U(t))_{t \in [0, T]}$ is called a *weak solution* of (SCP) $_{(A, B)}$ if the E_{-1} -valued process $i_{-1}U = (i_{-1}U(t))_{t \in [0, T]}$ is a weak solution of (SCP) $_{(A_{-1}, B)}$, i.e., for all $x_{-1}^* \in D(A_{-1}^*)$ the function $t \mapsto \langle i_{-1}U(t), A_{-1}^*x_{-1}^* \rangle$ is integrable almost surely and if for each $t \in [0, T]$ we have, almost surely,

$$\langle i_{-1}U(t), x_{-1}^* \rangle = \int_0^t \langle i_{-1}U(s), A_{-1}^*x_{-1}^* \rangle ds + W_H(t)B^*x_{-1}^*.$$

An E -valued process U is called a *mild solution* of (SCP) $_{(A, B)}$ if the E_{-1} -valued process $i_{-1}U$ is a mild solution of (SCP) $_{(A_{-1}, B)}$, i.e., if the function $t \mapsto S_{-1}(t)B$ is stochastically integrable in E_{-1} with respect to W_H and if for each $t \in [0, T]$ we have, almost surely,

$$i_{-1}U(t) = \int_0^t S_{-1}(t-s)B dW_H(s). \quad (2.6)$$

The following proposition is an extension of the main result of [24] (where the case $B \in \mathcal{L}(H, E)$ was considered).

Proposition 2.4. *Under the above assumptions, for an E -valued process U the following assertions are equivalent:*

- (a) U is weak solution of (SCP) $_{(A, B)}$;
- (b) U is mild solution of (SCP) $_{(A, B)}$;

(c) there exists an operator $R_T \in \gamma(L^2(0, T; H), E)$ such that for all $x_{-1}^* \in E_{-1}^*$

$$R_T^*(i_{-1}^* x_{-1}^*) = B^* S_{-1}^*(\cdot) x_{-1}^* \quad \text{in } L^2(0, T; H). \quad (2.7)$$

Proof. Let us prove the equivalence (b) \Leftrightarrow (c), because this is what we need in the sequel. The proof of (a) \Leftrightarrow (b) is left to the reader.

(b) \Rightarrow (c): By assumption there is a strongly measurable random variable $U(T) : \Omega \rightarrow E$ such that in E_{-1} we have

$$i_{-1} U(T) = \int_0^T S_{-1}(T-s) B dW_H(s).$$

For all $x_{-1}^* \in E_{-1}^*$, the random variable $\langle U(T), i_{-1}^* x_{-1}^* \rangle$ is Gaussian. Since $F := \{i_{-1}^* x_{-1}^* : x_{-1}^* \in E_{-1}^*\}$ is weak*-dense in E^* and the range of $U(T)$ is separable up to a null set, from [1, Corollary 1.3] it follows that $\langle U(T), x^* \rangle$ is Gaussian for all $x^* \in E^*$, i.e., $U(T)$ is Gaussian distributed.

By the results of [24] the operator $R_{-1, T} : L^2(0, T; H) \rightarrow E_{-1}$, defined by

$$R_{-1, T} f = \int_0^T S_{-1}(T-s) B f(s) ds,$$

belongs to $\gamma(L^2(0, T; H), E_{-1})$. Define the linear operator $R_T^* : F \rightarrow L^2(0, T; H)$ by

$$R_T^* i_{-1}^* x_{-1}^* := R_{-1, T} x_{-1}^*.$$

Then,

$$\begin{aligned} \|R_T^* i_{-1}^* x_{-1}^*\|_{L^2(0, T; H)}^2 &= \|R_{-1, T} x_{-1}^*\|_{L^2(0, T; H)}^2 \\ &= \int_0^T \|B^* S_{-1}^*(T-s) x_{-1}^*\|_H^2 ds \\ &= \mathbb{E} \left| \int_0^T B^* S_{-1}^*(T-s) x_{-1}^* dW_H(s) \right|^2 \\ &= \mathbb{E} |\langle U(T), i_{-1}^* x_{-1}^* \rangle|^2 = \|i_T^* i_{-1}^* x_{-1}^*\|_{\mathcal{H}_T}^2, \end{aligned} \quad (2.8)$$

where i_T is the canonical inclusion mapping of the reproducing kernel Hilbert space \mathcal{H}_T , associated with the Gaussian random variable $U(T)$, into E . This shows that R_T^* is well-defined and bounded on F .

At this point we would like to use a density argument to infer that R_T^* extends to a bounded operator from E^* into $L^2(0, T; H)$ which satisfies

$$\|R_T^* x^*\|_{L^2(0, T; H)}^2 = \|i_T^* x^*\|_{\mathcal{H}_T}^2, \quad \forall x^* \in E^*. \quad (2.9)$$

However, this will not work, since F is only weak*-dense in E^* . The correct way to proceed is as follows. The injectivity of $i_{-1} \circ i_T$ implies that $i_T^* \circ i_{-1}^*$ has weak*-dense range in \mathcal{H}_T . As \mathcal{H}_T is reflexive, this range is weakly dense and therefore, by the Hahn-Banach theorem, it is dense. Fixing an arbitrary $x^* \in E^*$, we may choose a sequence $(x_{-1, n}^*)_{n \geq 1}$ in E_{-1}^* such that $i_T^* i_{-1}^* x_{-1, n}^* \rightarrow i_T^* x^*$ in \mathcal{H}_T . By (2.8) the sequence $(R_T^* i_{-1}^* x_{-1, n}^*)_{n \geq 1}$ is Cauchy in $L^2(0, T; H)$ and converges to some $f_{x^*} \in L^2(0, T; H)$. It is routine to check that f_{x^*} is independent of the approximating sequence. Thus we may extend the R_T^* to E^* by putting

$$R_T^* x^* := f_{x^*}.$$

Clearly, for this extended operator the identity (2.9) is obtained.

We claim that its adjoint $R_T^{**} : L^2(0, T; H) \rightarrow E^{**}$ actually takes values in E , and that this operator is the one we are looking for.

First, for $f = \mathbf{1}_{(a, b)} \otimes h$ and $x^* \in E^*$ of the form $x^* = i_{-1}^* x_{-1}^*$ we have

$$\langle x^*, R_T^{**} f \rangle = [R_T^* i_{-1}^* x_{-1}^*, f]_{L^2(0, T; H)}$$

$$= \int_a^b \langle S_{-1}(T-s)Bh, x_{-1}^* \rangle ds = \langle i_{-1}y, x_{-1}^* \rangle = \langle y, x^* \rangle,$$

where $y = \int_a^b S_{-1}(T-s)Bh ds$ belongs to $D(A_{-1}) = E$. It follows that R_T^{**} maps the dense subspace of all H -valued step functions into E , and therefore it maps all of $L^2(0, T; H)$ into E .

Viewing $R_T := R_T^{**}$ as an operator from $L^2(0, T; H)$ to E , we finally note that the identity (2.9) exhibits $R_T \circ R_T^* = i_T \circ i_T^*$ as the covariance operator of the E -valued Gaussian random variable $U(T)$. This means that R_T is γ -radonifying as an operator from $L^2(0, T; H)$ to E (see, e.g., [22, Theorem 7.4]).

(c) \Rightarrow (b): We follow the ideas of [24]. We have $L^2(0, T; H) = \mathbf{N}(R_T) \oplus \overline{\mathbf{R}(R_T^*)}$. By the general theory of γ -radonifying operators, $G := \overline{\mathbf{R}(R_T^*)}$ is separable (see [22, Equation (3.1)]). By a Gram-Schmidt argument we may select a sequence $(x_{-1,n}^*)_{n \geq 1}$ in E_{-1}^* such that $(g_n)_{n \geq 1} := (R_T^* i_{-1}^* x_{-1,n}^*)_{n \geq 1}$ is an orthonormal basis for G . Then the Gaussian random variables

$$\gamma_n := \int_0^T B^* S_{-1}^*(T-s)x_{-1,n}^* dW_H(s)$$

are independent and normalised. Since R_T is γ -radonifying, the E -valued random variable

$$U(T) := \sum_{n \geq 1} \gamma_n R_T g_n$$

is well-defined, and it is easy to check that it satisfies (2.6) with t replaced by T . By well-known routine arguments, this is enough to assure that $(\text{SCP})_{(A,B)}$ has a mild solution U in E . \square

Suppose now that the problem $(\text{SCP})_{(A_{-1},B)}$ admits a mild solution U_{-1} in E_{-1} and let $\mu_{-1,t}$ denote the distribution of the random variable $U_{-1}(t)$. The weak limit $\mu_{-1,\infty}$ of these measures, if it exists, is called the (minimal) *invariant measure* associated with $(\text{SCP})_{(A_{-1},B)}$. Thus, by definition, the invariant measure, if it exists, is the unique Radon probability measure on E_{-1} which satisfies

$$\int_{E_{-1}} f d\mu_{-1,\infty} = \lim_{t \rightarrow \infty} \int_{E_{-1}} f d\mu_{-1,t}, \quad \forall f \in C_b(E_{-1}).$$

For an explanation of this terminology and a more systematic approach we refer the reader to [4]. This reference deals with Hilbert spaces E ; extensions of the linear theory to the Banach space setting were presented in [7, 25].

A Radon probability measure μ on E is an *invariant measure* for $(\text{SCP})_{(A,B)}$ if the image measure $i_{-1}(\mu)$ on E_{-1} is an invariant measure for $(\text{SCP})_{(A_{-1},B)}$. Extending a result from [25] (where the case $B \in \mathcal{L}(H, E)$ was considered) we have the following result. A proof is obtained along the same line of reasoning as in the previous proposition and is left as an exercise to the reader.

Proposition 2.5. *Under the above assumptions, for a Radon probability measure μ on E the following assertions are equivalent:*

- (a) $(\text{SCP})_{(A,B)}$ admits an invariant measure;
- (b) there exists an operator $R_\infty \in \gamma(L^2(\mathbb{R}_+; H), E)$ such that for all $x_{-1}^* \in E_{-1}^*$

$$R_\infty^*(i_{-1}^* x_{-1}^*) = B^* S_{-1}^*(\cdot) x_{-1}^* \text{ in } L^2(\mathbb{R}_+; H). \quad (2.10)$$

Formally, (2.7) and (2.10) express that the operators R_T and R_∞ are integral operators with kernels $S(\cdot)B$. Strictly speaking this makes no sense, since B maps into E_{-1} rather than into E . It will be convenient, however, to refer to R_T and R_∞ as the operators ‘associated with $S(\cdot)B$ ’ and we shall do so in the sequel without further warning.

2.6. Sectorial operators and H^∞ -calculus. For $\theta \in (0, \pi)$ let

$$\Sigma_\theta := \{z \in \mathbb{C} \setminus \{0\} : |\arg(z)| < \theta\}$$

denote the open sector of angle θ . A densely defined closed linear operator $-A$ in a Banach space E is called *sectorial (of angle $\theta \in (0, \pi)$)* if the spectrum of $-A$ is contained in $\overline{\Sigma_\theta}$ and

$$\sup_{z \notin \overline{\Sigma_\theta}} \|z(z+A)^{-1}\| < \infty.$$

The infimum of all $\theta \in (0, \pi)$ such that $-A$ is sectorial of angle θ is called the *angle of sectoriality* of $-A$.

It is well known (see [6, Theorem II.4.6]) that $-A$ is sectorial of angle less than $\pi/2$ if and only if A generates a strongly continuous bounded analytic semigroup on E .

Following [16] we denote by $S(E)$ the set of all densely defined, closed, injective operators in E that are sectorial of angle less than $\pi/2$ and have dense range. The injectivity and dense range conditions are not very restrictive: if A is a sectorial operator on a reflexive Banach space E , then we have the direct sum decomposition

$$E = N(A) \oplus \overline{R(A)}$$

in terms of the null space and closure of the range of A . In that case, the part of A in $\overline{R(A)}$ is sectorial and satisfies the additional injectivity and dense range conditions.

Let $-A \in S(E)$ be sectorial of angle $\theta \in (0, \pi/2)$ and fix $\eta \in (\theta, \pi/2)$. We denote by $H_0^\infty(\Sigma_\eta)$ the linear space of all bounded analytic functions $f : \Sigma_\eta \rightarrow \mathbb{C}$ with some power type decay at zero and infinity, i.e., for which there exists an $\varepsilon > 0$ such that

$$|f(z)| \leq C|z|^\varepsilon/(1+|z|)^{2\varepsilon}, \quad \forall z \in \Sigma_\eta.$$

For such functions we may define a bounded operator

$$f(-A) = \frac{1}{2\pi i} \int_{\partial \Sigma_{\eta'}} f(z)(z+A)^{-1} dz,$$

with $\eta' \in (\theta, \eta)$. The operator $-A$ is said to have a bounded H^∞ -calculus if there exists a constant C , independent of f , such that

$$\|f(-A)\| \leq C\|f\|_\infty, \quad \forall f \in H_0^\infty(\Sigma_\eta).$$

The infimum of all admissible η is called the *angle* of the H^∞ -calculus of $-A$.

Examples of operators A for which $-A$ has a bounded H^∞ -calculus of angle less than $\pi/2$ are generators of strongly continuous analytic contraction semigroups on Hilbert spaces and second order elliptic operators on L^p -spaces whose coefficients satisfy mild regularity assumptions. We refer to [5, 9, 17] for more details and examples.

If $-A \in S(E)$ has a bounded H^∞ -calculus, the mapping $f \mapsto f(-A)$ extends (uniquely, in some natural sense discussed in [17]) to a bounded algebra homomorphism from $H^\infty(\Sigma_\eta)$ into $\mathcal{L}(E)$ of norm at most C . A proof of the following result can be found in [13, Theorem 5.3].

Proposition 2.6. *Suppose that $-A \in S(E)$ admits a bounded H^∞ -calculus of angle $\eta < \pi/2$ and let $\eta < \eta' < \pi/2$. If E has property (α) , then the family*

$$\{f(-A) : f \in H^\infty(\Sigma_{\eta'}), \|f\|_\infty \leq 1\}$$

is γ -bounded. In particular, $-A$ is γ -sectorial of any angle $\eta < \eta' < \pi/2$, i.e., the family

$$\{z(z+A)^{-1} : z \notin \overline{\Sigma_{\eta'}}\}$$

is γ -bounded.

Remark 2.7. For the γ -sectoriality of $-A$ it suffices that E should have the so-called property (Δ) (see [13, Section 3]). Examples of spaces with this property include all UMD Banach spaces and all Banach spaces with property (α) . As we will not have any application for this, we refer the interested reader to [13, 17] for fuller discussions.

We will also need the following result.

Proposition 2.8. *Suppose that $-A \in S(E)$ has a bounded H^∞ -calculus of angle $\omega < \pi/2$ on a Banach space E with property (α) and let $\theta \in (\omega, \pi)$.*

- (a) *For all $\phi \in H_0^\infty(\Sigma_\theta)$, the function $t \mapsto \phi(-tA)x$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), E)$ for all $x \in X$ and*

$$\|t \mapsto \phi(-tA)x\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), E)} \lesssim \|x\|$$

with implied constant independent of $x \in X$.

- (b) *Given any two $\phi, \psi \in H_0^\infty(\Sigma_\theta)$, for all $x \in X$ we have*

$$\|t \mapsto \phi(-tA)x\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), E)} \approx \|t \mapsto \psi(-tA)x\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), E)}$$

with implied constants independent of $x \in X$.

This is a Banach space version of the classical square function characterisation of the H^∞ -calculus due to McIntosh [21] (see also [3, 9, 17]) and was first obtained in [15, Section 7]. As yet the manuscript [15] remains unpublished, but a proof can be obtained by following the lines of [17, Theorem 11.9]. An explicit proof of (a) is contained in [20, Theorem 6.3]; another proof will be contained in a forthcoming paper by Haak and Haase. The special case of (b) for spaces $E = L^p$ is older and can be found in [19, Corollary 2.3]; again the argument can be extended to yield the general case.

2.7. Rademacher interpolation. If $-A$ is a sectorial operator on E , then for $\theta \in \mathbb{R}$ we may define the Banach space \dot{E}_θ as the completion of $D((-A)^\theta)$ with respect to the norm

$$\|x\|_{\dot{E}_\theta} := \|(-A)^\theta x\|.$$

Note that $(-A)^\theta$ extends uniquely to an isomorphism from \dot{E}_θ onto E ; with some abuse of notation this extension will also be denoted by $(-A)^\theta$. In particular, \dot{E}_{-1} is the completion of the range $R(A)$ with respect to the norm

$$\|Ax\|_{\dot{E}_{-1}} := \|x\|.$$

The identity maps on E and $R(A)$ extend uniquely to continuous inclusions $E \hookrightarrow E_{-1}$ and $\dot{E}_{-1} \hookrightarrow E_{-1}$, and under these identifications we actually have

$$E + \dot{E}_{-1} = E_{-1} \tag{2.11}$$

with equivalent norms. For the reader's convenience we include the short proof. We trivially have $E \hookrightarrow E_{-1}$, and the embedding $\dot{E}_{-1} \hookrightarrow E_{-1}$ is a consequence of the fact that for all $x \in R(A)$, say $x = Ay$, we have

$$\|x\|_{E_{-1}} \leq C\|(I - A)^{-1}x\| = C\|A(I - A)^{-1}\| \|y\| = C\|A(I - A)^{-1}\| \|x\|_{\dot{E}_{-1}}.$$

It follows that $E + \dot{E}_{-1} \hookrightarrow E_{-1}$ with continuous inclusion. Since $I - A$ is surjective from E onto E_{-1} , every $x \in E_{-1}$ is of the form $x = y - Ay$ for some $y \in E$, which implies that $x \in E + \dot{E}_{-1}$. It follows that the inclusion $E + \dot{E}_{-1} \hookrightarrow E_{-1}$ is surjective, and the claim now follows from the open mapping theorem.

Let (X_0, X_1) be an interpolation couple of Banach spaces. Let $(r_n)_{n \in \mathbb{Z}}$ be a Rademacher sequence on a probability space (Ω, \mathbb{P}) . For $0 < \theta < 1$ the *Rademacher*

interpolation space $\langle X_0, X_1 \rangle_\theta$ consists of all $x \in X_0 + X_1$ which can be represented as a sum

$$x = \sum_{n \in \mathbb{Z}} x_n, \quad x_n \in X_0 \cap X_1, \quad (2.12)$$

convergent in $X_0 + X_1$, such that

$$\begin{aligned} \mathcal{C}_0((x_n)_{n \in \mathbb{Z}}) &:= \sup_{N \geq 0} \mathbb{E} \left(\left\| \sum_{n=-N}^N r_n 2^{-n\theta} x_n \right\|_{X_0}^2 \right)^{1/2} < \infty, \\ \mathcal{C}_1((x_n)_{n \in \mathbb{Z}}) &:= \sup_{N \geq 0} \mathbb{E} \left(\left\| \sum_{n=-N}^N r_n 2^{n(1-\theta)} x_n \right\|_{X_1}^2 \right)^{1/2} < \infty. \end{aligned}$$

The norm of an element $x \in \langle X_0, X_1 \rangle_\theta$ is defined as

$$\|x\|_{\langle X_0, X_1 \rangle_\theta} := \inf \left(\max \{ \mathcal{C}_0((x_n)_{n \in \mathbb{Z}}), \mathcal{C}_1((x_n)_{n \in \mathbb{Z}}) \} \right),$$

where the infimum extends over all representations (2.12). This interpolation method was introduced by Kalton, Kunstmann and Weis, who proved that if $-A$ admits a bounded H^∞ -calculus (of any angle $< \pi$), then for all $0 < \theta < 1$ and real numbers $\alpha < \beta$ one has

$$\langle \dot{E}_\alpha, \dot{E}_\beta \rangle_\theta = \dot{E}_{(1-\theta)\alpha + \theta\beta}$$

with equivalent norms [14, Theorem 7.4]. Applying this to the induced operator $I \otimes A$ on $L^2(\Omega; E)$, defined by $(I \otimes A)(f \otimes x) := f \otimes Ax$ for $f \in L^2(\Omega)$ and vectors $x \in \mathcal{D}(A)$, we obtain the following vector-valued extension of this result:

Proposition 2.9. *If $-A \in S(E)$ admits a bounded H^∞ -calculus, then*

$$\langle L^2(\Omega; \dot{E}_\alpha), L^2(\Omega; \dot{E}_\beta) \rangle_\theta = L^2(\Omega; \dot{E}_{(1-\theta)\alpha + \theta\beta}).$$

3. PROOF OF THEOREM 1.1

We begin with a useful observation.

Lemma 3.1. *Let A generate a strongly continuous semigroup on E and suppose that the equivalent conditions of Proposition 2.5 be satisfied. Then for all $\lambda \in \varrho(A)$ there exists an operator $\widehat{S}(\lambda)B \in \gamma(H, E)$ such that*

$$i_{-1} \circ \widehat{S}(\lambda)B = R(\lambda, A_{-1}) \circ B.$$

Proof. It suffices to prove this for one $\lambda \in \varrho(A)$; then, by the resolvent identity, this holds for all $\lambda \in \varrho(A)$.

Fix an arbitrary $\lambda > \omega_0(S_{-1})$, the exponential growth bound of $(S_{-1}(t))_{t \geq 0}$. By assumption there exists an operator $R_\infty \in \gamma(L^2(\mathbb{R}_+; H), E)$ such that for all $x_{-1}^* \in E_{-1}^*$ we have $R_\infty^*(i_{-1}^* x_{-1}^*) = B^* S_{-1}^*(\cdot) x_{-1}^*$ in $L^2(\mathbb{R}_+; H)$. The operator $\widehat{S}(\lambda)B : H \rightarrow E$ given by

$$\widehat{S}(\lambda)Bh := R_\infty(e^{-\lambda \cdot} \otimes h)$$

is γ -radonifying and satisfies, for all $x_{-1}^* \in E_{-1}^*$,

$$\langle i_{-1} \widehat{S}(\lambda)Bh, x_{-1}^* \rangle = \int_0^\infty e^{-\lambda t} \langle S_{-1}(t)Bh, x_{-1}^* \rangle dt = \langle R(\lambda, A_{-1})Bh, x_{-1}^* \rangle.$$

Hence by the Hahn-Banach theorem, $\widehat{S}(\lambda)B$ satisfies the desired identity. \square

The resolvent $R(\lambda, A_{-1})$ maps E_{-1} into $D(A_{-1}) = E$ and therefore we may interpret $R(\lambda, A_{-1})B$ as an operator from H to E . By the injectivity of i_{-1} this operator equals $\widehat{S}(\lambda)B$. From now on we simply write

$$R(\lambda, A)B := \widehat{S}(\lambda)B$$

to denote this operator.

Proposition 3.2. *Suppose that $-A \in S(E)$ has a bounded H^∞ -calculus of angle $\omega < \pi/2$ on a Banach space E with property (α) . Then for all $B \in \mathcal{L}(H, E_{-1})$ and $\theta \in (\omega, \pi)$ the following assertions are equivalent:*

- (a) $B \in \gamma(H, \dot{E}_{-\gamma/2})$;
- (b) $t \mapsto \phi(-tA)B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\gamma/2})$ for all $\phi \in H_0^\infty(\Sigma_\theta)$;
- (c) $t \mapsto \psi(-tA)B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\gamma/2})$, with $\psi(z) = z^{1/2}/(1+z)^{3/2}$.

In this situation, for any two $\phi, \tilde{\phi} \in H_0^\infty(\Sigma_\theta)$ satisfying

$$\int_0^\infty \phi(t) \frac{dt}{t} = \int_0^\infty \tilde{\phi}(t) \frac{dt}{t} = 1$$

we have an equivalence of norms

$$\|t \mapsto \phi(-tA)B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\gamma/2})} \approx \|t \mapsto \tilde{\phi}(-tA)B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\gamma/2})} \quad (3.1)$$

with implied constants independent of ϕ and $\tilde{\phi}$.

Proof. We shall prove the implications (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (a).

(a) \Rightarrow (b): Using property (α) we have a natural identification

$$\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\gamma/2}) = \gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), \gamma(H, \dot{E}_{-\gamma/2}))$$

with equivalent norms. Hence the implication follows from Proposition 2.8 (1) applied to (the extension of A to) the Banach space $\gamma(H, \dot{E}_{-\gamma/2})$.

(b) \Rightarrow (c): This is trivial, as ψ belongs to $H_0^\infty(\Sigma_\theta)$ for all $\theta < \pi$;

(c) \Rightarrow (a): Let $(r_j)_{j \geq 1}$ be a Rademacher sequence on a probability space (Ω, \mathbb{P}) and let $(h_j)_{j=1}^k$ be an orthonormal system in H . Using that $\psi \in H_0^\infty(\Sigma_\theta)$, from [9, Theorem 5.2.6] we obtain

$$\begin{aligned} \sum_{j=1}^k r_j B h_j &= \sum_{j=1}^k r_j \int_0^\infty (-tA)^{3/2} (1-tA)^{-3} B h_j \frac{dt}{t} \\ &= \sum_{j=1}^k \sum_{n \in \mathbb{Z}} r_j \int_{2^n}^{2^{n+1}} (-tA)^{3/2} (1-tA)^{-3} B h_j \frac{dt}{t} \end{aligned}$$

with convergence in $L^2(\Omega; E_{-1}) = L^2(\Omega; \dot{E}_{-1}) + L^2(\Omega; E)$ (cf. (2.11)). Defining the vectors $x_n \in L^2(\Omega; E) \cap L^2(\Omega; \dot{E}_{-1})$ by

$$x_n := \sum_{j=1}^k r_j \int_{2^n}^{2^{n+1}} (-tA)^{3/2} (1-tA)^{-3} B h_j \frac{dt}{t}$$

and setting $m_N(t) = (2^{-n}t)^{1/2}$ for $t \in [2^n, 2^{n+1})$, $n = -N, \dots, N$, and $m_N(t) = 0$ for $t \notin [2^{-N}, 2^{N+1})$, we obtain (relative to the spaces $X_0 = L^2(\Omega; \dot{E}_{-1})$ and $X_1 = L^2(\Omega; E)$)

$$\begin{aligned} &\mathcal{C}_0((x_n)_{n \in \mathbb{Z}})^2 \\ &= \sup_{N \geq 1} \tilde{\mathbb{E}} \left\| \sum_{j=1}^k \sum_{n=-N}^N r_j \tilde{r}_n 2^{-n/2} \int_{2^n}^{2^{n+1}} (-tA)^{3/2} (1-tA)^{-3} B h_j \frac{dt}{t} \right\|_{L^2(\Omega; \dot{E}_{-1})}^2 \\ &= \sup_{N \geq 1} \tilde{\mathbb{E}} \left\| \sum_{j=1}^k \sum_{n=-N}^N r_j \tilde{r}_n \int_{2^n}^{2^{n+1}} (2^{-n}t)^{1/2} (-tA) (1-tA)^{-3} B h_j \frac{dt}{t} \right\|_{L^2(\Omega; \dot{E}_{-\gamma/2})}^2 \\ &= \sup_{N \geq 1} \tilde{\mathbb{E}} \left\| \sum_{j=1}^k \sum_{n=-N}^N r_j \tilde{r}_n \int_0^\infty m_N(t) (-tA) (1-tA)^{-3} \mathbf{1}_{(2^n, 2^{n+1})}(t) B h_j \frac{dt}{t} \right\|_{L^2(\Omega; \dot{E}_{-\gamma/2})}^2 \end{aligned}$$

$$\approx \sup_{N \geq 1} \mathbb{E}' \left\| \sum_{j=1}^k \sum_{n=-N}^N r'_{jn} \int_0^\infty m_N(t) (-tA)(1-tA)^{-3} \mathbf{1}_{(2^n, 2^{n+1})}(t) B h_j \frac{dt}{t} \right\|_{\dot{E}_{-1/2}}^2.$$

In the last step, property (α) was used to pass from double Rademacher sums (on $(\Omega, \mathbb{P}) \times (\tilde{\Omega}, \tilde{\mathbb{P}})$) to doubly indexed Rademacher sums (on some other probability space (Ω', \mathbb{P}')). Now, estimating Rademacher sums in terms of Gaussian sums we have

$$\begin{aligned} & \mathcal{C}_0((x_n)_{n \in \mathbb{Z}})^2 \\ & \approx \sup_{N \geq 1} \mathbb{E}' \left\| \sum_{j=1}^k \sum_{n=-N}^N \gamma'_{jn} \int_0^\infty m_N(t) (-tA)(1-tA)^{-3} \mathbf{1}_{(2^n, 2^{n+1})}(t) B h_j \frac{dt}{t} \right\|_{\dot{E}_{-1/2}}^2 \end{aligned}$$

Since the functions $\mathbf{1}_{(2^n, 2^{n+1})} \otimes h_j$ in $L^2(\mathbb{R}_+, \frac{dt}{t}; H)$ are orthonormal (up to the numerical constant $(\ln 2)^{1/2}$), one may estimate the above right-hand side by

$$\lesssim \sup_{N \geq 1} \|t \mapsto m_N(t) \phi(-tA) B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-1/2})}^2$$

where $\phi \in H_0^\infty(\Sigma_\theta)$ is given by $\phi(z) = z/(1+z)^3$. Finally, using the Kalton–Weis γ -multiplier theorem and the γ -boundedness of the operators $(-tA)^{1/2}(1-tA)^{-3/2}$, $t > 0$, (which follows from Proposition 2.6) we conclude that

$$\begin{aligned} \mathcal{C}_0((x_n)_{n \in \mathbb{Z}})^2 & \lesssim \|t \mapsto \phi(-tA) B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-1/2})}^2 \\ & \lesssim \|t \mapsto \psi(-tA) B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-1/2})}^2 \end{aligned}$$

with $\psi(z) = z^{1/2}/(1+z)^{3/2}$.

Similarly,

$$\begin{aligned} & \mathcal{C}_1((x_n)_{n \in \mathbb{Z}})^2 \\ & = \sup_{N \geq 1} \tilde{\mathbb{E}} \left\| \sum_{j=1}^k \sum_{n=-N}^N r_j \tilde{r}_n 2^{n/2} \int_{2^n}^{2^{n+1}} (-tA)^{3/2} (1-tA)^{-3} B h_j \frac{dt}{t} \right\|_{L^2(\Omega; E)}^2 \\ & = \sup_{N \geq 1} \tilde{\mathbb{E}} \left\| \sum_{j=1}^k \sum_{n=-N}^N r_j \tilde{r}_n \right. \\ & \quad \times \left. \int_0^\infty (2^{-n}t)^{-1/2} (-tA)^2 (1-tA)^{-3} \mathbf{1}_{(2^n, 2^{n+1})}(t) B h_j \frac{dt}{t} \right\|_{L^2(\Omega; \dot{E}_{-1/2})}^2 \\ & \lesssim_E \|t \mapsto \tilde{\phi}(-tA) B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-1/2})}^2 \\ & \lesssim_E \|t \mapsto \psi(-tA) B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-1/2})}^2 \end{aligned}$$

with $\tilde{\phi}(z) = z^2/(1+z)^3$ and $\psi(z) = z^{1/2}/(1+z)^{3/2}$ as before.

By Proposition 2.9 and estimating Gaussian sums by Rademacher sums, this proves that

$$\begin{aligned} \left\| \sum_{j=1}^k \gamma_j B h_j \right\|_{L^2(\Omega; \dot{E}_{-1/2})} & \approx_E \left\| \sum_{j=1}^k r_j B h_j \right\|_{L^2(\Omega; \dot{E}_{-1/2})} \\ & \lesssim_E \|t \mapsto \psi(-tA) B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-1/2})}. \end{aligned}$$

Taking the supremum over all finite orthonormal systems in H and using that E has property (α) and therefore does not contain an isomorphic copy of c_0 , we obtain (using a theorem of Hoffmann-Jørgensen and Kwapien, see [22, Theorem 4.3]) that

B is γ -radonifying as an operator from H into $\dot{E}_{-\frac{1}{2}}$ and

$$\|B\|_{\gamma(H, \dot{E}_{-\frac{1}{2}})} \lesssim \|t \mapsto \psi(-tA)B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\frac{1}{2}})}.$$

We have now proved the equivalences (a) \Leftrightarrow (b) \Leftrightarrow (c). It remains to check that these equivalent conditions imply the norm equivalence (3.1). Let μ be the centred Gaussian measure on $\dot{E}_{-\frac{1}{2}}$ associated with the γ -radonifying operator $B \in \gamma(H, \dot{E}_{-\frac{1}{2}})$. Suppose $\phi, \tilde{\phi} \in H_0^\infty(\Sigma_\theta)$ are nonzero functions. By Proposition 2.3, assertion (a) implies

$$\begin{aligned} \|t \mapsto \phi(-tA)B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\frac{1}{2}})} &\approx \int_{\dot{E}_{-\frac{1}{2}}} \|t \mapsto \phi(-tA)x\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\frac{1}{2}})} d\mu(x) \\ &\stackrel{(1)}{\approx} \int_{\dot{E}_{-\frac{1}{2}}} \|t \mapsto \tilde{\phi}(-tA)x\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\frac{1}{2}})} d\mu(x) \\ &\approx \|t \mapsto \tilde{\phi}(-tA)B\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\frac{1}{2}})}. \end{aligned}$$

Here, step (1) follows from Proposition 2.8 (2). The implied constants are independent of ϕ and $\tilde{\phi}$ under the normalisation as stated in the proposition. \square

Remark 3.3. The only step in the proof where we made use of the boundedness of the functional calculus is the Rademacher interpolation argument. For all other parts, γ -sectoriality of angle less than $\frac{\pi}{2}$ is sufficient. However, one actually needs only the continuous embedding

$$\langle L^2(\Omega; E), L^2(\Omega; \dot{E}_{-1}) \rangle_{\frac{1}{2}} \hookrightarrow L^2(\Omega; \dot{E}_{-\frac{1}{2}})$$

instead of an equality. As in Proposition 2.9 this boils down to having the embedding for the underlying Banach spaces $\langle E, \dot{E}_{-1} \rangle_{\frac{1}{2}} \hookrightarrow \dot{E}_{-\frac{1}{2}}$. An inspection of the proof of [14, Theorems 4.1 and 7.4] shows that the latter embedding does not require the full power of the boundedness of the functional calculus but merely a (discrete dyadic) square function estimate of the form

$$\sup_{\epsilon_k = \pm 1} \left\| \sum_k \epsilon_k \varphi(2^k A^\sharp) x \right\| \lesssim \|x\|$$

for some $\varphi \in H_0^\infty(\Sigma_\theta)$ for $\theta \in (0, \pi)$, where A^\sharp denotes the part of A^* in $E^\sharp = \overline{D(A^*)} \cap \overline{R(A^*)}$ (the closures are taken in the strong topology of E^*). These ‘dual’ square function estimates match the hypothesis in Le Merdy’s theorem on the Weiss conjecture [18, Theorem 4.1] in the sense that Le Merdy treats observation operators and requires upper square function estimates for A whereas we treat control operators and therefore need ‘dual’ square function estimates. The construction of A^\sharp instead of A^* is needed when non-reflexive Banach spaces are considered. On reflexive spaces one has $A^\sharp = A^*$, and the explained duality with Le Merdy’s result is more apparent.

In the next lemma, \widehat{f} denotes the Laplace transform of a function f .

Lemma 3.4 (Laplace transforms). *For all $f \in L^2(\mathbb{R}_+, \frac{dt}{t}; H)$, the function $Lf(t) := t\widehat{f}(t)$ belongs to $L^2(\mathbb{R}_+, \frac{dt}{t}; H)$ and*

$$\|Lf\|_{L^2(\mathbb{R}_+, \frac{dt}{t}; H)} \leq \|f\|_{L^2(\mathbb{R}_+, \frac{dt}{t}; H)}.$$

Proof. By the Cauchy-Schwarz inequality,

$$\begin{aligned} \int_0^\infty t^2 \|\widehat{f}(t)\|_H^2 \frac{dt}{t} &= \int_0^\infty \left\| \int_0^\infty f(s) te^{-st} ds \right\|_H^2 \frac{dt}{t} \\ &\leq \int_0^\infty \int_0^\infty \|f(s)\|_H^2 te^{-st} ds \frac{dt}{t} \end{aligned}$$

$$= \int_0^\infty \int_0^\infty \|f(s)\|_H^2 e^{-st} dt ds = \int_0^\infty \|f(s)\|_H^2 \frac{ds}{s}. \quad \square$$

As a consequence, the mapping $L : f \mapsto Lf$ is a contraction on $L^2(\mathbb{R}_+, \frac{dt}{t}; H)$. By the Kalton–Weis extension theorem, L extends to a linear contraction on the space $\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), E)$, for any Banach space E .

Proof of the equivalences (a) \Leftrightarrow (b) \Leftrightarrow (c) of Theorem 1.1. (a) \Rightarrow (b): By assumption, $t \mapsto S(t)B$ belongs to $\gamma(L^2(\mathbb{R}_+; H), E)$. It follows that $t \mapsto \eta(-tA)B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\frac{1}{2}})$, with $\eta(z) = z^{\frac{1}{2}} \exp(-z)$. The Laplace transform of $t \mapsto (tz)^{\frac{1}{2}} \exp(-tz)$ equals $\lambda \mapsto \frac{1}{2} \sqrt{\pi} z^{\frac{1}{2}} (\lambda + z)^{-\frac{3}{2}}$. Hence, by [17, Lemma 9.12] or by using the Phillips calculus (see [9]),

$$\frac{1}{2} \sqrt{\pi} (-A)^{\frac{1}{2}} (\lambda - A)^{-\frac{3}{2}} B = \int_0^\infty e^{-\lambda t} (-tA)^{\frac{1}{2}} S(t)B dt,$$

or, equivalently,

$$\frac{1}{2} \sqrt{\pi} (-A/\lambda)^{\frac{1}{2}} (1 - A/\lambda)^{-\frac{3}{2}} B = \lambda \int_0^\infty e^{-\lambda t} \eta(-tA)B dt.$$

By Lemma 3.4 and the remark following it, we obtain that $\lambda \mapsto (-A/\lambda)^{\frac{1}{2}} (1 - A/\lambda)^{-\frac{3}{2}} B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{d\lambda}{\lambda}; H), \dot{E}_{-\frac{1}{2}})$. Upon substituting $1/\lambda = \mu$ we find that $\mu \mapsto \psi(-\mu A)B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{d\mu}{\mu}; H), \dot{E}_{-\frac{1}{2}})$ with $\psi(z) = z^{\frac{1}{2}}/(1+z)^{\frac{3}{2}}$. Now (b) follows as an application of Proposition 3.2.

(b) \Rightarrow (c): From Proposition 3.2 we get that $t \mapsto (-tA)^{\frac{1}{2}} (1 - tA)^{-1} B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\frac{1}{2}})$, or equivalently, that $t \mapsto t^{\frac{1}{2}} (1 - tA)^{-1} B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), E)$. Substituting $t = 1/s$ we obtain that $s \mapsto s^{\frac{1}{2}} (s - A)^{-1} B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{ds}{s}; H), E)$.

(c) \Rightarrow (b): By substituting $t = 1/s$ the assumption implies that $s \mapsto s^{\frac{1}{2}} (1 - sA)^{-1} B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{ds}{s}; H), E)$, or equivalently, that $s \mapsto (-sA)^{\frac{1}{2}} (1 - sA)^{-1} B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{ds}{s}; H), \dot{E}_{-\frac{1}{2}})$. Then by the γ -multiplier lemma (using that the operators $(1 - sA)^{-\frac{1}{2}}$, $s > 0$, are γ -bounded by Proposition 2.6), we obtain that assumption (c) of Proposition 3.2 is satisfied.

(b) \Rightarrow (a): By Proposition 3.2, $t \mapsto (-tA)^{\frac{1}{2}} \exp(tA)B = (-tA)^{\frac{1}{2}} S(t)B$ belongs to $\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), \dot{E}_{-\frac{1}{2}})$. This is equivalent to saying that $t \mapsto S(t)B$ belongs to $\gamma(L^2(\mathbb{R}_+; H), E)$. \square

Remark 3.5. The following direct proof of the implication (a) \Rightarrow (b) of Theorem 1.1, suggested to us by Mark Veraar, avoids the use of Rademacher interpolation.

Without loss of generality we may assume that H is separable. Let $(h_n)_{n \geq 1}$ be an orthonormal basis of H and let P_n denote the orthogonal projection onto the span of h_1, \dots, h_n . Let $C_m = AR(\frac{1}{m}, A)$, so that $C_m x \rightarrow x$ strongly for all $x \in E$ (see [17, Proposition 9.4]). Set $B_{mn} := C_m B P_n$. By the γ -convergence lemma [23, Proposition 2.4] it suffices to prove the

$$\|(-A)^{-\frac{1}{2}} B_{mn}\|_{\gamma(H, E)} \lesssim \|S(\cdot) B_{mn}\|_{\gamma(L^2(\mathbb{R}_+; H), E)}, \quad (3.2)$$

with implied constant independent of m and n . The idea is to extend the action of the semigroup S from E to $\gamma(H, E)$ by putting $S(h \otimes x) := h \otimes S(t)x$. The generator of this extended semigroup, $I_H \otimes A$, belongs to $S(\gamma(H, E))$ and has a bounded H^∞ -calculus of angle $< \frac{\pi}{2}$. Hence, by Proposition 2.8 and property (α) ,

$$\begin{aligned} \|(-A)^{-\frac{1}{2}} B_{mn}\|_{\gamma(H, E)} &= \|(-A)^{\frac{1}{2}} R(\frac{1}{m}, A) B P_n\|_{\gamma(H, E)} \\ &\approx \|(-tA)^{\frac{1}{2}} S(\cdot) [A^{\frac{1}{2}} R(\frac{1}{m}, A) B P_n]\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), \gamma(H, E))} \end{aligned}$$

$$\begin{aligned}
&= \|S(\cdot)B_{mn}\|_{\gamma(L^2(\mathbb{R}_+), \gamma(H, E))} \\
&\approx \|S(\cdot)B_{mn}\|_{\gamma(L^2(\mathbb{R}_+; H), E)}.
\end{aligned}$$

This proves (3.2).

For the proofs of the implications (b) \Rightarrow (d) \Rightarrow (c) of Theorem 1.1 we need some further preparations.

An interval in \mathbb{R}_+ will be called *dyadic* (with respect to the measure $\frac{dt}{t}$) if it is of the form $[2^{k/2^M}, 2^{(k+1)/2^M})$ with $M \in \mathbb{N}$ and $k \in \mathbb{Z}$.

Lemma 3.6. *Let $-A \in S(E)$ be γ -sectorial, let $F \subseteq \mathbb{Z}$ be a finite subset, and let I_n , $n \in F$, be dyadic intervals. For any choice of the numbers $s_n, t_n \in I_n$ we have the equivalence*

$$\left\| \sum_{n \in F} \gamma_n s_n^{1/2} R(s_n, A) B \right\|_{L^2(\Omega; \gamma(H, E))} \approx \left\| \sum_{n \in F} \gamma_n t_n^{1/2} R(t_n, A) B \right\|_{L^2(\Omega; \gamma(H, E))}$$

with constants independent of the set F , the intervals I_n , and the choice of s_n, t_n .

Proof. First note that, since I_n is dyadic, $|s_n^{1/2} \pm t_n^{1/2}| \leq 4 \max\{s_n^{1/2}, t_n^{1/2}\}$.

We have, using the resolvent identity, the γ -boundedness of the operators $tR(t, A)$ for $t > 0$, and the contraction principle,

$$\begin{aligned}
&\left\| \sum_{n \in F} \gamma_n (s_n^{1/2} R(s_n, A) - t_n^{1/2} R(t_n, A)) B \right\|_{L^2(\Omega; \gamma(H, E))} \\
&\leq \left\| \sum_{n \in F} \gamma_n \frac{t_n - s_n}{t_n^{1/2} s_n^{1/2}} s_n R(s_n, A) t_n^{1/2} R(t_n, A) B \right\|_{L^2(\Omega; \gamma(H, E))} \\
&\quad + \left\| \sum_{n \in F} \gamma_n \frac{s_n^{1/2} - t_n^{1/2}}{t_n^{1/2}} t_n^{1/2} R(t_n, A) B \right\|_{L^2(\Omega; \gamma(H, E))} \\
&\lesssim \left\| \sum_{n \in F} \gamma_n t_n^{1/2} R(t_n, A) B \right\|_{L^2(\Omega; \gamma(H, E))}.
\end{aligned}$$

By the triangle inequality in $L^2(\Omega; \gamma(H, E))$ it then follows that

$$\left\| \sum_{n \in F} \gamma_n s_n^{1/2} R(s_n, A) B \right\|_{L^2(\Omega; \gamma(H, E))} \lesssim \left\| \sum_{n \in F} \gamma_n t_n^{1/2} R(t_n, A) B \right\|_{L^2(\Omega; \gamma(H, E))}.$$

The converse inequality is obtained by reversing the roles of s_n and t_n . \square

Lemma 3.7. *Let $f : \Sigma_\theta \rightarrow H$ be a bounded analytic function and suppose that, for some $0 < \eta < \theta$, the functions $t \mapsto f(e^{\pm i\eta} t)$ belong to $L^2(\mathbb{R}_+, \frac{dt}{t}; H)$. Then*

$$\sum_{n \in \mathbb{Z}} \|f(2^n)\|_H^2 < \infty.$$

Proof. Since f is continuous we may suppose that H is separable. By expanding the values of f with respect to an orthonormal basis in H , it suffices to prove the lemma for the case when H equals the scalar field.

By considering $g(z) = f(\exp(z))$, we may reformulate the problem on the strip $S_\theta = \{z \in \mathbb{C} : |\operatorname{Im} z| < \theta\}$. The objective is then to show that if the restriction of a bounded analytic function g on S_θ to the lines $\operatorname{Im} z = \pm\eta$ belongs to $L^2(\mathbb{R})$, then $\sum_{n \in \mathbb{Z}} |g(n \ln 2)|^2 < \infty$. The proof of this uses the following standard technique. By the Poisson formula for the strip we have

$$\sup_{|\zeta| < \eta} \|g|_{\{\operatorname{Im} z = \zeta\}}\|_2 < \infty$$

and therefore $g|_{S_\eta} \in L^2(S_\eta)$. For $0 < \delta < \eta$ consider the discs

$$Q_n = \{z \in \mathbb{C} : |z - n \ln 2| < \delta\}, \quad n \in \mathbb{Z},$$

centred around $n \in \mathbb{Z}$. Taking δ small enough, the functions $\phi_n = |Q_n|^{-1/2} \mathbf{1}_{Q_n}$ have disjoint support and are hence orthonormal in $L^2(S_\eta)$. By the mean value theorem we obtain

$$\begin{aligned} \sum_{n \in \mathbb{Z}} |g(n \ln 2)|^2 &= \sum_{n \in \mathbb{Z}} \left| \frac{1}{|Q_n|} \int_{Q_n} g(x + iy) dx dy \right|^2 \\ &= \frac{1}{\pi \delta^2} \sum_{n \in \mathbb{Z}} \left| \int_{S_\eta} g(x + iy) \phi_n(x + iy) dx dy \right|^2 \\ &\leq \frac{1}{\pi \delta^2} \|g|_{S_\eta}\|_{L^2(S_\eta)}^2. \quad \square \end{aligned}$$

This lemma can be restated as saying that the mapping $f \mapsto (f(2^n))_{n \in \mathbb{Z}}$ is bounded from the weighted Hardy space $H^2(\Sigma_\eta, \mu; H)$ to $\ell^2(H)$, where μ is the image on the sector Σ_η of the Lebesgue measure on the strip S_η under the exponential mapping; note that Lebesgue measure on horizontal lines in the strip S_η is mapped to the measure dt/t on rays emanating from the origin in the sector Σ_η .

By the Kalton–Weis extension theorem, this mapping extends to a bounded operator from $\gamma(H^2(\Sigma_\eta, \mu; H), E)$ to $\gamma(\ell^2(H), E)$, for any Banach space E . This is what will be needed below.

End of the proof of Theorem 1.1. We shall now prove the remaining implications (b) \Rightarrow (d) \Rightarrow (c).

We begin with the proof of (b) \Rightarrow (d). First of all, Lemma 3.1 implies that $R(t, A)B \in \gamma(H, E)$ for all $t > 0$. By the implication (b) \Rightarrow (c) applied to the operators $e^{\pm i\theta} A$ for a sufficiently small $\theta > 0$ we find that the functions

$$t \mapsto t^{1/2} R(t, e^{\pm i\theta} A)B = e^{\mp i\theta} t^{1/2} R(te^{\mp i\theta}, A)B$$

belong to $\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), E)$. By Lemma 3.7 and the remark following it, we obtain that the sequence $(2^{n/2} R(2^n, A)B)_{n \in \mathbb{Z}}$ belongs to $\gamma(\ell^2(H), E)$. But this is the same as saying that (d) holds.

We turn to the proof of (d) \Rightarrow (c). Let $S_{nm}^{(M)}$ denote the average of $t^{1/2} R(t, A)$ (with respect to dt/t) over the dyadic interval $I_{nm}^{(M)} = [2^{n+m2^{-M}}, 2^{n+(m+1)2^{-M}})$. Let $t_{nm}^{(M)} = 2^{n+m2^{-M}}$ be the left endpoint of the interval $I_{nm}^{(M)}$. Then, writing $f_I = \frac{1}{|I|} \int_I$ for the average over an interval I , we have

$$\begin{aligned} S_{nm}^{(M)} B &= \int_{I_{nm}^{(M)}} t^{1/2} R(t, A)B \frac{dt}{t} \\ &= \int_{I_{nm}^{(M)}} t^{1/2} (R(t, A)(t_{nm}^{(M)} - A)) R(t_{nm}^{(M)}, A)B \frac{dt}{t} \\ &= \left(\int_{I_{nm}^{(M)}} \frac{t^{1/2}}{(t_{nm}^{(M)})^{1/2}} \left(\frac{t_{nm}^{(M)}}{t} \cdot tR(t, A) - AR(t, A) \right) \frac{dt}{t} \right) \circ [(t_{nm}^{(M)})^{1/2} R(t_{nm}^{(M)}, A)B] \\ &=: U_{nm}^{(M)} \circ [(t_{nm}^{(M)})^{1/2} R(t_{nm}^{(M)}, A)B]. \end{aligned}$$

Since $t/t_{nm}^{(M)} \in [1, 2]$ on $I_{nm}^{(M)}$, the operators $U_{nm}^{(M)}$ belong (up to a constant) to the closure of the absolute convex hull of $\{AR(t, A), tR(t, A) : t > 0\}$. By γ -sectoriality of A (which follows from Proposition 2.6) this family is γ -bounded.

Fix a finite set $F \subseteq \mathbb{Z}$. Then,

$$\left\| \sum_{n \in F} \sum_{m=0}^{2^M-1} \mathbf{1}_{I_{nm}^{(M)}} \otimes S_{nm}^{(M)} B \right\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}; H), E)}$$

$$\begin{aligned}
& \stackrel{(1)}{\approx} \left\| \sum_{n \in F} \sum_{m=0}^{2^M-1} \mathbf{1}_{I_{nm}^{(M)}} \otimes S_{nm}^{(M)} B \right\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), \gamma(H, E))} \\
& \stackrel{(2)}{\approx} \frac{1}{2^{M/2}} \left\| \sum_{n \in F} \sum_{m=0}^{2^M-1} \gamma_{nm} S_{nm}^{(M)} B \right\|_{L^2(\Omega; \gamma(H, E))} \\
& \stackrel{(3)}{\lesssim} \frac{1}{2^{M/2}} \left\| \sum_{n \in F} \sum_{m=0}^{2^M-1} \gamma_{nm} (t_{nm}^{(M)})^{1/2} R(t_{nm}^{(M)}, A) B \right\|_{L^2(\Omega; \gamma(H, E))} \\
& \stackrel{(4)}{\approx} \frac{1}{2^{M/2}} \left\| \sum_{n \in F} \sum_{m=0}^{2^M-1} \gamma_{nm} 2^{n/2} R(2^n, A) B \right\|_{L^2(\Omega; \gamma(H, E))} \\
& \stackrel{(5)}{=} \left\| \sum_{n \in F} \sum_{m=0}^{2^M-1} \mathbf{1}_{I_{nm}^{(M)}} \otimes 2^{n/2} R(2^n, A) B \right\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), \gamma(H, E))} \\
& = \left\| \sum_{n \in F} \mathbf{1}_{I_n} \otimes 2^{n/2} R(2^n, A) B \right\|_{\gamma(L^2(\mathbb{R}_+, \frac{dt}{t}), \gamma(H, E))} \\
& \stackrel{(6)}{\approx} \left\| \sum_{n \in F} \gamma_n 2^{n/2} R(2^n, A) B \right\|_{L^2(\Omega; \gamma(H, E))}
\end{aligned}$$

with implicit constants independent of F and M . In this computation, (1) follows from property (α) ; (2), (5), (6) from the identity (2.1) along with the fact that the dyadic interval $I_{nm}^{(M)}$ has dt/t -measure $\approx 2^{-M}$; Estimate (3) follows from the γ -boundedness of the operators $U_{nm}^{(M)}$; and (4) from Lemma 3.6 applied to the points $s_n=2^n$ and $t_{nm}^{(M)}$ in $I_n = [2^n, 2^{n+1})$.

By the γ -Fatou lemma (see (2.3)), the above estimate implies (c). \square

REFERENCES

- [1] Z. Brzeźniak and J.M.A.M. van Neerven. Stochastic convolution in separable Banach spaces and the stochastic linear Cauchy problem. *Studia Math.*, 143(1):43–74, 2000.
- [2] P. Clément, B. de Pagter, F.A. Sukochev, and H. Witvliet. Schauder decompositions and multiplier theorems. *Studia Math.*, 138(2):135–163, 2000.
- [3] M. Cowling, I. Doust, A. McIntosh, and A. Yagi. Banach space operators with a bounded H^∞ functional calculus. *J. Austral. Math. Soc. Ser. A*, 60(1):51–89, 1996.
- [4] G. Da Prato and J. Zabczyk. *Ergodicity for infinite-dimensional systems*, volume 229 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1996.
- [5] R. Denk, M. Hieber, and J. Prüss. R -boundedness, Fourier multipliers and problems of elliptic and parabolic type. *Mem. Amer. Math. Soc.*, 166(788), 2003.
- [6] K.-J. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*, volume 194 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000.
- [7] B. Goldys and J.M.A.M. van Neerven. Transition semigroups of Banach space-valued Ornstein-Uhlenbeck processes. *Acta Appl. Math.*, 76(3):283–330, 2003.
- [8] B.H. Haak and J.M.A.M. van Neerven. Uniformly γ -radonifying families of operators and the stochastic Weiss conjecture. *Operators and Matrices*, 6(4):767–792, 2012.
- [9] M.H.A. Haase. *The functional calculus for sectorial operators*, volume 169 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 2006.
- [10] B. Jacob and J.R. Partington. The Weiss conjecture on admissibility of observation operators for contraction semigroups. *Integral Eq. Operator Th.*, 40(2):231–243, 2001.
- [11] B. Jacob, J.R. Partington, and S. Pott. Admissible and weakly admissible observation operators for the right shift semigroup. *Proc. Edinb. Math. Soc. (2)*, 45(2):353–362, 2002.
- [12] B. Jacob and H.J. Zwart. Counterexamples concerning observation operators for C_0 -semigroups. *SIAM J. Control Optim.*, 43(1):137–153, 2004.
- [13] N. J. Kalton and L. Weis. The H^∞ -calculus and sums of closed operators. *Math. Ann.*, 321(2):319–345, 2001.

- [14] N.J. Kalton, P.C. Kunstmann, and L. Weis. Perturbation and interpolation theorems for the H^∞ -calculus with applications to differential operators. *Math. Ann.*, 336:747–801, 2006.
- [15] N.J. Kalton and L. Weis. The H^∞ -calculus and square function estimates. In preparation.
- [16] P.C. Kunstmann and L. Weis. Perturbation theorems for maximal L_p -regularity. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 30(2):415–435, 2001.
- [17] P.C. Kunstmann and L. Weis. Maximal L_p -regularity for parabolic equations, Fourier multiplier theorems and H^∞ -functional calculus. In *Functional Analytic Methods for Evolution Equations*, volume 1855 of *Lecture Notes in Math.*, pages 65–311. Springer, Berlin, 2004.
- [18] C. Le Merdy. The Weiss conjecture for bounded analytic semigroups. *J. London Math. Soc. (2)*, 67(3):715–738, 2003.
- [19] C. Le Merdy. On square functions associated to sectorial operators. *Bull. Soc. Math. France*, 132(1):137–156, 2004.
- [20] C. Le Merdy. H^∞ Functional calculus and square function estimates for Ritt operators. *arXiv:1202.0768*, 2012.
- [21] A. McIntosh. Operators which have an H_∞ functional calculus. In *Miniconference on operator theory and partial differential equations (North Ryde, 1986)*, volume 14 of *Proc. Centre Math. Anal. Austral. Nat. Univ.*, pages 210–231. Austral. Nat. Univ., Canberra, 1986.
- [22] J.M.A.M. van Neerven. γ -Radonifying operators—a survey. In *The AMSI-ANU Workshop on Spectral Theory and Harmonic Analysis*, volume 44 of *Proc. Centre Math. Appl. Austral. Nat. Univ.*, pages 1–61. Austral. Nat. Univ., Canberra, 2010.
- [23] J.M.A.M. van Neerven, M.C. Veraar, and L. Weis. Stochastic integration in UMD Banach spaces. *Ann. Probab.*, 35(4):1438–1478, 2007.
- [24] J.M.A.M. van Neerven and L. Weis. Stochastic integration of functions with values in a Banach space. *Studia Math.*, 166(2):131–170, 2005.
- [25] J.M.A.M. van Neerven and L. Weis. Invariant measures for the linear stochastic Cauchy problem and R -boundedness of the resolvent. *J. Evolution Equ.*, 6(2):205–228, 2006.
- [26] J.M.A.M. van Neerven and L. Weis. Stochastic integration of operator-valued functions with respect to Banach space-valued Brownian motion. *Potential Anal.*, 29(1):65–88, 2008.
- [27] G. Pisier. Some results on Banach spaces without local unconditional structure. *Compositio Math.*, 37(1):3–19, 1978.
- [28] L. Weis. Operator-valued Fourier multiplier theorems and maximal L_p -regularity. *Math. Ann.*, 319(4):735–758, 2001.
- [29] G. Weiss. Two conjectures on the admissibility of control operators. In *Estimation and Control of Distributed Parameter Systems (Vorau, 1990)*, volume 100 of *Internat. Ser. Numer. Math.*, pages 367–378. Birkhäuser, Basel, 1991.
- [30] H. Zwart, B. Jacob, and O. Staffans. Weak admissibility does not imply admissibility for analytic semigroups. *Systems Control Lett.*, 48(3-4):341–350, 2003.

INSTITUTO DE MATEMÁTICA, ESTATÍSTICA E COMPUTAÇÃO CIENTÍFICA, UNIVERSIDADE ESTADUAL DE CAMPINAS, RUA SÉRGIO BUARQUE DE HOLANDA, 651, CAMPINAS SP, BRAZIL
E-mail address: `jamil@ime.unicamp.br`

INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UNIVERSITÉ BORDEAUX 1, 351, COURS DE LA LIBÉRATION, F-33405 TALENCE CEDEX, FRANCE
E-mail address: `bernhard.haak@math.u-bordeaux1.fr`

DELFT INSTITUTE OF APPLIED MATHEMATICS, DELFT UNIVERSITY OF TECHNOLOGY, P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS
E-mail address: `J.M.A.M.vanNeerven@TUDelft.nl`