



Spinc-quantization and the K-multiplicities of the discrete series

Paul-Emile Paradan

► To cite this version:

Paul-Emile Paradan. Spinc-quantization and the K-multiplicities of the discrete series. *Annales Scientifiques de l'École Normale Supérieure*, 2003, 36 (5), pp.805-845. 10.1016/j.ansens.2003.03.001 . hal-00773251

HAL Id: hal-00773251

<https://hal.science/hal-00773251>

Submitted on 12 Jan 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SPIN^c-QUANTIZATION AND THE K -MULTIPLICITIES OF THE DISCRETE SERIES

PAUL-EMILE PARADAN

UMR 5582, Institut Fourier, B.P. 74, 38402, Saint-Martin-d'Hères cedex, France
e-mail: Paul-Emile.Paradan@ujf-grenoble.fr

November 2001

ABSTRACT. We express the K -multiplicities of a representation of the discrete series associated to a coadjoint orbit \mathcal{O} in terms of Spin^c-index on symplectic reductions of \mathcal{O} .

CONTENTS

1. Introduction and statement of the results	1
2. Spin ^c -quantization of compact Hamiltonian K -manifolds	7
3. Quantization of non-compact Hamiltonian K -manifolds	11
4. Quantization commutes with reduction	20
5. Quantization and the discrete series	33
References	41

1. INTRODUCTION AND STATEMENT OF THE RESULTS

The purpose of this paper is to show that the ‘*quantization commutes with reduction*’ principle of Guillemin-Sternberg [16] holds for the coadjoint orbits that parametrize the discrete series of a real connected semi-simple Lie group.

1.1. Discrete series and K -multiplicities. Let G be a connected, real, semisimple Lie group with finite center. By definition, the *discrete series* of G is the set of isomorphism classes of irreducible, square integrable, unitary representations of G . Let K be a maximal compact subgroup of G , and T be a maximal torus in K . Harish-Chandra has shown that G has a discrete series if and only if T is a Cartan subgroup of G [19]. For the remainder of this paper, we may therefore assume that T is a Cartan subgroup of G .

Let us fix some notation. We denote by $\mathfrak{g}, \mathfrak{k}, \mathfrak{t}$ the Lie algebras of G, K, T , and by $\mathfrak{g}^*, \mathfrak{k}^*, \mathfrak{t}^*$ their duals. Let $\Lambda^* \subset \mathfrak{t}^*$ be the set of real weights: $\alpha \in \Lambda^*$ if $i\alpha$ is the differential of a character of T . Let $\mathfrak{R}_c \subset \mathfrak{R} \subset \Lambda^*$ be respectively the set of roots

Keywords : moment map, reduction, geometric quantization, discrete series, transversally elliptic symbol.

1991 Mathematics Subject Classification : 58F06, 57S15, 19L47.

for the action of T on $\mathfrak{k} \otimes \mathbb{C}$ and $\mathfrak{g} \otimes \mathbb{C}$. We choose a system of positive roots \mathfrak{R}_c^+ for \mathfrak{R}_c . We denote by \mathfrak{t}_+^* the corresponding Weyl chamber, and we let ρ_c be half the sum of the elements of \mathfrak{R}_c^+ . The set $\Lambda_+^* := \Lambda^* \cap \mathfrak{t}_+^*$ parametrizes the unitary dual of K . For $\mu \in \Lambda_+^*$, let χ_μ^κ be the character of the irreducible K -representation with highest weight μ .

Harish-Chandra parametrizes the discrete series by a discrete subset \widehat{G}_d of regular elements of the Weyl chamber \mathfrak{t}_+^* [19]. He associates to any $\lambda \in \widehat{G}_d$ an invariant eigendistribution on G , denoted by Θ_λ , which is shown to be the global trace of an irreducible, square integrable, unitary representation \mathcal{H}_λ of G . It is a generalized function on G , invariant by conjugation, which admits a restriction to K denoted by $\Theta_\lambda|_K$. The distribution $\Theta_\lambda|_K$ corresponds to the global trace of the induced representation of K on \mathcal{H}_λ . It admits a decomposition

$$\Theta_\lambda|_K = \sum_{\mu \in \Lambda_+^*} m_\mu(\lambda) \chi_\mu^\kappa,$$

where the integers $m_\mu(\lambda)$ satisfy certain combinatorial identities called the Blattner formulas [20].

The main goal of the paper is to relate the multiplicities $m_\mu(\lambda)$ to the geometry of the coadjoint orbit $G \cdot \lambda \subset \mathfrak{g}^$ as predicted by the Guillemin-Sternberg principle evoked above.*

Before stating our result we recall how a representation belonging to the discrete series can be realized as the quantization of a coadjoint orbit.

1.2. Realisation of the discrete series. In the 60's, Kostant and Langlands conjectured realisations of the discrete series in terms of L^2 cohomology that fit into the general framework of quantization. The proof of this conjecture was given by Schmid some years later [34, 35]. Let us recall the procedure for a fixed $\lambda \in \widehat{G}_d$.

The manifold $G \cdot \lambda$ carries several G -invariant complex structures. For convenience we work with the complex structure J defined by the following condition: each weight α for the T -action on the tangent space $(\mathbf{T}_\lambda(G \cdot \lambda), J)$ satisfies $(\alpha, \lambda) > 0$.

Let $\mathfrak{R}^+ \subset \mathfrak{R}$ be the set of positive roots defined by λ : $\alpha \in \mathfrak{R}^+ \iff (\alpha, \lambda) > 0$. Let ρ be half the sum of the elements of \mathfrak{R}^+ . The condition $\lambda \in \widehat{G}_d$ imposes that $\lambda - \rho$ is a weight for T , so we can consider the line bundle

$$\tilde{L} := G \times_T \mathbb{C}_{\lambda - \rho}$$

over $G \cdot \lambda \simeq G/T$: this line bundle carries a canonical holomorphic structure. Let $\Omega^k(\tilde{L})$ be the space of \tilde{L} -valued $(0, k)$ forms on $G \cdot \lambda$, and $\bar{\partial}_{\tilde{L}} : \Omega^k(\tilde{L}) \rightarrow \Omega^{k+1}(\tilde{L})$ be the Dolbeault operator. The choice of G -invariant hermitian metrics on $G \cdot \lambda$ and on \tilde{L} give meaning to the formal adjoint $\bar{\partial}_{\tilde{L}}^*$ of the $\bar{\partial}_{\tilde{L}}$ operator, and to the Dolbeault-Dirac operator $\bar{\partial}_{\tilde{L}} + \bar{\partial}_{\tilde{L}}^*$.

The L^2 cohomology of \tilde{L} , which we denote by $H_{(2)}^*(G \cdot \lambda, \tilde{L})$, is equal to the kernel of the differential operator $\bar{\partial}_{\tilde{L}} + \bar{\partial}_{\tilde{L}}^*$ acting on the subspace of $\Omega^*(\tilde{L})$ formed by the square integrable elements.

Theorem 1.1. (*Schmid*).

Let $\lambda \in \widehat{G}_d$.

(i) $H_{(2)}^k(G \cdot \lambda, \tilde{L}) = 0$ if $k \neq \frac{\dim(G/K)}{2}$.

(ii) If $k = \frac{\dim(G/K)}{2}$, then $H_{(2)}^k(G \cdot \lambda, \tilde{L})$ is the irreducible representation \mathcal{H}_λ .

So, the representation \mathcal{H}_λ is the quantization of the coadjoint orbit $G \cdot \lambda$ being the index of the Dolbeault-Dirac operator $\bar{\partial}_L + \bar{\partial}_L^*$ (in the L^2 sense and modulo $(-1)^{\frac{\dim(G/K)}{2}}$). In the next subsection, we briefly recall the ‘quantization commutes with reduction’ principle of Guillemin-Sternberg, and in subsection 1.4 we state our main result.

1.3. Quantization commutes with reduction. Let M be a Hamiltonian K -manifold with symplectic form ω and moment map $\Phi : M \rightarrow \mathfrak{k}^*$. The coadjoint orbits $G \cdot \lambda$ introduced earlier are the key examples here. Each is equipped with its Kirillov-Kostant-Souriau symplectic form ω , and the action of G is Hamiltonian with moment map $G \cdot \lambda \hookrightarrow \mathfrak{g}^*$ equal to the inclusion. Let K be the maximal compact Lie subgroup of G introduced in subsection 1.2. The induced action of K on $G \cdot \lambda$ is Hamiltonian, and the corresponding moment map $\Phi : G \cdot \lambda \rightarrow \mathfrak{k}^*$ is equal to the composition of the inclusion $G \cdot \lambda \hookrightarrow \mathfrak{g}^*$ with the projection $\mathfrak{g}^* \rightarrow \mathfrak{k}^*$.

In the process of quantization one tries to associate a unitary representation of K to the data (M, ω, Φ) . In this general framework, when M is *compact* and under certain integrability conditions, we associate to these data a virtual representation of K defined as the equivariant index of a Spin^c Dirac operator: it’s the Spin^c quantization. We need two auxilliary data :

- (i) A *prequantum line bundle* $L \rightarrow M$: it is a K -equivariant Hermitian line bundle equipped with K -invariant connection whose curvature form is $-i\omega$.
- (ii) A K -invariant almost complex structure J on M , *compatible* with the symplectic structure: $(v, w) \mapsto \omega(v, Jw)$ defines a metric.

One considers then the K -equivariant Spin^c Dirac operator D_L corresponding to the Spin^c structure on M defined by J , and twisted by the line bundle L [25, 14]. The Spin^c-quantization of (M, ω, Φ) is the equivariant index of the differential operator \mathcal{D}_L

$$RR^K(M, L) := \text{Index}_M^K(\mathcal{D}_L) \in R(K),$$

where $R(K)$ is the representation ring of K . When K is reduced to $\{e\}$, the Spin^c-quantization of (M, ω) is just an integer: $RR(M, L) \in \mathbb{Z}$.

A fundamental result of Marsden-Weinstein asserts that if $\xi \in \mathfrak{k}^*$ is a regular value of the moment map Φ , the *reduced space*

$$M_\xi := \Phi^{-1}(\xi)/K_\xi \cong \Phi^{-1}(K \cdot \xi)/K$$

is an orbifold equipped with a symplectic structure ω_ξ (which one calls also symplectic quotient). For any dominant weight $\mu \in \Lambda_+^*$ which is a regular value of Φ ,

$$L_\mu := (L|_{\Phi^{-1}(\mu)} \otimes \mathbb{C}_{-\mu})/K_\mu$$

is a prequantum orbifold-line bundle over (M_μ, ω_μ) . The definition of Spin^c-index carries over to the orbifold case, hence $RR(M_\mu, L_\mu) \in \mathbb{Z}$ is defined. In [29], this is extended further to the case of singular symplectic quotients, using partial (or shift) desingularization. So the integer $RR(M_\mu, L_\mu) \in \mathbb{Z}$ is well defined for every $\mu \in \Lambda_+^*$.

The following Theorem was conjectured by Guillemin-Sternberg [16] and is known as “quantization commutes with reduction” [28, 29].

Theorem 1.2. (*Meinrenken, Meinrenken-Sjamaar*). *Let (M, ω, Φ) be a compact Hamiltonian K -manifold prequantized by L . Let $RR^K(M, -)$ be the equivariant Riemann-Roch character defined by means of a compatible almost complex structure on M . We have the following equality in $R(K)$*

$$RR^K(M, L) = \sum_{\mu \in \Lambda_+^*} RR(M_\mu, L_\mu) \chi_\mu^K .$$

Remark 1.3. *For a compact Hamiltonian K -manifold (M, ω, Φ) , the Convexity Theorem [23] asserts that $\Delta := \Phi(M) \cap \mathfrak{t}_+^*$ is a convex rational polytope. In Theorem 1.2, we have $RR(M_\mu, L_\mu) = 0$ if $\mu \notin \Delta$.*

Other proofs can be found in [33, 40]. For an introduction and further references see [37, 43].

A natural question is to extend Theorem 1.2 to the *non-compact* Hamiltonian K -manifolds which admit a *proper* moment map. In this situation, the reduced space $M_\xi := \Phi^{-1}(\xi)/K_\xi$ is compact for every $\xi \in \mathfrak{t}^*$, so the integer $RR(M_\mu, L_\mu) \in \mathbb{Z}$, $\mu \in \Lambda_+^*$, can be defined like before.

Conjecture 1.4. *Let (M, ω, Φ) be a Hamiltonian K -manifold with proper moment map, and prequantized by L . Let $\bar{\partial}_L + \bar{\partial}_L^*$ be the Dolbeault-Dirac operator defined by means of a K -invariant compatible almost complex structure, and K -invariant metric on M and L . Then*

$$L^2 - \text{Index}^K(\bar{\partial}_L + \bar{\partial}_L^*) = \sum_{\mu \in \Lambda_+^*} RR(M_\mu, L_\mu) \chi_\mu^K .$$

We present in the next subsection the central result of this paper that shows that Conjecture 1.4 is true, apart from a ρ_c -shift, for the coadjoint orbits that parametrize the discrete series.

1.4. The results. Consider the Hamiltonian action of K on the coadjoint orbit $G \cdot \lambda$. Since $G \cdot \lambda$ is closed in \mathfrak{g}^* , the moment map $\Phi : G \cdot \lambda \rightarrow \mathfrak{t}^*$ is *proper* [32]. Our main Theorem can be stated roughly as follows.

Theorem 1.5. *Let $m_\mu(\lambda), \mu \in \Lambda_+^*$, be the K -multiplicities of the representation $\mathcal{H}_\lambda|_K$. For $\mu \in \Lambda_+^*$ we have:*

(i) *If $\mu + \rho_c$ is a regular value of Φ , the orbifold $(G \cdot \lambda)_{\mu + \rho_c} := \Phi^{-1}(\mu + \rho_c)/T$, oriented by its symplectic form $\omega_{\mu + \rho_c}$, carries a Spin^c structure such that*

$$m_\mu(\lambda) = \mathcal{Q}((G \cdot \lambda)_{\mu + \rho_c}) ,$$

where the RHS is the index of the corresponding Spin^c Dirac operator on the reduced space $(G \cdot \lambda)_{\mu + \rho_c}$.

(ii) *In general, one can define an integer $\mathcal{Q}((G \cdot \lambda)_{\mu + \rho_c}) \in \mathbb{Z}$, as the index of a Spin^c Dirac operator on a reduced space $(G \cdot \lambda)_\xi$ where ξ is a regular value of Φ , close enough to $\mu + \rho_c$. We still have $m_\mu(\lambda) = \mathcal{Q}(M_{\mu + \rho_c})$.*

Our Theorem states that the decomposition of $\Theta_\lambda|_K$ into K -irreducible components follows the philosophy of Guillemin-Sternberg:

$$(1) \quad \Theta_\lambda|_K = \sum_{\mu \in \Lambda_+^*} \mathcal{Q}((G \cdot \lambda)_{\mu + \rho_c}) \chi_\mu^K .$$

We know from Theorem 1.1 that $\Theta_\lambda|_K = (-1)^{\frac{\dim(G/K)}{2}} L^2 - \text{Index}^K(\bar{\partial}_{\tilde{L}} + \bar{\partial}_{\tilde{L}}^*)$, hence Theorem 1.5 states also that

$$(2) \quad L^2 - \text{Index}^K(\bar{\partial}_{\tilde{L}} + \bar{\partial}_{\tilde{L}}^*) = (-1)^{\frac{\dim(G/K)}{2}} \sum_{\mu \in \Lambda_+^*} \mathcal{Q}((G \cdot \lambda)_{\mu+\rho_c}) \chi_\mu^K.$$

The main difference between Conjecture 1.4 and Theorem 1.5 is the ρ_c -shift and the choices of Spin^c structure on the symplectic quotients M_μ and $(G \cdot \lambda)_{\mu+\rho_c}$.

The ρ_c -shift is due to the fact that the line bundle \tilde{L} is not a prequantum line bundle over $(G \cdot \lambda, \omega)$. The difference on the choice of Spin^c structure comes from the fact that the complex structure J on $G \cdot \lambda$ is not compatible with the symplectic structure (unless $G = K$ is compact). Hence J does not descend to the symplectic reductions $(G \cdot \lambda)_{\mu+\rho_c}$ in general: the choice of the Spin^c structure on them need some care (see Propositions 4.10 and 4.11).

Remark 1.6. *For a Hamiltonian K -manifold M with proper moment map Φ , the Convexity Theorem [23, 26, 38] asserts that $\Delta := \Phi(M) \cap \mathfrak{t}_+^*$ is a convex rational polyhedron. In Theorem 1.5, we have $\mathcal{Q}((G \cdot \lambda)_{\mu+\rho_c}) = 0$ if $\mu + \rho_c$ does not belong to the relative interior of Δ (see Prop. 2.4).*

1.5. Outline of the Proof. We have to face the following difficulties:

- [1] The symplectic manifold $G \cdot \lambda$ is not compact.
- [2] The complex structure on $G \cdot \lambda$ is not compatible with the symplectic form ω . In other words, the Kirillov-Kostant-Souriau symplectic form does not define a Kähler structure on $G \cdot \lambda$ unless $G = K$ is compact.
- [3] The line bundle \tilde{L} is not a prequantum line bundle over $(G \cdot \lambda, \omega)$. It's what we call in the rest of this paper a κ -prequantum¹ line bundle over $(G \cdot \lambda, \omega, J)$: if κ denotes the canonical line bundle of $(G \cdot \lambda, J)$, the tensor product $\tilde{L}^2 \otimes \kappa^{-1}$ is a prequantum line bundle over $(G \cdot \lambda, 2\omega)$.

The first step of the proof is to solve the difficulties [2] and [3] in the *compact* situation. In Section 2, we give a modified version of Theorem 1.2 when (M, ω, Φ) is a *compact* Hamiltonian K -manifold which is equipped with an almost complex structure J - not necessarily compatible with ω - and a κ -prequantum line bundle \tilde{L} .

Theorem 1.7. *Let $RR^K(M, -)$ be the Riemann-Roch character defined by J . If the infinitesimal stabilizers for the action of K on M are Abelian, we have*

$$(3) \quad RR^K(M, \tilde{L}) = \varepsilon \sum_{\mu \in \Lambda_+^*} Q(M_{\mu+\rho_c}) \chi_\mu^K,$$

where $\varepsilon = \pm 1$ is the ‘quotient’ of the orientations induced by the almost complex structure, and the symplectic form.

In (3), the integer $Q(M_{\mu+\rho_c})$ are computed like in Theorem 1.5 (see Def. 2.4 for a more precise definition).

In the second step of the proof, we extend (3) to a non-compact setting. Instead of working with the L^2 -Index, we define in Section 3 a *generalized Riemann-Roch character* $RR_\Phi^K(M, -)$ when (M, ω, Φ) is a Hamiltonian manifold such that the

¹Formally, \tilde{L} is the tensor product of a prequantum line bundle over $(G \cdot \lambda, \omega)$ with a square root of κ .

function $\|\Phi\|^2: M \rightarrow \mathbb{R}$ has a *compact* set of critical points. For every K -vector bundle $E \rightarrow M$, the distribution $RR_\Phi^K(M, E)$ is defined as the index of a transversally elliptic operator on M [1]. When the manifold is compact, the maps $RR_\Phi^K(M, -)$ and $RR^K(M, -)$ coincide.

We prove in Section 4 that Theorem 1.7 generalizes to

Theorem 1.8. *Let (M, ω, Φ) be a Hamiltonian K -manifold with proper moment map and such that the function $\|\Phi\|^2: M \rightarrow \mathbb{R}$ has a compact set of critical points. If the infinitesimal stabilizers are Abelian, and under Assumption 3.6, we have*

$$(4) \quad RR_\Phi^K(M, \tilde{L}) = \varepsilon \sum_{\mu \in \Lambda_+^*} \mathcal{Q}(M_{\mu+\rho_c}) \chi_\mu^K,$$

for every κ -prequantum line bundle.

In contrast to (3), the RHS of (4) is in general an infinite sum. Assumption 3.6 is needed to control the data on the non-compact manifold M : it asserts in particular that for any coadjoint orbit \mathcal{O} of K , the square of the moment map $\Phi_{\mathcal{O}}: M \times \mathcal{O} \rightarrow \mathfrak{k}^*$, $(m, \xi) \mapsto \Phi(m) - \xi$, has a *compact* set of critical points.

In the final section we consider, for $\lambda \in \widehat{G}_d$, the case of the coadjoint orbit $G \cdot \lambda$ with the Hamiltonian K -action. The moment map Φ is *proper* and the critical set of $\|\Phi\|^2$ coincides with $K \cdot \lambda$, hence is compact. Thus the generalized Riemann-Roch character $RR_\Phi^K(G \cdot \lambda, -)$ is well defined, and we want to investigate the index $RR_\Phi^K(M, \tilde{L})$ for the κ -prequantum line bundle $\tilde{L} := G \times_T \mathbb{C}_{\lambda-\rho}$.

On one hand we are able to compute $RR_\Phi^K(G \cdot \lambda, \tilde{L})$ explicitly in term of the holomorphic induction map Hol_T^K . Let \mathfrak{p} be the orthogonal complement of \mathfrak{k} in \mathfrak{g} . It inherits a complex structure and an action of the torus T . The element $\wedge_{\mathbb{C}}^\bullet \mathfrak{p} \in R(T)$ admits a polarized inverse $[\wedge_{\mathbb{C}}^\bullet \mathfrak{p}]_\lambda^{-1} \in R^{-\infty}(T)$ (see [33][Section 5]). In Subsection 5.2 we prove that

$$(5) \quad RR_\Phi^K(G \cdot \lambda, \tilde{L}) = (-1)^{\frac{\dim(G/K)}{2}} \text{Hol}_T^K \left(t^{\lambda-\rho_c+\rho_n} [\wedge_{\mathbb{C}}^\bullet \mathfrak{p}]_\lambda^{-1} \right),$$

where $\rho_n = \rho - \rho_c$ is half the sum of the non-compact roots. On the other hand, we show (Lemma 5.4) that the Blattner formulas can be reinterpreted through Hol_T^K as follows:

$$(6) \quad \Theta_\lambda|_K = \text{Hol}_T^K \left(t^{\lambda-\rho_c+\rho_n} [\wedge_{\mathbb{C}}^\bullet \mathfrak{p}]_\lambda^{-1} \right).$$

From (5) and (6) we obtain

$$(7) \quad RR_\Phi^K(G \cdot \lambda, \tilde{L}) = (-1)^{\frac{\dim(G/K)}{2}} \Theta_\lambda|_K = L^2 - \text{Index}^K(\bar{\partial}_{\tilde{L}} + \bar{\partial}_{\tilde{L}}^*).$$

Since in this context $\varepsilon = (-1)^{\frac{\dim(G/K)}{2}}$, the Theorem follows from (4) and (7), provided one verifies that Assumption 3.6 holds for $G \cdot \lambda$. This is done in the final subsection of this paper.

Acknowledgments. I would like to thank Michèle Vergne for suggesting this problem, and helpful discussions.

Notation

Throughout the paper, K will denote a compact, connected Lie group, and \mathfrak{k} its Lie algebra. In Sections 2, 3, and 4, we consider a K -Hamiltonian action on a manifold M . And we use there the following notation.

- T : maximal torus of K with Lie algebra \mathfrak{t}
- W : Weyl group of (K, T)
- $\Lambda = \ker(\exp : \mathfrak{t} \rightarrow T)$: integral lattice of \mathfrak{t}
- $\Lambda^* = \text{hom}(\Lambda, 2\pi\mathbb{Z})$: real weight lattice
- \mathfrak{t}_+^*, ρ_c : Weyl chamber and corresponding half sum of the positive roots
- $\Lambda_+^* = \Lambda^* \cap \mathfrak{t}_+^*$: set of positive weights
- χ_μ^K : character of the irreducible K -representation with highest weight $\mu \in \Lambda_+^*$
- \mathbb{T}_β : subtorus of T generated by $\beta \in \mathfrak{t}$
- M^γ : submanifold of points fixed by $\gamma \in \mathfrak{k}$
- TM : tangent bundle of M
- $\mathbf{T}_K M$: set of tangent vectors orthogonal to the K -orbits in M
- Φ : moment map
- \tilde{L} : κ -prequantum line bundle
- $\tilde{\mathbb{C}}_{[\mu]} = K \times_T \mathbb{C}_\mu$: κ -prequantum line bundle over the coadjoint orbit $K \cdot (\mu + \rho_c)$
- $\text{Cr}(\|\Phi\|^2)$: critical set of the function $\|\Phi\|^2$
- $\Delta = \Phi(M) \cap \mathfrak{t}_+^*$: moment polytope
- \mathcal{H} : vector field generated by Φ
- $m_\mu(E)$: multiplicity of $RR_\Phi^K(M, E)$ relatively to $\mu \in \Lambda_+^*$.

In the final section, we consider the particular case of the K -action on $M := G \cdot \lambda$. Here G is a connected real semi-simple Lie group with finite center admitting K as a maximal compact subgroup, and T as a compact Cartan subgroup.

Let us recall the definition of the holomorphic induction map Hol_T^K . Every $\mu \in \Lambda^*$ defines a 1-dimensional T -representation, denoted \mathbb{C}_μ , where $t = \exp X$ acts by $t^\mu := e^{i\langle \mu, X \rangle}$. We denote by $R(K)$ (resp. $R(T)$) the ring of characters of finite-dimensional K -representations (resp. T -representations). We denote $R^{-\infty}(K)$ (resp. $R^{-\infty}(T)$) the set of generalized characters of K (resp. T). An element $\chi \in R^{-\infty}(K)$ is of the form $\chi = \sum_{\mu \in \Lambda_+^*} m_\mu \chi_\mu^K$, where $\mu \mapsto m_\mu, \Lambda_+^* \rightarrow \mathbb{Z}$ has at most polynomial growth. Likewise, an element $\chi \in R^{-\infty}(T)$ is of the form $\chi = \sum_{\mu \in \Lambda^*} m_\mu t^\mu$, where $\mu \mapsto m_\mu, \Lambda^* \rightarrow \mathbb{Z}$ has at most polynomial growth. We denote $w \circ \mu = w(\mu + \rho_c) - \rho_c$ the affine action of the Weyl group on Λ^* . The holomorphic induction map

$$\text{Hol}_T^K : R^{-\infty}(T) \longrightarrow R^{-\infty}(K)$$

is characterized by the following properties:

- i) $\text{Hol}_T^K(t^\mu) = \chi_\mu^K$ for every $\mu \in \Lambda_+^*$,
- ii) $\text{Hol}_T^K(t^{w \circ \mu}) = (-1)^w \text{Hol}_T^K(t^\mu)$ for every $w \in W$ and $\mu \in \Lambda^*$,
- iii) $\text{Hol}_T^K(t^\mu) = 0$ if $W \circ \mu \cap \Lambda_+^* = \emptyset$.

2. Spin^c-QUANTIZATION OF COMPACT HAMILTONIAN K -MANIFOLDS

In this Section we give a modified version of the ‘quantization commutes with reduction’ principle.

Let M be a compact Hamiltonian K -manifold with symplectic form ω and moment map $\Phi : M \rightarrow \mathfrak{k}^*$ characterized by the relation $d\langle \Phi, X \rangle = -\omega(X_M, -)$, where X_M is the vector field on M generated by $X \in \mathfrak{k} : X_M(m) := \frac{d}{dt} \exp(-tX).m|_{t=0}$, for $m \in M$.

Let J be a K -invariant almost complex structure on M which is not assumed to be compatible with the symplectic form. We denote $RR^K(M, -)$ the Riemann-Roch character defined by J . Let us recall the definition of this map.

Let $E \rightarrow M$ be a complex K -vector bundle. The almost complex structure on M gives the decomposition $\wedge \mathbf{T}^*M \otimes \mathbb{C} = \oplus_{i,j} \wedge^{i,j} \mathbf{T}^*M$ of the bundle of differential forms. Using Hermitian structure in the tangent bundle $\mathbf{T}M$ of M , and in the fibers of E , we define a Dolbeault-Dirac operator $\bar{\partial}_E + \bar{\partial}_E^* : \mathcal{A}^{0,even}(M, E) \rightarrow \mathcal{A}^{0,odd}(M, E)$, where $\mathcal{A}^{i,j}(M, E) := \Gamma(M, \wedge^{i,j} \mathbf{T}^*M \otimes_{\mathbb{C}} E)$ is the space of E -valued forms of type (i, j) . The Riemann-Roch character $RR^K(M, E)$ is defined as the index of the elliptic operator $\bar{\partial}_E + \bar{\partial}_E^*$:

$$RR^K(M, E) = \text{Index}_M^K(\bar{\partial}_E + \bar{\partial}_E^*) \in R(K)$$

viewed as an element of $R(K)$, the character ring of K . An alternative definition goes as follows. The almost complex structure defines a canonical invariant Spin^c structure². The Spin^c Dirac operator of M with coefficient in E has the same principal symbol as $\sqrt{2}(\bar{\partial}_E + \bar{\partial}_E^*)$ (see e.g. [14]), and therefore has the same equivariant index.

In the Kostant-Souriau framework, M is prequantized if there is a K -equivariant Hermitian line bundle L with a K -invariant Hermitian connection ∇^L of curvature $-i\omega$. The line bundle L is called a prequantum line bundle for the Hamiltonian K -manifold (M, ω, Φ) . Recall that the data (∇^L, Φ) are related by the Kostant formula

$$(2.1) \quad \mathcal{L}^L(X) - \nabla_{X_M}^L = i\langle \Phi, X \rangle, \quad X \in \mathfrak{k}.$$

Here $\mathcal{L}^L(X)$ is the infinitesimal action of X on the section of $L \rightarrow M$.

The tangent bundle $\mathbf{T}M$ endowed with J is a complex vector bundle over M , and we consider its complex dual $\mathbf{T}_{\mathbb{C}}^*M := \text{hom}_{\mathbb{C}}(\mathbf{T}M, \mathbb{C})$. We suppose first that the canonical line bundle $\kappa := \det \mathbf{T}_{\mathbb{C}}^*M$ admits a K -equivariant square root $\kappa^{1/2}$. If M is prequantized by L , a standard procedure in the geometric quantization literature is to tensor L by the bundle of half-forms $\kappa^{1/2}$ [45]. We consider the index $RR^K(M, L \otimes \kappa^{1/2})$ instead of $RR^K(M, L)$. In many contexts, the tensor product $\tilde{L} = L \otimes \kappa^{1/2}$ has a meaning even if L nor $\kappa^{1/2}$ exist.

Definition 2.1. *An Hamiltonian K -manifold (M, ω, Φ) , equipped with an almost complex structure, is κ -prequantized by an equivariant line bundle \tilde{L} if $L_{2\omega} := \tilde{L}^2 \otimes \kappa^{-1}$ is a prequantum line bundle for $(M, 2\omega, 2\Phi)$.*

The basic examples are the regular coadjoint orbits of K . For any $\mu \in \Lambda_+^*$, consider the regular coadjoint orbit $\mathcal{O}^{\mu+\rho_c} := K \cdot (\mu + \rho_c)$ with the compatible complex structure. The line bundle $\tilde{\mathbb{C}}_{[\mu]} = K \times_T \mathbb{C}_{\mu}$ is a κ -prequantum line bundle over $\mathcal{O}^{\mu+\rho_c}$, and we have

$$(2.2) \quad RR^K(\mathcal{O}^{\mu+\rho_c}, \tilde{\mathbb{C}}_{[\mu]}) = \chi_{\mu}^K$$

²See subsection 4.1 for a short review on the notion of Spin^c structure.

for any $\mu \in \Lambda_+^*$.

Definition 2.1 can be rewritten in the Spin^c setting (see subsection 4.1 for a brief review on Spin^c-structures). The almost complex structure induces a Spin^c structure P with canonical line bundle $\det_{\mathbb{C}} \mathbf{T}M = \kappa^{-1}$. If (M, ω, J) is κ -prequantized by \tilde{L} one can twist P by \tilde{L} , and then define a new Spin^c structure with canonical line bundle $\kappa^{-1} \otimes \tilde{L}^2 = L_{2\omega}$ (see Lemma 4.2).

Definition 2.2. *A symplectic manifold (M, ω) is Spin^c-prequantized if there exists a Spin^c structure with canonical line bundle $L_{2\omega}$ which is a prequantum line bundle on $(M, 2\omega)$. If a compact Lie group acts on M , the Spin^c-structure is required to be equivariant. Here we take the symplectic orientation on M .*

When (M, ω, J) is κ -prequantized by \tilde{L} , one wants to compute the K -multiplicities of $RR^K(M, \tilde{L})$ in geometrical terms, like in Theorem 1.2.

Definition 2.3. *An element $\xi \in \mathfrak{k}^*$ is a quasi-regular value of Φ if all the K_ξ -orbits in $\Phi^{-1}(\xi)$ have the same dimension. A quasi-regular value is generic if the submanifold $\Phi^{-1}(\xi)$ is of maximal dimension.*

For any quasi-regular value $\xi \in \mathfrak{k}^*$, the reduced space $M_\xi := \Phi^{-1}(\xi)/K_\xi$ is an orbifold equipped with a symplectic structure ω_ξ . Let \tilde{L} be a κ -prequantum line bundle over M , and let $L_{2\omega} := \tilde{L}^2 \otimes \kappa^{-1}$ be the corresponding prequantum line bundle for $(M, 2\omega)$. For any dominant weight $\mu \in \Lambda_+^*$ such that $\mu + \rho_c$ is a quasi-regular value of Φ ,

$$(L_{2\omega}|_{\Phi^{-1}(\mu+\rho_c)} \otimes \mathbb{C}_{-2(\mu+\rho_c)})/T$$

is a prequantum orbifold-line bundle over $(M_{\mu+\rho_c}, 2\omega_{\mu+\rho_c})$.

The following Proposition is the main point for computing the K -multiplicities of $RR^K(M, \tilde{L})$ in terms of the reduced spaces $M_{\mu+\rho_c} := \Phi^{-1}(\mu + \rho_c)/T$, $\mu \in \Lambda_+^*$. It deals with the coherence of the definition of an integer valued map $\mu \in \Lambda_+^* \mapsto \mathcal{Q}(M_{\mu+\rho_c})$. In the next proposition we suppose that (M, ω, Φ) is a Hamiltonian K -manifold with *proper* moment map. The set $\Phi(M) \cap \mathfrak{t}_+^*$ is denoted by Δ . By the Convexity Theorem [23, 26, 38] it is a convex rational polyhedron, referred to as the *moment polyhedron*.

Definition-Proposition 2.4. *Let (M, ω, Φ) be a Hamiltonian K -manifold, with proper moment map. We denote Δ° the relative interior of the moment polyhedron $\Delta := \Phi(M) \cap \mathfrak{t}_+^*$. Let \tilde{L} be a κ -prequantum line bundle relative to an almost complex structure J . Let $\mu \in \Lambda_+^*$.*

- *If $\mu + \rho_c \notin \Delta^\circ$, we set $\mathcal{Q}(M_{\mu+\rho_c}) = 0$.*
- *If $\mu + \rho_c$ is a generic quasi-regular value of Φ , then the Spin^c prequantization defined by the data (J, \tilde{L}) induces a Spin^c prequantization on the symplectic quotient $(M_{\mu+\rho_c}, \omega_{\mu+\rho_c})$. We denote $\mathcal{Q}(M_{\mu+\rho_c}) \in \mathbb{Z}$ the index of the corresponding Spin^c Dirac operator.*
- *If $\mu + \rho_c \in \Delta^\circ$, we take ξ generic and quasi-regular sufficiently close to $\mu + \rho_c$. The reduced space $M_\xi := \Phi^{-1}(\xi)/T$ inherits a Spin^c-structure with canonical line bundle $(L_{2\omega}|_{\Phi^{-1}(\xi)} \otimes \mathbb{C}_{-2(\mu+\rho_c)})/T$. The index $\mathcal{Q}(M_\xi)$ of the corresponding Spin^c Dirac operator on M_ξ does not depend of ξ , when ξ is sufficiently close to $\mu + \rho_c$: it is denoted $\mathcal{Q}(M_{\mu+\rho_c})$.*

When $\xi = \mu + \rho_c$ is a generic quasi-regular of Φ , the line bundle $(L_{2\omega}|_{\Phi^{-1}(\xi)} \otimes \mathbb{C}_{-2(\mu+\rho_c)})/T$ is a prequantum line bundle over $(M_{\mu+\rho_c}, 2\omega_{\mu+\rho_c})$: so the second point of this ‘definition’ is in fact a particular case of the third point. But we prefer to keep it since it outlines the main point: Spin^c prequantization is preserved under symplectic reductions.

The existence of Spin^c -structures on symplectic quotient is proved in Subsection 4.2. The hard part is to show that the index $\mathcal{Q}(M_\xi)$ does not depend of ξ , for ξ sufficiently close to $\mu + \rho_c$: it is done in Subsection 4.3.

Note that Definition 2.4 becomes trivial when Δ^o is not included in the interior of the Weyl chamber: $\mathcal{Q}(M_{\mu+\rho_c}) = 0$ for all $\mu \in \Lambda_+^*$. However, in this paper we work under the assumption that the infinitesimal stabilizers for the K -action are *Abelian*. And that imposes $\Delta^o \subset \text{Interior}\{\text{Weyl chamber}\}$ (see Lemma 4.9).

The following ‘*quantization commutes with reduction*’ Theorem holds for the κ -prequantum line bundles.

Theorem 2.5. *Let (M, ω, Φ) be a compact Hamiltonian K -manifold equipped with an almost complex structure J . Let \tilde{L} be a κ -prequantum line bundle over M , and let $RR^K(M, -)$ be the Riemann-Roch character defined by J . If the infinitesimal stabilizers for the action of K on M are Abelian, we have the following equality in $R(K)$*

$$(2.3) \quad RR^K(M, \tilde{L}) = \varepsilon \sum_{\mu \in \Lambda_+^*} \mathcal{Q}(M_{\mu+\rho_c}) \chi_\mu^K,$$

where $\varepsilon = \pm 1$ is the ‘quotient’ of the orientations defined by the almost complex structure, and by the symplectic form.

Theorem 2.5 will be proved in a stronger form in Section 4.

Let us now give an example where the stabilizers for the action of K on M are *not Abelian*, and where (2.3) does not hold. Suppose that the group K is not Abelian, so we can consider a face $\sigma \neq \{0\}$ of the Weyl chamber. Let $\rho_{c,\sigma}$ be half the sum of the positive roots which vanish on σ , and consider the coadjoint orbit $M := K \cdot (\rho_c - \rho_{c,\sigma})$ equipped with its compatible complex structure. Since $\rho_c - \rho_{c,\sigma}$ belongs to σ , the trivial line bundle $M \times \mathbb{C} \rightarrow M$ is κ -prequantum, and the image of the moment map $\Phi : M \rightarrow \mathfrak{k}^*$ does not intersect the interior of the Weyl chamber. So $M_{\mu+\rho_c} = \emptyset$ for every μ , thus the RHS of (2.3) is equal to zero. But the LHS of (2.3) is $RR^K(M, \mathbb{C})$ which is equal to 1, the character of the trivial representation. \square

Theorem 2.5 can be extended in two directions. First one can bypass the condition on the stabilizers by the following trick. Starting from a κ -prequantum line bundle $\tilde{L} \rightarrow M$, one can form the product $M \times (K \cdot \rho_c)$ with the coadjoint orbit through ρ_c . The Kunnetth formula gives

$$RR^K(M \times (K \cdot \rho_c), \tilde{L} \boxtimes \mathbb{C}) = RR^K(M, \tilde{L}) \otimes RR^K(K \cdot \rho_c, \mathbb{C}) = RR^K(M, \tilde{L})$$

since $RR^K(K \cdot \rho_c, \mathbb{C}) = 1$. Now we can apply Theorem 2.5 to compute the multiplicities of $RR^K(M \times (K \cdot \rho_c), \tilde{L} \boxtimes \mathbb{C})$ since $\tilde{L} \boxtimes \mathbb{C}$ is a κ -prequantum line bundle over $M \times (K \cdot \rho_c)$, and the stabilizers for the K -action on $M \times (K \cdot \rho_c)$ are Abelian. Finally we see that the multiplicity of the irreducible representation with highest weight μ in $RR^K(M, \tilde{L})$ is equal to $\varepsilon \mathcal{Q}((M \times (K \cdot \rho_c))_{\mu+\rho_c})$.

On the other hand, we can extend Theorem 2.5 to the Spin^c setting. It will be treated in a forthcoming paper.

3. QUANTIZATION OF NON-COMPACT HAMILTONIAN K -MANIFOLDS

In this section (M, ω, Φ) denotes a Hamiltonian K -manifold, not necessarily compact, but with *proper* moment map Φ . Let J be an almost complex structure on M , and let \tilde{L} be a κ -prequantum line bundle over (M, ω, J) (see Def. 2.1). From Proposition 2.4 the infinite sum

$$(3.4) \quad \sum_{\mu \in \Lambda_+^*} \mathcal{Q}(M_{\mu+\rho_c}) \chi_\mu^K$$

is a well defined element of $\widehat{R}(K) := \text{hom}_{\mathbb{Z}}(R(K), \mathbb{Z})$.

The aim of this section is to realize this sum as the index of a *transversally elliptic* symbol naturally associated to the data (M, Φ, J, \tilde{L}) .

3.1. Transversally elliptic symbols. Here we give the basic definitions of the theory of transversally elliptic symbols (or operators) defined by Atiyah in [1]. For an axiomatic treatment of the index morphism see Berline-Vergne [9, 10] and for a short introduction see [33].

Let M be a *compact* K -manifold. Let $p : \mathbf{T}M \rightarrow M$ be the projection, and let $(-, -)_M$ be a K -invariant Riemannian metric. If E^0, E^1 are K -equivariant vector bundles over M , a K -equivariant morphism $\sigma \in \Gamma(\mathbf{T}M, \text{hom}(p^*E^0, p^*E^1))$ is called a *symbol*. The subset of all $(m, v) \in \mathbf{T}M$ where $\sigma(m, v) : E_m^0 \rightarrow E_m^1$ is not invertible is called the *characteristic set* of σ , and is denoted by $\text{Char}(\sigma)$.

Let $\mathbf{T}_K M$ be the following subset of $\mathbf{T}M$:

$$\mathbf{T}_K M = \{(m, v) \in \mathbf{T}M, (v, X_M(m))_M = 0 \text{ for all } X \in \mathfrak{k}\}.$$

A symbol σ is *elliptic* if σ is invertible outside a compact subset of $\mathbf{T}M$ ($\text{Char}(\sigma)$ is compact), and is *transversally elliptic* if the restriction of σ to $\mathbf{T}_K M$ is invertible outside a compact subset of $\mathbf{T}_K M$ ($\text{Char}(\sigma) \cap \mathbf{T}_K M$ is compact). An elliptic symbol σ defines an element in the equivariant K -theory of $\mathbf{T}M$ with compact support, which is denoted by $\mathbf{K}_K(\mathbf{T}M)$, and the index of σ is a virtual finite dimensional representation of K [4, 5, 6, 7].

A *transversally elliptic* symbol σ defines an element of $\mathbf{K}_K(\mathbf{T}_K M)$, and the index of σ is defined as a trace class virtual representation of K (see [1] for the analytic index and [9, 10] for the cohomological one). Remark that any elliptic symbol of $\mathbf{T}M$ is transversally elliptic, hence we have a restriction map $\mathbf{K}_K(\mathbf{T}M) \rightarrow \mathbf{K}_K(\mathbf{T}_K M)$, and a commutative diagram

$$(3.5) \quad \begin{array}{ccc} \mathbf{K}_K(\mathbf{T}M) & \longrightarrow & \mathbf{K}_K(\mathbf{T}_K M) \\ \text{Index}_M^K \downarrow & & \downarrow \text{Index}_M^K \\ R(K) & \longrightarrow & R^{-\infty}(K) . \end{array}$$

Using the *excision property*, one can easily show that the index map $\text{Index}_{\mathcal{U}}^K : \mathbf{K}_K(\mathbf{T}_K \mathcal{U}) \rightarrow R^{-\infty}(K)$ is still defined when \mathcal{U} is a K -invariant relatively compact open subset of a K -manifold (see [33][section 3.1]).

3.2. Thom symbol deformed by the moment map. To a K -invariant almost complex structure J one associates the Thom symbol $\text{Thom}_K(M, J)$, and the corresponding Riemann-Roch character RR^K when M is compact [33]. Let us recall the definitions.

Consider a K -invariant Riemannian metric q on M such that J is orthogonal relatively to q , and let h be the Hermitian structure on $\mathbf{T}M$ defined by : $h(v, w) = q(v, w) - iq(Jv, w)$ for $v, w \in \mathbf{T}M$. The symbol

$$\text{Thom}_K(M, J) \in \Gamma(M, \text{hom}(p^*(\wedge_{\mathbb{C}}^{\text{even}} \mathbf{T}M), p^*(\wedge_{\mathbb{C}}^{\text{odd}} \mathbf{T}M)))$$

at $(m, v) \in \mathbf{T}M$ is equal to the Clifford map

$$(3.6) \quad \text{Cl}_m(v) : \wedge_{\mathbb{C}}^{\text{even}} \mathbf{T}_m M \longrightarrow \wedge_{\mathbb{C}}^{\text{odd}} \mathbf{T}_m M,$$

where $\text{Cl}_m(v).w = v \wedge w - c_h(v).w$ for $w \in \wedge_{\mathbb{C}}^{\bullet} \mathbf{T}_m M$. Here $c_h(v) : \wedge_{\mathbb{C}}^{\bullet} \mathbf{T}_m M \rightarrow \wedge_{\mathbb{C}}^{\bullet-1} \mathbf{T}_m M$ denotes the contraction map relative to h . Since the map $\text{Cl}_m(v)$ is invertible for all $v \neq 0$, the symbol $\text{Thom}_K(M, J)$ is elliptic when M is compact.

The important point is that for any K -vector bundle E , $\text{Thom}_K(M, J) \otimes p^*E$ corresponds to the *principal symbol* of the twisted Spin^c Dirac operator \mathcal{D}_E [14]. So, when M is a compact manifold, the Riemann-Roch character $RR^K(M, -) : \mathbf{K}_K(M) \rightarrow R(K)$ is defined by the following relation

$$(3.7) \quad RR^K(M, E) = \text{Index}_M^K(\text{Thom}_K(M, J) \otimes p^*E) .$$

Since the class of $\text{Thom}_K(M, J)$ in $\mathbf{K}_K(\mathbf{T}M)$ is independent of the choice of the Riemannian structure, the Riemann-Roch character $RR^K(M, -)$ also does not depend on this choice.

Consider now the case of a *non-compact* Hamiltonian K -manifold (M, ω, Φ) . We choose a K -invariant scalar product on \mathfrak{k}^* , and we consider the function $\|\Phi\|^2 : M \rightarrow \mathbb{R}$. Let \mathcal{H} be the Hamiltonian vector field for $\frac{1}{2}\|\Phi\|^2$, i.e. the contraction of the symplectic form by \mathcal{H} is equal to the 1-form $\frac{1}{2}d\|\Phi\|^2$. In fact the vector field \mathcal{H} only depends on Φ . The scalar product on \mathfrak{k}^* gives an identification $\mathfrak{k}^* \simeq \mathfrak{k}$, hence Φ can be consider as a map from M to \mathfrak{k} . We have then

$$(3.8) \quad \mathcal{H}_m = (\Phi(m))_M|_m, \quad m \in M ,$$

where $(\Phi(m))_M$ is the vector field on M generated by $\Phi(m) \in \mathfrak{k}$.

Definition 3.1. *The Thom symbol deformed by the moment map, which is denoted by $\text{Thom}_K^{\Phi}(M, J)$, is defined by the relation*

$$\text{Thom}_K^{\Phi}(M, J)(m, v) := \text{Thom}_K(M, J)(m, v - \mathcal{H}_m)$$

for any $(m, v) \in \mathbf{T}M$. Likewise, any equivariant map $S : M \rightarrow \mathfrak{k}$ defines a Thom symbol $\text{Thom}_K^S(M, J)$ deformed by the vector field $S_M : m \rightarrow S(m)_M|_m$: $\text{Thom}_K^S(M, J)(m, v) := \text{Thom}_K(M, J)(m, v - S_M(m))$.

Atiyah first proposed to ‘deform’ the symbol of an elliptic operator by the vector field induced by an S^1 -action in order to localize its index on the fixed point submanifold, giving then another proof of the Lefschetz fixed-point theorem [1][Lecture 6]. Afterwards the idea was exploited by Vergne to give a proof of the ‘quantization commutes with reduction’ theorem in the case of an S^1 -action [42]. In [33], we extended this procedure for an action of a compact Lie group. Here, we use this idea to produce a transversally elliptic symbol on a non-compact manifold.

The characteristic set of $\text{Thom}_K^\Phi(M, J)$ corresponds to $\{(m, v) \in \mathbf{T}M, v = \mathcal{H}_m\}$, the graph of the vector field \mathcal{H} . Since \mathcal{H} belongs to the set of tangent vectors to the K -orbits, we have

$$\begin{aligned} \text{Char}\left(\text{Thom}_K^\Phi(M, J)\right) \cap \mathbf{T}_K M &= \{(m, 0) \in \mathbf{T}M, \mathcal{H}_m = 0\} \\ &\cong \{m \in M, d\|\Phi\|_m^2 = 0\}. \end{aligned}$$

Therefore the symbol $\text{Thom}_K^\Phi(M, J)$ is transversally elliptic if and only if the set $\text{Cr}(\|\Phi\|^2)$ of critical points of the function $\|\Phi\|^2$ is compact.

Definition 3.2. Let (M, ω, Φ) be a Hamiltonian K -manifold with $\text{Cr}(\|\Phi\|^2)$ compact. For any invariant almost complex structure J , the symbol $\text{Thom}_K^\Phi(M, J)$ is transversally elliptic. For any K -vector bundle $E \rightarrow M$, the tensor product $\text{Thom}_K^\Phi(M, J) \otimes p^*E$ is transversally elliptic and we denote by

$$RR_\Phi^K(M, E) \in R^{-\infty}(K)$$

its index³. In the same way, an equivariant map $S : M \rightarrow \mathfrak{k}$ defines a transversally elliptic symbol $\text{Thom}_K^S(M, J)$ if and only if $\{m \in M, S_M(m) = 0\}$ is compact. If this holds one defines the localized Riemann-Roch character $RR_S^K(M, E) := \text{Index}_M^K(\text{Thom}_K^S(M) \otimes p^*E)$.

Remark 3.3. If M is compact the symbols $\text{Thom}_K(M, J)$ and $\text{Thom}_K^\Phi(M, J)$ are homotopic as elliptic symbols, thus the maps $RR^K(M, -)$ and $RR_\Phi^K(M, -)$ coincide (see section 4 of [33]).

We end up this subsection with some technical remarks about the symbols $\text{Thom}_K^S(M, J)$ associated to an equivariant map $S : M \rightarrow \mathfrak{k}$, and an almost complex structure.

Let \mathcal{U} be a K -invariant open subspace of M . The restriction $\text{Thom}_K^S(M, J)|_{\mathcal{U}} = \text{Thom}_K^S(\mathcal{U}, J)$ is transversally elliptic if and only if $\{m \in M, S_M(m) = 0\} \cap \mathcal{U}$ is compact. Let $j^{\mathcal{U}, \mathcal{V}} : \mathcal{U} \hookrightarrow \mathcal{V}$ be two K -invariant open subspaces of M , where $j^{\mathcal{U}, \mathcal{V}}$ denotes the inclusion. If $\{m \in M, S_M(m) = 0\} \cap \mathcal{U} = \{m \in M, S_M(m) = 0\} \cap \mathcal{V}$ is compact, the excision property tells us that

$$j_*^{\mathcal{U}, \mathcal{V}}\left(\text{Thom}_K^S(\mathcal{U}, J)\right) = \text{Thom}_K^S(\mathcal{V}, J),$$

where $j_*^{\mathcal{U}, \mathcal{V}} : \mathbf{K}_K(\mathbf{T}_K \mathcal{U}) \rightarrow \mathbf{K}_K(\mathbf{T}_K \mathcal{V})$ is the pushforward map (see [33][Section 3]).

Lemma 3.4. (1) If $\{m \in M, S_M(m) = 0\} \cap \mathcal{U}$ is compact, then the class defined by $\text{Thom}_K^S(\mathcal{U}, J)$ in $\mathbf{K}_K(\mathbf{T}_K \mathcal{U})$ does not depend on the choice of a Riemannian metric.

³Here we take a K -invariant relatively compact open subset \mathcal{U} of M such that $\text{Cr}(\|\Phi\|^2) \subset \mathcal{U}$. Then the restriction of $\text{Thom}_K^\Phi(M, J)$ to \mathcal{U} defines a class $\text{Thom}_K^\Phi(M, J)|_{\mathcal{U}} \in \mathbf{K}_K(\mathbf{T}_K \mathcal{U})$. Since the index map is well defined on \mathcal{U} , one sets $RR_\Phi^K(M, E) := \text{Index}_\mathcal{U}^K(\text{Thom}_K^\Phi(M, J)|_{\mathcal{U}} \otimes p^*E|_{\mathcal{U}})$. A simple application of the excision property shows us that the definition does not depend on the choice of \mathcal{U} . In order to simplify our notation (when the almost complex structure is understood), we write $RR_\Phi^K(M, E) := \text{Index}_M^K(\text{Thom}_K^\Phi(M) \otimes p^*E)$.

- (2) Let $S^0, S^1 : M \rightarrow \mathfrak{k}$ be two equivariant maps. Suppose there exist an open subset $\mathcal{U} \subset M$, and a vector field θ on \mathcal{U} such that $(S_M^0, \theta)_M$ and $(S_M^1, \theta)_M$ are > 0 outside a compact subset \mathcal{K} of \mathcal{U} . Then, the equivariant symbols $\text{Thom}_K^{S^1}(\mathcal{U}, J)$ and $\text{Thom}_K^{S^0}(\mathcal{U}, J)$ are transversally elliptic and define the same class in $\mathbf{K}_K(\mathbf{T}_K \mathcal{U})$.
- (3) Let J^0, J^1 be two almost complex structures on \mathcal{U} , and suppose that $\{m \in M, S_M(m) = 0\} \cap \mathcal{U}$ is compact. The transversally elliptic symbols $\text{Thom}_K^S(\mathcal{U}, J^0)$ and $\text{Thom}_K^S(\mathcal{U}, J^1)$ define the same class if there exists a homotopy $J^t, t \in [0, 1]$ of K -equivariant almost complex structures between J^0 and J^1 .

Proof. Two J -invariant Riemannian metrics q_0, q_1 are connected by $q_t := (1-t)q_0 + tq_1$. Hence the transversally elliptic symbols $\text{Thom}_K^S(\mathcal{U}, J, q_0)$ and $\text{Thom}_K^S(\mathcal{U}, J, q_1)$ are tied by the homotopy $t \mapsto \text{Thom}_K^S(\mathcal{U}, J, q_t)$. The point 1. is then proved. The proof of 2. is similar to our deformation process in [31]. Here we consider the maps $S^t := tS^1 + (1-t)S^0, t \in [0, 1]$, and the corresponding symbols $\text{Thom}_K^{S^t}(\mathcal{U}, J)$. The vector field θ , ensures that $\text{Char}(\text{Thom}_K^{S^t}(\mathcal{U}, J)) \cap \mathbf{T}_K \mathcal{U} \subset \mathcal{K}$ is compact. Hence $t \rightarrow \text{Thom}_K^{S^t}(\mathcal{U}, J)$ defines a homotopy of transversally elliptic symbols. The proof of 3. is identical to the proof of Lemma 2.2 in [33]. \square

Corollary 3.5. *When $\{m \in M, S_M(m) = 0\}$ is compact, the generalized Riemann-Roch character $RR_S^K(M, -)$ does not depend on the choice of a Riemannian metric. $RR_S^K(M, -)$ does not change either if the almost complex structure is deformed smoothly and equivariantly in a neighborhood of $\{m \in M, S_M(m) = 0\}$.*

In Subsections 3.3 and 3.4, we set up the technical preliminaries that are needed to compute the K -multiplicity of $RR_\Phi^K(M, \tilde{L})$.

In Section 4, we compute the K -multiplicity of $RR_\Phi^K(M, \tilde{L})$, when the moment map is *is proper*, in terms of the symplectic quotients $M_{\mu+\rho_c}, \mu \in \Lambda_+^*$.

3.3. Counting the K -multiplicities. Let E be a K -vector bundle over a Hamiltonian manifold (M, ω, Φ) and suppose that $\text{Cr}(\|\Phi\|^2)$ is compact. One wants to compute the K -multiplicities of $RR_\Phi^K(M, E) \in R^{-\infty}(K)$, i.e. the integers $m_\mu(E) \in \mathbb{Z}, \mu \in \Lambda_+^*$ such that

$$(3.9) \quad RR_\Phi^K(M, E) = \sum_{\mu \in \Lambda_+^*} m_\mu(E) \chi_\mu^K.$$

For this purpose one use the classical ‘shifting trick’. By definition, one has $m_\mu(E) = [RR_\Phi^K(M, E) \otimes V_\mu^*]^K$, where V_μ is the irreducible K -representation with highest weight μ , and V_μ^* is its dual. We know from (2.2) that the K -trace of V_μ is $\chi_\mu^K = RR^\kappa(\mathcal{O}^{\tilde{\mu}}, \tilde{\mathbb{C}}_{[\mu]}),$ where

$$(3.10) \quad \tilde{\mu} = \mu + \rho_c.$$

Hence the K -trace of the dual V_μ^* is equal to $RR^\kappa(\overline{\mathcal{O}^{\tilde{\mu}}}, \tilde{\mathbb{C}}_{[-\mu]}),$ where $\overline{\mathcal{O}^{\tilde{\mu}}}$ is the coadjoint orbit $\mathcal{O}^{\tilde{\mu}}$ with opposite symplectic structure and opposite complex structure. Let $\text{Thom}_K(\overline{\mathcal{O}^{\tilde{\mu}}})$ be the equivariant Thom symbol on $\overline{\mathcal{O}^{\tilde{\mu}}}$. Then the trace of V_μ^* is equal to $\text{Index}_{\mathcal{O}^{\tilde{\mu}}}^K(\text{Thom}_K(\overline{\mathcal{O}^{\tilde{\mu}}}) \otimes \tilde{\mathbb{C}}_{[-\mu]}),$ and finally the multiplicative property of

the index [1][Theorem 3.5] gives

$$m_\mu(E) = \left[\text{Index}_{M \times \mathcal{O}^{\tilde{\mu}}}^K \left((\text{Thom}_K^\Phi(M) \otimes p^*E) \odot (\text{Thom}_K(\overline{\mathcal{O}^{\tilde{\mu}}}) \otimes \tilde{\mathbb{C}}_{[-\mu]}) \right) \right]^K.$$

See [1, 33], for the definition of the exterior product $\odot : \mathbf{K}_K(\mathbf{T}_K M) \times \mathbf{K}_K(\mathbf{T}\mathcal{O}^{\tilde{\mu}}) \rightarrow \mathbf{K}_K(\mathbf{T}_K(M \times \mathcal{O}^{\tilde{\mu}}))$.

The moment map relative to the Hamiltonian K -action on $M \times \overline{\mathcal{O}^{\tilde{\mu}}}$ is

$$(3.11) \quad \begin{aligned} \Phi_{\tilde{\mu}} : M \times \overline{\mathcal{O}^{\tilde{\mu}}} &\longrightarrow \mathfrak{k}^* \\ (m, \xi) &\longmapsto \Phi(m) - \xi \end{aligned}$$

For any $t \in \mathbb{R}$, we consider the map $\Phi_{t\tilde{\mu}} : M \times \overline{\mathcal{O}^{\tilde{\mu}}} \rightarrow \mathfrak{k}^*$, $\Phi_{t\tilde{\mu}}(m, \xi) := \Phi(m) - t\xi$.

Assumption 3.6. *There exists a compact subset $\mathcal{K} \subset M$, such that, for every $t \in [0, 1]$, the critical set of the function $\|\Phi_{t\tilde{\mu}}\|^2 : M \times \mathcal{O}^{\tilde{\mu}} \rightarrow \mathbb{R}$ is contained in $\mathcal{K} \times \mathcal{O}^{\tilde{\mu}}$.*

If M satisfies Assumption 3.6 at $\tilde{\mu}$, one has a generalized Riemann-Roch character $RR_{\Phi_{\tilde{\mu}}}^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, -)$ since $\text{Cr}(\|\Phi_{\tilde{\mu}}\|^2)$ is compact.

Proposition 3.7. *Let $m_\mu(E)$ be the multiplicity of $RR_{\Phi}^K(M, E)$ relatively to the highest weight $\mu \in \Lambda_+^*$. If M satisfies Assumption 3.6 at $\tilde{\mu}$, then*

$$m_\mu(E) = \left[RR_{\Phi_{\tilde{\mu}}}^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, E \boxtimes \tilde{\mathbb{C}}_{[-\mu]}) \right]^K.$$

Proof. One has to show that the transversally elliptic symbols $\text{Thom}_K^\Phi(M) \odot \text{Thom}_K(\overline{\mathcal{O}^{\tilde{\mu}}})$ and $\text{Thom}_K^{\Phi_{\tilde{\mu}}}(M \times \overline{\mathcal{O}^{\tilde{\mu}}})$ define the same class in $\mathbf{K}_K(\mathbf{T}_K(M \times \mathcal{O}^{\tilde{\mu}}))$ when M satisfies Assumption 3.6 at $\tilde{\mu}$. Let σ_1, σ_2 be respectively the Thom symbols $\text{Thom}_K^\Phi(M)$ and $\text{Thom}_K(\overline{\mathcal{O}^{\tilde{\mu}}})$. The symbol $\sigma_I = \text{Thom}_K^\Phi(M) \odot \text{Thom}_K(\overline{\mathcal{O}^{\tilde{\mu}}})$ is defined by

$$\sigma_I(m, \xi, v, w) = \sigma_1(m, v - \mathcal{H}_m) \odot \sigma_2(\xi, w),$$

where $(m, v) \in \mathbf{T}M$, $(\xi, w) \in \mathbf{T}\mathcal{O}^{\tilde{\mu}}$, and \mathcal{H} is defined in (3.8). Let \mathcal{H}^t be the vector field on $M \times \mathcal{O}^{\tilde{\mu}}$ generated by the map $\Phi_{t\tilde{\mu}} : M \times \mathcal{O}^{\tilde{\mu}} \rightarrow \mathfrak{k}$. For $(m, \xi) \in M \times \mathcal{O}^{\tilde{\mu}}$, we have $\mathcal{H}_{(m, \xi)}^t = (\mathcal{H}_{(m, \xi)}^{a, t}, \mathcal{H}_{(m, \xi)}^{b, t})$ where $\mathcal{H}_{(m, \xi)}^{a, t} \in \mathbf{T}_m M$ and $\mathcal{H}_{(m, \xi)}^{b, t} \in \mathbf{T}_\xi \mathcal{O}^{\tilde{\mu}}$. The symbol $\sigma_{II} = \text{Thom}_K^{\Phi_{\tilde{\mu}}}(M \times \mathcal{O}^{\tilde{\mu}})$ is defined by

$$\sigma_{II}(m, \xi, v, w) = \sigma_1(m, v - \mathcal{H}_{m, \xi}^{a, 1}) \odot \sigma_2(\xi, w - \mathcal{H}_{(m, \xi)}^{b, 1}).$$

We connect σ_I and σ_{II} through two homotopies. First we consider the symbol A on $[0, 1] \times \mathbf{T}(M \times \mathcal{O}^{\tilde{\mu}})$ defined by

$$A(t; m, \xi, v, w) = \sigma_1(m, v - \mathcal{H}_{m, \xi}^{a, t}) \odot \sigma_2(\xi, w - \mathcal{H}_{(m, \xi)}^{b, t}),$$

for $t \in [0, 1]$, and $(m, \xi, v, w) \in \mathbf{T}(M \times \mathcal{O}^{\tilde{\mu}})$. We have $\text{Char}(A) = \{(t; m, \xi, v, w) \mid v = \mathcal{H}_{m, \xi}^{a, t}, \text{ and } w = \mathcal{H}_{(m, \xi)}^{b, t}\}$ and

$$\begin{aligned} \text{Char}(A) \cap [0, 1] \times \mathbf{T}_K(M \times \mathcal{O}^{\tilde{\mu}}) &= \{(t; m, \xi, 0, 0) \mid (m, \xi) \in \text{Cr}(\|\Phi_{t\tilde{\mu}}\|^2)\} \\ &\subset [0, 1] \times \mathcal{K} \times \mathcal{O}^{\tilde{\mu}}, \end{aligned}$$

where $\mathcal{K} \subset M$ is the compact subset of Assumption 3.6. Thus A defines a homotopy of transversally elliptic symbols. The restriction of A to $t = 1$ is equal to σ_{II} . The restriction of A to $t = 0$ defines the following transversally elliptic symbol

$$\sigma_{III}(m, \xi, v, w) = \sigma_1(m, v - \mathcal{H}_m) \odot \sigma_2(\xi, w - \mathcal{H}_{(m, \xi)}^{b,0})$$

since $\mathcal{H}_{m, \xi}^{a,0} = \mathcal{H}_m$ for every $(m, \xi) \in M \times \mathcal{O}^{\tilde{\mu}}$. Next, we consider the symbol B on $[0, 1] \times \mathbf{T}(M \times \mathcal{O}^{\tilde{\mu}})$ defined by

$$B(t; m, \xi, v, w) = \sigma_1(m, v - \mathcal{H}_m) \odot \sigma_2(\xi, w - t \mathcal{H}_{(m, \xi)}^{b,0}).$$

We have $\text{Char}(B) = \{(t; m, \xi, v, w) \mid v = \mathcal{H}_m, \text{ and } w = t \mathcal{H}_{(m, \xi)}^{b,0}\}$ and

$$\begin{aligned} \text{Char}(B) \cap [0, 1] \times \mathbf{T}_K(M \times \mathcal{O}^{\tilde{\mu}}) \subset \\ \left\{ (t; m, \xi, v = \mathcal{H}_m, w = t \mathcal{H}_{(m, \xi)}^{b,0}) \quad , \quad \| \mathcal{H}_m \|^2 + t \| \mathcal{H}_{(m, \xi)}^{b,0} \|^2 = 0 \right\}. \end{aligned}$$

In particular $\text{Char}(B) \cap [0, 1] \times \mathbf{T}_K(M \times \mathcal{O}^{\tilde{\mu}})$ is contained in $\{(t; m, \xi, 0, w = t \mathcal{H}_{(m, \xi)}^{b,0}), m \in \text{Cr}(\| \Phi \|^2)\}$ which is compact since $\text{Cr}(\| \Phi \|^2)$ is compact. So, B defines a homotopy of transversally elliptic symbols between $\sigma_I = B|_{t=0}$ and $\sigma_{III} = B|_{t=1}$. We have finally proved that $\sigma_I, \sigma_{II}, \sigma_{III}$ define the same class in $\mathbf{K}_K(\mathbf{T}_K(M \times \mathcal{O}^{\tilde{\mu}}))$. \square

When $E = \tilde{L}$ is a κ -prequantum line bundle over M , the line bundle $\tilde{L} \boxtimes \tilde{\mathbb{C}}_{[-\mu]}$ is a κ -prequantum line bundle over $M \times \overline{\mathcal{O}^{\tilde{\mu}}}$. Therefore Proposition 3.7 shows that under Assumption 3.6 the K -multiplicities of $RR_{\Phi}^K(M, \tilde{L})$ have the form

$$(3.12) \quad \left[RR_{\Phi}^K(\mathcal{X}, \tilde{L}_{\mathcal{X}}) \right]^K,$$

where $(\mathcal{X}, \omega_{\mathcal{X}}, \Phi)$ is a Hamiltonian K -manifold with $\text{Cr}(\| \Phi \|^2)$ compact, and $\tilde{L}_{\mathcal{X}}$ is a κ -prequantum line bundle over \mathcal{X} relative to a K -invariant almost complex structure. In order to compute the quantity (3.12), we exploit in the next subsection the localization techniques developed in [33].

3.4. Localization of the map RR_{Φ}^K . For a detailed account on the procedure of localization that we use here, see Sections 4 and 6 of [33]. In this section $(\mathcal{X}, \omega_{\mathcal{X}}, \Phi)$ is a Hamiltonian K -manifold which is equipped with a K -invariant almost complex structure, and a κ -prequantum line bundle \tilde{L} . We suppose furthermore that $\text{Cr}(\| \Phi \|^2)$ is compact. We give here a condition under which $[RR_{\Phi}^K(\mathcal{X}, \tilde{L})]^K$ depends only on the data in the neighborhood of $\Phi^{-1}(0)$.

For any $\beta \in \mathfrak{k}$, let \mathcal{X}^{β} be the symplectic submanifold of points of \mathcal{X} fixed by the torus \mathbb{T}_{β} generated by β . Following Kirwan [22], the critical set $\text{Cr}(\| \Phi \|^2)$ decomposes as

$$(3.13) \quad \text{Cr}(\| \Phi \|^2) = \bigcup_{\beta \in \mathcal{B}} C_{\beta}^K, \quad \text{with} \quad C_{\beta}^K = K \cdot (\mathcal{X}^{\beta} \cap \Phi^{-1}(\beta)),$$

where \mathcal{B} is the subset of \mathfrak{k}_+^* defined by $\mathcal{B} := \{\beta \in \mathfrak{k}_+^*, \mathcal{X}^{\beta} \cap \Phi^{-1}(\beta) \neq \emptyset\}$. Since $\text{Cr}(\| \Phi \|^2)$ is supposed to be compact, \mathcal{B} is finite.

For each $\beta \in \mathcal{B}$, let $\mathcal{U}^{\beta} \hookrightarrow \mathcal{X}$ be a K -invariant *relatively compact* open neighborhood of C_{β}^K such that $\overline{\mathcal{U}^{\beta}} \cap \text{Cr}(\| \Phi \|^2) = C_{\beta}^K$. The restriction of the transversally elliptic symbol $\text{Thom}_{\kappa}^{\Phi}(\mathcal{X})$ to the subset \mathcal{U}^{β} defines $\text{Thom}_{\kappa}^{\Phi}(\mathcal{U}^{\beta}) \in \mathbf{K}_K(\mathbf{T}_K \mathcal{U}^{\beta})$.

Definition 3.8. For every $\beta \in \mathcal{B}$, we denote by $RR_\beta^K(\mathcal{X}, -)$ the Riemann-Roch character localized near C_β^K , which is defined by

$$RR_\beta^K(\mathcal{X}, E) = \text{Index}_{\mathcal{U}^\beta}^K \left(\text{Thom}_\kappa^\Phi(\mathcal{U}^\beta) \otimes p^* E|_{\mathcal{U}^\beta} \right),$$

for every K -vector bundle $E \rightarrow \mathcal{X}$.

The excision property tells us that

$$(3.14) \quad RR_\Phi^K(\mathcal{X}, E) = \sum_{\beta \in \mathcal{B}} RR_\beta^K(\mathcal{X}, E)$$

for every K -vector bundle $E \rightarrow \mathcal{X}$ (see [33][Section 4]). In particular, $[RR_\Phi^K(\mathcal{X}, \tilde{L})]^K = \sum_\beta [RR_\beta^K(\mathcal{X}, \tilde{L})]^K$, and our main point here is to find suitable conditions under which $[RR_\beta^K(\mathcal{X}, \tilde{L})]^K = 0$ for $\beta \neq 0$.

Let β be a non-zero element in \mathfrak{k} . For every connected component \mathcal{Z} of \mathcal{X}^β , let $\mathcal{N}_\mathcal{Z}$ be the normal bundle of \mathcal{Z} in \mathcal{X} . Let $\alpha_1^\mathcal{Z}, \dots, \alpha_l^\mathcal{Z}$ be the real infinitesimal weights for the action of \mathbb{T}_β on the fibers of $\mathcal{N}_\mathcal{Z} \otimes \mathbb{C}$. The infinitesimal action of β on $\mathcal{N}_\mathcal{Z} \otimes \mathbb{C}$ is a linear map with trace equal to $\sqrt{-1} \sum_i \langle \alpha_i^\mathcal{Z}, \beta \rangle$.

Definition 3.9. Let us denote by $\text{Tr}_\beta|\mathcal{N}_\mathcal{Z}|$ the following positive number

$$\text{Tr}_\beta|\mathcal{N}_\mathcal{Z}| := \sum_{i=1}^l |\langle \alpha_i^\mathcal{Z}, \beta \rangle|,$$

where $\alpha_1^\mathcal{Z}, \dots, \alpha_l^\mathcal{Z}$ are the real infinitesimal weights for the action of \mathbb{T}_β on the fibers of $\mathcal{N}_\mathcal{Z} \otimes \mathbb{C}$. For any \mathbb{T}_β -equivariant real vector bundle $\mathcal{V} \rightarrow \mathcal{Z}$ (resp. real \mathbb{T}_β -equivariant real vector space E), we define in the same way $\text{Tr}_\beta|\mathcal{V}| \geq 0$ (resp. $\text{Tr}_\beta|E| \geq 0$).

Remark 3.10. If $\mathcal{V} = \mathcal{V}^1 \oplus \mathcal{V}^2$, we have $\text{Tr}_\beta|\mathcal{V}| = \text{Tr}_\beta|\mathcal{V}^1| + \text{Tr}_\beta|\mathcal{V}^2|$, and if \mathcal{V}' is an equivariant real subbundle of \mathcal{V} , we get $\text{Tr}_\beta|\mathcal{V}| \geq \text{Tr}_\beta|\mathcal{V}'|$. In particular one see that $\text{Tr}_\beta|\mathcal{N}_\mathcal{Z}| = \text{Tr}_\beta|\mathbf{T}\mathcal{X}|_\mathcal{Z}|$, and then, if $E_m \subset \mathbf{T}_m\mathcal{X}$ is a \mathbb{T}_β -invariant real vector subspace for some $m \in \mathcal{Z}$, we have $\text{Tr}_\beta|\mathcal{N}_\mathcal{Z}| \geq \text{Tr}_\beta|E_m|$.

The following Proposition and Corollary give us an essential condition under which the number $[RR_\Phi^K(\mathcal{X}, \tilde{L})]^K$ only depends on data localized in a neighborhood of $\Phi^{-1}(0)$.

Proposition 3.11. Let \tilde{L} be a κ -prequantum line bundle over \mathcal{X} . The multiplicity of the trivial representation in $RR_\beta^K(\mathcal{X}, \tilde{L})$ is equal to zero if

$$(3.15) \quad \|\beta\|^2 + \frac{1}{2} \text{Tr}_\beta|\mathcal{N}_\mathcal{Z}| - 2(\rho_c, \beta) > 0$$

for every connected component \mathcal{Z} of \mathcal{X}^β which intersects $\Phi^{-1}(\beta)$. Condition (3.15) always holds if $\beta \in \mathfrak{k} - \{0\}$ is K -invariant or if $\|\beta\| > \|\rho_c\|$.

Since every $\beta \in \mathcal{B}$ belongs to the Weyl chamber, we have $2(\rho_c, \beta) = \text{Tr}_\beta|\mathfrak{k}/\mathfrak{t}|$, and then (3.15) can be rewritten as $\|\beta\|^2 + \frac{1}{2} \text{Tr}_\beta|\mathcal{N}_\mathcal{Z}| - \text{Tr}_\beta|\mathfrak{k}/\mathfrak{t}| > 0$. From (3.14), we get

Corollary 3.12. *If condition (3.15) holds for all non-zero $\beta \in \mathcal{B}$, we have*

$$\left[RR_{\Phi}^K(\mathcal{X}, \tilde{L}) \right]^K = \left[RR_0^K(\mathcal{X}, \tilde{L}) \right]^K$$

where $RR_0^K(\mathcal{X}, -)$ is the Riemann-Roch character localized near $\Phi^{-1}(0)$ (see Definition 3.8). In particular, $[RR_{\Phi}^K(\mathcal{X}, \tilde{L})]^K = 0$ if (3.15) holds for all non-zero $\beta \in \mathcal{B}$, and $0 \notin \text{Image}(\Phi)$.

3.5. Proof of Proposition 3.11. When $\beta \in \mathfrak{k}$ is K -invariant, the scalar product (ρ_c, β) vanishes and then (3.15) trivially holds. Let us show that (3.15) holds when $\|\beta\| > \|\rho_c\|$. Let \mathcal{Z} a connected component of \mathcal{X}^β which intersects $\Phi^{-1}(\beta)$. Let $m \in \Phi^{-1}(\beta) \cap \mathcal{Z}$, and let $E_m \subset \mathbf{T}_m \mathcal{X}$ be the subspace spanned by $X_{\mathcal{X}}(m)$, $X \in \mathfrak{k}$. We have $E_m \simeq \mathfrak{k}/\mathfrak{k}_m$, where $\mathfrak{k}_m := \{X \in \mathfrak{k}, X_{\mathcal{X}}(m) = 0\}$. Since $\Phi(m) = \beta$, and Φ is equivariant $\mathfrak{k}_m \subset \mathfrak{k}_\beta := \{X \in \mathfrak{k}, [X, \beta] = 0\}$, so $\mathbf{T}_m \mathcal{X}$ contains a \mathbb{T}_β -equivariant subspace isomorphic to $\mathfrak{k}/\mathfrak{k}_\beta$. So we have $\text{Tr}_\beta |\mathcal{N}_{\mathcal{Z}}| \geq \text{Tr}_\beta |\mathfrak{k}/\mathfrak{k}_\beta| = 2(\rho_c, \beta)$, and then

$$\begin{aligned} \|\beta\|^2 + \frac{1}{2} \text{Tr}_\beta |\mathcal{N}_{\mathcal{Z}}| - 2(\rho_c, \beta) &\geq \|\beta\|^2 - (\rho_c, \beta) \\ &> 0 \end{aligned}$$

since $\|\beta\| > \|\rho_c\|$. \square

We prove now that condition (3.15) forces $[RR_{\beta}^K(\mathcal{X}, \tilde{L})]^K$ to be equal to 0. Let $m_{\beta, \mu}(E) \in \mathbb{Z}$ be the K -multiplicities of the localized Riemann-Roch character $RR_{\beta}^K(\mathcal{X}, E)$ introduced in Definition 3.8 : $RR_{\beta}^K(\mathcal{X}, E) = \sum_{\mu \in \Lambda_+^*} m_{\beta, \mu}(E) \chi_{\mu}^K$. We show now that $m_{\beta, 0}(\tilde{L}) = 0$, by using the formulas of localization that we proved in [33] for the maps $RR_{\beta}^K(\mathcal{X}, -)$.

First case : $\beta \in \mathcal{B}$ is a non-zero K -invariant element of \mathfrak{k}^ .*

We show here the following relation for the multiplicities $m_{\beta, \mu}(\tilde{L})$:

$$(3.16) \quad m_{\beta, \mu}(\tilde{L}) \neq 0 \implies (\mu, \beta) \geq \|\beta\|^2 + \frac{1}{2} \text{Tr}_\beta |\mathcal{N}_{\mathcal{Z}}| \quad \text{for some } \mathcal{Z} \subset \mathcal{X}^\beta,$$

in particular $m_{\beta, 0}(\tilde{L}) = 0$.

Since \mathbb{T}_β belongs to the center of K , \mathcal{X}^β is a symplectic K -invariant submanifold of \mathcal{X} . Let \mathcal{N} be the normal bundle of \mathcal{X}^β in \mathcal{X} . The K -invariant almost complex structure of \mathcal{X} induces a K -invariant almost complex structure on \mathcal{X}^β , and a complex structure on the fibers of $\mathcal{N} \rightarrow \mathcal{X}^\beta$. Then we have a Riemann-Roch character $RR_{\beta}^K(\mathcal{X}^\beta, -)$ localized along $\mathcal{X}^\beta \cap \Phi^{-1}(\beta)$ with the decomposition $RR_{\beta}^K(\mathcal{X}^\beta, F) = \sum_{\mathcal{Z}} RR_{\beta}^K(\mathcal{Z}, F|_{\mathcal{Z}})$, where the sum is taken over the connected components $\mathcal{Z} \subset \mathcal{X}^\beta$ which intersect $\Phi^{-1}(\beta)$. The torus \mathbb{T}_β acts linearly on the fibers of the complex vector bundle \mathcal{N} , thus we can associate the polarized complex K -vector bundle $\mathcal{N}^{+, \beta}$ and $(\mathcal{N} \otimes \mathbb{C})^{+, \beta}$ (see Definition 5.5 in [33]): for any real \mathbb{T}_β -weight α on $\mathcal{N}^{+, \beta}$, or on $(\mathcal{N} \otimes \mathbb{C})^{+, \beta}$, we have

$$(3.17) \quad (\alpha, \beta) > 0.$$

We proved the following localization formula in Section 6.2 of [33] which holds in $\hat{R}(K)$ for any K -vector bundle E over \mathcal{X} :

$$(3.18) \quad RR_{\beta}^K(\mathcal{X}, E) = (-1)^{r_{\mathcal{N}}} \sum_{k \in \mathbb{N}} RR_{\beta}^K(\mathcal{X}^\beta, E|_{\mathcal{X}^\beta} \otimes \det \mathcal{N}^{+, \beta} \otimes S^k((\mathcal{N} \otimes \mathbb{C})^{+, \beta})) .$$

Here $r_{\mathcal{N}}$ is the locally constant function on \mathcal{X}^β equal to the complex rank of $\mathcal{N}^{+, \beta}$, and $S^k(-)$ is the k -th symmetric product over \mathbb{C} .

Let $i : \mathfrak{t}_\beta \hookrightarrow \mathfrak{t}$ be the inclusion of the Lie algebra of \mathbb{T}_β , and let $i^* : \mathfrak{t}^* \rightarrow \mathfrak{t}_\beta^*$ be the canonical dual map. Let us recall the basic relationship between the \mathbb{T}_β -weight on the fibers of a K -vector bundle $F \rightarrow \mathcal{X}^\beta$ and the K -multiplicities of $RR_\beta^K(\mathcal{X}^\beta, F) \in \widehat{R}(K)$: if the irreducible representation V_μ occurs in $RR_\beta^K(\mathcal{X}^\beta, F)$, then $i^*(\mu)$ is a \mathbb{T}_β -weight on the fibers of F (see Appendix B in [33]).

If one now uses (3.18), one sees that $m_{\mu, \beta}(\tilde{L}) \neq 0$ only if $i^*(\mu)$ is a \mathbb{T}_β -weight on the fibers of some $\tilde{L}|_{\mathcal{Z}} \otimes \det \mathcal{N}_{\mathcal{Z}}^{+, \beta} \otimes S^k((\mathcal{N}_{\mathcal{Z}} \otimes \mathbb{C})^{+, \beta})$. Since $(i^*(\mu), \beta) = (\mu, \beta)$, (3.16) will be proved if one shows that each \mathbb{T}_β -weight $\gamma_{\mathcal{Z}}$ on $\tilde{L}|_{\mathcal{Z}} \otimes \det \mathcal{N}_{\mathcal{Z}}^{+, \beta} \otimes S^k((\mathcal{N}_{\mathcal{Z}} \otimes \mathbb{C})^{+, \beta})$ satisfies

$$(3.19) \quad (\gamma_{\mathcal{Z}}, \beta) \geq \|\beta\|^2 + \frac{1}{2} \text{Tr}_\beta |\mathcal{N}_{\mathcal{Z}}|.$$

Let $\alpha_{\mathcal{Z}}$ be the \mathbb{T}_β -weight on the fiber of the line bundle $\tilde{L}|_{\mathcal{Z}} \otimes \det \mathcal{N}_{\mathcal{Z}}^{+, \beta}$. Since any \mathbb{T}_β -weight on $S^k((\mathcal{N} \otimes \mathbb{C})^{+, \beta})$ satisfies (3.17), (3.19) holds if

$$(3.20) \quad (\alpha_{\mathcal{Z}}, \beta) \geq \|\beta\|^2 + \frac{1}{2} \text{Tr}_\beta |\mathcal{N}_{\mathcal{Z}}|$$

for every $\mathcal{Z} \subset \mathcal{X}^\beta$ which intersects $\Phi^{-1}(\beta)$. Let $L_{2\omega}$ be the prequantum line bundle on $(M, 2\omega, 2\Phi)$ such that $\tilde{L} = L_{2\omega} \otimes \kappa$ (where κ is by definition equal to $\det(\mathbf{T}_{\mathbb{C}}^* \mathcal{X}) \cong \det(\mathbf{T} \mathcal{X})^{-1}$). We have

$$(\tilde{L}|_{\mathcal{Z}} \otimes \det(\mathcal{N}^{+, \beta}))^2 = L_{2\omega}|_{\mathcal{Z}} \otimes \det(\mathbf{T} \mathcal{X})^{-1}|_{\mathcal{Z}} \otimes \det(\mathcal{N}^{+, \beta})^2.$$

So $2\alpha_{\mathcal{Z}} = \alpha_1 + \alpha_2$ where α_1, α_2 are respectively \mathbb{T}_β -weights on $L_{2\omega}|_{\mathcal{Z}}$ and $\det(\mathbf{T} \mathcal{X})^{-1}|_{\mathcal{Z}} \otimes \det(\mathcal{N}^{+, \beta})^2$. The Kostant formula (2.1) on $L_{2\omega}|_{\mathcal{Z}}$ gives $(\alpha_1, X) = 2(\beta, X)$ for every $X \in \mathfrak{t}_\beta$, in particular

$$(3.21) \quad (\alpha_1, \beta) = 2 \|\beta\|^2.$$

On \mathcal{Z} , the complex vector bundle $\mathbf{T} \mathcal{X}$ has the following decomposition, $\mathbf{T} \mathcal{X}|_{\mathcal{Z}} = \mathbf{T} \mathcal{Z} \oplus \mathcal{N}^{-, \beta} \oplus \mathcal{N}^{+, \beta}$, where $\mathcal{N}^{-, \beta}$ is the orthogonal complement of $\mathcal{N}^{+, \beta}$ in \mathcal{N} : every \mathbb{T}_β -weight δ on $\mathcal{N}^{-, \beta}$ verifies $(\delta, \beta) < 0$. So we get the decomposition $\det(\mathbf{T} \mathcal{X})^{-1}|_{\mathcal{Z}} \otimes \det(\mathcal{N}^{+, \beta})^2 = \det(\mathbf{T} \mathcal{Z}) \otimes \det(\mathcal{N}^{-, \beta})^{-1} \otimes \det(\mathcal{N}^{+, \beta})$, which gives

$$(3.22) \quad (\alpha_2, \beta) = \text{Tr}_\beta |\mathcal{N}_{\mathcal{Z}}|$$

since \mathbb{T}_β acts trivially on $\mathbf{T} \mathcal{Z}$. Finally (3.20) follows trivially from (3.21) and (3.22).

Second case : $\beta \in \mathcal{B}$ such that $K_\beta \neq K$.

Consider the induced Hamiltonian action of K_β on \mathcal{X} , with moment map $\Phi_{K_\beta} : \mathcal{X} \rightarrow \mathfrak{k}_\beta^*$. Let \mathcal{B}' be the indexing set for the critical point of $\|\Phi_{K_\beta}\|^2$ (see (3.13)). Following Definition 3.8, for each $\beta' \in \mathcal{B}'$ we consider the K_β -Riemann-Roch character $RR_{\beta'}^{K_\beta}(\mathcal{X}, -)$ localised along $C_{\beta'}^{K_\beta} = K_\beta \cdot (\mathcal{X}^{\beta'} \cap \Phi_{K_\beta}^{-1}(\beta'))$. Here β is a K_β -invariant element of \mathcal{B}' with $C_{\beta}^{K_\beta} = \mathcal{X}^\beta \cap \Phi^{-1}(\beta)$.

Let $\text{Hol}_T^K : R^{-\infty}(T) \rightarrow R^{-\infty}(K)$, $\text{Hol}_T^{K_\beta} : R^{-\infty}(T) \rightarrow R^{-\infty}(K_\beta)$, and $\text{Hol}_{K_\beta}^K : R^{-\infty}(K_\beta) \rightarrow R^{-\infty}(K)$ be the holomorphic induction maps (see Appendix B in [33]). Recall that $\text{Hol}_T^K = \text{Hol}_{K_\beta}^K \circ \text{Hol}_T^{K_\beta}$. The choice of a Weyl chamber determines a complex structure on the real vector space $\mathfrak{k}/\mathfrak{k}_\beta$. We denote by $\overline{\mathfrak{k}/\mathfrak{k}_\beta}$ the vector space endowed with the opposite complex structure.

The induction formula that we proved in [33][Section 6] states that

$$(3.23) \quad RR_{\beta}^K(\mathcal{X}, E) = \text{Hol}_{K_{\beta}}^K \left(RR_{\beta}^{K_{\beta}}(\mathcal{X}, E) \wedge_{\mathbb{C}}^{\bullet} \overline{\mathfrak{k}/\mathfrak{k}_{\beta}} \right)$$

for every equivariant vector bundle E . Let us first write the decomposition $RR_{\beta}^{K_{\beta}}(\mathcal{X}, \tilde{L}) = \sum_{\mu \in \Lambda_{\beta}^+} m_{\beta, \mu}(\tilde{L}) \chi_{\mu}^{K_{\beta}}$ into irreducible characters of K_{β} . Since β is K_{β} -invariant we can use the result of the First case. In particular (3.16) tells us that

$$(3.24) \quad m_{\beta, \mu}(\tilde{L}) \neq 0 \implies (\mu, \beta) \geq \|\beta\|^2 + \frac{1}{2} \text{Tr}_{\beta} |\mathcal{N}_{\mathcal{Z}}|$$

for some connected component $\mathcal{Z} \subset \mathcal{X}^{\beta}$ which intersects $\Phi^{-1}(\beta)$.

Each irreducible character $\chi_{\mu}^{K_{\beta}}$ is equal to $\text{Hol}_T^{K_{\beta}}(t^{\mu})$, so from (3.23) we get $RR_{\beta}^K(M, \tilde{L}) = \text{Hol}_T^K \left((\sum_{\mu} m_{\beta, \mu}(\tilde{L}) t^{\mu}) \prod_{\alpha \in \mathfrak{R}^+(\mathfrak{k}/\mathfrak{k}_{\beta})} (1 - t^{-\alpha}) \right)$ where $\mathfrak{R}^+(\mathfrak{k}/\mathfrak{k}_{\beta})$ is the set of positive T -weights on $\mathfrak{k}/\mathfrak{k}_{\beta}$: so $\langle \alpha, \beta \rangle > 0$ for all $\alpha \in \mathfrak{R}^+(\mathfrak{k}/\mathfrak{k}_{\beta})$. Finally, we see that $RR_{\beta}^K(M, \tilde{L})$ is a sum of terms of the form $m_{\beta, \mu}(\tilde{L}) \text{Hol}_T^K(t^{\mu - \alpha_I})$ where $\alpha_I = \sum_{\alpha \in I} \alpha$ and I is a subset of $\mathfrak{R}^+(\mathfrak{k}/\mathfrak{k}_{\beta})$. We know that $\text{Hol}_T^K(t^{\mu'})$ is either 0 or the character of an irreducible representation (times ± 1); in particular $\text{Hol}_T^K(t^{\mu'})$ is equal to ± 1 only if $(\mu', X) \leq 0$ for every $X \in \mathfrak{t}_+$ (see Appendix B in [33]). So $[RR_{\beta}^K(M, \tilde{L})]^K \neq 0$ only if there exists a weight μ such that $m_{\beta, \mu}(\tilde{L}) \neq 0$ and that $\text{Hol}_T^K(t^{\mu - \alpha_I}) = \pm 1$. The first condition imposes $(\mu, \beta) \geq \|\beta\|^2 + \frac{1}{2} \text{Tr}_{\beta} |\mathcal{N}_{\mathcal{Z}}|$ for some connected component $\mathcal{Z} \subset \mathcal{X}^{\beta}$, and the second one gives $(\mu, \beta) \leq (\alpha_I, \beta)$. Combining the two we end up with

$$\|\beta\|^2 + \frac{1}{2} \text{Tr}_{\beta} |\mathcal{N}_{\mathcal{Z}}| \leq (\alpha_I, \beta) \leq \sum_{\alpha \in \mathfrak{R}^+(\mathfrak{k}/\mathfrak{k}_{\beta})} (\alpha, \beta) = 2(\rho_c, \beta),$$

for some connected component $\mathcal{Z} \subset \mathcal{X}^{\beta}$ which intersects $\Phi^{-1}(\beta)$. This completes the proof that $[RR_{\beta}^K(M, \tilde{L})]^K = 0$ if $\|\beta\|^2 + \frac{1}{2} \text{Tr}_{\beta} |\mathcal{N}_{\mathcal{Z}}| > 2(\rho_c, \beta)$ for every component $\mathcal{Z} \subset \mathcal{X}^{\beta}$ which intersects $\Phi^{-1}(\beta)$. \square

4. QUANTIZATION COMMUTES WITH REDUCTION

Let (M, ω, Φ) be a Hamiltonian K -manifold equipped with an almost complex structure J . In this section, we assume that the moment map Φ is *proper* and that the set $\text{Cr}(\|\Phi\|^2)$ of critical points of $\|\Phi\|^2: M \rightarrow \mathbb{R}$ is *compact*. We denote the corresponding Riemann-Roch character by $RR_{\Phi}^K(M, -)$ (see Definition 3.2). Let $\Delta := \Phi(M) \cap \mathfrak{t}_+^*$ be the moment polyhedron.

The main result of this section is the following

Theorem 4.1. *Suppose that M satisfies Assumption 3.6 at every $\tilde{\mu} \in \mathfrak{t}^*$, and that the infinitesimal stabilizers for the K -action on M are Abelian. If \tilde{L} is a κ -prequantum line bundle over (M, ω, Φ, J) , we have*

$$(4.25) \quad RR_{\Phi}^K(M, \tilde{L}) = \varepsilon \sum_{\mu \in \Lambda_+^*} \mathcal{Q}(M_{\mu + \rho_c}) \chi_{\mu}^K,$$

where $\varepsilon = \pm 1$ is the ‘quotient’ of the orientation $o(J)$ defined by the almost complex structure and the orientation $o(\omega)$ defined by the symplectic form. Here the integer

$\mathcal{Q}(M_{\mu+\rho_c})$ is computed by Proposition 2.4. In particular, $\mathcal{Q}(M_{\mu+\rho_c}) = 0$ if $\mu + \rho_c$ does not belong to the relative interior of Δ .

The same result holds in the traditional ‘prequantum’ case. Suppose that M satisfies Assumption 3.6 at every $\mu \in \mathfrak{t}^*$, and that the almost complex structure J is compatible with ω . If L is prequantum line bundle over (M, ω, Φ) , we have $RR_\Phi^K(M, L) = \sum_{\mu \in \Lambda_+^*} RR(M_\mu, L_\mu) \chi_\mu^K$.

The next Lemma is the first step in computing the K -multiplicities $m_\mu(\tilde{L})$ of $RR_\Phi^K(M, \tilde{L})$. Since (M, Φ) satisfies Assumption 3.6 at every $\tilde{\mu}$, we know from Proposition 3.7 that $m_\mu(\tilde{L}) = [RR_{\Phi_{\tilde{\mu}}}^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, \tilde{L} \boxtimes \tilde{\mathbb{C}}_{[-\mu]})]^K$ for every $\mu \in \Lambda_+^*$.

Let $RR_0^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, -)$ be the Riemann-Roch character localized near $\Phi_{\tilde{\mu}}^{-1}(0) \simeq \Phi^{-1}(\mu + \rho_c)$ (see Definition 3.8). This map is the *zero map* if $\Phi^{-1}(\mu + \rho_c) = \emptyset$.

Lemma 4.2. *Let \tilde{L} be a κ -prequantum line bundle over M . Suppose that the infinitesimal stabilizers for the K -action are Abelian and that Assumption 3.6 is satisfied at $\tilde{\mu}$. We have then*

$$(4.26) \quad m_\mu(\tilde{L}) = \left[RR_0^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, \tilde{L} \boxtimes \tilde{\mathbb{C}}_{[-\mu]}) \right]^K.$$

In particular $m_\mu(\tilde{L}) = 0$ if $\mu + \rho_c$ does not belong to the moment polyhedron Δ .

Proof. The lemma follows from Corollary 3.12, applied to the Hamiltonian manifold $\mathcal{X} := M \times \overline{\mathcal{O}^{\tilde{\mu}}}$, with moment map $\Phi_{\tilde{\mu}}$ and κ -prequantum line bundle $\tilde{L} \boxtimes \tilde{\mathbb{C}}_{[-\mu]}$. Let $\beta \neq 0$ such that $\mathcal{X}^\beta \cap \Phi_{\tilde{\mu}}^{-1}(\beta) \neq \emptyset$. Let \mathcal{N} be the normal bundle of \mathcal{X}^β in \mathcal{X} , and let $x \in \mathcal{X}^\beta \cap \Phi_{\tilde{\mu}}^{-1}(\beta)$. From the criterion of Proposition 3.11, it is sufficient to show that

$$(4.27) \quad \|\beta\|^2 + \frac{1}{2} \mathbf{Tr}_\beta |\mathcal{N}_x| - 2(\rho_c, \beta) > 0.$$

Write $x = (m, \xi)$ with $m \in M^\beta$ and $\xi \in (\mathcal{O}^{\tilde{\mu}})^\beta$. We know that $\mathbf{Tr}_\beta |\mathcal{N}_x| = \mathbf{Tr}_\beta |\mathbf{T}_x \mathcal{X}| = \mathbf{Tr}_\beta |\mathbf{T}_m M| + \mathbf{Tr}_\beta |\mathbf{T}_\xi \mathcal{O}^{\tilde{\mu}}|$.

Since the stabilizer $\mathfrak{k}_\xi \simeq \mathfrak{t}$ is Abelian and $\beta \in \mathfrak{k}_\xi$ we have $\mathfrak{k}_\xi \subset \mathfrak{k}_\beta$. Then the tangent space $\mathbf{T}_\xi \mathcal{O}^{\tilde{\mu}} \simeq \mathfrak{k}/\mathfrak{k}_\xi$ contains a copy of $\mathfrak{k}/\mathfrak{k}_\beta$, so $\mathbf{Tr}_\beta |\mathbf{T}_\xi \mathcal{O}^{\tilde{\mu}}| \geq \mathbf{Tr}_\beta |\mathfrak{k}/\mathfrak{k}_\beta| = 2(\rho_c, \beta)$. On the other hand, $\mathbf{T}_m M$ contains the vector space $E_m \simeq \mathfrak{k}/\mathfrak{k}_m$ spanned by $X_M(m)$, $X \in \mathfrak{k}$. We have assume that the stabilizer subalgebra \mathfrak{k}_m is Abelian, and since $\beta \in \mathfrak{k}_m$, we get $\mathfrak{k}_m \subset \mathfrak{k}_\beta$. Thus $\mathfrak{k}/\mathfrak{k}_\beta \subset E_m \subset \mathbf{T}_m M$ and $\mathbf{Tr}_\beta |\mathbf{T}_m M| \geq 2(\rho_c, \beta)$. Finally (4.27) is proved since $\frac{1}{2}(\mathbf{Tr}_\beta |\mathbf{T}_m M| + \mathbf{Tr}_\beta |\mathbf{T}_\xi \mathcal{O}^{\tilde{\mu}}|) \geq 2(\rho_c, \beta)$. \square

The remaining part of this section is devoted to the proof of Theorem 4.1. Following the preceding Lemma we have to show that

$$(4.28) \quad \left[RR_0^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, \tilde{L} \boxtimes \tilde{\mathbb{C}}_{[-\mu]}) \right]^K = \varepsilon \mathcal{Q}(M_{\mu+\rho_c}),$$

where $\mathcal{Q}(M_{\mu+\rho_c})$ is defined in Proposition 2.4.

In Subsection 4.1, we recall the basic notions about Spin^c-structures. The existence of induced Spin^c-structures on symplectic quotient is proved in Subsection 4.2. The proof of (4.28) is settled in Subsection 4.3. We give in the same time the proof of the ‘hard part’ of Proposition 2.4: the fact that the index $\mathcal{Q}(M_\xi)$ does not depend on ξ , for ξ sufficiently close to $\mu + \rho_c$.

4.1. Spin^c structures and symbols. We refer to Lawson-Michelsohn [25] for background on Spin^c -structures, and to Duistermaat [14] for a discussion of the symplectic case.

The group Spin_n is the connected double cover of the group SO_n . Let $\eta : \text{Spin}_n \rightarrow \text{SO}_n$ be the covering map, and let ε be the element who generates the kernel. The group Spin_n^c is the quotient $\text{Spin}_n \times_{\mathbb{Z}_2} \text{U}_1$, where \mathbb{Z}_2 acts by $(\varepsilon, -1)$. There are two canonical group homomorphisms

$$\eta : \text{Spin}_n^c \rightarrow \text{SO}_n \quad , \quad \text{Det} : \text{Spin}_n^c \rightarrow \text{U}_1 \quad .$$

Note that $\eta^c = (\eta, \text{Det}) : \text{Spin}_n^c \rightarrow \text{SO}_n \times \text{U}_1$ is a double covering map.

Let $p : E \rightarrow M$ be a oriented Euclidean vector bundle of rank n , and let $\text{P}_{\text{SO}}(E)$ be its bundle of oriented orthonormal frames. A Spin^c -structure on E is a Spin_n^c principal bundle $\text{P}_{\text{Spin}^c}(E) \rightarrow M$, together with a Spin^c -equivariant map $\text{P}_{\text{Spin}^c}(E) \rightarrow \text{P}_{\text{SO}}(E)$. The line bundle

$$(4.29) \quad \mathbb{L} := \text{P}_{\text{Spin}^c}(E) \times_{\text{Det}} \mathbb{C}$$

is called the *canonical line bundle* associated to $\text{P}_{\text{Spin}^c}(E)$. We have then a double covering map⁴

$$(4.30) \quad \eta_E^c : \text{P}_{\text{Spin}^c}(E) \longrightarrow \text{P}_{\text{SO}}(E) \times \text{P}_{\text{U}}(\mathbb{L}) \quad ,$$

where $\text{P}_{\text{U}}(\mathbb{L}) := \text{P}_{\text{Spin}^c}(E) \times_{\text{Det}} \text{U}_1$ is the associated U_1 -principal bundle over M .

A Spin^c -structure on a oriented Riemannian manifold is a Spin^c -structure on its tangent bundle. If a group K acts on the bundle E , preserving the orientation and the Euclidean structure, we define a K -equivariant Spin^c -structure by requiring $\text{P}_{\text{Spin}^c}(E)$ to be a K -equivariant principal bundle, and (4.30) to be $(K \times \text{Spin}_n^c)$ -equivariant.

Let Δ_{2m} be the complex Spin representation of Spin_{2m}^c . Recall that $\Delta_{2m} = \Delta_{2m}^+ \oplus \Delta_{2m}^-$ inherits a canonical Clifford action $\mathbf{c} : \mathbb{R}^{2m} \rightarrow \text{End}_{\mathbb{C}}(\Delta_{2m})$ which is Spin_{2m}^c -equivariant, and which interchanges the grading : $\mathbf{c}(v) : \Delta_{2m}^{\pm} \rightarrow \Delta_{2m}^{\mp}$, for every $v \in \mathbb{R}^{2m}$. Let

$$(4.31) \quad \mathcal{S}(E) := \text{P}_{\text{Spin}^c}(E) \times_{\text{Spin}_{2m}^c} \Delta_{2m}$$

be the spinor bundle over M , with the grading $\mathcal{S}(E) := \mathcal{S}(E)^+ \oplus \mathcal{S}(E)^-$. Since $E = \text{P}_{\text{Spin}^c}(E) \times_{\text{Spin}_{2m}^c} \mathbb{R}^{2m}$, the bundle $p^*\mathcal{S}(E)$ is isomorphic to $\text{P}_{\text{Spin}^c}(E) \times_{\text{Spin}_{2m}^c} (\mathbb{R}^{2m} \oplus \Delta_{2m})$.

Let \overline{E} be the bundle E with opposite orientation. A Spin^c structure on E induces a Spin^c on \overline{E} , with the same canonical line bundle, and such that $\mathcal{S}(\overline{E})^{\pm} = \mathcal{S}(E)^{\mp}$.

Definition 4.3. Let $\text{S-Thom}(E) : p^*\mathcal{S}(E)^+ \rightarrow p^*\mathcal{S}(E)^-$ be the symbol defined by

$$\begin{aligned} \text{P}_{\text{Spin}^c}(E) \times_{\text{Spin}_{2m}^c} (\mathbb{R}^{2m} \oplus \Delta_{2m}^+) &\longrightarrow \text{P}_{\text{Spin}^c}(E) \times_{\text{Spin}_{2m}^c} (\mathbb{R}^{2m} \oplus \Delta_{2m}^-) \\ [p; v, w] &\longmapsto [p, v, \mathbf{c}(v)w] \quad . \end{aligned}$$

When E is the tangent bundle of a manifold M , the symbol $\text{S-Thom}(E)$ is denoted by $\text{S-Thom}(M)$. If a group K acts equivariantly on the Spin^c -structure, we denote by $\text{S-Thom}_K(E)$ the equivariant symbol.

⁴If P, Q are principal bundle over M respectively for the groups G and H , we denote simply by $P \times Q$ their fibering product over M which is a $G \times H$ principal bundle over M .

The characteristic set of $\text{S-Thom}(E)$ is $M \simeq \{\text{zero section of } E\}$, hence it defines a class in $\mathbf{K}(E)$ if M is compact (this class is a free generator of the $\mathbf{K}(M)$ -module $\mathbf{K}(E)$ [3]). When $E = \mathbf{T}M$, the symbol $\text{S-Thom}(M)$ corresponds to the *principal symbol* of the Spin^c Dirac operator associated to the Spin^c -structure [14]. If moreover M is compact, the number $\mathcal{Q}(M) \in \mathbb{Z}$ is defined as the index of $\text{S-Thom}(M)$. If we change the orientation, note that $\mathcal{Q}(\overline{M}) = -\mathcal{Q}(M)$.

Remark 4.4. *It should be noted that the choice of the metric on the fibers of E is not essential in the construction. Let g_0, g_1 be two metric on the fibers of E , and suppose that (E, g_0) admits a Spin^c -structure denoted by $\text{P}_{\text{Spin}^c}(E, g_0)$. The trivial homotopy $g_t = (1-t) \cdot g_0 + t \cdot g_1$ between the metrics, induces a homotopy between the principal bundles $\text{P}_{\text{SO}}(E, g_0), \text{P}_{\text{SO}}(E, g_1)$ which can be lifted to a homotopy between $\text{P}_{\text{Spin}^c}(E, g_0)$ and a Spin^c -bundle over (E, g_1) . When the base M is compact, the corresponding symbols $\text{S-Thom}(E, g_0)$ and $\text{S-Thom}(E, g_1)$ define the same class in $\mathbf{K}(E)$.*

These notions extend to the orbifold case. Let M be a manifold with a locally free action of a compact Lie group H . The quotient $\mathcal{X} := M/H$ is an orbifold, a space with finite quotient singularities. A Spin^c structure on \mathcal{X} is by definition a H -equivariant Spin^c structure on the bundle $\mathbf{T}_H M \rightarrow M$, where $\mathbf{T}_H M$ is identified with the pullback of $\mathbf{T}\mathcal{X}$ via the quotient map $\pi : M \rightarrow \mathcal{X}$. We define in the same way $\text{S-Thom}(\mathcal{X}) \in \mathbf{K}_{\text{orb}}(\mathbf{T}\mathcal{X})$, such that $\pi^* \text{S-Thom}(\mathcal{X}) = \text{S-Thom}_H(\mathbf{T}_H M)$. Here \mathbf{K}_{orb} denotes the K -theory of *proper* vector bundles [21]. The pullback by π induces an isomorphism $\pi^* : \mathbf{K}_{\text{orb}}(\mathbf{T}\mathcal{X}) \simeq \mathbf{K}_H(\mathbf{T}_H M)$. The number $\mathcal{Q}(\mathcal{X}) \in \mathbb{Z}$ is defined as the index of $\text{S-Thom}(\mathcal{X})$, or equivalently as the multiplicity of the trivial representation in $\text{Index}_M^H(\text{S-Thom}_H(\mathbf{T}_H M))$.

Consider now the case of a *Hermitian* vector bundle $E \rightarrow M$, of complex rank m . The orientation on the fibers of E is given by the complex structure J . Let $\text{P}_U(E)$ be the bundle of unitary frames on E . We denote by $i : U_m \hookrightarrow \text{SO}_{2m}$ the canonical inclusion map. We have a morphism $j : U_m \rightarrow \text{Spin}_{2m}^c$ which makes the diagram

$$(4.32) \quad \begin{array}{ccc} U_m & \xrightarrow{j} & \text{Spin}_{2m}^c \\ & \searrow i \times \det & \downarrow \eta^c \\ & & \text{SO}_{2m} \times U_1 \end{array}$$

commutative [25]. Then

$$(4.33) \quad \text{P}_{\text{Spin}^c}(E) := \text{Spin}_{2m}^c \times_j \text{P}_U(E)$$

defines a Spin^c -structure over E , with canonical line bundle equal to $\det_{\mathbb{C}} E$.

Lemma 4.5. *Let M be a manifold equipped with an almost complex structure J . The symbol $\text{S-Thom}(M)$ defined by the Spin^c -structure (4.33), and the Thom symbol $\text{Thom}(M, J)$ defined in Section 3.2 coincide.*

Proof. The Spinor bundle \mathcal{S} is of the form $\text{P}_{\text{Spin}^c}(\mathbf{T}M) \times_{\text{Spin}_{2m}^c} \Delta_{2m} = \text{P}_U(\mathbf{T}M) \times_{U_m} \Delta_{2m}$. The map $\mathbf{c} : \mathbb{R}^{2m} \rightarrow \text{End}_{\mathbb{C}}(\Delta_{2m})$, when restricted to the U_m -equivariant action through j , is equivalent to the Clifford map $\text{Cl} : \mathbb{R}^{2m} \rightarrow \text{End}_{\mathbb{C}}(\wedge^{\mathbb{C}^m})$ (with the canonical action of U_m on \mathbb{R}^{2m} and $\wedge^{\mathbb{C}^m}$). Then $\mathcal{S} = \wedge_{\mathbb{C}} \mathbf{T}M$ endowed with the Clifford action. \square

Lemma 4.6. *Let P be a Spin^c -structure over M , with bundle of spinors \mathcal{S} , and canonical line bundle \mathbb{L} . For every Hermitian line bundle $L \rightarrow M$, there exists a unique Spin^c -structure P_L with bundle of spinors $\mathcal{S} \otimes L$, and canonical line bundle $\mathbb{L} \otimes L^2$ (P_L is called the Spin^c -structure P twisted by L).*

Proof. Take $P_L = P \times_{\text{U}_1} \text{P}_\text{U}(L)$.

We finish this subsection with the following definitions. Let (M, o) be an oriented manifold. Suppose that

- a connected compact Lie K acts on M
- (M, o, K) carries a K -equivariant Spin^c -structure
- one has an equivariant map $\Psi : M \rightarrow \mathfrak{k}$.

Suppose first that M is compact. The symbol $\text{S-Thom}_K(M)$ is then elliptic and defines a map

$$\mathcal{Q}^K(M, -) : \mathbf{K}_K(M) \rightarrow R(K)$$

by the relation $\mathcal{Q}^K(M, V) := \text{Index}_M^K(\text{S-Thom}_K(M) \otimes V)$. Thus $\mathcal{Q}^K(M, V)$ is the equivariant index of the Spin^c Dirac operator on M twisted by V .

Let Ψ_M be the equivariant vector field on M defined by $\Psi_M(m) := \Psi(m)_M|_m$.

Definition 4.7. *The symbol $\text{S-Thom}_K(M)$ deformed by the map Ψ , which is denoted by $\text{S-Thom}_K^\Psi(M)$, is defined by the relation*

$$\text{S-Thom}_K^\Psi(M)(m, v) := \text{S-Thom}_K(M)(m, v - \Psi_M(m))$$

for any $(m, v) \in \mathbf{T}M$. The symbol $\text{S-Thom}_K^\Psi(M)$ is transversally elliptic if and only if $\{m \in M, \Psi_M(m) = 0\}$ is compact. When this holds one defines the localized map $\mathcal{Q}_\Psi^K(M, V) := \text{Index}_M^K(\text{S-Thom}_K^\Psi(M) \otimes V)$.

We finish this section with an adaptation of Lemma 9.4 and Corollary 9.5 of [33][Appendix B] to the localized map $\mathcal{Q}_\Psi^K(M, -)$. Let $\beta \in \mathfrak{k}_+^*$ be a non-zero element in the center of the Lie algebra $\mathfrak{k} \cong \mathfrak{k}^*$ of K . We suppose here that the subtorus $i : \mathbb{T}_\beta \hookrightarrow K$, which is equal to the closure of $\{\exp(t\beta), t \in \mathbb{R}\}$, acts trivially on M . Let $m_\mu(V)$, $\mu \in \Lambda_+^*$ be the K -multiplicities of $\mathcal{Q}_\Psi^K(M, V)$.

Lemma 4.8. *If $m_\mu(V) \neq 0$, $i^*(\mu)$ is a weight for the action of \mathbb{T}_β on $V \otimes \mathbb{L}^{\frac{1}{2}}$. If each weight α for the action of \mathbb{T}_β on $V \otimes \mathbb{L}^{\frac{1}{2}}$ satisfies $(\alpha, \beta) > 0$, then $[\mathcal{Q}_\Psi^K(M, V)]^K = 0$.*

4.2. Spin^c structures on symplectic reductions. Let (M, ω, Φ) be a Hamiltonian K -manifold, such that Φ is proper. Let J be a K -invariant almost complex structure on M . And let \tilde{L} be a κ -prequantum line bundle over (M, ω, J) . Since we do not impose a compatibility condition between J and ω , the almost complex structure does not descend to the symplectic quotients in general. Nevertheless we prove in this section that the Spin^c prequantization defined by the data (\tilde{L}, J) induces a Spin^c prequantization on the symplectic quotients $M_{\mu+\rho_c}$.

Let \mathcal{Y} be the subset $\Phi^{-1}(\text{interior}(\mathfrak{k}_+^*))$. When $\mathcal{Y} \neq \emptyset$, the Principal-cross-section Theorem tells us that \mathcal{Y} is a Hamiltonian T -submanifold of M , with moment map the restriction of Φ to \mathcal{Y} [26].

Lemma 4.9. *If the infinitesimal stabilizers for the K -action on M are Abelian, the symplectic slice \mathcal{Y} is not empty.*

Proof. There exists a unique relatively open face τ of the Weyl chamber \mathfrak{t}_+^* such that $\Phi(M) \cap \tau$ is dense in $\Phi(M) \cap \mathfrak{t}_+^*$. The face τ is called the principal face of (M, Φ) [26]. All points in the open face τ have the same connected centralizer K_τ . The Principal-cross-section Theorem tells us that $\mathcal{Y}_\tau := \Phi^{-1}(\tau)$ is a Hamiltonian K_τ -manifold, where $[K_\tau, K_\tau]$ acts trivially [26]. Here we have assumed that the subalgebras $\mathfrak{k}_m := \{X \in \mathfrak{k}, X_M(m) = 0\}$, $m \in M$, are Abelian. Hence $[\mathfrak{k}_\tau, \mathfrak{k}_\tau] \subset \mathfrak{k}_m$ for every $m \in \mathcal{Y}_\tau$, and this imposes $[\mathfrak{k}_\tau, \mathfrak{k}_\tau] = 0$. Therefore the subgroup K_τ is Abelian, and this is the case only if τ is the interior of the Weyl chamber. \square

For the remaining of this section, we assume that $\mathcal{Y} \neq \emptyset$, so that the relative interior Δ° of the moment polyhedron is a dense subset of $\Phi(\mathcal{Y})$. On M , we have the orientation $o(J)$ defined by the almost complex structure and the orientation $o(\omega)$ defined by the symplectic form. We denote their ‘quotient’ by $\varepsilon = \pm 1$. On the symplectic quotients we will have also two orientations, one induces by ω , and the other induces by J , with the same ‘quotient’ ε .

Proposition 4.10. *The almost complex structure J induces*

- i) an orientation $o(\mathcal{Y})$ on \mathcal{Y} , and*
- ii) a T -equivariant Spin^c structure on $(\mathcal{Y}, o(\mathcal{Y}))$ with canonical line bundle $\det_{\mathbb{C}}(\mathbf{T}M|_{\mathcal{Y}}) \otimes \mathbb{C}_{-2\rho_c}$.*

Proof. On \mathcal{Y} , we have the decomposition $\mathbf{T}M|_{\mathcal{Y}} = \mathbf{T}\mathcal{Y} \oplus [\mathfrak{k}/\mathfrak{t}]$, where $[\mathfrak{k}/\mathfrak{t}]$ denotes the trivial bundle $\mathcal{Y} \times \mathfrak{k}/\mathfrak{t}$ corresponding of the subspace of $\mathbf{T}M|_{\mathcal{Y}}$ formed by the vector fields generated by the infinitesimal action of $\mathfrak{k}/\mathfrak{t}$. The choice of the Weyl chamber induces a complex structure on $\mathfrak{k}/\mathfrak{t}$, and hence an orientation $o([\mathfrak{k}/\mathfrak{t}])$. This orientation can be also defined by a symplectic form of the type $\omega_{\mathfrak{k}/\mathfrak{t}}(X, Y) = \langle \xi, [X, Y] \rangle$, where ξ belongs to the interior \mathfrak{t}_+^* . Let $o(\mathcal{Y})$ be the orientation on \mathcal{Y} defined by $o(J)|_{\mathcal{Y}} = o(\mathcal{Y})o([\mathfrak{k}/\mathfrak{t}])$. On \mathcal{Y} , we have also the orientation $o(\omega_{\mathcal{Y}})$ defined by the symplectic form $\omega_{\mathcal{Y}}$. Note that if $o(J) = \varepsilon o(\omega)$, we have also $o(\mathcal{Y}) = \varepsilon o(\omega_{\mathcal{Y}})$.

Let $P := \text{Spin}_{2n}^c \times_{U_n} P_U(\mathbf{T}M)$ be the Spin^c structure on M induced by J (see (4.33)). When restricted to \mathcal{Y} , $P|_{\mathcal{Y}}$ defines a Spin^c structure on the bundle $\mathbf{T}\mathcal{Y} \oplus [\mathfrak{k}/\mathfrak{t}]$. Let q be a T -invariant Riemannian structure on $\mathbf{T}\mathcal{Y} \oplus [\mathfrak{k}/\mathfrak{t}]$ such that $\mathbf{T}\mathcal{Y}$ is orthogonal with $[\mathfrak{k}/\mathfrak{t}]$, and q equals the Killing form on $[\mathfrak{k}/\mathfrak{t}]$. Following Remark 4.4, $P|_{\mathcal{Y}}$ induces a Spin^c structure P' on $(\mathbf{T}\mathcal{Y} \oplus [\mathfrak{k}/\mathfrak{t}], q)$, with the same canonical line bundle $\mathbb{L} = \det_{\mathbb{C}}(\mathbf{T}M|_{\mathcal{Y}})$. Since the $\text{SO}_{2k} \times U_l$ -principal bundle $P_{\text{SO}}(\mathbf{T}\mathcal{Y}) \times U(\mathfrak{k}/\mathfrak{t})$ is a reduction⁵ of the SO_{2n} principal bundle $P_{\text{SO}}(\mathbf{T}\mathcal{Y} \oplus [\mathfrak{k}/\mathfrak{t}])$, we have the commutative diagram

$$(4.34) \quad \begin{array}{ccc} Q & \longrightarrow & P_{\text{SO}}(\mathbf{T}\mathcal{Y}) \times U(\mathfrak{k}/\mathfrak{t}) \times P_U(\mathbb{L}) \\ \downarrow & & \downarrow \\ P' & \longrightarrow & P_{\text{SO}}(\mathbf{T}\mathcal{Y} \oplus [\mathfrak{k}/\mathfrak{t}]) \times P_U(\mathbb{L}), \end{array}$$

where Q is a $(\eta^c)^{-1}(\text{SO}_{2k} \times U_l) \simeq \text{Spin}_{2k}^c \times U_l$ -principal bundle. Finally we see that $Q' = Q/U_l$ is a Spin^c structure on $\mathbf{T}\mathcal{Y}$. Since $(U(\mathfrak{k}/\mathfrak{t}) \times P_U(\mathbb{L}))/U_l \simeq P_U(\mathbb{L} \otimes \mathbb{C}_{-2\rho_c})$, the corresponding canonical line bundle is $\mathbb{L}' = \mathbb{L} \otimes \mathbb{C}_{-2\rho_c}$. \square

Let $\text{Aff}(\Delta)$ be the affine subspace generated by moment polyhedron Δ , and let $\overrightarrow{\Delta}$ be the subspace of \mathfrak{t}^* generated by $\{m - n \mid m, n \in \Delta\}$. Let T_{Δ} the subtorus

⁵Here $2n = \dim M$, $2k = \dim \mathcal{Y}$ and $2l = \dim(\mathfrak{k}/\mathfrak{t})$, so $n = k + l$.

of T with Lie algebra \mathfrak{t}_Δ equal to the orthogonal (for the duality) of $\vec{\Delta}$. It is not difficult to see that T_Δ corresponds to the connected component of the principal stabilizer for the T -action on \mathcal{Y} .

Here we consider the symplectic quotient $M_\xi := \Phi^{-1}(\xi)/T$ for *generic quasi-regular values* $\xi \in \Delta^\circ$ (see Definition 2.3). For such ξ , the fiber $\Phi^{-1}(\xi)$ is a smooth submanifold of M , with a locally free action of T/T_Δ , and with a tubular neighborhood equivariantly diffeomorphic to $\Phi^{-1}(\xi) \times \vec{\Delta}$. Recall that M_ξ inherits a canonical symplectic form ω_ξ .

Proposition 4.11. *Let $\mu \in \Lambda_+^*$ such that $\tilde{\mu} = \mu + \rho_c$ belongs to Δ . Let \tilde{L} be a κ -prequantum line bundle. For every generic quasi-regular value $\xi \in \Delta^\circ$, the Spin^c structures on \mathcal{Y} , when twisted by $\tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu}$, induces a Spin^c structure on the reduced space $M_\xi := \Phi^{-1}(\xi)/T$ with canonical line bundle $(L_{2\omega}|_{\Phi^{-1}(\xi)} \otimes \mathbb{C}_{-2\tilde{\mu}})/T$. Here we have two choices for the orientations : $o(M_\xi)$ induced by $o(\mathcal{Y})$, and $o(\omega_\xi)$ defined by the symplectic form ω_ξ . They are related by $o(M_\xi) = \varepsilon o(\omega_\xi)$.*

Remark 4.12. *The preceding Proposition will be used*

- i) when $\xi = \mu + \rho_c$ is a generic quasi-regular value of Φ : the symplectic quotient $(M_{\mu+\rho_c}, \omega_{\mu+\rho_c})$ is then Spin^c prequantized. Or
- ii) for general $\mu + \rho_c \in \Delta$. One takes then ξ generic quasi-regular close enough to $\mu + \rho_c$.

Proof of the Proposition. Let $\xi \in \Delta^\circ$ be a generic quasi-regular value of Φ , and $\mathcal{Z} := \Phi^{-1}(\xi)$. This is a submanifold of \mathcal{Y} with a trivial action of T_Δ and a locally free action of T/T_Δ . We denote the quotient map by $\pi : \mathcal{Z} \rightarrow M_\xi$. We identify $\pi^*(\mathbf{T}M_\xi)$ with the orthogonal complement (with respect to a Riemannian metric) of the trivial bundle $[\mathfrak{t}/\mathfrak{t}_\Delta]$ formed by the vector fields generated by the infinitesimal action of $\mathfrak{t}/\mathfrak{t}_\Delta$. On the other hand the tangent bundle $\mathbf{T}\mathcal{Y}$, when restricted to \mathcal{Z} , decomposes as $\mathbf{T}\mathcal{Y}|_{\mathcal{Z}} = \mathbf{T}\mathcal{Z} \oplus [\vec{\Delta}]$, so we have

$$\begin{aligned} \mathbf{T}\mathcal{Y}|_{\mathcal{Z}} &= \pi^*(\mathbf{T}M_\xi) \oplus [\mathfrak{t}/\mathfrak{t}_\Delta] \oplus [\vec{\Delta}] \\ (4.35) \quad &= \pi^*(\mathbf{T}M_\xi) \oplus [\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C}] , \end{aligned}$$

with the convention $\mathfrak{t}/\mathfrak{t}_\Delta = \mathfrak{t}/\mathfrak{t}_\Delta \otimes i\mathbb{R}$ and $\vec{\Delta} = \mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{R}$. Since $\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C}$ is canonically oriented by the complex multiplication by i , the orientation $o(\mathcal{Y})$ determines an orientation $o(M_\xi)$ on $\mathbf{T}M_\xi$ through (4.35).

Now we proceed like the proof of Proposition 4.11. Let Q' be the Spin^c structure on \mathcal{Y} introduced in Proposition 4.11, and let Q^μ be Q' twisted by the line bundle $\tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu}$: its canonical line bundle is $\det_{\mathbb{C}}(\mathbf{T}M)|_{\mathcal{Y}} \otimes \mathbb{C}_{-2\rho_c} \otimes (\tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu})^2 = L_{2\omega}|_{\mathcal{Y}} \otimes \mathbb{C}_{-2\tilde{\mu}}$. The $\text{SO}_{2k'} \times \text{U}_{l'}$ -principal bundle $\text{P}_{\text{SO}}(\pi^*(\mathbf{T}M_\xi)) \times \text{U}(\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C})$ is a reduction⁶ of the SO_{2k} principal bundle $\text{P}_{\text{SO}}(\pi^*(\mathbf{T}M_\xi) \oplus [\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C}])$; we have the commutative diagram

$$(4.36) \quad \begin{array}{ccc} Q'' & \longrightarrow & \text{P}_{\text{SO}}(\pi^*(\mathbf{T}M_\xi)) \times \text{U}(\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C}) \times \text{P}_{\text{U}}(\mathbb{L}|_{\mathcal{Z}}) \\ \downarrow & & \downarrow \\ Q^\mu|_{\mathcal{Z}} & \longrightarrow & \text{P}_{\text{SO}}(\pi^*(\mathbf{T}M_\xi) \oplus [\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C}]) \times \text{P}_{\text{U}}(\mathbb{L}|_{\mathcal{Z}}) , \end{array}$$

⁶Here $2k = \dim \mathcal{Y}$, $2k' = \dim M_\xi$ and $l' = \dim(\mathfrak{t}/\mathfrak{t}_\Delta)$, so $k = k' + l'$.

where $\mathbb{L} = L_{2\omega}|_{\mathcal{Y}} \otimes \mathbb{C}_{-2\tilde{\mu}}$. Here Q'' is a $(\eta^c)^{-1}(\mathrm{SO}_{2k'} \times \mathrm{U}_{l'}) \simeq \mathrm{Spin}_{2k'}^c \times \mathrm{U}_{l'}$ -principal bundle. The Kostant formula (2.1) tells us that the action of T_{Δ} is trivial on $\mathbb{L}|_{\mathcal{Z}}$, since $\xi - \tilde{\mu} \in \vec{\Delta}$. Thus the action of T_{Δ} is trivial on Q'' . Finally we see that $Q_{\xi} = Q''/(\mathrm{U}_{l'} \times T)$ is a Spin^c structure on M_{ξ} with canonical line bundle $\mathbb{L}_{\xi} = (L_{2\omega}|_{\mathcal{Z}} \otimes \mathbb{C}_{-2\tilde{\mu}})/T$. \square

4.3. Definition of $\mathcal{Q}(M_{\mu+\rho_c})$. First we give three different ways to define the quantity $\mathcal{Q}(M_{\mu+\rho_c}) \in \mathbb{Z}$ for any $\mu \in \Lambda_+^*$. The compatibility of these different definitions proves Theorem 4.1 and the ‘hard part’ of Proposition 2.4 simultaneously. First of all $\mathcal{Q}(M_{\mu+\rho_c}) = 0$ if $\mu + \rho_c \notin \Delta$.

First definition.

If $\mu + \rho_c \in \Delta^o$ is a *generic quasi regular* value of Φ , $M_{\mu+\rho_c} := \Phi^{-1}(\mu + \rho_c)/T$ is a symplectic orbifold. We know from Proposition 4.11 that $M_{\mu+\rho_c}$ inherits Spin^c -structures, with the same canonical line bundle $(L_{2\omega}|_{\Phi^{-1}(\tilde{\mu})} \otimes \mathbb{C}_{-2\tilde{\mu}})/T$, for the two choices of orientation $o(M_{\mu+\rho_c})$ and $o(\omega_{\mu+\rho_c})$. We denote the index of the Spin^c Dirac operator associated to the Spin^c structure on $(M_{\mu+\rho_c}, o(\omega_{\mu+\rho_c}))$ by $\mathcal{Q}(M_{\mu+\rho_c}) \in \mathbb{Z}$ and the index of the Spin^c Dirac operator associated to the Spin^c structure on $(M_{\mu+\rho_c}, o(M_{\mu+\rho_c}))$ by $\mathcal{Q}(M_{\mu+\rho_c}, o(M_{\mu+\rho_c}))$. Since $o(M_{\mu+\rho_c}) = \varepsilon o(\omega_{\mu+\rho_c})$, we have $\mathcal{Q}(M_{\mu+\rho_c}) = \varepsilon \mathcal{Q}(M_{\mu+\rho_c}, o(M_{\mu+\rho_c}))$.

Second definition.

We can also define $\mathcal{Q}(M_{\mu+\rho_c})$ by shift ‘desingularization’ as follows. If $\mu + \rho_c \in \Delta$, one considers generic quasi regular values $\xi \in \Delta^o$, close enough to $\mu + \rho_c$. Following Proposition 4.11, $M_{\xi} = \Phi^{-1}(\xi)/T$ inherits a Spin^c structure, with canonical line bundle $(L_{2\omega}|_{\Phi^{-1}(\xi)} \otimes \mathbb{C}_{-2\tilde{\mu}})/T$. Then we set $\mathcal{Q}(M_{\mu+\rho_c}) := \mathcal{Q}(M_{\xi})$, where the RHS is the index of the Spin^c Dirac operator associated to the Spin^c structure on $(M_{\xi}, o(\omega_{\xi}))$. If we take the orientation $o(M_{\xi})$ induced by $o(\mathcal{Y})$ we have another index $\mathcal{Q}(M_{\xi}, o(M_{\xi})) = \varepsilon \mathcal{Q}(M_{\xi})$. Here one has to show that these quantities does not depend on the choice of ξ when ξ is close enough to $\mu + \rho_c$. We will see that $\mathcal{Q}(M_{\mu+\rho_c}) = 0$ when $\mu + \rho_c \notin \Delta^o$.

Third definition.

We can use the characterization of the multiplicity $m_{\mu}(\tilde{L})$ given in Lemma 4.2. The number $\mathcal{Q}(M_{\mu+\rho_c})$ is the multiplicity of the trivial representation in $\varepsilon RR_0^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, \tilde{L} \otimes \tilde{\mathbb{C}}_{[-\mu]})$.

We have to show the compatibility of these definitions, that is

$$(4.37) \quad \text{if } \mu + \rho_c \in \Delta^o : \left[RR_0^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, \tilde{L} \otimes \tilde{\mathbb{C}}_{[-\mu]}) \right]^K = \mathcal{Q}(M_{\xi}, o(M_{\xi}))$$

for any generic quasi regular value $\xi \in \Delta^o$ close enough to $\mu + \rho_c$. And

$$(4.38) \quad \text{if } \mu + \rho_c \notin \Delta^o : \left[RR_0^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, \tilde{L} \otimes \tilde{\mathbb{C}}_{[-\mu]}) \right]^K = 0.$$

We have proved already (Lemma 4.2) that $[RR_0^K(M \times \overline{\mathcal{O}^{\tilde{\mu}}}, \tilde{L} \otimes \tilde{\mathbb{C}}_{[-\mu]})]^K = 0$ if $\mu + \rho_c \notin \Delta$.

We work now with a fixed element $\mu \in \Lambda_+^*$ such that $\tilde{\mu} = \mu + \rho_c$ belongs to Δ . During the remaining part of this section, \mathcal{Y} will denote a small T -invariant open neighborhood of $\Phi^{-1}(\mu + \rho_c)$ in the symplectic slice $\Phi^{-1}(\text{interior}(\mathfrak{t}_+^*))$.

We will check in Subsection 4.5, that the functions $\|\Phi - \tilde{\mu}\|^2$ and $\|\Phi - \xi\|^2$ have compact critical set on \mathcal{Y} when $\xi \in \text{Aff}(\Delta)$ is close enough to $\tilde{\mu}$. Since the manifold $(\mathcal{Y}, o(\mathcal{Y}))$ carries a T -invariant Spin^c -structure, we consider the localized maps $\mathcal{Q}_{\Phi-\tilde{\mu}}^T(\mathcal{Y}, -)$ and $\mathcal{Q}_{\Phi-\xi}^T(\mathcal{Y}, -)$ (see Definition 4.7). The proof of (4.37) and (4.38) is divided into two steps. We first relate the maps $RR_0^K(M \times \overline{\mathcal{O}^\mu}, -)$ and $\mathcal{Q}_{\Phi-\tilde{\mu}}^T(\mathcal{Y}, -)$ through the induction map

$$(4.39) \quad \text{Ind}_T^K : \mathcal{C}^{-\infty}(T) \longrightarrow \mathcal{C}^{-\infty}(K)^K.$$

Here $\mathcal{C}^{-\infty}(T)$, $\mathcal{C}^{-\infty}(K)$ denote respectively the set of generalized functions on T and K , and the K invariants are taken with the conjugation action. The map Ind_T^K is defined as follows : for $\phi \in \mathcal{C}^{-\infty}(T)$, we have $\int_K \text{Ind}_T^K(\phi)(k)f(k)dk = \frac{\text{vol}(K, dk)}{\text{vol}(T, dt)} \int_T \phi(t)f|_T(t)dt$, for every $f \in \mathcal{C}^\infty(K)^K$.

Proposition 4.13. *Let E and F be respectively K -equivariant complex vector bundles over M and $\mathcal{O}^{\tilde{\mu}}$. We have the following equality*

$$RR_0^K(M \times \overline{\mathcal{O}^\mu}, E \boxtimes F) = \text{Ind}_T^K \left(\mathcal{Q}_{\Phi-\tilde{\mu}}^T(\mathcal{Y}, E|_{\mathcal{Y}} \otimes F|_{\bar{e}}) \right)$$

in $R^{-\infty}(K)$. It gives in particular that

$$\left[RR_0^K(M \times \overline{\mathcal{O}^\mu}, E \boxtimes F) \right]^K = \left[\mathcal{Q}_{\Phi-\tilde{\mu}}^T(\mathcal{Y}, E|_{\mathcal{Y}} \otimes F|_{\bar{e}}) \right]^T.$$

After we compute the map $\mathcal{Q}_{\Phi-\tilde{\mu}}^T(\mathcal{Y}, -)$ by making the shift $\tilde{\mu} \rightarrow \xi$.

Proposition 4.14. *Suppose $\xi \in \text{Aff}(\Delta)$ is close enough to $\tilde{\mu}$. Then*

- i) *the maps $\mathcal{Q}_{\Phi-\tilde{\mu}}^T(\mathcal{Y}, -)$ and $\mathcal{Q}_{\Phi-\xi}^T(\mathcal{Y}, -)$ are equal,*
- ii) *if furthermore $\xi \in \Delta^o$ is a generic quasi-regular value of Φ we get*

$$\left[\mathcal{Q}_{\Phi-\xi}^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu}) \right]^T = \mathcal{Q}(M_\xi, o(M_\xi)),$$

- iii) *and if $\xi \notin \Delta$, $[\mathcal{Q}_{\Phi-\xi}^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu})]^T = 0$.*

Finally, if $\xi \in \text{Aff}(\Delta)$ is close enough to $\tilde{\mu}$, Propositions 4.13 and 4.14 give

$$(4.40) \quad \begin{aligned} \left[RR_0^K(M \times \overline{\mathcal{O}^\mu}, \tilde{L} \otimes \tilde{\mathbb{C}}_{[-\mu]}) \right]^K &= \left[\mathcal{Q}_{\Phi-\tilde{\mu}}^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu}) \right]^T \\ &= \left[\mathcal{Q}_{\Phi-\xi}^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu}) \right]^T. \end{aligned}$$

If $\mu + \rho_c \in \Delta^o$, we choose $\xi \in \Delta^o$ close to $\mu + \rho_c$: in this case (4.37) follows from (4.40) and the point ii) of Proposition 4.14. If $\mu + \rho_c \notin \Delta^o$, we choose ξ close to $\mu + \rho_c$ and not in Δ : in this case (4.38) follows from (4.40) and the point iii) of Proposition 4.14.

Propositions 4.13 and 4.14 are proved in the next subsections.

4.4. Proof of Proposition 4.13. The induction formula of Proposition 4.13 is essentially identical to the one we proved in [33]. The main difference is that the almost complex structure is not assumed to be compatible with the symplectic structure.

We identify the coadjoint orbit $\mathcal{O}^{\tilde{\mu}}$ with K/T . Let $\mathcal{H}^{\tilde{\mu}}$ be the Hamiltonian vector field of the function $\frac{-1}{2} \|\Phi_{\tilde{\mu}}\|^2 : M \times K/T \rightarrow \mathbb{R}$. Here \mathcal{Y} denotes a small

neighborhood of $\Phi^{-1}(\tilde{\mu})$ in the symplectic slice $\Phi^{-1}(\text{interior}(\mathfrak{t}_+^*))$ such that the open subset $\mathcal{U} := (K \times_T \mathcal{Y}) \times \overline{K/T}$ is a neighborhood of $\Phi_{\tilde{\mu}}^{-1}(0) = K \cdot (\Phi^{-1}(\tilde{\mu}) \times \{\bar{e}\})$ which verifies $\overline{\mathcal{U}} \cap \{\mathcal{H}^{\tilde{\mu}} = 0\} = \Phi_{\tilde{\mu}}^{-1}(0)$.

From Definition 3.8, the localized Riemann-Roch character $RR_0^K(M \times \overline{K/T}, -)$ is computed by means of the Thom class $\text{Thom}_K^{\Phi_{\tilde{\mu}}}(\mathcal{U}) \in \mathbf{K}_K(\mathbf{T}_K \mathcal{U})$. On the other hand, the localized map $\mathcal{Q}_{\Phi_{\tilde{\mu}}}^T(\mathcal{Y}, -)$ is computed by means of the class $\text{S-Thom}_T^{\Phi_{\tilde{\mu}}}(\mathcal{Y}) \in \mathbf{K}_T(\mathbf{T}_T \mathcal{Y})$ (see Definition 4.7). Proposition 4.13 will follow from a simple relation between these two transversally elliptic symbols.

First, we consider the isomorphism $\phi : \mathcal{U} \rightarrow \mathcal{U}'$, $\phi([k; y], [h]) = [k; [k^{-1}h], y]$, where $\mathcal{U}' := K \times_T (\overline{K/T} \times \mathcal{Y})$. Let $\phi^* : \mathbf{K}_K(\mathbf{T}_K \mathcal{U}') \rightarrow \mathbf{K}_K(\mathbf{T}_K \mathcal{U})$ be the induced isomorphism. Then one considers the inclusion $i : T \hookrightarrow K$ which induces an isomorphism $i_* : \mathbf{K}_T(\mathbf{T}_T(\overline{K/T} \times \mathcal{Y})) \rightarrow \mathbf{K}_K(\mathbf{T}_K \mathcal{U}')$ (see [1, 33]). Let $j : \mathcal{Y} \hookrightarrow \overline{K/T} \times \mathcal{Y}$ be the T -invariant inclusion map defined by $j(y) := (\bar{e}, y)$. We have then a pushforward map $j_! : \mathbf{K}_T(\mathbf{T}_T \mathcal{Y}) \rightarrow \mathbf{K}_T(\mathbf{T}_T(\overline{K/T} \times \mathcal{Y}))$. Finally we get a map

$$\Theta := \phi^* \circ i_* \circ j_! : \mathbf{K}_T(\mathbf{T}_T \mathcal{Y}) \longrightarrow \mathbf{K}_K(\mathbf{T}_K \mathcal{U}) ,$$

such that $\text{Index}_{\mathcal{U}}^K(\Theta(\sigma)) = \text{Ind}_T^K(\text{Index}_{\mathcal{Y}}^T(\sigma))$ for every $\sigma \in \mathbf{K}_T(\mathbf{T}_T \mathcal{Y})$ (see Section 3 in [33]).

Proposition 4.13 is an immediate consequence of the following

Lemma 4.15. *We have the equality*

$$\Theta \left(\text{S-Thom}_T^{\Phi_{\tilde{\mu}}}(\mathcal{Y}) \right) = \text{Thom}_K^{\Phi_{\tilde{\mu}}}(\mathcal{U}) .$$

Proof. Let \mathcal{S} be the bundle of spinors on $K \times_T \mathcal{Y}$: $\mathcal{S} = \mathbf{P} \times_{\text{Spin}_{2k}^c} \Delta_{2k}$, where $\mathbf{P} \rightarrow \mathbf{P}_{\text{SO}}(\mathbf{T}(K \times_T \mathcal{Y})) \times \mathbf{P}_{\text{U}}(\mathbb{L})$ is the Spin^c structure induced by the complex structure. From Proposition 4.10, we have the reductions

$$(4.41) \quad \begin{array}{ccc} \mathbf{Q} & \longrightarrow & \mathbf{P}_{\text{SO}}(\mathbf{T}\mathcal{Y}) \times \mathbf{U}(\mathfrak{k}/\mathfrak{t}) \times \mathbf{P}_{\text{U}}(\mathbb{L}|_{\mathcal{Y}}) \\ \downarrow & & \downarrow \\ \mathbf{P}|_{\mathcal{Y}} & \longrightarrow & \mathbf{P}_{\text{SO}}(\mathbf{T}\mathcal{Y} \oplus [\mathfrak{k}/\mathfrak{t}]) \times \mathbf{P}_{\text{U}}(\mathbb{L}|_{\mathcal{Y}}) \\ \downarrow & & \downarrow \\ \mathbf{P} & \longrightarrow & \mathbf{P}_{\text{SO}}(\mathbf{T}(K \times_T \mathcal{Y})) \times \mathbf{P}_{\text{U}}(\mathbb{L}) . \end{array}$$

Here \mathbf{Q}/\mathbf{U}_l is the induced Spin^c -structure on \mathcal{Y} . Let us denote by $p : \mathbf{T}(K \times_T \mathcal{Y}) \rightarrow K \times_T \mathcal{Y}$, $p_{\mathcal{Y}} : \mathbf{T}\mathcal{Y} \rightarrow \mathcal{Y}$ and $p_{K/T} : \mathbf{T}(K/T) \rightarrow K/T$ the canonical projections. Using (4.41), we see that

$$p^* \mathcal{S} = (K \times_T p_{\mathcal{Y}}^* \mathcal{S}(\mathcal{Y})) \otimes p_{K/T}^* \wedge_{\mathbb{C}} \mathbf{T}(K/T) ,$$

where $\mathcal{S}(\mathcal{Y})$ is the spinor bundle on \mathcal{Y} . Hence we get the decomposition

$$\text{S-Thom}_K(K \times_T \mathcal{Y}) = \text{Thom}_K(K/T) \odot K \times_T \text{S-Thom}_T(\mathcal{Y}) .$$

The transversally elliptic symbol $\text{Thom}_K^{\Phi_{\tilde{\mu}}}(\mathcal{U})$ is equal to

$$\left[\text{Thom}_K(K/T) \odot \text{Thom}_K(\overline{K/T}) \odot K \times_T \text{S-Thom}_T(\mathcal{Y}) \right]_{\text{deformed by } \mathcal{H}^{\tilde{\mu}}} ,$$

hence $\sigma_1 := (\phi^{-1})^* \text{Thom}_K^{\Phi_{\tilde{\mu}}}(\mathcal{U})$ is equal to

$$\left[\text{Thom}_K(K/T) \odot K \times_T \left(\text{Thom}_T(\overline{K/T}) \odot \text{S-Thom}_T(\mathcal{Y}) \right) \right]_{\text{deformed by } \mathcal{H}'},$$

where $\mathcal{H}' = \phi_*(\mathcal{H}_{\tilde{\mu}})$.

Using the decomposition $\mathbf{T}\mathcal{U}' \simeq K \times_T (\mathfrak{k}/\mathfrak{t} \oplus K \times_T (\overline{\mathfrak{k}/\mathfrak{t}} \oplus \mathbf{T}\mathcal{Y}))$, a small computation gives $\mathcal{H}'(m) = pr_{\mathfrak{k}/\mathfrak{t}}(h\tilde{\mu}) + R(m) + \mathcal{H}_{\tilde{\mu}}(y) + S(m)$ for $m = [k; [h], y] \in \mathcal{U}'$, where⁷ $R(m) \in \overline{\mathfrak{k}/\mathfrak{t}}$ and $S(m) \in \mathbf{T}_y\mathcal{Y}$ vanishes when $m \in K \times_T (\{\bar{e}\} \times \mathcal{Y})$, i.e. when $[h] = \bar{e}$. Here $\mathcal{H}_{\tilde{\mu}}$ is the Hamiltonian vector field of the function $\frac{-1}{2} \|\Phi - \tilde{\mu}\|^2: \mathcal{Y} \rightarrow \mathbb{R}$.

The transversally elliptic symbol σ_1 is equal to the exterior product

$$\sigma_1(m, \xi_1 + \xi_2 + v) = \mathbf{c}(\xi_1 - pr_{\mathfrak{k}/\mathfrak{t}}(h\tilde{\mu})) \odot \mathbf{c}(\xi_2 - R(m)) \odot \mathbf{c}(v - \mathcal{H}_{\tilde{\mu}} - S(m)),$$

with $\xi_1 \in \mathfrak{k}/\mathfrak{t}$, $\xi_2 \in \overline{\mathfrak{k}/\mathfrak{t}}$, and $v \in \mathbf{T}\mathcal{Y}$.

Now, we simplify the symbol σ_1 without changing its K -theoretic class. Since $\text{Char}(\sigma_1) \cap \mathbf{T}_K\mathcal{U}' = K \times_T (\{\bar{e}\} \times \mathcal{Y})$, we transform σ_1 through the K -invariant diffeomorphism $h = e^X$ from a neighborhood of 0 in $\mathfrak{k}/\mathfrak{t}$ to a neighborhood of \bar{e} in $\overline{K/T}$. That gives $\sigma_2 \in K_K(\mathbf{T}_K(K \times_T (\mathfrak{k}/\mathfrak{t} \times \mathcal{Y})))$ defined by

$$\begin{aligned} \sigma_2([k; X, y], \xi_1 + \xi_2 + v) = \\ \mathbf{c}(\xi_1 - pr_{\mathfrak{k}/\mathfrak{t}}(e^X \tilde{\mu})) \odot \mathbf{c}(\xi_2 - R(m)) \odot \mathbf{c}(v - \mathcal{H}_{\tilde{\mu}} - S(m)). \end{aligned}$$

Now trivial homotopies link σ_2 with the symbol

$$\sigma_3([k; X, y], \xi_1 + \xi_2 + v) = \mathbf{c}(\xi_1 - [X, \tilde{\mu}]) \odot \mathbf{c}(\xi_2) \odot \mathbf{c}(v - \mathcal{H}_{\tilde{\mu}}),$$

where we have removed the terms $R(m)$ and $S(m)$, and where we have replaced $pr_{\mathfrak{k}/\mathfrak{t}}(e^X \tilde{\mu}) = [X, \tilde{\mu}] + o([X, \tilde{\mu}])$ by the term $[X, \tilde{\mu}]$. Now we see that $\sigma_3 = i_*(\sigma_4)$ where the symbol $\sigma_4 \in K_T(\mathbf{T}_T(\mathfrak{k}/\mathfrak{t} \times \mathcal{Y}))$ is defined by

$$\sigma_4(X, y; \xi_2 + v) = \mathbf{c}(-[X, \tilde{\mu}]) \odot \mathbf{c}(\xi_2) \odot \mathbf{c}(v - \mathcal{H}_{\tilde{\mu}}).$$

So σ_4 is equal to the exterior product of $(y, v) \rightarrow \mathbf{c}(v - \mathcal{H}_{\tilde{\mu}})$, which is $\text{S-Thom}_T^{\Phi - \tilde{\mu}}(\mathcal{Y})$, with the transversally elliptic symbol on $\mathfrak{k}/\mathfrak{t}$: $(X, \xi_2) \rightarrow \mathbf{c}(-[X, \tilde{\mu}]) \odot \mathbf{c}(\xi_2)$. But the K -theoretic class of this former symbol is equal to $k_!(\mathbb{C})$, where $k: \{0\} \hookrightarrow \mathfrak{k}/\mathfrak{t}$ (see subsection 5.1 in [33]). This shows that

$$\sigma_4 = k_!(\mathbb{C}) \odot \text{S-Thom}_T^{\Phi - \tilde{\mu}}(\mathcal{Y}) = j_!(\text{S-Thom}_T^{\Phi - \tilde{\mu}}(\mathcal{Y})).$$

□

4.5. Proof of Proposition 4.14. In this subsection, $\tilde{\mu} = \mu + \rho_c$ is fixed, and is assumed to belong to Δ . The induced moment map on the symplectic slice $\Phi^{-1}(\text{Interior}(\mathfrak{k}_+^*))$ is still denoted by Φ . Let $r > 0$ be the smallest non zero critical value of $\|\Phi - \tilde{\mu}\|$, and let $\mathcal{Y} = \Phi^{-1}\{\xi \in \text{Aff}(\Delta), \|\xi - \tilde{\mu}\| < \frac{r}{2}\}$.

For $\xi \in \text{Aff}(\Delta)$, we consider $\xi_t = t\xi + (1-t)\tilde{\mu}$, $0 \leq t \leq 1$. If one shows that there exists a compact subset $\mathcal{K} \subset \mathcal{Y}$ such that $\text{Cr}(\|\Phi - \xi_t\|^2) \cap \mathcal{Y} \subset \mathcal{K}$, the family $\text{S-Thom}_T^{\Phi - \xi_t}(\mathcal{Y})$, $0 \leq t \leq 1$, defines then a homotopy of transversally elliptic symbols between $\text{S-Thom}_T^{\Phi - \tilde{\mu}}(\mathcal{Y})$ and $\text{S-Thom}_T^{\Phi - \xi}(\mathcal{Y})$. It shows that $\mathcal{Q}_{\Phi - \tilde{\mu}}^T(\mathcal{Y}, -)$ and $\mathcal{Q}_{\Phi - \xi}^T(\mathcal{Y}, -)$ are equal.

We describe now $\text{Cr}(\|\Phi - \xi_t\|^2) \cap \mathcal{Y}$ using a parametrization introduced in [30][Section 6]. Let \mathcal{B} be the collection of affine subspaces of \mathfrak{k}^* generated by the

⁷A small computation shows that $R(m) = pr_{\mathfrak{k}/\mathfrak{t}}(h^{-1}(pr_{\mathfrak{k}}(h\tilde{\mu}) - \Phi(y)))$, and $S(m) = [\tilde{\mu} - pr_{\mathfrak{k}}(h\tilde{\mu})]_{\mathcal{Y}}(y)$.

image under Φ of submanifolds \mathcal{Z} of the following type: \mathcal{Z} is a connected component of \mathcal{Y}^H which intersects $\Phi^{-1}(\tilde{\mu})$, H being a subgroup of T . The set \mathcal{B} is finite since $\Phi^{-1}(\tilde{\mu})$ is compact and thus has a finite number of stabilizers for the T action. Note that \mathcal{B} is reduced to $\text{Aff}(\Delta)$ if $\tilde{\mu}$ is a generic quasi regular value of Φ . For $A \in \mathcal{B}$, we denote by $\beta(-, A)$ the orthogonal projection on A . Let $\mathcal{B}_\xi = \{\beta(\xi, A) - \xi \mid A \in \mathcal{B}\}$. Like in [31][Section 4.3], we see that

$$(4.42) \quad \text{Cr}(\|\Phi - \xi\|^2) \cap \mathcal{Y} = \bigcup_{\beta \in \mathcal{B}_\xi} (\mathcal{Y}^\beta \cap \Phi^{-1}(\beta + \xi))$$

if $\|\xi - \tilde{\mu}\| < \frac{r}{2}$. If we take $\mathcal{K} := \Phi^{-1}\{\xi \in \text{Aff}(\Delta), \|\xi - \tilde{\mu}\| \leq \frac{r}{3}\}$, we have $\text{Cr}(\|\Phi - \xi\|^2) \cap \mathcal{Y} \subset \mathcal{K}$ for $\|\xi - \tilde{\mu}\| \leq \frac{r}{3}$. Thus point *i*) of Proposition 4.14 is proved.

Now we fix $\xi \in \text{Aff}(\Delta)$ close enough to $\tilde{\mu}$. And for each $\beta \in \mathcal{B}_\xi$, we denote by $\mathcal{Q}_\beta^T(\mathcal{Y}, -)$ the map localized near $\mathcal{Y}^\beta \cap \Phi^{-1}(\beta + \xi)$. The excision property tells us, like in (3.14), that

$$\mathcal{Q}_{\Phi-\xi}^T(\mathcal{Y}, -) = \sum_{\beta \in \mathcal{B}_\xi} \mathcal{Q}_\beta^T(\mathcal{Y}, -).$$

Note that $0 \in \mathcal{B}_\xi$ if and only if $\Phi^{-1}(\xi) \neq \emptyset$. Point iii) of Proposition 4.14 will follow from the following

Lemma 4.16. *Let $\xi \in \text{Aff}(\Delta)$ close enough to $\tilde{\mu}$, and let β be a non-zero element of \mathcal{B}_ξ . Then $[\mathcal{Q}_\beta^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C})]^T = 0$. Hence $[\mathcal{Q}_{\Phi-\xi}^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu})]^T = 0$, if $\xi \notin \Delta$.*

For the point *ii*) of Proposition 4.14, we also need the

Lemma 4.17. *If $\xi \in \Delta^\circ$ is a generic quasi regular value of Φ , we have $[\mathcal{Q}_0^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu})]^T = \mathcal{Q}(M_\xi, o(M_\xi))$.*

Other versions of Lemmas 4.17 and 4.16 are already known : in the Spin-case for an S^1 -action by Vergne [42], and by the author [33] when the Spin^c-structure comes from an almost complex structure.

We review briefly the arguments, since they work in the same way. We consider the Spin^c structure on \mathcal{Y} defined in Proposition 4.10, that we twist by the line bundle $\tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu}$: it defines a Spin^c structure \mathcal{Q}^μ on \mathcal{Y} with canonical line bundle $\mathbb{L}^\mu := L_{2\omega} \otimes \mathbb{C}_{-2\tilde{\mu}}$. We consider then the symbol S-Thom $_{T,\mu}^{\Phi-\xi}(\mathcal{Y})$ constructed with \mathcal{Q}^μ (see Definition 4.7). For $\beta \in \mathcal{B}_\xi$, the term $\mathcal{Q}_\beta^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu})$ is by definition the T -index of S-Thom $_{T,\mu}^{\Phi-\xi}(\mathcal{Y})|_{\mathcal{U}^\beta}$, where \mathcal{U}^β is a sufficiently small open neighborhood of $\mathcal{Y}^\beta \cap \Phi^{-1}(\beta + \xi)$ in \mathcal{Y} .

Proof of Lemma 4.17. A neighborhood \mathcal{U}^0 of $\mathcal{Z} := \Phi^{-1}(\xi)$ is diffeomorphic to a neighborhood of \mathcal{Z} in $\mathcal{Z} \times \overrightarrow{\Delta}$, where $\Phi - \xi : \mathcal{Z} \times \overrightarrow{\Delta} \rightarrow \overrightarrow{\Delta}$ is the projection to the second factor. Let $pr : \mathcal{Z} \times \overrightarrow{\Delta} \rightarrow \mathcal{Z}$ be the projection to the first factor. We still denote by \mathcal{Q}^μ the Spin^c-structure on $\mathcal{Z} \times \overrightarrow{\Delta}$ equal to $pr^*(\mathcal{Q}^\mu|_{\mathcal{Z}})$. We easily show that $\mathcal{Q}_\beta^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu})$ is equal to the T -index of $\sigma_{\mathcal{Z}} = \text{S-Thom}_{T,\mu}^{\Phi-\xi}(\mathcal{Z} \times \overrightarrow{\Delta})$. Let \mathcal{Q}'' be the reduction of $\mathcal{Q}^\mu|_{\mathcal{Z}}$ introduced in (4.36). Since $\mathcal{Q}^\mu|_{\mathcal{Z}} = \text{Spin}_{2k}^c \times_{(\text{Spin}_{2k'}^c \times \text{U}_{2l'})} \mathcal{Q}''$, the bundle of spinors \mathcal{S} over $\mathcal{Z} \times \overrightarrow{\Delta}$ decomposes as

$$\mathcal{S} = pr^*\left(\pi^*\mathcal{S}(M_\xi) \otimes \mathcal{Z} \times \wedge(\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C})\right).$$

Here $\mathcal{S}(M_\xi)$ is the bundle of spinors on M_ξ induces by the Spin^c -structure $Q''/U_{2\nu}$, and $\pi : \mathcal{Z} \rightarrow M_\xi$ is the quotient map. Inside the trivial bundle $\mathcal{Z} \times (\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C})$, we have identified $\mathcal{Z} \times (\mathfrak{t}/\mathfrak{t}_\Delta \otimes i\mathbb{R})$ with the subspace of $\mathbf{T}\mathcal{Z}$ formed by the vector fields generated by the infinitesimal action of $\mathfrak{t}/\mathfrak{t}_\Delta$, and $\mathcal{Z} \times (\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{R})$ with $\mathcal{Z} \times \vec{\Delta} \subset \mathbf{T}(\mathcal{Z} \times \vec{\Delta})|_{\mathcal{Z}}$. For $(z, f) \in \mathcal{Z} \times \vec{\Delta}$, let us decompose $v \in \mathbf{T}_{(z,f)}(\mathcal{Z} \times \vec{\Delta})$ as $v = v_1 + X + iY$, where $v_1 \in \pi^*(\mathbf{T}M_\xi)$, and $X + iY \in \mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C}$. The map $\sigma_{\mathcal{Z}}(z, f; v_1 + X + iY)$ acts on $\mathcal{S}(M_\xi)_z \otimes \wedge(\mathfrak{t}/\mathfrak{t}_\Delta \otimes \mathbb{C})$ as the product

$$\mathbf{c}_z(v_1) \odot \mathbf{c}(X + i(Y - f)) ,$$

which is homotopic⁸ to the transversally elliptic symbol

$$\mathbf{c}_z(v_1) \odot \mathbf{c}(f + iX) .$$

So we have proved that $\sigma_{\mathcal{Z}} = j_! \circ \pi^*(\text{S-Thom}(M_\xi))$, where $j_! : K_T(\mathbf{T}_T\mathcal{Z}) \rightarrow K_T(\mathbf{T}_T(\mathcal{Z} \times \vec{\Delta}))$ is induced by the inclusion $j : \mathcal{Z} \hookrightarrow \mathcal{Z} \times \vec{\Delta}$. The last equality finishes the proof (see [33][Section 6.1]).

Proof of Lemma 4.16. The equality $[\mathcal{Q}_\beta^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu})]^T = 0$ comes from a localization formula on the submanifold \mathcal{Y}^β for the map $\mathcal{Q}_\beta^T(\mathcal{Y}, -)$ (see [33, 42]). The normal bundle \mathcal{N} of \mathcal{Y}^β in \mathcal{Y} carries a complex structure $J_{\mathcal{N}}$ on the fibers such that each \mathbb{T}_β -weight α on $(\mathcal{N}, J_{\mathcal{N}})$ satisfies $(\alpha, \beta) > 0$. The principal bundle Q^μ , when restricted to \mathcal{Y}^β admits the reduction

$$(4.43) \quad \begin{array}{ccc} Q' & \longrightarrow & \text{P}_{\text{SO}}(\mathbf{T}\mathcal{Y}^\beta) \times \text{P}_{\text{U}}(\mathcal{N}) \times \text{P}_{\text{U}}(\mathbb{L}|_{\mathcal{Y}^\beta}) \\ \downarrow & & \downarrow \\ Q^\mu|_{\mathcal{Y}^\beta} & \longrightarrow & \text{P}_{\text{SO}}(\mathbf{T}\mathcal{Y}^\beta \oplus \mathcal{N}) \times \text{P}_{\text{U}}(\mathbb{L}^\mu|_{\mathcal{Y}^\beta}) , \end{array}$$

Hence $Q^\beta := Q'/U(l)$ is a Spin^c -structure on \mathcal{Y}^β with canonical line bundle equal to $\mathbb{L}^\beta := \mathbb{L}^\mu|_{\mathcal{Y}^\beta} \otimes (\det \mathcal{N})^{-1} = L_{2\omega}|_{\mathcal{Y}^\beta} \otimes \mathbb{C}_{-2\tilde{\mu}} \otimes (\det \mathcal{N})^{-1}$. Let $\mathcal{Q}_\beta^T(\mathcal{Y}^\beta, -)$ be the map defined by Q^β and localized near $\Phi^{-1}(\beta + \xi) \cap \mathcal{Y}^\beta$ by $\Phi - \xi$. Following the argument of [33][Section 6] one obtains

$$\mathcal{Q}_\beta^T(\mathcal{Y}, \tilde{L}|_{\mathcal{Y}} \otimes \mathbb{C}_{-\mu}) = (-1)^l \sum_{k \in \mathbb{N}} \mathcal{Q}_\beta^T(\mathcal{Y}^\beta, \det \mathcal{N} \otimes S^k(\mathcal{N})) ,$$

where $S^k(\mathcal{N})$ is the k -th symmetric product of \mathcal{N} , and $l = \text{rank}_{\mathbb{C}} \mathcal{N}$. Thus, it is sufficient to prove that $[\mathcal{Q}_\beta^T(\mathcal{Y}^\beta, \det \mathcal{N} \otimes S^k(\mathcal{N}))]^T = 0$ for every $k \in \mathbb{N}$. For this purpose, we use Lemma 4.8. Let α be the \mathbb{T}_β -weight on $\det \mathcal{N}$. From the Kostant formula (2.1), the \mathbb{T}_β -weight on $L_{2\omega}|_{\mathcal{Y}^\beta}$ is equal to $2(\beta + \xi)$. Hence any \mathbb{T}_β -weight γ on $\det \mathcal{N} \otimes S^k(\mathcal{N}) \otimes (\mathbb{L}^\beta)^{1/2}$ is of the form

$$\gamma = \beta + \xi - \tilde{\mu} + \frac{1}{2}\alpha + \delta$$

where δ is a \mathbb{T}_β -weight on $S^k(\mathcal{N})$. So $(\gamma, \beta) = (\beta + \xi - \tilde{\mu}, \beta) + \frac{1}{2}(\alpha, \beta) + (\delta, \beta)$. But the \mathbb{T}_β -weights on \mathcal{N} are ‘positive’ for β , so $(\alpha, \beta) > 0$ and $(\delta, \beta) \geq 0$. On the other hand, $\beta + \xi = \beta(\xi, A)$ is the orthogonal projection of ξ on some affine subspace $A \subset \mathfrak{t}_+^*$ which contains $\tilde{\mu}$: hence $(\beta + \xi - \tilde{\mu}, \beta) = 0$. This proves that $(\gamma, \beta) > 0$. \square

⁸See [33][Section 6.1].

5. QUANTIZATION AND THE DISCRETE SERIES

In this section we apply Theorem 4.1 to the coadjoint orbits that parametrize the discrete series of a real, connected, semi-simple Lie group G , with finite center. Nice references on the subject of ‘the discrete series’ are [36, 12].

Let K be a maximal compact subgroup of G , and T be a maximal torus in K . For the remainder of this section, we assume that T is a Cartan subgroup of G . The discrete series of G is then non-empty and is parametrized by a subset \widehat{G}_d in the dual \mathfrak{t}^* of the Lie algebra of T [19].

Let us fix some notation. Let $\mathfrak{R}_c \subset \mathfrak{R} \subset \Lambda^*$ be respectively the set of (real) roots for the action of T on $\mathfrak{k} \otimes \mathbb{C}$ and $\mathfrak{g} \otimes \mathbb{C}$. We choose a system of positive roots \mathfrak{R}_c^+ for \mathfrak{R}_c , we denote by \mathfrak{t}_+^* the corresponding Weyl chamber, and we let ρ_c be half the sum of the elements of \mathfrak{R}_c^+ . We denote by B the Killing form on \mathfrak{g} . It induces a scalar product (denoted by $(-, -)$) on \mathfrak{t} , and then on \mathfrak{t}^* . An element $\lambda \in \mathfrak{t}^*$ is called *regular* if $(\lambda, \alpha) \neq 0$ for every $\alpha \in \mathfrak{R}$, or equivalently, if the stabilizer subgroup of λ in G is T . Given a system of positive roots \mathfrak{R}^+ for \mathfrak{R} , consider the subset $\Lambda^* + \frac{1}{2} \sum_{\alpha \in \mathfrak{R}^+} \alpha$ of \mathfrak{t}^* . This does not depend on the choice of \mathfrak{R}^+ , and we denote it by Λ_ρ^* [12].

The discrete series of G are parametrized by

$$(5.44) \quad \widehat{G}_d := \{ \lambda \in \mathfrak{t}^*, \lambda \text{ regular} \} \cap \Lambda_\rho^* \cap \mathfrak{t}_+^* .$$

When $G = K$ is compact, the set \widehat{G}_d equals $\Lambda_+^* + \rho_c$, and it parametrizes the set of irreducible representations of K . Harish-Chandra has associated to any $\lambda \in \widehat{G}_d$ an invariant eigendistribution on G , denoted by Θ_λ , which is shown to be the global trace of an irreducible, square integrable, unitary representation of G .

On the other hand we associate to $\lambda \in \widehat{G}_d$, the regular coadjoint orbit $M := G \cdot \lambda$. It is a symplectic manifold with a Hamiltonian action of K . Since the vectors $X_M, X \in \mathfrak{g}$, span the tangent space at every $\xi \in M$, the symplectic 2-form is determined by

$$\omega(X_M, Y_M)_\xi = \langle \xi, [X, Y] \rangle .$$

The corresponding moment map $\Phi : M \rightarrow \mathfrak{k}^*$ for the K -action is the composition of the inclusion $i : M \hookrightarrow \mathfrak{g}^*$ with the projection $\mathfrak{g}^* \rightarrow \mathfrak{k}^*$. The vector λ determines a choice $\mathfrak{R}^{+, \lambda}$ of positive roots for the T -action on $\mathfrak{g} \otimes \mathbb{C} : \alpha \in \mathfrak{R}^{+, \lambda} \iff (\alpha, \lambda) > 0$. We recall now how the choice of $\mathfrak{R}^{+, \lambda}$ determines a complex structure on M . First take the decomposition $\mathfrak{g} \otimes \mathbb{C} = \mathfrak{t} \otimes \mathbb{C} \oplus \sum_{\alpha \in \mathfrak{R}} \mathfrak{g}_\alpha$ where $\mathfrak{g}_\alpha := \{ v \in \mathfrak{g} \otimes \mathbb{C} \mid \exp(X) \cdot v = e^{i\langle \alpha, X \rangle} v \text{ for any } X \in \mathfrak{t} \}$. It gives the following T -equivariant decomposition of the complexified tangent space of M at λ :

$$\mathbf{T}_\lambda M \otimes \mathbb{C} = \sum_{\alpha \in \mathfrak{R}} \mathfrak{g}_\alpha = \mathfrak{n} \oplus \bar{\mathfrak{n}} ,$$

with $\mathfrak{n} = \sum_{\alpha \in \mathfrak{R}^{+, \lambda}} \mathfrak{g}_\alpha$. We have then a T -equivariant isomorphism $\mathcal{I} : \mathbf{T}_\lambda M \rightarrow \mathfrak{n}$ equal to the composition of the inclusion $\mathbf{T}_\lambda M \hookrightarrow \mathbf{T}_\lambda M \otimes \mathbb{C}$ with the projection $\mathfrak{n} \oplus \bar{\mathfrak{n}} \rightarrow \mathfrak{n}$. The T -equivariant complex structure J_λ on $\mathbf{T}_\lambda M$ is determined by the relation $\mathcal{I}(J_\lambda v) = i\mathcal{I}(v)$. Hence, the set of real infinitesimal weights for the T -action on $(\mathbf{T}_\lambda M, J_\lambda)$ is $\mathfrak{R}^{+, \lambda}$. Since M is a homogeneous space, J_λ defines an invariant almost complex structure J on M , which is in fact integrable. Through the isomorphism $M \cong G/T$, the canonical line bundle $\kappa = \det_{\mathbb{C}}(\mathbf{T}M)^{-1}$ is equal to $\kappa = G \times_T \mathbb{C}_{-2\rho}$ with $\rho = \frac{1}{2} \sum_{\alpha \in \mathfrak{R}^{+, \lambda}} \alpha$.

If $\lambda \in \widehat{G}_d$, then $\lambda - \rho$ is a weight, and

$$(5.45) \quad \tilde{L} := G \times_T \mathbb{C}_{\lambda - \rho} \rightarrow G/T$$

is a κ -prequantum line bundle over (M, ω, J) . We have shown in [32], that $\text{Cr}(\|\Phi\|^2)$ is compact, equal to the K -orbit $K \cdot \lambda$. Then the generalized Riemann-Roch character $RR_\Phi^K(M, -)$ is well defined (see Definition 3.2). The main result of this section is the following

Theorem 5.1. *We have the following equality of tempered distributions on K*

$$\Theta_\lambda|_K = (-1)^{\frac{\dim(G/K)}{2}} RR_\Phi^K(G \cdot \lambda, \tilde{L}) ,$$

where $\Theta_\lambda|_K$ is the restriction of the eigendistribution Θ_λ to the subgroup K .

The proof of Theorem 5.1 is given in Subsection 5.2. It uses the Blattner formulas in an essential way (see Subsection 5.1).

With Theorem 5.1 at our disposal we can exploit the result of Theorem 4.1 to compute the K -multiplicities, $m_\mu(\lambda) \in \mathbb{N}$, of $\Theta_\lambda|_K$ in term of the reduced spaces. By definition we have

$$(5.46) \quad \Theta_\lambda|_K = \sum_{\mu \in \Lambda_+^*} m_\mu(\lambda) \chi_\mu^K \quad \text{in } R^{-\infty}(K) .$$

The moment map $\Phi : G \cdot \lambda \rightarrow \mathfrak{k}^*$ is *proper* since the $G \cdot \lambda$ is closed in \mathfrak{g}^* [32]. We show (Lemma 5.5) that the moment polyhedron $\Delta = \Phi(G \cdot \lambda) \cap \mathfrak{t}_+^*$ is of dimension $\dim T$. Thus on the relative interior Δ° of the moment polyhedron, the notions of *generic quasi-regular values* and *regular values* coincide : they concern the elements $\xi \in \Delta^\circ$ such that $\Phi^{-1}(\xi)$ is a smooth submanifold with a locally free action of T . We have shown (Subsection 4.3) how to define the quantity $Q((G \cdot \lambda)_{\mu + \rho_c}) \in \mathbb{Z}$ as the index of a suitable Spin^c Dirac operator on $\Phi^{-1}(\xi)/T$, where $\xi \in \Delta^\circ$ is a regular value of Φ close enough to $\mu + \rho_c$.

Proposition 5.2. *For every $\mu \in \Lambda_+^*$, we have*

$$m_\mu(\lambda) = \mathcal{Q}((G \cdot \lambda)_{\mu + \rho_c}) .$$

In particular $m_\mu(\lambda) = 0$ if $\mu + \rho_c$ does not belong to the relative interior of the moment polyhedron Δ .

Proof. A small check of orientations shows that $\varepsilon = (-1)^{\frac{\dim(G/K)}{2}}$, thus this proposition follows from Theorems 4.1 and 5.1 if one checks that the following holds: $(G \cdot \lambda, \Phi)$ satisfies Assumption 3.6, and the infinitesimal K -stabilizers are Abelian. The first point will be handled in Subsection 5.3. The second point is obvious since $M \cong G/T$: all the G -stabilizers are conjugate to T , so all the K -stabilizers are Abelian. \square

5.1. Blattner formulas. In this section, we fix $\lambda \in \widehat{G}_d$. Let $\mathfrak{R}^{+, \lambda}$ be the system of positive roots defined $\lambda: \alpha \in \mathfrak{R}^{+, \lambda} \iff (\alpha, \lambda) > 0$. Then $\mathfrak{R}_c^+ \subset \mathfrak{R}^{+, \lambda}$, and $\rho = \frac{1}{2} \sum_{\alpha \in \mathfrak{R}^{+, \lambda}} \alpha$ decomposes in $\rho = \rho_c + \rho_n$ where $\rho_n = \frac{1}{2} \sum_{\alpha \in \mathfrak{R}_n^+} \alpha$ and $\mathfrak{R}_n^+ = \mathfrak{R}^{+, \lambda} - \mathfrak{R}_c^+$.

Let $\mathcal{P}: \Lambda^* \mapsto \mathbb{N}$ be the partition function associated to the set \mathfrak{R}_n^+ : for $\mu \in \Lambda^*$, $\mathcal{P}(\mu)$ is the number of distinct ways we can write $\mu = \sum_{\alpha \in \mathfrak{R}_n^+} k_\alpha \alpha$ with $k_\alpha \in \mathbb{N}$ for all α . The following Theorem is known as the Blattner formulas and was first proved by Hecht and Schmid [20].

Theorem 5.3. *For $\mu \in \Lambda_+^*$, we have*

$$m_\mu(\lambda) = \sum_{w \in W} (-1)^w \mathcal{P}(w(\mu + \rho_c) - (\mu_\lambda + \rho_c)) ,$$

where⁹ $\mu_\lambda := \lambda - \rho_c + \rho_n$. Here W is the Weyl group of (K, T) .

Using Theorem 5.3, we can describe $\Theta_\lambda|_K$ through the holomorphic induction map $\text{Hol}_T^K: R^{-\infty}(T) \rightarrow R^{-\infty}(K)$. Recall that Hol_T^K is characterized by the following properties: i) $\text{Hol}_T^K(t^\mu) = \chi_\mu^K$ for every dominant weight $\mu \in \Lambda_+^*$; ii) $\text{Hol}_T^K(t^{w \circ \mu}) = (-1)^w \text{Hol}_T^K(t^\mu)$ for every $w \in W$ and $\mu \in \Lambda^*$; iii) $\text{Hol}_T^K(t^\mu) = 0$ if $W \circ \mu \cap \Lambda_+^* = \emptyset$. Using these properties we have

$$(5.47) \quad \sum_{\mu \in \Lambda^*} \mathcal{R}(\mu) \text{Hol}_T^K(t^\mu) = \sum_{\mu \in \Lambda_+^*} \left[\sum_{w \in W} (-1)^w \mathcal{R}(w \circ \mu) \right] \chi_\mu^K ,$$

for every map $\mathcal{R}: \Lambda^* \rightarrow \mathbb{Z}$.

For a weight $\alpha \in \Lambda^*$, with $(\lambda, \alpha) \neq 0$, let us denote the oriented inverse of $(1 - t^\alpha)$ in the following way

$$[1 - t^\alpha]_\lambda^{-1} = \begin{cases} \sum_{k \in \mathbb{N}} t^{k\alpha} , & \text{if } (\lambda, \alpha) > 0 \\ -t^{-\alpha} \sum_{k \in \mathbb{N}} t^{-k\alpha} , & \text{if } (\lambda, \alpha) < 0 . \end{cases}$$

Let $A = \{\alpha_1, \dots, \alpha_l\}$ be a set of weights with $(\lambda, \alpha_i) \neq 0, \forall i$. We denote by $A^+ = \{\varepsilon_1 \alpha_1, \dots, \varepsilon_l \alpha_l\}$ the corresponding set of polarized weights: $\varepsilon_i = \pm 1$ and $(\lambda, \varepsilon_i \alpha_i) > 0$ for all i . The product $\prod_{\alpha \in A} [1 - t^\alpha]_\lambda^{-1}$ is well defined in $R^{-\infty}(T)$, and is denoted by $[\prod_{\alpha \in A} (1 - t^\alpha)]_\lambda^{-1}$. A small computation shows that

$$(5.48) \quad \begin{aligned} \left[\prod_{\alpha \in A} (1 - t^\alpha) \right]_\lambda^{-1} &= (-1)^r t^{-\gamma} \left[\prod_{\alpha \in A^+} (1 - t^\alpha) \right]_\lambda^{-1} \\ &= (-1)^r t^{-\gamma} \sum_{\mu \in \Lambda^*} \mathcal{P}_{A^+}(\mu) t^\mu . \end{aligned}$$

Here $\mathcal{P}_{A^+}: \Lambda^* \mapsto \mathbb{N}$ is the partition function associated to A^+ , $\gamma = \sum_{(\lambda, \alpha) < 0} \alpha$, and $r = \#\{\alpha \in A, (\lambda, \alpha) < 0\}$. This notation is compatible with the one we used in [33][Section 5]. If V is a complex T -vector space where the subspace fixed by λ is reduced to $\{0\}$, then $\wedge_{\mathbb{C}}^\bullet V \in R(T)$ admits a polarized inverse $[\wedge_{\mathbb{C}}^\bullet V]_\lambda^{-1} = [\prod_{\alpha \in \mathfrak{R}(V)} (1 - t^\alpha)]_\lambda^{-1}$, where $\mathfrak{R}(V)$ is the set of real infinitesimal T -weights on V .

⁹We shall note that $\mu_\lambda \in \Lambda_+^*$ (see [12], section 5).

Lemma 5.4. *We have the following equality in $R^{-\infty}(K)$*

$$\Theta_\lambda|_K = \text{Hol}_T^K \left(t^{\mu_\lambda} \left[\Pi_{\alpha \in \mathfrak{R}_n^+} (1 - t^\alpha) \right]_\lambda^{-1} \right).$$

Proof. Let $\Theta \in R^{-\infty}(K)$ be the RHS in the equality of the Lemma. From (5.48), we have $\Theta = \sum_{\mu \in \Lambda^*} \mathcal{P}(\mu) \text{Hol}_T^K(t^{\mu+\mu_\lambda}) = \sum_{\mu \in \Lambda^*} \mathcal{P}(\mu - \mu_\lambda) \text{Hol}_T^K(t^\mu)$. If we use now (5.47), we see that multiplicity of Θ relative to the highest weight $\mu \in \Lambda_+^*$ is $\sum_{w \in W} (-1)^w \mathcal{P}(w(\mu + \rho_c) - (\mu_\lambda + \rho_c))$. From Theorem 5.3 we conclude that $\Theta_\lambda|_K = \Theta$. \square

5.2. Proof of Theorem 5.1. In Lemma 5.4 we have used the Blattner formulas to write $\Theta_\lambda|_K$ in term of the holomorphic induction map Hol_T^K . Theorem 5.1 is then proved if one shows that $RR_\Phi^K(G \cdot \lambda, \tilde{L}) = (-1)^r \text{Hol}_T^K(t^{\mu_\lambda} [\Pi_{\alpha \in \mathfrak{R}_n^+} (1 - t^\alpha)]_\lambda^{-1})$, with $\mu_\lambda = \lambda - \rho_c + \rho_n$, and $r = \frac{1}{2} \dim(G/K)$. More generally, we show in this section that for any K -equivariant vector bundle $V \rightarrow G \cdot \lambda$

$$(5.49) \quad RR_\Phi^K(G \cdot \lambda, V) = (-1)^r \text{Hol}_T^K \left(V_\lambda \cdot t^{2\rho_n} \cdot \left[\Pi_{\alpha \in \mathfrak{R}_n^+} (1 - t^\alpha) \right]_\lambda^{-1} \right),$$

where $V_\lambda \in R(T)$ is the fiber of V at λ .

First we recall why $\text{Cr}(\|\Phi\|^2) = K \cdot \lambda$ in $M := G \cdot \lambda$ (see [32] for the general case of closed coadjoint orbits). One can work with an adjoint orbit $M := G \cdot \tilde{\lambda}$ through the G -identification $\mathfrak{g}^* \simeq \mathfrak{g}$ given by the Killing form; then $\Phi : M \rightarrow \mathfrak{k}$ is just the restriction on M of the (orthogonal) projection $\mathfrak{g} \rightarrow \mathfrak{k}$. Let \mathfrak{p} be the orthogonal complement of \mathfrak{k} in \mathfrak{g} . Every $m \in M$ decomposes as $m = x_m + y_m$, with $x_m = \Phi(m)$ and $y_m \in \mathfrak{p}$. The Hamiltonian vector field of $\frac{-1}{2} \|\Phi\|^2$ is $\mathcal{H}_m = [x_m, m] = [x_m, y_m]$ (see 3.8). Thus

$$\text{Cr}(\|\Phi\|^2) = \{\mathcal{H} = 0\} = \{m \in M, [x_m, y_m] = 0\}.$$

Now, since $\tilde{\lambda}$ is elliptic, every $m \in M$ is also elliptic. If $m \in \text{Cr}(\|\Phi\|^2)$, $[m, x_m] = 0$ and m, x_m are elliptic, hence $y_m = m - x_m$ is elliptic and so is equal to 0. Finally $\text{Cr}(\|\Phi\|^2) = G \cdot \tilde{\lambda} \cap \mathfrak{k} = K \cdot \tilde{\lambda}$.

According to Definition 3.2, the computation of $RR_\Phi^K(G \cdot \lambda, \tilde{L})$ holds on a small K -invariant neighborhood of $K \cdot \lambda$ of $G \cdot \lambda$. Our model for the computation will be

$$\tilde{M} := K \times_T \mathfrak{p}$$

endowed with the canonical K -action. The tangent bundle $\mathbf{T}\tilde{M}$ is isomorphic to $K \times_T (\mathfrak{r} \oplus \mathbf{T}\mathfrak{p})$ where \mathfrak{r} is the T -invariant complement of \mathfrak{k} in \mathfrak{k} . One has a symplectic form $\tilde{\Omega}$ on \tilde{M} defined by $\tilde{\Omega}_m(V, V') = \langle \lambda, [X, X'] + [v, v'] \rangle$. Here $m = [k, x] \in K \times_T \mathfrak{p}$, and $V = [k, x; X + v]$, $X' = [k, x; X' + v']$ are two tangent vectors, with $X, X' \in \mathfrak{r}$ and $v, v' \in \mathfrak{p}$. A small computation shows that the K -action on $(K \times_T \mathfrak{p}, \tilde{\Omega})$ is Hamiltonian with moment map $\tilde{\Phi} : \tilde{M} \rightarrow \mathfrak{k}^*$ defined by

$$\tilde{\Phi}([k, x]) = k \cdot \left(\lambda - \frac{1}{2} pr_{\mathfrak{k}^*}(\lambda \circ \text{ad}(x) \circ \text{ad}(x)) \right).$$

Here $\text{ad}(x)$ is the adjoint action of x , and $pr_{\mathfrak{k}^*} : \mathfrak{g}^* \rightarrow \mathfrak{k}^*$ is the projection. Note first that the tangent space $\mathbf{T}_\lambda M$ and $\mathbf{T}_{[1,0]} \tilde{M}$ are canonically isomorphic to $\mathfrak{r} \oplus \mathfrak{p}$.

Lemma 5.5. *There exists a K -Hamiltonian isomorphism $\Upsilon : \mathcal{U} \simeq \tilde{\mathcal{U}}$, where \mathcal{U} is a K -invariant neighborhood of $K \cdot \lambda$ in M , $\tilde{\mathcal{U}}$ is a K -invariant neighborhood of K/T in \tilde{M} , and $\Upsilon(\lambda) = [1, 0]$. We impose furthermore that the differential of Υ at λ is the identity.*

Corollary 5.6. *The cone $\lambda + \sum_{\alpha \in \mathfrak{R}_n^+} \mathbb{R}^+ \alpha$ coincides with $\Delta = (G \cdot \lambda) \cap \mathfrak{t}_+^*$ in a neighborhood of λ . The polyhedral set Δ is of dimension $\dim T$.*

Proof of the Corollary. The first assertion is an immediate consequence of Lemma 5.5 and of the convexity Theorem [26]. Let $X_o \in \mathfrak{t}$ such that $\xi(X_o) = 0$ for all $\xi \in \vec{\Delta}$, that is $\alpha(X_o) = 0$ for all $\alpha \in \mathfrak{R}_n^+ : X_o$ commutes with all elements in \mathfrak{p} . Let \mathfrak{a} be a maximal Abelian subalgebra of \mathfrak{p} , and let Σ be the set of weights for the adjoint action of \mathfrak{a} on $\mathfrak{g} : \mathfrak{g} = \sum_{\alpha \in \Sigma} \mathfrak{g}_\alpha$, where $\mathfrak{g}_\alpha = \{Z \in \mathfrak{g}, [X, Z] = \alpha(X)Z \text{ for all } X \in \mathfrak{a}\}$. Since $[X_o, \mathfrak{a}] = 0$, we have $[X_o, \mathfrak{g}_\alpha] \subset \mathfrak{g}_\alpha$ for all $\alpha \in \Sigma$. But since $[X_o, \mathfrak{p}] = 0$ and $\mathfrak{g}_\alpha \cap \mathfrak{k} = 0$ for all $\alpha \neq 0$, we see that $[X_o, \mathfrak{g}_\alpha] = 0$ for all $\alpha \neq 0$. But X_o belongs to the Abelian subalgebra \mathfrak{g}_0 , so $[X_o, \mathfrak{g}_\alpha] = 0$. We see finally that X_o belongs to the center of \mathfrak{g} , and that implies $X_o = 0$ since G has a *finite center*. We have proved that $\vec{\Delta}^\perp = 0$, or equivalently $\vec{\Delta} = \mathfrak{t}$. \square

Proof of Lemma 5.5. The *symplectic cross-section Theorem* [17] asserts that the pre-image $\mathcal{Y} := \Phi^{-1}(\text{interior}(\mathfrak{t}_+^*))$ is a symplectic submanifold provided with a Hamiltonian action of T . The restriction $\Phi|_{\mathcal{Y}}$ is the moment map for the T -action on \mathcal{Y} . Moreover, the set $K \cdot \mathcal{Y}$ is a K -invariant neighborhood of $K \cdot \lambda$ in M diffeomorphic to $K \times_T \mathcal{Y}$. Since λ is a fixed T -point of \mathcal{Y} , a Hamiltonian model for $(\mathcal{Y}, \omega|_{\mathcal{Y}}, \Phi|_{\mathcal{Y}})$ in a neighborhood of λ is $(\mathbf{T}_\lambda \mathcal{Y}, \omega_\lambda, \Phi_\lambda)$ where ω_λ is the linear symplectic form of the tangent space $\mathbf{T}_\lambda M$ restricted to $\mathbf{T}_\lambda \mathcal{Y}$, and $\Phi_\lambda : \mathbf{T}_\lambda \mathcal{Y} \rightarrow \mathfrak{t}^*$ is the unique moment map with $\Phi_\lambda(0) = \lambda$. A small computation shows that $x \rightarrow \lambda \circ \text{ad}(x)$ is an isomorphism from \mathfrak{p} to $\mathbf{T}_\lambda \mathcal{Y}$, and $\Phi_\lambda(x) = \lambda - \frac{1}{2} \text{pr}_{\mathfrak{t}^*}(\lambda \circ \text{ad}(x) \circ \text{ad}(x))$. \square

We still denote the almost complex structure transported on $\tilde{\mathcal{U}} \subset K \times_T \mathfrak{p}$ through Υ by J . Since $d\Upsilon(\lambda)$ is the identity, $J_{[1,0]} : \mathfrak{t} \oplus \mathfrak{p} \rightarrow \mathfrak{t} \oplus \mathfrak{p}$ is equal to J_λ . Let $\pi : K \times_T \mathfrak{p} \rightarrow K/T$, and $\pi_{\tilde{\mathcal{U}}} : \tilde{\mathcal{U}} \rightarrow K/T$ be the fibering maps. Remark that for any equivariant vector bundle V over M the vector bundle $(\Upsilon^{-1})^*(V|_{\mathcal{U}}) \rightarrow \tilde{\mathcal{U}}$ is isomorphic to $\pi_{\tilde{\mathcal{U}}}^*(K \times_T V_\lambda)$. At this stage, we have according to Definition 3.2

$$(5.50) \quad RR_\Phi^K(G \cdot \lambda, V) = \text{Index}_{\tilde{\mathcal{U}}}^K \left(\text{Thom}_{\tilde{\mathcal{U}}}^{\tilde{\Phi}}(\tilde{\mathcal{U}}, J) \otimes \pi_{\tilde{\mathcal{U}}}^*(K \times_T V_\lambda) \right).$$

With the help of Lemma 3.4, we define now a simpler representative of the class defined by $\text{Thom}_{\tilde{\mathcal{U}}}^{\tilde{\Phi}}(\tilde{\mathcal{U}}, J)$ in $\mathbf{K}_K(\mathbf{T}_K \tilde{\mathcal{U}})$. Consider the map

$$\begin{aligned} \underline{\lambda} : K \times_T \mathfrak{p} &\longrightarrow \mathfrak{t}^* \\ (k, x) &\longmapsto k \cdot \lambda, \end{aligned}$$

and let $\underline{\lambda}_{\tilde{M}}$ be the vector field on \tilde{M} generated $\underline{\lambda}$ (see (3.8)). Note that $\underline{\lambda}_{\tilde{M}}$ never vanishes outside the zero section of $K \times_T \mathfrak{p}$. Let $(-, -)_{\tilde{M}}$ be the Riemannian metric on \tilde{M} defined by $(V, V')_{\tilde{M}} = (X, X') + (v, v')$ for $V = [k, x; X+v]$, $V' = [k, x; X'+v']$. A small computation shows that

$$(\tilde{\mathcal{H}}, \underline{\lambda}_{\tilde{M}})_{\tilde{M}} = \|\underline{\lambda}_{\tilde{M}}\|^2 + o(\|\underline{\lambda}_{\tilde{M}}\|^2)$$

in the neighborhood of the zero section in $K \times_T \mathfrak{p}$. Hence, if we take $\tilde{\mathcal{U}}$ small enough, $(\tilde{\mathcal{H}}, \underline{\Delta}_{\tilde{M}})_{\tilde{M}} > 0$ on $\tilde{\mathcal{U}} - \{\text{zero section}\}$, hence $\text{Thom}_{\tilde{K}}^{\tilde{\Phi}}(\tilde{\mathcal{U}}, J) = \text{Thom}_{\tilde{K}}^{\underline{\Delta}}(\tilde{\mathcal{U}}, J)$ in $\mathbf{K}_K(\mathbf{T}_K \tilde{\mathcal{U}})$ (see Lemma 3.4).

Let \tilde{J} be the K -invariant almost complex structure on \tilde{M} , constant on the fibers of $\tilde{M} \rightarrow K/T$, and equal to J_λ at $[1, 0]$ so that for $[k, x] \in K \times_T \mathfrak{p}$, $\tilde{J}_{[k, x]}(V) = [k, x, J_\lambda(X + v)]$ for $V = [k, x, X + v]$. Since the set $\{\underline{\Delta}_{\tilde{M}} = 0\} = K/T$ is compact, using \tilde{J} and the map $\underline{\Delta}$, one defines the localized Thom symbol

$$\text{Thom}_{\tilde{K}}^{\underline{\Delta}}(\tilde{M}, \tilde{J}) \in \mathbf{K}_K(\mathbf{T}_K \tilde{M}) .$$

Through the canonical identification of the tangent spaces at $[k, x]$ and $[k, 0]$, one can write $\tilde{J}_{[k, x]} = \tilde{J}_{[k, 0]} = J_{[k, 0]}$ for any $[k, x] \in \tilde{\mathcal{U}}$. We note that J and \tilde{J} are related on $\tilde{\mathcal{U}}$ by the homotopy J^t of almost complex structures: $J_{[k, x]}^t := J_{[k, tx]}$ for $[k, x] \in \tilde{\mathcal{U}}$. From Lemma 3.4, we conclude that the localized Thom symbols $\text{Thom}_{\tilde{K}}^{\underline{\Delta}}(\tilde{\mathcal{U}}, J)$ and $\text{Thom}_{\tilde{K}}^{\underline{\Delta}}(\tilde{M}, \tilde{J})|_{\tilde{\mathcal{U}}}$ define the same class in $\mathbf{K}_K(\mathbf{T}_K \tilde{\mathcal{U}})$, thus (5.50) becomes

$$(5.51) \quad RR_{\Phi}^K(G \cdot \lambda, V) = \text{Index}_M^K \left(\text{Thom}_{\tilde{K}}^{\underline{\Delta}}(\tilde{M}, \tilde{J}) \otimes \pi^*(K \times_T V_\lambda) \right) .$$

In order to compute (5.51), we use the induction morphism

$$i_* : \mathbf{K}_T(\mathbf{T}_T \mathfrak{p}) \longrightarrow \mathbf{K}_K(\mathbf{T}_K(K \times_T \mathfrak{p}))$$

defined by Atiyah in [1] (see [33][Section 3]). The map i_* enjoys two properties: first, i_* is an isomorphism and the K -index of $\sigma \in \mathbf{K}_K(\mathbf{T}_K(K \times_T \mathfrak{p}))$ can be computed with the T -index of $(i_*)^{-1}(\sigma)$.

Let $\sigma : p^*(E^+) \rightarrow p^*(E^-)$ be a K -transversally elliptic symbol on $K \times_T \mathfrak{p}$, where $p : \mathbf{T}(K \times_T \mathfrak{p}) \rightarrow K \times_T \mathfrak{p}$ is the projection, and E^+, E^- are equivariant vector bundles over $K \times_T \mathfrak{p}$. So for any $[k, x] \in K \times_T \mathfrak{p}$, we have a collection of linear maps $\sigma([k, x, X + v]) : E_{[k, x]}^+ \rightarrow E_{[k, x]}^-$ depending on the tangent vectors $X + v$. The symbol $(i_*)^{-1}(\sigma)$ is defined by

$$(5.52) \quad (i_*)^{-1}(\sigma)(x, v) = \sigma([1, x, 0 + v]) : E_{[1, x]}^+ \longrightarrow E_{[1, x]}^- \quad \text{for any } (x, v) \in \mathbf{T}\mathfrak{p}.$$

For $\sigma = \text{Thom}_{\tilde{K}}^{\tilde{\Phi}}(\tilde{M}, \tilde{J})$, the vector bundle E^+ (resp. E^-) is $\wedge_{\mathbb{C}}^{\text{odd}} \mathbf{T}\tilde{M}$ (resp. $\wedge_{\mathbb{C}}^{\text{even}} \mathbf{T}\tilde{M}$). Since the complex structure leaves $\mathfrak{t} \cong \mathfrak{k}/\mathfrak{t}$ and \mathfrak{p} invariant one gets

$$(i_*)^{-1}(\text{Thom}_{\tilde{K}}^{\underline{\Delta}}(\tilde{M}, \tilde{J})) = \text{Thom}_T^{\lambda}(\mathfrak{p}, J_\lambda) \wedge_{\mathbb{C}}^{\bullet} \mathfrak{k}/\mathfrak{t} ,$$

and

$$(5.53) \quad (i_*)^{-1} \left(\text{Thom}_{\tilde{K}}^{\underline{\Delta}}(\tilde{M}, \tilde{J}) \otimes \pi^*(K \times_T V_\lambda) \right) = \text{Thom}_T^{\lambda}(\mathfrak{p}, J_\lambda) V_\lambda \wedge_{\mathbb{C}}^{\bullet} \mathfrak{k}/\mathfrak{t} ,$$

where $\text{Thom}_T^{\lambda}(\mathfrak{p}, J_\lambda)$ is the T -equivariant Thom symbol on the complex vector space $(\mathfrak{p}, J_\lambda)$ deformed by the constant map $\mathfrak{p} \rightarrow \mathfrak{t}$, $x \mapsto \lambda$. In (5.53), our notation uses the structure of $R(T)$ -module for $\mathbf{K}_T(\mathbf{T}_T \mathfrak{p})$, hence we can multiply $\text{Thom}_T^{\lambda}(\mathfrak{p}, J_\lambda)$ by $V_\lambda \wedge_{\mathbb{C}}^{\bullet} \mathfrak{k}/\mathfrak{t}$.

Theorem 4.1 of Atiyah in [1] tells us that

$$(5.54) \quad \begin{array}{ccc} K_T(\mathbf{T}_T \mathfrak{p}) & \xrightarrow{i_*} & K_K(\mathbf{T}_K \tilde{M}) \\ \text{Index}_{\mathfrak{p}}^T \downarrow & & \downarrow \text{Index}_{\tilde{M}}^K \\ \mathcal{C}^{-\infty}(T) & \xrightarrow[\text{Ind}_T^K]{} & \mathcal{C}^{-\infty}(K)^K. \end{array}$$

is a commutative diagram, with $\tilde{M} = K \times_T \mathfrak{p}$, and where Ind_T^K is the induction map (see (4.39)). In other words, $\text{Index}_{\tilde{M}}^K(\sigma) = \text{Ind}_T^K(\text{Index}_{\mathfrak{p}}^T((i_*)^{-1}(\sigma)))$. With (5.51), (5.53), and (5.54), we find

$$\begin{aligned} RR_{\Phi}^K(G \cdot \lambda, V) &= \text{Ind}_T^K \left(\text{Index}_{\mathfrak{p}}^T(\text{Thom}_T^{\lambda}(\mathfrak{p}, J_{\lambda})) V_{\lambda} \wedge_{\mathbb{C}}^{\bullet} \mathfrak{k}/\mathfrak{t} \right) \\ &= \text{Hol}_T^K \left(\text{Index}_{\mathfrak{p}}^T(\text{Thom}_T^{\lambda}(\mathfrak{p}, J_{\lambda})) V_{\lambda} \right). \end{aligned}$$

(See the Appendix in [33] for the relation $\text{Hol}_T^K(-) = \text{Ind}_T^K(- \wedge_{\mathbb{C}}^{\bullet} \mathfrak{k}/\mathfrak{t})$.) But the index $\text{Index}_{\mathfrak{p}}^T(\text{Thom}_T^{\lambda}(\mathfrak{p}, J_{\lambda}))$ is computed in Section 5 of [33]:

$$\begin{aligned} \text{Index}_{\mathfrak{p}}^T(\text{Thom}_T^{\lambda}(\mathfrak{p}, J_{\lambda})) &= \left[\Pi_{\alpha \in \mathfrak{R}_n^+} (1 - t^{-\alpha}) \right]_{\lambda}^{-1} \\ &= (-1)^r t^{2\rho_n} \left[\Pi_{\alpha \in \mathfrak{R}_n^+} (1 - t^{\alpha}) \right]_{\lambda}^{-1}, \end{aligned}$$

with $r = \frac{1}{2} \dim(G/K)$. Equality (5.49) is then proved.

5.3. $(G \cdot \lambda, \Phi)$ satisfies Assumption 3.6. Let M be a regular elliptic coadjoint orbit for G , with the canonical Hamiltonian K -action. The goal of this section is to show that M satisfies Assumption 3.6 at every μ .

Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition of \mathfrak{g} . The Killing form B provides a G -equivariant identification $\mathfrak{g} \simeq \mathfrak{g}^*$ and K -equivariant identifications $\mathfrak{k} \simeq \mathfrak{k}^*$, $\mathfrak{p} \simeq \mathfrak{p}^*$. The Killing form B provides also a K -invariant Euclidean structure on \mathfrak{g} such that $B(X, X) = -\|X_1\|^2 + \|X_2\|^2$ and $\|X\|^2 = \|X_1\|^2 + \|X_2\|^2$, for $X = X_1 + X_2$, with $X_1 \in \mathfrak{k}$, $X_2 \in \mathfrak{p}$.

Hence we can and we shall consider M as a adjoint orbit of G : $M = G \cdot \lambda$ where $\lambda \in \mathfrak{k}$ is a regular element, i.e. $G_{\lambda} = K_{\lambda}$ is a maximal torus in K (in this section \cdot means the adjoint action). The moment map $\Phi : M \rightarrow \mathfrak{k}$ is then the restriction to M of the orthogonal projection $\mathfrak{k} \oplus \mathfrak{p} \rightarrow \mathfrak{k}$. For $\mu \in \mathfrak{k}$, we consider the map $\Phi_{\mu} : M \times K \cdot \mu \rightarrow \mathfrak{k}$, $(m, n) \mapsto \Phi(m) - n$.

This section is devoted to the proof of the following

Proposition 5.7. *The set $\text{Cr}(\|\Phi_{\mu}\|^2)$ of critical points of $\|\Phi_{\mu}\|^2$ is a compact subset of $M \times K \cdot \mu$. More precisely, for any $r \geq 0$, there exists $c(r) > 0$ such that*

$$\text{Cr}(\|\Phi_{\mu}\|^2) \subset \left(M \cap \left\{ \xi \in \mathfrak{g}, \|\xi\| \leq c(r) \right\} \right) \times K \cdot \mu,$$

whenever $\|\mu\| \leq r$.

Note that $M \cap \{\xi \in \mathfrak{g}, \|\xi\| \leq c(r)\}$ is compact, thus Proposition 5.7 shows that M satisfies Assumption 3.6 at every μ . Let \mathfrak{a} be a maximal Abelian subalgebra of \mathfrak{p} , and consider the map

$$F^{\mu} : (K \cdot \lambda) \times (K \cdot \mu) \times \mathfrak{a} \rightarrow \mathbb{R}$$

defined by $F^\mu(m, n, X) = \frac{1}{2} \|e^X \cdot m\|^2 - 2 \langle e^X \cdot m, n \rangle$.

Proposition 5.8. *For any $r \geq 0$, there exists $c(r) > 0$ such that*

$$(m, n, X) \in \text{Cr}(F^\mu) \implies \|e^X \cdot m\| \leq c(r),$$

whenever $\|\mu\| \leq r$.

We first show that Proposition 5.8 implies Proposition 5.7, and then we concentrate on the proof of Proposition 5.8.

Proposition 5.8 \implies Proposition 5.7. Consider the map $\Phi - \mu : M \rightarrow \mathfrak{k}$. One easily sees that $\text{Cr}(\|\Phi_\mu\|^2) = K \cdot (\text{Cr}(\|\Phi_\mu\|^2) \cap (M \times \{\mu\}))$, and $\text{Cr}(\|\Phi_\mu\|^2) \cap (M \times \{\mu\}) \subset \text{Cr}(\|\Phi - \mu\|^2) \times \{\mu\}$. Thus Proposition 5.7 is proved if one shows that for any $r \geq 0$, there exists $c(r) > 0$ such that

$$\text{Cr}(\|\Phi - \mu\|^2) \subset M \cap \left\{ \xi \in \mathfrak{g}, \|\xi\| \leq c(r) \right\},$$

whenever $\|\mu\| \leq r$. Since the bilinear form B is G -invariant, the map $m \rightarrow B(m, m)$ is constant on M , equal to $-\|\lambda\|^2$, and thus $\|\Phi(m)\|^2 = \frac{1}{2} \|m\|^2 + \frac{1}{2} \|\lambda\|^2$ for any $m \in M$. Finally we have on M the equality

$$\|\Phi(m) - \mu\|^2 = \frac{1}{2} \|m\|^2 - 2 \langle m, \mu \rangle + \text{cst}$$

where $\text{cst} = \frac{1}{2} \|\lambda\|^2 + \|\mu\|^2$. If we use now the Cartan decomposition $G = K \cdot \exp(\mathfrak{p})$, and the fact that $\mathfrak{p} = \cup_{k \in K} k \cdot \mathfrak{a}$, we see that every element M is of the form $m = (k_1^{-1} e^X k_2) \cdot \lambda$ with $k_1, k_2 \in K$ and $X \in \mathfrak{a}$. It follows that $\|\Phi((k_1^{-1} e^X k_2) \cdot \lambda) - \mu\|^2 = F^\mu(m', n, X) + \text{cst}$ with $m' = k_2 \cdot \lambda$, $n = k_1 \cdot \mu$. It is now obvious that if $m = (k_1^{-1} e^X k_2) \cdot \lambda \in \text{Cr}(\|\Phi - \mu\|^2)$ then $(k_2 \cdot \lambda, k_1 \cdot \mu, X) \in \text{Cr}(F^\mu)$. Finally, if Proposition 5.8 holds we get $\|m\| = \|e^X \cdot m'\| \leq c(r)$. \square

Proof of Proposition 5.8. Let $(m, n, X) \in \text{Cr}(F^\mu)$. Then, the identity $\frac{d}{dt} F^\mu(m, n, X + tX)|_{t=0} = 0$ gives

$$(5.55) \quad \langle e^X \cdot m, e^X \cdot [X, m] \rangle = 2 \langle e^X \cdot [X, m], n \rangle.$$

The proof of Proposition 5.8 is then reduced to the

Lemma 5.9. i) *For any $r \geq 0$, there exists $d(r) > 0$ such that $\|e^X \cdot [X, m]\| \leq d(r) \|X\|$ holds for every $(m, n, X) \in K \cdot \lambda \times \mathfrak{k} \times \mathfrak{a}$ satisfying (5.55) and $\|n\| \leq r$.*
ii) *For any $d > 0$ there exists $c > 0$, such that for every $(m, X) \in K \cdot \lambda \times \mathfrak{a}$, we have $\|e^X \cdot [X, m]\| \leq d \|X\| \implies \|e^X \cdot m\| \leq c$.*

Proof of i). Let Σ be the set of weights for the adjoint action of \mathfrak{a} on \mathfrak{g} : $\mathfrak{g} = \sum_{\alpha \in \Sigma} \mathfrak{g}_\alpha$, where $\mathfrak{g}_\alpha = \{Z \in \mathfrak{g}, [X, Z] = \alpha(X)Z \text{ for all } X \in \mathfrak{a}\}$. Each $m \in \mathfrak{g}$ admits a decomposition $m = \sum_\alpha m_\alpha$, with $m_\alpha \in \mathfrak{g}_\alpha$, which is stable relatively to the Cartan involution:

$$(5.56) \quad \theta(m_\alpha) = m_{-\alpha}, \quad \text{for every } m \in \mathfrak{g}.$$

Suppose now that $v := (m, n, X) \in K \cdot \lambda \times \mathfrak{k} \times \mathfrak{a}$ satisfies (5.55). We decompose $m \in K \cdot \lambda$ into $m = \sum_\alpha m_\alpha$ with $m_\alpha \in \mathfrak{g}_\alpha$. Let $\Sigma_v^\pm := \{\alpha \in \Sigma, m_\alpha \neq 0 \text{ and } \pm$

$\alpha(X) > 0\}$. The LHS of (5.55) decomposes in $\text{LHS} = \sum_{\alpha} e^{2\alpha(X)} \alpha(X) \|m_{\alpha}\|^2$, and

$$\begin{aligned} \text{LHS} &= \sum_{\alpha \in \Sigma_v^+} e^{2\alpha(X)} \alpha(X) \|m_{\alpha}\|^2 + \sum_{\alpha \in \Sigma_v^-} e^{2\alpha(X)} \alpha(X) \|m_{\alpha}\|^2 \\ &\geq \sum_{\alpha \in \Sigma_v^+} e^{2\alpha(X)} \frac{\alpha(X)^2}{R \|X\|} \|m_{\alpha}\|^2 - R \|X\| \sum_{\alpha \in \Sigma_v^-} \|m_{\alpha}\|^2 \quad [1] \end{aligned}$$

with $R := \sup_{\alpha, \|X\| \leq 1} |\alpha(X)|$. But

$$\begin{aligned} \sum_{\alpha \in \Sigma_v^+} e^{2\alpha(X)} \alpha(X)^2 \|m_{\alpha}\|^2 &= \|e^X \cdot [X, m]\|^2 - \sum_{\alpha \in \Sigma_v^-} e^{2\alpha(X)} \alpha(X)^2 \|m_{\alpha}\|^2 \\ &\geq \|e^X \cdot [X, m]\|^2 - R^2 \|X\|^2 \sum_{\alpha \in \Sigma_v^-} \|m_{\alpha}\|^2 \quad [2] \end{aligned}$$

Since $\alpha \in \Sigma_v^+ \Leftrightarrow -\alpha \in \Sigma_v^-$, we have $2 \sum_{\alpha \in \Sigma_v^-} \|m_{\alpha}\|^2 \leq \sum_{\alpha \in \Sigma} \|m_{\alpha}\|^2 = \|m\|^2 = \|\lambda\|^2$. So, the inequalities [1] and [2] give

$$(5.57) \quad \text{LHS} \geq \frac{\|e^X \cdot [X, m]\|^2}{R \|X\|} - R \|X\| \cdot \|\lambda\|^2.$$

Since the RHS of (5.55) satisfies obviously $\text{RHS} \leq 2 \|e^X \cdot [X, m]\| \cdot \|n\|$, (5.55) and (5.57) yield

$$2 \|e^X \cdot [X, m]\| \cdot \|n\| \geq \frac{\|e^X \cdot [X, m]\|^2}{R \|X\|} - R \|X\| \cdot \|\lambda\|^2.$$

In other words $E := \|e^X \cdot [X, m]\|$ satisfies the polynomial inequality $E^2 - 2aE - b^2 \leq 0$, with $b = R \|X\| \cdot \|\lambda\|$ and $a = R \|X\| \cdot \|n\|$. A direct computation gives

$$\|e^X \cdot [X, m]\| \leq d \|X\|,$$

with $d = R(\|n\| + \sqrt{\|n\|^2 + \|\lambda\|^2})$. \square

Proof of ii). Suppose that ii) does not hold. So there is a sequence $(m_i, X_i)_{i \in \mathbb{N}}$ in $K \cdot \lambda \times \mathfrak{a}$ such that $\|e^{X_i} \cdot [X_i, m_i]\| \leq d \|X_i\|$ but $\lim_{i \rightarrow \infty} \|e^{X_i} \cdot m_i\| = \infty$. We write $X_i = t_i v_i$ with $t_i \geq 0$ and $\|v_i\| = 1$. We can assume moreover that $v_i \rightarrow v_{\infty} \in \mathfrak{a}$ with $\|v_{\infty}\| = 1$, and $m_i \rightarrow m_{\infty} \in K \cdot \lambda$ when $i \rightarrow \infty$.

But $m_{\infty} \in K \cdot \lambda$ is a regular element of G , and $\text{rank}(G) = \text{rank}(K)$, thus $[v_{\infty}, m_{\infty}] = \sum_{\alpha \in \Sigma} \alpha(v_{\infty}) m_{\infty, \alpha} \neq 0$: there exists $\alpha_o \in \Sigma$ such that $\alpha_o(v_{\infty}) m_{\infty, \alpha_o} \neq 0$, and then also $\alpha_o(v_{\infty}) m_{\infty, -\alpha_o} \neq 0$ (see (5.56)).

On one hand the sequence $e^{t_i v_i} \cdot m_i = \sum_{\alpha} e^{t_i \alpha(v_i)} m_{i, \alpha}$ diverges. Hence $(t_i)_{i \in \mathbb{N}}$ is not bounded and so can be assumed to be divergent. On the other hand $e^{t_i v_i} \cdot [v_i, m_i] = \sum_{\alpha} e^{t_i \alpha(v_i)} \alpha(v_i) m_{i, \alpha}$ is bounded, so the sequences $e^{t_i \pm \alpha_o(v_i)} \alpha(v_i) m_{i, \pm \alpha_o}$ are also bounded. But $\lim_{i \rightarrow \infty} \alpha(v_i) m_{i, \pm \alpha_o} = \alpha_o(v_{\infty}) m_{\infty, \pm \alpha_o} \neq 0$, hence the sequences $e^{t_i \pm \alpha_o(v_i)}$ are bounded. This contradicts the fact that $\lim_{i \rightarrow \infty} t_i = +\infty$ and $\lim_{i \rightarrow \infty} \alpha_o(v_i) \neq 0$. \square

REFERENCES

- [1] M.F. ATIYAH, Elliptic operators and compact groups, Springer, 1974. Lecture notes in Mathematics, **401**.
- [2] M.F. ATIYAH, Convexity and commuting hamiltonians, *Bull. London Math. Soc.* **14**, 1982, p. 1-15.
- [3] M.F. ATIYAH, R. BOTT and A. SHAPIRO, Clifford modules, *Topology* **3**, Suppl. 1, p. 3-38.

- [4] M.F. ATIYAH, G.B. SEGAL, The index of elliptic operators II, *Ann. Math.* **87**, 1968, p. 531-545.
- [5] M.F. ATIYAH, I.M. SINGER, The index of elliptic operators I, *Ann. Math.* **87**, 1968, p. 484-530.
- [6] M.F. ATIYAH, I.M. SINGER, The index of elliptic operators III, *Ann. Math.* **87**, 1968, p. 546-604.
- [7] M.F. ATIYAH, I.M. SINGER, The index of elliptic operators IV, *Ann. Math.* **93**, 1971, p. 139-141.
- [8] N. BERLINE, E. GETZLER and M. VERGNE, *Heat kernels and Dirac operators*, Grundlehren, vol. 298, Springer, Berlin, 1991.
- [9] N. BERLINE and M. VERGNE, The Chern character of a transversally elliptic symbol and the equivariant index, *Invent. Math.* **124**, 1996, p. 11-49.
- [10] N. BERLINE and M. VERGNE, L'indice équivariant des opérateurs transversalement elliptiques, *Invent. Math.* **124**, 1996, p. 51-101.
- [11] A. CANNAS DA SILVA, Y. KARSHON and S. TOLMAN, Quantization of presymplectic manifolds and circle actions, *Trans. Amer. Math. Soc.* **352**, 2000, p. 525-552.
- [12] M. DUFLO, Représentations de carré intégrable des groupes semi-simples réels, *Séminaire Bourbaki*, Vol. 1977/78, Exposé No.508, Lect. Notes Math. 710, 22-40 (1979).
- [13] M. DUFLO, G. HECKMAN and M. VERGNE, Projection d'orbites, formule de Kirillov et formule de Blattner, *Mémoires de la S.M.F.* **15**, 1984, p. 65-128.
- [14] J. J. DUISTERMAAT, *The heat lefschetz fixed point formula for the $Spin^c$ -Dirac operator*, Progress in Nonlinear Differential Equation and Their Applications, vol. 18, Birkhauser, Boston, 1996.
- [15] V. GUILLEMIN and S. STERNBERG, Convexity properties of the moment mapping, *Invent. Math.* **67**, 1982, p. 491-513.
- [16] V. GUILLEMIN and S. STERNBERG, Geometric quantization and multiplicities of group representations, *Invent. Math.* **67**, 1982, p. 515-538.
- [17] V. GUILLEMIN and S. STERNBERG, A normal form for the moment map, in *Differential Geometric Methods in Mathematical Physics*(S. Sternberg, ed.), Reidel Publishing Company, Dordrecht, 1984.
- [18] L. JEFFREY and F. KIRWAN, Localization and quantization conjecture, *Topology* **36**, 1997, p. 647-693.
- [19] HARISH-CHANDRA Discrete series for semi-simple Lie group, I and II, *Acta Mathematica*, **113** (1965) p. 242-318, and **116** (1966) p. 1-111.
- [20] H. HECHT and W. SCHMID, A proof of Blattner's conjecture, *Invent. Math.*, **31**, 1975, p. 129-154.
- [21] T.KAWASAKI, The index of elliptic operators over V-manifolds, *Nagoya Math. J.*, **84**, 1981, p. 135-157.
- [22] F. KIRWAN, *Cohomology of quotients in symplectic and algebraic geometry*, Princeton Univ. Press, Princeton, 1984.
- [23] F. KIRWAN, Convexity properties of the moment mapping III, *Invent. Math.* **77**, 1984, p. 547-552.
- [24] B. KOSTANT, Quantization and unitary representations, in *Modern Analysis and Applications*, Lecture Notes in Math., Vol. 170, Spinger-Verlag, 1970, p. 87-207.
- [25] H. LAWSON and M.-L. MICHELSON, *Spin geometry*, Princeton Math. Series, 38. Princeton Univ. Press, Princeton, 1989.
- [26] E. LERMAN, E. MEINRENKEN, S. TOLMAN and C. WOODWARD, Non-Abelian convexity by symplectic cuts, *Topology* **37**, 1998, p. 245-259.
- [27] E. MEINRENKEN, On Riemann-Roch formulas for multiplicities, *J. Amer. Math. Soc.*, **9**, 1996, p. 373-389.
- [28] E. MEINRENKEN, Symplectic surgery and the $Spin^c$ -Dirac operator, *Adv. in Math.* **134**, 1998, p. 240-277.
- [29] E. MEINRENKEN, S. SJAMAAR, Singular reduction and quantization, *Topology* **38**, 1999, p. 699-762.
- [30] P-E. PARADAN, Formules de localisation en cohomologie équivariante, *Compositio Mathematica* **117**, 1999, p. 243-293.
- [31] P-E. PARADAN, The moment map and equivariant cohomology with generalized coefficient, *Topology*, **39**, 2000, p. 401-444.

- [32] P-E. PARADAN, The Fourier transform of semi-simple coadjoint orbits, *J. Funct. Anal.* **163**, 1999, p. 152-179.
- [33] P-E. PARADAN, Localization of the Riemann-Roch character, *J. Funct. Anal.* **187**, 2001, p. 442-509..
- [34] W. SCHMID, On a conjecture of Langlands, *Ann. of Math.* **93**, 1971, p. 1-42.
- [35] W. SCHMID, L^2 -cohomology and the discrete series, *Ann. of Math.* **103**, 1976, p. 375-394.
- [36] W. SCHMID, Discrete Series, *Proc. of Symp. in Pure Math.* **61**, 1997, p. 83-113.
- [37] R. SJAMAAR, Symplectic reduction and Riemann-Roch formulas for multiplicities, *Bull. Amer. Math. Soc.* **33**, 1996, p. 327-338.
- [38] R. SJAMAAR, Convexity properties of the moment mapping re-examined, *Adv. Math.* **138**, 1998, p. 46-91.
- [39] G. SEGAL, Equivariant K-Theory, *Publ. Math. IHES* **34**, 1968, p. 129-151.
- [40] Y. TIAN, W. ZHANG, An analytic proof of the geometric quantization conjecture of Guillemin-Sternberg, *Invent. Math.* **132**, 1998, p. 229-259.
- [41] M. VERGNE, Geometric quantization and equivariant cohomology, First European Congress in Mathematics, vol. 1, *Progress in Mathematics* **119**, Birkhauser, Boston, 1994, p. 249-295.
- [42] M. VERGNE, Multiplicity formula for geometric quantization, Part I, Part II, and Part III, *Duke Math. Journal* **82**, 1996, p. 143-179, p 181-194, p 637-652.
- [43] M. VERGNE, Quantification géométrique et réduction symplectique, *Séminaire Bourbaki* **888**, 2001.
- [44] E. WITTEN, Two dimensional gauge theories revisited, *J. Geom. Phys.* **9**, 1992, p. 303-368.
- [45] WOODHOUSE, *Geometric quantization*, 2nd ed. Oxford Mathematical Monographs. Oxford: Clarendon Press, 1997.