



HAL
open science

Exact simulation for SDE with discontinuous drift

Pierre Etoré, Miguel Martinez

► **To cite this version:**

Pierre Etoré, Miguel Martinez. Exact simulation for SDE with discontinuous drift. 2013. hal-00775871v1

HAL Id: hal-00775871

<https://hal.science/hal-00775871v1>

Preprint submitted on 14 Jan 2013 (v1), last revised 4 Oct 2013 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exact simulation for solutions of one-dimensional Stochastic Differential Equations with discontinuous drift

Pierre Étoré^{a,1,*}, Miguel Martinez^b

^aLaboratoire Jean Kuntzmann- Tour IRMA 51, rue des Mathématiques, 38041 Grenoble Cedex 9, France.

^bUniversité Paris-Est, Laboratoire d'Analyse et de Mathématiques Appliquées, UMR 8050, 5 Bld Descartes, Champs-sur-marne, 77454 Marne-la-Vallée Cedex 2, France.

Abstract

In this note we propose an exact simulation algorithm for the solution of

$$dX_t = dW_t + \bar{b}(X_t)dt, \quad X_0 = x, \quad (1)$$

where \bar{b} is a smooth real function except at point 0 where $\bar{b}(0+) \neq \bar{b}(0-)$. The main idea is to sample an exact skeleton of X using an algorithm deduced from the convergence of the solutions of the skew perturbed equation

$$dX_t^\beta = dW_t + \bar{b}(X_t^\beta)dt + \beta dL_t^0(X^\beta), \quad X_0 = x \quad (2)$$

towards X solution of (1) as $\beta \neq 0$ tends to 0.

In this note, we show that this convergence induces the convergence of exact simulation algorithms proposed by the authors in [7] for the solutions of (2) towards a limit algorithm. Thanks to stability properties of the rejection procedures involved as β tends to 0, we prove that this limit algorithm is an exact simulation algorithm for the solution of the limit equation (1). Numerical examples are shown to illustrate the performance of this exact simulation algorithm.

Keywords:

Exact simulation methods ; Brownian motion with Two-Valued Drift ; One-dimensional diffusion ; Local Time.

1. Introduction

1.1. Motivations and exposition of the problem

Exact simulation methods for trajectories of one-dimensional SDEs has been a subject of much interest in the last years : see for example [6], [3], [4], [12], [14]. Unlike the classical simulation methods, which all involve some kind of discretization error (see for example [1] for the Euler Scheme), the exact simulation methods are constructed in such a way that they do not present any discretization error (under the strong hypothesis that the diffusion coefficient is constant and equal to one). In the last years, the original method presented in the fundamental article [3] has been extended to overcome various limitations of the initial algorithm ; it has been generalized to include the cases of unbounded drifts ([4], [5]) and extended to various 'non classical' type of SDE ([7]).

In this paper, we present an attempt for the adaptation of the exact simulation methods of [3] to one-dimensional SDE that possess a discontinuous drift at point 0. Namely, our object of study is $(X_t)_{t \geq 0}$ solution of

$$dX_t = W_t + \bar{b}(X_t)dt, \quad X_0 = x, \quad (3)$$

*Corresponding author

Email addresses: pierre.etore@imag.fr (Pierre Étoré), miguel.martinez@univ-mlv.fr (Miguel Martinez)

¹Phone: + 33 (0)4 76 51 45 57; Fax: +33 (0)4 76 63 12 63

where \bar{b} is a smooth real function except possibly at point 0 where $\bar{b}(0+) \neq \bar{b}(0-)$.

The simplest case of a process solution of an equation of type (3) is surely the so-called 'Brownian motion with two valued drift' solution of

$$dX_t = W_t + (\theta_0 1_{X_t > 0} + \theta_1 1_{X_t < 0}) dt, \quad X_0 = x, \quad (4)$$

where $(\theta_0, \theta_1) \in \mathbb{R}^2$. For a general reference concerning these type of motions, we refer to [10] p.440-441 or [9]. These motions appear in stochastic control problems (see for example [2], [9]) and also theoretical studies concerning representations of reflected Brownian motion with drift (see [8] in the case $\theta_0 = -\theta_1$). Even though there exists explicit representation formulae for the densities of such Brownian motions with two valued drift in terms of combination of convolution integrals (see [10] p.440-441), up to our knowledge there is no exact numerical simulation algorithm for such motions available in the literature. The algorithm presented in this paper gives an answer to this question.

1.2. Main ideas of the paper

In [7], the authors manage to adapt the exact simulation methods of [3] to the case of one-dimensional SDEs that possess an additional term involving the local time of the unknown process at point 0. Namely, the exact simulation methods of [3] are modified in [7] to include the case where $(X_t^\beta)_{t \geq 0}$ is the solution of

$$dX_t^\beta = W_t + \bar{b}(X_t^\beta) dt + \beta dL_t^0(X^\beta), \quad X_0 = x. \quad (5)$$

In this situation $0 \neq |\beta| < 1$ (with $\beta \neq 0$), $L_t^0(X)$ denotes the symmetric local time of X^β in zero at time t , and \bar{b} is still allowed to be discontinuous at 0.

The main idea in [7] was to propose an exact rejection simulation algorithm for the solutions of (5) using as sampling reference measure the law of some drifted skew Brownian motion with prescribed terminal distribution and with drift of magnitude $1/\beta$ avoiding the case where $\beta = 0$, for which we propose a proper treatment here. Our contribution in [7] deals mainly on the simulation of such bridges of drifted skew Brownian motions using a classical rejection procedure and looking for tractable rejection functions.

Unfortunately, a direct exact simulation method along the same lines as [7] cannot be properly defined in the case where $\beta = 0$. However, we know from Le Gall in [11] that X^β solution of (5) tends strongly to X the solution of (3) as β tends to 0 on each time interval $[0, T]$. This leads us to examine what happens at the level of the algorithms proposed in [7] as β tends to 0.

In fact, we check here by computations that there is indeed a convergence phenomenon at the level of rejection functions and rejection sets involved in the exact simulation algorithms given in [7]. This convergence gives rise naturally to a nice and implementable limiting algorithm.

The main problem becomes then to prove rigorously that this limiting algorithm is indeed an exact simulation algorithm for the solution of (3). In particular, as far as we see, the direct interpretation of this limiting algorithm is not clear ; for the time being, we have to confess that we really understand the construction of the limiting algorithm exposed in this paper only *via* the convergence procedure explained above. Let us also emphasize that this new algorithm is still a rejection algorithm, and one may naturally ask for a direct interpretation of its corresponding reference measure. In Remark 1 we give an interpretation of the reference measure (corresponding to the limit rejection algorithm) in terms of a standard Brownian motion conditioned on prescribed laws for its final position and its local time at 0 at time horizon T .

1.3. Outline of the paper

The paper is organized as follows. In the preliminary Section 2, we explain the convergence of rejection sampling algorithms in a general framework. The result exposed in this section will be used

to justify that our limiting algorithm is indeed an exact simulation algorithm for the solution of (3). The exact simulation problem treated here is presented in Section 3, where we explain the manner in which we adapt the exact simulation methods of [3] to our situation. Yet, the resulting algorithm adapted from [3] is not directly implementable in our context because we have to sample from a complicated reference probability measure $\widehat{\mathbb{Z}}$. The sections 4 and 5 are devoted to the interpretation of $\widehat{\mathbb{Z}}$ as a limit of some sequence $(\widehat{\mathbb{Z}}_n)$ of better known probability measures. Finally in Section 6, we apply the results of the preliminary Section 2 to the sequence $(\widehat{\mathbb{Z}}_n)$. This gives rise to a directly implementable limit algorithm for the exact simulation of a skeleton along the reference probability $\widehat{\mathbb{Z}}$. We end up the article with numerical results and illustrative examples shown in Section 7.

2. Preliminary : convergence of abstract rejection sampling algorithms

Proposition 1 *i) Assume that we have a sequence (ξ_n) of probability measures on a measurable space (S, \mathcal{S}) , and ξ_{dom} a probability measure on (S, \mathcal{S}) , satisfying for any $n \in \mathbb{N}$*

$$\frac{d\xi_n}{d\xi_{dom}} = \frac{1}{\varepsilon_n} f_n,$$

with $\varepsilon_n > 0$ and $0 \leq f_n \leq 1$.

Assume that $f_n \rightarrow f$ as $n \rightarrow \infty$ point-wise on S and that $f_n \leq M$ for any $n \in \mathbb{N}$, with $M < \infty$.

Then, (ξ_n) converges towards a probability measure ξ satisfying

$$\frac{d\xi}{d\xi_{dom}} = \frac{1}{\varepsilon} f, \tag{6}$$

with $\varepsilon = \lim_{n \rightarrow \infty} \varepsilon_n$.

ii) Moreover, let $(Y_k, I_k)_{k \geq 1}$ be a sequence of i.i.d. random elements taking values in $S \times \{0, 1\}$ such that $Y_1 \sim \xi_{dom}$ and $\mathbb{P}[I_1 = 1 | Y_1 = y] = f(y)$ for all $y \in S$. Define $\tau = \min(k \geq 1 = I_k = 1)$. Then, $\mathbb{P}(Y_\tau \in dy) = \xi(dy)$.

Proof. For any $A \in \mathcal{S}$ we have $\xi_n(A) = \frac{1}{\varepsilon_n} \int_A f_n(z) \xi_{dom}(dz)$. By dominated convergence we have

$$\int_A f_n(z) \xi_{dom}(dz) \xrightarrow{n \rightarrow \infty} \int_A f(z) \xi_{dom}(dz).$$

Taking $A = S$, and as $\xi_n(S) = 1$ for any $n \in \mathbb{N}$, we have

$$\varepsilon_n = \frac{1}{\int_S f_n(z) \xi_{dom}(dz)} \xrightarrow{n \rightarrow \infty} \frac{1}{\int_S f(z) \xi_{dom}(dz)} =: \varepsilon.$$

Setting now for any $A \in \mathcal{S}$, $\xi(A) = \frac{1}{\varepsilon} \int_A f(z) \xi_{dom}(dz)$, it is clear that

$$\forall A \in \mathcal{S}, \quad \xi_n(A) \xrightarrow{n \rightarrow \infty} \xi(A).$$

Then ξ is a probability measure on (S, \mathcal{S}) . It satisfies (6) by construction. This proves point i). For the proof of point ii), see Proposition 1 in [6]. \square

3. Exact sampling algorithm for an SDE with discontinuous drift (inspired by the seminal paper [6])

3.1. Assumptions

The function $\bar{b} : \mathbb{R} \rightarrow \mathbb{R}$ is bounded with bounded first derivative on $\mathbb{R}^{*,+}$ and $\mathbb{R}^{*, -}$ with a possible discontinuity at point $\{0\}$. We suppose that both limits $\lim_{z \rightarrow 0+} \bar{b}(z) = \bar{b}(0+)$ and $\lim_{z \rightarrow 0-} \bar{b}(z) = \bar{b}(0-)$ exist and are finite. The value $\bar{b}(0)$ of the function \bar{b} at 0 is of no importance and can be fixed arbitrarily to some constant (possibly different from either $\bar{b}(0+)$ or $\bar{b}(0-)$).

We introduce the notation

$$\theta = \frac{\bar{b}(0+) - \bar{b}(0-)}{2}. \quad (7)$$

3.2. Change of probability

Let $0 < T < \infty$. Denote $C = C([0, T], \mathbb{R})$ the set of continuous mappings from $[0, T]$ to \mathbb{R} and \mathcal{C} the Borel σ -field on C induced by the supreme norm.

Let \mathbb{P} be a probability measure on (C, \mathcal{C}) and W a Brownian motion under \mathbb{P} together with its completed natural filtration $(\mathcal{F}_t)_{t \geq 0}$. We will denote $\mathbb{P}^x = \mathbb{P}(\cdot | W_0 = x)$. When necessary we will denote by $\omega = (\omega_t)_{0 \leq t \leq T}$ the coordinate process.

We consider the following SDE

$$dX_t = dW_t + \bar{b}(X_t)dt, \quad X_0 = x. \quad (8)$$

Our objective is to sample along X_T .

Let us define on (C, \mathcal{C}) the probability measure \mathbb{W} by

$$\frac{d\mathbb{W}}{d\mathbb{P}} = \exp \left\{ - \int_0^T \bar{b}(X_t) dW_t - \frac{1}{2} \int_0^T \bar{b}^2(X_t) dt \right\}.$$

Under \mathbb{W} the process X is a Brownian motion and we have,

$$\frac{d\mathbb{P}}{d\mathbb{W}} = \exp \left\{ \int_0^T \bar{b}(X_t) dX_t - \frac{1}{2} \int_0^T \bar{b}^2(X_t) dt \right\}.$$

Thus for any bounded measurable functional $F : (C, \mathcal{C}) \rightarrow \mathbb{R}$ we have,

$$\mathbb{E}_{\mathbb{P}}^x[F(X)] = \mathbb{E}_{\mathbb{W}}^x[F(X) \exp \left\{ \int_0^T \bar{b}(X_t) dX_t - \frac{1}{2} \int_0^T \bar{b}^2(X_t) dt \right\}]. \quad (9)$$

We set $B(x) = \int_0^x \bar{b}(y) dy$. Using the symmetric Itô-Tanaka formula (see Exercise VI-1-25 in [13]), and the Occupation times formula ([13]) we get

$$B(X_T) - B(X_0) = \int_0^T \bar{b}(X_t) dX_t + \frac{1}{2} \int_0^T \bar{b}'(X_t) dt + \frac{\bar{b}(0+) - \bar{b}(0-)}{2} L_T^0(X),$$

Thus (9) becomes,

$$\mathbb{E}_{\mathbb{P}}^x[F(X)] = \mathbb{E}_{\mathbb{W}}^x[F(X) \exp \left\{ B(X_T) - B(x) - \theta L_T^0(X) - \int_0^T \phi(X_t) dt \right\}],$$

where we have set

$$\phi(x) = \frac{\bar{b}^2(x) + \bar{b}'(x)}{2}.$$

Setting now

$$\tilde{\phi}(x) = \phi(x) - \inf_{x \in \mathbb{R}} \phi(x),$$

we finally get that for any bounded and measurable functional $F : (C, \mathcal{C}) \rightarrow \mathbb{R}$ we have,

$$\mathbb{E}_{\mathbb{P}}^x[F(X)] \propto \mathbb{E}_{\mathbb{W}}^x[F(X) \exp \{B(X_T) - B(x) - \theta L_T^0(X)\} \exp \left\{ - \int_0^T \tilde{\phi}(X_t) dt \right\}].$$

Let us now introduce the probability measure \mathbb{Z} on (C, \mathcal{C}) defined in the following way

$$\frac{d\mathbb{Z}}{d\mathbb{W}}(\omega) \propto \exp \{B(X_T(\omega)) - B(x) - \theta L_T^0(X)(\omega)\}.$$

We have

$$\mathbb{E}_{\mathbb{P}}^x[F(X)] \propto \mathbb{E}_{\mathbb{Z}}^x[F(X) \exp \left\{ - \int_0^T \tilde{\phi}(X_t) dt \right\}].$$

In the sequel we note $\widehat{\mathbb{Z}}$ the probability measure induced on (C, \mathcal{C}) by the law of X under \mathbb{Z} .

Remark 1 (Interpretation of the probability $\widehat{\mathbb{Z}}$)

Recall that under \mathbb{W} the process X is a Brownian motion and that, by definition,

$$\frac{d\mathbb{Z}}{d\mathbb{W}}(\omega) \propto \exp \{B(X_T(\omega)) - B(x) - \theta L_T^0(X)(\omega)\}.$$

In particular, under the probability \mathbb{Z} , X is a Brownian motion conditioned on $(X_T, L_T^0) \sim h(y, \ell) dy d\ell$ with

$$h(y, \ell) dy d\ell \propto \exp(B(y) - B(x) - \theta \ell) \mathbb{W}^x(X_T \in dy, L_T^0 \in d\ell).$$

This makes it difficult to sample exactly X_t under \mathbb{Z} for $t \in (0, T)$.

3.3. Exact simulation algorithm for the solution of (8) starting from x

Let us denote by K an upper bound for $\tilde{\phi}(x)$. Following the spirit of [3] we can thus sample from X_T using the following algorithm.

EXACT SIMULATION ALGORITHM FOR THE SOLUTION OF (8) STARTING FROM x .

1. SIMULATE A POISSON POINT PROCESS WITH UNIT DENSITY ON $[0, T] \times [0, K]$. THE RESULT IS A RANDOM NUMBER N OF POINTS OF COORDINATES $(t_1, z_1), \dots, (t_N, z_N)$.
 2. SIMULATE A SKELETON $(\omega_{t_1}, \dots, \omega_{t_N}, \omega_T)$ WHERE $\omega \sim \widehat{\mathbb{Z}}$.
 3. IF $\forall i \in \{1, \dots, N\} \tilde{\phi}(\omega_{t_i}) \leq z_i$ ACCEPT THE SKELETON. ELSE RETURN TO STEP 1.
-

This algorithm produces an exact sampling of X_T under \mathbb{P} : it is the final instance ω_T of an accepted skeleton.

The main issue in the above algorithm is to sample a skeleton of the canonical process under $\widehat{\mathbb{Z}}$ (Step 2).

Remark 2 (Other exact simulation algorithms ?)

Other probability changes might be performed in order to try to tackle the exact simulation problem presented in the introduction. For example (though we will not prove it here) it is possible to swap to a probability measure \mathbb{S} under which X has the law of a some Brownian motion with a symmetric two valued drift (solution of equation (4) in the case where $\theta_0 = -\theta_1$) with some prescribed terminal law. Although the density probability distribution of such bridges may be explicitly computed, because of their particular form, it seems difficult to find tractable general rejection bounds for these laws.

4. Recalls on the skew Brownian motion with drift

In this section, we recall some basic facts concerning the skew Brownian motion with constant drift. Although these facts seem at first quite far away from our purpose, they will be used in the sequel in order to justify that the limit rejection algorithm presented in Section 6 returns an exact sampling under $\widehat{\mathbb{Z}}$. At the end of this section, we give an algorithm for the simulation of bridges of SBM with constant drift that will be used as a basic building block in the sequel.

4.1. The transition function of the skew Brownian motion with drift

Let us recall that the Skew Brownian Motion (SBM) with constant drift component $\mu \in \mathbb{R}$, denoted by $B^{\beta,\mu}$, solves

$$dB_t^{\beta,\mu} = dW_t + \mu dt + \beta dL_t^0(B^{\beta,\mu}).$$

This SDE with local time has a unique strong solution as soon as $|\beta| < 1$ (see [11]). The process $B^{\beta,\mu}$ enjoys the homogeneous Markov property. We shall denote by $p^{\beta,\mu}(t, x, y)$ its transition function.

Let us introduce the function $v^{\beta,\mu}(t, x, y)$ defined by

$$\begin{aligned} v^{\beta,\mu}(t, x, y) &= (1 - e^{-2xy/t})\mathbf{1}_{xy>0} \\ &\quad + (1 + \text{Sgn}(y)\beta)e^{-2xy/t} \left[1 - \beta\mu\sqrt{2\pi t} \exp\left\{\frac{(|x|+|y|+t\beta\mu)^2}{2t}\right\} N^c\left(\frac{\beta\mu t + |x| + |y|}{\sqrt{t}}\right) \right], \end{aligned} \quad (10)$$

where $N^c(y) = \frac{1}{\sqrt{2\pi}} \int_y^\infty e^{-z^2/2} dz$.

With this notation we can rewrite the expression of $p^{\beta,\mu}(t, x, y)$ given in [7].

Proposition 2 *We have for all $t > 0$, all $x, y \in \mathbb{R}$,*

$$p^{\beta,\mu}(t, x, y) = p^{0,\mu}(t, x, y)v^{\beta,\mu}(t, x, y). \quad (11)$$

Proof. See [7] □

4.2. Bounds for the transition function of the SBM with drift

In this paragraph, we give bounds on the transition function of the SBM with drift. These bounds will be used in the sequel to find tractable rejection bounds for our algorithm.

Let us set $\bar{\alpha} = \max(\frac{1+\beta}{2}, \frac{1-\beta}{2})$ and

$$\gamma^{\beta,\mu}(t, z) = 1 - \beta\mu\sqrt{2\pi t} \exp\left(\frac{(z + t\beta\mu)^2}{2t}\right) N^c\left(\frac{\beta\mu t + z}{\sqrt{t}}\right). \quad (12)$$

We also set

$$c_{t,x}^{\beta,\mu} = \begin{cases} 2\bar{\alpha} & \text{if } \beta\mu \geq 0 \\ 2\bar{\alpha}\gamma^{\beta,\mu}(t, |x|) & \text{if } \beta\mu < 0. \end{cases} \quad (13)$$

We have the following result.

Lemma 1 *Let $(\beta, \mu) \in (-1, 1) \times \mathbb{R}$. We have*

$$v^{\beta,\mu}(t, x, y) \leq c_{t,x}^{\beta,\mu}, \quad \forall x, y \in \mathbb{R}. \quad (14)$$

Proof. Equation (14) comes from (11) and the fact that, if $\beta\mu \geq 0$, we have $p^{\beta,\mu}(t, x, y) \leq 2\bar{\alpha}p^{0,\mu}(t, x, y)$ for all $x, y \in \mathbb{R}$, and if $\beta\mu < 0$, we have $p^{\beta,\mu}(t, x, y) \leq 2\bar{\alpha}\gamma^{\beta,\mu}(t, |x|)p^{0,\mu}(t, x, y)$ for all $x, y \in \mathbb{R}$ (see, in [7], Lemma 4 and its proof). □

We also have the following lemma.

Lemma 2 Let $(\beta, \mu) \in (-1, 1) \times \mathbb{R}$. We have

$$v^{\beta, \mu}(t, x, y) \leq c_{t, y}^{\beta, \mu}, \quad \forall x, y \in \mathbb{R}. \quad (15)$$

Proof. This comes again from (11), together with the fact that $p^{\beta, \mu}(t, x, y) \leq c_{t, y}^{\beta, \mu} p^{0, \mu}(t, x, y)$, for all $x, y \in \mathbb{R}$ (see again, in [7], Lemma 4, especially the proof of Equation (19)). \square

Remark 3 Note that $v^{\beta, \mu}(t, x, y) > 0$ and $\gamma^{\beta, \mu}(t, z) > 0$ for any $t \in \mathbb{R}^{*, +}$, $x, y, z \in \mathbb{R}$, even for large values of μ (see Remark 6 in [7]).

4.3. Sampling bridges of the SBM with drift

We denote by $q^{\beta, \mu}(t, T, a, b, y)$ the density defined (for $t < T$) by

$$\mathbb{P}[B_t^{\beta, \mu} \in dy \mid B_0^{\beta, \mu} = a, B_T^{\beta, \mu} = b] = q^{\beta, \mu}(t, T, a, b, y) dy.$$

The function $(t, y) \mapsto q^{\beta, \mu}(t, T, a, b, y)$ is the transition density function of a bridge of an SBM with drift relating points a and b in T unit time.

As $q^{0, \mu}(t, T, a, b, y) = q^{0, 0}(t, T, a, b, y)$, by Proposition 2 we get,

$$q^{\beta, \mu}(t, T, a, b, y) = q^{0, 0}(t, T, a, b, y) \frac{v^{\beta, \mu}(t, a, y) v^{\beta, \mu}(T - t, y, b)}{v^{\beta, \mu}(T, a, b)}. \quad (16)$$

Let us set

$$C_{t, T, a, b}^{\beta, \mu} = \begin{cases} 4\bar{\alpha}^2 & \text{if } \beta\mu \geq 0 \\ 4\bar{\alpha}^2 \gamma^{\beta, \mu}(t, |a|) \gamma^{\beta, \mu}(T - t, |b|) & \text{if } \beta\mu < 0. \end{cases} \quad (17)$$

We have

$$\frac{q^{\beta, \mu}(t, T, a, b, y)}{q^{0, 0}(t, T, a, b, y)} = \frac{C_{t, T, a, b}^{\beta, \mu}}{v^{\beta, \mu}(T, a, b)} f_{a, b}^{\beta, \mu}(y),$$

with

$$f_{a, b}^{\mathfrak{B}, \beta, \mu}(y) = \frac{v^{\beta, \mu}(t, a, y) v^{\beta, \mu}(T - t, y, b)}{C_{t, T, a, b}^{\beta, \mu}}, \quad (18)$$

where the superscript \mathfrak{B} appears for the wording Bridge.

Considering (13), (14), (15) and (17) it is clear that

$$f_{a, b}^{\mathfrak{B}, \beta, \mu}(y) \leq 1, \quad \forall y \in \mathbb{R}.$$

We thus propose the following rejection algorithm in order to sample along $q^{\beta, \mu}(t, T, a, b, y) dy$.

AUXILIARY ALGORITHM 1: SAMPLING ALONG $q^{\beta, \mu}(t, T, a, b, y) dy$

1. SAMPLE A BROWNIAN BRIDGE Y ALONG $q^{0, 0}(t, T, a, b, y)$.

2. EVALUATE

$$f_{a, b}^{\mathfrak{B}, \beta, \mu}(Y) \leq 1.$$

3. DRAW $U \sim \mathcal{U}([0, 1])$. IF $U \leq f_{a, b}^{\mathfrak{B}, \beta, \mu}(Y)$ ACCEPT THE PROPOSED VALUE Y . ELSE RETURN TO STEP 2.

Remark 4 Note that the quantities $v^{\beta, \mu}$, $\gamma^{\beta, \mu}$, $c_{t, x}^{\beta, \mu}$, $C_{t, T, a, b}^{\beta, \mu}$, and $f_{a, b}^{\mathfrak{B}, \beta, \mu}$ defined respectively in (10), (12), (13) (17), and (18) involved in the above algorithm depend only on β and μ through the product $\beta\mu$. This computational fact gives the key ensuring the construction of the limit algorithm by convergence performed at the beginning of Section 6.

5. Convergence of a sequence of probability measures towards $\widehat{\mathbb{Z}}$

In this section, for any $n \in \mathbb{N}$ we denote by X^n the solution of

$$dX_t^n = dW_t + \bar{b}(X_t^n)dt + \frac{1}{n}L_t^0(X^n), \quad X_0^n = x. \quad (19)$$

For the existence and uniqueness of solutions to (19) see [11].

5.1. A strong convergence result

Theorem 1 (Le Gall [11], 1984) *Let X be the solution of (8) and (X^n) the sequence of solutions of (19). We have for all $0 < t < T$,*

$$\mathbb{E}\left[\sup_{0 \leq s \leq t} |X_s - X_s^n|\right] \xrightarrow{n \rightarrow \infty} 0.$$

Proof. We use the notations of [11]. Using the Occupation times formula we can rewrite Equation (19) as

$$dX_t^n = dW_t + \int_{\mathbb{R}} \nu_n(dy) dL_t^y(X^n), \quad X_0^n = x,$$

with $\nu_n(dy) = \bar{b}(y)dy + \frac{1}{n}\delta_0(dy)$. Lemma 2.1 in [11] asserts that there is for each $n \in \mathbb{N}$ a function f_{ν_n} , unique up to a multiplicative constant, satisfying $f'_{\nu_n}(dy) + (f_{\nu_n}(y) + f_{\nu_n}(y-))\nu_n(dy)$, where the notation $f'_{\nu_n}(dy)$ is for the derivative of f_{ν_n} in the generalized sense. Lemma 2.1 in [11] also asserts that if we require that $f_{\nu_n}(x) \rightarrow 1$ as $x \rightarrow -\infty$ then,

$$f_{\nu_n}(y) = \exp\left(-2 \int_{-\infty}^y \bar{b}(z)dz\right) \times \mathbf{1}_{y \geq 0} \frac{1 - 1/n}{1 + 1/n}.$$

The sequence of functions $(f_{\nu_n})_n$ clearly converges point-wise to $f(y) = \exp\left(-2 \int_{-\infty}^y \bar{b}(z)dz\right)$. By dominated convergence we have for all $K > 0$ that $\int_{-K}^K |f_{\nu_n} - f|(y)dy \rightarrow 0$ as $n \rightarrow \infty$. Thus Theorem 3.1 in [11] asserts that

$$\mathbb{E}\left[\sup_{0 \leq s \leq t} |X_s - X_s^n|\right] \xrightarrow{n \rightarrow \infty} 0,$$

with X the solution of

$$dX_t = dW_t + \int_{\mathbb{R}} \nu(dy) dL_t^y(X), \quad X_0 = x,$$

where $\nu(dy) = -\frac{f'(dy)}{f(y)+f(y-)} = -\frac{1}{2}\frac{f'(y)}{f(y)}dy = \bar{b}(y)dy$. That is to say X is the solution of (8). \square

5.2. The probability measure $\widehat{\mathbb{Z}}$ as a limit of a sequence $(\widehat{\mathbb{Z}}_n)_n$

Recall the definition (7) of θ . Let us set

$$\mu_n = \frac{1 + 1/n}{2/n} \bar{b}(0+) - \frac{1 - 1/n}{2/n} \bar{b}(0-) = \frac{\bar{b}(0+) + \bar{b}(0-)}{2} + \theta n, \quad (20)$$

and $b_n(x) = \bar{b}(x) - \mu_n$.

We have,

$$X_T^n = x + W_T^{SD,n} + \mu_n T + \frac{1}{n}L_T^0(X^n),$$

where the process $W^{SD,n}$ defined by

$$dW_t^{SD,n} = dW_t + b_n(X_t^n)dt$$

is a Brownian motion under the probability measure $\mathbb{W}^{SD,n}$ defined by

$$\frac{d\mathbb{W}^{SD,n}}{d\mathbb{P}} = \exp \left\{ - \int_0^T b_n(X_t^n) dW_t - \frac{1}{2} \int_0^T b_n^2(X_t^n) dt \right\}.$$

Let us now set $B_n(x) = \int_0^x b_n(z) dz$. As shown in [7] using a symmetric Itô-Tanaka formula we have for any bounded measurable functional $F : (C, \mathcal{C}) \rightarrow \mathbb{R}$,

$$\mathbb{E}_{\mathbb{P}}^x [F(X^n)] = \mathbb{E}_{\mathbb{W}^{SD,n}}^x [F(X^n) \exp \{ B_n(X_T^n) - B_n(x) - \int_0^T \phi_n(X_t^n) dt \}],$$

where $\phi_n(z) = \frac{b_n^2(z) + b_n'(z) + 2\mu_n b_n(z)}{2}$. Notice that the law of X^n under $\mathbb{W}^{SD,n}$ is the one of a SBM with drift.

Let us define the probability measure \mathbb{Z}_n on (C, \mathcal{C}) by

$$\frac{d\mathbb{Z}_n}{d\mathbb{W}^{SD,n}}(\omega) \propto \exp \{ B_n(X_T^n(\omega)) - B_n(x) \},$$

and $\widehat{\mathbb{Z}}_n$ the probability measure induced on (C, \mathcal{C}) by the law of X^n under \mathbb{Z}_n .

The law $\widehat{\mathbb{Z}}_n$ can be well described (see Subsection 5.3). Thus in [7] we managed to sample exactly along (19) using skeletons under $\widehat{\mathbb{Z}}_n$ as proposals and $\exp \{ - \int_0^T (\phi_n(\omega_t) - \inf \phi_n) dt \}$ for a rejection rule.

For our present purpose it turns out that we have the following result.

Proposition 3 *We have*

$$\widehat{\mathbb{Z}}_n \xrightarrow[n \rightarrow \infty]{w} \widehat{\mathbb{Z}}.$$

Proof. Let be $F : (C, \mathcal{C}) \rightarrow \mathbb{R}$ a bounded measurable functional. We have

$$\begin{aligned} \mathbb{E}_{\widehat{\mathbb{Z}}_n}^x [F(\omega)] &= \mathbb{E}_{\mathbb{Z}_n}^x [F(X^n)] \\ &= c_n \mathbb{E}_{\mathbb{W}^{SD,n}}^x [F(X^n) \exp \{ B_n(X_T^n) - B_n(x) \}] \\ &= c_n \mathbb{E}_{\mathbb{P}}^x [F(X^n) \exp \{ B(X_T^n) - B(x) \} \exp \{ -\mu_n(X_T^n - x) - \int_0^T b_n(X_t^n) dW_t - \frac{1}{2} \int_0^T b_n^2(X_t^n) dt \}]. \end{aligned}$$

But

$$\begin{aligned} & -\mu_n(X_T^n - x) - \int_0^T b_n(X_t^n) dW_t - \frac{1}{2} \int_0^T b_n^2(X_t^n) dt \\ &= -\mu_n W_T - \int_0^T \mu_n \bar{b}(X_t^n) dt - \frac{1}{n} \mu_n L_T^0(X^n) - \int_0^T \bar{b}(X_t^n) dW_t + \mu_n W_T \\ & - \frac{1}{2} \int_0^T \bar{b}^2(X_t^n) dt + \int_0^T \mu_n \bar{b}(X_t^n) dt - \frac{1}{2} \mu_n^2 T \\ &= -\frac{1}{n} \mu_n L_T^0(X^n) - \int_0^T \bar{b}(X_t^n) dW_t - \frac{1}{2} \int_0^T \bar{b}^2(X_t^n) dt - \frac{1}{2} \mu_n^2 T. \end{aligned}$$

Thus,

$$\begin{aligned} \mathbb{E}_{\widehat{\mathbb{Z}}_n}^x [F(\omega)] &= c_n e^{-\frac{\mu_n^2 T}{2}} \mathbb{E}_{\mathbb{P}}^x [F(X^n) \exp \{ B(X_T^n) - B(x) - \frac{\mu_n}{n} L_T^0(X^n) \} \exp \{ - \int_0^T \bar{b}(X_t^n) dW_t - \frac{1}{2} \int_0^T \bar{b}^2(X_t^n) dt \}]. \end{aligned}$$

From (20), we see that $\frac{\mu_n}{n} \rightarrow \theta$ as $n \rightarrow \infty$, and combining the results of Theorem 1 and Lebesgue's convergence theorem for stochastic integrals (see for example Theorem (2.12) p.142 in [13]), we have

$$\begin{aligned} & \lim_{n \rightarrow \infty} \mathbb{E}_{\widehat{\mathbb{Z}}_n} [F(\omega)] \\ &= \lim_{n \rightarrow \infty} (c_n e^{-\frac{\mu_n^2 T}{2}}) \mathbb{E}_{\mathbb{P}}^x [F(X) \exp\{B(X_T) - B(x) - \theta L_T^0(X)\} \exp\{-\int_0^T \bar{b}(X_t) dW_t - \frac{1}{2} \int_0^T \bar{b}^2(X_t) dt\}] \end{aligned}$$

and by definition of \mathbb{W} ,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \mathbb{E}_{\widehat{\mathbb{Z}}_n} [F(\omega)] \\ &= \lim_{n \rightarrow \infty} (c_n e^{-\frac{\mu_n^2 T}{2}}) \mathbb{E}_{\mathbb{W}}^x [F(X) \exp\{B(X_T) - B(x) - \theta L_T^0(X)\}] \\ &= \lim_{n \rightarrow \infty} (c_n e^{-\frac{\mu_n^2 T}{2}}) \mathbb{E}_{\mathbb{Z}}^x [F(X)] \\ &= \lim_{n \rightarrow \infty} (c_n e^{-\frac{\mu_n^2 T}{2}}) \mathbb{E}_{\mathbb{Z}}^x [F(\omega)]. \end{aligned}$$

For the special case $F \equiv 1$, the above computation ensures that $\lim_{n \rightarrow \infty} (c_n e^{-\frac{\mu_n^2 T}{2}}) = 1$. Thus, for any bounded and measurable $F : (C, \mathcal{C}) \rightarrow \mathbb{R}$ we have

$$\lim_{n \rightarrow \infty} \mathbb{E}_{\widehat{\mathbb{Z}}_n}^x [F(\omega)] = \mathbb{E}_{\mathbb{Z}}^x [F(\omega)]$$

and the proposition is proved. □

5.3. Sampling a skeleton under $\widehat{\mathbb{Z}}_n$

We have the following proposition.

Proposition 4 *For any $n \in \mathbb{N}$ the law $\widehat{\mathbb{Z}}_n$ is the one of a SBM with drift $B_n^{\frac{1}{n}, \mu_n}$ conditionally on $B_T^{\frac{1}{n}, \mu_n} \sim h_n$ with*

$$h_n(y) = C_n \exp(B_n(y) - B_n(x)) p_n^{\frac{1}{n}, \mu_n}(T, x, y),$$

where C_n is the normalizing constant such that $\int h_n(y) dy = 1$.

Proof. See [3]. □

Let n_0 be fixed and $0 = t_0 < t_1 < \dots < t_{n_0} < T$. Set $y_0 = x$ to simplify the notations, the law of $(\omega_{t_1}, \dots, \omega_{t_{n_0}}, \omega_T)$ under $\widehat{\mathbb{Z}}_n$ is given by

$$h_n(y) \prod_{i=0}^{n_0-1} q_n^{\frac{1}{n}, \mu_n}(t_{i+1} - t_i, T - t_i, y_i, y, y_{i+1}) dy_1 \dots dy_{n_0} dy. \quad (21)$$

Once ω_T has been sampled along $h_n(y) dy$, we can sample ω_{t_1} along $q_n^{\frac{1}{n}, \mu_n}(t_1, T, x, \omega_T, y_1) dy_1$ and each $\omega_{t_{i+1}}$ along $q_n^{\frac{1}{n}, \mu_n}(t_{i+1} - t_i, T - t_i, \omega_{t_i}, \omega_T, y_{i+1})$, using the Auxiliary Algorithm 1.

In order to sample along $h_n(y) dy$ we make use of the following considerations. We have

$$\begin{aligned} h_n(y) &= C_n \exp(B_n(y) - B_n(x)) p^{0, \mu_n}(T, x, y) v_n^{\frac{1}{n}, \mu_n}(T, x, y) \\ &= C_n \exp\left(-\mu_n(y - x) + \int_x^y \bar{b}(z) dz\right) \times \exp\left(+\mu_n(y - x) - \frac{\mu_n^2}{2} T\right) p^{0,0}(T, x, y) v_n^{\frac{1}{n}, \mu_n}(T, x, y) \\ &= C_n e^{-\frac{\mu_n^2}{2} T} \exp(B(y) - B(x)) v_n^{\frac{1}{n}, \mu_n}(T, x, y) p^{0,0}(T, x, y). \end{aligned}$$

Let us denote by M an upper bound for the function $z \mapsto |\bar{b}|(z)$ (see our assumptions in § 3.1). Then, using the result of Lemma 1 and performing easy computations, we easily see that for any $0 < \delta < 1$:

$$\frac{h_n(y)}{p^{0,0}(T/(1-\delta), x, y)} = C_n \frac{e^{T^2 M^2 / \delta}}{\sqrt{1-\delta}} e^{-\frac{\mu_n^2}{2} T} c_{T,x}^{\frac{1}{n}, \mu_n} f_\delta^{\frac{1}{n}, \mu_n}(y),$$

with

$$f_\delta^{\frac{1}{n}, \mu_n}(y) = \sqrt{1-\delta} \exp\left(B(y) - B(x) - \frac{T^2 M^2}{\delta}\right) \frac{p^{0,0}(T, x, y)}{p^{0,0}(T/(1-\delta), x, y)} \frac{v^{\frac{1}{n}, \mu_n}(T, x, y)}{c_{T,x}^{\frac{1}{n}, \mu_n}}.$$

Using (14) one may easily check that $f_\delta^{\frac{1}{n}, \mu_n}(y) \leq 1$ for any $y \in \mathbb{R}$. One might then optimize w.r.t. $\delta \in (0, 1)$ in order to find $f_\delta^{\frac{1}{n}, \mu_n}$ closest to 1.

Let us set for simplicity, $f^{\frac{1}{n}, \mu_n} = f_{1/2}^{\frac{1}{n}, \mu_n}$. We deduce therefore the following procedure in order to sample along $h_n(y)dy$.

AUXILIARY ALGORITHM 2: SAMPLING ALONG $h_n(y)dy$

1. SAMPLE $Y \sim \mathcal{N}(x, 2T)$.
2. EVALUATE

$$f^{\frac{1}{n}, \mu_n}(Y) \leq 1.$$

3. DRAW $U \sim \mathcal{U}([0, 1])$. IF $U \leq f^{\frac{1}{n}, \mu_n}(Y)$ ACCEPT THE PROPOSED VALUE Y . ELSE RETURN TO STEP 1.
-

6. Direct exact sampling of a skeleton under $\widehat{\mathbb{Z}}$ (performing Step 2 of the Exact Simulation Algorithm)

Proposition 1 will now play a crucial role.

Recall the definition (7) of θ . Let us denote

$$v^\theta(t, x, y) = (1 - e^{-2xy/t}) \mathbf{1}_{xy > 0} + e^{-2xy/t} [1 - \theta \sqrt{2\pi t} \exp\left\{\frac{(|x| + |y| + t\theta)^2}{2t}\right\} N^c\left(\frac{\theta t + |x| + |y|}{\sqrt{t}}\right)],$$

$$\gamma^\theta(t, z) = 1 - \theta \sqrt{2\pi t} \exp\left(\frac{(z + t\theta)^2}{2t}\right) N^c\left(\frac{\theta t + z}{\sqrt{t}}\right),$$

$$c_{t,x}^\theta = \begin{cases} 1 & \text{if } \theta \geq 0 \\ \gamma^\theta(t, |x|) & \text{if } \theta < 0, \end{cases} \quad \text{and} \quad C_{t,T,a,b}^\theta = \begin{cases} 1 & \text{if } \theta \geq 0 \\ \gamma^\theta(t, |a|) \gamma^\theta(T-t, |b|) & \text{if } \theta < 0. \end{cases}$$

From the fact that $\frac{1}{n} \mu_n \rightarrow \theta$ (as n tends to $+\infty$) and Remark 4, we clearly have that

$$v^{\frac{1}{n}, \mu_n}(t, x, y) \xrightarrow{n \rightarrow \infty} v^\theta(t, x, y) \quad \forall (t, x, y) \in \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R},$$

$$\gamma^{\frac{1}{n}, \mu_n}(t, z) \xrightarrow{n \rightarrow \infty} \gamma^\theta(t, z) \quad \forall (t, z) \in \mathbb{R}^+ \times \mathbb{R},$$

$$c_{t,x}^{\frac{1}{n}, \mu_n} \xrightarrow{n \rightarrow \infty} c_{t,x}^\theta \quad \forall (t, x) \in \mathbb{R}^+ \times \mathbb{R},$$

$$C_{t,T,a,b}^{\frac{1}{n}, \mu_n} \xrightarrow{n \rightarrow \infty} C_{t,T,a,b}^\theta \quad \forall (t, T, a, b) \in \mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}.$$

Let us now examine the sequence $(f_{\delta}^{h, \frac{1}{n}, \mu_n})$ of the rejection functions used in the Auxiliary Algorithm 2. From the same reasons as above, it is clear that $(f_{\delta}^{h, \frac{1}{n}, \mu_n})$ converges towards

$$f_{\delta}^{h, \theta}(y) = \sqrt{1 - \delta} \exp \left(B(y) - B(x) - \frac{T^2 M^2}{\delta} \right) \frac{p^{0,0}(T, x, y)}{p^{0,0}(T/(1 - \delta), x, y)} \frac{v^{\theta}(T, x, y)}{c_{T,x}^{\theta}} \leq 1,$$

this convergence being dominated. Thus, applying the result of Proposition 1, the sequence of laws $(h_n(y)dy)$ converges to some limit law $h_{\theta}(y)dy$.

In the same manner, for any fixed $a, b \in \mathbb{R}$, the sequence $(f_{a,b}^{\mathfrak{B}, \frac{1}{n}, \mu_n})$ of rejection functions used in Auxiliary Algorithm 1 converges towards

$$f_{a,b}^{\mathfrak{B}, \theta}(y) = \frac{v^{\theta}(t, a, y)v^{\theta}(T - t, y, b)}{C_{t,T,a,b}^{\theta}} \leq 1,$$

this convergence being dominated.

Consequently, the law $q^{\frac{1}{n}, \mu_n}(t, T, a, b, y)dy$ converges towards a limit law $q^{\theta}(t, T, a, b, y)dy$.

Let again n_0 be fixed and $0 < t_1 < \dots < t_{n_0} < T$. Passing to the limit in (21) we get that the law of $(\omega_{t_1}, \dots, \omega_{t_{n_0}}, \omega_T)$ under $\widehat{\mathbb{Z}}_n$ converges (with $y_0 = x$) towards

$$h_{\theta}(y) \prod_{i=0}^{n_0-1} q^{\theta}(t_{i+1} - t_i, T - t_i, y_i, y, y_{i+1})dy_1 \dots dy_{n_0}dy. \quad (22)$$

Consequently, from Proposition 3, we conclude that the law given by (22) is nothing else than the law of $(\omega_{t_1}, \dots, \omega_{t_{n_0}}, \omega_T)$ under $\widehat{\mathbb{Z}}$.

Using again Proposition 1 and the above considerations we can propose the expected algorithm in order to sample skeletons under $\widehat{\mathbb{Z}}$. It will use the two following Limit Auxiliary Algorithms.

LIMIT AUXILIARY ALGORITHM 1: SAMPLING ALONG $h_{\theta}(y)dy$

1. SAMPLE $Y \sim \mathcal{N}(x, 2T)$.
2. EVALUATE

$$f^{h, \theta}(Y) \leq 1.$$

3. DRAW $U \sim \mathcal{U}([0, 1])$. IF $U \leq f^{h, \theta}(Y)$ ACCEPT THE PROPOSED VALUE Y . ELSE RETURN TO STEP 1.

LIMIT AUXILIARY ALGORITHM 2: SAMPLING ALONG $q^{\theta}(t, T, a, b, y)dy$

1. SAMPLE A BROWNIAN BRIDGE Y ALONG $q^{0,0}(t, T, a, b, y)$.
2. EVALUATE

$$f_{a,b}^{\mathfrak{B}, \theta}(Y) \leq 1.$$

3. DRAW $U \sim \mathcal{U}([0, 1])$. IF $U \leq f_{a,b}^{\mathfrak{B}, \theta}(Y)$ ACCEPT THE PROPOSED VALUE Y . ELSE RETURN TO STEP 2.




Figure 1: Brownian motion with two-valued drift, case $\theta_0 = -\theta_1 = 1$ ($T = 1$).

PERFORMING STEP 2 OF THE EXACT SIMULATION ALGORITHM.
 SAMPLING $(\omega_{t_1}, \dots, \omega_{t_{n_0}}, \omega_T)$ UNDER $\hat{\mathbb{Z}}$ (STARTING FROM x)

1. SAMPLE ω_T ALONG $h_\theta(y)dy$ USING THE LIMIT AUXILIARY ALGORITHM 1.
 2. SAMPLE ω_{t_1} ALONG $q^\theta(t_1, T, x, \omega_T, y)dy$ USING THE LIMIT AUXILIARY ALGORITHM 2.
 3. FOR $i = 2, \dots, n_0$, SAMPLE $\omega_{t_{i+1}}$ ALONG $q^\theta(t_{i+1} - t_i, T - t_i, \omega_{t_i}, \omega_T, y)dy$ USING THE LIMIT AUXILIARY ALGORITHM 2.
-

7. Numerical Experiments

7.1. Exact simulation of a Brownian motion with two-valued (or alternate) drift

In this paragraph, we exhibit numerical results obtained with the exact limit algorithm for the simplest non-trivial cases

$$dX_t = dW_t \pm \text{sgn}(X_t)dt, \quad X_0 = 0,$$

corresponding to either $\theta_0 = -\theta_1 = \pm 1$ in (4) ($\bar{b}(y) = \pm \text{sgn}(y)$ in (1)).

We draw the renormalized histogram of 10^6 samples of X_T and compare it to the explicit density of X_T given in [10] p. 440-441 (Figure 1 for the outgoing case $\theta_0 = 1$ and Figure 2 for the incoming case $\theta_0 = -1$).




Figure 2: Brownian motion with two-valued drift, case $\theta_0 = -\theta_1 = -1$ ($T = 1$).

7.2. Exact simulation of an SDE with a discontinuous drift coefficient

We consider now the SDE (1) with

$$\bar{b}(x) = \begin{cases} -\frac{\pi}{2} \cos\left(\frac{\pi}{5}x\right) & \text{if } x \geq 0 \\ \frac{3\pi}{2} - \frac{\pi}{2} \cos\left(\frac{\pi}{5}x\right) & \text{if } x < 0. \end{cases} \quad (23)$$

Let $0 < T < \infty$. We wish to sample along X_T .

We have $\theta = -3\pi/4$ and

$$\tilde{\phi}(x) = \frac{\bar{b}^2(x) + \bar{b}'(x)}{2} + \frac{\pi^2}{20}.$$

We take $K = 2\pi^2 + \frac{\pi^2}{10}$ as an upper bound for $\tilde{\phi}$. This allows to use the limit Algorithm.

Figure 3 shows a comparison between a renormalized histogram of 10^6 samples of X_T obtained with the exact limit algorithm, and a renormalized histogram of 10^6 samples of X_T^Δ , where (X^Δ) denotes an Euler Scheme with time step Δ . We chose time-steps $\Delta = T \cdot 10^{-1}$ and $\Delta = T \cdot 10^{-5}$.

References

- [1] V. Bally and D. Talay. The law of the Euler scheme for stochastic differential equations: I. Convergence rate of the distribution function. *Probab. Theory Related Fields*, 104(1):43–60, 1996.
- [2] V. E. Beneš, L. A. Shepp, and H. S. Witsenhausen. Some solvable stochastic control problems. In *Analysis and optimisation of stochastic systems (Proc. Internat. Conf., Univ. Oxford, Oxford, 1978)*, pages 3–10. Academic Press, London, 1980.
- [3] A. Beskos, O. Papaspiliopoulos, and G.O Roberts. Retrospective exact simulation of diffusion sample paths with applications. *Bernoulli*, 12(6):1077–1098, 2006.




Figure 3: Limit algorithm v.s. Euler Scheme for the case where \bar{b} is given by (23) ($T = 1$).

- [4] A. Beskos, O. Papaspiliopoulos, and G.O. Roberts. A factorisation of diffusion measure and finite sample path constructions. *Methodol. Comput. Appl. Probab.*, 10(1):85–104, 2008.
- [5] A. Beskos, G. Roberts, A. Stuart, and J. Voss. MCMC methods for diffusion bridges. *Stoch. Dyn.*, 8(3):319–350, 2008.
- [6] A. Beskos and G.O. Roberts. Exact simulation of diffusions. *Ann. Appl. Probab.*, 15(4):2422–2444, 2005.
- [7] P. Etoré and M. Martinez. Exact simulation of one-dimensional stochastic differential equations involving the local time at zero of the unknown process. *To appear in Monte-Carlo Methods and Applications*, 2013.
- [8] S.E. Graversen and A. N. Shiryaev. An extension of P. Lévy’s distributional properties to the case of a Brownian motion with drift. *Bernoulli*, 6(4):615–620, 2000.
- [9] I. Karatzas and S. E. Shreve. Trivariate density of Brownian motion, its local and occupation times, with application to stochastic control. *Ann. Probab.*, 12(3):819–828, 1984.
- [10] I. Karatzas and S.E. Shreve. *Brownian motion and stochastic calculus. 2nd ed.* Graduate Texts in Mathematics, 113. New York etc.: Springer-Verlag. xxiii, 470 p. , 1991.
- [11] J.-F. Le Gall. One-dimensional stochastic differential equations involving the local times of the unknown process. In *Stochastic analysis and applications (Swansea, 1983)*, volume 1095 of *Lecture Notes in Math.*, pages 51–82. Springer, Berlin, 1984.
- [12] V. Reutenauer and E. Tanré. Exact simulation of prices and greeks: application to cir. *Preprint*.
- [13] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*. 3rd ed, Springer-Verlag, 1999.

- [14] M. Sbai. *Modélisation de la dépendance et simulation de processus en finance*. PhD thesis, CERMICS - Centre d'Enseignement et de Recherche en Mathématiques et Calcul Scientifique, 2009.