



HAL
open science

A global approach to the Schrödinger-Poisson system: An Existence result in the case of infinitely many states

Otared Kavian, Stéphane Mischler

► To cite this version:

Otared Kavian, Stéphane Mischler. A global approach to the Schrödinger-Poisson system: An Existence result in the case of infinitely many states. *Journal de Mathématiques Pures et Appliquées*, 2015, 104 (5), pp.942-964. 10.1016/j.matpur.2015.05.009 . hal-00787323v3

HAL Id: hal-00787323

<https://hal.science/hal-00787323v3>

Submitted on 4 Mar 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A global approach to the
Schrödinger-Poisson system:
An existence result in the case
of infinitely many states

Otared Kavian

Département de Mathématiques
Université de Versailles
45, avenue des États-Unis
78035 Versailles Cedex, France.
kavian@math.uvsq.fr

Stéphane Mischler

Ceremade
Université de Paris–Dauphine
place du Maréchal de Lattre de Tassigny
75116 Paris, France
stephane.mischler@dauphine.fr

March 4, 2015

Abstract

In this paper we prove the existence of a solution to a nonlinear Schrödinger–Poisson eigenvalue problem in dimension N , with $N \leq 6$. Our proof is based on a global approach to the determination of eigenvalues and eigenfunctions which allows us to characterize the complete sequence of eigenvalues and eigenfunctions at once, via a variational approach, and thus differs from the usual and less general proofs developed for similar problems in the literature. Our method seems to be new for the determination of the spectrum and eigenfunctions for compact and self-adjoint operators, even in a finite dimensional setting.

Keywords: Schrödinger–Poisson, variational methods, unitary operators, Palais–Smale condition.

2010 AMS Subject Classification: Primary: 49J35, 35A15, 35S15; Secondary: 47G20, 45G05.

1 Introduction

In this paper we are interested in the study of the following stationary Schrödinger system of equations: for a given bounded Lipschitz domain

$\Omega \subset \mathbb{R}^N$, find an infinite sequence $(\lambda_m, u_m)_{m \geq 1}$ and a potential V satisfying

$$\begin{cases} -\Delta u_m + \tilde{V}_0 u_m + V u_m = \lambda_m u_m & \text{in } \Omega \\ -\Delta V = \sum_{m=1}^{\infty} \rho_m |u_m|^2 & \text{in } \Omega \end{cases} \quad (1a)$$

$$\quad (1b)$$

where, in addition, we require that

$$\begin{cases} u_m \in H_0^1(\Omega), \text{ for all } m \geq 1, & V \in H_0^1(\Omega), \\ (u_m)_{m \geq 1} \text{ is a Hilbert basis for } L^2(\Omega). \end{cases} \quad (2a)$$

$$\quad (2b)$$

Here we assume that the external potential \tilde{V}_0 and the positive numbers $(\rho_m)_{m \geq 1}$ are given.

The system (1a)-(1b) appears in the modeling of nanoscale semiconductor devices as part of the so-called ‘‘quantum-kinetic subband model’’, which itself is a simplified model of the full evolution 3D-Schrödinger-Poisson equation, see N. Ben Abdallah & F. Méhats [1, 5]. In order to improve the cost of numerical simulation of the evolution 3D-Schrödinger-Poisson equation and taking advantage of the extreme confinement of the electrons in one direction transverse to the transport directions, one can perform a block diagonalisation of the electron Hamiltonian, thanks to a separation of the confinement and transport directions. This reduction process leads to replace the 3D-Schrödinger-Poisson equation by a system of 1D stationary Schrödinger equations (for the confinement) coupled to a 2D equation (for the transport). In that reduced model, the system of 1D stationary Schrödinger equations is nothing but (1a)-(1b) where the parameters $(\rho_m)_{m \geq 1}$ are the sequence of occupation numbers, which may depend on time and space, and which are given by the above mentioned 2D equation (for the transport).

On the other hand, as it is outlined in P. Zweifel [18], the Schrödinger-Poisson system of equations derives from a quantum transport equation, the Poisson-Wigner system. After performing the Wigner transform to the former system one ends up with the following system of evolution equations

$$\begin{cases} i \partial_t \psi_m = -\Delta \psi_m + \tilde{V}_0 \psi_m + V \psi_m & (3a) \\ \psi_m(0, x) = \psi_{0m}(x) & (3b) \\ -\Delta V = \sum_{m=1}^{\infty} \rho_m |\psi_m|^2, & (3c) \end{cases}$$

with the condition $(\psi_{0m} | \psi_{0k})_{L^2} = \delta_{km}$ for any $k, m \geq 1$. Seeking standing-wave solutions of the form

$$\psi_m(t, x) := e^{-i \lambda_m t} u_m(x)$$

leads to the equations (1a)-(1b).

Existence of solutions to the system (1a)-(1b) has been proved in R. Illner, O. Kavian, H. Lange [2] in the case when $\rho_m = 0$ for any $m \geq 2$, and the same method extends to the case when $\rho_m = 0$ for any $m \geq M$, for a given $M \geq 2$. On the other hand, a different but similar eigenvalue problem has been considered by F. Nier [8, 10, 9].

In order to solve this system of equations, taking into account the fact that the family $(u_m)_{m \geq 1}$ must be contained in $H_0^1(\Omega)$ and, at the same time, has to be a Hilbert basis of $L^2(\Omega)$, we observe the following. Let us consider a fixed Hilbert basis of $L^2(\Omega)$, denoted by $(e_m)_{m \geq 1}$, such that $e_m \in H_0^1(\Omega)$ for all $m \geq 1$. For instance such a basis may be given by the eigenfunctions of the Laplace operator on $H_0^1(\Omega)$, that is a family satisfying

$$-\Delta e_m = \mu_m e_m, \quad e_m \in H_0^1(\Omega), \quad \int_{\Omega} e_\ell(x) e_m(x) dx = \delta_{\ell m}, \quad (4)$$

where the sequence of eigenvalues of the Laplace operator, with Dirichlet boundary conditions, is denoted by $(\mu_m)_{m \geq 1}$. Now, saying that the family $(u_m)_{m \geq 1}$ satisfies condition (2b) means that the linear operator U acting on $L^2(\Omega)$ and defined by

$$\text{for all } m \geq 1, \quad U e_m := u_m, \quad U f := \sum_{m \geq 1} (f | e_m) u_m \quad \text{for } f \in L^2(\Omega), \quad (5)$$

is a unitary operator, that is $U^*U = UU^* = I$. Therefore determining the whole family $(u_m)_{m \geq 1}$ satisfying equations (1a)-(2b) is equivalent to find a linear operator U defined on $L^2(\Omega)$ verifying

$$U^*U = UU^* = I, \quad U e_j \in H_0^1(\Omega) \quad \text{for } j \geq 1, \quad (6)$$

and such that the family $u_j := U e_j$ is the family of normalized eigenfunctions of (1a) where V is given by (1b).

In this paper we give a variational formulation of the system (1a)-(2b), yielding a solution in terms of critical points of a real valued functional defined on a subset of the group of unitary operators. More precisely we define a subset \mathbb{S} of unitary operators on $L^2(\Omega)$ as

$$\mathbb{S} := \{U : L^2(\Omega) \longrightarrow L^2(\Omega) ; U \text{ satisfies (6)}\}, \quad (7)$$

and then we define a functional J on \mathbb{S} by setting

$$J_0(U) := \sum_{m \geq 1} \rho_m \int_{\Omega} \left(|\nabla u_m(x)|^2 + \tilde{V}_0(x) u_m^2(x) \right) dx, \quad (8)$$

and then

$$J(U) := J_0(U) + \frac{1}{2} \int_{\Omega} |\nabla V[U](x)|^2 dx, \quad (9)$$

where $u_m := Ue_m$ and $V[U]$ is the solution of (1b). The main purpose of this paper is to show that critical points of J on \mathbb{S} yield solutions of the Schrödinger–Poisson system and that the minimum of J is achieved on \mathbb{S} .

It is clear that in order to define the potential $V[U]$ and the functional J on the manifold \mathbb{S} some conditions must be imposed on the sequence $(\rho_m)_{m \geq 1}$.

Our main result concerning the system of equations (1a)–(2b) is the following (regarding the condition $1 \leq N \leq 6$ see Remark 9 below):

Theorem 1. *Let $\Omega \subset \mathbb{R}^N$ be a bounded domain and $1 \leq N \leq 6$. Assume that the sequence $(\rho_m)_{m \geq 1}$ satisfies*

$$\rho_m > 0, \quad \sum_{m \geq 1} m^{2/N} \rho_m < \infty, \quad (10)$$

and

$$\tilde{V}_0^+ \in L^1(\Omega), \quad \tilde{V}_0^- \in L^{p_0}(\Omega),$$

for some $p_0 > N/2$ (or $p_0 = 1$ if $N = 1$). Then the Schrödinger–Poisson system of equations (1a)–(2b) has a solution obtained as the minimum over \mathbb{S} of the functional J defined in (9).

In order to give a clear exposition of our global approach to the determination of a system of eigenvectors in terms of unitary operators U , in this introduction we give an outline of our approach, getting rid of technicalities inherent to an infinite dimensional Hilbert space, and to nonlinear problems.

Thus, in a first step, assume that we are given a finite dimensional (complex) Hilbert space H of dimension $n \geq 2$, its scalar product being denoted by $(\cdot|\cdot)$. If $A : H \rightarrow H$ is a self-adjoint, nonnegative operator (matrix), our aim is to define a procedure in which all the eigenvectors of A are determined at once, to compare with a step by step construction of eigenvalues (and eigenvectors) through the construction of critical values of the Rayleigh quotient

$$\frac{(Au|u)}{(u|u)}, \quad u \neq 0,$$

by a min-max procedure. To this end, for $u_1, \dots, u_n \in H$ let us define the functional

$$F(u_1, \dots, u_n) := \sum_{j=1}^n \rho_j (Au_j|u_j),$$

where, as above, we assume that the coefficients ρ_j verify $\rho_j > 0$, and also, for the sake of simplicity of exposition (see below for the general case), here assume moreover that

$$\rho_i \neq \rho_j \quad \text{for } i \neq j.$$

Define the subset (or manifold) $S \subset H^n$ by

$$S := \{ (u_1, \dots, u_n) \in H^n ; (u_i | u_j) = \delta_{ij} \}$$

We claim that upon maximizing or minimizing F on the manifold $S \subset H^n$, all the eigenvectors of A can be determined (as a matter of fact, any critical point of F yields such a result).

We begin by observing that an orthonormal basis e_1, \dots, e_n of H being given once and for all, the manifold S can be identified with the set of unitary operators U on H such that $U^*U = UU^* = I$, where I is the identity operator on H : indeed it is enough to see u_j as the j -th column of the matrix representation of U , that is to set $u_j := Ue_j$. Then we see that $(Au_j | u_j) = (U^*AUe_j | e_j)$, and denoting by D the diagonal matrix $D := \text{diag}(\rho_1, \dots, \rho_n)$, that is the matrix defined by $De_j = \rho_j e_j$, we check easily that

$$F(u_1, \dots, u_n) = \sum_{j \geq 1} \rho_j (Au_j | u_j) = \sum_{j \geq 1} \rho_j (U^*AUe_j | e_j) = \text{tr}(DU^*AU),$$

where $\text{tr}(B)$ denotes the trace of the operator (or matrix) B . Finally, considering for instance the minimization of F , this can be reformulated in the following way:

$$\text{minimize } J(U) := \text{tr}(DU^*AU) \text{ under the constraint } U^*U = I.$$

Clearly J is C^∞ (in fact analytic) and positive on the set

$$\mathbf{U}(n) := \mathbf{U}(n, H) := \{U : H \longrightarrow H ; U^*U = I\},$$

which is a smooth and compact manifold: therefore J achieves its minimum at some point $U_0 \in \mathbf{U}(n)$. Now we have to show that the vectors $u_j := U_0 e_j$ are indeed the eigenvectors of A .

Let $M : H \longrightarrow H$ be skew-adjoint (that is $M^* = -M$) and consider the one parameter group $U(t) = \exp(tM)U_0$ for $t \in \mathbb{R}$; note that since $M^* = -M$, one has $\exp(tM)^* = \exp(-tM)$ and thus one checks easily that

having $U_0^*U_0 = I$, then for all $t \in \mathbb{R}$ one has $U(t) \in \mathbf{U}(n)$, and consequently $J(U_0) \leq J(U(t))$ for all $t \in \mathbb{R}$. Now

$$J(U(t)) = \operatorname{tr}(DU_0^* \exp(-tM)A \exp(tM)U_0) \quad \text{and} \quad \left. \frac{dJ(U(t))}{dt} \right|_{t=0} = 0,$$

so that, after a straightforward calculation, we obtain

$$\begin{cases} \text{for all } M \text{ such that } M^* = -M, \text{ we have} \\ \operatorname{tr}(DU_0^*MAU_0) = \operatorname{tr}(DU_0^*AMU_0). \end{cases}$$

Setting

$$B := U_0DU_0^*,$$

and using the fact that for two given matrices K, L we have $\operatorname{tr}(KL) = \operatorname{tr}(LK)$, we observe that $\operatorname{tr}(DU_0^*MAU_0) = \operatorname{tr}(MAB)$, and that

$$\operatorname{tr}(DU_0^*AMU_0) = \operatorname{tr}(BAM) = \operatorname{tr}(MBA).$$

Summing up, we conclude that

$$\text{for all } M \text{ such that } M^* = -M, \quad \text{we have} \quad \operatorname{tr}(M(AB - BA)) = 0.$$

Taking $M := (AB - BA)^*$, we conclude that $BA = AB$, that is

$$U_0DU_0^*A = AU_0DU_0^*.$$

Applying this equality to the vector $u_j := U_0e_j$, and taking into account the definition of the diagonal operator D , we obtain (recall that $U_0^*U_0 = I$)

$$(U_0DU_0^*)Au_j = AU_0DU_0^*u_j = AU_0De_j = \rho_jAU_0e_j = \rho_jAu_j,$$

that is $(U_0DU_0^*)Au_j = \rho_jAu_j$, which means that Au_j is an eigenvector of $U_0DU_0^*$. This implies that $D(U_0^*Au_j) = \rho_j(U_0^*Au_j)$, and we see that $U_0^*Au_j$ is an eigenvector of D for the eigenvalue ρ_j , which is a simple eigenvalue of D , corresponding to the eigenvector e_j . This means that there exists $\lambda_j \in \mathbb{C}$ such that $U_0^*Au_j = \lambda_j e_j$, that is

$$Au_j = \lambda_j u_j.$$

As a matter of fact one sees that $\lambda_j \in \mathbb{R}$, while u_j is an eigenvector of A and U_0 is a diagonalization operator for A , which consists in the matrix whose columns are the eigenvectors u_j . Actually this procedure allows us to construct all the eigenvectors of A through the minimization of a unique

functional defined on the group $\mathbf{U}(n)$. Also, since $F(U_0) = \sum_{j \geq 1} \lambda_j \rho_j$, one easily sees that different choices in ordering the numbers ρ_j yield different ordering of eigenvalues and eigenvectors of A : for instance one may check that if the ρ_j 's are decreasing, that is if $\rho_j > \rho_{j+1}$ for $1 \leq j \leq n-1$, then one obtains the eigenvalues of A in a non decreasing order, that is $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. While if $\rho_j < \rho_{j+1}$ for $1 \leq j \leq n-1$, then one obtains the eigenvalues in a non increasing order, that is $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. (Had we began by maximizing J , the conclusion would be somehow reversed but analogous: if the ρ_j 's are decreasing then the λ_j 's would be non increasing).

In the next section of this paper we will show that, for a certain class of self-adjoint operators A , the eigenvectors and eigenvalues of A can be obtained through the minimization of the functional

$$J_0(U) := \text{tr}(DU^*AU)$$

on an appropriate subset of unitary operators U : this is precisely stated and proved in section 2. In section 3 we gather a certain number of preliminary results used in section 4, after stating the assumptions on the domain Ω and on the sequence $(\rho_m)_m$, we prove Theorem 1, as well as slightly more general variants of the Schrödinger–Poisson systems (see Theorem 15 in section 4). In section 5 we shall discuss some generalizations and state a few remarks about the results presented here.

2 Global determination of eigenvectors and eigenvalues

In this section we consider an infinite dimensional, separable, complex Hilbert space H whose scalar product is denoted by $(\cdot|\cdot)$ and its norm by $\|\cdot\|$. We shall make the following assumptions:

Hypothesis 1. We assume that $(A, D(A))$ is a densely defined, selfadjoint positive operator acting on the Hilbert space H , and that the domain $D(A)$ equipped with its graph norm is compactly imbedded in H , so that A has a compact resolvent and A possesses a sequence of eigenvalues $(\mu_j)_{j \geq 1}$ such that $0 \leq \mu_j < \mu_{j+1}$, each eigenvalue having finite multiplicity $m_j \geq 1$, and $\mu_j \rightarrow +\infty$ as $j \rightarrow \infty$, H being infinite dimensional.

We denote by $D(A^{1/2})$ the domain of $A^{1/2}$, that is the subspace of H obtained upon the completion of $D(A)$ with the scalar product $(u, v) \mapsto (u|v) + (Au|v)$, and we recall that $D(A^{1/2})$ is dense in H . Hence we can introduce the next assumption:

Hypothesis 2. We consider a fixed Hilbert basis of H , denoted by $(e_j)_{j \geq 1}$, such that $e_j \in D(A^{1/2})$ for each $j \geq 1$.

With the Hilbert basis $(e_j)_{j \geq 1}$ given by hypothesis 2, we consider a sequence $(\rho_j)_{j \geq 1}$ of real numbers such that

$$\rho_j > 0, \quad \sum_{j \geq 1} \rho_j \|e_j\|_{D(A^{1/2})}^2 < \infty, \quad (11)$$

and we denote by D the diagonal operator defined by

$$De_j := \rho_j e_j, \quad \text{for } j \geq 1. \quad (12)$$

Note that since H is infinite dimensional and A has a compact resolvent, while $e_j \in D(A^{1/2})$, we have $\|e_j\|_{D(A^{1/2})} \rightarrow \infty$ as $j \rightarrow \infty$. Indeed, otherwise, the sequence $(e_j)_j$ would be bounded in $D(A^{1/2})$, and the imbedding $D(A^{1/2}) \subset H$ being compact, one would extract a subsequence $(e_{j_k})_{k \geq 1}$ such that $e_{j_k} \rightharpoonup f$ in $D(A^{1/2})$ and $e_{j_k} \rightarrow f$ strongly in H ; in particular $\|f\| = 1$, since $(e_j)_j$ is a Hilbert basis of H . But we have also $e_j \rightarrow 0$ in H , and thus we should have $f = 0$. This contradiction shows that $(e_j)_j$ cannot contain any bounded sequence in $D(A^{1/2})$. As a consequence we have $\rho_j \rightarrow 0$ and D is a compact operator.

Next we shall consider unitary operators $U : H \rightarrow H$ which satisfy the following condition (this expresses the fact that the operator DU^*AU is of trace class, see M. Reed & B. Simon [14], volume 1, section VI.6)

$$\begin{cases} U^*U = UU^* = I, & Ue_j \in D(A^{1/2}) \text{ for } j \geq 1, \\ \sum_{j \geq 1} \rho_j (U^*AUe_j | e_j) < \infty. \end{cases} \quad (13)$$

and we define the set \mathbb{S} through

$$\mathbb{S} := \{U : H \rightarrow H ; U \text{ satisfies (13)}\}. \quad (14)$$

Remark 2. Let us point out that such operators U exist, that is \mathbb{S} is not empty: indeed, not only for $t \in \mathbb{R}$ we have $e^{itA} \in \mathbb{S}$, but also for any $\lambda > 0$, the operator U_λ (the so-called Cayley transform of λA , see K. Yosida [17]) defined by

$$U_\lambda := (I + i\lambda A)(I - i\lambda A)^{-1}$$

is a bounded operator on H and one checks easily that

$$U_\lambda^* = (I - i\lambda A)(I + i\lambda A)^{-1},$$

so that $U_\lambda^* U_\lambda = I$. Moreover, for any $f \in D(A^{1/2})$ we have $(I + i\lambda A)^{-1} f \in D(A^{3/2})$, and thus $U_\lambda f \in D(A^{1/2})$. As a matter of fact, not only U_λ is a unitary operator on H , but one has also $\|U_\lambda f\|_{D(A^{1/2})} = \|f\|_{D(A^{1/2})}$. Therefore, since the sequence $(\rho_j)_j$ satisfies (11), one sees that U_λ satisfies (13) and $U_\lambda \in \mathbb{S}$. \square

For a unitary operator $U : H \rightarrow H$ satisfying (13), we define $J_0(U)$ by

$$J_0(U) := \operatorname{tr}(DU^*AU) := \sum_{j \geq 1} \rho_j(U^*AUe_j|e_j) \quad (15)$$

The following result concerns eigenvectors of A :

Theorem 3. *Assume that the hypotheses 1 and 2, as well as condition (11) are satisfied. Then the functional J_0 defined in (15) achieves its minimum on \mathbb{S} defined by (14). Then there exists $\widehat{U}_0 \in \mathbb{S}$ such that*

$$J_0(\widehat{U}_0) = \min_{U \in \mathbb{S}} J_0(U),$$

and for each $j \geq 1$, the vector $\varphi_j := \widehat{U}_0 e_j$ is an eigenvector of A corresponding to the eigenvalue $\lambda_j := (A\varphi_j|\varphi_j)$.

Remark 4. It is clear that the eigenvalues $(\lambda_j)_{j \geq 1}$ are independent of the choice of the sequence $(\rho_j)_{j \geq 1}$. However, as pointed out in the introduction, if one assumes that the sequence $(\rho_j)_{j \geq 1}$ is decreasing, then the eigenvalues $(\lambda_j)_{j \geq 1}$ are ordered in a non decreasing order, that is $\lambda_j \leq \lambda_{j+1}$ for $j \geq 1$. Thus for different choices of the sequence $(\rho_j)_{j \geq 1}$ one may obtain different diagonalization operators \widehat{U}_0 for A . \square

We split the proof of this result into a couple of lemmas.

Lemma 5. *The functional J_0 achieves its minimum on \mathbb{S} at a certain $U_0 \in \mathbb{S}$.*

Proof. Indeed consider the infimum $\alpha := \inf_{U \in \mathbb{S}} J_0(U)$. Since $\mathbb{S} \neq \emptyset$, we have $0 \leq \alpha < \infty$. Consider a minimizing sequence $(U_n)_{n \geq 1} \in \mathbb{S}$, such that for instance $\alpha \leq J(U_n) \leq \alpha + 1/n$. Then for each fixed $j \geq 1$, setting $u_j^n := U_n e_j$, we have for all $n \geq 1$

$$\|u_j^n\|_{D(A^{1/2})}^2 = 1 + (Au_j^n|u_j^n) \leq 1 + \frac{\alpha + 1}{\rho_j}.$$

Thus, since the inclusion $D(A^{1/2}) \subset H$ is compact, upon extracting subsequences through Cantor's diagonal scheme, and denoting again this diagonal subsequence by $(u_j^n)_n$, we may assume that for all $j \geq 1$ there exist a family $(u_j)_j$ such that for $j \geq 1$ fixed

$$u_j^n \rightharpoonup u_j \text{ weakly in } D(A^{1/2}), \quad u_j^n \rightarrow u_j \text{ strongly in } H$$

as $n \rightarrow \infty$. Setting $U_0 e_j := u_j$, one checks easily that U_0 can be extended by linearity to the subspace $\text{span}\{e_j ; j \geq 1\}$, and that for $f \in \text{span}\{e_j ; j \geq 1\}$ we have

$$\|U_0 f\|^2 = \lim_{n \rightarrow \infty} \|U_n f\|^2 = \|f\|^2.$$

In other words U_0 is a unitary operator on (the algebraic) $\text{span}\{e_j ; j \geq 1\}$, and therefore can be extended as such to the whole space H . Since for any $m \geq 1$ we have

$$\sum_{j=1}^m \rho_j(Au_j | u_j) \leq \liminf_{n \rightarrow \infty} \sum_{j=1}^m \rho_j(Au_j^n | u_j^n) \leq \liminf_{n \rightarrow \infty} J_0(U_n) = \alpha,$$

upon letting $m \rightarrow \infty$ we conclude that $J_0(U_0) \leq \alpha$. Thus, having $U_0^* U_0 = I$ and $U_0 e_j \in D(A^{1/2})$ for all $j \geq 1$, and $J_0(U_0) < \infty$, we have $U_0 \in \mathbb{S}$ and $J_0(U_0) = \alpha$. \square

Next we show that U_0 , given by Lemma 5 is a diagonalization operator for A .

Lemma 6. *Under the assumptions of Theorem 3, let U_0 be given by Lemma 5, and set $u_j := U_0 e_j$, for $j \geq 1$.*

(i) *Assume that $k \geq 1$ is such that*

$$\rho_\ell \neq \rho_k \quad \text{for } \ell \neq k. \tag{16}$$

Then there exist $\lambda_k \in \mathbb{R}_+$ such that $Au_k = \lambda_k u_k$.

(ii) *Assume that $k \geq 1$ is such that for some $m \geq 2$*

$$\begin{cases} \rho_k = \rho_{k+\ell} & \text{for } 0 \leq \ell \leq m-1, \\ \rho_k \neq \rho_n & \text{for } n \notin \{k+\ell ; 0 \leq \ell \leq m-1\}. \end{cases} \tag{17}$$

Then there exists a unitary transformation U_k of the m -dimensional space $H_k := \text{span}\{U_0 e_{k+\ell} ; 0 \leq \ell \leq m-1\}$ such that if

$$\widehat{u}_{k+\ell} := U_k U_0 e_{k+\ell},$$

then there exists $\lambda_{k+\ell} \in \mathbb{R}_+$ such that $A\widehat{u}_{k+\ell} = \lambda_{k+\ell} \widehat{u}_{k+\ell}$ for $0 \leq \ell \leq m-1$.

Proof. First let $M : H \rightarrow H$ be a bounded skewadjoint operator such that $M : D(A^{1/2}) \rightarrow D(A^{1/2})$ is also bounded. Indeed such operators do exist (consider for instance $i(I + \lambda A)^{-1}$ for $\lambda > 0$). Setting $U(t) := \exp(-tM)U_0$ for $t \in \mathbb{R}$, one checks easily that, since $M^* = -M$, one has $U(t) \in \mathbb{S}$ for all t , and thus the function $g(t) := J_0(U(t))$ is well defined, is of class C^1 and achieves its minimum at $t = 0$. However since

$$g(t) = \text{tr}(DU_0^* \exp(tM)A \exp(-tM)U_0),$$

one concludes that

$$g'(0) = \text{tr}(DU_0^* MAU_0) - \text{tr}(DU_0^* AMU_0) = 0 \quad (18)$$

for all bounded operators $M : H \rightarrow H$ such that $M^* = -M$ and M is also bounded from $D(A^{1/2})$ into itself. In the same way, if we consider a bounded operator $L : H \rightarrow H$ such that $L = L^*$ and L is also bounded from $D(A^{1/2})$ into itself, upon setting $M := iL$, we conclude that (18) yields

$$\text{tr}(DU_0^* LAU_0) = \text{tr}(DU_0^* ALU_0), \quad (19)$$

for all such operators L .

Note that the above relation (18) yields that

$$\begin{aligned} \sum_{j \geq 1} \rho_j(U_0^* MAU_0 e_j | e_j) - \sum_{j \geq 1} \rho_j(U_0^* AMU_0 e_j | e_j) &= 0 \\ \sum_{j \geq 1} \rho_j(MAu_j | u_j) - \sum_{j \geq 1} \rho_j(AMu_j | u_j) &= 0, \end{aligned}$$

that is, since $M^* = -M$,

$$-\sum_{j \geq 1} \rho_j(Au_j | Mu_j) - \sum_{j \geq 1} \rho_j(AMu_j | u_j) = 0 \iff \text{Re} \sum_{j \geq 1} \rho_j(Au_j | Mu_j) = 0. \quad (20)$$

Analogously using (19) one obtains in the same way

$$\sum_{j \geq 1} \rho_j(Au_j | Lu_j) = \sum_{j \geq 1} \rho_j(ALu_j | u_j) \iff \text{Im} \sum_{j \geq 1} \rho_j(Au_j | Lu_j) = 0. \quad (21)$$

At this point, in a first step, assume that the integer k is such that condition (16) is fulfilled. Consider an integer $n \neq k$, so that $\rho_n \neq \rho_k$, and define the operators M and L in the following way

$$\begin{cases} Mu_k := u_n, & Mu_n := -u_k, \\ Lu_k := u_n, & Lu_n := u_k, \\ Lu_j = Mu_j = 0 & \text{for } j \notin \{k, n\}. \end{cases} \quad (22)$$

Clearly M and L satisfy the required conditions above, and using (20), with our choice of the operator M , we get $(\rho_n - \rho_k)\text{Re}(Au_k|u_n) = 0$, that is, since $\rho_k - \rho_n \neq 0$,

$$\text{Re}(Au_k|u_n) = 0.$$

Upon using (21), with our above choice of the operator L and the fact that $\rho_n - \rho_k \neq 0$, analogously we have that

$$\text{Im}(Au_k|u_n) = 0.$$

So, from the above two relations, we infer that $(Au_k|u_n) = 0$ for all n such that $\rho_n \neq \rho_k$, that is

$$Au_k \in \text{span}\{u_n ; n \neq k\}^\perp = \text{span}\{u_k\},$$

where we use the fact that the family $(u_j)_j$ is a Hilbert basis of H , being the image of the Hilbert basis $(e_j)_j$ under the unitary operator U_0 . This means that $Au_k = \lambda_k u_k$ for some $\lambda_k \in \mathbb{C}$, but since A is a nonnegative self-adjoint operator, as a matter of fact we have $\lambda_k \geq 0$.

Next assume that the integer k is such that the coefficient ρ_k has multiplicity $m \geq 2$, that is condition (17) is satisfied. Arguing as above, we consider the following operators M and L : for $n \notin \{k+j ; 0 \leq j \leq m-1\}$ and $0 \leq \ell \leq (m-1)$ fixed, set

$$\begin{cases} Mu_{k+\ell} := u_n, & Mu_n := -u_{k+\ell}, \\ Lu_{k+\ell} := u_n, & Lu_n := u_{k+\ell} \\ Lu_j = Mu_j = 0 & \text{for all } j \notin \{n, k+\ell\}. \end{cases} \quad (23)$$

Then, proceeding as above, we conclude that $(Au_{k+\ell}|u_n) = 0$ for all $n \notin \{k+j ; 0 \leq j \leq m-1\}$, that is:

$$Au_{k+\ell} \in (\text{span}\{u_n ; n \neq k+j, 0 \leq j \leq m-1\})^\perp$$

that is

$$Au_{k+\ell} \in \text{span}\{u_{k+i} ; 0 \leq i \leq m-1\}.$$

This means that if we set $H_k := \text{span}\{u_{k+i} ; 0 \leq i \leq m-1\}$, then $A : H_k \rightarrow H_k$ is a self-adjoint operator on the finite dimensional space H_k . Therefore there exists a unitary operator U_k , acting on this space, such that if for $0 \leq \ell \leq m-1$ we set $\hat{u}_{k+\ell} = U_k u_{k+\ell} = U_k U_0 e_{k+\ell}$, we have $A\hat{u}_{k+\ell} = \lambda_{k+\ell} \hat{u}_{k+\ell}$ for some $\lambda_{k+\ell} \geq 0$. \square

As we may see from the above analysis, when all the ρ_j 's are distinct, then U_0 , any unitary operator which minimizes J_0 , is a diagonalization operator for A . However in the general case, when some of the coefficients ρ_k have multiplicity $m_k \geq 2$, it is possible that one has to impose a unitary transformation U_k in the space

$$H_k := \text{span}\{U_0 e_{k+\ell} ; 0 \leq \ell \leq m_k - 1\}$$

in order to have the operator A diagonalized. In other words, one may find a unitary operator U_k on H_k such that if $A_k := A|_{H_k}$ is the trace of A on H_k , the operator $U_k^* A_k U_k$ is diagonal. Thus since $\rho_k = \rho_{k+\ell}$ for $0 \leq \ell \leq m_k - 1$, if we denote by \widehat{U} the unitary operator obtained through the composition of all such operators U_k and U_0 , one has $J_0(\widehat{U}) = J_0(U_0)$. More precisely, we can state the following corollary, which ends the proof of Theorem 3:

Corollary 7. *Under the assumptions of Theorem 3, let U_k be given by Lemma 6 when $k \geq 1$ is such that (17) is satisfied. Then the operator \widehat{U}_0 defined by $\widehat{U}_0 e_k = U_0 e_k$ when k satisfies (16), and*

$$\widehat{U}_0 e_{k+\ell} := U_k U_0 e_{k+\ell}, \quad \text{for } 0 \leq \ell \leq m_k - 1, \quad \text{when (17) is satisfied,}$$

belongs to \mathbb{S} , while $J_0(\widehat{U}_0) = J_0(U_0)$ and $\widehat{U}_0^ A \widehat{U}_0$ is diagonal. Setting $\varphi_j := \widehat{U}_0 e_j$ for $j \geq 1$, then there exists $\lambda_j \geq 0$ such that $A\varphi_j = \lambda_j \varphi_j$.*

Remark 8. Regarding the finite dimensional case, after the completion of this work and its submission for publication, R.V. Kohn, in a private communication, pointed to one of the authors (O.K.) that L. Mirsky [6], develops a result of J. von Neumann [7] stating that

$$|\text{tr}(AB)| \leq \sum_{j=1}^n \sigma_j(A) \sigma_j(B)$$

for two $n \times n$ matrices A, B , where $(\sigma_j(M))_{1 \leq j \leq n}$ denotes the decreasing singular values of a matrix M . From this L. Mirsky concludes that actually one has

$$\sup_{U, V} |\text{tr}(BUAV)| = \sum_{j=1}^n \sigma_j(A) \sigma_j(B),$$

which is another result due to J. von Neumann. In this respect, in the finite dimensional case our result can be compared to the above result in the particular case of self-adjoint matrices, and moreover the approach given here characterizes the diagonalization matrix by a variational method. \square

3 Preliminary results for Schrödinger–Poisson system

In this section we prove an existence result regarding the system of equations (1a)–(2b). We shall assume that

$$\Omega \subset \mathbb{R}^N \text{ is a bounded domain and that } N \leq 6, \quad (24)$$

and we endow the (complex) space $L^2(\Omega)$ with its scalar product denoted by $(\cdot|\cdot)$ and its norm $\|\cdot\|$. Let \tilde{V}_0 be a real valued potential such that

$$\tilde{V}_0^+ \in L^1(\Omega), \quad \tilde{V}_0^- \in L^{p_0}(\Omega) \text{ for some } p_0 > \frac{N}{2} \text{ and } p_0 \geq 1. \quad (25)$$

Then we define an unbounded operator $(A, D(A))$ by setting

$$D(A) := \left\{ u \in H_0^1(\Omega) ; -\Delta u + \tilde{V}_0 u \in L^2(\Omega) \right\}, \quad Au := -\Delta u + \tilde{V}_0 u. \quad (26)$$

This operator is self-adjoint, has a compact resolvent, and there exists a Hilbert basis of eigensystem denoted by $(\lambda_m, \varphi_m)_{m \geq 1}$, that is (here δ_{mn} being the Kronecker symbol)

$$-\Delta \varphi_m + \tilde{V}_0 \varphi_m = \lambda_m \varphi_m, \quad \varphi_m \in H_0^1(\Omega), \quad \int_{\Omega} \varphi_m(x) \varphi_n(x) dx = \delta_{mn}.$$

It is well-known that by Weyl's theorem there exist two positive constants c_1, c_2 , depending on Ω and \tilde{V}_0 , and $m_0 \geq 1$ large enough such that for all integers $m \geq m_0$ one has

$$c_1 m^{2/N} \leq \lambda_m \leq c_2 m^{2/N}.$$

(See for instance [3], chapter 5, § 3, where the case of Neumann boundary conditions is also treated). For this reason, as far as the sequence $(\rho_m)_{m \geq 1}$ is concerned, in order to ensure the finiteness of the functionals we are going to minimize, we assume that the growth condition (10) is satisfied.

For a given unitary operator $U : L^2(\Omega) \rightarrow L^2(\Omega)$ such that moreover $U : H_0^1(\Omega) \rightarrow H_0^1(\Omega)$ is also a bounded operator, we shall denote by $V := V[U]$ the potential defined by the Poisson equation (here $|U\varphi_j|$ denotes the modulus of the function $U\varphi_j$)

$$-\Delta V = \sum_{j \geq 1} \rho_j |U\varphi_j|^2 \text{ in } \Omega, \quad V \in H_0^1(\Omega). \quad (27)$$

Remark 9. We wish to explain here the limitation $N \leq 6$ in Theorem 1. Indeed, thanks to the Sobolev imbedding theorem we have $H_0^1(\Omega) \subset L^{2^*}(\Omega)$, where $2^* = 2N/(N-2)$ when $N \geq 3$, while 2^* can be any finite exponent if $N = 2$, and $2^* = \infty$ if $N = 1$. This means that the right hand side of the above equation (27), that is

$$f := \sum_{j \geq 1} \rho_j |U \varphi_j|^2,$$

belongs to $L^q(\Omega)$ where $q := N/(N-2)$ if $N \geq 3$, or $q < \infty$ arbitrary if $N = 2$, or $q = \infty$ if $N = 1$. Since $q \geq (2^*)' := 2N/(N+2)$ if and only if $N \leq 6$, we conclude that for such N 's we have $f \in H^{-1}(\Omega) \cap L^{2N/(N+2)}(\Omega)$: this ensures that equation (27) has a unique solution $V := V[U] \in H_0^1(\Omega)$ when $N \leq 6$ and thus the functional

$$U \mapsto J_1(U) := \frac{1}{2} \int_{\Omega} |\nabla V[U]|^2 dx = \frac{1}{2} \sum_{j \geq 1} \rho_j \int_{\Omega} V[U] |U \varphi_j|^2 dx$$

is well defined (this functional J_1 is used in the minimization procedure, see below). Moreover while for $N \leq 5$ one may show that J_1 is, in an appropriate sense, weakly sequentially continuous, the case $N = 6$ is a limit case and we are only able to prove that J_1 is weakly sequentially lower semi-continuous. Thus for $N \leq 6$ we shall show that J_1 is weakly sequentially lower semi-continuous, and this allows us to proceed with our minimization procedure.

Finally, when $N \geq 7$ the functional J_1 is not well defined for an arbitrary unitary operator $U : L^2(\Omega) \rightarrow L^2(\Omega)$ which is also a bounded operator on $H_0^1(\Omega)$, and the method we use here has to be modified by considering other classes of operators U . \square

Remark 10. Observe that if $V[U]$ satisfies (27), then by the maximum principle we have $V[U] > 0$ in Ω . Note also that when $N \leq 3$, we have $f \in L^q(\Omega)$ for some $q > N/2$. Then a classical regularity result (see for instance [4], or G. Stampacchia [16]) states that there exists a constant $c > 0$ such that if $V \in H_0^1(\Omega)$ satisfies $-\Delta V = f$ and $f \in L^q(\Omega)$, then

$$\|V\|_{\infty} \leq c \|f\|_q. \quad (28)$$

Therefore, if $V := V[U]$ is given by (27) we have $V \in L^{\infty}(\Omega)$ when $N \leq 3$.

When $N = 4$, we have $f \in L^2(\Omega)$, and in this case, since $N/2 = 2$, we have that $V \in L^p(\Omega)$ for all $p < \infty$. Moreover for a constant $c(p)$ depending on Ω we have:

$$\|V\|_p \leq c(p) \|f\|_2. \quad (29)$$

Finally, when $5 \leq N \leq 6$, we have $f \in L^q(\Omega)$ with $1 < q = N/(N-2) < N/2$: using a regularity result, we have $V \in L^p(\Omega)$ where p is given by

$$\frac{1}{p} = \frac{1}{q} - \frac{2}{N},$$

that is $p := N/(N-4)$ (again cf. [4], or G. Stampacchia [16]). In this case there exists a constant $c > 0$ depending on Ω such that

$$\|V\|_{N/(N-4)} \leq c \|f\|_{N/(N-2)}. \quad (30)$$

These observations will allow us to show that the functional J_1 is, in an appropriate sense, weakly sequentially lower semi-continuous (see below Lemma 13). \square

It is convenient to consider the Sobolev space \mathbb{H}_1 endowed with the norm $\|\cdot\|_{\mathbb{H}_1}$:

$$\mathbb{H}_1 := \left\{ u \in H_0^1(\Omega) ; \|u\|_{\mathbb{H}_1}^2 := \|\nabla u\|^2 + \int_{\Omega} \tilde{V}_0^+(x)|u(x)|^2 dx < \infty \right\}.$$

The imbedding $\mathbb{H}_1 \subset L^2(\Omega)$ is compact. Note that since the eigenfunctions φ_m belong to $L^\infty(\Omega)$, we have $\varphi_m \in \mathbb{H}_1$.

Regarding the manifold \mathbb{S} defined in (7), we have to modify it slightly, as we did in section § 2. More precisely we shall consider unitary operators $U : L^2(\Omega) \rightarrow L^2(\Omega)$ such that

$$U^*U = UU^* = I, \quad U\varphi_j \in \mathbb{H}_1 \quad \text{for } j \geq 1, \quad \sum_{j \geq 1} \rho_j(U^*AU\varphi_j|\varphi_j) < \infty, \quad (31)$$

and we consider the manifold defined by

$$\mathbb{S} := \{U : L^2(\Omega) \rightarrow L^2(\Omega) ; U \text{ satisfies (31)}\}. \quad (32)$$

We denote by D the diagonal operator acting on $L^2(\Omega)$ defined by $D\varphi_j = \rho_j\varphi_j$, and for $U \in \mathbb{S}$ we define the functionals J_0 and J_1 as follows:

$$J_0(U) := \text{tr}(DU^*AU) = \sum_{j \geq 1} \rho_j \int_{\Omega} \left(|\nabla U\varphi_j|^2(x) + \tilde{V}_0(x)|U\varphi_j|^2(x) \right) dx \quad (33)$$

and

$$\begin{aligned} J_1(U) &:= \frac{1}{2} \int_{\Omega} |\nabla V[U]|^2(x) dx \\ &= \left\langle -\frac{1}{2} \Delta V[U], V[U] \right\rangle \\ &= \frac{1}{2} \sum_{j \geq 1} \rho_j (V[U]U\varphi_j|U\varphi_j) = \frac{1}{2} \text{tr}(DU^*V[U]U), \end{aligned} \quad (34)$$

where, in the last equality of (34), by an abuse of notation, we denote by $V[U]$ the (linear) multiplication operator $f \mapsto V[U]f$. Since we assume $N \leq 6$, as explained in the above Remark 9, the functional J_1 is well defined on \mathbb{S} .

Note that here the potential \tilde{V}_0 may have a negative part, so at some point we will need to ensure that the functional J_0 is bounded below, that it is *coercive* in some sense. More precisely we have:

Lemma 11. *There exists $C \geq 0$ such that for any $U \in \mathbb{S}$ one has*

$$J_0(U) \geq \frac{1}{2} \sum_{j \geq 1} \rho_j \int_{\Omega} \left(|\nabla U \varphi_j|^2 + 2\tilde{V}_0^+ |U \varphi_j|^2 \right) dx - C.$$

Proof. Assume that $N \geq 3$ (the case $N \leq 2$ can be handled in a similar way). For $t > 0$ and $u \in H_0^1(\Omega)$ such that $\|u\| = 1$ we have

$$\begin{aligned} \int_{\Omega} \tilde{V}_0^- |u|^2 dx &= \int_{[\tilde{V}_0^- > t]} \tilde{V}_0^- |u|^2 dx + \int_{[\tilde{V}_0^- \leq t]} \tilde{V}_0^- |u|^2 dx \\ &\leq \int_{\Omega} 1_{[\tilde{V}_0^- > t]} \tilde{V}_0^- |u|^2 dx + t \int_{\Omega} |u|^2 dx \\ &\leq \|1_{[\tilde{V}_0^- > t]} \tilde{V}_0^-\|_{L^{N/2}} \|u\|_{L^{2^*}}^2 + t \\ &\leq C_1(N) \text{meas}([\tilde{V}_0^- > t])^{\theta} \|\tilde{V}_0^-\|_{L^{p_0}} \|\nabla u\|^2 + t \end{aligned}$$

where we have used Hölder's inequality twice (once with $N/(N-2)$ and $(N/(N-2))' = N/2$, once with p_0 and $N/2$, where $\theta = (2/N) - (1/p_0) = (2/3) - (1/p_0) > 0$). We used also Sobolev's inequality $\|u\|_{2^*} \leq C \|\nabla u\|$. Now, since $\tilde{V}_0^- \in L^{p_0}(\Omega)$, we know that $\text{meas}([\tilde{V}_0^- > t]) \rightarrow 0$ as $t \rightarrow +\infty$. We choose $t > 0$ large enough to ensure that

$$C_1(N) \text{meas}([\tilde{V}_0^- > t])^{\theta} \|\tilde{V}_0^-\|_{L^{p_0}} \leq \frac{1}{2}.$$

Then we have for all $u \in \mathbb{H}_1$

$$\int_{\Omega} |\nabla u|^2 dx + \int_{\Omega} \tilde{V}_0 |u|^2 dx \geq \frac{1}{2} \|\nabla u\|^2 + \int_{\Omega} \tilde{V}_0^+ |u|^2 dx - t.$$

Applying this to $u := U \varphi_j$, multiplying by $\rho_j > 0$ and calculating the sum over j yields the inequality claimed by our lemma, with $C := t \sum_{j \geq 1} \rho_j$. \square

It is well known that the fact that the functional $u \mapsto \|\nabla u\|^2$ is weakly sequentially lower semi-continuous (l.s.c.) on $H_0^1(\Omega)$ plays a crucial role in many minimization problems. Regarding the functional J_0 we need an analogous property which is stated below:

Lemma 12. *The functional J_0 is “weakly sequentially lower semi-continuous” in the following sense: let $(U_n)_{n \geq 1}$ be a sequence in \mathbb{S} such that for some $R > 0$ and all $n \geq 1$ one has $J_0(U_n) \leq R$. Then there exists a subsequence $(U_{n_k})_k$ such that for any fixed $j \geq 1$ one has $U_{n_k} \varphi_j \rightharpoonup u_j$ in \mathbb{H}_1 as $k \rightarrow +\infty$, and if we set $u_j^{n_k} := U_{n_k} \varphi_j$ and we define a linear operator U by setting $U \varphi_j := u_j$ we have $u_j^{n_k} \rightarrow u_j$ strongly in $L^2(\Omega)$ and almost everywhere on Ω , and*

$$\lim_{k \rightarrow \infty} \int_{\Omega} \tilde{V}_0^- |u_j^{n_k}|^2 dx = \int_{\Omega} \tilde{V}_0^- |u_j|^2 dx, \quad U \in \mathbb{S}, \quad J_0(U) \leq \liminf_{k \rightarrow \infty} J_0(U_{n_k}).$$

Proof. Assume $N \geq 3$ (the case $N \leq 2$ being analgous). Thanks to Lemma 11, we know that

$$\sum_{j \geq 1} \rho_j \int_{\Omega} (|\nabla U_n \varphi_j|^2 + 2\tilde{V}_0^+ |U_n \varphi_j|^2) dx \leq 2R + 2C =: C_1.$$

This implies that for each $j \geq 1$ fixed the sequence $(u_j^n)_n := (U_n \varphi_j)_n$ is bounded in \mathbb{H}_1 , more precisely $\|u_j^n\|_{\mathbb{H}_1}^2 \leq C/\rho_j$. By using Cantor’s diagonal scheme and the compactness of the imbedding $\mathbb{H}_1 \subset L^2(\Omega)$, we may extract a subsequence denoted by $(u_j^{n_k})_{k \geq 1}$ such that

$$\begin{cases} u_j^{n_k} \rightharpoonup u_j & \text{weakly in } \mathbb{H}_1, \\ u_j^{n_k} \rightarrow u_j & \text{strongly in } L^2(\Omega), \\ u_j^{n_k} \rightarrow u_j & \text{a.e. in } \Omega, \end{cases}$$

as $k \rightarrow \infty$. For any $m \geq 1$ fixed, we have

$$\sum_{j=1}^m \rho_j \|u_j\|_{\mathbb{H}_1}^2 \leq \liminf_{k \rightarrow \infty} \sum_{j=1}^m \rho_j \|u_j^{n_k}\|_{\mathbb{H}_1}^2 \leq \liminf_{k \rightarrow \infty} \sum_{j=1}^{\infty} \rho_j \|u_j^{n_k}\|_{\mathbb{H}_1}^2 \leq C,$$

and finally

$$\sum_{j=1}^{\infty} \rho_j \|u_j\|_{\mathbb{H}_1}^2 \leq \liminf_{k \rightarrow \infty} \sum_{j=1}^{\infty} \rho_j \|u_j^{n_k}\|_{\mathbb{H}_1}^2 \leq C. \quad (35)$$

Setting $U \varphi_j := u_j$, one checks easily that U can be extended by linearity to the subspace $\text{span}\{\varphi_j ; j \geq 1\}$, and that for $f \in \text{span}\{\varphi_j ; j \geq 1\}$ we have

$$\|Uf\|^2 = \lim_{n \rightarrow \infty} \|U_n f\|^2 = \|f\|^2.$$

In other words U is a unitary operator on (the algebraic) $\text{span}\{\varphi_j ; j \geq 1\}$, and therefore can be extended as such to the whole space $L^2(\Omega)$. Then (35) shows that $U \in \mathbb{S}$.

We note also that in particular we have

$$\sum_{j \geq 1} \rho_j \int_{\Omega} \tilde{V}_0^-(x) |u_j(x)|^2 dx < \infty. \quad (36)$$

Since we assume $N \geq 3$, the strong convergence of $u_j^{n_k} \rightarrow u_j$ in $L^2(\Omega)$ implies (through Hölder's inequality, or interpolation between $L^2(\Omega)$ and $L^{2N/(N-2)}(\Omega)$) that for any fixed $p < N/(N-2)$, and any $j \geq 1$ we have that $u_j^{n_k} \rightarrow u_j$ strongly in $L^{2p}(\Omega)$ and a.e. in Ω , and thus $|u_j^{n_k}|^2 \rightarrow |u_j|^2$ strongly in $L^p(\Omega)$. Since $\tilde{V}_0^- \in L^{p_0}(\Omega)$ and $p_0 > N/2$, taking $p := p'_0 = p_0/(p_0 - 1)$, so that $p < N/(N-2)$, we conclude first that

$$\lim_{k \rightarrow \infty} \int_{\Omega} \tilde{V}_0^- |u_j^{n_k}|^2 dx = \int_{\Omega} \tilde{V}_0^- |u_j|^2 dx,$$

and then thanks to (36) and the monotone convergence theorem,

$$\lim_{k \rightarrow \infty} \sum_{j \geq 1} \rho_j \int_{\Omega} \tilde{V}_0^- |u_j^{n_k}|^2 dx = \sum_{j \geq 1} \rho_j \int_{\Omega} \tilde{V}_0^- |u_j|^2 dx.$$

From this and (35) it is clear that

$$\begin{aligned} \sum_{j=1}^{\infty} \rho_j \int_{\Omega} (|\nabla u_j|^2 + \tilde{V}_0^+(x) |u_j|) dx &\leq \liminf_{k \rightarrow \infty} J_0(U_{n_k}) \\ &+ \lim_{k \rightarrow \infty} \sum_{j \geq 1} \rho_j \int_{\Omega} \tilde{V}_0^- |u_j^{n_k}|^2 dx \\ &\leq \liminf_{k \rightarrow \infty} J_0(U_{n_k}) \\ &+ \sum_{j \geq 1} \rho_j \int_{\Omega} \tilde{V}_0^- |u_j|^2 dx, \end{aligned}$$

which means that $J_0(U) \leq \liminf_{k \rightarrow \infty} J_0(U_{n_k})$, as claimed. \square

Regarding the functional J_1 we have the following result:

Lemma 13. *Let $1 \leq N \leq 6$. The functional J_1 is “weakly sequentially continuous” in the following sense: let $(U_n)_{n \geq 1}$ be a sequence in \mathbb{S} such that for some $R > 0$ and all $n \geq 1$ one has*

$$\sum_{j \geq 1} \rho_j \|U_n \varphi_j\|_{\mathbb{H}_1}^2 \leq R,$$

and such that for any fixed $j \geq 1$ one has

$$U_n \varphi_j \rightarrow u_j \text{ in } \mathbb{H}_1, \quad U_n \varphi_j \rightarrow u_j \text{ in } L^2(\Omega) \text{ and a.e. on } \Omega$$

as $n \rightarrow +\infty$. If we set $u_j^n := U_n \varphi_j$ and we define a linear operator U by setting $U \varphi_j := u_j$ we have

$$J_1(U) \leq \liminf_{n \rightarrow \infty} J_1(U_n).$$

Proof. This property is due to the positivity of the Green function associated to the Dirichlet problem. Indeed it is well-known that for $f \in L^p(\Omega)$ where $p := 2N/(N+2)$ if $N \geq 3$, or $p > 1$ if $N = 2$, or $p = 1$ if $N = 1$, the solution of the Dirichlet problem

$$-\Delta w = f \quad \text{in } \Omega, \quad w \in H_0^1(\Omega)$$

may be represented with the Green kernel

$$w(x) = \int_{\Omega} K(x, y) f(y) dy$$

where $K(x, y) > 0$ on $\Omega \times \Omega$. In particular we have $\|\nabla w\|^2 = \int_{\Omega} f(x) w(x) dx$, and thus

$$\|\nabla w\|^2 = \int_{\Omega \times \Omega} K(x, y) f(y) f(x) dx dy.$$

Applying this to $f := f^n := \sum_{j \geq 1} \rho_j |u_j^n|^2$ and $w := w^n := V[U_n]$, we see that

$$J_1(U_n) = \frac{1}{2} \sum_{j, k \geq 1} \rho_j \rho_k \int_{\Omega \times \Omega} K(x, y) |u_j^n(x)|^2 |u_k^n(y)|^2 dx dy.$$

Since and $K(x, y) > 0$ and $u_j^n(x) \rightarrow u_j(x)$ a.e. on Ω , for any fixed $m \geq 1$, by Fatou's lemma we have

$$\begin{aligned} \frac{1}{2} \sum_{j, k=1}^m \rho_j \rho_k \int_{\Omega \times \Omega} K(x, y) |u_j(x)|^2 |u_k(y)|^2 dx dy &\leq \\ \frac{1}{2} \liminf_{n \rightarrow \infty} \sum_{j, k=1}^m \rho_j \rho_k \int_{\Omega \times \Omega} K(x, y) |u_j^n(x)|^2 |u_k^n(y)|^2 dx dy &\leq \\ \frac{1}{2} \liminf_{n \rightarrow \infty} \sum_{j, k \geq 1} \rho_j \rho_k \int_{\Omega \times \Omega} K(x, y) |u_j^n(x)|^2 |u_k^n(y)|^2 dx dy &= \liminf_{n \rightarrow \infty} J_1(U_n) \end{aligned}$$

Finally letting $m \rightarrow \infty$, we get $J_1(U) \leq \liminf_{n \rightarrow \infty} J_1(U_n)$. \square

Remark 14. As a matter of fact when $N \leq 5$, we have $J_1(U) = \lim_{n \rightarrow \infty} J_1(U_n)$, that is J_1 is weakly sequentially continuous. Indeed let us assume $3 \leq N \leq 5$, as the case $N \leq 2$ can be easily handled analogously. Since for a fixed $j \geq 1$ we have $u_j^n \rightharpoonup u_j$ in \mathbb{H}_1 , and $u_j^n \rightarrow u_j$ strongly in $L^2(\Omega)$, and a.e. on Ω , as $n \rightarrow \infty$, we infer that for $2 < p < 2^* = 2N/(N-2)$ we have $u_j^n \rightarrow u_j$ strongly in $L^p(\Omega)$ and that $|u_j^n|^2 \rightarrow |u_j|^2$ in $L^{p/2}(\Omega)$ as $n \rightarrow \infty$. On the other hand, by Hölder's inequality and Sobolev imbedding theorem, if $v \in H_0^1(\Omega)$ and $\|v\|_1 = 1$, for $1 < p/2 < N/(N-2)$ we have

$$\|v^2\|_{p/2} \leq \|v^2\|_1^{1-\theta} \|v^2\|_{N/(N-2)}^\theta = \|v\|_{2^*}^{2\theta} \leq c \|\nabla v\|^{2\theta},$$

with $0 < \theta < 1$ given by

$$\frac{2}{p} = \frac{1-\theta}{1} + \frac{\theta(N-2)}{N}.$$

Applying this to $v := |u_j^n|^2$ yields

$$\| |u_j^n|^2 \|_{p/2} \leq c \|\nabla u_j^n\|^{2\theta}.$$

Now we infer first that for any $m \geq 1$ fixed, upon setting $q_0 := 1/\theta > 1$ and $q'_0 = q_0/(q_0 - 1) < \infty$ we have

$$\begin{aligned} \left\| \sum_{j>m} \rho_j |u_j^n|^2 \right\|_{p/2} &\leq \sum_{j>m} \rho_j \|u_j^n\|_p^2 \leq c \sum_{j>m} \rho_j \|\nabla u_j^n\|^{2\theta} \\ &\leq c \left(\sum_{j>m} \rho_j \right)^{1/q'_0} \left(\sum_{j>m} \rho_j \|\nabla u_j^n\|^2 \right)^{1/q_0} \leq c R^{1/q} \left(\sum_{j>m} \rho_j \right)^{1/q'}. \end{aligned}$$

From this it follows that

$$f_n := \sum_{j \geq 1} \rho_j |u_j^n|^2 \rightarrow f := \sum_{j \geq 1} \rho_j |u_j|^2 \quad \text{strongly in } L^{p/2}(\Omega). \quad (37)$$

Since here we are assuming that $N \leq 5$, we have $N/(N-2) > 2N/(N+2) = (2^*)'$ and thus we may fix $p > 2$ such that $2N/(N+2) \leq p/2 < N/(N-2)$: therefore $f_n \rightarrow f$ in $H^{-1}(\Omega)$, since $L^{p/2}(\Omega) \subset H^{-1}(\Omega)$ with continuous imbedding (the functions f_n and f are defined in (37)). Therefore, since $-\Delta V[U_n] = f_n$ and $V[U_n] \in H_0^1(\Omega)$, clearly we can deduce that $V[U_n] \rightarrow V[U]$ strongly in $H_0^1(\Omega)$ and finally

$$\int_{\Omega} |\nabla V[U_n]|^2 dx \rightarrow \int_{\Omega} |\nabla V[U]|^2 dx,$$

as claimed. \square

4 Existence of solutions for the Schrödinger–Poisson system

In this section we solve the following Schrödinger–Poisson problem:

$$\begin{cases} -\Delta u_m + \tilde{V}_0 u_m + V u_m = \lambda_m u_m & \text{in } \Omega \\ -\Delta V = \sum_{m=1}^{\infty} \rho_m |u_m|^2 & \text{in } \Omega \end{cases} \quad (38a)$$

$$(38b)$$

and moreover

$$\begin{cases} u_m \in H_0^1(\Omega), \text{ for all } m \geq 1, & V \in H_0^1(\Omega), \\ (u_m)_{m \geq 1} \text{ is a Hilbert basis for } L^2(\Omega). \end{cases} \quad (39a)$$

$$(39b)$$

The main result of this section is:

Theorem 15. *Assume that the hypotheses (24)–(25), as well as condition (10) are satisfied. The functionals J_0 and J_1 being defined in (33)–(34), we set $J(U) := J_0(U) + J_1(U)$ for $U \in \mathbb{S}$ given by (32). Then J achieves its minimum on \mathbb{S} and there exists $\hat{U}_0 \in \mathbb{S}$ such that $J_0(\hat{U}_0) = \min_{U \in \mathbb{S}} J_0(U)$, and the family $u_j := \hat{U}_0 \varphi_j$ is solution to (38a)–(39b). Moreover if $V \in H_0^1(\Omega)$ satisfies $-\Delta V = \sum_{m \geq 1} \rho_m |u_m|^2$, then the eigenvalues λ_j satisfy*

$$\lambda_j := (-\Delta u_j + \tilde{V}_0 u_j + V u_j |u_j) = \int_{\Omega} |\nabla u_j|^2 dx + \int_{\Omega} (\tilde{V}_0 + V) u_j(x)^2 dx.$$

We split the proof of this theorem into several lemmas. First we show that J achieves indeed its minimum.

Lemma 16. *The functional J achieves its minimum on \mathbb{S} at a certain $U_0 \in \mathbb{S}$.*

Proof. Since $\mathbb{S} \neq \emptyset$ and J_0 is bounded below (see Lemma 11), so is J and the infimum

$$\alpha := \inf_{U \in \mathbb{S}} J(U)$$

is finite. Consider a minimizing sequence $(U_n)_{n \geq 1} \in \mathbb{S}$, such that for instance $\alpha \leq J(U_n) \leq \alpha + 1/n$. In particular $J_0(U_n) \leq 1 + \alpha$, and thanks to Lemma 12, there exists a subsequence (which denote again by U_n) such that if we set $u_j^n := U_n \varphi_j$ for each fixed $j \geq 1$, we have, for all $2 \leq p < 2^*$,

$$\begin{cases} u_j^n \rightharpoonup u_j & \text{weakly in } \mathbb{H}_1, \\ u_j^n \rightarrow u_j & \text{strongly in } L^p(\Omega), \\ u_j^n \rightarrow u_j & \text{a.e. in } \Omega, \end{cases}$$

and the operator U_0 being defined by $U_0\varphi_j := u_j$, we have $U_0 \in \mathbb{S}$ and

$$J_0(U_0) \leq \liminf_{n \rightarrow \infty} J_0(U_n).$$

On the other hand, thanks to Lemma 13, we know that J_1 is weakly sequentially lower semi-continuous, that is $J_1(U_0) \leq \liminf_{n \rightarrow \infty} J_1(U_n)$. Thus

$$J(U_0) = J_0(U_0) + J_1(U_0) \leq \liminf_{n \rightarrow \infty} J_0(U_n) + \liminf_{n \rightarrow \infty} J_1(U_n) \leq \liminf_{n \rightarrow \infty} J(U_n),$$

that is $J(U_0) \leq \alpha$. Since $U_0 \in \mathbb{S}$, we conclude that $J(U_0) = \alpha$ that is J achieves its minimum at U_0 . \square

Lemma 17. *Let U_0 be given by Lemma 16, and let $M : H \rightarrow H$ be a bounded skewadjoint operator such that $M : \mathbb{H}_1 \rightarrow \mathbb{H}_1$ is also bounded. Set $U(t) := \exp(-tM)U_0$ for $t \in \mathbb{R}$, and $g_0(t) := J_0(U(t))$. Then g_0 is of class C^1 and*

$$g'_0(0) = -2\operatorname{Re} \sum_{j \geq 1} \rho_j(AU_0\varphi_j | MU_0\varphi_j). \quad (40)$$

Proof. First one checks easily that, since $M^* = -M$, one has $U(t) \in \mathbb{S}$ for all t , and thus the function

$$g_0(t) := J_0(U(t)) = \sum_{j \geq 1} \rho_j(AU(t)\varphi_j | U(t)\varphi_j)$$

is well defined and is of class C^1 . Since $U'(t) := dU(t)/dt = -M \exp(-tM)U_0$, one sees that

$$g'_0(t) = \sum_{j \geq 1} \rho_j(AU'(t)\varphi_j | U(t)\varphi_j) + \sum_{j \geq 1} \rho_j(AU(t)\varphi_j | U'(t)\varphi_j),$$

and finally,

$$g'_0(0) = - \sum_{j \geq 1} \rho_j(AMU_0\varphi_j | U_0\varphi_j) - \sum_{j \geq 1} \rho_j(AU_0\varphi_j | MU_0\varphi_j),$$

which yields our claim since $A^* = A$. \square

We have an analogous result concerning the functional J_1 : before showing this, we need to show that the mapping $U \mapsto V[U]$ is smooth.

Lemma 18. *Let U_0 be given by Lemma 16, and let $M : H \rightarrow H$ be a bounded skewadjoint operator such that $M : \mathbb{H}_1 \rightarrow \mathbb{H}_1$ is also bounded. Set $U(t) := \exp(-tM)U_0$ for $t \in \mathbb{R}$. Denoting by $V(t)$ the mapping $t \mapsto V[U(t)]$,*

then $t \mapsto V(t)$ is of class C^1 from \mathbb{R} into $L^\infty(\Omega) \cap H_0^1(\Omega)$ and denoting by W the solution of

$$-\Delta W = -2\operatorname{Re} \sum_{j \geq 1} \rho_j (MU_0 \varphi_j) \overline{U_0 \varphi_j}, \quad W \in H_0^1(\Omega),$$

we have $V'(0) = W$.

Proof. The fact that for all $T > 0$ and $t \in [-T, T]$ we have

$$\|M \exp(-tM)U_0 \varphi_j\|_{\mathbb{H}_1} \leq \|M\|_{\mathcal{L}(\mathbb{H}_1)} \exp(T\|M\|_{\mathcal{L}(\mathbb{H}_1)}) \|U_0 \varphi_j\|_{\mathbb{H}_1},$$

shows that for any p such that $1 < p/2 < N/(N-2)$ (see the proof of Lemma 13; here assume that $3 \leq N \leq 6$, the case $N \leq 2$ being treated analogously) the mapping

$$t \mapsto \sum_{j \geq 1} \rho_j |U(t) \varphi_j|^2$$

is of class C^1 from $(-T, T) \rightarrow L^{p/2}(\Omega) \subset H^{-1}(\Omega)$, and thus using (28) or (29) or (30), the mapping $t \mapsto V(t) := V[U(t)]$ is of class C^1 from $(-T, T)$ into $L^{p_1}(\Omega) \cap H_0^1(\Omega)$, where $p_1 := \infty$ if $N \leq 3$, and $p_1 < \infty$ arbitrary if $N = 4$, and $p_1 := N/(N-4)$ if $5 \leq N \leq 6$ (cf. Remark 10 above). The calculation of $V'(0)$ is straightforward. \square

Now we can state the following result, which will allow us to characterize U_0 given by Lemma 16.

Lemma 19. *Let U_0 be given by Lemma 16, and let $M : H \rightarrow H$ be a bounded skewadjoint operator such that $M : \mathbb{H}_1 \rightarrow \mathbb{H}_1$ is also bounded. Set $U(t) := \exp(-tM)U_0$ for $t \in \mathbb{R}$, and $g_1(t) := J_1(U(t))$. Then g_1 is of class C^1 and*

$$g_1'(0) = -2\operatorname{Re} \sum_{j \geq 1} \rho_j (V[U_0]U_0 \varphi_j | MU_0 \varphi_j). \quad (41)$$

Proof. Thanks to Lemma 18 one checks easily that the function

$$g_1(t) := J_1(U(t)) = \frac{1}{2} \int_{\Omega} |\nabla V[U(t)]|^2 dx = \frac{1}{2} (-\Delta V[U(t)] | V[U(t)])$$

is C^1 , and that denoting by $W := V'(0)$, we have (with $V_0 := V[U_0]$)

$$\begin{aligned} g_1'(0) &= (-\Delta W | V_0) = -2\operatorname{Re} \sum_{j \geq 1} \rho_j \int_{\Omega} (MU_0 \varphi_j) \overline{U_0 \varphi_j} V_0 dx \\ &= -2\operatorname{Re} \sum_{j \geq 1} \rho_j (MU_0 \varphi_j | V_0 U_0 \varphi_j), \end{aligned}$$

where we use the fact that $V_0 := V[U_0]$ is real valued. \square

The following result is analogous to Lemma 6: the only difference is that due to the presence of the nonlinear term we have to check that when some ρ_k has multiplicity $m \geq 2$, we can still proceed as before.

Lemma 20. *Under the assumptions of Theorem 15, let U_0 be given by lemma 16, and set $u_j := U_0\varphi_j$, for $j \geq 1$. Then the conclusions of lemma 6 hold.*

Proof. The proof is very much the same as in Lemma 6, so we give only the outline and the changes to be made. With the notations of Lemmas 17 and 19, we set $g(t) := J(U(t)) = g_0(t) + g_1(t)$. Since $g(0) \leq g(t)$ for all $t \in \mathbb{R}$, we have $g'(0) = 0$, that is

$$\operatorname{Re} \sum_{j \geq 1} \rho_j (AU_0\varphi_j | MU_0\varphi_j) + \operatorname{Re} \sum_{j \geq 1} \rho_j (V[U_0]U_0\varphi_j | MU_0\varphi_j) = 0 \quad (42)$$

for all bounded skew-adjoint operators M such that $M : \mathbb{H}_1 \rightarrow \mathbb{H}_1$ is also bounded.

In the same way, if we consider a bounded adjoint operator L such that $L : \mathbb{H}_1 \rightarrow \mathbb{H}_1$ is also bounded, we may set $M := iL$ and conclude that (42) yields

$$\operatorname{Im} \sum_{j \geq 1} \rho_j (AU_0\varphi_j | LU_0\varphi_j) + \operatorname{Im} \sum_{j \geq 1} \rho_j (V[U_0]U_0\varphi_j | LU_0\varphi_j) = 0 \quad (43)$$

for all such operators L .

At this point, in a first step, assume that the integer k is such that condition (16) is fulfilled. Choosing M and L as in (22), and proceeding exactly as in the proof of Lemma 6, using the fact that $\rho_k - \rho_n \neq 0$, we conclude that

$$(Au_k + V[U_0]u_k | u_n) = 0,$$

that is

$$Au_k + V[U_0]u_k \in \operatorname{span}\{u_n ; n \neq k\}^\perp = \operatorname{span}\{u_k\}.$$

This means that

$$Au_k + V[U_0]u_k = \lambda_k u_k$$

for some $\lambda_k \in \mathbb{C}$, but since $A + V[U_0]$ is a self-adjoint operator, as a matter of fact we have $\lambda_k \in \mathbb{R}$.

Next assume that the integer k is such that the coefficient ρ_k has multiplicity $m \geq 2$, that is condition (17) is satisfied. Arguing as above, we choose the operators M and L as in (23), and conclude that

$$(Au_{k+\ell} + V[U_0]u_{k+\ell}|u_n) = 0$$

for all $n \notin \{k+j; 0 \leq j \leq m-1\}$, that is:

$$Au_{k+\ell} + V[U_0]u_{k+\ell} \in (\text{span}\{u_n; n \neq k+j, 0 \leq j \leq m-1\})^\perp.$$

This means that if we set $H_k := \text{span}\{u_{k+i}; 0 \leq i \leq m-1\}$, and

$$A_0 u := -\Delta u + (\tilde{V}_0 + V[U_0]) u$$

then $A_0 : H_k \rightarrow H_k$ is a self-adjoint operator on the finite dimensional space H_k . Therefore there exists a unitary operator U_k , acting on this space, such that if for $0 \leq \ell \leq m-1$ we set $\hat{u}_{k+\ell} = U_k u_{k+\ell} = U_k U_0 \varphi_{k+\ell}$, we have $A_0 \hat{u}_{k+\ell} = \lambda_{k+\ell} \hat{u}_{k+\ell}$ for some $\lambda_{k+\ell} \in \mathbb{R}$.

However, since U_k is a unitary operator on H_k we have

$$\sum_{\ell=0}^{m-1} |U_0 \varphi_{k+\ell}|^2 = \sum_{\ell=0}^{m-1} |U_k U_0 \varphi_{k+\ell}|^2$$

and thus

$$\sum_{j=k}^{k+m-1} \rho_j |U_0 \varphi_j|^2 = \rho_k \sum_{\ell=0}^{m-1} |U_0 \varphi_{k+\ell}|^2 = \rho_k \sum_{\ell=0}^{m-1} |U_k U_0 \varphi_{k+\ell}|^2.$$

This means that if we set $\tilde{U}_k \varphi_j := U_0 \varphi_j$ if $j \notin \{k+\ell; 0 \leq \ell \leq m-1\}$ and $\tilde{U}_j = U_k U_0 \varphi_j$ if $k \leq j \leq k+m-1$, we have $V[U_0] = V[\tilde{U}_k]$, and finally this implies that

$$A_0 \hat{u}_{k+\ell} = -\Delta \hat{u}_{k+\ell} + (\tilde{V}_0 + V[\tilde{U}_k]) \hat{u}_{k+\ell} = \lambda_{k+\ell} \hat{u}_{k+\ell}.$$

□

As we may see from the above analysis, when all the ρ_j 's are distinct, then any unitary operator U_0 which minimizes J , yields a solution to the Schrödinger–Poisson system. However in the general case, when some of the coefficients ρ_k have multiplicity $m_k \geq 2$, it is possible that one has to impose a unitary transformation U_k in the space

$$H_k := \text{span}\{U_0 \varphi_{k+\ell}; 0 \leq \ell \leq m_k - 1\}$$

in order to obtain a solution (note that these unitary transformations do not change the value of $V[U_0]$).

In other words, one may find a unitary operator U_k on H_k such that if $A_k := A|_{H_k}$ is the trace of A on H_k , the operator $U_k^* A_k U_k$ is diagonal. Thus since $\rho_k = \rho_{k+\ell}$ for $0 \leq \ell \leq m-1$, if we denote by \widehat{U} the unitary operator obtained through the composition of all such operators U_k and U_0 , one has $J(\widehat{U}) = J(U_0)$. More precisely, we can state the following corollary, which ends the proof of Theorem 20:

Corollary 21. *Under the assumptions of Theorem 15, let U_k be given by Lemma 20 when $k \geq 1$ is such that (17) is satisfied. Define the operator \widehat{U}_0 by $\widehat{U}_0 \varphi_k = U_0 \varphi_k$ when k satisfies (16), and*

$$\widehat{U}_0 \varphi_{k+\ell} := U_k U_0 \varphi_{k+\ell}, \quad \text{for } 0 \leq \ell \leq m-1, \quad \text{when (17) is satisfied.}$$

Then \widehat{U}_0 belongs to \mathbb{S} , while $J(\widehat{U}_0) = J(U_0)$ and $V[\widehat{U}_0] = V[\widehat{U}_0]$. Moreover setting $u_j := \widehat{U}_0 \varphi_j$ for $j \geq 1$, there exists $\lambda_j \in \mathbb{R}$ such that

$$-\Delta u_j + \left(\widetilde{V}_0 + V[\widehat{U}_0] \right) u_j = \lambda_j u_j, \quad u_j \in \mathbb{H}_1, \quad (u_j | u_k) = \delta_{jk}$$

and moreover $V := V[\widehat{U}_0]$ satisfies

$$-\Delta V = \sum_{j \geq 1} \rho_j |u_j|^2, \quad V \in H_0^1(\Omega).$$

5 Further remarks

The one dimensional case $d = 1$ is particularly simple to handle, using a completely different method. Indeed we point out that for a given potential $V \in C([0, 1])$ the spectral sequence $(\lambda_k, \varphi_k)_{k \geq 1}$

$$-\varphi_k'' + (V + \widetilde{V})\varphi_k = \lambda_k \varphi_k, \quad \varphi_k(0) = \varphi_k(1) = 0, \quad \varphi_k'(0) > 0,$$

is well defined, and $\int_0^1 \varphi_k \varphi_j dx = \delta_{kj}$, each eigenvalue λ_k being simple. Using the simplicity of the eigenvalues, it is known that the mapping $V \mapsto \varphi_k$ is continuous from $C([0, 1])$ into $L^2(0, 1)$ (see J. Pöschel & E. Trubowitz [11]). If the coefficients $(\rho_j)_{j \geq 1}$ satisfy

$$\rho_j > 0, \quad \sum_{j \geq 1} \rho_j =: M < \infty$$

one can easily see that the mapping $F : C([0, 1]) \longrightarrow L^1(0, 1)$

$$V \mapsto F(V) := \sum_{k \geq 1} \rho_k |\varphi_k|^2$$

is continuous.

Now consider the mapping $B : L^1(0, 1) \longrightarrow L^1(0, 1)$ defined by $Bf := v$ where $v \in C^1([0, 1])$ is given by

$$-v'' = f \quad \text{in } (0, 1), \quad v(0) = v(1) = 0.$$

Clearly B can also be considered as a linear mapping on $C([0, 1])$, and B is compact. Also observe that the second equation in (1b) is equivalent to find $V \in C([0, 1])$ such that

$$V - BF(V) = 0.$$

Denoting by $T(V) := BF(V)$, we know that $I - T$ is a compact perturbation of the identity on $C([0, 1])$ and one can check easily that there exists $R > 0$ such that for any $\theta \in [0, 1]$

$$V - \theta T(V) = 0 \implies \|V\|_\infty < R.$$

Thus the invariance by homotopy of the Leray–Schauder topological degree implies that for all $\theta \in [0, 1]$ we have

$$\deg(I - \theta T, B(0, R), 0) = \deg(I, B(0, R), 0) = 1,$$

which means in particular that $\deg(I - T, B(0, R), 0) = 1$. Therefore there exists at least one $V \in C([0, 1])$ such that $V - T(V) = 0$, that is the system (1a)–(1b) has at least one solution, when $\Omega = (0, 1)$.

We point out also that in the case in which ρ_m is a function of λ_m , for instance $\rho_m := \exp(-\lambda_m)$ (see F. Nier [10]), the same approach can be applied. In dimensions $N \geq 1$, J.Ph. Solovej [15] considers a *one-particle density matrix* defined by $f \mapsto \gamma f := \sum_{m \geq 1} \rho_m(u_m | f) u_m$ and minimizes a functional depending on γ under the constraint $\text{spectrum}(\gamma) = \{\rho_m ; m \geq 1\}$. See also E. Prodan [12], E. Prodan & P. Nordlander [13] where Hartree-Fock approximations are considered.

References

- [1] N. Ben Abdallah and F. Méhats. On a Vlasov-Schrödinger-Poisson model. *Comm. Partial Differential Equations*, 29(1-2):173–206, 2004.

- [2] R. Illner, O. Kavian, and H. Lange. Stationary solutions of quasi-linear Schrödinger-Poisson systems. *J. Differential Equations*, 145(1):1–16, 1998.
- [3] O. Kavian. *Introduction à la théorie des points critiques et applications aux problèmes elliptiques*, volume 13 of *Mathématiques & Applications (Berlin) [Mathematics & Applications]*. Springer-Verlag, Paris, 1993.
- [4] O. Kavian. Remarks on regularity theorems for solutions to elliptic equations via the ultracontractivity of the heat semigroup. In *Contributions to nonlinear analysis*, volume 66 of *Progr. Nonlinear Differential Equations Appl.*, pages 321–350. Birkhäuser, Basel, 2006.
- [5] F. Méhats. Analysis of a quantum subband model for the transport of partially confined charged particles. *Monatsh. Math.*, 147(1):43–73, 2006.
- [6] L. Mirsky. A trace inequality of John von Neumann. *Monatsh. Math.*, 79(4):303–306, 1975.
- [7] J. von Neumann. Some matrix-inequalities and metrization of matrix-space. *Tomsk Univ. Rev.* 1, 286–300 (1937). Reprinted in *Collected Works* (Pergamon Press, 1962), iv, 205–219.
- [8] F. Nier. A stationary Schrödinger-Poisson system arising from the modelling of electronic devices. *Forum Math.*, 2(5):489–510, 1990.
- [9] F. Nier. Schrödinger-Poisson systems in dimension $d \leq 3$: the whole-space case. *Proc. Roy. Soc. Edinburgh Sect. A*, 123(6):1179–1201, 1993.
- [10] F. Nier. A variational formulation of Schrödinger-Poisson systems in dimension $d \leq 3$. *Comm. Partial Differential Equations*, 18(7-8):1125–1147, 1993.
- [11] J. Pöschel and E. Trubowitz. *Inverse spectral theory*, volume 130 of *Pure and Applied Mathematics*. Academic Press Inc., Boston, MA, 1987.
- [12] E. Prodan. Symmetry breaking in the self-consistent Kohn-Sham equations. *J. Phys. A*, 38(25):5647–5657, 2005.
- [13] E. Prodan and P. Nordlander. Hartree approximation. I. The fixed point approach. *J. Math. Phys.*, 42(8):3390–3406, 2001.

- [14] M. Reed and B. Simon. *Methods of modern mathematical physics. I.* Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, second edition, 1980. Functional analysis.
- [15] Jan Philip Solovej. Proof of the ionization conjecture in a reduced Hartree-Fock model. *Invent. Math.*, 104(2):291–311, 1991.
- [16] G. Stampacchia. Équations Elliptiques du Second Ordre à Coefficients Discontinus. Presses de l'Université de Montréal, série “Séminaires de Mathématiques Supérieures”, # 16, Montréal, 1965.
- [17] K. Yosida. *Functional analysis.* Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the sixth (1980) edition.
- [18] P. F. Zweifel. The Poisson-Wigner system: a quantum transport equation. In *Proceedings of the XXV Symposium on Mathematical Physics (Toruń, 1992)*, volume 33, pages 317–323, 1993.