



Constructible characters and b-invariants

Cédric Bonnafé

► To cite this version:

| Cédric Bonnafé. Constructible characters and b-invariants. 2013. hal-00787350v1

HAL Id: hal-00787350

<https://hal.science/hal-00787350v1>

Preprint submitted on 12 Feb 2013 (v1), last revised 8 Apr 2015 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONSTRUCTIBLE CHARACTERS AND b -INVARIANT

by

CÉDRIC BONNAFÉ

Let (W, S) be a finite Coxeter system and let $\varphi : S \rightarrow \mathbb{R}_{>0}$ be a *weight function* that is, a map such that $\varphi(s) = \varphi(t)$ whenever s and t are conjugate in W . Associated with this datum, G. Lusztig has defined [Lu3, §22] a notion of *constructible characters* of W : it is conjectured that a character is constructible if and only if it is the character afforded by a Kazhdan-Lusztig left cell (defined using the weight function φ). These constructible characters depend heavily on φ so we will call them the φ -*constructible characters* of W : the set of φ -constructible characters will be denoted by $\text{Cons}_{\varphi}^{\text{Lus}}(W)$. We shall also define a graph $\mathcal{G}_{W, \varphi}^{\text{Lus}}$ as follows: the vertices of $\mathcal{G}_{W, \varphi}^{\text{Lus}}$ are the irreducible characters and two irreducible characters χ and χ' are joined in this graph if there exists a φ -constructible character γ of W such that χ and χ' both occur as constituents of γ . The connected components of $\mathcal{G}_{W, \varphi}^{\text{Lus}}$ (viewed as subsets of $\text{Irr}(W)$) will be called the *Lusztig φ -families*: the set of Lusztig φ -families will be denoted by $\text{Fam}_{\varphi}^{\text{Lus}}(W)$. If $\mathcal{F} \in \text{Fam}_{\varphi}^{\text{Lus}}(W)$, we denote by $\text{Cons}_{\varphi}^{\text{Lus}}(\mathcal{F})$ the set of φ -constructible characters of W all of whose irreducible components belong to \mathcal{F} .

On the other hand, using the theory of rational Cherednik algebras at $t = 0$ and the geometry of the Calogero-Moser space associated with (W, φ) , R. Rouquier and the author (see [BoRo1] and [BoRo2]) have defined a notion of *Calogero-Moser φ -cells* of W , a notion of *Calogero-Moser φ -cellular characters* of W (whose set is denoted by $\text{Cell}_{\varphi}^{\text{CM}}(W)$) and a notion of *Calogero-Moser φ -families* (whose set is denoted by $\text{Fam}_{\varphi}^{\text{CM}}(W)$).

Conjecture (see [BoRo1], [BoRo2] and [GoMa]). *With the above notation,*

$$\text{Cons}_{\varphi}^{\text{Lus}}(W) = \text{Cell}_{\varphi}^{\text{CM}}(W) \quad \text{and} \quad \text{Fam}_{\varphi}^{\text{Lus}}(W) = \text{Fam}_{\varphi}^{\text{CM}}(W)$$

for every weight function $\varphi : S \rightarrow \mathbb{R}_{>0}$.

The statement about families in this conjecture holds for classical Weyl groups thanks to a case-by-case analysis relying on [Lu3, §22] (for the computation of Lusztig φ -families), [GoMa] (for the computation of Calogero-Moser φ -families in type A and B) and [Be2] (for the computation of the Calogero-Moser φ -families in type D). It also holds whenever $|S| = 2$ (see [Lu3, §17 and Lemma 22.2] and [Be1, §6.10]).

The statement about constructible characters is much more difficult to establish, as the computation of Calogero-Moser φ -cellular characters is at that time out of reach: it has been proved whenever the Calogero-Moser space associated with (W, S, φ) is smooth [BoRo2, Theorem 14.4.1] (it has also been checked if W is of type $B_2 \dots$).

Our aim in this paper is to show that this conjecture is compatible with properties of the b -invariant (as defined below). With each irreducible character χ of W is associated its *fake degree* $f_\chi(\mathbf{t})$, using the invariant theory of W (see for instance [BoRo2, Definition 1.5.7]). Let us denote by b_χ the valuation of $f_\chi(\mathbf{t})$: b_χ is called the b -invariant of χ . For instance, $b_1 = 0$ and b_ε is the number of reflections of W (here, $\varepsilon : W \rightarrow \{1, -1\}$ denotes the sign character). Also, $b_\chi = 1$ if and only if χ is an irreducible constituent of the canonical reflection representation of W . The following result is proved in [BoRo2, Theorems 9.6.1 and 12.3.14]:

Theorem CM. *Let $\varphi : S \rightarrow \mathbb{R}_{>0}$ be a weight function. Then:*

- (a) *If $\mathcal{F} \in \text{Fam}_\varphi^{\text{CM}}(W)$, then there exists a unique $\chi \in \mathcal{F}$ with minimal b -invariant.*
- (b) *If $\gamma \in \text{Cons}_\varphi^{\text{CM}}(W)$, then there exists a unique irreducible constituent χ of γ with minimal b -invariant.*

The next theorem is proved in [Lu2, Theorem 5.25 and its proof] (see also [Lu1] for the first occurrence of the *special* representations):

Theorem (Lusztig). *Assume that φ is constant. Then:*

- (a) *If $\mathcal{F} \in \text{Fam}_\varphi^{\text{Lus}}(W)$, then there exists a unique $\chi_{\mathcal{F}} \in \mathcal{F}$ with minimal b -invariant ($\chi_{\mathcal{F}}$ is called the **special** character of \mathcal{F}).*
- (b) *If $\gamma \in \text{Cons}_\varphi^{\text{Lus}}(\mathcal{F})$, then $\chi_{\mathcal{F}}$ is an irreducible constituent of γ (and, of course, among the irreducible constituents of γ , $\chi_{\mathcal{F}}$ is the unique one with minimal b -invariant).*

It turns out that, for general φ , there might exist Lusztig φ -families \mathcal{F} such that no element of \mathcal{F} occurs as an irreducible constituent of *all* the φ -constructible characters in $\text{Cons}_\varphi^{\text{Lus}}(\mathcal{F})$ (this already occurs in type B_3 , and the reader can also check this fact in type F_4 , using the tables given by Geck [Ge, Table 2]). Nevertheless, we will prove in this paper the following result, which is compatible with the above conjecture and the above theorems:

Theorem L. *Let $\varphi : S \rightarrow \mathbb{R}_{>0}$ be a weight function. Then:*

- (a) *If $\mathcal{F} \in \text{Fam}_{\varphi}^{\text{Lus}}(W)$, then there exists a unique $\chi \in \mathcal{F}$ with minimal b -invariant.*
- (b) *If $\gamma \in \text{Cons}_{\varphi}^{\text{Lus}}(W)$, then there exists a unique irreducible constituent χ of γ with minimal b -invariant.*

The proof of Theorem CM is general and conceptual, while our proof of Theorem L goes through a case-by-case analysis, based on Lusztig's description of φ -constructible characters and Lusztig φ -families [Lu3, §22].

REMARK 0 - Let γ_{χ} denote the coefficient of $\mathbf{t}^{b_{\chi}}$ in $F_{\chi}(\mathbf{t})$. Then it has been noticed by Lusztig [Lu1, §2, Page 325] that $\gamma_{\chi} = 1$ whenever χ is special.

As the only irreducible Coxeter systems affording possibly unequal parameters are of type $I_2(2m)$, F_4 or B_n , and as $\gamma_{\chi} = 1$ for any character χ in these groups, we can conclude that, in general (equal or unequal parameters), $\gamma_{\chi} = 1$ for all the characters χ with minimal b -invariant constructed in Theorem L (for both (a) and (b)).

The same property holds for the characters χ with minimal b -invariant constructed in Theorem CM (in this case, the proof is again general and conceptual [BoRo2]). ■

1. Proof of Theorem L

1.A. Reduction. — It is easily seen that the proof of Theorem L may be reduced to the case where (W, S) is irreducible. If W is of type A_n , D_n , E_6 , E_7 , E_8 , H_3 or H_4 , then φ is necessarily constant and Theorem L follows immediately from Lusztig's Theorem. If W is dihedral, then Theorem L is easily checked using [Lu3, §17 and Lemma 22.2]. If W is of type F_4 , then Theorem L follows from inspection of [Ge, Table 2]. Therefore, this shows that we may, and we will, assume that W is of type B_n , with $n \geq 2$. Write $S = \{t, s_1, s_2, \dots, s_{n-1}\}$ in such a way that the Dynkin diagram of (W, S) is

$$(\#) \quad \begin{array}{ccccccc} & t & s_1 & s_2 & & \dots & s_{n-1} \\ & \circ & \circ & \circ & \text{---} & \dots & \text{---} \circ \end{array}$$

Write $b = \varphi(t)$ and $a = \varphi(s_1) = \varphi(s_2) = \dots = \varphi(s_{n-1})$. If $b \notin a\mathbb{N}^*$, then $\text{Cons}_{\varphi}^{\text{Lus}}(W) = \text{Irr}(W)$ (see [Lu3, Proposition 22.25]) and Theorem L becomes obvious. So we may assume that $b = ra$ with $r \in \mathbb{N}^*$. To summarize:

Hypothesis. *From now on, and until the end of this section, we will assume that (W, S) is of type B_n , with $n \geq 2$, that $S = \{t, s_1, s_2, \dots, s_{n-1}\}$ is such that the Dynkin diagram of (W, S) is given by (#), that $\varphi(t) = r\varphi(s_1) = r\varphi(s_2) = \dots = r\varphi(s_{n-1}) = 1$ with $r \in \mathbb{N}^*$.*

1.B. Admissible involutions. — Let $l \geq 0$ and let Z be a totally ordered set of size $2l + r$. We shall define by induction on l what is an r -admissible involution of Z . Let $\iota : Z \rightarrow Z$ be an involution. Then ι is said *r -admissible* if it has r fixed points and, if $l \geq 1$, there exist two consecutive elements b and c of Z such that $\iota(b) = c$ and the restriction of ι to $Z \setminus \{b, c\}$ is r -admissible.

Note that, if ι is an r -admissible involution and if $\iota(b) = c > b$ and $\iota(z) = z$, then $z < b$ or $z > c$ (this is easily proved by induction on $|Z|$).

1.C. Symbols. — We shall denote by $\mathbf{Sym}_k(r)$ the set of *symbols* $\Lambda = \begin{pmatrix} \beta \\ \gamma \end{pmatrix}$ where $\beta = (\beta_1 < \beta_2 < \dots < \beta_{k+r})$ and $\gamma = (\gamma_1 < \gamma_2 < \dots < \gamma_k)$ are increasing sequences of *non-zero* natural numbers. We set

$$|\Lambda| = \sum_{i=1}^{k+r} (\beta_i - i) + \sum_{j=1}^k (\gamma_j - j)$$

and
$$\mathbf{b}(\Lambda) = \sum_{i=1}^{k+r} (2k + 2r - 2i)(\beta_i - i) + \sum_{j=1}^k (2k + 1 - 2j)(\gamma_j - j).$$

The number $\mathbf{b}(\Lambda)$ will be called the *\mathbf{b} -invariant* of Λ . For simplifying our arguments, we shall define

$$\nabla_{k,r} = \sum_{i=1}^{k+r} (2k + 2r - 2i)i + \sum_{j=1}^k (2k + 1 - 2j)j$$

so that

$$\mathbf{b}(\Lambda) = \sum_{i=1}^{k+r} (2k + 2r - 2i)\beta_i + \sum_{j=1}^k (2k + 1 - 2j)\gamma_j - \nabla_{k,r}.$$

By abuse of notation, we denote by $\beta \cap \gamma$ the set $\{\beta_1, \beta_2, \dots, \beta_{k+r}\} \cap \{\gamma_1, \gamma_2, \dots, \gamma_k\}$ and by $\beta \cup \gamma$ the set $\{\beta_1, \beta_2, \dots, \beta_{k+r}\} \cup \{\gamma_1, \gamma_2, \dots, \gamma_k\}$. We also set $\beta \dot{+} \gamma = (\beta \cup \gamma) \setminus (\beta \cap \gamma)$.

Now, let $\mathbf{z}'(\Lambda) = (\beta_1, \beta_2, \dots, \beta_r, \gamma_1, \beta_{r+1}, \gamma_2, \beta_{r+2}, \dots, \gamma_k, \beta_{r+k})$. We shall write

$$\mathbf{z}'(\Lambda) = (z'_1(\Lambda), z'_2(\Lambda), \dots, z'_{2k+r}(\Lambda)),$$

so that

$$\begin{aligned} \mathbf{b}(\Lambda) &= \sum_{i=1}^r (2k + 2r - 2i)z'_i(\Lambda) + \sum_{i=r+1}^{2k+r} (2k + r - i)z'_i(\Lambda) - \nabla_{k,r} \\ (\clubsuit) \quad &= \sum_{i=1}^r (r - i)z'_i(\Lambda) + \sum_{i=1}^{2k+r} (2k + r - i)z'_i(\Lambda) - \nabla_{k,r} \\ &= \sum_{i=1}^r \left(\sum_{j=1}^i z'_j(\Lambda) \right) + \sum_{i=1}^{2k+r-1} \left(\sum_{j=1}^i z'_j(\Lambda) \right) - \nabla_{k,r}. \end{aligned}$$

1.D. Families of symbols. — We denote by $\mathbf{z}(\Lambda)$ the sequence $z_1 \leq z_2 \leq \dots \leq z_{2k+r}$ obtained after rewriting the sequence $(\beta_1, \beta_2, \dots, \beta_{k+r}, \gamma_1, \gamma_2, \dots, \gamma_k)$ in non-decreasing order.

REMARK 1 - Note that the sequence $\mathbf{z}'(\Lambda)$ determines the symbol Λ , contrarily to the sequence $\mathbf{z}(\Lambda)$. However, $\mathbf{z}(\Lambda)$ determines completely $|\Lambda|$ thanks to the formula $|\Lambda| = \sum_{z \in \mathbf{z}(\Lambda)} z - r(r+1)/2 - (k+r)(k+r+1)/2$. \square

We say that two symbols $\Lambda = \begin{pmatrix} \beta \\ \gamma \end{pmatrix}$ and $\Lambda' = \begin{pmatrix} \beta' \\ \gamma' \end{pmatrix}$ in $\mathbf{Sym}_k(r)$ are in the same family if $\mathbf{z}(\Lambda) = \mathbf{z}(\Lambda')$. Note that this is equivalent to say that $\beta \cap \gamma = \beta' \cap \gamma'$ and $\beta \cup \gamma = \beta' \cup \gamma'$. If \mathcal{F} is the family of Λ , we set $X_{\mathcal{F}} = \beta \cap \gamma$ and $Z_{\mathcal{F}} = \beta \dot{+} \gamma$: note that $X_{\mathcal{F}}$ and $Z_{\mathcal{F}}$ depend only on \mathcal{F} (and not on the particular choice of $\Lambda \in \mathcal{F}$).

If ι is an r -admissible involution of $Z_{\mathcal{F}}$, we denote by \mathcal{F}_{ι} the set of symbols $\Lambda = \begin{pmatrix} \beta \\ \gamma \end{pmatrix}$ in \mathcal{F} such that $|\beta \cap \omega| = 1$ for all ι -orbits ω .

1.E. Lusztig families, constructible characters. — Let $\Lambda \in \mathbf{Sym}_k(r)$ be such that $|\Lambda| = n$. Let $\text{Bip}(n)$ be the set of bipartitions of n . We set

$$\lambda_1(\Lambda) = (\beta_{k+r} - (k+r) \geq \dots \geq \beta_2 - 2 \geq \beta_1 - 1),$$

$$\lambda_2(\Lambda) = (\gamma_k - k \geq \dots \geq \gamma_2 - 2 \geq \gamma_1 - 1)$$

and

$$\lambda(\Lambda) = (\lambda_1(\Lambda), \lambda_2(\Lambda)).$$

Then $\lambda(\Lambda)$ is a bipartition of n . We denote by χ_{Λ} the irreducible character of W denoted by $\chi_{\lambda(\Lambda)}$ in [Lu3, §22] or in [GePf, §5.5.3]. Then [GePf, §5.5.3]

$$(\diamond) \quad b_{\chi_{\Lambda}} = \mathbf{b}(\Lambda).$$

With these notation, Lusztig described the φ -constructible characters in [Lu3, Proposition 22.24], from which the description of Lusztig φ -families follow by using [Lu3, Lemma 22.22]:

Theorem 2 (Lusztig). *Let \mathcal{F}_{Lus} be a Lusztig φ -family and let $\gamma \in \text{Cons}_{\varphi}^{\text{Lus}}(\mathcal{F}_{\text{Lus}})$. If we choose k sufficiently large, then:*

(a) *There exists a family \mathcal{F} of symbols in $\mathbf{Sym}_k(r)$ such that*

$$\mathcal{F}_{\text{Lus}} = \{\chi_{\Lambda} \mid \Lambda \in \mathcal{F}\}.$$

(b) *There exists an r -admissible involution ι of $Z_{\mathcal{F}}$ such that*

$$\gamma = \sum_{\Lambda \in \mathcal{F}_{\iota}} \chi_{\Lambda}.$$

Definition 3. The symbol Λ is said *special* if $\mathbf{z}(\Lambda) = \mathbf{z}'(\Lambda)$.

REMARK 4 - According to Remark 1, there is a unique special symbol in each family. It will be denoted by $\Lambda_{\mathcal{F}}$. Finally, note that, if Λ, Λ' belong to the same family, then $|\Lambda| = |\Lambda'|$. \square

Now, Theorem L follows from Theorem 2, Formula (\diamond) and the following next Theorem:

Theorem 5. Let \mathcal{F} be a family of symbols in $\mathbf{Sym}_k(r)$, let ι be an r -admissible involution of $Z_{\mathcal{F}}$ and let $\Lambda \in \mathcal{F}$. Then:

- (a) $\mathbf{b}(\Lambda) \geq \mathbf{b}(\Lambda_{\mathcal{F}})$ with equality if and only if $\Lambda = \Lambda_{\mathcal{F}}$.
- (b) There is a unique symbol $\Lambda_{\mathcal{F}, \iota}$ in \mathcal{F}_{ι} such that, if $\Lambda \in \mathcal{F}_{\iota}$, then $\mathbf{b}(\Lambda) \geq \mathbf{b}(\Lambda_{\mathcal{F}, \iota})$, with equality if and only if $\Lambda = \Lambda_{\mathcal{F}, \iota}$.

1.F. Proof of Theorem 5(a). — First, note that $\mathbf{z}(\Lambda) = \mathbf{z}(\Lambda_{\mathcal{F}}) = \mathbf{z}'(\Lambda_{\mathcal{F}})$. As $\mathbf{z}'(\Lambda)$ is a permutation of the non-decreasing sequence $\mathbf{z}'(\Lambda_{\mathcal{F}})$, we have

$$\sum_{j=1}^i z'_j(\Lambda) \geq \sum_{j=1}^i z'_j(\Lambda_{\mathcal{F}})$$

for all $i \in \{1, 2, \dots, 2k+r\}$. So, it follows from (\clubsuit) that

$$\mathbf{b}(\Lambda) - \mathbf{b}(\Lambda_{\mathcal{F}}) = \sum_{i=1}^{r-1} \left(\sum_{j=1}^i (z'_j(\Lambda) - z'_j(\Lambda_{\mathcal{F}})) \right) + \sum_{i=1}^{2k+r-1} \left(\sum_{j=1}^i (z'_j(\Lambda) - z'_j(\Lambda_{\mathcal{F}})) \right).$$

So $\mathbf{b}(\Lambda) \geq \mathbf{b}(\Lambda_{\mathcal{F}})$ with equality only whenever $\sum_{j=1}^i z'_j(\Lambda) = \sum_{j=1}^i z'_j(\Lambda_{\mathcal{F}})$ for all $i \in \{1, 2, \dots, 2k+r\}$. The proof of Theorem 5(a) is complete.

1.G. Reduction for the proof of Theorem 5(b). — First, assume that $X_{\mathcal{F}} \neq \emptyset$. Let $b \in X_{\mathcal{F}}$ and let $\Lambda = \begin{pmatrix} \beta \\ \gamma \end{pmatrix} \in \mathcal{F}$. Then $b \in \beta \cap \gamma = X_{\mathcal{F}}$ and we denote by $\beta[b]$ the sequence obtained by removing b to β . Similarly, let $\Lambda[b] = \begin{pmatrix} \beta[b] \\ \gamma[b] \end{pmatrix}$.

Then $\Lambda[b] \in \mathbf{Sym}_{k-1}(r)$ and

$$(\heartsuit) \quad \mathbf{b}(\Lambda) = \mathbf{b}(\Lambda[b]) + \nabla_{k,r} - \nabla_{k-1,r} + b \left(4k + 2r + 1 - \sum_{\substack{z \in \mathbf{z}(\Lambda) \\ z \leq b}} 2 \right) + 2 \sum_{\substack{z \in \mathbf{z}(\Lambda) \\ z < b}} z.$$

Proof of (\heartsuit). Let i_0 and j_0 be such that $\beta_{i_0} = b$ and $\gamma_{j_0} = b$. Then

$$\mathbf{b}(\Lambda) - \mathbf{b}(\Lambda[b]) = \nabla_{k,r} - \nabla_{k-1,r} + (2k + 2r - 2i_0)b + \sum_{i=1}^{i_0-1} 2\beta_i + (2k + 1 - 2j_0)b + \sum_{j=1}^{j_0-1} 2\gamma_j.$$

But the numbers $\beta_1, \beta_2, \dots, \beta_{i_0}, \gamma_1, \gamma_2, \dots, \gamma_{j_0}$ are exactly the elements of the sequence $\mathbf{z}(\Lambda)$ which are $\leq b$. So

$$i_0 + j_0 = \sum_{\substack{z \in \mathbf{z}(\Lambda) \\ z \leq b}} 1$$

and

$$\sum_{i=1}^{i_0-1} \beta_i + \sum_{j=1}^{j_0-1} \gamma_j = \sum_{\substack{z \in \mathbf{z}(\Lambda) \\ z < b}} z.$$

This shows (\heartsuit) . ■

Now, the family of $\Lambda[b]$ depends only on the family of Λ (and not on Λ itself): indeed, $\mathbf{z}(\Lambda[b])$ is obtained from $\mathbf{z}(\Lambda)$ by removing the two entries equal to b . We will denote by $\mathcal{F}[b]$ the family of $\Lambda[b]$. Moreover, $Z_{\mathcal{F}[b]} = Z_{\mathcal{F}}$ and the map $\Lambda \mapsto \Lambda[b]$ induces a bijection between \mathcal{F} and $\mathcal{F}[b]$, and also induces a bijection between \mathcal{F}_i and $\mathcal{F}[b]_i$.

Moreover, the formula (\heartsuit) shows that the difference between $\mathbf{b}(\Lambda)$ and $\mathbf{b}(\Lambda[b])$ depends only on b and \mathcal{F} , so proving Lemma 6 for the pair (\mathcal{F}, ι) is equivalent to proving Lemma 6 for the pair $(\mathcal{F}[b], \iota)$. By applying several times this principle if necessary, this means that we may, and we will, assume that

$$X_{\mathcal{F}} = \emptyset.$$

1.H. Proof of Theorem 5(b). — We denote by $f_r < \dots < f_1$ the elements of $Z_{\mathcal{F}}$ which are fixed by ι . We also set $f_{r+1} = 0$ and $f_0 = \infty$. As ι is r -admissible, the set $Z_{\mathcal{F}}^{(d)} = \{z \in Z_{\mathcal{F}} \mid f_{d+1} < z < f_d\}$ is ι -stable and contains no ι -fixed point (for $d \in \{0, 1, \dots, r\}$). Let $k_d = |Z_{\mathcal{F}}^{(d)}|/2$ and let ι_d be the restriction of ι to $Z_{\mathcal{F}}^{(d)}$. Then ι_d is a 0-admissible involution of $Z_{\mathcal{F}}^{(d)}$.

If $\Lambda = \begin{pmatrix} \beta \\ \gamma \end{pmatrix} \in \mathcal{F}_{\iota}$, we set $\beta^{(d)} = \beta \cap Z_{\mathcal{F}}^{(d)}$, $\gamma^{(d)} = \gamma \cap Z_{\mathcal{F}}^{(d)}$ and $\Lambda^{(d)} = \begin{pmatrix} \beta^{(d)} \\ \gamma^{(d)} \end{pmatrix}$. Then $\Lambda^{(d)} \in \mathbf{Sym}_{k_d}(0)$ and, if $\mathcal{F}^{(d)}$ denotes the family of $\Lambda^{(d)}$, then $\Lambda^{(d)} \in \mathcal{F}_{\iota_d}^{(d)}$.

Now, if $\Lambda' = \begin{pmatrix} \beta' \\ \gamma' \end{pmatrix} \in \mathbf{Sym}_{k'}(0)$, we set

$$\mathbf{b}_d(\Lambda') = \sum_{i=1}^{k'} (2k' + 2d - 2i) \beta'_i + \sum_{j=1}^{k'} (2k' + 1 - 2j) \gamma'_j.$$

The number $\mathbf{b}_d(\Lambda')$ is called the \mathbf{b}_d -invariant of Λ' . It then follows from the definition of \mathbf{b} and $\nabla_{k,r}$ that

$$(\spadesuit) \quad \mathbf{b}(\Lambda) = \sum_{d=0}^r \mathbf{b}_d(\Lambda^{(d)}) - \nabla_{k,r} + \sum_{d=1}^r 2(k_0 + k_1 + \dots + k_{d-1}) \left(f_d + \sum_{z \in Z^{(d)}} z \right).$$

Since the map

$$\begin{aligned} \mathcal{F}_\iota &\longrightarrow \prod_{d=0}^r \mathcal{F}_{\iota_d}^{(d)} \\ \Lambda &\longmapsto (\Lambda^{(0)}, \Lambda^{(1)}, \dots, \Lambda^{(d)}) \end{aligned}$$

is bijective and since $\mathbf{b}(\Lambda) - \sum_{d=0}^r \mathbf{b}_d(\Lambda^{(d)})$ depends only on (\mathcal{F}, ι) and not on Λ (as shown by the formula (\spadesuit)), Theorem 5(b) will follow from the following lemma :

Lemma 6. *There exists a unique symbol in $\mathcal{F}_{\iota_d}^{(d)}$ with minimal \mathbf{b}_d -invariant.*

The proof of Lemma 6 will be given in the next section.

2. Minimal \mathbf{b}_d -invariant

For simplifying notation, we set $Z = Z_{\mathcal{F}}^{(d)}$, $l = k_d$, $\mathcal{G} = \mathcal{F}^{(d)}$ and $J = \iota_d$. Let us write $Z = \{z_1, z_2, \dots, z_{2l}\}$ with $z_1 < z_2 < \dots < z_{2l}$. Recall from the previous section that J is a 0-admissible involution of Z .

2.A. Construction. — We shall define by induction on $l \geq 0$ a symbol $\Lambda_J^{(d)}(Z) \in \mathcal{G}_J$. If $l = 0$, then $\Lambda_J^{(d)}(Z)$ is obviously empty. So assume now that, for any set of non-zero integers Z' of order $2(l-1)$, for any 0-admissible involution J' of Z' and any $d' \geq 0$, we have defined a symbol $\Lambda_{J'}^{(d')}(Z')$. Then $\Lambda_J^{(d)}(Z) = \begin{pmatrix} \beta_J^{(d)}(Z) \\ \gamma_J^{(d)}(Z) \end{pmatrix}$ is defined as follows: let $Z' = Z \setminus \{z_1, \iota(z_1)\}$, J' the restriction of J to Z' and let

$$d' = \begin{cases} d-1 & \text{if } d \geq 1, \\ 1 & \text{if } d = 0. \end{cases}$$

Then $|Z'| = 2(l-1)$ and J' is 0-admissible. So $\Lambda_{J'}^{(d')}(Z') = \begin{pmatrix} \beta_{J'}^{(d')}(Z') \\ \gamma_{J'}^{(d')}(Z') \end{pmatrix}$ is well-defined by the induction hypothesis. We then set

$$\beta_J^{(d)}(Z) = \begin{cases} \beta_{J'}^{(d')}(Z') \cup \{z_1\} & \text{if } d \geq 1, \\ \beta_{J'}^{(d')}(Z') \cup \{J(z_1)\} & \text{if } d = 0, \end{cases}$$

and

$$\gamma_J^{(d)}(Z) = \begin{cases} \gamma_{J'}^{(d')}(Z') \cup \{J(z_1)\} & \text{if } d \geq 1, \\ \gamma_{J'}^{(d')}(Z') \cup \{z_1\} & \text{if } d = 0. \end{cases}$$

Then Lemma 6 is implied by the next lemma :

Lemma 6⁺. *Let $\Lambda \in \mathcal{G}_J$. Then $\mathbf{b}_d(\Lambda) \geq \mathbf{b}_d(\Lambda_J^{(d)}(Z))$ with equality if and only if $\Lambda = \Lambda_J^{(d)}(Z)$.*

The rest of this section is devoted to the proof of Lemma 6⁺. We will first prove Lemma 6⁺ whenever $d \in \{0, 1\}$ using Lusztig's Theorem. We will then turn to the general case, which will be handled by induction on $l = |Z|/2$. We fix $\Lambda = \begin{pmatrix} \beta \\ \gamma \end{pmatrix} \in \mathcal{G}_l$.

2.B. Proof of Lemma 6⁺ whenever $d = 1$. — Let z be a natural number strictly bigger than all the elements of Z . Let $\tilde{\Lambda} = \begin{pmatrix} \beta \cup \{z\} \\ \gamma \end{pmatrix} \in \mathbf{Sym}_k(1)$. Then $\mathbf{b}_1(\Lambda) = \mathbf{b}(\tilde{\Lambda}) + C$, where C depends only on Z . Let $\tilde{\Lambda}_0 = \begin{pmatrix} z_1, z_3, \dots, z_{2l-1}, z \\ z_2, \dots, z_{2l} \end{pmatrix}$. Since j is 0-admissible, it is easily seen that, if $j(z_i) = z_j$, then $j - i$ is odd. So $\tilde{\Lambda}_0 \in \mathcal{G}_j$. But, by [Lu1, §5], $\mathbf{b}(\tilde{\Lambda}) \geq \mathbf{b}(\tilde{\Lambda}_0)$ with equality if and only if $\tilde{\Lambda} = \tilde{\Lambda}_0$. So it is sufficient to notice that $\overline{\Lambda_j^{(1)}(Z)} = \tilde{\Lambda}_0$, which is easily checked.

2.C. Proof of Lemma 6⁺ whenever $d = 0$. — Assume in this subsection, and only in this subsection, that $d = 0$ or 1. We denote by $\Lambda^{\text{op}} = \begin{pmatrix} \gamma \\ \beta \end{pmatrix} \in \mathcal{G}_j$. It is readily seen from the construction that $\Lambda_j^{(0)}(Z)^{\text{op}} = \Lambda_j^{(1)}(Z)$ and that

$$\mathbf{b}_1(\Lambda) = \mathbf{b}_0(\Lambda^{\text{op}}) + \sum_{z \in Z} z.$$

So Lemma 6⁺ for $d = 0$ follows from Lemma 6⁺ for $d = 1$.

2.D. Proof of Lemma 6⁺ whenever $d \geq 2$. — Assume now, and until the end of this section, that $d \geq 2$. We shall prove Lemma 6⁺ by induction on $l = |Z|/2$. The result is obvious if $l = 0$, as well as if $l = 1$. So we assume that $l \geq 2$ and that Lemma 6⁺ holds for $l' \leq l - 1$. Write $j(z_1) = z_{2m}$, where $m \leq l$ (note that $j(z_1) \notin \{z_1, z_3, z_5, \dots, z_{2l-1}\}$ since j is 0-admissible).

Assume first that $m < l$. Then Z can be written as the union $Z = Z^+ \dot{\cup} Z^-$, where $Z^+ = \{z_1, z_2, \dots, z_{2m}\}$ and $Z^- = \{z_{2m+1}, z_{2m+2}, \dots, z_{2l}\}$ are j -stable (since j is 0-admissible). If $\varepsilon \in \{+, -\}$, let j^ε denote the restriction of j to Z^ε , let $\beta^\varepsilon = \beta \cap Z^\varepsilon$, $\gamma^\varepsilon = \gamma \cap Z^\varepsilon$ and $\Lambda^\varepsilon = \begin{pmatrix} \beta^\varepsilon \\ \gamma^\varepsilon \end{pmatrix}$, and let \mathcal{G}^ε denote the family of Λ^ε . Then it is easily seen that $\Lambda^\varepsilon \in \mathcal{G}_{j^\varepsilon}^\varepsilon$, that $\mathbf{b}_d(\Lambda) - (\mathbf{b}_d(\Lambda^+) + \mathbf{b}_d(\Lambda^-))$ depends only on (\mathcal{G}, j) and that $\Lambda_j^{(d)}(Z)^\varepsilon = \Lambda_{j^\varepsilon}^{(d)}(Z^\varepsilon)$. By the induction hypothesis, $\mathbf{b}_d(\Lambda^\varepsilon) \geq \mathbf{b}_d(\Lambda_{j^\varepsilon}^{(d)}(Z^\varepsilon))$ with equality if and only if $\Lambda^\varepsilon = \Lambda_{j^\varepsilon}^{(d)}(Z^\varepsilon)$. So the result follows in this case. This means that we may, and we will, work under the following hypothesis:

Hypothesis. From now on, and until the end of this section, we assume that $j(z_1) = z_{2l}$.

As in the construction of $\Lambda_j^{(d)}(Z)$, let $Z' = Z \setminus \{z_1, z_{2l}\} = \{z_2, z_3, \dots, z_{2l-1}\}$, let j' denote the restriction of j to Z' and let

$$d' = \begin{cases} d-1 & \text{if } d \geq 1, \\ 1 & \text{if } d = 0. \end{cases}$$

Then $|Z'| = 2(l-1)$ and j' is 0-admissible. Let $\Lambda' = \begin{pmatrix} \beta' \\ \gamma' \end{pmatrix}$ where $\beta' = \beta \setminus \{z_1, z_{2l}\}$ and $\gamma' = \gamma \setminus \{z_1, z_{2l}\}$. Since $d \geq 2$, we have $z_1 \in \beta_j^{(d)}(Z)$ and $z_{2l} \in \gamma_j^{(d)}(Z)$. This implies that

$$(\star) \quad \mathbf{b}_d(\Lambda_j^{(d)}(Z)) = \mathbf{b}_{d-1}(\Lambda_{j'}^{(d-1)}(Z')) + z_{2l} + 2(l+d)z_1 + 2 \sum_{z \in Z'} z.$$

If $z_1 \in \beta$, then $\Lambda = \Lambda_j^{(d)}(Z)$ if and only if $\Lambda' = \Lambda_{j'}^{(d')}(Z')$ and again

$$\mathbf{b}_d(\Lambda) = \mathbf{b}_{d-1}(\Lambda') + z_{2l} + 2(l+d)z_1 + 2 \sum_{z \in Z'} z.$$

So the result follows from (\star) and from the induction hypothesis.

This means that we may, and we will, assume that $z_1 \in \gamma$. In this case,

$$\mathbf{b}_d(\Lambda) = \mathbf{b}_{d+1}(\Lambda') + 2dz_{2l} + (2l+1)z_1.$$

Then it follows from (\star) that

$$\mathbf{b}_d(\Lambda) - \mathbf{b}_d(\Lambda_j^{(d)}(Z)) = \mathbf{b}_{d+1}(\Lambda') - \mathbf{b}_{d-1}(\Lambda_{j'}^{(d-1)}(Z')) + (2d-1)(z_{2l} - z_1) - 2 \sum_{z \in Z'} z.$$

So, by the induction hypothesis,

$$\mathbf{b}_d(\Lambda) - \mathbf{b}_d(\Lambda_j^{(d)}(Z)) \geq \mathbf{b}_{d+1}(\Lambda_{j'}^{(d+1)}(Z')) - \mathbf{b}_{d-1}(\Lambda_{j'}^{(d-1)}(Z')) + (2d-1)(z_{2l} - z_1) - 2 \sum_{z \in Z'} z.$$

Since $z_{2l} - z_1 > z_{2l-1} - z_2$, it is sufficient to show that

$$(?) \quad \mathbf{b}_{d+1}(\Lambda_{j'}^{(d+1)}(Z')) - \mathbf{b}_{d-1}(\Lambda_{j'}^{(d-1)}(Z')) \geq -(2d-1)(z_{2l-1} - z_2) + 2 \sum_{z \in Z'} z.$$

This will be proved by induction on the size of Z' . First, if $j(z_2) < z_{2l}$, then we can separate Z' into two j' -stable subsets and a similar argument as before allows to conclude thanks to the induction hypothesis.

So we assume that $j'(z_2) = z_{2l-1}$. Let $Z'' = Z' \setminus \{z_2, z_{2l-1}\}$ and let j'' denote the restriction of j' to Z'' . Since $z_2 \in \beta_{j'}^{(d+1)}(Z')$, we can apply (\star) one step further to get

$$\begin{aligned} \mathbf{b}_{d+1}(\Lambda_{j'}^{(d+1)}(Z')) - \mathbf{b}_{d-1}(\Lambda_{j'}^{(d-1)}(Z')) &= \mathbf{b}_d(\Lambda_{j''}^{(d)}(Z'')) + z_{2l-1} + 2(l+d)z_2 + 2 \sum_{z \in Z''} z \\ &\quad - (\mathbf{b}_{d-2}(\Lambda_{j''}^{(d-2)}(Z'')) + z_{2l-1} + 2(l+d-2)z_2 + 2 \sum_{z \in Z''} z) \\ &= \mathbf{b}_d(\Lambda_{j''}^{(d)}(Z'')) - \mathbf{b}_{d-2}(\Lambda_{j''}^{(d-2)}(Z'')) + 4z_2. \end{aligned}$$

So, by the induction hypothesis,

$$\begin{aligned}
\mathbf{b}_{d+1}(\Lambda_{j'}^{(d+1)}(Z')) - \mathbf{b}_{d-1}(\Lambda_{j'}^{(d-1)}(Z')) &\geq -(2d-3)(z_{2l-2} - z_3) + 2 \sum_{z \in Z''} z + 4z_2 \\
&\geq -(2d-3)(z_{2l-1} - z_2) + 2 \sum_{z \in Z'} z + 2z_2 - 2z_{2l-1} \\
&= -(2d-1)(z_{2l-1} - z_2) + 2 \sum_{z \in Z'} z,
\end{aligned}$$

as desired. This shows (?) and completes the proof of Lemma 6⁺.

3. Complex reflection groups

If \mathcal{W} is a complex reflection group, then R. Rouquier and the author have also defined Calogero-Moser cellular characters and Calogero-Moser families (see [BoRo1] or [BoRo2]). If \mathcal{W} is of type $G(l, 1, n)$ (in Shephard-Todd classification), then Leclerc and Miyachi [LeMi, §6.3] proposed, in link with canonical bases of $U_v(\mathfrak{sl}_\infty)$ -modules, a family of characters that could be good analogue of constructible characters: let us call them the *Leclerc-Miyachi constructible characters* of $G(l, 1, n)$. If $l = 2$, then they coincide with constructible characters [LeMi, Theorem 10].

Of course, it would be interesting to know if Calogero-Moser cellular characters coincide with the Leclerc-Miyachi ones: this seems rather complicated but it should be at least possible to check if the Leclerc-Miyachi constructible characters satisfy the analogous properties with respect to the b -invariant.

References

- [Be1] G. BELLAMY, Generalized Calogero-Moser spaces and rational Cherednik algebras, PhD thesis, University of Edinburgh (2010).
- [Be2] G. BELLAMY, The Calogero-Moser partition for $G(m, d, n)$, *Nagoya Math. J.* **207** (2012), 47-77.
- [BoRo1] C. BONNAFÉ & R. ROUQUIER, Calogero-Moser versus Kazhdan-Lusztig cells, to appear in *Pacific J. Math.*, preprint (2012), [arXiv:1201.0585](#).
- [BoRo2] C. BONNAFÉ & R. ROUQUIER, Cellules de Calogero-Moser, preprint (2013), available on [arXiv](#).
- [Ge] M. GECK, Computing Kazhdan-Lusztig cells for unequal parameters *J. Algebra* **281** (2004) 342-365.
- [GePf] M. GECK & G. PFEIFFER, *Characters of finite Coxeter groups and Iwahori-Hecke algebras*, London Mathematical Society Monographs, New Series **21**, The Clarendon Press, Oxford University Press, New York, 2000, xvi+446 pp.
- [GoMa] I. G. GORDON & M. MARTINO, Calogero-Moser space, restricted rational Cherednik algebras and two-sided cells, *Math. Res. Lett.* **16** (2009), 255-262.

- [LeMi] B. LECLERC & H. MIYACHI, Constructible characters and canonical bases, *J. Algebra* **277** (2004), 298-317.
- [Lu1] G. LUSZTIG, A class of irreducible representations of a Weyl group, *Indag. Math.* **41** (1979), 323-335.
- [Lu2] G. LUSZTIG, *Characters of reductive groups over a finite field*, Annals of Mathematics Studies, **107**. Princeton University Press, Princeton, NJ, 1984, xxi+384 pp.
- [Lu3] G. LUSZTIG, *Hecke algebras with unequal parameters*, CRM Monograph Series **18**, American Mathematical Society, Providence, RI (2003), 136 pp.

February 12, 2013

CÉDRIC BONNAFÉ, Institut de Mathématiques et de Modélisation de Montpellier (CNRS: UMR 5149), Université Montpellier 2, Case Courrier 051, Place Eugène Bataillon, 34095 MONTPELLIER Cedex, FRANCE • E-mail : `cedric.bonnafe@univ-montp2.fr`