



HAL
open science

Poisson ensembles of loops of one-dimensional diffusions

Titus Lupu

► **To cite this version:**

| Titus Lupu. Poisson ensembles of loops of one-dimensional diffusions. 2013. hal-00788901v3

HAL Id: hal-00788901

<https://hal.science/hal-00788901v3>

Preprint submitted on 11 Jan 2014 (v3), last revised 7 Jan 2019 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Titus Lupu

**POISSON ENSEMBLES OF LOOPS OF
ONE-DIMENSIONAL DIFFUSIONS**

Titus Lupu

Laboratoire de Mathématiques, Bât.425, Université Paris-Sud XI, 91405 Orsay
(France).

E-mail : `titus.lupu@math.u-psud.fr`

2000 Mathematics Subject Classification. — 60-02, 60G15, 60G17, 60G55, 60G60, 60J55, 60J60, 60J65, 60J80.

Key words and phrases. — Poisson ensembles of Markov loops, "loop soup", one-dimensional diffusion processes, Vervaat's transformation, continuous state branching processes with immigration, Gaussian Free Field, Dynkin's isomorphism, Wilson's algorithm, Uniform Spanning Tree, determinantal point processes .

POISSON ENSEMBLES OF LOOPS OF ONE-DIMENSIONAL DIFFUSIONS

Titus Lupu

Abstract. — We study the analogue of Poisson ensembles of Markov loops (“loop soups”) in the setting of one-dimensional diffusions. We give a detailed description of the corresponding intensity measure. The properties of this measure on loops lead us to an extension of Vervaat’s bridge-to-excursion transformation that relates the bridges conditioned by their minimum and the excursions of all the diffusion we consider and not just the Brownian motion. Further we describe the Poisson point process of loops, their occupation fields and explain how to sample these Poisson ensembles of loops using two-dimensional Markov processes. Finally we introduce a couple of interwoven determinantal point processes on the line which is a dual through Wilson’s algorithm of Poisson ensembles of loops and study the properties of these determinantal point processes.

Résumé (Ensemble poissonien de boucles des diffusions unidimensionnelles)

Nous étudions l’analogie des ensembles poissonniens de boucles markoviennes (“loop soups”) dans le cadre des diffusions unidimensionnelles. Nous donnons une description détaillée de la mesure d’intensité correspondante. Les propriétés de cette mesure sur les boucles nous amènent à une extension de la transformation de Vervaat pont-excursion qui relie les ponts conditionnés par leur minimum et les excursions de toutes les diffusions que nous considérons et non juste ceux du mouvement Brownien. Ensuite nous décrivons le processus ponctuels de Poisson des boucles, leurs champs d’occupation et expliquons comment séquencer ces ensembles poissonniens de boucles à partir de processus de Markov bidimensionnels. Enfin nous introduisons un couple de processus ponctuels déterminantaux sur la droite, entrelacés, qui est un dual, à travers l’algorithme de Wilson, de l’ensemble poissonien de boucles, et étudions les propriétés de ces processus ponctuels déterminantaux.

CONTENTS

1. Introduction	1
2. Preliminaries on generators and semi-groups	5
2.1. A second order ODE.....	5
2.2. One-dimensional diffusions.....	11
2.3. "Generators" with creation of mass.....	15
3. Measure on loops and its basic properties	23
3.1. Spaces of loops.....	23
3.2. Measures $\mu^{x,y}$ on finite life-time paths.....	24
3.3. The measure μ^* on unrooted loops.....	31
3.4. Multiple local times.....	34
3.5. A disintegration of μ^* induced by the Vervaat's transformation	38
3.6. A generalization of the Vervaat's transformation.....	41
3.7. Restricting loops to a discrete subset.....	46
3.8. Measure on loops in case of creation of mass.....	48
4. Occupation fields of the Poisson ensembles of Markov loops	51
4.1. Inhomogeneous continuous state branching processes with immigration ..	51
4.2. Occupation field.....	53
4.3. Dynkin's isomorphism.....	64
5. Decomposing paths into Poisson ensembles of loops	67
5.1. Glueing together excursions ordered by their minima.....	67
5.2. Loops represented as excursions and glued together.....	69
5.3. The case $\alpha = 1$	76
6. Wilson's algorithm in dimension one	81
6.1. Description of the algorithm.....	81
6.2. The erased paths.....	84
6.3. Determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$: Brownian case.....	85

6.4. Determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$: general case.....	104
7. Monotone couplings for the point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$.....	107
7.1. Conditioning.....	107
7.2. Couplings.....	121
Bibliography.....	139

CHAPTER 1

INTRODUCTION

Lawler and Werner introduced in [19] the notion of Poisson ensemble of Markov loops ("loop soup") for planar Brownian motion. In [26] it was used by Sheffield and Werner to construct the Conformal Loops Ensemble (CLE). Le Jan studied in [13] the analogue of the Poissonian ensembles of Markov loops in the setting of a symmetric Markov jump process on a finite graph. In both cases one defines an infinite measure μ^* on time-parametrized unrooted loops (i.e. loops parametrized by a circle where it is not specified when the cut between the beginning and the end occurs) and considers the Poisson point ensemble of intensity $\alpha\mu^*$, $\alpha > 0$, denoted here \mathcal{L}_α . In both cases the ensemble \mathcal{L}_1 (where $\alpha = 1$) is related to the loops erased during the loop-erasure procedure applied to Markovian sample paths. In particular in the discrete setting Wilson's algorithm ([31]) leads to a duality between \mathcal{L}_1 and the Uniform Spanning Trees. In [13] Le Jan also studied the occupation field of \mathcal{L}_α , that is the sum of the occupation times in a given vertex of the graph of individual loops. In case $\alpha = \frac{1}{2}$ he found that it is the square of a Gaussian Free Field and related it to the Dynkin's Isomorphism ([8]).

The analogue of the measure μ^* can be defined for a much larger class of Markov processes ([15], [10]). The aim of this essay is on one hand to study the measure μ^* and the Poisson ensembles of Markov loops \mathcal{L}_α in the setting of one-dimensional, not necessarily conservative, diffusion processes, and on the other hand to define and study some determinantal point processes on \mathbb{R} that are analogous to Uniform Spanning Trees and dual to \mathcal{L}_1 . The diffusion processes we consider take values on a subinterval I of \mathbb{R} , are always killed at hitting a boundary point of I , and may be killed by a killing measure on the interior of I . One can transform a diffusion process into another applying a change of scale, a random change of time, a restriction to a subinterval, an increase of the killing measure or an h-transform. The measure μ^* is covariant with all these transformations on Markov processes. In other words the map diffusion to measure on loops is a covariant functor. Moreover we will show that μ^*

is invariant by h-transform on underlying diffusions. We will also extend the scope of our study by associating a measure on loops to "generators" which contain a creation of mass term: If $L = L^{(0)} + \nu$ where $L^{(0)}$ is a second order differential operator on I and ν is a signed measure, and if one sets zero Dirichlet boundary conditions for L , one can define in a consistent way a measure on loops related to L even in case the semi-group $(e^{tL})_{t \geq 0}$ does not make sense. This extended definition of μ^* will be particularly handy for computing the exponential moments of the Poissonian ensemble of Markov loops.

The layout of this paper is the following: In chapter 2 we will recall some facts on one-dimensional diffusions and set the important notations. We will further consider "generators" with creation of mass term and characterize a class of such operators which up to an h-transform are equivalent to the generators of diffusions. In chapter 3 we will define the measure μ^* and point out different covariance and invariance properties. Further we will make a connection between the Brownian measure on loops and the Levy-Itô measure on Brownian excursion using the Vervaat's bridge-to-excursion transformation. This in turn will lead us to a conditioned version of Vervaat's transformation that holds for any one-dimensional diffusion process, that is an absolute continuity relation between the bridge conditioned to have a given minimum and an excursion of the same duration above this minimum. The Vervaat's transformation is deeply related to the measure on loops μ^* : The loops are unrooted, so one can freely chose a moment separating the end from the start. If one chooses this moment uniformly over the life-time of the loop, then the loop under the measure μ^* looks in some sense like a bridge. If one chooses this moment when the loop hits its minimum, then it looks like an excursion. In chapter 4 we will study the occupation field of the Poisson ensemble of Markov loops. Each loop is endowed with a family of local times. The occupation field is the sum of local times over the loops. We will identify its law as an non-homogeneous continuous state branching process with immigration parametrised by the position points in I . In case $\alpha = \frac{1}{2}$ we will identify it as the square of a Gaussian Free Field and show how it is possible to derive particular versions of the Dynkin's Isomorphism using this fact and Palm's identity for Poissonian ensembles. In chapter 5 we will root each loop in \mathcal{L}_α at its minimum and obtain this way a collection of positive excursions. Then we will order this excursions in the decreasing sense of their minima and glue them together. We will obtain this way a continuous path which can be described using two-dimensional Markov processes. This is a way to sample \mathcal{L}_α . In the particular case $\alpha = 1$ the path we obtain is the sample path of an one-dimensional diffusion. This is the analogue in our setting of the relation between \mathcal{L}_1 and the loop-erasure procedure observed in the setting of the two-dimensional Brownian motion or of the symmetric Markov jump processes on graphs. In chapter 6 we will apply an extension of Wilson's algorithm to transient one-dimensional diffusions and obtain a couple of interwoven determinantal

point processes on \mathbb{R} which is dual to \mathcal{L}_1 . In chapter 7 we will prove some monotone coupling properties for the determinantal point processes introduced in chapter 6.

The author thanks Yves Le Jan for fruitful discussions and its helpful advice in relation with this work.

CHAPTER 2

PRELIMINARIES ON GENERATORS AND SEMI-GROUPS

2.1. A second order ODE

In this chapter we will introduce the one-dimensional diffusions we will consider throughout this work (section 2.2). In the section 2.3 we will extend the framework to the "generators" containing a mass-creation term. In the section 2.1 we will prove or recall some facts on the functions harmonic for these generators.

Let I be an open interval of \mathbb{R} and ν a signed measure on I . By signed measure we mean that the total variation $|\nu|$ is a positive Radon measure, but not necessarily finite, and $\nu(dx) = \epsilon(x)|\nu|(dx)$ where ϵ takes values in $\{\pm 1\}$. We look for the solutions of the linear second order differential equation on I :

$$(2.1.1) \quad \frac{d^2 u}{dx^2} + u\nu = 0$$

Given a solution u of (2.1.1) we will write $\frac{du}{dx}(x^+)$ and $\frac{du}{dx}(x^-)$ for the right-hand side respectively left-hand side derivative of u at x . The two are related by

$$\frac{du}{dx}(x^+) - \frac{du}{dx}(x^-) = -u(x)\nu(\{x\})$$

Using a standard fixed point argument one can show that (2.1.1) satisfies a Cauchy-Lipschitz principle: if $x_0 \in I$ and $u_0, v_0 \in \mathbb{R}$, there is a unique solution u of (2.1.1), continuous on I , satisfying $u(x_0) = u_0$ and $\frac{du}{dx}(x_0^+) = v_0$. Let $x_1 \in I \cap (x_0, +\infty)$. A continuous function u on $[x_0, x_1]$ is solution of (2.1.1) with previous initial conditions at x_0 if and only if it is a fixed point of the affine operator \mathfrak{J} on $\mathcal{C}([x_0, x_1])$ defined as

$$(\mathfrak{J}u)(x) := u_0 + (x - x_0)v_0 - \int_{(x_0, x]} (x - y)u(y)\nu(dy)$$

The Lipschitz norm of \mathfrak{J}^n is smaller or equal to $\frac{|\nu|([x_0, x_1])^n (x_1 - x_0)^n}{n!}$. So for n large enough \mathfrak{J}^n is contracting and thus \mathfrak{J} has a unique fixed point in $\mathcal{C}([x_0, x_1])$.

Let $W(u_1, u_2)(x)$ be the Wronskian of two functions u_1, u_2 :

$$W(u_1, u_2)(x) := u_1(x) \frac{du_2}{dx}(x^+) - u_2(x) \frac{du_1}{dx}(x^+)$$

If u_1, u_2 are both solutions of (2.1.1), $W(u_1, u_2)$ is constant on I . Using this fact we get a results which is similar to Sturm's separation theorem for the case of a measure ν with a continuous density with respect to the Lebesgue measure (see theorem 7, section 2.6 in [4]):

Property 2.1. — Given $x_0 < x_1$ be two points in I :

- (i) Let u_1 be a solution of (2.1.1) satisfying $u_1(x_0) = 0$, $\frac{du_1}{dx}(x_0^+) > 0$, and u_2 a solution such that $u_2(x_0) > 0$. Assume that $u_2 \geq 0$ on $[x_0, x_1]$. Then $u_1 > 0$ on $(x_0, x_1]$.
- (ii) Let u_1, u_2 be two solutions such that $u_1(x_0) = u_2(x_0) > 0$ and $\frac{du_1}{dx}(x_0^+) > \frac{du_2}{dx}(x_0^+)$. Assume that $u_2 \geq 0$ on $[x_0, x_1]$. Then $u_1 > u_2$ on $(x_0, x_1]$.
- (iii) If there is a solution u to (2.1.1) positive on (x_0, x_1) and zero at x_0 and x_1 then any other linearly independent solution of (2.1.1) has exactly one zero in (x_0, x_1) .

Next we prove a lemma that will be useful in the section 2.3.

Lemma 2.2. — Let ν_+ be the positive part of ν . Let $x_0 < x_1 \in I$. Let f be a continuous positive function on $[x_0, x_1]$ such that $\min_{[x_0, x_1]} f > \nu_+([x_0, x_1])^2$. Then the equation

$$(2.1.2) \quad \frac{d^2u}{dx^2} + u\nu - uf = 0$$

has a positive solution that is non-decreasing on $[x_0, x_1]$.

Proof. — Set $a := \min_{[x_0, x_1]} f$. Let u be the solution to (2.1.2) with the initial values $u(x_0) = 1$, $\frac{du}{dx}(x_0^+) = \sqrt{a}$. We will show that u is non-decreasing on $[x_0, x_1]$. Assume that this is not the case. This means that $\frac{du}{dx}(x^+)$ takes negative values somewhere in $[x_0, x_1]$. Let

$$x_2 := \inf \left\{ x \in [x_0, x_1] \mid \frac{du}{dx}(x^+) \leq 0 \right\}$$

Since $\frac{du}{dx}(x^+)$ is right-continuous, $\frac{du}{dx}(x_2^+) \leq 0$. Let $r(x) := \frac{1}{u(x)} \frac{du}{dx}(x^+)$. u is positive on $[x_0, x_2]$ hence r is defined $[x_0, x_2]$. $r(x_0) = \sqrt{a}$. r is cadlag and satisfies the equation

$$dr = (f - r^2)dx - d\nu$$

Let $x_3 := \sup\{x \in [x_0, x_2] \mid r(x) \geq \sqrt{a}\}$. We have

$$r(x_2) = r(x_3^-) + \int_{x_3}^{x_2} (f(x) - r^2(x))dx - \nu([x_3, x_2])$$

By construction $r(x_3^-) \geq \sqrt{a}$. By definition $f - r^2 \geq 0$ on $(x_3, x_2]$. Thus

$$r(x_2) \geq \sqrt{a} - \nu([x_3, x_2]) > 0$$

It follows that $r(x_2) > 0$, which is absurd. \square

In the case $\nu = -2\kappa$ where κ is a non-zero positive Radon measure, the equation (2.1.1) becomes:

$$(2.1.3) \quad \frac{1}{2} \frac{d^2 u}{dx^2} - u\kappa = 0$$

It commonly appears when studying the Brownian motion with a killing measure κ . In this case the two-dimensional linear space of solutions is spanned by two convex positive solutions u_\uparrow and u_\downarrow , u_\uparrow being non-decreasing and u_\downarrow non-increasing. Given $x_0 \in I$, we can construct u_\uparrow as the limit when $x_1 \rightarrow \inf I$ of the unique solution which equals 0 in x_1 and 1 in x_0 . For u_\downarrow we take the limit as $x_1 \rightarrow \sup I$. u_\uparrow and u_\downarrow are defined up to a positive multiplicative constant. See [5], section 16.11, or [24], Appendix 8, for more details. Next we give equivalent conditions on the asymptotic behaviour of u_\uparrow and u_\downarrow that will be used in chapter 6.

Proposition 2.3. — *In case $[0, +\infty) \subseteq I$, the following four conditions are equivalent:*

- (i) $\int_{(0, +\infty)} x\kappa(dx) < +\infty$
- (ii) $u_\downarrow(+\infty) > 0$
- (iii) *There is $C > 0$ such that for all $x \geq 1$, $u_\uparrow(x) \leq Cx$*
- (iv) $\int_{(0, +\infty)} u_\uparrow(x)u_\downarrow(x)\kappa(dx) < +\infty$

Proof. — We will prove in order that (ii) implies (i), (iii) implies (i), (i) implies (ii), (i) implies (iii) and (iv) implies (ii). (iv) is obviously implied by the combination of (i), (ii) and (iii).

(ii) implies (i): For all $x \in [0, +\infty)$:

$$-\frac{du_\downarrow}{dx}(x^+) = 2 \int_{(x, +\infty)} u_\downarrow(y)\kappa(dy) \leq 2u_\downarrow(+\infty)\kappa((x, +\infty))$$

$-\frac{du_\downarrow}{dx}(x^+)$ is integrable on $(0, +\infty)$. Since $u_\downarrow(+\infty) > 0$, this implies that:

$$\int_{(0, +\infty)} \kappa((x, +\infty))dx < +\infty$$

But

$$\int_{(0, +\infty)} \kappa((x, +\infty))dx = \int_{(0, +\infty)} y\kappa(dy)$$

and hence (i).

(iii) implies (i): If (iii) holds then for all $x \in [0, +\infty)$, $\frac{du_\uparrow}{dx}(x^+) \leq C$. But

$$\frac{du_\uparrow}{dx}(x^+) = \frac{du_\uparrow}{dx}(0^+) + 2 \int_{(0, x]} u_\uparrow(y)\kappa(dy)$$

This implies that

$$\int_{(0,+\infty)} u_{\uparrow}(y)\kappa(dy) < +\infty$$

Since u_{\uparrow} is convex, $u_{\uparrow}(y) \geq u_{\uparrow}(0) + \frac{du_{\uparrow}}{dy}(0^+)y$. So (i) is satisfied.

(i) implies (ii): For all $y \in [0, +\infty)$:

$$u_{\downarrow}(y) - u_{\downarrow}(+\infty) = 2 \int_y^{+\infty} \int_{(z,+\infty)} u_{\downarrow}(x)\kappa(dx)dz \leq 2u_{\downarrow}(y) \int_{(y,+\infty)} (x-y)\kappa(dx)$$

Condition (i) implies that:

$$\lim_{y \rightarrow +\infty} 2 \int_{(y,+\infty)} (x-y)\kappa(dx) = 0$$

So for y large enough, $u_{\downarrow}(y) - u_{\downarrow}(+\infty) < u_{\downarrow}(y)$. Necessarily $u_{\downarrow}(+\infty) > 0$.

(i) implies (iii): For all $y < x \in [0, +\infty)$:

$$(2.1.4) \quad \frac{du_{\uparrow}}{dx}(x^+) = \frac{du_{\uparrow}}{dy}(y^+) + 2u_{\uparrow}(y)\kappa((y, x]) + 2 \int_{(y,x]} (u_{\uparrow}(z) - u_{\uparrow}(y))\kappa(dz)$$

Let y be large enough such that:

$$2 \int_{(y,+\infty)} (z-y)\kappa(dz) < 1$$

Then there is $C > 0$ large enough such that:

$$(2.1.5) \quad C > \frac{du_{\uparrow}}{dy}(y^+) + 2u_{\uparrow}(y)\kappa((y, +\infty)) + 2C \int_{(y,+\infty)} (z-y)\kappa(dz)$$

Assume that there is $x \in [0, +\infty)$ such that $\frac{du_{\uparrow}}{dx}(x^+) \geq C$. Let

$$x_0 := \inf \left\{ x \geq y \mid \frac{du_{\uparrow}}{dx}(x^+) \geq C \right\}$$

$x \mapsto \frac{du_{\uparrow}}{dx}(x^+)$ is right-continuous. Thus $\frac{du_{\uparrow}}{dx}(x_0^+) \geq C$. By definition, for all $z \in [y, x_0]$, $\frac{du_{\uparrow}}{dz}(z^+) \leq C$ and hence $u_{\uparrow}(z) - u_{\uparrow}(y) \leq C(z-y)$. But then (2.1.4) and (2.1.5) imply that $\frac{du_{\uparrow}}{dx}(x_0^+) < C$ which is contradictory. It follows that $\frac{du_{\uparrow}}{dx}(x^+)$ is bounded by C , which implies property (iii).

(iv) implies (ii): Applying integration by parts we get that for all $x > 0$:

$$\begin{aligned} 2 \int_{(0,x]} u_{\uparrow}(y)u_{\downarrow}(y)\kappa(dy) &= \int_{(0,x]} u_{\downarrow}(y)d\left(\frac{du_{\uparrow}}{dy}\right)(dy) \\ &= \frac{du_{\uparrow}}{dx}(x^+)u_{\downarrow}(x) - \frac{du_{\uparrow}}{dx}(0^+)u_{\downarrow}(0) - \int_0^x \frac{du_{\downarrow}}{dy}(y^+)\frac{du_{\uparrow}}{dy}(y^+)dy \end{aligned}$$

$\frac{du_{\uparrow}}{dx}(x^+)u_{\downarrow}(x)$ is positive. We get that:

$$(2.1.6) \quad - \int_0^{+\infty} \frac{du_{\downarrow}}{dy}(y^+)\frac{du_{\uparrow}}{dy}(y^+)dy \leq 2 \int_{(0,+\infty)} u_{\uparrow}(y)u_{\downarrow}(y)\kappa(dy) + \frac{du_{\uparrow}}{dx}(0^+)u_{\downarrow}(0) < +\infty$$

Next

$$(2.1.7) \quad \begin{aligned} \frac{du_\uparrow}{dx}(x^+)(u_\downarrow(x) - u_\downarrow(+\infty)) &= -\frac{du_\uparrow}{dx}(x^+) \int_x^{+\infty} \frac{du_\downarrow}{dy}(y^+) dy \\ &\leq -\int_x^{+\infty} \frac{du_\downarrow}{dy}(y^+) \frac{du_\uparrow}{dy}(y^+) dy \end{aligned}$$

Assume that $u_\downarrow(+\infty) = 0$. Then (2.1.7) implies that:

$$\lim_{x \rightarrow +\infty} \frac{du_\uparrow}{dx}(x^+) u_\downarrow(x) = 0$$

and

$$(2.1.8) \quad \lim_{x \rightarrow +\infty} -\frac{du_\downarrow}{dx}(x^+) u_\uparrow(x) = W(u_\downarrow, u_\uparrow) - \lim_{x \rightarrow +\infty} \frac{du_\uparrow}{dx}(x^+) u_\downarrow(x) = W(u_\downarrow, u_\uparrow)$$

(2.1.6) together with (2.1.8) imply that

$$\int_0^{+\infty} \frac{1}{u_\uparrow(y)} \frac{du_\uparrow}{dy}(y^+) dy < +\infty$$

But this is impossible because $\log(u_\uparrow(+\infty)) = +\infty$. Thus $u_\downarrow(+\infty) > 0$. \square

Next we deal with the continuity of u_\uparrow and u_\downarrow with respect the measure κ . We will write $u_{\kappa, \uparrow}$ and $u_{\kappa, \downarrow}$ to denote the dependence on κ .

Lemma 2.4. — *Let $x_0 \in I$. Let $(\kappa_n)_{n \geq 0}$ be a sequence of non-zero positive Radon measures on I converging vaguely (i.e. against functions with compact support) to κ . Then $\frac{u_{\kappa_n, \uparrow}}{u_{\kappa_n, \uparrow}(x_0)}$ converges to $\frac{u_{\kappa, \uparrow}}{u_{\kappa, \uparrow}(x_0)}$, $\frac{u_{\kappa_n, \downarrow}}{u_{\kappa_n, \downarrow}(x_0)}$ converges to $\frac{u_{\kappa, \downarrow}}{u_{\kappa, \downarrow}(x_0)}$ and the convergences are uniform on compact subsets of I .*

Proof. — We will deal with the convergence of $\frac{u_{\kappa_n, \downarrow}}{u_{\kappa_n, \downarrow}(x_0)}$, the other one being similar. To simplify notations we will chose the normalization $u_{\kappa, \downarrow}(x_0) = u_{\kappa_n, \downarrow}(x_0) = 1$. Without loss of generality we will also assume that $\kappa(\{x_0\}) = 0$. The proof will be made of two parts. First we will show that if u is the solution of (2.1.3) and u_n solution of

$$(2.1.9) \quad \frac{1}{2} \frac{d^2 u}{dx^2} - u \kappa_n = 0$$

and if $u_n(x_0) = u(x_0) = 1$ and $\frac{du}{dx}(x_0^+) = \lim_{n \rightarrow +\infty} \frac{du_n}{dx}(x_0^+)$ then u_n converges to u uniformly on compact subsets of I . After that we will show that $\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+)$ converges to $\frac{du_{\kappa, \downarrow}}{dx}(x_0^+)$.

Let $x_1 \in I \cap (x_0, +\infty)$. Let $(v_n)_{n \geq 0}$ be a sequence in \mathbb{R} converging to v . Let \mathfrak{J}_n respectively \mathfrak{J} be the following affine operators on $\mathcal{C}([x_0, x_1])$:

$$\begin{aligned} (\mathfrak{J}_n f)(x) &:= 1 + (x - x_0)v_n + 2 \int_{(x_0, x]} (x - y)f(y)\kappa_n(dy) \\ (\mathfrak{J} f)(x) &:= 1 + (x - x_0)v + 2 \int_{(x_0, x]} (x - y)f(y)\kappa(dy) \end{aligned}$$

Let u_n respectively u be the fixed points of \mathfrak{J}_n respectively \mathfrak{J} . Let $\varepsilon \in (0, 1)$. The Lipschitz norm of \mathfrak{J}_n^j is bounded by $\frac{2^j}{j!} \kappa_n([x_0, x_1])^j (x_1 - x_0)^j$. For $j \geq j_\varepsilon$, for all $n \in \mathbb{N}$, this norm is less than ε . Then

$$\begin{aligned} \max_{[x_0, x_1]} |u_n - u| &= \max_{[x_0, x_1]} |\mathfrak{J}_n^{j_\varepsilon} u_n - \mathfrak{J}^{j_\varepsilon} u| \leq \max_{[x_0, x_1]} |\mathfrak{J}_n^{j_\varepsilon} u - \mathfrak{J}^{j_\varepsilon} u| + \max_{[x_0, x_1]} |\mathfrak{J}_n^{j_\varepsilon} u_n - \mathfrak{J}_n^{j_\varepsilon} u| \\ &\leq \max_{[x_0, x_1]} |\mathfrak{J}_n^{j_\varepsilon} u - \mathfrak{J}^{j_\varepsilon} u| + \varepsilon \max_{[x_0, x_1]} |u_n - u| \end{aligned}$$

Hence

$$(2.1.10) \quad \max_{[x_0, x_1]} |u_n - u| \leq \frac{1}{1 - \varepsilon} \max_{[x_0, x_1]} |\mathfrak{J}_n^{j_\varepsilon} u - \mathfrak{J}^{j_\varepsilon} u|$$

For $y < x \in I$ and $i \in \mathbb{N}^*$ let

$$\begin{aligned} f_{n,i}(y, x) &:= \int_{y < y_1 < \dots < y_{i-1} < x} (x - y_{i-1}) \dots (y_2 - y_1)(y_1 - y) \kappa_n(dy_1) \dots \kappa_n(dy_{i-1}) \\ f_i(y, x) &:= \int_{y < y_1 < \dots < y_{i-1} < x} (x - y_{i-1}) \dots (y_2 - y_1)(y_1 - y) \kappa(dy_1) \dots \kappa(dy_{i-1}) \end{aligned}$$

and $f_{0,i}(y, x) = f_0(y, x) = x - y$. $f_{n,i}$ and f_i are continuous functions. Moreover the vague convergence of κ_n to κ ensures that if $(y_n, x_n)_{n \geq 0}$ is a sequence converging to (y, x) then $f_{n,i}(y_n, x_n)$ converges to $f_i(y, x)$.

$$\begin{aligned} (\mathfrak{J}_n^{j_\varepsilon} u)(x) &= 1 + (x - x_0)v_n + \sum_{i=0}^{j_\varepsilon-2} \int_{x_0}^x (1 + (y - x_0)v_n) f_{n,i}(y, x) \kappa_n(dy) \\ &\quad + \int_{x_0}^x u(y) f_{n, j_\varepsilon-1}(y, x) \kappa_n(dy) \\ (\mathfrak{J}^{j_\varepsilon} u)(x) &= 1 + (x - x_0)v + \sum_{i=0}^{j_\varepsilon-2} \int_{x_0}^x (1 + (y - x_0)v) f_i(y, x) \kappa(dy) \\ &\quad + \int_{x_0}^x u(y) f_{j_\varepsilon-1}(y, x) \kappa(dy) \end{aligned}$$

For fixed x , the functions $y \mapsto 1_{x_0 < y < x} f_{n,i}(y, x)$ and $y \mapsto 1_{x_0 < y < x} f_i(y, x)$ have a compact support but are discontinuous at x_0 . If $(z_n)_{n \geq 0}$ is a sequence in $[x_0, x_1]$ converging to z , then the convergence of v_n to v , the weak convergence of κ_n to κ and the condition $\kappa(\{x_0\}) = 0$ ensure that $(\mathfrak{J}_n^{j_\varepsilon} u)(z_n)$ converges to $(\mathfrak{J}^{j_\varepsilon} u)(z)$. This implies the uniform convergence of $\mathfrak{J}_n^{j_\varepsilon} u$ to $\mathfrak{J}^{j_\varepsilon} u$ on $[x_0, x_1]$. From (2.1.10) follows that u_n converges uniformly to u on $[x_0, x_1]$. The situation is similar for $x_1 < x_0$ and we get the uniform convergence on compact sets of u_n to u .

Let

$$\underline{v} := \liminf_{n \rightarrow +\infty} \frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) \quad \bar{v} := \limsup_{n \rightarrow +\infty} \frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+)$$

Let $v < \frac{du_{\kappa, \downarrow}}{dx}(x_0^+)$. There is $x_1 \in I \cap (x_0, +\infty)$ such that the solution of (2.1.3) with initial conditions $u(x_0) = 1$, $\frac{du}{dx}(x^+) = v$ is zero at x_1 since $u_{\kappa_n, \downarrow}$ converges to $u_{\kappa, \downarrow}$

uniformly on $[x_0, x_1]$ and $u_{\kappa, \downarrow}$ is positive on $[x_0, x_1]$, we get that for n large enough, $u_{\kappa_n, \downarrow}$ is positive on $[x_0, x_1]$ and $\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) > v$. Thus $\underline{v} \geq \frac{du_{\kappa, \downarrow}}{dx}(x_0^+)$.

Conversely, let $v < \bar{v}$. Let u_n be the solution of (2.1.9) with initial conditions $u_n(x_0) = 1$, $\frac{du_n}{dx}(x_0^+) = v$. If $\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) > v$, then for any $x \in I \cap [x_0, +\infty)$

$$\begin{aligned} \frac{du_n}{dx}(x^+) &\leq \frac{du_{\kappa_n, \downarrow}}{dx}(x^+) - \left(\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) - v \right) \leq - \left(\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) - v \right) \\ u_n(x) &\leq u_{\kappa_n, \downarrow} - \left(\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) - v \right) (x - x_0) \end{aligned}$$

If $\sup I < +\infty$ then by convexity of $u_{\kappa_n, \downarrow}$:

$$u_n(x) \leq \frac{\sup I - x}{\sup I - x_0} - \left(\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) - v \right) (x - x_0)$$

and $u_n(z_n) \leq 0$ where

$$z_n := \frac{\sup I + \left(\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) - v \right) x_0 (\sup I - x_0)}{1 + \left(\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) - v \right) (\sup I - x_0)}$$

This is also true if $\sup I = +\infty$ and in this case $z_n = x_0 + \left(\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+) - v \right)^{-1}$. Let u be the solution of (2.1.3) with initial conditions $u(x_0) = 1$, $\frac{du}{dx}(x^+) = v$ and

$$z_\infty := \frac{\sup I + (\bar{v} - v)x_0(\sup I - x_0)}{1 + (\bar{v} - v)(\sup I - x_0)}$$

Considering a subsequence along which $\frac{du_{\kappa_n, \downarrow}}{dx}(x_0^+)$ converges to \bar{v} , we get by uniform convergence of u_n to u on compact sets that $u(z_\infty) \geq 0$. It follows that $\frac{du_{\kappa, \downarrow}}{dx}(x_0^+) \geq v$. Hence $\frac{du_{\kappa, \downarrow}}{dx}(x_0^+) \geq \bar{v}$.

Finally $\underline{v} = \bar{v} = \frac{du_{\kappa, \downarrow}}{dx}(x_0^+)$ and this implies the uniform convergence on compact subsets of $u_{\kappa_n, \downarrow}$ to $u_{\kappa, \downarrow}$. \square

2.2. One-dimensional diffusions

In this subsection we will describe the kind of linear diffusion we are interested in, recall some facts and introduce notations that will be used subsequently. For a detailed presentation of one-dimensional diffusions see [12] and [5], chapter 16.

Let I be an open interval of \mathbb{R} , m and w continuous positive functions on I . We consider a diffusion $(X_t)_{0 \leq t < \zeta^{(0)}}$ on I with generator

$$L^{(0)} := \frac{1}{m(x)} \frac{d}{dx} \left(\frac{1}{w(x)} \frac{d}{dx} \right)$$

and killed as it hits the boundary of I . In case I is unbounded, we also allow for X to blow up to infinity in finite time. $\zeta^{(0)}$ is the first time X either hits the boundary or explodes. To avoid some technicalities we will assume that $\frac{dw}{dx}$ is locally bounded, although this condition is not essential. Given such a diffusion, the speed measure

$m(x)dx$ and the scale measure $w(x)dx$ are defined up to a positive multiplicative constant, but the product mw is uniquely defined. A primitive S of w is a natural scale function of X . Consider the random time change $d\tilde{t} = \frac{1}{m(X_t)}dt$. Then $(\frac{1}{2}S(X_{\tilde{t}}))_{0 \leq \tilde{t} < \tilde{\zeta}(0)}$ is a standard Brownian motion on $S(I)$ killed when it first hits the boundary of $S(I)$. For all f, g smooth, compactly supported in I ,

$$\int_I (L^{(0)}f)(x)g(x)m(x)dx = \int_I f(x)(L^{(0)}g)(x)m(x)dx$$

The diffusion X has a family of local times $(\ell_t^x(X))_{x \in I, t \geq 0}$ with respect to the measure $m(x)dx$ such that $(x, t) \mapsto \ell_t^x(X)$ is continuous. We can further consider diffusions with killing measures. Let κ be a non-negative Radon measure on I . We kill X as soon as $\int_I \ell_t^x(X)m(x)\kappa(dx)$ hits an independent exponential time with parameter 1. The corresponding generator is

$$(2.2.1) \quad L = \frac{1}{m(x)} \frac{d}{dx} \left(\frac{1}{w(x)} \frac{d}{dx} \right) - \kappa$$

Let $(X_t)_{0 \leq t < \zeta}$ be the diffusion of generator (2.2.1), which is killed either by hitting ∂I , or by exploding, or by the killing measure κ . For $x \in I$ let $\eta_{exc}^{>x}$ and $\eta_{exc}^{<x}$ be the excursion measures of X above and below the level x up to the last time X visits x . The behaviour of X from the first to the last time it visits x is a Poisson point process with intensity $\eta_{exc}^{>x} + \eta_{exc}^{<x}$, parametrized by the local time at x up to the value $\ell_t^x(X)$. $\eta_{exc}^{>x}$ and $\eta_{exc}^{<x}$ are obtained from the Levy-Itô measure on Brownian excursions through scale change, time change and multiplication by a density function accounting for the killing. See [25] for details on excursion measures in case of recurrent diffusions.

If X is transient the Green's function of L ,

$$G(x, y) := \mathbb{E}_x[\ell_t^y(X)]$$

is finite, continuous and symmetric. For $x \leq y$ it can be written

$$G(x, y) = u_{\uparrow}(x)u_{\downarrow}(y)$$

where $u_{\uparrow}(x)$ and $u_{\downarrow}(y)$ are positive, respectively non-decreasing and non-increasing solutions to the equation $Lu = 0$, which through a change of scale reduces to an equation of form (2.1.3). If S is bounded from below, $u_{\uparrow}(\inf I^+) = 0$. If S is bounded from above, $u_{\downarrow}(\sup I^-) = 0$. $u_{\uparrow}(x)$ and $u_{\downarrow}(y)$ are each determined up to a multiplication by a positive constant, but when entering the expression of G , the two constants are related. For $x \leq y \in I$:

$$\frac{u_{\uparrow}(x)}{u_{\uparrow}(y)} = \mathbb{P}_y(X \text{ hits } x \text{ before time } \zeta) \quad \frac{u_{\downarrow}(y)}{u_{\downarrow}(x)} = \mathbb{P}_x(X \text{ hits } y \text{ before time } \zeta)$$

See [12] or [5], chapter 16, for details. Let $W(u_{\downarrow}, u_{\uparrow})$ be the Wronskian of u_{\downarrow} and u_{\uparrow} :

$$W(u_{\downarrow}, u_{\uparrow})(x) := u_{\downarrow}(x) \frac{du_{\uparrow}}{dx}(x^+) - u_{\uparrow}(x) \frac{du_{\downarrow}}{dx}(x^+)$$

This Wronskian is actually the density of the scale measure: $W(u_\downarrow, u_\uparrow) \equiv w$. We may write G_L when there is an ambiguity on L .

If the killing measure κ is non zero, then the probability that X , starting from x , gets killed by κ before reaching a boundary of I or exploding equals $\int_I G(x, y)m(y)\kappa(dy)$. Conditionally on this event, the distribution of X_{ζ^-} is:

$$\frac{1_{z \in I} G(x, z)m(z)\kappa(dz)}{\int_I G(x, y)m(y)\kappa(dy)}$$

Indeed, let f be a non-negative compactly supported measurable function on I and

$$\tau_l := \inf \left\{ t \in [0, \zeta^{(0)} \mid \int_I \ell_t^y(X)m(y)\kappa(dy) > l \right\}$$

Then by definition

$$\mathbb{E}_x[f(X_{\zeta^-})] = \int_0^{+\infty} e^{-l} \mathbb{E}_x[f(X_{\tau_l \wedge \zeta^{(0)}})] dl = \int_0^{+\infty} dv e^{-v} \mathbb{E}_x \left[\int_0^v f(X_{\tau_l \wedge \zeta^{(0)}}) dl \right]$$

But

$$\int_0^v f(X_{\tau_l \wedge \zeta^{(0)}}) dl = \int_I \ell_{\tau_v \wedge \zeta^{(0)}}^y(X)m(y)\kappa(dy)$$

(see corollary 2.13, chapter X in [24]). It follows that

$$\mathbb{E}[f(X_{\zeta^-})] = \int_I f(y) \left[\int_0^{+\infty} e^{-v} \ell_{\tau_v \wedge \zeta^{(0)}}^y(X) dv \right] m(y)\kappa(dy) = \int_I f(y) G(x, y)m(y)\kappa(dy)$$

The semi-group of L has positive transition densities $p_t(x, y)$ with respect to the speed measure $m(y)dy$ and $(t, x, y) \mapsto p_t(x, y)$ is continuous on $(0, +\infty) \times I \times I$. McKean gives a proof of this in [21] in case when the killing measure k has a continuous density with respect to the Lebesgue measure. If this is not the case, we can take u a positive continuous solution to $Lu = 0$ and consider the h-transform of L by u : $u^{-1}Lu$. The latter is the generator of a diffusion without killing measure and by [21] this diffusion has continuous transition densities $\tilde{p}_t(x, y)$ with respect to $m(y)dy$. Then $u(x)\tilde{p}_t(x, y)\frac{1}{u(y)}$ are the transition densities of the semi-group of L . Transition densities with respect to the speed measure are symmetric: $p_t(x, y) = p_t(y, x)$. For all $x, y \in I$ and $t \geq 0$ the following equality holds:

$$(2.2.2) \quad \mathbb{E}_x[\ell_{t \wedge \zeta}^y(X)] = \int_0^t p_s(x, y) ds$$

Next we deal with bridge probability measures.

Proposition 2.5. — *The bridge probability measures $\mathbb{P}_{x,y}^t(\cdot)$ (bridge of X from x to y in time t conditioned neither to die nor to explode in the interval) satisfy: for all $x \in I$ the map $(x, y, t) \mapsto \mathbb{P}_{x,y}^t(\cdot)$ is continuous for the weak topology on probability measures on continuous paths.*

Proof. — Our proof mainly relies on absolute continuity arguments of [22] and [6], and the time reversal argument of [22]. [6] gives a proof of weak continuity of bridges for conservative Feller cadlag processes on second countable locally compact spaces. But since the proof contains an error and we do not restrict to conservative diffusions, we give here accurate arguments for the weak continuity.

First we can restrict to the case $\kappa = 0$. Otherwise consider u a solution to $Lu = 0$, positive on I . The generator of the h-transform of L by u is

$$\frac{1}{u(x)^2 m(x)} \frac{d}{dx} \left(\frac{u(x)^2}{w(x)} \frac{d}{dx} \right)$$

and does not contain any killing measure. The h-transform preserves the bridge measures and changes the density functions relatively to $m(y)dy$ to $\frac{1}{u(x)} p_t(x, y) u(y)$, and thus preserves their continuity.

Then we normalise the length of bridges: if $(X_s^{(x, y, t)})_{0 \leq s \leq t}$ is a path under the law $\mathbb{P}_{x, y}^t(\cdot)$, let $\tilde{\mathbb{P}}_{x, y}^t(\cdot)$ be the law of $(X_{rt}^{(x, y, t)})_{0 \leq r \leq 1}$. It is sufficient to prove that $(x, y, t) \mapsto \tilde{\mathbb{P}}_{x, y}^t(\cdot)$ is continuous. For $v \in [0, 1]$, let $\tilde{\mathbb{P}}_{x, y}^{t, v}(\cdot)$ be the law of $(X_{rt}^{(x, y, t)})_{0 \leq r \leq v}$. Let $\tilde{\mathbb{P}}_x^{t, v}(\cdot)$ be the law of the Markovian path $(X_{rt})_{0 \leq r \leq v}$ starting from x . For $v \in [0, 1]$ we have the following absolute continuity relationship:

$$(2.2.3) \quad d\tilde{\mathbb{P}}_{x, y}^{t, v} = 1_{vt < \zeta} \frac{p_{(1-v)t}(X_{vt}, y)}{p_t(x, y)} d\tilde{\mathbb{P}}_x^{t, v}$$

Let $(J_n)_{n \geq 0}$ be an increasing sequence of compact subintervals of I such that $I = \bigcup_{n \geq 0} J_n$. Let T_n be the first exit time from J_n . Let f_n be continuous compactly supported function on I such that $0 \leq f_n \leq 1$ and $f_n|_{J_n} \equiv 1$. We can further assume that the sequence $(f_n)_{n \geq 0}$ is non-decreasing. The map

$$(x, y, t) \mapsto f_n(\sup_{[0, vt]} X) f_n(\inf_{[0, vt]} X) d\tilde{\mathbb{P}}_x^{t, v}$$

is weakly continuous. Let $(x_j, y_j, t_j)_{j \geq 0}$ be a sequence converging to (x, y, t) . Let F be a continuous bounded functional on $\mathcal{C}([0, v])$. Then applying (2.2.3) we get:

$$(2.2.4) \quad \tilde{\mathbb{P}}_{x_j, y_j}^{t_j, v}(f_n(\sup_{[0, v]} \gamma) f_n(\inf_{[0, v]} \gamma) F(\gamma)) - \tilde{\mathbb{P}}_{x, y}^{t, v}(f_n(\sup_{[0, v]} \gamma) f_n(\inf_{[0, v]} \gamma) F(\gamma)) =$$

$$(2.2.5) \quad \tilde{\mathbb{P}}_{x_j}^{t_j, v} \left(\frac{p_{(1-v)t_j}(\gamma(v), y)}{p_{t_j}(x_j, y)} f_n(\sup_{[0, v]} \gamma) f_n(\inf_{[0, v]} \gamma) F(\gamma) \right)$$

$$(2.2.6) \quad - \tilde{\mathbb{P}}_x^{t, v} \left(\frac{p_{(1-v)t}(\gamma(v), y)}{p_t(x, y)} f_n(\sup_{[0, v]} \gamma) f_n(\inf_{[0, v]} \gamma) F(\gamma) \right)$$

$$(2.2.7) \quad + \tilde{\mathbb{P}}_{x_j}^{t_j, v} \left(\frac{p_{(1-v)t_j}(\gamma(v), y_j)}{p_{t_j}(x_j, y_j)} f_n(\sup_{[0, v]} \gamma) f_n(\inf_{[0, v]} \gamma) F(\gamma) \right)$$

$$(2.2.8) \quad -\tilde{\mathbb{P}}_{x_j}^{t_j, v} \left(\frac{p_{(1-v)t}(\gamma(v), y)}{p_t(x, y)} f_n(\sup_{[0, v]} \gamma) f_n(\inf_{[0, v]} \gamma) F(\gamma) \right)$$

Since $\frac{p_{(1-v)t}(\cdot, y)}{p_t(x, y)}$ is continuous and bounded on J_n , (2.2.5)–(2.2.6) converges to 0. Moreover for j large enough, $\frac{p_{(1-v)t_j}(\cdot, y_j)}{p_{t_j}(x_j, y_j)}$ is uniformly close on J_n to $\frac{p_{(1-v)t}(\cdot, y)}{p_t(x, y)}$. Thus (2.2.7)–(2.2.8) converges to 0 and finally (2.2.4) converges to 0. Let $n_0 \in \mathbb{N}$ and $n \geq n_0$. Then

$$\begin{aligned} \tilde{\mathbb{P}}_{x_j, y_j}^{t_j, v} (1 - f_n(\sup_{[0, v]} \gamma) f_n(\inf_{[0, v]} \gamma)) &= 1 - \tilde{\mathbb{P}}_{x_j, y_j}^{t_j, v} (f_n(\sup_{[0, v]} \gamma) f_n(\inf_{[0, v]} \gamma)) \\ &\leq 1 - \tilde{\mathbb{P}}_{x_j, y_j}^{t_j, v} (f_{n_0}(\sup_{[0, v]} \gamma) f_{n_0}(\inf_{[0, v]} \gamma)) \rightarrow 1 - \tilde{\mathbb{P}}_{x, y}^{t, v} (f_{n_0}(\sup_{[0, v]} \gamma) f_{n_0}(\inf_{[0, v]} \gamma)) \end{aligned}$$

and consequently

$$\lim_{n \rightarrow +\infty} \limsup_{j \rightarrow +\infty} \tilde{\mathbb{P}}_{x_j, y_j}^{t_j, v} (1 - f_n(\sup_{[0, v]} \gamma) f_n(\inf_{[0, v]} \gamma)) = 0$$

It follows that

$$\lim_{j \rightarrow +\infty} \tilde{\mathbb{P}}_{x_j, y_j}^{t_j, v} (F(\gamma)) = \tilde{\mathbb{P}}_{x, y}^{t, v} (F(\gamma))$$

From this we get that the law of any finite-dimensional family of marginals of $\tilde{\mathbb{P}}_{x, y}^t(\cdot)$ depends continuously on (x, y, t) . To conclude we need a tightness result for $(x, y, t) \mapsto \tilde{\mathbb{P}}_{x, y}^t(\cdot)$. We have already tightness for $(x, y, t) \mapsto \tilde{\mathbb{P}}_{x, y}^{t, v}(\cdot)$. The image of $\tilde{\mathbb{P}}_{x, y}^t(\cdot)$ through time reversal is $\tilde{\mathbb{P}}_{y, x}^t(\cdot)$. So we also have tightness on intervals $[1 - v', 1]$ where $0 < v' < 1$. But if $v + v' > 1$, tightness on $[0, v]$ and on $[1 - v', 1]$ implies tightness on $[0, 1]$. This concludes. The article [6] contains an error in the proof of the tightness of bridge measures in the neighbourhood of the endpoint. \square

2.3. "Generators" with creation of mass

In this section we consider more general operators

$$(2.3.1) \quad L = \frac{1}{m(x)} \frac{d}{dx} \left(\frac{1}{w(x)} \frac{d}{dx} \right) + \nu$$

with zero Dirichlet boundary conditions on ∂I , where ν is a signed measure on I which is no longer assumed to be negative. We set

$$L^{(0)} := L - \nu$$

In the sequel we may call L "generator" even in case the semi-group $(e^{tL})_{t \geq 0}$ does not make sense. Our main goal in this subsection is to characterize through a positivity condition the subclass of operators of form (2.3.1) that are equivalent up to an h -transform to the generator of a diffusion of form (2.2.1).

We will consider several kinds of transformations on operators of the form (2.3.1). First, the h-transform: Let h be a positive continuous function on I such that $\frac{d^2 h}{dx^2}$ is a signed measure. We call $Conj(h, L)$ the operator

$$Conj(h, L) = \frac{1}{h(x)^2 m(x)} \frac{d}{dx} \left(\frac{h(x)^2}{w(x)} \frac{d}{dx} \right) + \nu + \frac{1}{h} L^{(0)} h$$

If f is smooth function compactly supported in I then

$$Conj(h, L)f = h^{-1} L(hf)$$

We will call $Conj(h, L)$ the h-transform of L by h even though h may not be harmonic ($Lh = 0$) or superharmonic ($Lh \leq 0$) and L is not necessarily the generator of a diffusion.

Second, the change of scale: If A is a C^1 function on I such that $\frac{dA}{dx} > 0$ and $\frac{d^2 A}{dx^2} \in \mathbb{L}_{loc}^\infty(I)$ and $(\gamma(t))_{0 \leq t \leq T}$ a continuous path in I , then we will set $Scale_A(\gamma)$ to be the continuous path $(A(\gamma(s)))_{0 \leq s \leq T}$ in $A(I)$. Let $Scale_A^{gen}(L)$ be the operator on functions on $A(I)$ with zero Dirichlet boundary conditions induced by this change of scale:

$$Scale_A^{gen}(L) = \frac{1}{m \circ A^{-1}(a)} \frac{d}{da} \left(\frac{1}{w \circ A^{-1}(a)} \frac{d}{da} \right) + A_* \nu$$

where $A_* \nu$ is the push-forward of the measure ν by A .

Third, the change of time: If V is positive continuous on I then we can consider the change of time $ds = V(\gamma(t))dt$. Let $Speed_V$ be the corresponding transformation on paths. The corresponding "generator" is $\frac{1}{V}L$.

Finally, the restriction: if \tilde{I} is an open subinterval of I then set $L|_{\tilde{I}}$ to be the operator L acting on functions supported in \tilde{I} and with zero Dirichlet conditions on $\partial\tilde{I}$.

For the analysis of L we will use a bit of spectral theory: If $[x_0, x_1]$ is a compact interval of \mathbb{R} and \tilde{m}, \tilde{w} are positive continuous functions on $[x_0, x_1]$, then the operator $\frac{1}{\tilde{m}(x)} \frac{d}{dx} \left(\frac{1}{\tilde{w}(x)} \frac{d}{dx} \right)$ with zero Dirichlet boundary conditions has a discrete spectrum of negative eigenvalues. Let $-\tilde{\lambda}_1$ be the first eigenvalue. It is simple. According to Sturm-Liouville theory (see for instance [29], section 5.5) we have the following picture:

Property 2.6. — Let $\lambda > 0$ and u a solution to

$$\frac{1}{\tilde{m}} \frac{d}{dx} \left(\frac{1}{\tilde{w}} \frac{d}{dx} \right) + \lambda u = 0$$

with initial conditions $u(x_0) = 0, \frac{du}{dx}(x_0) > 0$.

- (i) If u is positive on (x_0, x_1) and $u(x_1) = 0$ then $\lambda = \tilde{\lambda}_1$ and u is the fundamental eigenfunction.
- (ii) If u is positive on $(x_0, x_1]$ then $\lambda < \tilde{\lambda}_1$

- (iii) If u changes sign on (x_0, x_1) then $\lambda > \tilde{\lambda}_1$

Next we state and prove the main result of this section.

Proposition 2.7. — *The following two conditions are equivalent:*

- (i) There is a positive continuous function u on I satisfying $Lu = 0$.
- (ii) For any f smooth compactly supported in I

$$(2.3.2) \quad \int_I (L^{(0)}f)(x)f(x)m(x)dx + \int_I f(x)^2m(x)\nu(dx) \leq 0$$

Proof. — (i) implies (ii): First observe that the equation $Lu = 0$ reduces through a change of scale to an equation of the form (2.1.1). Let u be given by condition (i). Let $\tilde{L} := \text{Conj}(u, L)$. Since $Lu = 0$, \tilde{L} is a generator of a diffusion without killing measure. Let $\tilde{m}(x) := u^2(x)m(x)$. Then for all g smooth compactly supported in I , $\int_I (\tilde{L}g)(x)g(x)\tilde{m}(x)dx \leq 0$. But

$$\int_I (\tilde{L}g)(x)g(x)\tilde{m}(x)dx = \int_I (L^{(0)}(ug))(x)(ug)(x)m(x)dx + \int_I (ug)(x)^2m(x)\nu(dx)$$

Thus (2.3.2) holds for all f positive compactly supported in I such that $u^{-1}f$ is smooth. By density arguments, this holds for general smooth f .

(ii) implies (i): First we will show that for every compact subinterval J of I there is a positive continuous function u_J on $\overset{\circ}{J}$ satisfying $Lu_J = 0$ on $\overset{\circ}{J}$. Let J be such an interval. According to lemma 2.2 there is $\lambda > 0$ and u_λ positive continuous on J satisfying $Lu_\lambda - \lambda u_\lambda = 0$ on J . Let $L_\lambda := \text{Conj}(u_\lambda, L|_J)$. Then

$$L_\lambda = \frac{1}{u_\lambda^2 m} \frac{d}{dx} \left(\frac{u_\lambda^2}{w} \frac{d}{dx} \right) + \lambda$$

Let $L_\lambda^{(0)} := L_\lambda - \lambda$. $L^{(0)}$ is the generator of a diffusion on $\overset{\circ}{J}$. We can apply the standard spectral theorem to $L_\lambda^{(0)}$. Let $-\lambda_1$ be its fundamental eigenvalue. $L_\lambda^{(0)} + \lambda = L_\lambda$ is a non-positive operator because it is an h-transform of $L|_J$ which satisfies condition (ii). This implies that $\lambda \leq \lambda_1$. Let \tilde{u} be a solution of $L_\lambda^{(0)}\tilde{u} + \lambda\tilde{u} = 0$ with initial conditions $\tilde{u}(\min J) = 0$ and $\frac{d\tilde{u}}{dx}(\min J) > 0$. Since $\lambda \leq \lambda_1$, according to property 2.6, \tilde{u} is positive on $\overset{\circ}{J}$. We set $u_J := u_\lambda\tilde{u}$. Then u_J is positive continuous on $\overset{\circ}{J}$ and satisfies $Lu_J = 0$. This finishes the proof of the first step.

Now consider a fixed point x_0 in I and $(J_n)_{n \geq 0}$ an increasing sequence of compact subintervals of I such that $x_0 \in \overset{\circ}{J}_0$ and $\bigcup_{n \geq 0} J_n = I$. Let u_{J_n} be a positive L -harmonic function on $\overset{\circ}{J}_n$. We may assume that $u_{J_n}(x_0) = 1$. The sequence $\left(\frac{du_{J_n}}{dx}(x_0^+) \right)_{n \geq 0}$ is bounded from below. Otherwise some of the u_{J_n} would change sign on $I \cap (x_0, +\infty)$. Similarly, since none of the u_{J_n} changes sign on $I \cap (-\infty, x_0)$,

$\left(\frac{du_{J_n}}{dx}(x_0^+)\right)_{n \geq 0}$ is bounded from above. Let v be an accumulation value of the sequence $\left(\frac{du_{J_n}}{dx}(x_0^+)\right)_{n \geq 0}$. Then the L -harmonic function satisfying the initial conditions $u(x_0) = 1$ and $\frac{du}{dx}(x_0^+) = v$ is positive on I .

We will divide the operators of the form (2.3.2) in two sets: $\mathfrak{D}^{0,-}$ for those that satisfies the constraints of the proposition 2.7 and \mathfrak{D}^+ for those that don't. $\mathfrak{D}^{0,-}$ is made exactly of operators that are equivalent up to an h -transform to the generator of a diffusion. We will subdivide the set $\mathfrak{D}^{0,-}$ in two: \mathfrak{D}^- for the operators that are an h -transform of the generator of a transient diffusion and \mathfrak{D}^0 for those that are an h -transform of the generator of a recurrent diffusion. These two subclasses are well defined since a transient diffusion can not be an h -transform of a recurrent one. Observe that each of $L \in \mathfrak{D}^-, \mathfrak{D}^0$ and \mathfrak{D}^+ is stable under h -transforms, changes of scale and of speed. Operators in \mathfrak{D}^- and \mathfrak{D}^0 do not need to be generators of transient or recurrent diffusions themselves. For instance consider on \mathbb{R}

$$L = \frac{1}{2} \frac{d^2}{dx^2} + a_+ \delta_1 - a_- \delta_{-1}$$

where $a_+, a_- > 0$. If $3a_+ - a_- > 0$ then $L \in \mathfrak{D}^+$, if $3a_+ - a_- = 0$ then $L \in \mathfrak{D}^0$, if $3a_+ - a_- < 0$ then $L \in \mathfrak{D}^-$. \square

If $L \in \mathfrak{D}^{0,-}$, the semi-group $(e^{tL})_{t \geq 0}$ is well defined. Indeed, let X be the diffusion on I of generator $L^{(0)}$ and ζ the first time it hits the boundary of I or blows up to infinity. Let u be a positive L -harmonic function and $\tilde{L} := \text{Conj}(u, L)$. \tilde{L} is the generator of a diffusion \tilde{X} on I without killing measure. Let $\tilde{\zeta}$ be the first time \tilde{X} hits the boundary of I or blows up to infinity. Using Girsanov's theorem, one can show that for any F positive measurable functional on paths, $x \in I$ and $t > 0$ the following equality holds:

$$\mathbb{E}_x \left[1_{t < \zeta} \exp \left(\int_I \ell_t^y(X) m(y) \nu(dy) \right) F((X_s)_{0 \leq s \leq t}) \right] = \frac{1}{u(x)} \mathbb{E}_x \left[1_{t < \tilde{\zeta}} u(\tilde{X}_t) F((\tilde{X}_s)_{0 \leq s \leq t}) \right]$$

In case $L \in \mathfrak{D}^-$, let $(G_{\tilde{L}}(x, y))_{x, y \in I}$ be the Green's function of \tilde{L} relatively to the measure $u(x)^2 m(x) dx$. Then L has a Green's function $(G_L(x, y))_{x, y \in I}$ that equals

$$G_L(x, y) = \mathbb{E}_x \left[\int_0^\zeta \exp \left(\int_I \ell_t^z(X) m(z) \nu(dz) \right) d_t \ell_t^y(X) \right] = u(x) u(y) G_{\tilde{L}}(x, y)$$

For $L \in \mathfrak{D}^-$, the Green's functions G_L satisfy the following resolvent identities

Lemma 2.8. — *If $L \in \mathfrak{D}^-$ and $\tilde{\nu}$ is a signed measure with compact support on I such that $L + \tilde{\nu} \in \mathfrak{D}^-$, then for all $x, y \in I$*

$$\begin{aligned} G_{L+\tilde{\nu}}(x, y) - G_L(x, y) &= \int_I G_{L+\tilde{\nu}}(x, z) G_L(z, y) m(z) \tilde{\nu}(dz) \\ &= \int_I G_L(x, z) G_{L+\tilde{\nu}}(z, y) m(z) \tilde{\nu}(dz) \end{aligned}$$

Proof. — We decompose L as $L = L^{(0)} + \nu$ where $L^{(0)}$ does not contain measures and ν is a signed measure on I . Let $(X_t)_{0 \leq t < \zeta}$ be the diffusion of generator $L^{(0)}$. Then

$$G_L(x, y) = \mathbb{E}_x \left[\int_0^\xi \exp \left(\int_I \ell_t^a(X) m(a) \nu(da) \right) d_t \ell_t^y(X) \right]$$

$$G_{L+\tilde{\nu}}(x, y) = \mathbb{E}_x \left[\int_0^\xi \exp \left(\int_I \ell_t^a(X) m(a) (\nu + \tilde{\nu})(da) \right) d_t \ell_t^y(X) \right]$$

and

$$\begin{aligned} &\exp \left(\int_I \ell_t^a(X) m(a) (\nu + \tilde{\nu})(da) \right) - \exp \left(\int_I \ell_t^a(X) m(a) \nu(da) \right) \\ &= \exp \left(\int_I \ell_t^a(X) m(a) \nu(da) \right) \times \left(\exp \left(\int_I \ell_t^a(X) m(a) \tilde{\nu}(da) \right) - 1 \right) \\ &= \exp \left(\int_I \ell_t^a(X) m(a) \nu(da) \right) \int_I \int_0^t \exp \left(\int_I \ell_s^a(X) m(a) \tilde{\nu}(da) \right) d_s \ell_s^z(X) m(z) \tilde{\nu}(dz) \end{aligned}$$

Thus $G_{L+\tilde{\nu}}(x, y) - G_L(x, y)$ equals

(2.3.3)

$$\mathbb{E}_x \left[\int_I \int_0^\xi \int_0^t \exp \left(\int_I m(a) (\ell_t^a(X) \nu(da) + \ell_s^a(X) \tilde{\nu}(da)) \right) d_s \ell_s^z(X) d_s \ell_t^y(X) m(z) \tilde{\nu}(dz) \right]$$

We would like to interchange $\mathbb{E}_x[\cdot]$ and $\int_I (\cdot) m(z) \tilde{\nu}(dz)$. Let $z \in I$ and $(X_t^{(x)})_{0 \leq t < \zeta_x}$, $(X_t^{(z)})_{0 \leq t < \zeta_z}$ be two independent diffusions of generator $L^{(0)}$ starting in x respectively z . Applying Markov property, we get

$$\begin{aligned} &\mathbb{E}_x \left[\int_0^\xi \int_0^t \exp \left(\int_I m(a) (\ell_t^a(X) \nu(da) + \ell_s^a(X) \tilde{\nu}(da)) \right) d_s \ell_s^z(X) d_s \ell_t^y(X) \right] \\ &= \mathbb{E} \left[\int_0^{\zeta_x} \int_0^{\zeta_z} \exp \left(\int_I m(a) (\ell_s^a(X^{(x)}) (\nu + \tilde{\nu})(da)) \right) \right. \\ &\quad \left. \times \exp \left(\int_I m(a) (\ell_u^a(X^{(z)}) \nu(da)) d_u \ell_u^z(X^{(z)}) d_s \ell_s^x(X^{(x)}) \right) \right] \\ &= G_{L+\tilde{\nu}}(x, z) G_L(z, y) \end{aligned}$$

Since $\tilde{\nu}$ has compact support

$$\begin{aligned} \mathbb{E}_x \left[\int_I \int_0^\xi \int_0^t \exp \left(\int_I m(a)(\ell_t^a(X)\nu(da) + \ell_s^a(X)\tilde{\nu}(da)) \right) d_s \ell_s^z(X) d_s \ell_t^y(X) m(z) |\tilde{\nu}|(dz) \right] \\ = \int_I \mathbb{E}_x \left[\int_0^\xi \int_0^t \exp \left(\int_I m(a)(\ell_t^a(X)\nu(da) + \ell_s^a(X)\tilde{\nu}(da)) \right) d_s \ell_s^z(X) d_s \ell_t^y(X) \right] m(z) |\tilde{\nu}|(dz) \\ = \int_I G_{L+\tilde{\nu}}(x, z) G_L(z, y) m(z) |\tilde{\nu}|(dz) < +\infty \end{aligned}$$

Thus in (2.3.3) we can interchange $\mathbb{E}_x[\cdot]$ and $\int_I(\cdot)m(z)\tilde{\nu}(dz)$ and get

$$G_{L+\tilde{\nu}}(x, y) - G_L(x, y) = \int_I G_{L+\tilde{\nu}}(x, z) G_L(z, y) m(z) \tilde{\nu}(dz)$$

Since L and $L + \tilde{\nu}$ play symmetric roles, we also have

$$G_L(x, y) - G_{L+\tilde{\nu}}(x, y) = \int_I G_L(x, z) G_{L+\tilde{\nu}}(z, y) m(z) (-\tilde{\nu})(dz)$$

□

The discrete analogue of the sets \mathfrak{D}^- , \mathfrak{D}^0 and \mathfrak{D}^+ are symmetric matrices with non-negative off-diagonal coefficients inducing a connected transition graph, with the highest eigenvalue that is respectively negative, zero and positive. However in continuous case the sets $L \in \mathfrak{D}^-$, \mathfrak{D}^0 and \mathfrak{D}^+ can not be defined spectrally because for operators from $L \in \mathfrak{D}^-$ and \mathfrak{D}^+ the maximum of the spectrum can also equal zero. However the next result shows that the sets \mathfrak{D}^- and \mathfrak{D}^+ are stable under small perturbations of the measure ν and that \mathfrak{D}^0 is not.

Proposition 2.9. — (i) If $L \in \mathfrak{D}^0$ and κ is a non-zero positive Radon measure on I then $L - \kappa \in \mathfrak{D}^-$ and $L + \kappa \in \mathfrak{D}^+$.

– (ii) If $L \in \mathfrak{D}^-$ and J is a compact subinterval of I then there is $K > 0$ such that for any positive measure κ supported in J satisfying $\kappa(J) < K$ we have $L + \kappa \in \mathfrak{D}^-$.

– (iii) If $L \in \mathfrak{D}^+$ then there is $K > 0$ such that for any positive finite measure κ satisfying $\kappa(I) < K$ we have $L - \kappa \in \mathfrak{D}^+$.

– (iv) If $L \in \mathfrak{D}^+$, there is a positive Radon measure κ on I such that $L - \kappa \in \mathfrak{D}^0$.

– (v) Let $L \in \mathfrak{D}^+$ and $x_0 < x_1 \in I$. Then $L|_{(x_0, x_1)} \in \mathfrak{D}^0$ if and only if there is an L -harmonic function u positive on (x_0, x_1) and zero in x_0 and x_1 .

Proof. — (i): Consider h positive continuous on I such that $Conj(h, L)$ is the generator of a recurrent diffusion. Since $Conj(h, L - \kappa) = Conj(h, L) - \kappa$, $Conj(h, L - \kappa)$ is the generator of a diffusion killed at rate κ and thus $L - \kappa \in \mathfrak{D}^-$. Similarly we can not have $L + \kappa \in \mathfrak{D}^{0,-}$ because this would mean $L = (L + \kappa) - \kappa \in \mathfrak{D}^-$.

(ii): Without loss of generality we may assume that L is the generator of a transient diffusion and that it is at natural scale, that is $L = \frac{1}{m(x)} \frac{d^2}{dx^2}$. Since the diffusion is transient, $I \neq \mathbb{R}$. We may assume that $x_0 := \inf I > -\infty$. Write $J = [x_1, x_2]$. Let κ be a positive measure supported in $[x_1, x_2]$. Let u be the solution to $Lu + u\kappa = 0$ with the initial conditions $u(x_0) = 0$, $\frac{du}{dx}(x_0^+) = 1$. u is affine on $[x_0, x_1]$ and on $[x_2, \sup I)$. On $[x_1, x_2]$ u is bounded from above by $x_2 - x_0$. Thus, if

$$\kappa([x_1, x_2]) \leq \frac{\min_{[x_1, x_2]} m}{(x_2 - x_0)}$$

then u is non-decreasing on I and hence positive. This implies that $L + \kappa \in \mathfrak{D}^{0,-}$. By the point (i) of current proposition, if $\kappa([x_1, x_2]) < \frac{\min_{[x_1, x_2]} m}{(x_2 - x_0)}$ then $L + \kappa \in \mathfrak{D}^-$.

(iii): By definition there is f smooth compactly supported in I such that (2.3.2) does not hold for f . Let U be the value of the left-hand side in (2.3.2). $U > 0$. If κ is a positive finite measure on I satisfying

$$\kappa(I) < \frac{U}{\|f\|_\infty^2 \max_{\text{Supp} f} m}$$

then if we replace ν by $\nu - \kappa$ in (2.3.2), keeping the same function f , we still get something positive. Thus $L - \kappa \in \mathfrak{D}^+$.

(iv): Let f be a smooth function compactly supported in I such that (2.3.2) does not hold for f . Let J be a compact subinterval of I containing the support of f . The set

$$\{s \in [0, 1] \mid L - \nu_+ + s 1_J \nu_+ \in \mathfrak{D}^-\}$$

is not empty because it contains 0, and open by proposition 2.9 (ii). Let s_{max} be its supremum. Then $s_{max} < 1$ and $L - \nu_+ + s_{max} 1_J \nu_+ \in \mathfrak{D}^0$. Then

$$\kappa := 1_{I \setminus J} \nu_+ + (1 - s_{max}) 1_J \nu_+$$

is appropriate.

(v): First assume that there is such a function u . Then by definition $L_{|(x_0, x_1)} \in \mathfrak{D}^{0,-}$. $\text{Conj}(u, L_{|(x_0, x_1)})$ does not have any killing measure and the derivative of its natural scale function is $\frac{u}{u^2}$. It is not integrable in the neighbourhood of x_0 or x_1 . Thus the corresponding diffusion never hits x_0 or x_1 . This means that it is recurrent. Conversely, assume that $L_{|(x_0, x_2)} \in \mathfrak{D}^0$. Let u be a solution to $Lu = 0$ satisfying $u(x_0) = 0$ and $\frac{du}{dx}(x_0^+) > 0$. If u changed its sign on (x_0, x_1) then according to the preceding we would have $L_{|(x_0, x_1)} \in \mathfrak{D}^+$. If u were positive on an interval larger than (x_0, x_1) we would have $L_{|(x_0, x_1)} \in \mathfrak{D}^-$. The only possibility is that u is positive on (x_0, x_1) and zero in x_1 . \square

CHAPTER 3

MEASURE ON LOOPS AND ITS BASIC PROPERTIES

3.1. Spaces of loops

In this chapter, in the section 3.3, we will introduce the infinite measure μ^* on loops which is at the center of this work. Prior to this, in the section 3.2 we will introduce measures $\mu^{x,y}$ on finite life-time paths which will be instrumental for defining μ^* . In the sections 3.4, 3.5, 3.7, 3.8 will be explored different aspects of μ^* . In the section 3.6 we will extend the Vervaat's Brownian bridge to Brownian excursion transformation to general diffusions. This generalisation can be easily interpreted in terms of measure μ^* and is related to the results of section 3.5. In the section 3.1 we will introduce the spaces of paths and loops on which will be defined the measures we will consider throughout the paper.

First we will consider continuous, time parametrized, paths on \mathbb{R} , $(\gamma(t))_{0 \leq t \leq T(\gamma)}$, with finite life-time $T(\gamma) \in (0, +\infty)$. Given two such paths $(\gamma(t))_{0 \leq t \leq T(\gamma)}$ and $(\gamma'(t))_{0 \leq t \leq T(\gamma')}$, a natural distance between them is

$$d_{paths}(\gamma, \gamma') := |\log(T(\gamma)) - \log(T(\gamma'))| + \max_{v \in [0,1]} |\gamma(vT(\gamma)) - \gamma'(vT(\gamma'))|$$

A rooted loop in \mathbb{R} will be a continuous finite life-time path $(\gamma(t))_{0 \leq t \leq T(\gamma)}$ such that $\gamma(T(\gamma)) = \gamma(0)$ and \mathfrak{L} will stand for the space of such loops. \mathfrak{L} endowed with the metric d_{paths} is a Polish space. In the sequel we will use the corresponding Borel σ -algebra, $\mathcal{B}_{\mathfrak{L}}$, for the definition of measures on \mathfrak{L} . For $v \in [0, 1]$ we define a parametrisation shift transformation $shift_v$ on \mathfrak{L} : $shift_v(\gamma) = \tilde{\gamma}$ where $T(\tilde{\gamma}) = T(\gamma)$ and

$$\tilde{\gamma}(t) = \begin{cases} \gamma(vT(\gamma) + t) & \text{if } t \leq (1-v)T(\gamma) \\ \gamma(t - (1-v)T(\gamma)) & \text{if } t \geq (1-v)T(\gamma) \end{cases}$$

We introduce an equivalence relation on \mathfrak{L} : $\gamma \sim \gamma'$ if $T(\gamma') = T(\gamma)$ and there is $v \in [0, 1]$ such that $\gamma' = shift_v(\gamma)$. We call the quotient space \mathfrak{L}/\sim the space of unrooted loops, or just loops, and denote it \mathfrak{L}^* . Let π be the projection $\pi : \mathfrak{L} \rightarrow \mathfrak{L}^*$.

There is a natural metric $\delta_{\mathfrak{L}^*}$ on \mathfrak{L}^* :

$$d_{\mathfrak{L}^*}(\pi(\gamma), \pi(\gamma')) := \min_{v \in [0,1]} d_{paths}(shift_v(\gamma), \gamma')$$

$(\mathfrak{L}^*, d_{\mathfrak{L}^*})$ is a Polish space and π is continuous. For defining measures on \mathfrak{L}^* we will use its Borel σ -algebra, $\mathcal{B}_{\mathfrak{L}^*}$. $\pi^{-1}(\mathcal{B}_{\mathfrak{L}^*})$, the inverse image of $\mathcal{B}_{\mathfrak{L}^*}$ by π , is a sub-algebra of $\mathcal{B}_{\mathfrak{L}}$.

In the sequel we will consider paths and loops that have a continuous family of local times $(\ell_t^x(\gamma))_{x \in \mathbb{R}, 0 \leq t \leq T(\gamma)}$ relatively to a measure $m(x)dx$ such that for any positive measurable function f on \mathbb{R} and any $t \in [0, T(\gamma)]$

$$\int_0^t f(\gamma(s))ds = \int_I \ell_t^x(\gamma)m(x)dx$$

We will simply write $\ell^x(\gamma)$ for $\ell_{T(\gamma)}^x(\gamma)$.

In the sequel we will also consider transformations on paths and loops and the images of different measures by these transformation. We will use everywhere the following notation: If \mathcal{E} and \mathcal{E}' are two measurable spaces, $\varphi : \mathcal{E} \mapsto \mathcal{E}'$ a measurable map and η a positive measure on \mathcal{E} , $\varphi_*\eta$ will be the measure on \mathcal{E}' obtained as the image of η through φ .

3.2. Measures $\mu^{x,y}$ on finite life-time paths

First we recall the framework that Le Jan used in [13]: $\mathbb{G} = (V, E)$ is a finite connected undirected graph. $L_{\mathbb{G}}$ is the generator of a symmetric Markov jump process with killing on \mathbb{G} . $m_{\mathbb{G}}$ is the duality measure for $L_{\mathbb{G}}$. $(p_t^{\mathbb{G}}(x, y))_{x, y \in V, t \geq 0}$ is the family of transition densities of the jump process and $(\mathbb{P}_{x,y}^{\mathbb{G}, t}(\cdot))_{x, y \in V, t \geq 0}$ the family of bridge probability measures. The measure on rooted loops associated with $L_{\mathbb{G}}$ is

$$(3.2.1) \quad \mu_{L_{\mathbb{G}}}(\cdot) = \int_{t>0} \sum_{x \in V} \mathbb{P}_{x,x}^{\mathbb{G}, t}(\cdot) p_t^{\mathbb{G}}(x, x) m_{\mathbb{G}}(x) \frac{dt}{t}$$

$\mu_{L_{\mathbb{G}}}^*$ is the image of $\mu_{L_{\mathbb{G}}}$ by the projection on unrooted loops. The definition of $\mu_{L_{\mathbb{G}}}^*$ is the exact formal analogue of the definition used in [19] for the loops of the two-dimensional Brownian motion. In [13] also appear variable life-time bridge measures $(\mu_{L_{\mathbb{G}}}^{x,y})_{x, y \in V}$ which are related to $\mu_{L_{\mathbb{G}}}^*$:

$$(3.2.2) \quad \mu_{L_{\mathbb{G}}}^{x,y}(\cdot) = \int_0^{+\infty} \mathbb{P}_{x,y}^{\mathbb{G}, t}(\cdot) p_t^{\mathbb{G}}(x, y) dt$$

In this subsection we will define and give the important properties of the formal analogue of the measures $\mu_{L_{\mathbb{G}}}^{x,y}$ in case of one-dimensional diffusions. In the next section 3.2 we will do the same with the measure on loops $\mu_{L_{\mathbb{G}}}^*$.

I is an open interval of \mathbb{R} . $(X_t)_{0 \leq t < \zeta}$ is a diffusion on I with a generator L of the form (2.2.1). We use the notations of the section 2.1. Let $x, y \in I$. Following the pattern of (3.2.2) we define:

Definition 3.1. —

$$\mu_L^{x,y}(\cdot) := \int_0^{+\infty} \mathbb{P}_{x,y}^t(\cdot) p_t(x,y) dt$$

We will write $\mu^{x,y}$ instead of $\mu_L^{x,y}$ whenever there is no ambiguity on L . The definition of $\mu^{x,y}$ depends on the choice of m , but $m(y)\mu^{x,y}$ does not. Measures $\mu^{x,y}$ were first introduced by Dynkin in [7] and enter the expression of Dynkin's isomorphism between the Gaussian Free Field and the local times of random paths. Pitman and Yor studied these measures in [23] in the setting of one-dimensional diffusions without killing measure ($\kappa = 0$). Next we give a handy representation of $\mu^{x,y}$ in the setting of one-dimensional diffusions. It was observed and proved by Pitman and Yor in case $\kappa = 0$. We consider the general case.

Proposition 3.2. — *Let F be a non-negative measurable functional on the space of variable life-time paths starting from x . Then*

$$(3.2.3) \quad \mu^{x,y}(F(\gamma)) = \mathbb{E}_x \left[\int_0^\zeta F((X_s)_{0 \leq s \leq t}) d_t \ell_t^y(X) \right]$$

Equivalently

$$\mu^{x,y}(F(\gamma)) = \mathbb{E}_x \left[\int_0^{\ell_\zeta^y(X)} F((X_s)_{0 \leq s \leq \tau_l^y}) dl \right]$$

where $\tau_l^y := \inf\{t \geq 0 \mid \ell_t^z(X) > l\}$.

Proof. — It is enough to prove this for F non-negative continuous bounded functional which takes value 0 if either the life-time of the paths exceeds some value $t_{max} < +\infty$ or if it is inferior to some value t_{min} or if the endpoint of the path lies out of a compact subinterval $[z_1, z_2]$ of I . For $j \leq n \in \mathbb{N}$, set $t_{j,n} := t_{min} + \frac{j(t_{max} - t_{min})}{n}$ and $\Delta t_n := \frac{t_{max} - t_{min}}{n}$. Almost surely $\int_0^\zeta F((X_s)_{0 \leq s \leq t}) d_t \ell_t^y$ is a limit as $n \rightarrow +\infty$ of

$$(3.2.4) \quad \sum_{j=0}^{n-1} F((X_s)_{0 \leq s \leq t_{j,n}}) (\ell_{t_{j+1,n} \wedge \zeta}^y(X) - \ell_{t_{j,n} \wedge \zeta}^y(X))$$

Moreover (3.2.4) is dominated by $\|F\|_\infty \ell_{t_{max} \wedge \zeta}^y$. It follows that the expectations converge too. Using the Markov property and (2.2.2), we get that the expectation of (3.2.4) equals

$$(3.2.5) \quad \sum_{j=0}^{n-1} \int_{z \in I} \int_0^{\Delta t_n} \mathbb{P}_{x,z}^{t_{j,n}}(F((X_s)_{0 \leq s \leq t_{j,n}})) p_{t_{j,n}}(x,z) p_r(z,y) dr m(z) dz$$

Using the fact that $p_r(\cdot, \cdot)$ is symmetric, we can rewrite (3.2.5) as

$$(3.2.6) \quad \int_{z_1}^{z_2} \left(\sum_{j=0}^{n-1} \Delta t_n \mathbb{P}_{x,z}^{t_{j,n}}(F((X_s)_{0 \leq s \leq t_{j,n}})) p_{t_{j,n}}(x,z) \right) \frac{1}{\Delta t_n} \int_0^{\Delta t_n} p_r(y,z) dr m(z) dz$$

As $n \rightarrow +\infty$ the measure $\frac{1}{\Delta t_n} \int_0^{\Delta t_n} p_r(y, z) dm(z) dz$ converges weakly to δ_y . Using the weak continuity of bridge probabilities (proposition 2.5) we get that (3.2.6) converges to

$$\int_{t_{\min}}^{t_{\max}} \mathbb{P}_{x,y}^t (F((X_s)_{0 \leq s \leq t})) p_t(x, y) dt$$

□

Proposition 3.2 also holds in case of a Markov jump processes on a graph, where the local time is replaced by the occupation time in a vertex divided by its weight. Proposition 3.2 shows that we can consider $\mu^{x,y}$ as a measure on paths $(\gamma(t))_{0 \leq t \leq T(\gamma)}$ endowed with continuous occupation densities $(\ell_t^z(\gamma))_{z \in I, 0 \leq t \leq T(\gamma)}$. Next we state several properties that either follow almost immediately from the definition 3.1 and proposition 3.2 or are already known.

Property 3.3. — (i) *The total mass of the measure $\mu^{x,y}$ is finite if and only if X is transient and then it equals $G(x, y)$. If it is the case, $\frac{1}{G(x,x)} \mu^{x,x}$ is the law of X , starting from $X(0) = x$, up to the last time it visits x . $\frac{1}{G(x,y)} \mu^{x,y}$ is the law of X , starting from $X(0) = x$, conditioned to visit y before ζ , up to the last time it visits y .*

- (ii) *The measure $\mu^{y,x}$ is image of the measure $\mu^{x,y}$ by time reversal.*
- (iii) *If \tilde{I} is an open subinterval of I then*

$$\mu_{L|_{\tilde{I}}}^{x,y}(d\gamma) = \mathbf{1}_{\gamma \text{ contained in } \tilde{I}} \mu_L^{x,y}(d\gamma)$$

- (iv) *If $\tilde{\kappa}$ is a positive Radon measure on I then*

$$\mu_{L-\tilde{\kappa}}^{x,y}(d\gamma) = \exp\left(-\int_I \ell^z(\gamma) m(z) \tilde{\kappa}(dz)\right) \mu_L^{x,y}(d\gamma)$$

- (v) *If A is a change of scale function then*

$$\mu_{\text{Scale}_A^{gen} L}^{A(x), A(y)} = \text{Scale}_{A*} \mu_L^{x,y}$$

- (vi) *If V is a positive continuous function on I then for the time changed diffusion of generator $\frac{1}{V}L$:*

$$\mu_{\frac{1}{V}L}^{x,y} = \text{Speed}_{V*} \mu_L^{x,y}$$

- (vii) *If h is a positive continuous function on I such that $\frac{d^2 h}{dx^2}$ is a signed measure and Lu is a negative measure then*

$$\mu_{\text{Conj}(h,L)}^{x,y} = \frac{1}{h(x)h(y)} \mu_L^{x,y}$$

- (viii) *Let X and \tilde{X} be two independent Markovian paths of generator L starting from $X(0) = x$ and $\tilde{X}(0) = y$. For $a \leq x \wedge y$, we introduce T_a and \tilde{T}_a the first time X respectively \tilde{X} hits a . Let $\mathbb{P}_x^{T_a}$ be the first passage bridge of X from x to a , conditioned by the event $T_a < \zeta$. Let $\tilde{\mathbb{P}}_y^{\tilde{T}_a}$ be the analogue for \tilde{X} . Let $\tilde{\mathbb{P}}_y^{\tilde{T}_a \wedge}$ be*

the image of $\tilde{\mathbb{P}}_y^{T_a}$ through time reversal and $\mathbb{P}_x^{T_a} \triangleleft \tilde{\mathbb{P}}_y^{T_a \wedge}$ the image of $\mathbb{P}_x^{T_a} \otimes \tilde{\mathbb{P}}_y^{T_a \wedge}$ through concatenation at a of two paths, one ending and the other starting in a . Then

$$\mu^{x,y}(\cdot) = \int_{a \in I, a \leq x \wedge y} \mathbb{P}_x(T_a < \zeta) \mathbb{P}_y(\tilde{T}_a < \tilde{\zeta}) \left(\mathbb{P}_x^{T_a} \triangleleft \tilde{\mathbb{P}}_y^{T_a \wedge} \right) (\cdot) w(a) da$$

Previous equalities depend on a particular choice of the speed measure for the modified generator. For (iv) we keep the measure $m(y)dy$. For (iii) we restrict $m(y)dy$ to \tilde{I} . For (v) we choose $(\frac{dA}{dx} \circ A^{-1})^{-1} m \circ A^{-1} da$. For (vi) we choose $\frac{1}{V(y)} m(y)dy$. For (vii) we choose $h(y)^2 m(y) dy$. Property (ii) follows from that $p_t(x, y) = p_t(y, x)$ and $\mathbb{P}_{y,x}^t(\cdot)$ is the image of $\mathbb{P}_{x,y}^t(\cdot)$ by time reversal. Property (vi) is not immediate from definition 1 because fixed times are transformed by time change in random times, but follows from proposition 3.2. Property (vii) follows from that an h-transform does not change bridge probability measures and changes the semi-group $(p_t(x, y)m(y)dy)_{t \geq 0, x \in I}$ to $(\frac{1}{u(x)} p_t(x, y)u(y)m(y)dy)_{t \geq 0, x \in I}$. Properties (ii) and (viii) were proved by Pitman and Yor in case $\kappa = 0$. See [23]. The case $\kappa \neq 0$ can be obtained through h-transforms.

Next property was given without proof by Dynkin in [7].

Lemma 3.4. — Assume $\kappa \neq 0$. Let $\mathbb{P}_x(\cdot)$ be the law of $(X_t)_{0 \leq t < \zeta}$ where $X(0) = x$. Then

$$\int_{y \in I} \mu^{x,y}(\cdot) m(y) \kappa(dy) = 1_X \text{ killed by } \kappa \mathbb{P}_x(\cdot)$$

Proof. — Let $0 < t_1 < t_2 < \dots < t_n$ and let $A_1, A_2, \dots, A_n, A_{n+1}$ be Borel subsets of I . The measure $\mu^{x,y}$ satisfies the following Markov property

$$\begin{aligned} \mu^{x,y}(T(\gamma) > t_n, \gamma(t_1) \in A_1, \dots, \gamma(t_n) \in A_n, \gamma(T(\gamma)) \in A_{n+1}) = \\ \int_{A_1 \times \dots \times A_n} p_{t_1}(x, x_1) m(x_1) \dots p_{t_n - t_{n-1}}(x_{n-1}, x_n) m(x_n) \mu^{x_n, y}(T(\gamma) \in A_{n+1}) dx_1 \dots dx_n \\ = 1_{y \in A_{n+1}} \int_{A_1 \times \dots \times A_n} p_{t_1}(x, x_1) m(x_1) \dots p_{t_n - t_{n-1}}(x_{n-1}, x_n) m(x_n) G(x_n, y) dx_1 \dots dx_n \end{aligned}$$

Hence

$$(3.2.7) \quad \int_{y \in I} \mu^{x,y}(T(\gamma) > t_n, \gamma(t_1) \in A_1, \dots, \gamma(t_n) \in A_n, \gamma(T(\gamma)) \in A_{n+1}) m(y) \kappa(dy) = \\ \int_{A_1 \times \dots \times A_{n+1}} p_{t_1}(x, x_1) m(x_1) \dots p_{t_n - t_{n-1}}(x_{n-1}, x_n) m(x_n) G(x_n, y) m(y) dx_1 \dots dx_n \kappa(dy)$$

From Markov property of X follows

$$\begin{aligned} \mathbb{P}_x(\zeta > t_n, X_{t_1} \in A_1, \dots, X_{t_n} \in A_n, X_{\zeta^-} \in A_{n+1}) = \\ \int_{A_1 \times \dots \times A_n} p_{t_1}(x, x_1) m(x_1) \dots p_{t_n - t_{n-1}}(x_{n-1}, x_n) m(x_n) P_{x_n}(X_{\zeta^-} \in A_{n+1}) dx_1 \dots dx_n \end{aligned}$$

Since the distribution of X_{ζ^-} on the event of X killed by κ is $1_{y \in I} G(X_0, y) m(y) \kappa(dy)$, we get

$$(3.2.8) \quad \mathbb{P}_x(\zeta > t_n, X_{t_1} \in A_1, \dots, X_{t_n} \in A_n, X_{\zeta^-} \in A_{n+1}) = \int_{A_1 \times \dots \times A_{n+1}} p_{t_1}(x, x_1) m(x_1) \dots p_{t_n - t_{n-1}}(x_{n-1}, x_n) m(x_n) G(x_n, y) m(y) dx_1 \dots dx_n \kappa(dy)$$

The equality between (3.2.7) and (3.2.8) implies the lemma. \square

Next we study the continuity of $(x, y) \mapsto \mu^{x,y}$.

Lemma 3.5. — *Let J be a compact subinterval of I . Then the family of local times of X satisfies: for every $\varepsilon > 0$*

$$\lim_{t \rightarrow 0^+} \sup_{x \in J} \mathbb{P}_x \left(\sup_{y \in I} \ell_{t \wedge \zeta}^y(X) > \varepsilon \right) = 0$$

Proof. — It is enough to prove it in case the killing measure κ is zero because adding a killing measure only lowers $\ell_{t \wedge \zeta}^y(X)$. Without loss of generality we may also assume that the diffusion is on its natural scale, that is to say $w \equiv 2$. Then X is just a time changed Brownian motion on some open subinterval of \mathbb{R} . For a Brownian motion $(B_t)_{t \geq 0}$ the statement is clear. In this case $\mathbb{P}_x \left(\sup_{y \in \mathbb{R}} \ell_{t \wedge \zeta}^y(B) > \varepsilon \right)$ does not depend on x and for a given x

$$\lim_{t \rightarrow 0^+} \mathbb{P}_x \left(\sup_{y \in \mathbb{R}} \ell_{t \wedge \zeta}^y(B) > \varepsilon \right) = 0$$

Otherwise let

$$\mathcal{I}_t := \int_0^t m(X_s) ds$$

Then given the time change that transforms X into a Brownian motion B , we have

$$\ell_t^y(X) = \ell_{\mathcal{I}_t}^y(B)$$

Let $J = [x_0, x_1]$. Let $x_{min} \in I$, $x_{min} < x_0$ and $x_{max} \in I$, $x_{max} > x_1$. Let $T_{x_{min}, x_{max}}$ the first time X hits either x_{min} or x_{max} . Let $s > 0$, $\varepsilon > 0$ and $x \in J$. If $t \leq \frac{s}{\max_{[x_{min}, x_{max}]} m}$ then on the event $T_{x_{min}, x_{max}} \geq t$, \mathcal{I}_t is less or equal to s . So for t small enough

$$\mathbb{P}_x \left(\sup_{y \in I} \ell_{t \wedge \zeta}^y(X) > \varepsilon \right) \leq \mathbb{P}_x \left(\sup_{y \in \mathbb{R}} \ell_s^y(B) > \varepsilon \right) + \mathbb{P}_x(T_{x_{min}, x_{max}} < t)$$

But

$$\mathbb{P}_x(T_{x_{min}, x_{max}} < t) = \mathbb{P}_{x_0}(T_{x_{min}, x_{max}} < t) + \mathbb{P}_{x_1}(T_{x_{min}, x_{max}} < t)$$

and

$$\lim_{t \rightarrow 0^+} \sup_{x \in J} \mathbb{P}_x(T_{x_{min}, x_{max}} < t) = 0$$

Thus

$$\limsup_{t \rightarrow 0^+} \sup_{x \in J} \mathbb{P}_x \left(\sup_{y \in I} \ell_{t \wedge \zeta}^y(X) > \varepsilon \right) \leq \mathbb{P}_x \left(\sup_{y \in \mathbb{R}} \ell_s^y(B) > \varepsilon \right)$$

Letting s go to 0 we get the statement of the lemma. \square

Proposition 3.6. — *Let $t_{max} > 0$. Let F be a bounded functional on finite life-time paths endowed with continuous local times that depends continuously on the path $(\gamma_t)_{0 \leq t \leq T(\gamma)}$ and on $(l_{T(\gamma)}^x(\gamma))_{x \in I}$ where we take the topology of uniform convergence for the occupation densities on I . On top of that we assume that F is zero if $T(\gamma) > t_{max}$. Then the function $(x, y) \mapsto \mu^{x,y}(F(\gamma))$ is continuous on $I \times I$.*

Proof. — If we had assumed that F does only depend on the path regardless to its occupation field then the continuity of $(x, y) \mapsto \mu^{x,y}(F(\gamma))$ would just be a consequence of the continuity of transition densities and of the weak continuity of bridge probability measures. For our proof we further assume that L does not contain any killing measure. If this is not the case, then we can consider a continuous positive L -harmonic function u . Then $Conj(u, L)$ does not contain any killing measure and up to a continuous factor $u(x)u(y)$ gives the same measure $\mu^{x,y}$ (property 3.3 (vii)). We will mainly rely on the representation given by proposition 3.2.

Let $x, y \in I$ and $(x_j, y_j)_{j \geq 0}$ a sequence in $I \times I$ converging to (x, y) . Without loss of generality we assume that $(x_j)_{j \geq 0}$ is increasing. We consider sample paths $(X_t)_{0 \leq t < \zeta}$ and $(X_t^{(j)})_{0 \leq t < \zeta_j}$ of the diffusion of generator L starting from x and each of x_j , coupled on a same probability space in the following way: First we sample X starting from x . Then we sample $X^{(0)}$ starting from x_0 . It starts independently from X until the first time $X_t^{(0)} = X_t$. After that time $X^{(0)}$ sticks to X . This two paths may never meet if one of them dies to early. If $X, X^{(0)}, \dots, X^{(j)}$ are already sampled, we start $X^{(j+1)}$ from x_{j+1} independently from the preceding sample paths until it meets one of them. After that time $X^{(j+1)}$ sticks to the path it has met. Let

$$T^{(j)} := \inf\{t \geq 0 \mid X_t^{(j)} = X_t\}$$

If $X^{(j)}$ does not meet X , we set $T^{(j)} = +\infty$. By construction, $(T^{(j)})_{j \geq 0}$ is a non-increasing sequence. Here we use that there is no killing measure. $T^{(j)}$ is equal in law to the first time two independent sample paths of the diffusion, one starting from x and the other from x_j , meet. Thus the sequence $(T^{(j)})_{j \geq 0}$ converges to 0 in probability. Since it is decreasing, it converges almost surely to 0.

We use reduction to absurdity. The sequence $(\mu^{x_j, y_j}(F(\gamma)))_{j \geq 0}$ is bounded because F is bounded and zero on paths with life-time greater then t_{max} . Assume that it does not converge to $\mu^{x,y}(F(\gamma))$. Then there is a subsequence that converges to a value other than $\mu^{x,y}(F(\gamma))$. We may as well assume that the whole sequence $(\mu^{x_j, y_j}(F(\gamma)))_{j \geq 0}$ converges to a value $v \neq \mu^{x,y}(F(\gamma))$. According to lemma 3.5, the sequence $((\ell_{T^{(j)}}^z(X^{(j)}))_{z \in I})_{j \geq 0}$ of occupation density functions

converges in probability to the null function. Thus there is an extracted subsequence $((\ell_{T^{(j_n)}}^z(X^{(j_n)}))_{z \in I})_{n \geq 0}$ that converges almost surely uniformly to the null function. We will show that $(\mu^{x_{j_n}, y_{j_n}}(F(\gamma)))_{n \geq 0}$ converges to $\mu^{x, y}(F(\gamma))$ and obtain a contradiction.

For $z \in I$ and $l > 0$ let

$$\tau_l^z := \inf\{t \geq 0 \mid \ell_t^z(X) > l\}$$

and

$$\tau_{j, l}^z := \inf\{t \geq 0 \mid \ell_t^z(X^{(j)}) > l\}$$

Then according to proposition 3.2

$$\begin{aligned} \mu^{x, y}(F(\gamma)) &= \mathbb{E} \left[\int_0^{\ell_{t_{max} \wedge \zeta}^y(X)} F((X_s)_{0 \leq s \leq \tau_l^y}) dl \right] \\ \mu^{x_j, y_j}(F(\gamma)) &= \mathbb{E} \left[\int_0^{\ell_{t_{max} \wedge \zeta_j}^{y_j}(X^{(j)})} F((X_s^{(j)})_{0 \leq s \leq \tau_{j, l}^{y_j}}) dl \right] \end{aligned}$$

For any $z \in I$, if $\tau_{j, l}^z \in [T^{(j)}, \zeta_j)$ then $\tau_{j, l}^z = \tau_{l'}^z$ where

$$l' = l + \ell_{T^{(j)}}^z(X) - \ell_{T^{(j)}}^z(X^{(j)})$$

Along the subset of indices $(j_n)_{n \geq 0}$, $\tau_{j_n, l}^{y_{j_n}}$ converges to τ_l^y for every $l \in (0, l_\zeta^y(X))$ except possibly the countable set of values of l where $l \mapsto \tau_{j, l}^y$ jumps. For any l such that $\tau_{j_n, l}^{y_{j_n}}$ converges to τ_l^y , the path $(X_s^{(j_n)})_{0 \leq s \leq \tau_{j_n, l}^{y_{j_n}}}$ converges to the path $(X_s)_{0 \leq s \leq \tau_l^y}$. Moreover for such l the occupation densities $(\ell_{\tau_{j_n, l}^{y_{j_n}}}^{z_{y_{j_n}}}(X^{(j_n)}))_{z \in I}$ converge uniformly to $(\ell_{\tau_l^y}^z(X))_{z \in I}$. Indeed

$$\ell_{\tau_{j_n, l}^{y_{j_n}}}^{z_{y_{j_n}}}(X^{(j_n)}) = \ell_{\tau_{j_n, l}^{y_{j_n}}}(X) - \ell_{T^{(j)}}^z(X) + \ell_{T^{(j)}}^z(X^{(j_n)})$$

Thus for all $l \in (0, l_\zeta^y(X))$, except possibly countably many,

$$\lim_{n \rightarrow +\infty} F((X_s^{(j_n)})_{0 \leq s \leq \tau_{j_n, l}^{y_{j_n}}}) = F((X_s)_{0 \leq s \leq \tau_l^y})$$

For n large enough, $\zeta_j = \zeta$ and $\ell_{t_{max} \wedge \zeta_{j_n}}^{y_{j_n}}(X^{(j_n)})$ converges to $\ell_{t_{max} \wedge \zeta}^y(X)$. It follows that the following almost sure convergence holds

(3.2.9)

$$\lim_{n \rightarrow +\infty} \int_0^{\ell_{t_{max} \wedge \zeta_{j_n}}^{y_{j_n}}(X^{(j_n)})} F((X_s^{(j_n)})_{0 \leq s \leq \tau_{j_n, l}^{y_{j_n}}}) dl = \int_0^{\ell_{t_{max} \wedge \zeta}^y(X)} F((X_s)_{0 \leq s \leq \tau_l^y}) dl$$

The left-hand side of (3.2.9) is dominated by $\|F\|_{+\infty} \ell_{t_{max} \wedge \zeta_{j_n}}^{y_{j_n}}(X^{(j_n)})$. In order to conclude that the almost sure convergence (3.2.9) is also an \mathbb{L}^1 convergence we need only to show that

$$(3.2.10) \quad \mathbb{E} \left[\left| \ell_{t_{max} \wedge \zeta_{j_n}}^{y_{j_n}}(X^{(j_n)}) - \ell_{t_{max} \wedge \zeta}^y(X) \right| \right] = 0$$

We already know that $\ell_{t_{max} \wedge \zeta_{j_n}}^{y_{j_n}}(X^{(j_n)})$ converges almost surely to $\ell_{t_{max} \wedge \zeta}^y(X)$. Moreover

$$\mathbb{E} \left[\ell_{t_{max} \wedge \zeta_{j_n}}^{y_{j_n}}(X^{(j_n)}) \right] = \int_0^{t_{max}} p_t(x_{j_n}, y_{j_n})$$

and

$$\mathbb{E} \left[\ell_{t_{max} \wedge \zeta}^y(X) \right] = \int_0^{t_{max}} p_t(x, y)$$

It follows that the expectations converge. By Scheffe's lemma, the \mathbb{L}^1 convergence (3.2.10) holds.

We have shown that there is always a subsequence $(\mu^{x_{j_n}, y_{j_n}}(F(\gamma)))_{n \geq 0}$ that converges to $\mu^{x, y}(F(\gamma))$ which contradict the convergence of $(\mu^{x_j, y_j}(F(\gamma)))_{j \geq 0}$ to a different value. \square

3.3. The measure μ^* on unrooted loops

The measure $\mu^{x, x}$ can be seen as a measure on the space of rooted loops \mathfrak{L} . Next we define a natural measure μ_L^* on \mathfrak{L}^* following the pattern (3.2.1)

Definition 3.7. — Let μ_L be the following measure on \mathfrak{L} :

$$\mu_L(d\gamma) := \int_{t>0} \int_{x \in I} \mathbb{P}_{x, x}^t(d\gamma) p_t(x, x) m(x) dx \frac{dt}{t} = \frac{1}{T(\gamma)} \int_{x \in I} \mu_L^{x, x}(d\gamma) m(x) dx$$

$\mu_L^* := \pi_* \mu_L$ is a measure on \mathfrak{L}^* .

We will drop the subscript L whenever there is no ambiguity on L . The definition 2 does not depend on the choice of the speed measure $m(x) dx$. The measures μ and μ^* are σ -finite but not finite. They satisfy the following elementary properties:

Property 3.8. — (i) μ is invariant by time reversal.

(ii) If \tilde{I} is an open subinterval of I then

$$\mu_{L|\tilde{I}}(d\gamma) = 1_{\gamma \text{ contained in } \tilde{I}} \mu_L(d\gamma)$$

(iii) If $\tilde{\kappa}$ is a positive Radon measure on I then

$$\mu_{L-\tilde{\kappa}}(d\gamma) = \exp \left(- \int_I \ell^z(\gamma) m(z) \tilde{\kappa}(dz) \right) \mu_L(d\gamma)$$

(iv) If A is a change of scale function then

$$\mu_{\text{Scale}_A^{\text{gen}} L} = \text{Scale}_{A^*} \mu_L$$

(v) If h is a positive continuous function on I such that $\frac{d^2 h}{dx^2}$ is a signed measure and Lu is a negative measure then

$$\mu_{\text{Conj}(h, L)} = \mu_L$$

Same properties hold for μ^* .

The measures μ and μ^* contain some information on the diffusion X but the invariance by h-transforms (property 3.8 (v)) shows that they do not capture its asymptotic behaviour. In the section 3.4 we will prove a converse to the property 3.8 (v). In our setting, most important examples of h-transforms are:

- The Bessel 3 process on $(0, +\infty)$ is an h-transform of the Brownian motion on $(0, +\infty)$, killed when hitting 0, through the function $x \mapsto x$.
- The Brownian motion on \mathbb{R} killed with uniform rate κdx (i.e. κ constant) is an h-transform of the drifted Brownian motion on \mathbb{R} with constant drift $\sqrt{2\kappa}$, through the function $x \mapsto e^{-\sqrt{2\kappa}x}$.

In the sequel we will be interested mostly in μ^* and not μ . As it will be clear from the next propositions, the measure μ^* has some nice features that μ does not.

Proposition 3.9. — *Let $v \in [0, 1]$. Then $shift_{v*}\mu = \mu$. In particular*

$$(3.3.1) \quad \mu(\cdot) = \int_{v \in [0,1]} shift_{v*}\mu(\cdot) dv$$

Proof. — For a rooted loop γ of life-time $T(\gamma)$ we will introduce γ_1 the path restricted to time interval $[0, vT(\gamma)]$ and γ_2 the path restricted to $[vT(\gamma), T(\gamma)]$. By bridge decomposition property, the measure $\mu(d\gamma_1, d\gamma_2)$ equals

$$\int_{t>0} \int_I \int_I \mathbb{P}_{x,y}^{vt} (d\gamma_1) \mathbb{P}_{y,x}^{(1-v)t} (d\gamma_2) p_{vt}(x, y) p_{(1-v)t}(y, x) m(y) dy m(x) dx \frac{dt}{t}$$

Since γ_1 and γ_2 play symmetric roles, changing the order of γ_1 and γ_2 does not change the measure μ . \square

Formula (3.3.1) shows that we can get back to the measure μ from the measure μ^* by cutting the circle parametrizing a loop in \mathfrak{L}^* in a point chosen uniformly on this circle, in order to separate the start from the end.

Corollary 3.10. — *Let F be a positive measurable functional on \mathfrak{L} . Then the map $\gamma \mapsto \int_0^1 F(shift_v(\gamma)) dv$ is $\pi^{-1}(\mathcal{B}_{\mathfrak{L}^*})$ -measurable and*

$$\frac{d(F(\gamma)\mu)}{d\mu} \Big|_{\pi^{-1}(\mathcal{B}_{\mathfrak{L}^*})} = \int_0^1 F(shift_v(\gamma)) dv$$

Proof. — We need only to show that for every F' measurable functional on \mathfrak{L}^* :

$$(3.3.2) \quad \int_{\mathfrak{L}} F(\gamma) F'(\pi(\gamma)) \mu(d\gamma) = \int_0^1 \int_{\mathfrak{L}} F(shift_v(\gamma)) F'(\pi(\gamma)) \mu(d\gamma) dv$$

From proposition 3.9 follows that for every $v \in [0, 1]$:

$$(3.3.3) \quad \int_{\mathfrak{L}} F(\gamma) F'(\pi(\gamma)) \mu(d\gamma) = \int_{\mathfrak{L}} F(shift_v(\gamma)) F'(\pi(\gamma)) \mu(d\gamma)$$

Integrating (3.3.3) on $[0, 1]$ leads to (3.3.2). \square

The next identity appears in [13] in the setting of Markov jump processes on graphs. It can be generalized to a wider class of Markov processes admitting local times (see lemma 2.2 in [10]). We will give a short proof that suits our framework.

Corollary 3.11. — *Let $x \in I$. Then*

$$(3.3.4) \quad \ell^x(\gamma)\mu^*(d\gamma) = \pi_*\mu^{x,x}(d\gamma)$$

For $l > 0$, let $\mathbb{P}_x^{\tau_l^x}(\cdot)$ be the law of the sample paths of a diffusion X of generator L , started from x , until the time τ_l^x when $\ell_t^x(X)$ hits l , conditioned by $\tau_l^x < \zeta$. Then

$$(3.3.5) \quad 1_{\gamma \text{ visits } x}\mu^*(d\gamma) = \int_0^{+\infty} \pi_*\mathbb{P}_x^{\tau_l^x}(d\gamma)e^{-\frac{l}{G(x,x)}}\frac{dl}{l}$$

Conventionally we set $G(x,x) = +\infty$ if X is recurrent.

Proof. — Let $\varepsilon > 0$ such that $[x - \varepsilon, x + \varepsilon] \subseteq I$. Let $T_{[x-\varepsilon, x+\varepsilon]}(\gamma)$ be the time a loop γ spends in $[x - \varepsilon, x + \varepsilon]$. From the identity (3.3.1) follows that

$$\frac{T_{[x-\varepsilon, x+\varepsilon]}(\gamma)}{T(\gamma)}\mu^*(d\gamma) = \frac{1}{T(\gamma)} \int_{x-\varepsilon}^{x+\varepsilon} \pi_*\mu^{z,z}(d\gamma)m(z)dz$$

and simplifying $T(\gamma)$:

$$T_{[x-\varepsilon, x+\varepsilon]}(\gamma)\mu^*(d\gamma) = \int_{x-\varepsilon}^{x+\varepsilon} \pi_*\mu^{z,z}(d\gamma)m(z)dz$$

Using local times we rewrite the previous expression as

$$(3.3.6) \quad \frac{\int_{x-\varepsilon}^{x+\varepsilon} \ell^z(\gamma)m(z)dz}{\int_{x-\varepsilon}^{x+\varepsilon} m(z)dz}\mu^*(d\gamma) = \frac{1}{\int_{x-\varepsilon}^{x+\varepsilon} m(z)dz} \int_{x-\varepsilon}^{x+\varepsilon} \pi_*\mu^{z,z}(d\gamma)m(z)dz$$

Let $\varepsilon_0 > 0$ such that $[x - \varepsilon_0, x + \varepsilon_0] \subseteq I$. Let F be a continuous bounded functional on loops endowed with continuous local times such that F is zero if the life-time of the loop exceeds $t_{max} > 0$ and if $\sup_{z \in [x-\varepsilon_0, x+\varepsilon_0]} \ell^z(\gamma)$ exceeds l_{max} . According to the proposition 2.5, the right-hand side of (3.3.6) applied to F converges as $\varepsilon \rightarrow 0$ to $(\pi_*\mu^{x,x})(F(\gamma))$. By dominated convergence it follows that the left-hand side of (3.3.6) applied to F converges as $\varepsilon \rightarrow 0$ to

$$\int_{\mathcal{L}^*} \ell^x(\gamma)F(\gamma)\mu^*(d\gamma)$$

Thus we have the equality

$$(3.3.7) \quad \int_{\mathcal{L}^*} \ell^x(\gamma)F(\gamma)\mu^*(d\gamma) = (\pi_*\mu^{x,x})(F(\gamma))$$

The set of test functionals F that satisfy (3.3.7) is large enough to deduce the equality (3.3.4) between measures.

From proposition 3.2 follows that

$$\mu^{x,x}(\cdot) = \int_0^{+\infty} \mathbb{P}_x^{\tau_l^x}(\cdot)e^{-\frac{l}{G(x,x)}}dl$$

Applying (3.3.4) to the above disintegration, we get (3.3.5). \square

Corollary 3.12. — *Let V be a positive continuous function on I . We consider a time change with speed V : $ds = V(x)dt$. Then*

$$(3.3.8) \quad \mu_{\frac{1}{V}L}^* = \text{Speed}_{V*} \mu_L^*$$

Proof. — By definition 3.7 and property 3.3 (vi):

$$\mu_{\frac{1}{V}L}(d\gamma) = \frac{1}{T(\gamma)} \int_0^{T(\gamma)} \frac{V(\gamma(0))}{V(\gamma(s))} ds \text{Speed}_{V*}(\mu_L(d\gamma))$$

Applying corollary 3.10 we obtain:

$$\frac{d\text{Speed}_{V*} \mu_L}{d\mu_{\frac{1}{V}L}} \Big|_{\pi^{-1}(\mathcal{B}_{\mathbb{R}^*})} = \frac{\int_0^1 V^{-1}(\gamma(vT(\gamma))) dv}{\frac{1}{T(\gamma)} \int_0^{T(\gamma)} V^{-1}(\gamma(s)) ds} = 1$$

This concludes. \square

In dimension two, the time change covariance of the measure μ^* on loops plays a key role for the construction of the Conformal Loop Ensembles (CLE) using loop soups as in [26]: Let D be an open domain of the complex plane, $(B_t)_{0 \leq t < \zeta}$ the two-dimensional standard Brownian motion in D killed when hitting ∂D and μ^* the corresponding measure on loops. If $f : D \rightarrow D$ is a conformal map, then $(f(B_t))_{0 \leq t < \zeta}$ is a time changed Brownian motion. If we consider μ^* not as a measure on loops parametrized by time but a measure on the geometrical drawings of loops, then μ^* is invariant by the transformation $(\gamma(t))_{0 \leq t \leq T(\gamma)} \mapsto (f(\gamma(t)))_{0 \leq t \leq T(\gamma)}$. This is proved in [19].

Given that μ^* is invariant through h-transforms and covariant with the change of scale and change of time, if X is a recurrent diffusion, then up to a change of scale and time, μ^* is the same as for the Brownian motion on \mathbb{R} , and if X is a transient diffusion, even if the killing measure κ is non-zero, then up to a change of scale and time, μ^* is the same as for the Brownian motion on a bounded interval, killed when it hits the boundary.

3.4. Multiple local times

In this subsection we define the multiple local time functional on loops. Corollary 3.11 gives a link between the measure μ^* and the measures $(\mu^{x,x})_{x \in I}$. Using multiple local times we will get a further relation between μ^* and $(\mu^{x,y})_{x,y \in I}$. This will allow us to prove a converse to the property 3.8 (v): two diffusions that have the same measure on unrooted loops are related through an h-transform.

Definition 3.13. — If $(\gamma(t))_{0 \leq t \leq T(\gamma)}$ is a continuous path in I having a family of local times $(\ell_t^x(\gamma))_{x \in I, 0 \leq t \leq T(\gamma)}$ relatively to the measure $m(x)dx$, we introduce multiple local times $\ell^{x_1, x_2, \dots, x_n}(\gamma)$ for $x_1, x_2, \dots, x_n \in I$:

$$\ell^{x_1, x_2, \dots, x_n}(\gamma) := \int_{0 \leq t_1 \leq t_2 \leq \dots \leq t_n \leq T(\gamma)} d_{t_1} \ell_{t_1}^{x_1}(\gamma) d_{t_2} \ell_{t_2}^{x_2}(\gamma) \dots d_{t_n} \ell_{t_n}^{x_n}(\gamma)$$

If $\gamma \in \mathfrak{L}$ and has local times, we introduce circular local times for γ :

$$\ell^{*x_1, x_2, \dots, x_n}(\gamma) := \sum_{\substack{c \text{ circular} \\ \text{permutation} \\ \text{of } \{1, 2, \dots, n\}}} \ell^{x_{c(1)}, x_{c(2)}, \dots, x_{c(n)}}(\gamma)$$

$\ell^{*x_1, x_2, \dots, x_n}$ being invariant under the transformations $(shift_v)_{v \in [0, 1]}$, we see it as a functional defined on \mathfrak{L}^* .

Multiple local times of the form $\ell^{x, x, \dots, x}(\gamma)$, called self intersection local times, were studied by Dynkin in [9]. Circular local times were introduced by Le Jan in [13].

Let $n \in \mathbb{N}^*$ and $p \in \{1, \dots, n\}$. Let $Shuffle_{p, n}$ be the set of permutations σ of $\{1, \dots, n\}$ such that for all $i \leq j \in \{1, \dots, p\}$, $\sigma(i) \leq \sigma(j)$ and for all $i \leq j \in \{p+1, \dots, n\}$, $\sigma(i) \leq \sigma(j)$. Permutations in $Shuffle_{p, n}$ are obtained by shuffling two card decks $\{1, \dots, p\}$ and $\{p+1, \dots, n\}$. Let $Shuffle'_{p, n}$ be the permutations of $\{1, \dots, n\}$ of the form $\sigma \circ c$ where c is a circular permutation of $\{p+1, \dots, n\}$ and $\sigma \in Shuffle_{p, n}$ satisfies $\sigma(1) = 1$. One can check that

Property 3.14. — For all $x_1, \dots, x_p, x_{p+1}, \dots, x_n \in I$:

– (i)

$$\ell^{x_1, \dots, x_p}(\gamma) \ell^{x_{p+1}, \dots, x_n}(\gamma) = \sum_{\sigma \in Shuffle_{p, n}} \ell^{x_{\sigma(1)}, \dots, x_{\sigma(p)}, x_{\sigma(p+1)}, \dots, x_{\sigma(n)}}(\gamma)$$

– (ii)

$$\ell^{*x_1, \dots, x_p}(\gamma) \ell^{*x_{p+1}, \dots, x_n}(\gamma) = \sum_{\sigma' \in Shuffle'_{p, n}} \ell^{x_{\sigma'(1)}, \dots, x_{\sigma'(p)}, x_{\sigma'(p+1)}, \dots, x_{\sigma'(n)}}(\gamma)$$

The equality 3.14 (ii) appears in [13]. It is also shown in [13] that for transient Markov jump processes:

$$(3.4.1) \quad \int \ell^{*x_1, x_2, \dots, x_n}(\gamma) \mu(d\gamma) = G(x_1, x_2) \times \dots \times G(x_{n-1}, x_n) \times G(x_n, x_1)$$

It turns out that we have more: We consider L a generator of a diffusion on I of form (2.2.1). If γ_i for $i \in \{1, 2, \dots, n-1\}$ is a continuous path from x_i to x_{i+1} , then we can concatenate $\gamma_1, \gamma_2, \dots, \gamma_{n-1}$ to obtain a continuous path $\gamma_1 \triangleleft \gamma_2 \triangleleft \dots \triangleleft \gamma_{n-1}$ from x_1 to x_n . Let $\mu^{x_1, x_2} \triangleleft \dots \triangleleft \mu^{x_{n-1}, x_n}$ be the image measure $\mu^{x_1, x_2} \otimes \dots \otimes \mu^{x_{n-1}, x_n}$ by this concatenation procedure.

Proposition 3.15. — The following absolute continuity relations hold:

- (i) $(\mu^{x_1, x_2} \triangleleft \dots \triangleleft \mu^{x_{n-1}, x_n})(d\gamma) = \ell^{x_2, \dots, x_{n-1}}(\gamma) \mu^{x_1, x_n}(d\gamma)$
- (ii) $\pi_*(\mu^{x_1, x_2} \triangleleft \dots \triangleleft \mu^{x_{n-1}, x_n} \triangleleft \mu^{x_n, x_1})(d\gamma) = \ell^{*x_1, x_2, \dots, x_n}(\gamma) \mu^*(d\gamma)$

Proof. — (i): Let $((X_t^{(j)})_{0 \leq t < \zeta_j})_{0 \leq j \leq n-1}$ be $n-1$ independent diffusions of generator L , with $X_0^{(j)} = x_j$. For $l \geq 0$, let

$$\tau_{j,l}^{x_{j+1}} := \inf \left\{ t_j \geq 0 \mid \ell_{t_j}^{x_{j+1}}(X^{(j)}) > l \right\}$$

According to proposition 3.2, $(\mu^{x_1, x_2} \triangleleft \dots \triangleleft \mu^{x_{n-1}, x_n})(F(\gamma))$ equals

$$(3.4.2) \quad \mathbb{E} \left[\int_{l_j < \ell_{\zeta_j}^{x_{j+1}}(X^{(j)}), 1 \leq j \leq n-1} F((X_t^{(1)})_{0 \leq t \leq \tau_{1,l_1}^{x_2}} \triangleleft \dots \triangleleft (X_t^{(n-1)})_{0 \leq t \leq \tau_{n-1, l_{n-1}}^{x_n}}) dl_1 \dots dl_{n-1} \right]$$

Let $(X_t)_{0 \leq t < \zeta}$ be an other diffusion of generator L . Let

$$\tau_{l_1} := \inf \{ t \geq 0 \mid l_t^{x_2}(X) > l_1 \}$$

and recursively defined

$$\tau_{l_1, \dots, l_{j-1}, l_j} := \inf \{ t \geq \tau_{l_1, \dots, l_{j-1}} \mid \ell_t^{x_{j+1}}(X) > l_j \}$$

Then by strong Markov property, (3.4.2) equals

$$\mathbb{E} \left[\int 1_{\tau_{l_1, \dots, l_{n-1}} \leq \zeta} F((X_t)_{0 \leq t \leq \tau_{l_1, \dots, l_{n-1}}}) dl_1 \dots dl_{n-1} \right]$$

which in turn equals

$$(3.4.3) \quad \mathbb{E} \left[\int 1_{\forall j, t_j < \zeta} F((X_t)_{0 \leq t \leq t_{n-1}}) d_{t_1} \ell_{t_1}^{x_2}(X) \dots d_{t_{n-1}} \ell_{t_{n-1}}^{x_n}(X) \right]$$

By proposition 3.2, (3.4.3) equals $\int \ell^{x_1, \dots, x_{n-1}}(\gamma) F(\gamma) \mu^{x_1, x_n}(d\gamma)$.

(ii): According to the identity (i) and corollary 3.10, we have

$$\pi_*(\mu^{x_1, x_2} \triangleleft \dots \triangleleft \mu^{x_{n-1}, x_n} \triangleleft \mu^{x_n, x_1})(d\gamma) = \int_0^1 \ell^{x_2, \dots, x_n}(\text{shift}_v(\gamma)) dv \pi_* \mu^{x_1, x_1}(d\gamma)$$

According to corollary 3.11

$$\int_0^1 \ell^{x_2, \dots, x_n}(\text{shift}_v(\gamma)) dv \pi_* \mu^{x_1, x_1}(d\gamma) = \ell^{x_1}(\gamma) \int_0^1 \ell^{x_2, \dots, x_n}(\text{shift}_v(\gamma)) dv \mu^*(d\gamma)$$

But

$$\ell^{x_1}(\gamma) \int_0^1 \ell^{x_2, \dots, x_n}(\text{shift}_v(\gamma)) dv = \ell^{*x_1, x_2, \dots, x_n}(\gamma)$$

which ends the proof. \square

The proposition 3.15 (ii) implies (3.4.1).

Proposition 3.16. — *If L and \tilde{L} are two generators of diffusions on I of the form (2.2.1) such that $\mu_L^* = \mu_{\tilde{L}}^*$, then there is a positive continuous function h on I such that $\frac{d^2 h}{dx^2}$ is a signed measure, Lh a negative measure and $\tilde{L} = \text{Conj}(h, L)$. If the diffusion of generator L is recurrent then $\tilde{L} = L$.*

Proof. — Let $m(x)dx$ be a speed measure for L and $\tilde{m}(x)dx$ be a speed measure for \tilde{L} . First let's assume that both L and \tilde{L} are generators of transient diffusions. Applying the identity (3.4.1) to $\int_{\mathfrak{G}^*} \ell^{*x,y}(\gamma) \mu^*(d\gamma)$ we get that for all $x, y \in I$:

$$(3.4.4) \quad G_{\tilde{L}}(x, y)G_{\tilde{L}}(y, x)\tilde{m}(x)\tilde{m}(y) = G_L(x, y)G_L(y, x)m(x)m(y)$$

and for all $x, y, z \in I$:

$$(3.4.5) \quad G_{\tilde{L}}(x, y)G_{\tilde{L}}(y, z)G_{\tilde{L}}(z, x)\tilde{m}(x)\tilde{m}(y)\tilde{m}(z) = G_L(x, y)G_L(y, z)G_L(z, x)m(x)m(y)m(z)$$

Fix $x_0 \in I$. Let h be

$$h(x) := \frac{G_{\tilde{L}}(x_0, x)\tilde{m}(x)}{G_L(x_0, x)m(x)}$$

h is positive and continuous. $\frac{1}{h(x)}G_L(x, y)h(y)m(y)$ equals:

$$(3.4.6) \quad \frac{G_L(x_0, x)G_L(x, y)G_L(y, x_0)m(x_0)m(x)m(y)}{G_{\tilde{L}}(x_0, x)G_{\tilde{L}}(x, y)G_{\tilde{L}}(y, x_0)\tilde{m}(x_0)\tilde{m}(x)\tilde{m}(y)} \\ \times \frac{G_{\tilde{L}}(x_0, y)G_{\tilde{L}}(y, x_0)\tilde{m}(x_0)\tilde{m}(y)}{G_L(x_0, y)G_L(y, x_0)m(x_0)m(y)} \times G_{\tilde{L}}(x, y)\tilde{m}(y)$$

Applying (3.4.4) and (3.4.5) to (3.4.6) we get that

$$(3.4.7) \quad \frac{1}{h(x)}G(x, y)h(y)m(y) = G_{\tilde{L}}(x, y)\tilde{m}(y)$$

Applying (3.4.7) once to (x, y) and once do (x, x) we get that

$$(3.4.8) \quad h(y) = h(x) \frac{G_{\tilde{L}}(x, y)}{G(x, y)} \frac{G(y, y)}{G_{\tilde{L}}(y, y)}$$

From (3.4.8) we deduce that $\frac{d^2 h}{dx^2}$ is a signed measure. From (3.4.7) we deduce that $\tilde{L} = Conj(h, L)$. $-Lh$ is the killing measure of \tilde{L} and is positive.

If we no longer assume that L and \tilde{L} generate transient diffusions then consider $\lambda > 0$. Then $\mu_{L-\lambda}^* = \mu_{\tilde{L}-\lambda}^*$. According to the above, there is h positive continuous function on I such that $\frac{d^2 h}{dx^2}$ is a signed measure and

$$\tilde{L} - \lambda = Conj(h, L - \lambda) = Conj(h, L) - \lambda$$

Then $\tilde{L} = Conj(h, L)$ and necessarily Lh is a negative measure.

The class of recurrent diffusions is preserved by h-transforms. So if L is the generator of a recurrent diffusion then so is \tilde{L} , and thus h is bound to satisfy $Lh = 0$. But since the diffusion of L is recurrent, the only solutions to $Lh = 0$ are constant functions. Thus $\tilde{L} = L$. \square

3.5. A disintegration of μ^* induced by the Vervaat's transformation

By conditioning the measure μ by the life-time of loops we get a sum of bridge measures. In this section we will disintegrate the measure μ^* as a measure on the minimal value of the loop and its behaviour above this value. By doing this way we will obtain a sum of excursion measures $\eta_{exc}^{>x}$. In case of Brownian loops on \mathbb{R} this disintegration will follow from the Vervaat's bridge to excursion transformation. The case of general diffusion will be obtained using covariance of the measure on loops by time and scale change, restriction to a subinterval, killing, as well as invariance by h-transforms.

Theorem(Vervaat). — ([30],[3]) *Let $(\gamma(s))_{0 \leq s \leq t}$ be a random path following the Brownian bridge probability measure $\mathbb{P}_{BM,0,0}^t(\cdot)$. Let $s_{min} := \operatorname{argmin} \gamma$. Then the path*

$$s \mapsto -\min \gamma + (\operatorname{shift}_{\frac{s_{min}}{t}} \gamma)(s)$$

has the law of a positive Brownian excursion of life-time t .

In the sequel if η is a measure on paths and $x \in \mathbb{R}$, we will write $(x + \eta)$ for the image of η by $\gamma \mapsto x + \gamma$. $\eta_{BM}^{>0}$ will be the Levy-Itô measure on positive Brownian excursions and $\eta_{t,BM}^{>0}$ the probability measure on positive Brownian excursions of duration t . Given a continuous loop $(\gamma_t)_{0 \leq t \leq T(\gamma)}$ and t_{min} the first time γ hits $\min \gamma$, let $\mathcal{V}(\gamma)$ be the transformation $\operatorname{shift}_{\frac{t_{min}}{T(\gamma)}}$. \mathcal{V} is $\mathcal{B}_{\mathcal{L}}$ -measurable.

Proposition 3.17. — *Let μ_{BM}^* be the measure on loops associated to the Brownian motion on \mathbb{R} . Then:*

$$(3.5.1) \quad \mu_{BM}^*(d\gamma) = 2 \int_{a \in \mathbb{R}} \pi_*(a + \eta_{BM}^{>0})(d\gamma) da$$

The measure on $(\min \gamma, \max \gamma)$ induced by μ_{BM}^ is $1_{a < b}(b - a)^{-2} da db$. Let $a < b \in \mathbb{R}$ and $\rho, \tilde{\rho}$ two independent Bessel 3 processes starting from 0. Let T_{b-a} and \tilde{T}_{b-a} be the first times ρ respectively $\tilde{\rho}$ hits $b - a$. Let $(\beta_t)_{0 \leq t \leq T_{b-a} + \tilde{T}_{b-a}}$ be the path*

$$\beta_t := \begin{cases} a + \rho_t & \text{if } t \leq T_{b-a} \\ a + \tilde{\rho}_{T_{b-a} + \tilde{T}_{b-a} - t} & \text{if } t \geq T_{b-a} \end{cases}$$

Then the law of $(\beta_t)_{0 \leq t \leq T_{b-a} + \tilde{T}_{b-a}}$ is the probability measure obtained by conditioning the measure μ_{BM}^ by $(\min \gamma, \max \gamma) = (a, b)$.*

Proof. — For the Brownian motion on \mathbb{R} , μ_{BM} writes

$$\mu_{BM}(\cdot) = \int_{x \in \mathbb{R}} \int_{t > 0} (x + \mathbb{P}_{BM,0,0}^t(\cdot)) \frac{dt}{\sqrt{2\pi t^3}} dx$$

Let $\chi(a)da$ be the law of the minimum of the bridge under $\mathbb{P}_{BM,0,0}^t$. Applying the Vervaat's transformation, we get that

$$\mathcal{V}_* \mu_{BM}(\cdot) = \int_{a \in \mathbb{R}} \int_{t > 0} \left(\int_{x > a} \chi(x-a) dx \right) (a + \eta_{t,BM}^{>0})(\cdot) \frac{dt}{\sqrt{2\pi t^3}} da$$

Since $\int_{x > a} \chi(x-a) dx = 1$, the right-hand side above equals

$$\int_{a \in \mathbb{R}} \int_{t > 0} (a + \eta_{t,BM}^{>0})(\cdot) \frac{dt}{\sqrt{2\pi t^3}} da$$

But

$$\int_{t > 0} (a + \eta_{t,BM}^{>0})(\cdot) \frac{dt}{\sqrt{2\pi t^3}} = 2(a + \eta_{BM}^{>0})(\cdot)$$

The equality (3.5.1) follows. The rest of the proposition is a consequence of the William's representation of Brownian excursions. \square

Corollary 3.18. — *Let I be an open interval of \mathbb{R} and $\lambda \geq 0$. Let μ^* be the measure on loops in I associated to the generator $\frac{1}{2} \frac{d^2}{dx^2} - \lambda$. Given a loop $(\gamma(t))_{0 \leq t \leq T(\gamma)}$, let $R(\gamma)$ be the loop*

$$R(\gamma) := (\max \gamma + \min \gamma - \gamma(t))_{0 \leq t \leq T(\gamma)}$$

that is the image of γ through reflection relatively to $\frac{\max \gamma + \min \gamma}{2}$. Then

$$R_* \mu^* = \mu^*$$

Proof. — It is enough to prove this in case $\lambda = 0$ and $I = \mathbb{R}$. Otherwise we multiply the measure μ_{BM}^* by a density function that is left invariant by R . Then we use the description of the measure μ_{BM}^* conditioned by the value of $(\min \gamma, \max \gamma)$ and the fact that if $a > 0$, $(\rho_t)_{t \geq 0}$ is a Bessel 3 process starting from 0 and T_b is the first time it hits b , then $(y - \rho_{T_b - t})_{0 \leq t \leq T_b}$ has the same law as $(\rho_t)_{0 \leq t \leq T_b}$ (see [24], chapter VII, §4). \square

Now we consider that L is a generator of a diffusion on I of form (2.2.1). Given a point $x_0 \in I$, u^{+,x_0} and u^{-,x_0} will be the L -harmonic functions satisfying the initial conditions $u^{+,x_0}(x_0) = u^{-,x_0}(x_0) = 0$, $\frac{du^{+,x_0}}{dx}(x_0^+) = 1$ and $\frac{du^{-,x_0}}{dx}(x_0^-) = -1$. If $x \leq y \in I$ then

$$(3.5.2) \quad w(y)u^{-,y}(x) = w(x)u^{+,x}(y)$$

Indeed, the Wronskian $W(u^{-,y}, u^{+,x})$ takes in x the value $u^{-,y}(x)$ and in y the value $u^{+,x}(y)$, and the ratio $\frac{1}{w(z)}W(u^{-,y}, u^{+,x})(z)$ is constant. If $\kappa = 0$, then the both sides of (3.5.2) equal $\int_x^y w(z)dz$. u^{+,x_0} is positive on $I \cap (x_0, +\infty)$ and u^{-,x_0} is positive on $I \cap (-\infty, x_0)$. Let L^{+,x_0} be $\text{Conj}(u^{+,x_0}, L)$ restricted to $I \cap (x_0, +\infty)$ and L^{-,x_0} be $\text{Conj}(u^{-,x_0}, L)$ restricted to $I \cap (-\infty, x_0)$. L^{+,x_0} and L^{-,x_0} are generators of transient diffusions without killing measures. If L is the generator of the Brownian motion on \mathbb{R} , then $L^{+,0}$ is just the generator of a Bessel 3 process. In general case, x_0 is an entrance boundary for L^{+,x_0} and L^{-,x_0} , that is to say a diffusion started from

$x \neq x_0$ will never reach the boundary at x_0 , and we can also start this diffusions at the boundary point x_0 , in which case it will be immediately repelled away from x_0 . Let $x \in I$ and $(\rho_t^{+,x})_{0 \leq t < \zeta^{+,x}}$ be a diffusion of generator $L^{+,x}$ starting from x . Let $y \in I$, $y > x$. Let $T_y^{+,x}$ be the first time $\rho^{+,x}$ hits y and $\widehat{T}_y^{+,x}$ the last time it visits y . Then $(\rho_{\widehat{T}_y^{+,x}+t}^{+,x})_{0 \leq t < \zeta^{+,x} - \widehat{T}_y^{+,x}}$ is a diffusion of generator $L^{+,y}$ starting from y . Let $(\rho_t^{-,y})_{0 \leq t < \zeta^{-,y}}$ be a diffusion of generator $L^{-,y}$ starting from y and $T_x^{-,y}$ the first time it hits x . Then $(\rho_t^{+,x})_{0 \leq t \leq T_y^{+,x}}$ and $(\rho_{T_x^{-,y}-t}^{-,y})_{0 \leq t \leq T_x^{-,y}}$ are equal in law: Indeed let C be the constant

$$C = \frac{w(z)}{W(u^{-,y}, u^{+,x})(z)}$$

The Green's operator of $\rho^{+,x}$ killed in y is

$$((-L_{|(x,y)}^{+,x})^{-1}f)(x') = C \int_x^y u^{+,x}(x' \wedge y') u^{-,y}(x' \vee y') \frac{u^{+,x}(y')}{u^{+,x}(x')} m(y') dy'$$

and the Green's operator of $\rho^{-,y}$ killed in x is

$$((-L_{|(x,y)}^{-,y})^{-1}f)(x') = C \int_x^y u^{+,x}(x' \wedge y') u^{-,y}(x' \vee y') \frac{u^{-,y}(y')}{u^{-,y}(x')} m(y') dy'$$

The potential measure of $(\rho_t^{+,x})_{0 \leq t \leq T_y^{+,x}}$ starting from x is

$$U(x') dx' = C u^{+,x}(x') u^{-,y}(x') m(x') dx'$$

and for any f, g bounded functions on (x, y)

$$(3.5.3) \quad \int_x^y ((-L_{|(x,y)}^{+,x})^{-1}f)(x') g(x') U(x') dx' = \int_x^y f(x') ((-L_{|(x,y)}^{-,y})^{-1}g)(x') U(x') dx'$$

The time reversal property for $(\rho_t^{+,x})_{0 \leq t \leq T_y^{+,x}}$ follows from the duality relation (3.5.3). See [24], chapter VII, §4 for details on time reversal.

Corollary 3.19. — *If L is a generator of a diffusion on I of form (2.2.1), then*

$$(3.5.4) \quad \mu^*(\cdot) = \int_{a \in I} \pi_* \eta^{>a}(\cdot) w(a) da$$

The measure on $(\min \gamma, \max \gamma)$ induced by μ^* is $1_{a < b \in I} \frac{da db}{u^{+,a}(b) u^{-,b}(a)}$. Let $a < b \in I$. Let $(\rho_t^{+,a})_{0 \leq t < \zeta^{+,a}}$ and $(\rho_t^{-,b})_{0 \leq t < \zeta^{-,b}}$ be two independent diffusion, the first of generator $L^{+,a}$ starting from a and the second of generator $L^{-,b}$ starting from b . Let $T_b^{+,a}$ be the first time $\rho^{+,a}$ hits b and $T_a^{-,b}$ the first time $\rho^{-,b}$ hits a . Let $(\beta_t)_{0 \leq t \leq T_b^{+,a} + T_a^{-,b}}$ be the path

$$\beta_t := \begin{cases} \rho_t^{+,a} & \text{if } t \leq T_b^{+,a} \\ \rho_{t-T_b^{+,a}}^{-,b} & \text{if } t \geq T_b^{+,a} \end{cases}$$

Then the law of $(\beta_t)_{0 \leq t \leq T_b^{+,a} + T_a^{-,b}}$ is the probability measure obtained by conditioning the measure μ^* by $(\min \gamma, \max \gamma) = (a, b)$.

Proof. — Both sides of (3.5.4) are covariant by scale and time change. Moreover both sides satisfy the property 3.8 (ii) for the restriction to a subinterval and the property 3.8 (iii) when adding a killing measure. Thus the general case (3.5.4) follows from the Brownian case (3.5.1) by this covariance properties.

If L is a generator without killing measure ($\kappa = 0$) then the description of the measure on $(\min \gamma, \max \gamma)$ and the probabilities obtained after conditioning by the value of $(\min \gamma, \max \gamma)$ follow through a change of scale and time from the analogous description in proposition 3.17. If $\kappa \neq 0$, then we can take u a positive L -harmonic function and deduce the result for L from the result for $Conj(u, L)$ using the fact that $\mu_L^* = \mu_{Conj(u, L)}^*$. \square

The relation between the measure on loops and the excursions measures in dimension 1 (identity (3.5.4)) is analogous to the relation between the measure on Brownian loops and the so called bubble measures observed by Lawler and Werner in dimension 2. See propositions 7 and 8 in [19].

3.6. A generalization of the Vervaat's transformation

In this subsection we will show a conditioned version of the Vervaat's transformation that holds for any one-dimensional diffusion of form (2.2.1) and not just for the Brownian motion. L will be a generator of a diffusion on I of form (2.2.1). From corollary 3.11 and identity (3.5.4) follows that for every $x \in I$:

$$(3.6.1) \quad \int_{t>0} \mathcal{V}_* \mathbb{P}_{x,x}^t(d\gamma) p_t(x, x) dt = \int_{a \in I, a < x} \ell^x(\gamma) \eta^{>a}(d\gamma) w(a) da$$

Let $\mathbb{P}_{x,x}^t(d\gamma | \min \gamma = a)$ be the bridge probability measure condition by the value of the minimum to equal a . Further we will show that there is a version that depends continuously on (a, t) . Let $\eta_t^{>a}$ the probability measure obtained from $\eta^{>a}$ by conditioning the excursion to have a life-time t . The identity (3.6.1) suggests the following:

Proposition 3.20. — *For every $a < x \in I$ and $t > 0$*

$$(3.6.2) \quad \mathcal{V}_* \mathbb{P}_{x,x}^t(d\gamma | \min \gamma = a) = \frac{\ell_t^x(\gamma) \eta_t^{>a}(d\gamma)}{\eta_t^{>a}(\ell_t^x(\gamma))}$$

The distribution of $\min \gamma$ under $\mathbb{P}_{x,x}^t$ equals

$$(3.6.3) \quad w(a) \eta_t^{>a}(\ell_t^x(\gamma)) \frac{1}{p_t(x, x)} \frac{\eta^{>a}(T(\gamma) \in (t, t + dt))}{dt} da$$

where $\frac{\eta^{>a}(T(\gamma) \in (t, t + dt))}{dt}$ is the density of the measure on the life-time of the excursion induced by $\eta^{>a}$. Given an excursion γ following the law $\frac{\ell_t^x(\gamma) \eta_t^{>a}(d\gamma)}{\eta_t^{>a}(\ell_t^x(\gamma))}$, the local time in x is a measure on $\{s \in [0, t] | \gamma(s) = x\}$. The transformation \mathcal{V} sends the starting

point of the bridge to a point $s \in [0, t]$ distributed conditionally on the excursion γ according the measure $\frac{d_s \ell_s^x(\gamma)}{\ell_t^x(\gamma)}$.

Identities (3.6.2) and (3.6.3) can be viewed as a conditioned analogue of the Verwaat's relation between the Brownian bridge and the Brownian excursion. The latter can be deduced from (3.6.2) and (3.6.3) using the translation invariance of the Brownian motion. From (3.6.2) we can only deduce that (3.6.2) and (3.6.3) hold for Lebesgue almost all t and a . We need to show the weak continuity in (a, t) of conditioned bridge probabilities and biased conditioned excursion probabilities to conclude. It is enough to prove the proposition 3.20 for L not containing any killing measure and such that for all $a < x \in I$, a diffusion starting from x reaches a almost surely. Indeed, for a general generator, $Conj(u_\downarrow, L)$ does satisfy the above constraints and if the proposition 3.20 is true for $Conj(u_\downarrow, L)$ then it is also true for L . From now on we assume that L satisfies the above constraints. Next we give a more constructive description of the conditioned bridges and biased conditioned excursions. We start with bridges.

Property 3.3 (viii) shows that the measure $\mathbb{P}_x^{T_a} \triangleleft \tilde{\mathbb{P}}_x^{\tilde{T}^a} \wedge$ conditioned on $T_a + \tilde{T}_a = t$ is a version of $\mathbb{P}_{x,x}^t(d\gamma | \min \gamma = a)$. Let $p_t^{(a \times)}(x, y)$ be the transition density on $I \cap (a, +\infty)$ relatively to $m(y)dy$ of the semi-group generated by $L|_{I \cap (a, +\infty)}$. Then $p_t^{(a \times)}(x, a^+) = 0$. According to [21], for all $t > 0$, $y \mapsto p_t^{(a \times)}(x, y)$ is \mathcal{C}^1 . Let $\partial_2 p_t^{(a \times)}(x, y)$ be the derivative relatively to y . It has a positive limit $\partial_2 p_t^{(a \times)}(x, a^+)$ as $y \rightarrow a^+$. Extended in this way, the map $(t, x, y) \mapsto \partial_2 p_t^{(a \times)}(x, y)$ is continuous on $(0, +\infty) \times I \cap (a, +\infty) \times I \cap [a, +\infty)$. The distribution of T_a under \mathbb{P}_x is (see [12], page 154):

$$\frac{1}{w(a)} \partial_2 p_t^{(a \times)}(x, a^+) dt$$

Let $\mathbb{P}_{x,y}^{(a \times), t}$ be the bridge probability measures of $L|_{I \cap (a, +\infty)}$. It has a weak limit $\mathbb{P}_{x, a^+}^{(a \times), t}$ as $y \rightarrow a^+$. Let \mathcal{F}_s be the sigma-algebra generated by the restriction of a continuous path to the time interval $[0, s]$. Let $\mathbb{P}_a^{+, a}$ be the law of $\rho^{+, a}$ starting from a . For all $s \in (0, t)$ we have the following absolute continuity relations:

$$(3.6.4) \quad \frac{d\mathbb{P}_{x, a^+}^{(a \times), t}}{d\mathbb{P}_x} \Big|_{\mathcal{F}_s} = 1_{s < T_a} \frac{\partial_2 p_{t-s}^{(a \times)}(X_s, a^+)}{\partial_2 p_t^{(a \times)}(x, a^+)}$$

and for the time reversed bridge

$$(3.6.5) \quad \frac{d\mathbb{P}_{x, a^+}^{(a \times), t \wedge}}{d\mathbb{P}_a^{+, a}} \Big|_{\mathcal{F}_s} = \frac{p_{t-s}^{(a \times)}(\rho_s^{+, a}, x)}{\partial_2 p_t^{(a \times)}(x, a^+)}$$

Using the absolute continuity relation (3.6.4) and (3.6.5) one can prove in a similar way as in proposition 2.5 that the map $(t, y) \mapsto \mathbb{P}_{x, a^+}^{(a \times), t}$ is continuous for the weak topology. The first passage bridge $\mathbb{P}_x^{T_a}$ disintegrates as follows

$$(3.6.6) \quad \mathbb{P}_x^{T_a}(\cdot) = \frac{1}{w(a)} \int_{t > 0} \mathbb{P}_{x, a^+}^{(a \times), t}(\cdot) \partial_2 p_t^{(a \times)}(x, a^+) dt$$

From the property 3.3 (viii) and (3.6.6) we get that

Property 3.21. — *The distribution of $\min \gamma$ under $P_{x,x}^t$ is*

$$(3.6.7) \quad \frac{da}{w(a)p_t(x,x)} \int_0^t \partial_2 p_s^{(a^\times)}(x, a^+) \partial_2 p_{t-s}^{(a)}(x, a^+) ds$$

There is a version of $\mathbb{P}_{x,x}^t(d\gamma | \min \gamma = a)$ that disintegrates as

$$(3.6.8) \quad \frac{\int_0^t \left(\mathbb{P}_{x,a^+}^{(a^\times),s} \triangleleft \mathbb{P}_{x,a^+}^{(a^\times),t-s\wedge} \right) (d\gamma) \partial_2 p_s^{(a^\times)}(x, a^+) \partial_2 p_{t-s}^{(a^\times)}(x, a^+) ds}{\int_0^t \partial_2 p_s^{(a^\times)}(x, a^+) \partial_2 p_{t-s}^{(a^\times)}(x, a^+) ds}$$

Next we show that the probability measure given by (3.6.8) depends continuously on (a, t) .

Lemma 3.22. — *The functions $(x, a, t) \mapsto p_t^{(a^\times)}(x, a^+)$ and $(x, a, t) \mapsto \partial_2 p_t^{(a^\times)}(x, a^+)$ are continuous on $\{(x, a) | x > a \in I\} \times (0, +\infty)$.*

Proof. — As in [21], we can use the eigendifferential expansion of L to express $p_t^{(a^\times)}(x, a^+)$ and $\partial_2 p_t^{(a^\times)}(x, a^+)$. Let x_0 . For $\lambda \in \mathbb{R}$ consider $e_1(\cdot, \lambda)$ and $e_2(\cdot, \lambda)$ two solutions to $Lu + \lambda u = 0$ with initial conditions

$$e_1(x_0, \lambda) = 1 \quad \frac{\partial e_1}{\partial x}(x_0, \lambda) = 0 \quad e_2(x_0, \lambda) = 0 \quad \frac{\partial e_2}{\partial x}(x_0, \lambda) = 1$$

Let $\mathbf{e}(x, \lambda)$ be the 2-vector whose entries are $e_1(x, \lambda)$ and $e_2(x, \lambda)$. According to theorems 3.2 and 4.3 in [21], for all $a \in I$ there is a Radon measure $\mathfrak{f}^{(a)}$ on $(-\infty, 0]$ with values in the space of 2×2 symmetric positive semi-definite matrices such that for all $x \in I \cap (a, +\infty)$

$$p_t^{(a^\times)}(x, a^+) = \int_{-\infty}^0 e^{t\lambda\tau} \mathbf{e}(x, \lambda) \mathfrak{f}^{(a^\times)}(d\lambda) \mathbf{e}(a, \lambda)$$

$$\partial_2 p_t^{(a^\times)}(x, a^+) = \int_{-\infty}^0 e^{t\lambda\tau} \mathbf{e}(x, \lambda) \mathfrak{f}^{(a^\times)}(d\lambda) \frac{\partial \mathbf{e}}{\partial x}(a, \lambda)$$

Let $x > a \in I$. Consider a two sequences $(x_n)_{n \geq 0}$ and $(a_n)_{n \geq 0}$ in $I \cap (-\infty, x)$ converging to x respectively a such that for all $n \geq 0$, $x_n > a_n$. Let $(b_j)_{j \geq 0}$ be an increasing sequence in $I \cap (x, \sup I)$ converging to $\sup I$. Let $\mathfrak{f}_{n,j}$ be the 2×2 -matrix valued measure on $(-\infty, 0]$ corresponding to the eigendifferential expansion of L restricted to (a_n, b_j) . $\mathfrak{f}_{n,j}$ charges only a discrete set of atoms. As shown in the proof of theorem 3.2 in [21], the total mass of the measures $1 \wedge |\lambda|^{-2} \|\mathfrak{f}_{n,j}\|(d\lambda)$, $1 \wedge |\lambda|^{-2} \|\mathfrak{f}^{(a_n^\times)}\|(d\lambda)$ and $1 \wedge |\lambda|^{-2} \|\mathfrak{f}^{(a^\times)}\|(d\lambda)$ is uniformly bounded. Moreover for a fixed n , as $j \rightarrow +\infty$, $1 \wedge |\lambda|^{-2} \mathfrak{f}_{n,j}(d\lambda)$ converges vaguely, that is against continuous functions vanishing at infinity, to the measure $1 \wedge |\lambda|^{-2} \mathfrak{f}^{(a_n^\times)}(d\lambda)$. Moreover, for any increasing integer-valued sequence $(j_n)_{n \geq 0}$ converging to $+\infty$, $1 \wedge |\lambda|^{-2} \mathfrak{f}_{n,j_n}(d\lambda)$ converges vaguely as $n \rightarrow +\infty$ to $1 \wedge |\lambda|^{-2} \mathfrak{f}^{(a^\times)}(d\lambda)$. Since the sequence $(j_n)_{n \geq 0}$

is arbitrary, this implies that $1 \wedge |\lambda|^{-2} \mathfrak{f}^{(a_n \times)}(d\lambda)$ converges vaguely as $n \rightarrow +\infty$ to $1 \wedge |\lambda|^{-2} \mathfrak{f}^{(a \times)}(d\lambda)$.

There are constants $C, c' > 0$ such that for all $\lambda \leq 0$ and $n \geq 0$

$$(3.6.9) \quad \|\mathfrak{e}(x_n, \lambda)\| \leq C e^{c' \sqrt{|\lambda|}} \quad \|\mathfrak{e}(a_n, \lambda)\| \leq C e^{c' \sqrt{|\lambda|}} \quad \left\| \frac{\partial \mathfrak{e}}{\partial x}(a_n, \lambda) \right\| \leq C e^{c' \sqrt{|\lambda|}}$$

Let $t > 0$ and $(t_n)_{n \geq 0}$ a sequence of times converging to t . From (3.6.9) follows that

$$\lim_{\lambda \rightarrow -\infty} \sup_{n \geq 0} |\lambda|^2 e^{t_n \lambda} \|\mathfrak{e}(x_n, \lambda)\| \times \|\mathfrak{e}(a_n, \lambda)\| = 0$$

$\lambda \mapsto 1 \vee |\lambda|^2 e^{t_n \lambda} (\mathfrak{e}(x_n, \lambda), \partial \mathfrak{e}(a_n, \lambda))$ vanishes at infinity and converges uniformly on $(-\infty, 0]$ to $\lambda \mapsto 1 \vee |\lambda|^2 e^{t \lambda} (\mathfrak{e}(x, \lambda), \mathfrak{e}(a, \lambda))$. The vague convergence of measures implies that

$$\lim_{n \rightarrow +\infty} \int_{-\infty}^0 e^{t_n \lambda} \mathfrak{e}(x_n, \lambda) \mathfrak{f}^{(a_n \times)}(d\lambda) \mathfrak{e}(a_n, \lambda) = \int_{-\infty}^0 e^{t \lambda} \mathfrak{e}(x, \lambda) \mathfrak{f}^{(a \times)}(d\lambda) \mathfrak{e}(a, \lambda)$$

Similarly $\partial_2 p_{t_n}^{(a_n \times)}(x_n, a_n^+)$ converges to $\partial_2 p_t^{(a \times)}(x, a^+)$. \square

Lemma 3.23. — *The map $a \mapsto \mathbb{P}_a^{+,a}$ is weakly continuous.*

Proof. — Let $a_0 \in I$. Consider the process $(\rho_t^{+,a_0})_{t \geq 0}$ following the law $\mathbb{P}_{a_0}^{+,a_0}$. For $a \in I \cap (a_0, +\infty)$, let \widehat{T}_a be the last time ρ^{+,a_0} visits a . Then $(\rho_{\widehat{T}_a+t}^{+,a_0})_{t \geq 0}$ follows the law $\mathbb{P}_a^{+,a}$. The process valued map $a \mapsto (\rho_{\widehat{T}_a+t}^{+,a_0})_{t \geq 0}$ is almost surely continuous on $I \cap (a_0, +\infty)$ and thus the laws depend weakly continuously on a . \square

Proposition 3.24. — *The version of $\mathbb{P}_{x,x}^t(d\gamma | \min \gamma = a)$ given by (3.6.8) is weakly continuous in (a, t) .*

Proof. — From the absolute continuity relations (3.6.4) for the bridge $\mathbb{P}_{x,a^+}^{(a \times),t}$ and (3.6.5) for its time reversal, together with the continuity of the densities which follows from lemma 3.22, and the weak continuity of $a \mapsto \mathbb{P}_a^{+,a}$, we can deduce in a very similar way as in proposition 2.5 that the map $(a, t) \mapsto \mathbb{P}_{x,a^+}^{(a \times),t}$ is weakly continuous on $(0, +\infty) \times I \cap (-\infty, x)$ and hence $(a, s, t) \mapsto \mathbb{P}_{x,a^+}^{(a \times),s} \triangleleft \mathbb{P}_{x,a^+}^{(a \times),t-s \wedge}$ is weakly continuous. Finally the densities that appear in expression (3.6.8) are continuous with respect to (a, s, t) . \square

Next we will give a decomposition of the measure $\eta^{>a}$ which is similar to the Bismut's decomposition of Brownian excursions (see [24], chapter XII, §4, theorem 4.7). Biane used this Bismut's decomposition to give an alternative proof for the Brownian Vervaat's transformation ([3]). $\partial_2 p_t^{(a \times)}(x, a^+)$ is \mathcal{C}^1 relatively to x and the derivative $\partial_{1,2} p_t^{(a \times)}(x, a^+)$ has a positive limit $\partial_{1,2} p_t^{(a \times)}(a^+, a^+)$ as $y \rightarrow a^+$. Moreover $t \mapsto \partial_{1,2} p_t^{(a \times)}(a^+, a^+)$ is continuous. The measure on the life-time of the excursion induced by $\eta^{>a}$ is (see [25]):

$$\frac{1}{w(a)^2} \partial_{1,2} p_t^{(a \times)}(a^+, a^+) dt$$

Let $s \in [0, t]$. The measure $\eta_t^{>a}(\cdot)$ disintegrates as (see [25]):

$$(3.6.10) \quad \int_{x \in I, x > a} \left(\mathbb{P}_{x, a^+}^{(a \times), s \wedge} \triangleleft \mathbb{P}_{x, a^+}^{(a \times), t-s} \right) (\cdot) \frac{\partial_2 p_s^{(a \times)}(x, a^+) \partial_2 p_{t-s}^{(a \times)}(x, a^+) m(y)}{\partial_{1,2} p_t^{(a \times)}(a^+, a^+)} dy$$

For every $s_1 < s_2 \in [0, s]$, under the bridge measure $\mathbb{P}_{y, z}^{(a \times), s}$

$$(3.6.11) \quad \mathbb{P}_{y, z}^{(a \times), t}(\ell_{s_2}^x(\gamma) - \ell_{s_1}^x(\gamma)) = \int_{s_1}^{s_2} \frac{p_r^{(a \times)}(y, x) p_{s-r}^{(a \times)}(x, z)}{p_s^{(a \times)}(y, z)} dr$$

and under the bridge measure $\mathbb{P}_{y, a^+}^{(a \times), s}$

$$(3.6.12) \quad \mathbb{P}_{y, a^+}^{(a \times), t}(\ell_{s_2}^x(\gamma) - \ell_{s_1}^x(\gamma)) = \int_{s_1}^{s_2} \frac{p_r^{(a \times)}(y, x) \partial_2 p_{s-r}^{(a \times)}(x, a^+)}{\partial_2 p_s^{(a \times)}(y, a^+)} dr$$

Combining (3.6.10) and (3.6.12) we get that for every $s_1 < s_2 \in [0, s]$:

$$(3.6.13) \quad \eta_t^{>a}(\ell_{s_2}^x(\gamma) - \ell_{s_1}^x(\gamma)) = \int_{s_1}^{s_2} \frac{\partial_2 p_s^{(a \times)}(x, a^+) \partial_2 p_{t-s}^{(a \times)}(x, a^+)}{\partial_{1,2} p_t^{(a \times)}(a^+, a^+)} ds$$

Proposition 3.25. — *Let F_1 and F_2 be two non-negative measurable functional on the paths with variable life-time. Then*

$$(3.6.14) \quad \eta_t^{>a} \left(\int_0^t F_1((\gamma(r))_{0 \leq r \leq s}) F_2((\gamma(s+r))_{0 \leq r \leq t-s}) ds \ell_s^x(\gamma) \right) = \int_0^t \mathbb{P}_{x, a^+}^{(a \times), s \wedge} (F_1) \mathbb{P}_{x, a^+}^{(a \times), t-s} (F_2) \frac{\partial_2 p_s^{(a \times)}(x, a^+) \partial_2 p_{t-s}^{(a \times)}(x, a^+)}{\partial_{1,2} p_t^{(a \times)}(a^+, a^+)} ds$$

In particular

$$(3.6.15) \quad \ell_t^x(\gamma) \eta_t^{>a}(d\gamma) = \int_0^t \left(\mathbb{P}_{x, a^+}^{(a \times), s \wedge} \triangleleft \mathbb{P}_{x, a^+}^{(a \times), t-s} \right) (d\gamma) \frac{\partial_2 p_s^{(a \times)}(x, a^+) \partial_2 p_{t-s}^{(a \times)}(x, a^+)}{\partial_{1,2} p_t^{(a \times)}(a^+, a^+)} ds$$

Proof. — It is enough to prove the result in case F_1 and F_2 are non-negative, continuous and bounded. On top of that we may assume that there are $s_{min} < s_{max} \in (0, t)$ such that F_1 respectively F_2 takes value 0 if the life-time of a path is smaller than s_{min} respectively $t - s_{max}$, and that there is $C \in I$, $C > a$, such that F_1 and F_2 take value 0 if $\max \gamma > C$. For $j \leq n \in \mathbb{N}$ set $\Delta s_n := \frac{1}{n}(s_{max} - s_{min})$ and $s_{j,n} := s_{min} + j \Delta s_n$. Then almost surely

$$(3.6.16) \quad \int_0^t F_1((\gamma(r))_{0 \leq r \leq s}) F_2((\gamma(s+r))_{0 \leq r \leq t-s}) ds \ell_s^x(\gamma) = \lim_{n \rightarrow +\infty} \sum_{j=0}^{n-1} F_1((\gamma(r))_{0 \leq r \leq s_{j,n}}) (\ell_{s_{j+1,n}}^x(\gamma) - \ell_{s_{j,n}}^x(\gamma)) F_2((\gamma(s_{j+1,n} + r))_{0 \leq r \leq t-s_{j+1,n}})$$

Moreover the right-hand side of (3.6.16) is dominated by $\ell_t^x(\gamma)\|F_1\|_\infty\|F_2\|_\infty$. Thus the $\eta_t^{>a}$ -expectation converges too. Applying (3.6.10) and (3.6.11) we get

$$\eta_t^{>a}\left(F_1((\gamma(r))_{0\leq r\leq s_{j,n}})(\ell_{s_{j+1,n}}^x(\gamma) - \ell_{s_{j,n}}^x(\gamma))F_2((\gamma(s_{j+1,n} + r))_{0\leq r\leq t-s_{j+1,n}})\right) = \int_0^{\Delta s_n} \int_{(a,C)^2} \mathbb{P}_{y,a^+}^{(a\times),s_{j,n}\wedge}(F_1)\mathbb{P}_{z,a^+}^{(a\times),t-s_{j+1,n}}(F_2)q_n(r,y,z)m(y)dy m(z)dz dr$$

where

$$q_n(r,y,z) = \frac{\partial_2 p_{s_{j,n}}^{(a\times)}(y,a^+)\partial_2 p_{t-s_{j+1,n}}^{(a\times)}(z,a^+)}{\partial_{1,2} p_t^{(a\times)}(a^+,a^+)} p_r^{(a\times)}(y,x) p_{\Delta s_n - r}^{(a\times)}(x,z)$$

The measure $1_{y,z>a\in I} \frac{1}{\Delta s_n} \int_0^{\Delta s_n} q_n(r,y,z) dr dy dz$ converges weakly as $n \rightarrow +\infty$ to $\delta_{(x,x)}$. The maps $(s,y) \mapsto \partial_2 p_s^{(a\times)}(x,a^+)$ and $(s,y) \mapsto \mathbb{P}_s^{(a\times),y,a^+}(\cdot)$ are continuous. Moreover $\partial_2 p_{s_{j,n}}^{(a\times)}(y,a^+)\partial_2 p_{t-s_{j+1,n}}^{(a\times)}(z,a^+)$ is uniformly bounded for $j \leq n \in \mathbb{N}$ and $y,z \in (a,C]$. All this ensures that the $\eta_t^{>a}$ -expectation of the right-hand side of (3.6.16) converges as $n \rightarrow +\infty$ to the right-hand side of (3.6.14). \square

Now we need only to match the preceding descriptions to prove proposition 3.20. (3.6.8) and (3.6.15) imply (3.6.2). (3.6.7) and (3.6.13) imply (3.6.3). The fact that the point where the excursion is split is distributed according to $\frac{d_s \ell_s^x(\gamma)}{\ell_t^x(\gamma)}$ follows from (3.6.14).

3.7. Restricting loops to a discrete subset

Let L be the generator of a diffusion on I of form (2.2.1) and $(X_t)_{0\leq t<\zeta}$ be the corresponding diffusion. Let \mathbb{J} be a countable discrete subset of I . A Markov jump process to the nearest neighbours on \mathbb{J} is naturally embedded in the diffusion X . In this section we will show that, given any $x,y \in \mathbb{J}$, the image of the measure $\mu_L^{x,y}$ through the restriction application that sends a sample paths of the diffusion $(X_t)_{0\leq t<\zeta}$ to a sample path of a Markov jump process on \mathbb{J} is a measure on \mathbb{J} -valued paths that follows the pattern (3.2.2). From this we will deduce that the image of the measure μ_L^* through the restriction to \mathbb{J} is a measure on \mathbb{J} -valued loops following the pattern (3.2.1) and which was studied in [13]. This property will be used in section 4.2 to express the law of finite-dimensional marginals of the occupation field of a Poisson ensemble of intensity $\alpha\mu_L^*$.

For a continuous path $(\gamma(t))_{0\leq t\leq T(\gamma)}$ in I , endowed with continuous local times, let

$$\mathcal{I}_t^{\mathbb{J}}(\gamma) := \sum_{x\in\mathbb{J}} \ell_t^x(\gamma)m(x)$$

For $s \geq 0$, we introduce the stopping time

$$\tau_s^{\mathbb{J}}(\gamma) := \inf\{t \geq 0 | \mathcal{I}_t^{\mathbb{J}}(\gamma) \geq s\}$$

We write $\gamma^{\mathbb{J}}$ for the path $(\gamma(\tau_s^{\mathbb{J}}))_{0 \leq s \leq \mathcal{I}_T^{\mathbb{J}}(\gamma)}$ on \mathbb{J} . Let $m_{\mathbb{J}}$ be the measure

$$m_{\mathbb{J}} := \sum_{x \in \mathbb{J}} m(x) \delta_x$$

The occupation measure of $\gamma^{\mathbb{J}}$ is

$$\sum_{x \in \mathbb{J}} \ell^x(\gamma) m(x) \delta_x$$

and $(\ell^x(\gamma))_{x \in \mathbb{J}}$ are also occupation densities of the restricted path $\gamma^{\mathbb{J}}$ with respect to $m_{\mathbb{J}}$.

The restricted diffusion $X^{\mathbb{J}}$ is a Markov jump process to nearest neighbours on \mathbb{J} , potentially with killing. If $x_0 < x_1$ are two consecutive points in \mathbb{J} , the jump rate from x_0 to x_1 is $\frac{1}{m(x_0)w(x_0)} \frac{1}{u^{+,x_0}(x_1)}$ and the jump rate from x_1 to x_0 is $\frac{1}{m(x_1)w(x_1)} \frac{1}{u^{-,x_1}(x_0)}$. If $x_0 < x_1 < x_2$ are three consecutive points in \mathbb{J} , then the rate of killing while in x_1 is

$$\frac{1}{m(x_1)w(x_1)} \left(\frac{W(u^{-,x_2}, u^{+,x_0})(x_1)}{u^{-,x_2}(x_1)u^{+,x_0}(x_1)} - \frac{1}{u^{-,x_1}(x_0)} - \frac{1}{u^{+,x_1}(x_2)} \right)$$

If \mathbb{J} has a minimum x_0 and x_1 is the second lowest point in \mathbb{J} , then the killing rate while in x_0 is

$$\frac{1}{m(x_0)w(x_0)} \left(\frac{W(u^{-,x_1}, u_{\uparrow})(x_0)}{u^{-,x_1}(x_0)u_{\uparrow}(x_0)} - \frac{1}{u^{+,x_0}(x_1)} \right)$$

An analogous expression holds for the killing rate while in a possible maximum of \mathbb{J} . $X^{\mathbb{J}}$ is transient if and only if X is. Let $L_{\mathbb{J}}$ be the generator of $X^{\mathbb{J}}$. $L_{\mathbb{J}}$ is symmetric relatively to $m_{\mathbb{J}}$. Its Green's function relatively to $m_{\mathbb{J}}$ is $(G(x, y))_{x, y \in \mathbb{J}}$, that is the restriction of the Green's function of L to $\mathbb{J} \times \mathbb{J}$. $X^{\mathbb{J}}$ may not be conservative even if the diffusion X is. In case if \mathbb{J} is not finite, $X^{\mathbb{J}}$ may blow up performing an infinite number of jumps in finite time. Measures $(\mu_L^{x, y})_{x, y \in I}$, μ_L and μ_L^* have discrete space analogues $(\mu_{L_{\mathbb{J}}}^{x, y})_{x, y \in \mathbb{J}}$, $\mu_{L_{\mathbb{J}}}$ and $\mu_{L_{\mathbb{J}}}^*$ as defined in [13], that follow the patterns (3.2.2) and (3.2.1).

Proposition 3.26. — *Let $x, y \in \mathbb{J}$. Then $\gamma \mapsto \gamma^{\mathbb{J}}$ transforms $\mu_L^{x, y}$ in $\mu_{L_{\mathbb{J}}}^{x, y}$ and μ_L^* in $\mu_{L_{\mathbb{J}}}^*$.*

Proof. — The representation (3.2.3) also holds for $\mu_{L_{\mathbb{J}}}^{x, y}$. For $l > 0$, let

$$\tau_l^y := \inf\{t \geq 0 \mid \ell_t^y(X) > l\}$$

and

$$\tau_l^{y, \mathbb{J}} := \inf\{s \geq 0 \mid \ell_s^y(X^{\mathbb{J}}) > l\}$$

Then for any non-negative measurable functional F

$$\mu_{L_{\mathbb{J}}}^{x, y}(F(\gamma)) = \int_0^{+\infty} dl \mathbb{E}_x \left[\mathbf{1}_{\tau_l^{y, \mathbb{J}} < \mathcal{I}_{\zeta}^{\mathbb{J}}} F((X_s^{\mathbb{J}})_{0 \leq s \leq \tau_l^{y, \mathbb{J}}}) \right]$$

But $(X_s^{\mathbb{J}})_{0 \leq s \leq \tau_\lambda^{y,\mathbb{J}}}$ is the image of $(X_t)_{0 \leq t \leq \tau_\lambda^y}$ by the map $\gamma \mapsto \gamma^{\mathbb{J}}$ and $\tau_l^{y,\mathbb{J}} < \mathcal{I}_\zeta^{\mathbb{J}}$ if and only if $\tau_l^y < \zeta$. Thus $\mu_{L_{\mathbb{J}}}^{x,y}$ is the image of $\mu_L^{x,y}$ through the restriction on path to \mathbb{J} . The second part of the proposition can be deduced from that for any $x \in \mathbb{J}$

$$\ell^x(\gamma)\mu_L^*(d\gamma) = \pi_*\mu_L^{x,x}(d\gamma)$$

and as noticed in [13]

$$\ell^x(\gamma)\mu_{L_{\mathbb{J}}}^*(d\gamma^{\mathbb{J}}) = \pi_*\mu_{L_{\mathbb{J}}}^{x,x}(d\gamma^{\mathbb{J}})$$

□

Previous restriction property and the time-change covariance of μ^* (corollary 3.12) can be treated in a unified framework of the time change by the inverse of a continuous additive functional. This is done in [10], section 7.

3.8. Measure on loops in case of creation of mass

We can further extend the definition of the measures $\mu^{x,y}$ on paths and μ and μ^* on loops to the case of L being a "generator" on I containing a creation of mass term as in (2.3.1). Doing so will enable us to emphasize further the h -transform invariance of the measure on loops and will be useful in section 4.2 to compute the exponential moments of the occupation field of Poisson ensembles of Markov loops. Let ν be signed measure on I . Let $L^{(0)} := \frac{1}{m(x)} \frac{d}{dx} \left(\frac{1}{w(x)} \frac{d}{dx} \right)$ and $L := L^{(0)} + \nu$.

Definition 3.27. —

- $\mu_L^{x,y}(d\gamma) := \exp \left(\int_I l^x(\gamma) m(x) \nu(dx) \right) \mu_{L^{(0)}}^{x,y}(d\gamma)$
- $\mu_L(d\gamma) := \exp \left(\int_I l^x(\gamma) m(x) \nu(dx) \right) \mu_{L^{(0)}}(d\gamma)$
- $\mu_L^* := \pi_* \mu_L$

Definition 3.27 is consistent with properties 3.3 (iv) and 3.8 (iii). If $\tilde{\nu}$ is any other signed measure on I , then

$$(3.8.1) \quad \mu_{L+\tilde{\nu}}^{x,y}(d\gamma) := \exp \left(\int_I \ell^x(\gamma) m(x) \tilde{\nu}(dx) \right) \mu_L^{x,y}(d\gamma)$$

Same holds for μ and μ^* . Under the extended definition, the measures $\mu^{x,y}$ still satisfy properties 3.3 (ii), (iii), (v) and (vi). Proposition 3.6 remains true. μ still satisfies properties 3.8 (i), (ii) and (iv). Proposition 3.9 and corollary 3.10 still hold. The identities (3.3.4) and (3.3.8) remain true for μ^* . Concerning the h -transforms, we have:

Proposition 3.28. — *Let h be a continuous positive function on I such that $\frac{d^2 h}{dx^2}$ is a signed measure. $h(x)^2 m(x) dx$ is a speed measure for $\text{Conj}(h, L)$. Then for all $x, y \in I$, $\mu_{\text{Conj}(h, L)}^{x,y} = \frac{1}{h(x)h(y)} \mu_L^{x,y}$, and $\mu_{\text{Conj}(h, L)} = \mu_L$. Conversely, if L and \tilde{L} are two "generators" with or without creation of mass such that $\mu_L = \mu_{\tilde{L}}$ then there is a positive continuous function h on I such that $\frac{d^2 h}{dx^2}$ is a signed measure and $\tilde{L} = \text{Conj}(h, L)$.*

Proof. — There is a positive Radon measure $\tilde{\kappa}$ on I such that both $L - \tilde{\kappa}$ and $\text{Conj}(h, L) - \tilde{\kappa}$ are generators of (killed) diffusions. But

$$\text{Conj}(h, L) - \tilde{\kappa} = \text{Conj}(h, L - \tilde{\kappa})$$

It follows that $\mu_{\text{Conj}(h, L) - \tilde{\kappa}}^{x, y} = \frac{1}{h(x)h(y)}\mu_{L - \tilde{\kappa}}^{x, y}$ and $\mu_{\text{Conj}(h, L) - \tilde{\kappa}} = \mu_{L - \tilde{\kappa}}$. Applying (3.8.1) we get the result.

If $\mu_L = \mu_{\tilde{L}}$, we can again consider $\tilde{\kappa}$ a positive Radon measure on I such that both $L - \tilde{\kappa}$ and $\tilde{L} - \tilde{\kappa}$ are generators of (killed) diffusions. Then according to proposition 3.16, there is a positive continuous function h on I such that $\frac{d^2 h}{dx^2}$ is a signed measure and $\tilde{L} - \tilde{\kappa} = \text{Conj}(h, L - \tilde{\kappa})$. Then $\tilde{L} = \text{Conj}(h, L)$. \square

Similarly to the case of generators of diffusions (section 3.5), one can consider L -harmonic functions $u^{-,x}$ and $u^{+,x}$ in case of L containing creation of mass. If $L \in \mathfrak{D}^+$, then $u^{-,x}$ respectively $u^{+,x}$ is not necessarily positive on $I \cap (-\infty, x)$ respectively $I \cap (x, +\infty)$. Let

$$M(x) := \sup\{y \in I, y \geq x \mid \forall z \in (x, y), u^{+,x}(z) > 0\} \in I \cup \{\sup I\}$$

If $L \in \mathfrak{D}^{0,-}$ then for all $x \in I$, $M(x) = \sup I$. Let $y \in I$, $y > x$. If $y < M(x)$, then $L|_{(x,y)} \in \mathfrak{D}^-$. If $y = M(x)$, then $L|_{(x,y)} \in \mathfrak{D}^0$. If $y > M(x)$, then $L|_{(x,y)} \in \mathfrak{D}^+$. The diffusion $\rho^{+,x}$ of generator $L^{+,x} = \text{Conj}(u^{+,x}, L|_{(x,M(x))}^{+,x})$ is defined on $(x, M(x))$. Similarly for $\rho^{-,y}$. Moreover if $M(x) \in I$, then $L|_{(x,M(x))}^{+,x} = L|_{(x,M(x))}^{-,M(x)}$.

If $L \in \mathfrak{D}^{0,-}$, the description of the measure on $(\min \gamma, \max \gamma)$ induced by μ^* as well as of the probability measures obtained by conditioning μ^* by the value of $(\min \gamma, \max \gamma)$ is the same as given by corollary 3.19, with the same formal expressions. Next we state what happens if $L \in \mathfrak{D}^+$:

Proposition 3.29. — *Let $L \in \mathfrak{D}^+$. The measure on $(\min \gamma, \max \gamma)$ induced by μ^* and restricted to the set $\{a \in I, b \in (a, M(a))\}$ is $1_{a \in I, b \in (a, M(a))} \frac{dadb}{u^{+,a}(b)u^{-,b}(a)}$. If $a < b < M(a)$, then the probability measure obtained through conditioning by $(\min \gamma, \max \gamma) = (a, b)$ has the same description as in corollary 3.19. Outside the set $\{a \in I, b \in (a, M(a))\}$, the measure on $(\min \gamma, \max \gamma)$ is not locally finite. That is to say that, if $a < b \in I$ and $b \geq M(a)$, then for all $\varepsilon > 0$.*

$$(3.8.2) \quad \mu^*(\{\min \gamma \in (a, a + \varepsilon), \max \gamma \in (b - \varepsilon, b)\}) = +\infty$$

Proof. — For the behaviour on $\{a \in I, b \in (a, M(a))\}$: There is a countable collection $(I_j)_{j \geq 0}$ of open subintervals of I such that

$$\{a \in I, b \in (a, M(a))\} = \bigcup_{j \geq 0} \{x < y \in I_j\}$$

Since for all j , $L|_{I_j} \in \mathfrak{D}^{0,-}$, corollary 3.19 applies to $L|_{I_j}$. Combining the descriptions on different $\{a < b \in I_j\}$, we get the description on $\{a \in I, b \in (a, M(a))\}$.

For the behaviour outside $\{a \in I, b \in (a, M(a))\}$: Let $A < B \in \mathbb{R}$. Then

$$(3.8.3) \quad \mu_{BM}^*(\{\min \gamma < A, \max \gamma > B\}) = \int_B^{+\infty} \int_{-\infty}^A \frac{dadb}{(b-a)^2} = +\infty$$

If $a < b \in I$ and $M(a) = b$, then $1_{a < \gamma < b} \mu^*$ is the image of μ_{BM}^* through a change of scale and time. In this case (3.8.2) follows from (3.8.3). If $b > M(a)$, then $L_{|(a,b)} \in \mathfrak{D}^+$. According to proposition 2.9 (iv), there is a positive measure Radon measure κ on (a, b) such that $L_{|(a,b)} - \kappa \in \mathfrak{D}^0$. From what precedes, (3.8.2) holds for $\mu_{L_{|(a,b)} - \kappa}^*$. Moreover, $\mu_{L_{|(a,b)}}^* \geq \mu_{L_{|(a,b)} - \kappa}^*$. So (3.8.2) holds for $\mu_{L_{|(a,b)}}^*$. \square

CHAPTER 4

OCCUPATION FIELDS OF THE POISSON ENSEMBLES OF MARKOV LOOPS

4.1. Inhomogeneous continuous state branching processes with immigration

We will identify the occupation fields of the Poisson ensembles of Markov loops as inhomogeneous continuous state branching processes with immigration. This will be done in section 4.2. In the section 4.1 we will give the basic properties of such processes. In section 4.3 we will deal with the particular case of the intensity being $\frac{1}{2}\mu^*$, in relation with Dynkin's isomorphism.

Let I be an open interval of \mathbb{R} . We will consider stochastic processes where $x \in I$ is the evolution variable. We do not call it time because in the sequel it will rather represent a space variable. Let $(\mathbb{B}_x)_{x \in \mathbb{R}}$ be a standard Brownian motion. Consider the following SDE:

$$(4.1.1) \quad d\tilde{Z}_x = \sigma(x)\sqrt{\tilde{Z}_x}d\mathbb{B}_x + b(x)\tilde{Z}_x dx$$

$$(4.1.2) \quad dZ_x = \sigma(x)\sqrt{Z_x}d\mathbb{B}_x + b(x)Z_x dx + c(x)dx$$

For our needs we will assume that σ is positive and continuous on I , that b and c are only locally bounded and that c is non negative. In this case existence and pathwise uniqueness holds for (4.1.1) and (4.1.2) (see [24], chapter IX, §3), and \tilde{Z} and Z take values in \mathbb{R}_+ . 0 is an absorbing state for \tilde{Z} .

(4.1.1) satisfies the branching property: if $\tilde{Z}^{(1)}$ and $\tilde{Z}^{(2)}$ are two independent processes solutions in law to (4.1.1), defined on $I \cap [x_0, +\infty)$, then $\tilde{Z}^{(1)} + \tilde{Z}^{(2)}$ is a solution in law to (4.1.1). If \tilde{Z} and Z are two independent processes, \tilde{Z} solution in law to (4.1.1) and Z solution in law to (4.1.2), defined on $I \cap [x_0, +\infty)$, then $Z + \tilde{Z}$ is a solution in law to (4.1.2). Solutions to (4.1.2) are (inhomogeneous) continuous state branching processes with immigration. The branching mechanism is given by (4.1.1) and the immigration measure is $c(x)dx$. The homogeneous case (σ , b and c constant) was extensively studied. See [17].

The case of inhomogeneous branching without immigration reduces to the homogeneous case as follows: Let $x_0 \in I$ and let

$$C(x) := \exp\left(-\int_{x_0}^x b(y) dy\right) \quad A(x) := \int_{x_0}^x \sigma(y)^2 C(y)^2 dy$$

If $(\tilde{Z}_x)_{x \in I}$ is a solution to (4.1.1), then $(C(A^{-1}(a))\tilde{Z}_{A^{-1}(a)})_{a \in A(I)}$ is a solution in law to

$$d\tilde{Z}_a = 2\sqrt{\tilde{Z}_a} d\mathbb{B}_a$$

Let \tilde{Z} be a solution to (4.1.1) defined on $I \cap [x_0, +\infty)$, starting at x_0 with the initial condition $\tilde{Z}_{x_0} = z_0 \geq 0$. Then, for $\lambda \geq 0$ and $x \in I$, $x \geq x_0$:

$$\mathbb{E}_{\tilde{Z}_{x_0}=z_0} \left[e^{-\lambda \tilde{Z}_x} \right] = e^{-z_0 \psi(x_0, x, \lambda)}$$

$\psi(x_0, x, \lambda)$ depends continuously on (x_0, x, λ) . If $x = x_0$ then

$$(4.1.3) \quad \psi(x_0, x_0, \lambda) = \lambda$$

If $x_0 \leq x_1 \leq x_2 \in I$ then

$$\psi(x_0, x_2, \lambda) = \psi(x_0, x_1, \psi(x_1, x_2, \lambda))$$

ψ satisfies the differential equation

$$(4.1.4) \quad \frac{\partial \psi}{\partial x_0}(x_0, x, \lambda) = \frac{\sigma(x_0)^2}{2} \psi(x_0, x, \lambda)^2 - b(x_0) \psi(x_0, x, \lambda)$$

If b is not continuous, equation (4.1.4) should be understood in the weak sense. If b is continuous, then (4.1.4) satisfies the Cauchy-Lipschitz conditions, and ψ is uniquely determined by (4.1.4) and the initial condition (4.1.3). This is also the case even if b is not continuous. Indeed, by considering $C(x)\tilde{Z}_x$ rather than \tilde{Z}_x , that is to say considering $\frac{C(x)}{C(x_0)}\psi(x_0, x, \lambda)$ rather than $\psi(x_0, x, \lambda)$, we get rid of b .

Inhomogeneous branching processes are related to the local times of general one-dimensional diffusions:

Proposition 4.1. — *Let $x_0 \in I$ and let $(X_t)_{0 \leq t < \zeta}$ be a diffusion on I of generator L of form (2.2.1) starting from x_0 . Let $z_0 > 0$ and*

$$\tau_{z_0}^{x_0} := \inf\{t \geq 0 \mid \ell_t^{x_0}(X) > z_0\}$$

Then conditionally on $\tau_{z_0}^{x_0} < \zeta$, $(\ell_{\tau_{z_0}^{x_0}}^x(X))_{x \in I, x \geq x_0}$ is a solution in law to the SDE:

$$(4.1.5) \quad d\tilde{Z}_x = \sqrt{2w(x)} \sqrt{\tilde{Z}_x} d\mathbb{B}_x + 2 \frac{d \log u_\downarrow}{dx}(x) \tilde{Z}_x dx$$

Proof. — If X is the Brownian motion on \mathbb{R} , then $w \equiv 2$ and u_\downarrow is constant. In this case the assertion is the second Ray-Knight theorem. See [24], chapter XI, §2. The equation (4.1.5) is then the equation of a square of Bessel 0 process. If $x_{min} < x_0$ and X is the Brownian motion on $(x_{min}, +\infty)$ killed in x_{min} then the law

of $(\ell_{\tau_{z_0}^{x_0}}^x(X))_{x \in I, x \geq x_0}$ conditionally on $\tau_{z_0}^{x_0} < \zeta$ does not depend on x_{min} and is the same as in case of the Brownian motion on \mathbb{R} . Equation (4.1.5) is still satisfied.

If X is a diffusion on I that satisfies that for all $x > a \in I$, starting from x , X reaches almost surely a , which is equivalent to u_\downarrow being constant, then through a change of scale and time X is the Brownian motion on some $(x_{min}, +\infty)$ where $x_{min} \in [-\infty, +\infty)$. Time change does not change the local times because we defined them relatively to the speed measure. Only the change of scale matters. If S is a primitive of w , then conditionally on $\tau_{z_0}^{x_0} < \zeta$, $(\ell_{\tau_{z_0}^{x_0}}^{S^{-1}(2y)}(X))_{y \geq \frac{1}{2}S(x_0)}$ is a square of Bessel 0 process. The equation (4.1.5) follows from the equation of the square of Bessel 0 process by deterministic change of variable $dy := \frac{1}{2}w(x)dx$.

Now the general case: let $(\tilde{X}_t)_{0 \leq t < \tilde{\zeta}}$ be the diffusion of generator $Conj(u_\downarrow, L)$. $\frac{w(x)}{u_\downarrow(x)^2}dx$ is the natural scale measure of \tilde{X} and $u_\downarrow(x)^2m(x)dx$ is its speed measure. We assume that both X and \tilde{X} start from x_0 . The law of \tilde{X} up to the last time it visits x_0 is the same as for X . Let

$$\tilde{\tau} := \inf \left\{ t \geq 0 \mid \ell_t^{x_0}(\tilde{X}) > \frac{1}{u_\downarrow(x_0)^2}z_0 \right\}$$

Then the law of $(\ell_{\tau_{z_0}^{x_0}}^x(X))_{x \in I, x \geq x_0}$ conditionally on $\tau_{z_0}^{x_0} < \zeta$ is the same as the law of $(u_\downarrow(x)^2 \ell_{\tilde{\tau}}^x(\tilde{X}))_{x \in I, x \geq x_0}$ conditionally on $\tilde{\tau} < \tilde{\zeta}$. The factor $u_\downarrow(x)^2$ comes from the fact that performing an h-transform we change the measure relatively to which the local times are defined. For any $a < x_0 \in I$, \tilde{X} reaches a a.s. Thus $(\ell_{\tilde{\tau}}^x(\tilde{X}))_{x \in I, x \geq x_0}$ satisfies the SDE

$$d\tilde{Z}_x = \frac{\sqrt{2w(x)}}{u_\downarrow(x)} \sqrt{\tilde{Z}_x} d\mathbb{B}_x$$

and $(u_\downarrow(x)^2 \ell_{\tau_{z_0}^{x_0}}^x(\tilde{X}))_{x \in I, x \geq x_0}$ satisfies (4.1.5). \square

If there is immigration: Let Z be a solution to (4.1.2) defined on $I \cap [x_0, +\infty)$, starting at x_0 with the initial condition $Z_{x_0} = z_0 \geq 0$. Then, for $\lambda \geq 0$ and $x \in I$, $x \geq x_0$:

$$(4.1.6) \quad \mathbb{E}_{Z_{x_0}=z_0} [e^{-\lambda Z_x}] = \exp \left(-z_0 \psi(x_0, x, \lambda) - \int_{x_0}^x \psi(y, x, \lambda) c(y) dy \right)$$

4.2. Occupation field

Let L be the generator of a diffusion on I of form (2.2.1). Let $\mathcal{L}_{\alpha, L}$ be a Poisson ensemble of intensity $\alpha \mu_L^*$. $\mathcal{L}_{\alpha, L}$ is a random infinite countable collection of unrooted loops supported in I . It is sometimes called "loop soup".

Definition 4.2. — The occupation field of $\mathcal{L}_{\alpha, L}$ is $(\hat{\mathcal{L}}_{\alpha, L}^x)_{x \in I}$ where

$$\hat{\mathcal{L}}_{\alpha, L}^x := \sum_{\gamma \in \mathcal{L}_{\alpha, L}} \ell^x(\gamma)$$

We will drop out the subscript L whenever there is no ambiguity on L . In this subsection we will identify the law of $(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ as an inhomogeneous continuous state branching process with immigration. If \mathbb{J} is a discrete subset of I , then applying proposition 3.26 we deduce that $(\widehat{\mathcal{L}}_\alpha^x)_{x \in \mathbb{J}}$ is the occupation field of the Poisson ensemble of discrete loops of intensity $\alpha \mu_{L_{\mathbb{J}}}^*$ as defined in [13], chapter 4. This fact allows us to apply the results of [13] in order to describe the finite-dimensional marginals of the occupation field. If the diffusion is recurrent, then for all $x \in I$, $\widehat{\mathcal{L}}_\alpha^x = +\infty$ a.s. If the diffusion is transient, then for all $x \in I$, $\widehat{\mathcal{L}}_\alpha^x < +\infty$ a.s. Next we state how does the occupation field behave if we apply various transformations on L .

Property 4.3. — *Let L be the generator of a transient diffusion.*

- (i) *If A is a change of scale function, then*

$$\widehat{\mathcal{L}}_{\alpha, \text{Scale}_A^{\text{gen}} L}^{A(x)} = \widehat{\mathcal{L}}_{\alpha, L}^x$$

- (ii) *If V is a positive continuous function on I , then*

$$\widehat{\mathcal{L}}_{\alpha, \frac{1}{V}}^x = \widehat{\mathcal{L}}_{\alpha, L}^x$$

- (iii) *If h is a positive continuous function on I such that Lh is a negative measure, then*

$$\widehat{\mathcal{L}}_{\alpha, \text{Conj}(h, L)}^x = \frac{1}{h(x)^2} \widehat{\mathcal{L}}_{\alpha, L}^x$$

Previous equalities depend on a particular choice of the speed measure for the modification of L . For (i) we choose $(\frac{dA}{dx} \circ A^{-1})^{-1} m \circ A^{-1} da$. For (ii) we choose $\frac{1}{V(x)} m(x) dx$. For (iii) we choose $h(x)^2 m(x) dx$. The fact that $\widehat{\mathcal{L}}_{\alpha, \text{Conj}(h, L)}^x \neq \widehat{\mathcal{L}}_{\alpha, L}^x$ despite $\mathcal{L}_{\alpha, \text{Conj}(h, L)} = \mathcal{L}_{\alpha, L}$ comes from a change of speed measure.

Next we characterize the finite-dimensional marginals of the occupation field by stating the results that appear in [13], chapter 4.

Property 4.4. — *The distribution of $\widehat{\mathcal{L}}_\alpha^x$ is*

$$\frac{(G_L(x, x))^\alpha}{\Gamma(\alpha)} l^{\alpha-1} \exp\left(-\frac{l}{G_L(x, x)}\right) 1_{l>0} dl$$

Let $x_1, x_2, \dots, x_n \in I$ and $\lambda_1, \lambda_2, \dots, \lambda_n \geq 0$. Then

$$(4.2.1) \quad \mathbb{E} \left[\exp\left(-\sum_{i=1}^n \lambda_i \widehat{\mathcal{L}}_\alpha^{x_i}\right) \right] = \left(\frac{\det(G_L - \sum_{i=1}^n \lambda_i \delta_{x_i})(x_i, x_j)_{1 \leq i, j \leq n}}{\det(G_L(x_i, x_j))_{1 \leq i, j \leq n}} \right)^\alpha$$

The moment $\mathbb{E} \left[\widehat{\mathcal{L}}_\alpha^{x_1} \widehat{\mathcal{L}}_\alpha^{x_2} \dots \widehat{\mathcal{L}}_\alpha^{x_n} \right]$ is an α -permanent:

$$\mathbb{E} \left[\widehat{\mathcal{L}}_\alpha^{x_1} \widehat{\mathcal{L}}_\alpha^{x_2} \dots \widehat{\mathcal{L}}_\alpha^{x_n} \right] = \sum_{\sigma \in \mathfrak{S}_n} \alpha^{\# \text{ cycles of } \sigma} \prod_{i=1}^n G(x_i, x_{\sigma(i)})$$

If \mathbb{J} is a discrete subset of I , then $(\widehat{\mathcal{L}}_\alpha^x)_{x \in \mathbb{J}}$, viewed as a stochastic process that evolves when x increases, is an inhomogeneous continuous state branching process with immigration defined on the discrete set \mathbb{J} . In particular, for any $x_1 \leq x_2 \leq \dots \leq x_n \in I$ and $p \in \{1, 2, \dots, n\}$, $(\widehat{\mathcal{L}}_\alpha^{x_1}, \widehat{\mathcal{L}}_\alpha^{x_2}, \dots, \widehat{\mathcal{L}}_\alpha^{x_p})$ and $(\widehat{\mathcal{L}}_\alpha^{x_p}, \widehat{\mathcal{L}}_\alpha^{x_{p+1}}, \dots, \widehat{\mathcal{L}}_\alpha^{x_n})$ are independent conditionally on $\widehat{\mathcal{L}}_\alpha^{x_p}$.

Next we show that the processes $x \mapsto \widehat{\mathcal{L}}_\alpha^x$ parametrized by $x \in I$, where x is assumed to increase, is an inhomogeneous branching process with immigration of form (4.1.2). In particular, it has a continuous version and is inhomogeneous Markov.

Proposition 4.5. — $(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ has the same finite-dimensional marginals as a solution to the stochastic differential equation

$$(4.2.2) \quad dZ_x = \sqrt{2w(x)}\sqrt{Z_x}d\mathbb{B}_x + 2\frac{d \log u_\downarrow}{dx}(x)Z_x dx + \alpha w(x)dx$$

If L is the generator of a Brownian motion on $(0, +\infty)$ killed when it hits 0, then $(\widehat{\mathcal{L}}_\alpha^x)_{x > 0}$ has the same law as the square of a Bessel process of dimension 2α starting from 0 at $x = 0$. If L is the generator of a Brownian motion on $(0, x_{max})$, killed when hitting the boundary, then $(\widehat{\mathcal{L}}_\alpha^x)_{0 < x < x_{max}}$ has the same law as the square of a Bessel bridge of dimension 2α from 0 at $x = 0$ to 0 at $x = x_{max}$.

Proof. — Let $x_0 < x \in I$ and $\lambda_0, \lambda \geq 0$. Applying the identity (4.2.1) to the case of two points, we get that

$$(4.2.3) \quad \mathbb{E} \left[\exp \left(-\lambda_0 \widehat{\mathcal{L}}_\alpha^{x_0} - \lambda \widehat{\mathcal{L}}_\alpha^x \right) \right] = \left((1 + \lambda_0 G(x_0, x_0))(1 + \lambda G(x, x)) - \lambda_0 \lambda (G(x_0, x))^2 \right)^{-\alpha}$$

Let

$$\Lambda(x_0, \lambda_0) := \mathbb{E} \left[e^{-\lambda_0 \widehat{\mathcal{L}}_\alpha^{x_0}} \right] = \left(\frac{G(x_0, x_0)}{G(x_0, x_0) + \lambda_0} \right)^\alpha$$

For $y \leq x$, let

$$\psi(y, x, \lambda) := \frac{G(x, y)G(y, x)\lambda}{G(y, y)(G(y, y) + \lambda \det_{y,x} G)}$$

$$\varphi(y, x, \lambda) := -\log \left(\frac{G(y, y)}{G(y, y) + \lambda \det_{y,x} G} \right)$$

One can check that the right-hand side of (4.2.3) equals

$$\Lambda(x_0, \lambda_0 + \psi(x_0, x, \lambda)) \exp(-\alpha \varphi(x_0, x, \lambda))$$

In particular for the conditional Laplace transform:

$$(4.2.4) \quad \mathbb{E} \left[\exp \left(-\lambda \widehat{\mathcal{L}}_\alpha^x \right) \middle| \widehat{\mathcal{L}}_\alpha^{x_0} \right] = \exp \left(-\widehat{\mathcal{L}}_\alpha^{x_0} \psi(x_0, x, \lambda) \right) \exp(-\alpha \varphi(x_0, x, \lambda)) \text{ a.s.}$$

Moreover

$$\begin{aligned} \frac{\partial \psi}{\partial y}(y, x, \lambda) &= W(u_\downarrow, u_\uparrow)(y) \psi(y, x, \lambda)^2 - \frac{2}{u_\downarrow(y)} \frac{du_\downarrow}{dy}(y) \psi(y, x, \lambda) \\ &= w(y) \psi(y, x, \lambda)^2 - 2 \frac{d \log u_\downarrow}{dy}(y) \psi(y, x, \lambda) \end{aligned}$$

and

$$\frac{\partial \varphi}{\partial y}(y, x, \lambda) = -W(u_\downarrow, u_\uparrow)(y) \psi(y, x, \lambda) = -w(y) \psi(y, x, \lambda)$$

and we have the initial conditions $\psi(x, x, \lambda) = \lambda$ and $\varphi(x, x, \lambda) = 0$. Thus (4.2.4) has the same form as (4.1.6) where $c(y) = \alpha w(y)$. Let $(Z_y)_{y \in I, y \geq x_0}$ be a solution to (4.2.2) with the initial condition Z_{x_0} being a gamma random variable of parameter α with mean $\alpha G(x_0, x_0)$. It follows from what precedes that $(\widehat{\mathcal{L}}_\alpha^{x_0}, \widehat{\mathcal{L}}_\alpha^x)$ has the same law as (Z_{x_0}, Z_x) . Using the conditional independence satisfied by the occupation field, we deduce that $(\widehat{\mathcal{L}}_\alpha^y)_{y \in I, y \geq x_0}$ has the same finite-dimensional marginals as $(Z_y)_{y \in I, y \geq x_0}$. Making x_0 converge to $\inf I$ along a countable subset, we get a consistent family of continuous stochastic processes, which induces a continuous stochastic process $(Z_y)_{y \in I}$ defined on whole I . It satisfies (4.2.2) and has the same finite-dimensional marginals as $(\widehat{\mathcal{L}}_\alpha^y)_{y \in I}$.

In case of a Brownian motion in $(0, +\infty)$ killed in 0, the equation (4.2.2) becomes

$$dZ_x = 2\sqrt{Z_x} d\mathbb{B}_x + 2\alpha dx$$

which is the SDE satisfied by the square of a Bessel process of dimension 2α . Moreover $(\widehat{\mathcal{L}}_\alpha^x)_{x>0}$ has the same one-dimensional marginals as the latter, more precisely $\widehat{\mathcal{L}}_\alpha^x$ is a gamma r.v. of parameter α with mean $2\alpha x$. This shows the equality in law.

In case of a Brownian motion in $(0, x_{max})$ killed in 0 and x_{max} the equation (4.2.2) becomes

$$dZ_x = 2\sqrt{Z_x} d\mathbb{B}_x + \frac{1}{x_{max} - x} Z_x dx + 2\alpha dx$$

which is the SDE satisfied by the square of a Bessel bridge of dimension 2α from 0 at $x = 0$ to 0 at $x = x_{max}$. Moreover the latter process and $(\widehat{\mathcal{L}}_\alpha^x)_{0 < x < x_{max}}$ have the same one-dimensional marginals, more precisely gamma r.v. of parameter α with mean $2\alpha(x_{max} - x) \frac{x}{x_{max}}$. Thus the two have the same law. \square

We showed that $(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ has the same finite-dimensional marginals as a continuous stochastic process. We will assume in the sequel and prove in section 5.2 that one can couple the Poisson ensemble \mathcal{L}_α and a continuous version of its occupation field $(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ on the same probability space. This does not follow trivially from the fact that the process $(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ has a continuous version. Consider the following counterexample: Let U be a uniform r.v. on $(0, 1)$. Let \mathcal{E} be a countable random set of Brownian excursions defined as follows: conditionally on U \mathcal{E} is a Poisson ensemble

with intensity $\eta_{BM}^{>U} + \eta_{BM}^{<U}$. Let $(\widehat{\mathcal{E}}_x)_{x \in \mathbb{R}}$ be the occupation field of \mathcal{E} . Then $\widehat{\mathcal{E}}$ is continuous on $(-\infty, U)$ and $(U, +\infty)$ but not at U . Indeed $\widehat{\mathcal{E}}_U = 0$ and

$$\lim_{x \rightarrow U^-} \widehat{\mathcal{E}}_x = \lim_{x \rightarrow U^+} \widehat{\mathcal{E}}_x = 1$$

Let $(\widehat{\mathcal{E}}'_x)_{x \in \mathbb{R}}$ be the field defined by: $\widehat{\mathcal{E}}'_x = \widehat{\mathcal{E}}_x$ if $x \neq U$ and $\widehat{\mathcal{E}}'_U = 1$. $(\widehat{\mathcal{E}}'_x)_{x \in \mathbb{R}}$ is continuous and for any fixed $x \in \mathbb{R}$ $\widehat{\mathcal{E}}'_x = \widehat{\mathcal{E}}_x$ a.s. Thus $(\widehat{\mathcal{E}}'_x)_{x \in \mathbb{R}}$ is a continuous version of the process $(\widehat{\mathcal{E}}_x)_{x \in \mathbb{R}}$ but it can not be implemented as a sum of local time across the excursions in \mathcal{E} . As we will show in section 5.2, such a difficulty does not arise in case of \mathcal{L}_α .

$(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ is an inhomogeneous continuous state branching with immigration. The branching mechanism is the same as for the local times of the diffusion X , given by (4.1). The immigration measure is $\alpha w(x)dx$. The interpretation is the following: given a loop in \mathcal{L}_α , its family of local times performs a branching according to the mechanism (4.1), independently from the other loops. The immigration between x and $x + \Delta x$ comes from the loops whose minima belong to $(x, x + \Delta x)$. It is remarkable that although the immigration measure is absolutely continuous with respect to Lebesgue measure, there is only a countable number of moments at which immigration occurs. These are the positions of the minima of loops in \mathcal{L}_α . Moreover the local time of each loop at its minimum is zero. For $x > a \in I$, let

$$\widehat{\mathcal{L}}_\alpha^{(a),x} := \sum_{\substack{\gamma \in \mathcal{L}_\alpha \\ \min \gamma > a}} \ell^x(\gamma)$$

Let $a < b \in I$. For $j \leq n \in \mathbb{N}$, let $\Delta x_n := \frac{1}{n}(b - a)$ and let $x_{j,n} := a + j\Delta x_n$. Then $(\widehat{\mathcal{L}}_\alpha^{(x_{j-1}, x_j)})_{1 \leq j \leq n}$ is a sequence of independent gamma r.v. of parameter α and the mean of $\widehat{\mathcal{L}}_\alpha^{(x_{j-1}, x_j)}$ is $\alpha \left(G(x_j, x_j) - \frac{G(x_{j-1}, x_j)G(x_j, x_{j-1})}{G(x_{j-1}, x_{j-1})} \right)$. For n large

$$G(x_j, x_j) - \frac{G(x_{j-1}, x_j)G(x_j, x_{j-1})}{G(x_{j-1}, x_{j-1})} = w(x_{j-1})\Delta x_n + o(\Delta x_n)$$

and $o(\Delta x_n)$ is uniform in j . Thus

$$\begin{aligned} \lim_{n \rightarrow +\infty} \mathbb{E} \left[\sum_{j=1}^n \widehat{\mathcal{L}}_\alpha^{(x_{j-1}, x_j)} \right] &= \\ \lim_{n \rightarrow +\infty} \alpha \sum_{j=1}^n \left(G(x_j, x_j) - \frac{G(x_{j-1}, x_j)G(x_j, x_{j-1})}{G(x_{j-1}, x_{j-1})} \right) &= \alpha \int_a^b w(x) dx \end{aligned}$$

and

$$\begin{aligned} \lim_{n \rightarrow +\infty} \text{Var} \left(\sum_{j=1}^n \widehat{\mathcal{L}}_{\alpha}^{(x_{j-1}), x_j} \right) &= \\ \lim_{n \rightarrow +\infty} \alpha \sum_{j=1}^n \left(G(x_j, x_j) - \frac{G(x_{j-1}, x_j)G(x_j, x_{j-1})}{G(x_{j-1}, x_{j-1})} \right)^2 &= 0 \end{aligned}$$

It follows that $\sum_{j=1}^n \widehat{\mathcal{L}}_{\alpha}^{(x_{j-1}), x_j}$ converges in probability to $\alpha \int_a^b w(x) dx$. This is consistent with our interpretation of immigration.

Next proposition deals with the zeroes of the occupation field.

Proposition 4.6. — *Let $x_0 \in I$. If $\int_{\inf I}^{x_0} w(x) dx < +\infty$ then*

$$\lim_{x \rightarrow \inf I} \widehat{\mathcal{L}}_{\alpha}^x = 0$$

Analogous result holds if $\int_{x_0}^{\sup I} w(x) dx < +\infty$.

If $\alpha \geq 1$, then the continuous process $(\widehat{\mathcal{L}}_{\alpha}^x)_{x \in I}$ stays almost surely positive on I . If $\alpha < 1$ then $(\widehat{\mathcal{L}}_{\alpha}^x)_{x \in I}$ hits 0 infinitely many times on I .

Proof. — If $\int_{\inf I}^{x_0} w(x) dx < +\infty$, then $L + \kappa$, where κ is the killing measure of L , is also the generator of a transient diffusion. We can couple $(\widehat{\mathcal{L}}_{\alpha, L}^x)_{x \in I}$ and $(\widehat{\mathcal{L}}_{\alpha, L+\kappa}^x)_{x \in I}$ on the same probability space such that a.s. for all $x \in I$, $\widehat{\mathcal{L}}_{\alpha, L}^x \leq \widehat{\mathcal{L}}_{\alpha, L+\kappa}^x$. But according to property 4.3 (i), $(\widehat{\mathcal{L}}_{\alpha, L+\kappa}^x)_{x \in I}$ is just a scale changed square of Bessel process starting from 0 or square of a Bessel bridge from 0 to 0. Thus

$$\lim_{x \rightarrow \inf I} \widehat{\mathcal{L}}_{\alpha, L}^x \leq \lim_{x \rightarrow \inf I} \widehat{\mathcal{L}}_{\alpha, L+\kappa}^x = 0$$

Regarding the number of zeros of $(\widehat{\mathcal{L}}_{\alpha}^x)_{x \in I}$ on I , property 4.3 ensures that it remains unchanged if we apply scale, time changes and h-transforms to L . Since any generator of a transient diffusion is equivalent through latter transformation to the generator of a Brownian motion on $(0, +\infty)$ killed in 0, the result on the number of zeros of $(\widehat{\mathcal{L}}_{\alpha}^x)_{x \in I}$ follows from standard properties of Bessel processes. \square

In [26] respectively [14] are studied the clusters of loops induced by a Poisson ensemble of loops in the setting of planar Brownian motion respectively Markovian jump processes on graphs. In our setting of one dimensional diffusions the description of such clusters is simple and is related to the zeros of the occupation field. We introduce an equivalence relation on the loops of \mathcal{L}_{α} : γ is in the same class as $\tilde{\gamma}$ if there is a chain of loops $\gamma_0, \gamma_1, \dots, \gamma_n$ in \mathcal{L}_{α} such that $\gamma_0 = \gamma$, $\gamma_n = \tilde{\gamma}$ and for all $i \in \{0, 1, \dots, n-1\}$, $\gamma_i([0, T(\gamma_i)]) \cap \gamma_{i+1}([0, T(\gamma_{i+1})]) \neq \emptyset$. A cluster is the union of all $\gamma([0, T(\gamma)])$ where the loops γ belong to the same equivalence class. It is a subinterval of I . By definition clusters corresponding to different equivalence classes are disjoint.

Proposition 4.7. — *Let L be the generator of a transient diffusion on I . If $\alpha \geq 1$, the loops in \mathcal{L}_α form a single cluster: I . If $\alpha \in (0, 1)$, there are infinitely many clusters. These are the maximal open intervals on which $(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ is positive. In case of the Brownian motion on $(0, +\infty)$ killed at 0, the clusters correspond to the jumps of a stable subordinator with index $1 - \alpha$. In case of a general diffusion, by performing a change of scale of derivative $\frac{1}{2} \frac{w}{u_\downarrow}$, we reduce the problem to the previous case. In case of the Brownian motion on $(0, +\infty)$ killed at 0 and with uniform killing κ , the clusters correspond to the jumps of a subordinator with Levy measure $1_{x>0} \frac{e^{2\sqrt{2\kappa}x} dx}{(e^{2\sqrt{2\kappa}x} - 1)^{2-\alpha}}$.*

Proof. — Assume that \mathcal{L}_α and a continuous version of $(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ are defined on the same probability space. Almost surely the following holds

- Given $\gamma \neq \gamma' \in \mathcal{L}_\alpha$, $\min \gamma \neq \max \gamma'$ and $\max \gamma \neq \min \gamma'$.
- For all $\gamma \in \mathcal{L}_\alpha$, $\ell^{\min \gamma}(\gamma) = \ell^{\max \gamma}(\gamma) = 0$ and $\ell^x(\gamma)$ is positive for $x \in (\min \gamma, \max \gamma)$.

Whenever the above two conditions hold it follows deterministically that the clusters are the intervals on which $(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ stays positive. We deduce then the number of clusters from proposition 4.6.

If L is the generator of the Brownian motion on $(0, +\infty)$ killed at 0, then $(\widehat{\mathcal{L}}_\alpha^x)_{x \in I}$ is the square of a Bessel process of dimension 2α and its excursions correspond to the jumps of a stable subordinator with index $1 - \alpha$.

In general a generator L has the same measure on loops as $Conj(u_\downarrow, L)$. A diffusion of generator $Conj(u_\downarrow, L)$ transforms through a change of time and a change of scale of density $\frac{1}{2} \frac{w}{u_\downarrow}$ into a Brownian motion on $(0, +\infty)$ killed at 0. For the clusters, the change of time does not matter.

In case of a Brownian motion on $(0, +\infty)$ killed at 0 and with uniform killing κ , we can take $u_\downarrow(x) = e^{-\sqrt{2\kappa}x}$. The scale function is then

$$S(x) = \int_0^x \frac{dy}{u_\downarrow(y)^2} = \int_0^x e^{2\sqrt{2\kappa}y} dy = \frac{1}{2\sqrt{2\kappa}} (e^{2\sqrt{2\kappa}x} - 1)$$

Let $(Y_t)_{t \geq 0}$ be an $1 - \alpha$ stable subordinator with Levy measure $1_{y>0} y^{-(2-\alpha)} dy$. The clusters of $\mathcal{L}_{\alpha, \frac{1}{2} \frac{d^2}{dx^2} - \kappa}$ correspond to the jumps of the process $(S^{-1}(Y_t))_{t \geq 0}$, which is not a subordinator. We will that nevertheless the latter process the same set of jumps as a subordinator with Levy measure $1_{x>0} \frac{e^{2\sqrt{2\kappa}x} dx}{(e^{2\sqrt{2\kappa}x} - 1)^{2-\alpha}}$. Let $\varepsilon > 0$ and $(Y_{\varepsilon,t})_{t \geq 0}$ be the process obtained from $(Y_t)_{t \geq 0}$ by removing all the jumps of height less than ε . By construction $Y_{\varepsilon,t} \leq Y_t$. $(S^{-1}(Y_{\varepsilon,t}))_{t \geq 0}$ is a Markov process: given the position of $S^{-1}(Y_{\varepsilon,t})$ at time t , the process waits an exponential holding time with inverse of the mean equal to

$$\int_\varepsilon^{+\infty} \frac{dy}{y^{2-\alpha}} = \frac{1}{(1-\alpha)\varepsilon^{1-\alpha}}$$

Once a jump occurs, the jump of Y_ε is distributed according to the probability

$$1_{y>\varepsilon}(1-\alpha)\varepsilon^{1-\alpha}\frac{dy}{y^{2-\alpha}}$$

The distribution of the corresponding jump of $S^{-1}(Y_{\varepsilon,t})$ is obtained by pushing forward the above probability by the map $y \mapsto S^{-1}(y + Y_{\varepsilon,t}) - S^{-1}(Y_{\varepsilon,t})$ which gives

$$\begin{aligned} & 1_{x>S^{-1}(\varepsilon+Y_{\varepsilon,t})-S^{-1}(Y_{\varepsilon,t})}(1-\alpha)\varepsilon^{1-\alpha}\frac{(2\sqrt{2\kappa})^{2-\alpha}e^{2\sqrt{2\kappa}(x+S^{-1}(Y_{\varepsilon,t}))}dx}{(e^{2\sqrt{2\kappa}(x+S^{-1}(Y_{\varepsilon,t}))}-e^{2\sqrt{2\kappa}S^{-1}(Y_{\varepsilon,t})})^{2-\alpha}} \\ &= 1_{x>S^{-1}(\varepsilon+Y_{\varepsilon,t})-S^{-1}(Y_{\varepsilon,t})}(1-\alpha)\varepsilon^{1-\alpha}(2\sqrt{2\kappa})^{2-\alpha}e^{-(1-\alpha)2\sqrt{2\kappa}S^{-1}(Y_{\varepsilon,t})}\frac{e^{2\sqrt{2\kappa}x}dx}{(e^{2\sqrt{2\kappa}x}-1)^{2-\alpha}} \\ &= 1_{x>S^{-1}(\varepsilon+Y_{\varepsilon,t})-S^{-1}(Y_{\varepsilon,t})}(1-\alpha)\varepsilon^{1-\alpha}\frac{(2\sqrt{2\kappa})^{2-\alpha}}{(1+2\sqrt{2\kappa}Y_{\varepsilon,t})^{1-\alpha}}\frac{e^{2\sqrt{2\kappa}x}dx}{(e^{2\sqrt{2\kappa}x}-1)^{2-\alpha}} \end{aligned}$$

Consider now the random time change

$$\tau_\varepsilon(v) := \inf \left\{ t \geq 0 \mid \int_0^t \frac{(2\sqrt{2\kappa})^{2-\alpha}}{(1+2\sqrt{2\kappa}Y_{\varepsilon,s})^{1-\alpha}} ds \geq v \right\}$$

and at the limit as $\varepsilon \rightarrow 0$

$$\tau(v) := \inf \left\{ t \geq 0 \mid \int_0^t \frac{(2\sqrt{2\kappa})^{2-\alpha}}{(1+2\sqrt{2\kappa}Y_{\varepsilon,s})^{1-\alpha}} ds \geq v \right\}$$

For the time-changed process $(S^{-1}(Y_{\varepsilon,\tau_\varepsilon(v)}))_{v \geq 0}$, the rate of jumps of height belonging to $[x, x + dx]$ is

$$\begin{cases} \frac{e^{2\sqrt{2\kappa}x}dx}{(e^{2\sqrt{2\kappa}x}-1)^{2-\alpha}} & \text{if } x > S^{-1}(\varepsilon + Y_{\varepsilon,\tau_\varepsilon(v)}) - S^{-1}(Y_{\varepsilon,\tau_\varepsilon(v)}) \\ 0 & \text{otherwise} \end{cases}$$

Thus, as ε goes to 0, on one hand the process $(S^{-1}(Y_{\varepsilon,\tau_\varepsilon(v)}))_{v \geq 0}$ converges in law to $(S^{-1}(Y_{\tau(v)}))_{v \geq 0}$ and on the other hand it converges in law to a subordinator with Levy measure $1_{x>0}\frac{e^{2\sqrt{2\kappa}x}dx}{(e^{2\sqrt{2\kappa}x}-1)^{2-\alpha}}$. \square

The clusters coalesce when α increases and fragment when α decreases. Some information on the coalescence of clusters delimited by the zeroes of Bessel processes is given in [2], section 3. These clusters can be obtained as a limit of clusters of discrete loops on discrete subsets. In case of a symmetric jump process to the nearest neighbours on $\varepsilon\mathbb{N}$, if $\alpha > 1$, there are finitely many clusters, and if $\alpha \in (0, 1)$, there are infinitely many clusters and these clusters are given by the holding times of a renewal process, which suitably normalized converges in law as $\varepsilon \rightarrow 0^+$ to the inverse of a stable subordinator with index $1 - \alpha$. See remark 3.3 in [14].

We can consider the occupation field $(\widehat{\mathcal{L}}_{\alpha,L}^x)_{x \in I}$ if L is not the generator of a diffusion but contains creation of mass as in (2.3.1). In this setting, if h is a positive

continuous function on I such that $\frac{d^2h}{dx^2}$ is a signed measure, then for all $x \in I$

$$\widehat{\mathcal{L}}_{\alpha, \text{Conj}(h,L)}^x = \frac{1}{h(x)^2} \widehat{\mathcal{L}}_{\alpha,L}^x$$

It follows that if $L \in \mathfrak{D}^-$ then for all $x \in I$, $\widehat{\mathcal{L}}_{\alpha,L}^x < +\infty$ a.s. and if $L \in \mathfrak{D}^0$ then for all $x \in I$, $\widehat{\mathcal{L}}_{\alpha,L}^x = +\infty$ a.s. If $L \in \mathfrak{D}^+$, then according to proposition 2.9 (iv), there is a positive Radon measure $\tilde{\kappa}$ such that $L - \tilde{\kappa} \in \mathfrak{D}^0$. Then for all $x \in I$, $\widehat{\mathcal{L}}_{\alpha,L}^x \geq \widehat{\mathcal{L}}_{\alpha,L-\tilde{\kappa}}^x = +\infty$. If $L \in \mathfrak{D}^-$, then properties 4.3 (i) and (ii) still hold. The description given by the property 4.4 of the finite-dimensional marginals of $(\widehat{\mathcal{L}}_{\alpha}^x)_{x \in I}$ is still true, although the case of creation of mass wasn't considered in [13]. $(\widehat{\mathcal{L}}_{\alpha}^x)_{x \in I}$ still satisfies the SDE (4.2.2).

Proposition 4.8. — *Let $L \in \mathfrak{D}^-$ and $\tilde{\nu}$ a finite signed measure with compact support in I . Then there is equivalence between*

- (i) $\mathbb{E} \left[\exp \left(\int_I \widehat{\mathcal{L}}_{\alpha,L}^x \tilde{\nu}(dx) \right) \right] < +\infty$
- (ii) $L + \tilde{\nu} \in \mathfrak{D}^-$

If $L + \tilde{\nu} \in \mathfrak{D}^-$ then for $s \in [0, 1]$

$$(4.2.5) \quad \mathbb{E} \left[\exp \left(\int_I \widehat{\mathcal{L}}_{\alpha,L}^x \tilde{\nu}(dx) \right) \right] = \exp \left(\alpha \int_0^1 \int_I G_{L+s\tilde{\nu}}(x, x) \tilde{\nu}(dx) ds \right)$$

Proof. — First observe that $\int_I \widehat{\mathcal{L}}_{\alpha,L}^x |\tilde{\nu}|(dx)$ is almost surely finite because $|\tilde{\nu}|$ is finite and has compact support and $(\widehat{\mathcal{L}}_{\alpha,L}^x)_{x \in I}$ is continuous. Also observe that \mathfrak{D}^- is convex. So if $L + \tilde{\nu} \in \mathfrak{D}^-$, then for all $s \in [0, 1]$, $L + s\tilde{\nu} \in \mathfrak{D}^-$.

(i) implies (ii): Let $\mathbb{P}_{\mathcal{L}_{\alpha,L}}$ be the law of $\mathcal{L}_{\alpha,L}$ and $\mathbb{P}_{\mathcal{L}_{\alpha,L+\tilde{\nu}}}$ be the law of $\mathcal{L}_{\alpha,L+\tilde{\nu}}$. There is an absolute continuity relation between the intensity measures:

$$\mu_{L+\tilde{\nu}}(d\gamma) = \exp \left(\int_I \ell^x(\gamma) \right) \mu_L(d\gamma)$$

In case (i) is true $\mathbb{P}_{\mathcal{L}_{\alpha,L+\tilde{\nu}}}$ is absolutely continuous with respect to $\mathbb{P}_{\mathcal{L}_{\alpha,L}}$ and

$$(4.2.6) \quad d\mathbb{P}_{\mathcal{L}_{\alpha,L+\tilde{\nu}}} = \frac{\exp \left(\int_I \widehat{\mathcal{L}}_{\alpha,L}^x \tilde{\nu}(dx) \right)}{\mathbb{E} \left[\exp \left(\int_I \widehat{\mathcal{L}}_{\alpha,L}^x \tilde{\nu}(dx) \right) \right]} d\mathbb{P}_{\mathcal{L}_{\alpha,L}}$$

But this can not be if $L + \tilde{\nu} \notin \mathfrak{D}^-$ because then for any $x \in I$, $\widehat{\mathcal{L}}_{\alpha,L}^x < +\infty$ and $\widehat{\mathcal{L}}_{\alpha,L+\tilde{\nu}}^x = +\infty$. Thus necessarily $L + \tilde{\nu} \in \mathfrak{D}^-$.

(ii) implies (i): We first assume that $\tilde{\nu}$ is a positive measure and $L + \tilde{\nu} \in \mathfrak{D}^-$. Then $\mathbb{P}_{\mathcal{L}_{\alpha,L}}$ is absolutely continuous with respect to $\mathbb{P}_{\mathcal{L}_{\alpha,L+\tilde{\nu}}}$ and

$$d\mathbb{P}_{\mathcal{L}_{\alpha,L}} = \frac{\exp \left(- \int_I \widehat{\mathcal{L}}_{\alpha,L+\tilde{\nu}}^x \tilde{\nu}(dx) \right)}{\mathbb{E} \left[\exp \left(- \int_I \widehat{\mathcal{L}}_{\alpha,L+\tilde{\nu}}^x \tilde{\nu}(dx) \right) \right]} d\mathbb{P}_{\mathcal{L}_{\alpha,L+\tilde{\nu}}}$$

Inverting the above absolute continuity relation, we get that

$$\mathbb{E} \left[\exp \left(\int_I \widehat{\mathcal{L}}_{\alpha, L}^x \tilde{\nu}(dx) \right) \right] = \mathbb{E} \left[\exp \left(- \int_I \widehat{\mathcal{L}}_{\alpha, L + \tilde{\nu}}^x \tilde{\nu}(dx) \right) \right]^{-1} < +\infty$$

If $\tilde{\nu}$ is not positive, let $\tilde{\nu}^+$ and $-\tilde{\nu}^-$ be its positive respectively negative part. Then

$$\begin{aligned} \mathbb{E} \left[\exp \left(\int_I \widehat{\mathcal{L}}_{\alpha, L}^x \tilde{\nu}(dx) \right) \right] &= \mathbb{E} \left[\exp \left(\int_I \widehat{\mathcal{L}}_{\alpha, L - \tilde{\nu}^-}^x \tilde{\nu}^+(dx) \right) \right] \mathbb{E} \left[\exp \left(- \int_I \widehat{\mathcal{L}}_{\alpha, L}^x \tilde{\nu}^-(dx) \right) \right] \\ &= \frac{\mathbb{E} \left[\exp \left(- \int_I \widehat{\mathcal{L}}_{\alpha, L}^x \tilde{\nu}^-(dx) \right) \right]}{\mathbb{E} \left[\exp \left(- \int_I \widehat{\mathcal{L}}_{\alpha, L + \tilde{\nu}}^x \tilde{\nu}^+(dx) \right) \right]} < +\infty \end{aligned}$$

For the expression (4.2.5) of exponential moments:

$$(4.2.7) \quad \frac{d}{ds} \mathbb{E} \left[\exp \left(s \int_I \widehat{\mathcal{L}}_{\alpha, L}^x \tilde{\nu}(dx) \right) \right] = \mathbb{E} \left[\int_I \widehat{\mathcal{L}}_{\alpha, L}^x \tilde{\nu}(dx) \exp \left(s \int_I \widehat{\mathcal{L}}_{\alpha, L}^x \tilde{\nu}(dx) \right) \right]$$

From the absolute continuity relation (4.2.6) follows that the right-hand side of (4.2.7) equals

$$\alpha \int_I G_{L+s\tilde{\nu}}(x, x) \tilde{\nu}(dx) \mathbb{E} \left[\exp \left(s \int_I \widehat{\mathcal{L}}_{\alpha, L}^x \tilde{\nu}(dx) \right) \right]$$

This implies (4.2.5). \square

As in discrete space case, the above exponential moments can be expressed using determinants. On the complex Hilbert space $\mathbb{L}^2(d|\tilde{\nu}|)$ define for $s \in [0, 1]$ the operators

$$(\mathfrak{G}_{s\tilde{\nu}} f)(x) := \int_I G_{L+s\tilde{\nu}}(x, y) f(y) \tilde{\nu}(dy)$$

$$(|\mathfrak{G}_{s\tilde{\nu}}^*| f)(x) := \int_I G_{L+s\tilde{\nu}}(x, y) f(y) |\tilde{\nu}|(dy)$$

The operator $|\mathfrak{G}_{s\tilde{\nu}}^*|$ is self-adjoint, positive semi-definite with continuous kernel function, and according to [27], theorem 2.12, it is trace class. Since trace class operators form a two-sided ideal in the algebra of bounded operators, $\mathfrak{G}_{s\tilde{\nu}}$ is also trace class. Moreover

$$(4.2.8) \quad \text{Tr}(\mathfrak{G}_{s\tilde{\nu}}) = \int_I G_{L+s\tilde{\nu}}(x, x) \tilde{\nu}(dx)$$

The determinant $\det(Id + \mathfrak{G}_{s\tilde{\nu}})$ is well defined as a converging product of its eigenvalues (see [27], chapter 3).

Proposition 4.9. —

$$\exp \left(\alpha \int_0^1 \int_I G_{L+s\tilde{\nu}}(x, x) \tilde{\nu}(dx) ds \right) = (\det(Id + \mathfrak{G}_{\tilde{\nu}}))^\alpha$$

Proof. — $\mathfrak{G}_{\tilde{\nu}}$ has only real eigenvalues. Indeed, let λ be such an eigenvalue and f a non zero eigenfunction for λ . The sign of $\tilde{\nu}$, $\text{sgn}(\tilde{\nu})$, is a $\{-1, +1\}$ -valued function defined $d|\tilde{\nu}|$ almost everywhere.

$$(4.2.9) \quad \int_I (\text{sgn}(\tilde{\nu})\bar{f})(x)|\mathfrak{G}_{\tilde{\nu}}|(\text{sgn}(\tilde{\nu})f)(x)|\tilde{\nu}|(dx) = \lambda \int_I |f|^2(x)\tilde{\nu}(dx)$$

The left-hand side of (4.2.9) is non-negative. If the right-hand side of (4.2.9) is non-zero, then λ is real. If it is zero, consider $f_\varepsilon := f + \varepsilon \text{sgn}(\tilde{\nu})f$. Then

$$\lambda = \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\varepsilon} \left(\int_I (\text{sgn}(\tilde{\nu})\bar{f}_\varepsilon)(x)|\mathfrak{G}_{\tilde{\nu}}|(\text{sgn}(\tilde{\nu})f_\varepsilon)(x)|\tilde{\nu}|(dx) \right) \left(\int_I |f|^2(x)|\tilde{\nu}|(dx) \right)^{-1}$$

and thus λ is real.

The operators $\mathfrak{G}_{s\tilde{\nu}}$ are compact and the characteristic space corresponding to each of their non-zero eigenvalue is of finite dimension. Let $(\lambda_i)_{i \geq 0}$ be the non-increasing sequence of positive eigenvalues of $\mathfrak{G}_{\tilde{\nu}}$. Each eigenvalue λ_i appears as many times as the dimension of its characteristic space $\ker(\mathfrak{G}_{\tilde{\nu}} - \lambda_i Id)^n$ (n large enough). Similarly let $(-\tilde{\lambda}_j)_{j \geq 0}$ be the non-decreasing sequence of the negative eigenvalues of $\mathfrak{G}_{\tilde{\nu}}$. Let $s \in [0, 1]$. According to the resolvent identity (lemma 2.8), the operators $\mathfrak{G}_{\tilde{\nu}}$ and $\mathfrak{G}_{s\tilde{\nu}}$ commute and satisfy the relation

$$(4.2.10) \quad \mathfrak{G}_{\tilde{\nu}}\mathfrak{G}_{s\tilde{\nu}} = \mathfrak{G}_{s\tilde{\nu}}\mathfrak{G}_{\tilde{\nu}} = \frac{1}{1-s}(\mathfrak{G}_{\tilde{\nu}} - \mathfrak{G}_{s\tilde{\nu}})$$

Since $\mathfrak{G}_{\tilde{\nu}}$ and $\mathfrak{G}_{s\tilde{\nu}}$ commute, these operators have common characteristic spaces. From (4.2.10) follows that $(\frac{\lambda_i}{1+(1-s)\lambda_i})_{i \geq 0}$ is a non-increasing sequence of positive eigenvalues of $\mathfrak{G}_{s\tilde{\nu}}$. If $\frac{-1}{1-s}$ is not an eigenvalue of $\mathfrak{G}_{\tilde{\nu}}$, then $(\frac{-\tilde{\lambda}_j}{1-(1-s)\tilde{\lambda}_j})_{j \geq 0}$ is also a sequence of eigenvalues of $\mathfrak{G}_{s\tilde{\nu}}$. But the family of operators $(\mathfrak{G}_{s\tilde{\nu}})_{s \in [0,1]}$ is bounded. Thus none of $\frac{-\tilde{\lambda}_j}{1-(1-s)\tilde{\lambda}_j}$ can blow up when s varies. So it turns out that $\mathfrak{G}_{\tilde{\nu}}$ has no eigenvalues in $(-\infty, -1]$. From (4.2.8) we get

$$\int_I G_{L+s\tilde{\nu}}(x, x)\tilde{\nu}(dx) = \sum_{i \geq 0} \frac{\lambda_i}{1+(1-s)\lambda_i} - \sum_{j \geq 0} \frac{\tilde{\lambda}_j}{1-(1-s)\tilde{\lambda}_j}$$

The above sum is absolutely convergent, uniformly for $s \in [0, 1]$. Integrating over $[0, 1]$ yields

$$\int_0^1 \int_I G_{L+s\tilde{\nu}}(x, x)\tilde{\nu}(dx)ds = \sum_{i \geq 0} \log(1 + \lambda_i) + \sum_{j \geq 0} \log(1 - \tilde{\lambda}_j)$$

This concludes the proof. \square

4.3. Dynkin's isomorphism

In this subsection we recall the equality in law observed in [13] between the occupation field $(\widehat{\mathcal{L}}_{\frac{x}{2}})_{x \in I}$ and the square of a Gaussian Free Field and show how to derive from this particular versions of Dynkin's isomorphism.

Let L be a generator of a transient diffusion on I of form (2.2.1). Let $(\phi_x)_{x \in I}$ be a centred Gaussian process with variance-covariance function:

$$\mathbb{E}[\phi_x \phi_y] = G(x, y)$$

$(\phi_x)_{x \in I}$ is the Gaussian Free Field associated to L . Let \tilde{S} be a primitive of $\frac{w}{u_{\downarrow}}$. Then $\tilde{S}(\sup I) = +\infty$. Moreover $\tilde{S}(\inf I) > -\infty$ because L is the generator of a transient diffusion. $\left(\frac{1}{u_{\downarrow}(\tilde{S}^{-1}(a))} \phi_{\tilde{S}^{-1}(a)}\right)_{a \in \tilde{S}(I)}$ is a standard Brownian motion starting from 0 at $\tilde{S}(\inf I)$. In particular $(\phi_x)_{x \in I}$ is inhomogeneous Markov and has continuous sample paths.

It was shown in [13], chapter 5, that when $\alpha = \frac{1}{2} (\widehat{\mathcal{L}}_{\frac{x}{2}})_{x \in I}$ has the same law as $(\frac{1}{2} \phi_x^2)_{x \in I}$. In case of a Brownian motion on $(0, +\infty)$ killed in 0, $(\widehat{\mathcal{L}}_{\frac{x}{2}})_{x > 0}$ is the square of a standard Brownian motion starting from 0. In case of a Brownian motion on $(0, x_{max})$ killed in 0 and x_{max} , $(\widehat{\mathcal{L}}_{\frac{x}{2}})_{0 < x < x_{max}}$ is the square of a standard Brownian bridge on $[0, x_{max}]$ from 0 to 0. In case of a Brownian motion on \mathbb{R} with constant killing rate κ , $(\widehat{\mathcal{L}}_{\frac{x}{2}})_{x \in \mathbb{R}}$ is the square of a stationary Ornstein–Uhlenbeck process.

The relation between the occupation field of a Poisson ensemble of Markov loops and the square of a Gaussian Fee Field extends the Dynkin's isomorphism which we state below (see [7] and [9]):

Theorem(Dynkin's Isomorphism). — *Let $x_1, x_2, \dots, x_{2n} \in I$. Then for any non-negative measurable functional F on continuous paths on I ,*

$$(4.3.1) \quad \mathbb{E}_{\phi} \left[\prod_{i=1}^{2n} \phi_{x_i} F\left(\left(\frac{1}{2} \phi_x^2\right)_{x \in I}\right) \right] = \sum_{\text{pairings}} \int \mathbb{E}_{\phi} \left[F\left(\left(\frac{1}{2} \phi_x^2 + \sum_{j=1}^n \ell^x(\gamma_j)\right)_{x \in I}\right) \right] \prod_{\text{pairs}} \mu^{y_j, z_j}(d\gamma_j)$$

where \sum_{pairings} means that the n pairs $\{y_j, z_j\}$ are formed with all $2n$ points x_i in all $\frac{(2n)!}{2^n n!}$ possible ways.

Next we will show that in case $x_i = x_{i+n}$, for $i \in \{1, \dots, n\}$, i.e. $\prod_{i=1}^{2n} \phi_{x_i}$ being a product of squares $\prod_{i=1}^n \phi_{x_i}^2$, one can deduce the Dynkin's isomorphism from the relation between the square of the Gaussian Free Field and the occupation field. In [15] and [10] this is only done in case $n = 1$ and $x_1 = x_2$ using the Palm's identity for Poissonian ensembles and the analogue of the relation (3.3.4). To generalize for

any n we will use an extended version of Palm's identity and the absolute continuity relation given by proposition 3.4.1 (ii).

Lemma 4.10. — *Let \mathcal{E} be an abstract Polish space. Let $\mathfrak{M}(\mathcal{E})$ be the space of locally finite measures on \mathcal{E} and let $\mathcal{M} \in \mathfrak{M}(\mathcal{E})$. Let Φ be a Poisson random measure of intensity \mathcal{M} . Let H be a positive measurable function on $\mathfrak{M}(\mathcal{E}) \times \mathcal{E}^n$. Let \mathfrak{P}_n be the set of partitions of $\{1, \dots, n\}$. If $\mathcal{P} \in \mathfrak{P}_n$ and $i \in \{1, \dots, n\}$, then $\mathcal{P}(i)$ will be the equivalence class of i under \mathcal{P} . The following identity holds:*

$$(4.3.2) \quad \mathbb{E} \left[\int_{\mathcal{E}^n} H(\Phi, q_1, \dots, q_n) \prod_{i=1}^n \Phi(dq_i) \right] = \sum_{\mathcal{P} \in \mathfrak{P}_n} \int_{\mathcal{E}^{\#\mathcal{P}}} \mathbb{E} \left[H(\Phi + \sum_{c \in \mathcal{P}} \delta_{q_c}, q_{\mathcal{P}(1)}, \dots, q_{\mathcal{P}(n)}) \right] \prod_{c \in \mathcal{P}} \mathcal{M}(dq_c)$$

Proof. — We will make a recurrence over n . If $n = 1$, (4.3.2) is the Palm's identity for Poisson random measures. Assume that $n \geq 2$ and that (4.3.2) holds for $n - 1$. We set

$$\tilde{H}(\Phi, q_1, \dots, q_{n-1}) := \int_{\mathcal{E}} H(\Phi, q_1, \dots, q_{n-1}, q_n) \Phi(dq_n)$$

Then

$$(4.3.3) \quad \begin{aligned} \mathbb{E} \left[\int_{\mathcal{E}^n} H(\Phi, q_1, \dots, q_{n-1}, q_n) \prod_{i=1}^n \Phi(dq_i) \right] &= \mathbb{E} \left[\int_{\mathcal{E}^{n-1}} \tilde{H}(\Phi, q_1, \dots, q_{n-1}) \prod_{i=1}^{n-1} \Phi(dq_i) \right] \\ &= \sum_{\mathcal{P}' \in \mathfrak{P}_{n-1}} \int_{\mathcal{E}^{\#\mathcal{P}'}} \mathbb{E} \left[\int_{\mathcal{E}} H(\Phi + \sum_{c' \in \mathcal{P}'} \delta_{q_{c'}}, q_{\mathcal{P}'(1)}, \dots, q_{\mathcal{P}'(n-1)}, q_n) \right. \\ &\quad \left. \times (\Phi(dq_n) + \sum_{c' \in \mathcal{P}'} \delta_{q_{c'}}(dq_n)) \right] \prod_{c' \in \mathcal{P}'} \mathcal{M}(dq_{c'}) \end{aligned}$$

Given a partition $\mathcal{P}' \in \mathfrak{P}_{n-1}$, one can extend it to a partition of $\{1, \dots, n-1, n\}$ either by deciding that n is single in its equivalence class or by choosing an equivalence class $c' \in \mathcal{P}'$ and adjoining n to it. In the identity (4.3.3) the first case corresponds to the integration with respect to $\Phi(dq_n)$, and according to Palm's identity

$$\begin{aligned} \mathbb{E} \left[\int_{\mathcal{E}} H(\Phi + \sum_{c' \in \mathcal{P}'} \delta_{q_{c'}}, q_{\mathcal{P}'(1)}, \dots, q_{\mathcal{P}'(n-1)}, q_n) \Phi(dq_n) \right] &= \\ \int_{\mathcal{E}} \mathbb{E} \left[H(\Phi + \sum_{c' \in \mathcal{P}'} \delta_{q_{c'}}, q_{\mathcal{P}'(1)}, \dots, q_{\mathcal{P}'(n-1)}, q_n) \right] \mathcal{M}(dq_n) \end{aligned}$$

The second case corresponds to the integration with respect to $\delta_{q_{c'}}(dq_n)$. Thus the right-hand side of (4.3.3) equals the right-hand side of (4.3.2). \square

Next we show how derive a particular case of Dynkin's isomorphism using the above extended Palm's formula. Since $(\widehat{\mathcal{L}}_{\frac{x}{2}})_{x \in I}$ and $(\frac{1}{2}\phi_x^2)_{x \in I}$ are equal in law:

$$\mathbb{E}_\phi \left[\prod_{i=1}^n \phi_{x_i}^2 F((\frac{1}{2}\phi_x^2)_{x \in I}) \right] = 2^n \mathbb{E}_{\mathcal{L}_{\frac{1}{2}}} \left[\prod_{i=1}^n \widehat{\mathcal{L}}_{\frac{x_i}{2}} F((\widehat{\mathcal{L}}_{\frac{x}{2}})_{x \in I}) \right]$$

Applying lemma 4.10 we get that

$$\mathbb{E}_{\mathcal{L}_{\frac{1}{2}}} \left[\prod_{i=1}^n \widehat{\mathcal{L}}_{\frac{x_i}{2}} F((\widehat{\mathcal{L}}_{\frac{x}{2}})_{x \in I}) \right] = \sum_{\mathcal{P} \in \mathfrak{P}_n} \int \prod_{i=1}^n \ell^{x_i}(\gamma_{\mathcal{P}(i)}) \mathbb{E} \left[F((\widehat{\mathcal{L}}_{\frac{x}{2}} + \sum_{c \in \mathcal{P}} \ell^x(\gamma_c))_{x \in I}) \right] \prod_{c \in \mathcal{P}} \frac{1}{2} \mu^*(d\gamma_c)$$

Let $\mathfrak{S}_n(\mathcal{P})$ be all the permutations σ of $\{1, \dots, n\}$ such that the classes of the partition \mathcal{P} are the supports of the disjoint cycles of σ . Given a class $c \in \mathcal{P}$, let j_c be its smallest element. From property 3.14 (ii) follows that

$$\prod_{i=1}^n \ell^{x_i}(\gamma_{\mathcal{P}(i)}) = \sum_{\sigma \in \mathfrak{S}_n(\mathcal{P})} \prod_{c \in \mathcal{P}} \ell^{*x_{j_c}, x_{\sigma(j_c)}, \dots, x_{\sigma^{|c|}(j_c)}}(\gamma_c)$$

Proposition 3.15 (ii) states that

$$\ell^{*x_{j_c}, x_{\sigma(j_c)}, \dots, x_{\sigma^{|c|}(j_c)}}(\gamma_c) \mu^*(d\gamma_c) = \pi_* (\mu^{j_c, \sigma(j_c)}(d\tilde{\gamma}_{j_c}) \triangleleft \dots \triangleleft \mu^{\sigma^{|c|-1}(j_c), \sigma^{|c|}(j_c)}(d\tilde{\gamma}_{\sigma^{|c|-1}(j_c)}) \triangleleft \mu^{\sigma^{|c|}(j_c), j_c}(d\tilde{\gamma}_{\sigma^{|c|}(j_c)}))$$

and if the loop γ_c is a concatenation of paths $\tilde{\gamma}_{j_c}, \dots, \tilde{\gamma}_{\sigma^{|c|-1}(j_c)}, \tilde{\gamma}_{\sigma^{|c|}(j_c)}$ then

$$\ell^x(\gamma_c) = \ell^x(\tilde{\gamma}_{j_c}) + \dots + \ell^x(\tilde{\gamma}_{\sigma^{|c|-1}(j_c)}) + \ell^x(\tilde{\gamma}_{\sigma^{|c|}(j_c)})$$

It follows that

$$(4.3.4) \quad 2^n \mathbb{E}_{\mathcal{L}_{\frac{1}{2}}} \left[\prod_{i=1}^n \widehat{\mathcal{L}}_{\frac{x_i}{2}} F((\widehat{\mathcal{L}}_{\frac{x}{2}})_{x \in I}) \right] = \sum_{\sigma \in \mathfrak{S}_n} 2^{n - \#cycles \text{ of } \sigma} \int \mathbb{E}_{\mathcal{L}_{\frac{1}{2}}} \left[F((\widehat{\mathcal{L}}_{\frac{x}{2}} + \sum_{i=1}^n \ell^x(\tilde{\gamma}_i))_{x \in I}) \right] \prod_{i=1}^n \mu^{i, \sigma(i)}(d\tilde{\gamma}_i)$$

But the right-hand side of (4.3.4) is just the same as the right-hand side of (4.3.1) in the specific case when for all $i \in \{1, \dots, n\}$, $x_{i+n} = x_i$. This finishes the derivation of the special case of Dynkin's isomorphism.

CHAPTER 5

DECOMPOSING PATHS INTO POISSON ENSEMBLES OF LOOPS

5.1. Glueing together excursions ordered by their minima

Let L be the generator of a diffusion on I of form (2.2.1). A loop of $\mathcal{L}_{\alpha,L}$ rooted at its minimal point is a positive excursion. For a given $x_0 \in I$, we will consider the loops $\gamma \in \mathcal{L}_{\alpha,L}$ such that $\min \gamma \in (\inf I, x_0]$. We will root these loops at their minima and then order the obtained excursions in the decreasing sense of their minima. Then we will glue all these excursions together and obtain a continuous paths $\xi_{\alpha,L}^{(x_0)}$. The law of this path can be described as a one-dimensional projection of a two-dimensional Markov process. Moreover this path contains all the information on the ensemble of loops $\mathcal{L}_{\alpha,L} \cap \{\gamma \in \mathfrak{L}^* \mid \min \gamma < x_0\}$. So this is a way to sample the latter ensemble of loops. In the particular case of $\alpha = 1$, $\xi_{1,L}^{(x_0)}$ is the sample paths of a one-dimensional diffusion. This is analogue of the link between \mathcal{L}_1 and the loop-erasure procedure already observed in [19] and in [13], chapter 8 and will be described in detail in section 5.3 In the section 5.1 we will consider generalities about glueing together excursions ordered by their minima and probability laws won't be involved. In the section 5.2 we will deal with $\xi_{\alpha,L}^{(x_0)}$ and identify its law. In the section 5.3 we will focus on the case $\alpha = 1$ and describe other ways of slicing sample paths of diffusions into Poisson ensembles of loops.

Let $x_0 \in \mathbb{R}$ and let \mathcal{Q} be a countable everywhere dense subset of $(-\infty, x_0)$. We consider a deterministic collection of excursions $(\mathbf{e}_q)_{q \in \mathcal{Q}}$ where $(\mathbf{e}_q(t))_{0 \leq t \leq T(\mathbf{e}_q)}$ is a continuous excursion above 0, $T(\mathbf{e}_q) > 0$ and

$$\begin{aligned} \mathbf{e}_q(0) &= \mathbf{e}_q(T(\mathbf{e}_q)) = 0 \\ \forall t \in (0, T(\mathbf{e}_q)), \quad \mathbf{e}_q(t) &> 0 \end{aligned}$$

We also assume that for all $C > 0$ and $a < x_0$, there are only finitely many $q \in \mathcal{Q} \cap (a, x_0)$ such that $\max \mathbf{e}_q > C$ and that for all $a < x_0$

$$(5.1.1) \quad \sum_{q \in \mathcal{Q} \cap (a, x_0)} T(\mathbf{e}_q) < +\infty$$

Let $\mathcal{T}(y)$ be the function defined on $[0, +\infty)$ by

$$\mathcal{T}(y) := \sum_{q \in \mathcal{Q} \cap (x_0 - y, x_0)} T(\mathbf{e}_q)$$

\mathcal{T} is a non-decreasing function. Since \mathcal{Q} is everywhere dense, \mathcal{T} is increasing. \mathcal{T} is right-continuous and jumps when $x_0 - y \in \mathcal{Q}$. The height of the jump is then $T(\mathbf{e}_{-y})$.

Let $T_{max} := \mathcal{T}(+\infty) \in (0, +\infty]$. For $t \in [0, T_{max})$ we define

$$\theta(t) := x_0 - \sup\{y \in [0, +\infty) | \mathcal{T}(y) > t\}$$

θ is a non-increasing function from $[0, T_{max})$ to $(-\infty, x_0]$. Since \mathcal{T} is increasing, θ is continuous. We define

$$b^-(t) = \inf\{s \in [0, T_{max}) | \theta(s) = \theta(t)\}$$

$$b^+(t) = \sup\{s \in [0, T_{max}) | \theta(s) = \theta(t)\}$$

$b^-(t) < b^+(t)$ if and only if $\theta(t) \in \mathcal{Q}$ and then $b^+(t) - b^-(t) = T(\mathbf{e}_{\theta(t)})$. We introduce the set

$$\mathfrak{b}^- := \{t \in [0, T_{max}) | \theta(t) \in \mathcal{Q}, b^-(t) = \theta(t)\}$$

\mathfrak{b}^- is in one to one correspondence with \mathcal{Q} by $t \mapsto \theta(t)$.

Finally we define on $[0, T_{max})$ the function ξ :

$$\xi(t) := \begin{cases} \theta(t) & \text{if } \theta(t) \notin \mathcal{Q} \\ \theta(t) + \mathbf{e}_{\theta(t)}(t - b^-(t)) & \text{if } \theta(t) \in \mathcal{Q} \end{cases}$$

Intuitively ξ is the function obtained by gluing together the excursions $(q + \mathbf{e}_q)_{q \in \mathcal{Q}}$ ordered in decreasing sense of their minima. See figure 1 for an example of ξ and θ .

Proposition 5.1. — ξ is continuous. For all $t \in [0, T_{max})$

$$(5.1.2) \quad \theta(t) = \inf_{[0, t]} \xi$$

The set \mathfrak{b}^- can be recovered from ξ as follows:

$$(5.1.3) \quad \mathfrak{b}^- = \{t \in [0, T_{max}) | \xi(t) = \inf_{[0, t]} \xi \text{ and } \exists \varepsilon > 0, \forall s \in (0, \varepsilon), \xi(t+s) > \xi(t)\}$$

If $t_0 \in \mathfrak{b}^-$ then

$$(5.1.4) \quad b^+(t_0) = \inf\{t \in [t_0, T_{max}) | \xi(t) < \xi(t_0)\}$$

Proof. — Let $t \in [0, T_{max})$. To prove the continuity of ξ at t we distinguish three case: the first case is when $\theta(t) \in \mathcal{Q}$ and $b^-(t) < t < b^+(t)$, the second case is when $\theta(t) \notin \mathcal{Q}$ and the third case is when $\theta(t) \in \mathcal{Q}$ and either $b^-(t) = t$ or $b^+(t) = t$.

In the first case, for all $s \in (b^-(t), b^+(t))$,

$$\xi(s) = \theta(t) + \mathbf{e}_{\theta(t)}(s - b^-(t))$$

$\mathbf{e}_{\theta(t)}$ being continuous, we get the continuity of ξ at t .

In the second case we consider a sequence $(t_n)_{n \geq 0}$ in $[0, T_{max})$ converging to t . Let $C > 0$. There are only finitely many $q \in \mathcal{Q}$ such that there is $n \geq 0$ such that $\theta(t_n) = q$

and $\max \mathbf{e}_q > C$. Moreover for any $q \in \mathcal{Q}$, there are only finitely many $n \geq 0$ such that $\theta(t_n) = q$. Thus there are only finitely many $n \geq 0$ such that $\theta(t_n) \in \mathcal{Q}$ and $\max \mathbf{e}_{\theta(t_n)} > C$. So for n large enough

$$(5.1.5) \quad \theta(t_n) \leq \xi(t_n) \leq \theta(t_n) + C$$

But $\xi(t) = \theta(t)$ and $\theta(t_n)$ converges to $\theta(t)$. Since we may take C arbitrarily small, (5.1.5) implies that $\xi(t_n)$ converges to $\theta(t)$.

Regarding the third case, assume for instance that $\theta(t) \in \mathcal{Q}$ and $t = b^-(t)$. The right-continuity of ξ at t follows from the same argument as in the first case and left-continuity from the same argument as in the second case.

By definition, for all $t \in [0, T_{max})$, $\theta(t) \leq \xi(t)$. θ being non-increasing, for all $t \in [0, T_{max})$

$$\theta(t) \leq \inf_{[0,t]} \xi$$

For the converse inequality, we have

$$\theta(t) = \xi(b^-(t)) \geq \inf_{[0,t]} \xi$$

Regarding (5.1.3) and (5.1.4) we have the following disjunction: if $\theta(t) \in \mathcal{Q}$ and $b^-(t) < t < b^+(t)$ then $\xi(t) > \theta(t)$. If $\theta(t) \in \mathcal{Q}$ and $t = b^-(t)$ then for all $s \in (0, b^+(t) - b^-(t))$, $\xi(t+s) > \xi(t)$. If either $\theta(t) \in \mathcal{Q}$ and $t = b^+(t)$ or $\theta(t) \notin \mathcal{Q}$ then $\xi(t) = \theta(t)$ and there is a positive sequence $(s_n)_{n \geq 0}$ decreasing to 0 such that $\theta(t+s_n) \notin \mathcal{Q}$ and $\xi(t+s_n) = \theta(t+s_n) < \theta(t)$. \square




Fig. 1 - Drawing of ξ (full line) and θ (dashed line).

Previous proposition shows that one can reconstruct \mathcal{Q} and the family of excursions $(\mathbf{e}_q)_{q \in \mathcal{Q}}$ only knowing ξ . (5.1.2) shows how to recover θ from ξ . (5.1.3) and (5.1.4) show how to recover the left and the right time boundaries of the excursions of ξ above θ . Also observe that the set defined by the right-hand side of (5.1.3) is countable whatever the continuous function ξ is, even if it is not obtained by glueing together excursions.

5.2. Loops represented as excursions and glued together

Let $\alpha > 0$ and $\mathcal{L}_{\alpha, BM}$ the Poisson ensemble of loops of intensity $\alpha \mu_{BM}^*$ where μ_{BM}^* is the measure on loops associated to the Brownian motion on \mathbb{R} . Let $x_0 \in \mathbb{R}$. We

consider the random countable set \mathcal{Q} :

$$\mathcal{Q} := \{\min \gamma \mid \gamma \in \mathcal{L}_{\alpha, BM}\} \cap (-\infty, x_0)$$

Almost surely \mathcal{Q} is everywhere dense in $(-\infty, x_0)$ and for every $q \in \mathcal{Q}$ there is only one $\gamma \in \mathcal{L}_{\alpha, BM}$ such that $\min \gamma = q$. Almost surely $\gamma \in \mathcal{L}_{\alpha, BM}$ reaches its minimum at one single moment. Given $q \in \mathcal{Q}$ and $\gamma \in \mathcal{L}_{\alpha, BM}$ such that $\min \gamma = q$ we consider \mathbf{e}_q to be the excursion above 0 equal to $\gamma - q$ where we root the unrooted loop γ at $\operatorname{argmin} \gamma$. Then the random set of excursions $(\mathbf{e}_q)_{q \in \mathcal{Q}}$ almost surely satisfies the assumptions of the section 5.1. In particular the condition (5.1.1) follows from the fact that, according to (3.5.1)

$$\int_{\mathcal{Q}^*} 1 \wedge T(\gamma) 1_{\min \gamma \in (a, x_0)} \mu_{BM}^*(d\gamma) = (x_0 - a) \int_0^{+\infty} \frac{t \wedge 1}{\sqrt{2\pi t^3}} dt < +\infty$$

Thus we can consider the random continuous function $(\xi_{\alpha, BM}(t)^{(x_0)})_{t \geq 0}$ constructed by glueing together the excursions $(q + \mathbf{e}_q)_{q \in \mathcal{Q}}$ in the way described in section 5.1. Let

$$\begin{aligned} \theta_{\alpha, BM}^{(x_0)}(t) &= \inf_{[0, t]} \xi_{\alpha, BM}^{(x_0)} \\ \Xi_{\alpha, BM}^{(x_0)}(t) &:= (\xi_{\alpha, BM}^{(x_0)}(t), \theta_{\alpha, BM}^{(x_0)}(t)) \end{aligned}$$

Next we will describe the law of the two-dimensional process $(\Xi_{\alpha, BM}^{(x_0)}(t))_{t \geq 0}$.

Proposition 5.2. — *Let $(\tilde{B}_t)_{t \geq 0}$ be a standard Brownian motion on \mathbb{R} starting from 0. $(\Xi_{\alpha, BM}^{(x_0)}(t))_{t \geq 0}$ has the same law as*

$$\left(x_0 + |\tilde{B}_t| - \frac{1}{\alpha} \ell_t^0(\tilde{B}), x_0 - \frac{1}{\alpha} \ell_t^0(\tilde{B}) \right)_{t \geq 0}$$

In particular for $\alpha = 1$, $(\xi_{1, BM}^{(x_0)}(t))_{t \leq 0}$ has the same law as a Brownian motion starting from x_0 .

Proof. — For $a < x_0$ let T_a be the first time $\theta_{\alpha, BM}^{(x_0)}$ hits a . For $l > 0$ let

$$\tilde{\tau}_l^0 := \inf\{t > 0 \mid \ell_t^0(\tilde{B}) > l\}$$

According to the disintegration (3.5.1) of the measure μ_{BM}^* in the proposition 3.17, for all $a < x_0$ the family $(\mathbf{e}_q)_{q \in \mathcal{Q} \cap (a, x_0)}$ of excursions above 0 is a Poisson point process of intensity $2\alpha\eta_{BM}^{>0}$. This implies the following equality in law

$$(\xi_{\alpha, BM}^{(x_0)}(t) - \theta_{\alpha, BM}^{(x_0)}(t))_{0 \leq t \leq T_a} \stackrel{(law)}{=} (|\tilde{B}_t|)_{0 \leq t \leq \tilde{\tau}_{\alpha(x_0 - a)}^0}$$

Since the above holds for all $a < x_0$, we have the following equality in law

$$(\xi_{\alpha, BM}^{(x_0)}(t) - \theta_{\alpha, BM}^{(x_0)}(t), \alpha(x_0 - \theta_{\alpha, BM}^{(x_0)}(t)))_{t \geq 0} \stackrel{(law)}{=} (|\tilde{B}_t|, \ell_t^0(\tilde{B}))_{t \geq 0}$$

which is exactly the equality in law we needed. Finally for $\alpha = 1$, $(x_0 + |\tilde{B}_t| - \ell_t^0(\tilde{B}))_{t \geq 0}$ has the law of a Brownian motion starting from x_0 . See [24], chapter VI, §2. \square

According to proposition 5.2 a Brownian sample path can be decomposed into a Poisson process of positive excursion with decreasing minima. This decomposition is for instance described in [16], section 6.2.D. In case $\alpha = 1$, proposition 4.5 states that the occupation field of a the Poisson ensemble of loops associated to the Brownian motion on $(0, +\infty)$ killed at 0 is the square of a Bessel process of dimension 2 starting from 0 at 0. This result can also be obtained using the fact that $(\xi_{1, BM}^{(x_0)}(t))_{t \leq 0}$ is a Brownian sample path and applying the first Ray-Knight theorem which gives the law of the occupation field of a Brownian path stopped upon hitting 0.

From proposition 5.2 follows in particular that $(\Xi_{\alpha, BM}^{(x_0)}(t))_{t \geq 0}$ is a sample path of a two-dimensional Feller process. Let

$$T^+(\mathbb{R}^2) := \{(x, a) \in \mathbb{R}^2 | x \geq a\} \quad \text{Diag}(\mathbb{R}^2) := \{(x, x) | x \in \mathbb{R}\}$$

For $(x_0, a_0) \in T^+(\mathbb{R}^2)$ we define the process

$$(5.2.1) \quad (\Xi_{\alpha, BM}^{(x_0, a_0)}(t))_{t \geq 0} = (\xi_{\alpha, BM}^{(x_0, a_0)}(t), \theta_{\alpha, BM}^{(x_0, a_0)}(t))_{t \geq 0} \\ := \left(a_0 + |x_0 - a_0 + \tilde{B}_t| - \frac{1}{\alpha} \ell_t^{a_0 - x_0}(\tilde{B}), a_0 - \frac{1}{\alpha} \ell_t^{a_0 - x_0}(\tilde{B}) \right)_{t \geq 0}$$

where $(\tilde{B}_t)_{t \geq 0}$ is a Brownian motion starting from 0. $\Xi_{\alpha, BM}^{(x_0, x_0)}$ has the same law as $\Xi_{\alpha, BM}^{(x_0)}$. The family of paths $(\Xi_{\alpha, BM}^{(x_0, a_0)})_{x_0 \geq a_0}$ are the sample paths of the same Feller semi-group on $T^+(\mathbb{R}^2)$ starting from all possible positions. Next we describe this semi-group in terms of generator and domain. Let f be a continuous function on $T^+(\mathbb{R}^2)$, \mathcal{C}^2 on the interior of $T^+(\mathbb{R}^2)$, such that all its second order derivatives extend continuously to $\text{Diag}(\mathbb{R}^2)$. This implies in particular that the first order derivatives also extend continuously to $\text{Diag}(\mathbb{R}^2)$. We write $\partial_1 f$, $\partial_2 f$ and $\partial_{1,1} f$ for the first order derivative relatively to the first variable, the second variable and the second order derivative relatively the first variable. Applying Itô-Tanaka's formula we get

$$f(\Xi_{\alpha, BM}^{(x_0, a_0)}(t)) = f(x_0, a_0) + \int_0^t \partial_1 f(\Xi_{\alpha, BM}^{(x_0, a_0)}(s)) \text{sgn}(x_0 - a_0 + \tilde{B}_s) d\tilde{B}_s + \\ \int_0^t \left(\left(1 - \frac{1}{\alpha}\right) \partial_1 - \frac{1}{\alpha} \partial_2 \right) f(\Xi_{\alpha, BM}^{(x_0, a_0)}(s)) d_s \ell_s^{a_0 - x_0}(\tilde{B}) + \frac{1}{2} \int_0^t \partial_{1,1} f(\Xi_{\alpha, BM}^{(x_0, a_0)}(s)) ds$$

Let $\mathcal{D}_{\alpha, BM}$ be the set of continuous functions f on $D_{\mathbb{R}}$, \mathcal{C}^2 on the interior of $T^+(\mathbb{R}^2)$, such that all the second order derivatives extend continuously to $\text{Diag}(\mathbb{R}^2)$ and that moreover satisfy the following constraints: f and $\partial_{1,1} f$ are uniformly continuous and bounded (which also implies that $\partial_1 f$ is bounded by the inequality $\|\partial_1 f\|_{\infty} \leq 2\sqrt{\|f\|_{\infty} \|\partial_{1,1} f\|_{\infty}}$) and on $\text{Diag}(\mathbb{R}^2)$ the following equality holds:

$$\left(\left(1 - \frac{1}{\alpha}\right) \partial_1 - \frac{1}{\alpha} \partial_2 \right) f(x, x) = 0$$

If $f \in \mathcal{D}_{\alpha, BM}$ then $\frac{1}{t}(\mathbb{E}[f(\Xi_{\alpha, BM}^{x_0, a_0}(t))] - f(x_0, a_0))$ converges as $t \rightarrow 0^+$, uniformly for $(x_0, a_0) \in T^+(\mathbb{R}^2)$, to $\frac{1}{2}\partial_{1,1}f(x_0, a_0)$. Moreover $\mathcal{D}_{\alpha, BM}$ is a core for $\frac{1}{2}\partial_{1,1}$ in the space of continuous bounded function on $T^+(\mathbb{R}^2)$.

Next we describe what we obtain if we glue together the loops, seen as excursion, ordered in the decreasing sense of their minima, where instead of $\mathcal{L}_{\alpha, BM}$ we use the Poisson ensemble of Markov loops associated to a general diffusion. Let I be an open interval of \mathbb{R} and \tilde{L} a generator on I of form

$$\tilde{L} = \frac{1}{\tilde{m}(x)} \frac{d}{dx} \left(\frac{1}{\tilde{w}(x)} \frac{d}{dx} \right)$$

with zero Dirichlet boundary conditions. Let \tilde{S} be a primitive of $\tilde{w}(x)$. We assume that $\tilde{S}(\sup I) = +\infty$. Let

$$T^+(I^2) := \{(x, a) \in I^2 | x \geq a\} \quad \text{Diag}(I^2) := \{(x, x) | x \in I\}$$

Let $\widehat{T^+(I^2)}$ be the closure of $T^+(I^2)$ in $(\inf I, \sup I)^2$.

Given any $x'_0 \geq a'_0 > \frac{1}{2}\tilde{S}(\inf I)$ let $\tilde{\zeta}_\alpha$ be the first time $\Xi_{\alpha, BM}^{(x'_0, a'_0)}$ hits $\frac{1}{2}\tilde{S}(\inf I)$. Let

$$\tilde{I}_t := \int_0^t \frac{1}{\tilde{m}}(\tilde{S}^{-1}(2\xi_{\alpha, BM}^{(x'_0, a'_0)}(s))) ds$$

Let $(\tilde{I}_t^{-1})_{0 \leq t < \tilde{I}_{\tilde{\zeta}_\alpha}}$ be the inverse function of $(\tilde{I}_t)_{0 \leq t < \tilde{\zeta}_\alpha}$. It is a family of stopping times for $\Xi_{\alpha, BM}^{(x'_0, a'_0)}$. For $x_0 \geq a_0 \in I$ and $t < \tilde{I}_{\tilde{\zeta}_\alpha}$ let

$$\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)}(t) = (\xi_{\alpha, \tilde{L}}^{(x_0, a_0)}(t), \theta_{\alpha, \tilde{L}}^{(x_0, a_0)}(t)) := \Xi_{\alpha, BM}^{(\tilde{S}(2x_0), \tilde{S}(2a_0))}(\tilde{I}_t^{-1})$$

If $\alpha = 1$ then $\xi_{\alpha, \tilde{L}}^{(x_0, a_0)}$ is just the sample paths starting x_0 of a diffusion of generator \tilde{L} . Let $\widehat{\mathcal{D}}_{\alpha, \tilde{L}}$ be the space of continuous functions f on $T^+(I^2)$ satisfying

- $f \circ \tilde{S}^{-1}$ is \mathcal{C}^2 on the interior of $T^+(I^2)$ and all the second order derivatives extend continuously to $\text{Diag}(I^2)$.
- $f(x, a)$ and $\frac{1}{\tilde{m}(x)}\partial_1 \left(\frac{1}{\tilde{w}(x)}\partial_1 f(x, a) \right)$ are bounded on $T^+(I^2)$ and extend continuously to $\widehat{T^+(I^2)}$.
- $f(x, a)$ and $\frac{1}{\tilde{m}(x)}\partial_1 \left(\frac{1}{\tilde{w}(x)}\partial_1 f(x, a) \right)$ converge to 0 as a converges to $\inf I$ uniformly in x .
- On $\text{Diag}(I^2)$ the following equality holds:

$$(5.2.2) \quad \left(\left(1 - \frac{1}{\alpha} \right) \partial_1 - \frac{1}{\alpha} \partial_2 \right) f(x, x) = 0$$

Lemma 5.3. — $(\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)})_{x_0 \geq a_0 \in I}$ is a family of sample path starting from all possible positions of the same Markovian or sub-Markovian semi-group on $T^+(I^2)$. The law of the path $\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)}$ depends weakly continuously on the starting point (x_0, a_0) .

The domain of the generator of this semi-group contains $\widehat{\mathcal{D}}_{\alpha, \tilde{L}}$, and on this space the generator equals

$$\frac{1}{\tilde{m}(x)} \partial_1 \left(\frac{1}{\tilde{w}(x)} \partial_1 \right)$$

Moreover there is only one Markovian or sub-Markovian semi-group with such generator on $\widehat{\mathcal{D}}_{\alpha, \tilde{L}}$.

Proof. — Since a change of scale does not alter the validity of the above statement, we can assume that $\tilde{w} \equiv 2$. Then $\sup I = +\infty$. $(\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)}(t))_{0 \leq t \leq \tilde{I}_{\tilde{\zeta}_\alpha}}$ is then obtained from $(\Xi_{\alpha, BM}^{(x_0, a_0)}(t))_{0 \leq t < \tilde{\zeta}_\alpha}$ by a random time change. The Markov property and the continuous dependence on the starting point for $\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)}$ follows from analogous properties for $\Xi_{\alpha, BM}^{(x_0, a_0)}$. If $f \in \widehat{\mathcal{D}}_{\alpha, \tilde{L}}$ then

$$\left(f(\Xi_{\alpha, BM}^{(x_0, a_0)}(\tilde{I}_t^{-1} \wedge \tilde{\zeta}_\alpha)) - \frac{1}{2} \int_0^{\tilde{I}_t^{-1} \wedge \tilde{\zeta}_\alpha} \partial_{1,1} f(\Xi_{\alpha, BM}^{(x_0, a_0)}(s)) ds \right)_{t \geq 0}$$

is a local martingale. We can rewrite it as

$$\left(f(\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)}(t \wedge \tilde{I}_{\tilde{\zeta}_\alpha})) - \int_0^t \frac{1}{2\tilde{m}(\xi_{\alpha, \tilde{L}}^{(x_0, a_0)}(s))} \partial_{1,1} f(\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)}(s)) 1_{s < \tilde{I}_{\tilde{\zeta}_\alpha}} ds \right)_{t \geq 0}$$

The above local martingale is bounded on all finite time intervals and thus is a true martingale. Since $\frac{1}{2\tilde{m}(x)} \partial_{1,1} f(x, a)$ converges to 0 as a converges to $\inf I$, uniformly in x , it follows that

$$f(\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)}(t \wedge \tilde{I}_{\tilde{\zeta}_\alpha})) = 1_{t < \tilde{I}_{\tilde{\zeta}_\alpha}} f(\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)}(t))$$

Thus

$$\lim_{t \rightarrow 0^+} \frac{1}{t} \left(\mathbb{E} \left[1_{t < \tilde{I}_{\tilde{\zeta}_\alpha}} f(\Xi_{\alpha, \tilde{L}}^{(x_0, a_0)}(t)) \right] - f(x_0, a_0) \right) = \frac{1}{2\tilde{m}(x_0)} \partial_{1,1} f(x_0, a_0)$$

Moreover the above convergence is uniform in (x_0, a_0) because $\frac{1}{2\tilde{m}(x)} \partial_{1,1} f(x, a)$ extends continuously to $T^+(\widehat{I^2})$.

To prove the uniqueness of the semi-group we need to show that there is $\lambda > 0$ such that

$$\left(\frac{1}{2\tilde{m}(x)} \partial_{1,1} - \lambda \right) (\widehat{\mathcal{D}}_{\alpha, \tilde{L}})$$

is sufficiently large, for instance that it contains all functions with compact support in $T^+(\widehat{I^2})$. Let g be such a function and $\lambda > 0$. Consider the equation

$$(5.2.3) \quad \frac{1}{2\tilde{m}(x)} \partial_{1,1} f(x, a) - \lambda f(x, a) = g(x, a)$$

Let $\tilde{u}_{\lambda, \downarrow}$ be a positive decreasing solution to

$$\frac{1}{2\tilde{m}(x)} \frac{d^2 u}{dx^2}(x) - \lambda u(x) = 0$$

Let

$$f_0(x, a) := \tilde{u}_{\lambda, \downarrow}(x) \int_x^{+\infty} \int_y^{+\infty} 2\tilde{m}(z)g(z, a)\tilde{u}_{\lambda, \downarrow}(z) dz \frac{dy}{\tilde{u}_{\lambda, \downarrow}(y)^2}$$

Then f_0 is a solution to (5.2.3) and it is compactly supported in $T^+(I^2)$. We look for the solutions to (5.2.3) of form

$$f(x, a) = f_0(x, a) + C(a)\tilde{u}_{\lambda, \downarrow}(x)$$

f satisfies the constraint (5.2.2) if and only if C satisfies

$$-\frac{1}{\alpha}\tilde{u}_{\lambda, \downarrow}(a)\frac{dC}{da}(a) + \left(1 - \frac{1}{\alpha}\right)\frac{d\tilde{u}_{\lambda, \downarrow}}{dx}(a)C(a) + h(a) = 0$$

where

$$h(a) = \left(\left(1 - \frac{1}{\alpha}\right)\partial_1 - \frac{1}{\alpha}\partial_2 \right) f_0(a, a)$$

h is compactly supported in I . We can set

$$C(a) = \tilde{u}_{\lambda, \downarrow}(a)^{\alpha-1} \int_{\inf I}^x \frac{h(y)}{\tilde{u}_{\lambda, \downarrow}(y)^\alpha} dy$$

C is zero in the neighbourhood of $\inf I$. Moreover $\tilde{u}_{\lambda, \downarrow}$ has a limit at $+\infty$. It follows that $f \in \widehat{\mathcal{D}}_{\alpha, \tilde{L}}$. \square

Let L be the generator of a diffusion on I of form (2.2.1). Let $x_0 \in I$. Consider the loops γ in $\mathcal{L}_{\alpha, L}$ such that $\min \gamma < x_0$, rooted at $\operatorname{argmin} \gamma$, seen as excursions. Let $(\xi_{\alpha, L}^{(x_0)}(t))_{0 \leq t < \zeta_\alpha}$ be the path on I obtained by glueing together these excursions ordered in the decreasing sense of their minima. Let

$$\theta_{\alpha, L}^{(x_0)}(t) := \min_{[0, t]} \xi_{\alpha, L}^{(x_0)}$$

$$\Xi_{\alpha, L}^{(x_0)} := (\xi_{\alpha, L}^{(x_0)}, \theta_{\alpha, L}^{(x_0)})$$

Proposition 5.4. — *Let $\tilde{L} := \operatorname{Conj}(u_\downarrow, L)$. Then $(\Xi_{\alpha, L}^{(x_0)}(t))_{0 \leq t < \zeta_\alpha}$ has the same law as $(\Xi_{\alpha, \tilde{L}}^{(x_0, x_0)}(t))_{0 \leq t < \tilde{\zeta}_\alpha}$. So it is a sample path of a two-dimensional Feller process.*

In particular for $\alpha = 1$, $\xi_{1, L}^{(x_0)}$ is the sample path of a diffusion of generator \tilde{L} . For all $\alpha > 0$

$$\liminf_{t \rightarrow \zeta_\alpha} \xi_{\alpha, L}^{(x_0)}(t) = \inf I$$

If L is the generator of a recurrent diffusion then

$$\limsup_{t \rightarrow \zeta_\alpha} \xi_{\alpha, L}^{(x_0)}(t) = \sup I$$

Otherwise

$$\limsup_{t \rightarrow \zeta_\alpha} \xi_{\alpha, L}^{(x_0)}(t) = \inf I$$

Proof. — First notice that if L is the generator of a recurrent diffusion then $\tilde{L} = L$. Otherwise a diffusion of generator $\tilde{L} = L$ is, put informally, a diffusion of generator L conditioned to converge to $\inf I$ (which may occur with zero probability). From h-transform invariance of the measure on loops follows that $\mathcal{L}_{\alpha,L} = \mathcal{L}_{\alpha,\tilde{L}}$. From property 3.8 (iv) and corollary 3.12 follows that $\Xi_{\alpha,L}^{(x_0)}$ is obtained from $\Xi_{\alpha,BM}$ by scale and time change in the same way as $\Xi_{\alpha,\tilde{L}}^{(x_0,x_0)}$ and thus $\Xi_{\alpha,L}^{(x_0)}$ and $\Xi_{\alpha,\tilde{L}}^{(x_0,x_0)}$ have the same law. Regarding the limits of $\xi_{\alpha,L}^{(x_0)}$ at ζ_α , we need just to observe that they hold if L is the generator of the Brownian motion on an interval of form $(a, +\infty)$, $a \in [-\infty, +\infty)$, and by time and scale change they hold in general. \square

As explained in the proposition 5.1, the knowledge of the path $(\xi_{\alpha,L}^{(x_0)}(t))_{0 \leq t < \zeta_\alpha}$ alone is enough to reconstruct $\mathcal{L}_{\alpha,L} \cap \{\gamma \in \mathfrak{L}^* \mid \min \gamma < x_0\}$. From this we deduce the following

Corollary 5.5. — *If L is the generator of a transient diffusion, it is possible to construct on the same probability space $\mathcal{L}_{\alpha,L}$ and a continuous version of the occupation field $(\hat{\mathcal{L}}_{\alpha,L}^x)_{x \in I}$.*

Proof. — By scale and time change covariance and h-transform invariance of the Poisson ensembles of loops, it is enough to prove the proposition in case of a Brownian motion on $(0, +\infty)$ killed at 0. Let $(x_n)_{n \geq 0}$ be an increasing sequence in $(0, +\infty)$ converging to $+\infty$. We consider a sequence of independent paths $(\xi_{\alpha,BM}^{(x_n,x_n)})_{n \geq 0}$ defined by (5.2.1). Let

$$T_{n,x_{n-1}} := \inf \{t \geq 0 \mid \xi_{\alpha,BM}^{(x_n,x_n)}(t) = x_{n-1}\}$$

where conventionally we set $x_{-1} := 0$. By decomposing on $[0, T_{n,x_{n-1}}]$ the restricted path $(\xi_{\alpha,BM}^{(x_n,x_n)}(t))_{0 \leq t < T_{n,x_{n-1}}}$ one can reconstruct a family of loops γ such that $\min \gamma \in (x_{n-1}, x_n)$: there is a random countable set \mathcal{B}_n of disjoint compact subintervals $[b^-, b^+]$ of $[0, T_{n,x_{n-1}}]$ such that

$$\{(\xi_{\alpha,BM}^{(x_n,x_n)}(b^- + t))_{0 \leq t \leq b^+ - b^-} \mid [b^-, b^+] \in \mathcal{B}_n\} = \mathcal{L}_{\alpha,BM} \cap \{\gamma \in \mathfrak{L}^* \mid \min \gamma \in (x_{n-1}, x_n)\}$$

(see (5.1.3)). The union of all previous families of loops for $n \geq 0$ is a Poisson ensemble of loops $\mathcal{L}_{\alpha,BM} \cap \{\gamma \in \mathfrak{L}^* \mid \min \gamma > 0\}$.

Each of $\xi_{\alpha,BM}^{(x_n,x_n)}$ is a semi-martingale and its quadratic variation is

$$\langle \xi_{\alpha,BM}^{(x_n,x_n)}, \xi_{\alpha,BM}^{(x_n,x_n)} \rangle_t = t$$

Moreover for all $x \in \mathbb{R}$

$$\int_0^t 1_{\xi_{\alpha,BM}^{x_n,x_n} = x} d\xi_{\alpha,BM}^{(x_n,x_n)}(s) = \left(1 - \frac{1}{\alpha}\right) \int_0^t 1_{\ell_s^0(\tilde{B}) = \alpha x} d_s \ell_s^0(\tilde{B}) = 0$$

From theorems 1.1 and 1.7 in [24], chapter VI, §1, follows that we can construct on the same probability space $\xi_{\alpha,BM}^{(x_n,x_n)}$ and a space-time continuous version of local times

$(\ell_t^x(\xi_{\alpha, BM}^{(x_n, x_n)}))_{x \in \mathbb{R}, t \geq 0}$ of $\xi_{\alpha, BM}^{(x_n, x_n)}$ relatively to the Lebesgue measure. In particular $x \mapsto \ell_{T_n, x_{n-1}}^x(\xi_{\alpha, BM}^{(x_n, x_n)})$ is continuous. If $[b^-, b^+] \in \mathcal{J}_n$, then

$$(\ell_{b^+}^x(\xi_{\alpha, BM}^{(x_n, x_n)}) - \ell_{b^-}^x(\xi_{\alpha, BM}^{(x_n, x_n)}))_{x > 0}$$

is the occupation field of the loop corresponding to the time interval $[b^-, b^+]$. We need to check that a.s

$$(5.2.4) \quad \forall x > 0, \ell_{T_n, x_{n-1}}^x(\xi_{\alpha, BM}^{(x_n, x_n)}) = \sum_{[b^-, b^+] \in \mathcal{B}_n} \ell_{b^+}^x(\xi_{\alpha, BM}^{(x_n, x_n)}) - \ell_{b^-}^x(\xi_{\alpha, BM}^{(x_n, x_n)})$$

For $x > 0$, consider the random set of times

$$(5.2.5) \quad \{t \in [0, T_n, x_{n-1}] | \xi_{\alpha, BM}^{(x_n, x_n)}(t) = x\} \setminus \bigcup_{[b^-, b^+] \in \mathcal{B}_n} [b^-, b^+]$$

If x is a minimum of a loop embedded in $(\xi_{\alpha, BM}^{(x_n, x_n)}(t))_{0 \leq t < T_n, x_{n-1}}$ or if $x \notin (x_{n-1}, x_n)$ then the set (5.2.5) is empty. Otherwise it is reduced to one point: the first hitting time of the level x . Almost surely, for all $x > 0$, the measure $d_t \ell_t^x(\xi_{\alpha, BM}^{(x_n, x_n)})$ is supported in $\{t \geq 0 | \xi_{\alpha, BM}^{(x_n, x_n)}(t) = x\}$ and has no atoms, and thus does not charge the set (5.2.5). This implies (5.2.4). Finally we can conclude that $(\ell_{T_n, x_{n-1}}^x(\xi_{\alpha, BM}^{(x_n, x_n)}))_{x > 0}$ is the occupation field of $\mathcal{L}_{\alpha, BM} \cap \{\gamma \in \mathcal{L}^* | \min \gamma \in (x_{n-1}, x_n)\}$.

The occupation field of $\mathcal{L}_{\alpha, BM} \cap \{\gamma \in \mathcal{L}^* | \min \gamma > 0\}$ is

$$\left(\sum_{n \geq 0} \ell_{T_n, x_{n-1}}^x(\xi_{\alpha, BM}^{(x_n, x_n)}) \right)_{x > 0}$$

The above sum is locally finite and thus varies continuously with x . \square

5.3. The case $\alpha = 1$

According to proposition 5.4 in case $\alpha = 1$ the Poisson ensemble of loops $\mathcal{L}_{1, L}$ can be recovered from sample paths of one-dimensional diffusions. A similar property was observed for loops of the two-dimensional Brownian Motion and of Markov jump processes on graphs. In [13], chapter 8, it is shown that by launching consecutively symmetric Markov jump processes from different vertices of a finite graph and applying the Wilson's algorithm ([31]), one can simultaneously construct a uniform spanning tree of the graph with prescribed weights on the edges and an independent Poisson ensemble of Markov loops of parameter $\alpha = 1$. If \mathbb{D} is a simply-connected open domain of \mathbb{C} other than \mathbb{C} , it was shown in [32] that one can couple a Brownian motion on \mathbb{D} , killed at hitting $\partial\mathbb{D}$, and a simple curve (SLE_2) with same extremal points such that the latter appears as the loop-erasure of the first. It is conjectured that given this loop-erased Brownian motion and an independent Poisson ensemble of Brownian loops of parameter 1, by attaching to the simple curve the loops that

cross it one reconstructs a Brownian sample path. See [19], conjecture 1, and [18], theorem 7.3.

in case of one-dimensional diffusions one can partially recover $\mathcal{L}_{1,L}$ from Markovian sample paths otherwise than slicing $\xi_{1,L}^{(x_0)}$ in excursions. The next result has an analogue for loops of Markov jump processes on graphs. See [13], remark 21.

Proposition 5.6. — *Assume that L is the generator of a transient diffusion. Let $x \in I$. Let $(X_t)_{0 \leq t < \zeta}$ be the sample path of a diffusion of generator L started from x . Let \widehat{T}_x the last time X visits x . For $l \geq 0$ let*

$$\tau_l^x := \{t \geq 0 | \ell_t^x(X) > l\}$$

Let $(q_j)_{j \in \mathbb{N}}$ be a Poisson-Dirichlet partition $PD(0,1)$ of $[0,1]$, independent from X , ordered in an arbitrary way. Let

$$l_j := \ell_{\zeta}^x(X) \sum_{i=0}^j q_i$$

The family of bridges $((X_t)_{\tau_{l_{j-1}}^x \leq t \leq \tau_{l_j}^x})_{j \geq 0}$ has, up to unrooting, the same law as the loops in

$$\mathcal{L}_{1,L} \cap \{\gamma \in \mathfrak{L}^* | x \in \gamma([0, T(\gamma)])\}$$

In particular $(X_t)_{0 \leq t \leq \widehat{T}_x}$ can be obtained through sticking together all the loops in $\mathcal{L}_{\alpha,L}$ that visit x .

Proof. — According to corollary 3.11, $(\ell^x(\gamma))_{\gamma \in \mathcal{L}_{\alpha,L,\gamma}}$ visits x is a Poisson ensemble of intensity $e^{-\frac{l}{G(x,x)}} \frac{dl}{l}$. Thus $\widehat{\mathcal{L}}_{\alpha,L}^x$ is an exponential r.v. with mean $G(x,x)$ and has the same law as $\ell_{\zeta}^x(X)$. Moreover the Poisson ensemble $(\ell^x(\gamma))_{\gamma \in \mathcal{L}_{\alpha,L,\gamma}}$ visits x has up to reordering the same law as $(l_j - l_{j-1})_{j \geq 0}$. Almost surely $l \mapsto \tau_l^x$ does not jump at any l_j . Conditionally on $(l_j)_{j \geq 0}$, $((X_t)_{\tau_{l_{j-1}}^x \leq t \leq \tau_{l_j}^x})_{j \geq 0}$ is an independent family of bridges and $(X_t)_{\tau_{l_{j-1}}^x \leq t \leq \tau_{l_j}^x}$ has the same law as $(X_t)_{0 \leq t \leq \tau_{l_j - l_{j-1}}^x}$. We conclude using identity (3.3.5) and the theory of marked Poisson ensembles. \square

Assume that L is the generator of a transient diffusion. Let $x \in I$ and let $(X_t)_{0 \leq t < \zeta}$ be a sample path starting from x of the diffusion corresponding to L . We will describe two different ways to slice $(X_t)_{0 \leq t < \zeta}$ so as to obtain the loops

$$\mathcal{L}_{1,L} \cap \{\gamma \in \mathfrak{L}^* | \gamma([0, T(\gamma)]) \cap [X(0), X(\zeta^-)] \text{ (or } [X(\zeta^-), X(0)]) \neq \emptyset\}$$

The first method corresponds to the "loop-erasure procedure" applied to $(X_t)_{0 \leq t < \zeta}$ and the second to the "loop-erasure procedure" applied to the time-reversed path $(X_{\zeta-t})_{0 < t \leq \zeta}$. Let \widehat{T}_x be the last time $(X_t)_{0 \leq t < \zeta}$ visits x . Let \widetilde{T} be the first time X hits X_{ζ^-} . If $X_{\zeta^-} \in \partial I$ then $\widetilde{T} = \zeta$. Let $(q_j)_{j \in \mathbb{N}}$ be a Poisson-Dirichlet partition $PD(0,1)$ of $[0,1]$, independent from X . The first method of decomposition is the following:

- The path $(X_t)_{0 \leq t \leq \hat{T}_x}$ is decomposed in bridges $((X_t)_{\tau_{i_j-1}^x \leq t \leq \tau_{i_j}^x})_{j \geq 0}$ from x to x by applying the Poisson-Dirichlet partition $(q_j)_{j \in \mathbb{N}}$ to $\ell_\zeta^x(X)$, as described in proposition 5.6.
- Given the path $(X_{\hat{T}_x+t})_{0 \leq t < \zeta - \hat{T}_x}$, if $X_{\zeta^-} < x$ we define

$$\mathfrak{b}^+ := \left\{ t \in [0, \zeta - \hat{T}_x) \mid X_{\hat{T}_x+t} = \sup_{s \in [t, \zeta - \hat{T}_x)} X_{\hat{T}_x+s} \right. \\ \left. \text{and } \exists \varepsilon \in (0, t) \text{ s.t. } \forall s \in (t - \varepsilon, t), X_{\hat{T}_x+s} < X_{\hat{T}_x+t} \right\}$$

\mathfrak{b}^+ is countable and we define on \mathfrak{b}^+ the map b^- :

$$b^-(t) := \sup \{ s \in [0, t) \mid X_{\hat{T}_x+s} = X_{\hat{T}_x+t} \}$$

$((X_{\hat{T}_x+b^-(t)+s})_{0 \leq s \leq t-b^-(t)})_{t \in \mathfrak{b}^+}$ is the family of negative excursions of the path $(X_{\hat{T}_x+t})_{0 \leq t < \zeta - \hat{T}_x}$ below $(\sup_{[\hat{T}_x+t, \zeta)} X)_{0 \leq t < \zeta - \hat{T}_x}$. If $X_{\zeta^-} > x$ then

$$\mathfrak{b}^+ := \left\{ t \in [0, \zeta - \hat{T}_x) \mid X_{\hat{T}_x+t} = \inf_{s \in [t, \zeta - \hat{T}_x)} X_{\hat{T}_x+s} \right. \\ \left. \text{and } \exists \varepsilon \in (0, t) \text{ s.t. } \forall s \in (t - \varepsilon, t), X_{\hat{T}_x+s} > X_{\hat{T}_x+t} \right\}$$

We define on \mathfrak{b}^+ the map b^- :

$$b^-(t) := \sup \{ s \in [0, t) \mid X_{\hat{T}_x+s} = X_{\hat{T}_x+t} \}$$

$((X_{\hat{T}_x+b^-(t)+s})_{0 \leq s \leq t-b^-(t)})_{t \in \mathfrak{b}^+}$ are the positive excursions of $(X_{\hat{T}_x+t})_{0 \leq t < \zeta - \hat{T}_x}$ above $(\inf_{[\hat{T}_x+t, \zeta)} X)_{0 \leq t < \zeta - \hat{T}_x}$.

- We denote $\mathcal{L}^1((X_t)_{0 \leq t < \zeta})$ the set of loops

$$\{(X_{\tau_{i_j-1}^x+s})_{0 \leq s \leq \tau_{i_j}^x - \tau_{i_j-1}^x} \mid j \geq 0\} \cup \{(X_{\hat{T}_x+b^-(t)+s})_{0 \leq s \leq t-b^-(t)} \mid t \in \mathfrak{b}^+\}$$

where the loops are considered to be unrooted.

The second method of decomposition is the following:

- If $X_{\zeta^-} < x$ we define

$$\mathfrak{b}^- := \left\{ t \in [0, \tilde{T}) \mid X_t = \inf_{[0, t]} X \text{ and } \exists \varepsilon > 0 \text{ s.t. } \forall s \in (t, t + \varepsilon), X_s > X_t \right\}$$

On \mathfrak{b}^- we define the map b^+ :

$$b^+(t) := \inf \{ s \in (t, \tilde{T}) \mid X_s = X_t \}$$

$((X_{t+s})_{0 \leq s \leq b^+(t)-t})_{t \in \mathfrak{b}^-}$ are the positive excursions of the path $(X_t)_{0 \leq t < \tilde{T}}$ above $(\inf_{[0, t]} X)_{0 \leq t < \tilde{T}}$. This is exactly the decomposition described in the previous section 5.2. If $X_{\zeta^-} > x$ then

$$\mathfrak{b}^- := \left\{ t \in [0, \tilde{T}) \mid X_t = \sup_{[0, t]} X \text{ and } \exists \varepsilon > 0 \text{ s.t. } \forall s \in (t, t + \varepsilon), X_s < X_t \right\}$$

The map b^+ defined on \mathfrak{b}^- is

$$b^+(t) := \inf\{s \in (t, \tilde{T}) \mid X_s = X_t\}$$

$((X_{t+s})_{0 \leq s \leq b^+(t)-t})_{t \in \mathfrak{b}^-}$ are the negative excursions of the path $(X_t)_{0 \leq t < \tilde{T}}$ below $(\sup_{[0,t]} X)_{0 \leq t \leq \tilde{T}}$.

– If $\tilde{T} < \zeta$ we introduce:

$$\tilde{l}_j := \ell_\zeta^{X_{\zeta^-}}(X) \sum_{i=0}^j q_i$$

and

$$\tau_{\tilde{l}_j} := \inf\{t \in [\tilde{T}, \zeta) \mid \ell_t^{X_{\zeta^-}}(X) > \tilde{l}_j\}$$

We decompose the path $(X_t)_{\tilde{T} \leq t < \zeta}$ in bridges $((X_t)_{\tau_{\tilde{l}_{j-1}} \leq t \leq \tau_{\tilde{l}_j}})_{j \geq 0}$ from X_{ζ^-} to X_{ζ^-} .

– We denote $\mathcal{L}^2((X_t)_{0 \leq t < \zeta})$ the set of loops

$$\{(X_{t+s})_{0 \leq s \leq b^+(t)-t} \mid t \in \mathfrak{b}^-\} \cup \{(X_{\tau_{\tilde{l}_{j-1}}+s})_{0 \leq s \leq \tau_{\tilde{l}_j}-\tau_{\tilde{l}_{j-1}}} \mid j \geq 0\}$$

where the loops are considered to be unrooted.

The loops in $\mathcal{L}^1((X_t)_{0 \leq t < \zeta})$ and $\mathcal{L}^2((X_t)_{0 \leq t < \zeta})$ are not the same but follow the same law.

Proposition 5.7. — $\mathcal{L}^1((X_t)_{0 \leq t < \zeta})$ and $\mathcal{L}^2((X_t)_{0 \leq t < \zeta})$, considered as collections of unrooted loops, have the same law. Let $\mathcal{L}_{1,L}$ be a Poisson ensemble of loops independent from X_{ζ^-} . Then $\mathcal{L}^1((X_t)_{0 \leq t < \zeta})$ and $\mathcal{L}^2((X_t)_{0 \leq t < \zeta})$ have the same law as

$$(5.3.1) \quad \mathcal{L}_{1,L} \cap \{\gamma \in \mathfrak{L}^* \mid \gamma([0, T(\gamma)]) \cap [X(0), X(\zeta^-)] \text{ (or } [X(\zeta^-), X(0)]) \neq \emptyset\}$$

Proof. — First we will prove that $\mathcal{L}^2((X_t)_{0 \leq t < \zeta})$ has the same law as (5.3.1). If $\mathbb{P}(X_{\zeta^-} = \inf I) > 0$, then conditionally on $X_{\zeta^-} = \inf I$, $(X_t)_{0 \leq t < \zeta}$ has the law of a sample path corresponding to the generator $\text{Conj}(u_\downarrow, L)$. If $y \in I \cap (-\infty, x]$ and y is in the support of κ (the killing measure in L) then conditionally on $X_{\zeta^-} = y$, $(X_t)_{0 \leq t < \zeta}$ is distributed according to the measure $\frac{1}{G(x,y)} \mu_L^{x,y}$ (property 3.3 (i)). According to the lemma 3.4, $(X_t)_{0 \leq t \leq \tilde{T}}$ and $(X_{\tilde{T}+t})_{0 \leq t \leq \zeta - \tilde{T}}$ are independent conditionally $X_{\zeta^-} = y$, $(X_t)_{0 \leq t \leq \tilde{T}}$ having the law of a sample path corresponding to the generator $\text{Conj}(u_\downarrow, L)$, run until hitting y , and $(X_{\tilde{T}+t})_{0 \leq t \leq \zeta - \tilde{T}}$ following the law $\frac{1}{G(y,y)} \mu_L^{y,y}$. From proposition 5.4 and 5.6 follows that $\mathcal{L}^2((X_t)_{0 \leq t < \zeta})$ and (5.3.1) have the same law on the event $X_{\zeta^-} \leq x$. Symmetrically this also true on the event $X_{\zeta^-} \geq x$.

The decomposition $\mathcal{L}^1((X_t)_{0 \leq t < \zeta})$ is obtained by first applying the decomposition \mathcal{L}^2 to the time-reversed path $(X_{\zeta-t})_{0 < t \leq \zeta}$ and then applying again the time-reversal to the obtained loops. The law of the loops in (5.3.1) is invariant by time-reversal. Let $y \in I$, y in the support of κ . Conditionally on $X_{\zeta^-} = y$, the law of $(X_{\zeta-t})_{0 < t \leq \zeta}$ is

$\frac{1}{G(x,y)}\mu^{y,x}$. So applying the decomposition \mathcal{L}^2 to the path $(X_{\zeta-t})_{0 < t \leq \zeta}$ conditioned by $X_{\zeta^-} = y$ gives

$$\mathcal{L}_{1,L} \cap \{\gamma \in \mathfrak{L}^* | \gamma([0, T(\gamma)]) \cap [y, x] \text{ (or } [x, y]) \neq \emptyset\}$$

If $\mathbb{P}(X_{\zeta^-} = \inf I) > 0$ then conditionally on $X_{\zeta^-} = \inf I$, the path $(X_t)_{0 \leq t < \zeta}$ is a limit as $y \rightarrow \inf I$ of paths following the law $\frac{1}{G(x,y)}\mu^{x,y}$ (i.e. the latter are restrictions of the former). Thus conditionally on $X_{\zeta^-} = \inf I$ $\mathcal{L}^1((X_t)_{0 \leq t < \zeta})$ is an increasing limit as $y \rightarrow \inf I$ of

$$\mathcal{L}_{1,L} \cap \{\gamma \in \mathfrak{L}^* | \gamma([0, T(\gamma)]) \cap [y, x] \neq \emptyset\}$$

which is

$$\mathcal{L}_{1,L} \cap \{\gamma \in \mathfrak{L}^* | \gamma([0, T(\gamma)]) \cap [\inf I, x] \neq \emptyset\}$$

Similar is true conditionally on $X_{\zeta^-} = \sup I$. □

CHAPTER 6

WILSON'S ALGORITHM IN DIMENSION ONE

6.1. Description of the algorithm

Given a finite undirected connected graph $\mathbb{G} = (V, E)$ and C a positive weight function on its edges, a Uniform Spanning Tree of the weighted graph \mathbb{G} is a random spanning tree with the occurrence probability of a spanning tree \mathcal{T} proportional to

$$\prod_{e \text{ edge of } \mathcal{T}} C(e)$$

The edges belonging to the Uniform Spanning Tree are a determinantal point process (transfer current theorem). In [31] Wilson showed how to sample a Uniform Spanning Tree using successive random walks to nearest neighbours, with transition probabilities proportional to C , starting from different vertices, and erasing the loops created by these random walks. The edges left after loop-erasure form a Uniform Spanning Tree. This is known as Wilson's algorithm. See [1] for a review. In [13], chapter 8, Le Jan shows that the loops erased during the execution of Wilson's algorithm are related to the Poisson ensemble of Markov loops of parameter 1.

In [13], chapter 10, Le Jan suggests that Wilson's algorithm can be adapted to the situation where the random walk on a graph is replaced by a transient diffusion on a subinterval I of \mathbb{R} . In this section we will describe the algorithm in the latter setting. The algorithm returns on one hand a sequence of one-dimensional paths which can be decomposed into a Poisson ensemble of Markov loops of parameter 1 (section 6.2), and on the other hand a pair of interwoven determinantal point processes on I , which may be interpreted as some kind of Uniform Spanning Tree. In section 6.3 we will derive the law of this pair of determinantal point processes in the setting where the underlying is a Brownian motion on \mathbb{R} with a killing measure. In section 6.4 we will give without proof the law in general case as it follows directly from the Brownian case.

Let I be a subinterval of \mathbb{R} and L a generator of a transient diffusion on I of form 2.2.1. Let κ be the killing measure in L , which may be zero. Let $(x_n)_{n \geq 1}$ be a sequence

of pairwise distinct points in I which is dense in I . Let $\left((X_t^{(x_n)})_{0 \leq t < \zeta_n}\right)_{n \geq 1}$ be a sequence of independent sample paths of the diffusion of generator L , with starting points $X_0^{(x_n)} = x_n$. In the first step of Wilson's algorithm we will recursively define sequences $(T_n)_{n \geq 1}$, $(\mathcal{Y}_n)_{n \geq 1}$ and $(\mathcal{J}_n)_{n \geq 1}$ where T_n is a killing time for $X^{(x_n)}$, \mathcal{Y}_n is a finite subset of $\text{Supp}(\kappa) \cup \partial I$ and \mathcal{J}_n is a finite set of disjoint compact subintervals of \bar{I} , some of which may be reduced to one point:

- $T_1 := \zeta_1$, $\mathcal{Y}_1 := \{X_{T_1^-}^{(x_1)}\}$, $\mathcal{J}_1 := \{[x_1, X_{T_1^-}^{(x_1)}]\}$ (or $\{[B_{T_1^-}^{(x_1)}, x_1]\}$).
- Assume that \mathcal{Y}_n and \mathcal{J}_n are constructed. If $x_{n+1} \in \bigcup_{J \subseteq \mathcal{J}_n} J$ then we set $T_{n+1} := 0$, $\mathcal{Y}_{n+1} := \mathcal{Y}_n$ and $\mathcal{J}_{n+1} := \mathcal{J}_n$. If $x_{n+1} \notin \bigcup_{J \subseteq \mathcal{J}_n} J$ then we define

$$T_{n+1} := \min\left(\zeta_n, \inf\left\{t \geq 0 \mid X_t^{(x_{n+1})} \in \bigcup_{J \subseteq \mathcal{J}_n} J\right\}\right)$$

If $X_{T_{n+1}^-}^{(x_{n+1})} \in \bigcup_{J \subseteq \mathcal{J}_n} J$ then there is a unique $J \in \mathcal{J}_n$ such that $X_{T_{n+1}^-}^{(x_{n+1})} \in J$. In this case we set $\mathcal{Y}_{n+1} := \mathcal{Y}_n$ and

$$\begin{aligned} \mathcal{J}_{n+1} := & (\mathcal{J}_n \setminus \{J\}) \cup \{J \cup [x_{n+1}, X_{T_{n+1}^-}^{(x_{n+1})}]\} \\ & \left(\text{or } (\mathcal{J}_n \setminus \{J\}) \cup \{J \cup [X_{T_{n+1}^-}^{(x_{n+1})}, x_{n+1}]\}\right) \end{aligned}$$

If $X_{T_{n+1}^-}^{(x_{n+1})} \notin \bigcup_{J \subseteq \mathcal{J}_n} J$ then we set $\mathcal{Y}_{n+1} := \mathcal{Y}_n \cup \{X_{T_{n+1}^-}^{(x_{n+1})}\}$ and

$$\mathcal{J}_{n+1} := \mathcal{J}_n \cup \{[x_{n+1}, X_{T_{n+1}^-}^{(x_{n+1})}]\} \left(\text{or } \mathcal{J}_n \cup \{[X_{T_{n+1}^-}^{(x_{n+1})}, x_{n+1}]\}\right)$$

It is immediate to check by induction the following facts:

- $\mathcal{Y}_n \subseteq \text{Supp}(\kappa) \cup \partial I$. More precisely $\mathcal{Y}_n \subseteq \text{Supp}(\kappa) \cup \{y \in \partial I \mid \mathbb{P}(X_{\zeta_n}^{(x_n)} = y) > 0\}$.
- The intervals in \mathcal{J}_n are pairwise disjoint.
- $\#\mathcal{Y}_n = \#\mathcal{J}_n \leq n$
- For every $y \in \mathcal{Y}_n$ there is one single $J \in \mathcal{J}_n$ such that $y \in J$.
- $\mathcal{Y}_n \subseteq \mathcal{Y}_{n+1}$
- If $n \leq n'$, then for every $J \in \mathcal{J}_n$ there is one single $J' \in \mathcal{J}_{n'}$ such that $J \subseteq J'$. We denote $\iota_{n,n'}$ the corresponding application from \mathcal{J}_n to $\mathcal{J}_{n'}$. The application $\iota_{n,n'}$ is injective. Trivially for $n \leq n' \leq n''$, $\iota_{n,n''} = \iota_{n',n''} \circ \iota_{n,n'}$
- For any $J \in \mathcal{J}_n$, $\partial J \subseteq \mathcal{Y}_n \cup \{x_1, \dots, x_n\}$.

In the second step of Wilson's algorithm we will take the limit of the sequence $((\mathcal{Y}_n, \mathcal{J}_n))_{n \geq 1}$ and define $(\mathcal{Y}_\infty, \mathcal{J}_\infty)$ as follows:

$$\mathcal{Y}_\infty := \bigcup_{n \geq 1} \mathcal{Y}_n \quad \mathcal{J}_\infty := \bigcup_{n \geq 1} \bigcup_{J \in \mathcal{J}_n} \left\{ \bigcup_{n' \geq n} \iota_{n,n'}(J) \right\}$$

\mathcal{Y}_∞ is a finite or countable subset of $\text{Supp}(\kappa) \cup \partial I$. \mathcal{J}_∞ is a finite or countable set of disjoint subintervals of \bar{I} , but these subintervals are not necessarily closed or bounded. For any $y \in \mathcal{Y}_\infty$, there is a single $J \in \mathcal{J}_\infty$ such that $y \in J$, and this induces a bijection

between \mathcal{Y}_∞ and \mathcal{J}_∞ . For any $J \in \mathcal{J}_n$, there is a single $J' \in \mathcal{J}_\infty$ such that $J \subseteq J'$. We define $\iota_{n,\infty}(J) = J'$. $\iota_{n,\infty}$ is injective. Trivially, for $n \leq n'$, $\iota_{n,\infty} = \iota_{n',\infty} \circ \iota_{n,n'}$. We will sometimes write $\mathcal{Y}_n(x_1, \dots, x_n)$, $\mathcal{J}_n(x_1, \dots, x_n)$, $\mathcal{Y}_\infty((x_n)_{n \geq 1})$ and $\mathcal{J}_\infty((x_n)_{n \geq 1})$ in order to emphasize the dependence on the starting points $(x_n)_{n \geq 1}$. In the sections 6.3 and 6.4 we will see that

- The set \mathcal{Y}_∞ is a.s. discrete.
- A.s. for any intervals $J \in \mathcal{J}_\infty$, $J \setminus \partial I$ is open
- The subset $I \setminus \bigcup_{J \in \mathcal{J}_\infty} J$ is a.s. discrete.
- The law of $(\mathcal{Y}_\infty, \mathcal{J}_\infty)$ does not depend on the choice of starting points $(x_n)_{n \geq 1}$.

We introduce $\mathcal{Z}_\infty := I \setminus \left(\bigcup_{J \in \mathcal{J}_\infty} J \right)$. We will further see that \mathcal{Y}_∞ and \mathcal{Z}_∞ are determinantal point processes.

The couple $(\mathcal{Y}_\infty, \mathcal{J}_\infty)$ may be interpreted as a spanning tree. Consider the following undirected "graph": Its set of "vertices" is $\bar{I} \cup \{\dagger\}$ where \dagger is a cemetery point outside of \bar{I} . Every point $x \in I$ is connected by an "edge" to its two infinitesimal neighbours $x-dx$ and $x+dx$. Every point in $Supp(\kappa)$ is connected by an "edge" to \dagger . Finally any point in $y \in \partial I$ such that $\mathbb{P}(X_{\zeta_n}^{(x_n)} = y) > 0$ is connected by an "edge" to \dagger . On this "graph" $(\mathcal{Y}_\infty, \mathcal{J}_\infty)$ induces the following "spanning tree": Each point in $\bigcup_{J \in \mathcal{J}_\infty} J$ is connected to its infinitesimal neighbours in I and \mathcal{Z}_∞ represents "edges" on I that are missing. Moreover every point in \mathcal{Y}_∞ is connected to \dagger .

There are two trivial cases in which $(\mathcal{Y}_\infty, \mathcal{J}_\infty)$ is deterministic. In the first one $\kappa = 0$ and I has one single regular or exit boundary point y characterized by $\mathbb{P}(X_{\zeta_n}^{(x_n)} = y) > 0$ (see [5], chapter 16, for the characterization of boundaries). Then \mathcal{Y}_∞ is made of this boundary point and \mathcal{J}_∞ contains one single interval $I \cup \mathcal{Y}_\infty$. \mathcal{Z}_∞ is empty. In the second case I does not have regular or exit boundaries and κ is proportional to a Dirac measure $c\delta_{y_0}$. Then $\mathcal{Y}_\infty = \{y_0\}$ and $\mathcal{J}_\infty = \{I\}$. \mathcal{Z}_∞ is again empty. In all other situation \mathcal{Z}_∞ is non-empty and random. See figure 2.a for an illustration of $(\mathcal{Y}_n, \mathcal{J}_n)$ for $1 \leq n \leq 5$ and figure 2.b for an illustration of $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$.




Fig. 2.a - Illustration of $((\mathcal{Y}_n, \mathcal{J}_n))_{1 \leq n \leq 5}$: x-dots represent the points of \mathcal{Y}_n and thick lines the intervals in \mathcal{J}_n .




Fig. 2.b - Illustration of $(\mathcal{Y}_\infty, \mathcal{J}_\infty)$: x-dots represent the points of \mathcal{Y}_∞ and diamonds the points of \mathcal{Z}_∞ .

6.2. The erased paths

During the execution of Wilson's algorithm we used the paths $\left((X_t^{(x_n)})_{0 \leq t < T_n} \right)_{n \geq 1}$. These paths can be further decomposed using the procedure described in the section 5.3.

Proposition 6.1. — *The family of unrooted loops*

$$\bigcup_{n \geq 1} \mathcal{L}^1 \left((X_t^{(x_n)})_{0 \leq t < T_n} \right)$$

has the same law as the Poisson ensemble $\mathcal{L}_{1,L}$. Moreover it is independent from $(\mathcal{Y}_\infty, \mathcal{J}_\infty)$.

Proof. — Let $\mathcal{L}_{1,L}$ be a Poisson ensemble of loops independent from the family of paths $\left((X_t^{(x_n)})_{0 \leq t < T_n} \right)_{n \geq 1}$. Using proposition 5.7 and induction is it immediate to show that the triple

$$\left(\mathcal{Y}_n, \mathcal{J}_n, \bigcup_{j=1}^n \mathcal{L}^1 \left((X_t^{(x_j)})_{0 \leq t < T_j} \right) \right)$$

has the same law as

$$\left(\mathcal{Y}_n, \mathcal{J}_n, \left\{ (\gamma(t))_{0 \leq t \leq T(\gamma)} \in \mathcal{L}_{1,L} \mid \gamma([0, T(\gamma)]) \cap \bigcup_{J \in \mathcal{J}_n} J \neq \emptyset \right\} \right)$$

Since $(\mathcal{Y}_\infty, \mathcal{J}_\infty)$ is by construction independent from $\left((X_t^{(x_j)})_{0 \leq t < T_j} \right)_{1 \leq j \leq n}$ conditionally on $(\mathcal{Y}_n, \mathcal{J}_n)$, we further get that the triple

$$\left(\mathcal{Y}_\infty, \mathcal{J}_\infty, \bigcup_{j=1}^n \mathcal{L}^1 \left((X_t^{(x_j)})_{0 \leq t < T_j} \right) \right)$$

has the same law as

$$\left(\mathcal{Y}_\infty, \mathcal{J}_\infty, \left\{ (\gamma(t))_{0 \leq t \leq T(\gamma)} \in \mathcal{L}_{1,L} \mid \gamma([0, T(\gamma)]) \cap \bigcup_{J \in \mathcal{J}_n} J \neq \emptyset \right\} \right)$$

Taking the limit of the third component as n tends to infinity we get that

$$\left(\mathcal{Y}_\infty, \mathcal{J}_\infty, \bigcup_{j \geq 1} \mathcal{L}^1 \left((X_t^{(x_j)})_{0 \leq t < T_j} \right) \right)$$

has the same law as

$$\left(\mathcal{Y}_\infty, \mathcal{J}_\infty, \left\{ (\gamma(t))_{0 \leq t \leq T(\gamma)} \in \mathcal{L}_{1,L} \mid \gamma([0, T(\gamma)]) \cap \bigcup_{J \in \mathcal{J}_\infty} J \neq \emptyset \right\} \right)$$

To conclude we need only to show that almost surely

$$\left\{ (\gamma(t))_{0 \leq t \leq T(\gamma)} \in \mathcal{L}_{1,L} \mid \gamma([0, T(\gamma)]) \cap \bigcup_{J \in \mathcal{J}_\infty} J \neq \emptyset \right\} = \mathcal{L}_{1,L}$$

The latter is equivalent to $\bigcup_{J \in \mathcal{J}_\infty} J$ being dense in I , which will be proved in the next section. \square

6.3. Determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$: Brownian case

In this section we will describe $(\mathcal{Y}_\infty, \mathcal{J}_\infty)$ in the Brownian case by giving the joint law of the point processes \mathcal{Y}_∞ and \mathcal{Z}_∞ . First we will study the case of a Brownian motion on a bounded interval (a, b) , killed upon hitting a or b , and without killing measure. Then we will study the case of the Brownian motion on \mathbb{R} with a non-zero Radon killing measure κ . We will write $(B_t^{(x_n)})_{0 \leq t < \zeta_n}$ instead of $(X_t^{(x_n)})_{0 \leq t < \zeta_n}$.

Proposition 6.2. — *In the case of a Brownian motion on a bounded interval (a, b) , killed upon hitting a or b , and without killing measure, \mathcal{Y}_∞ is deterministic and equals $\{a, b\}$ and \mathcal{Z}_∞ is made of a single point distributed uniformly on (a, b) .*

Proof. — For $n \geq 1$ we define $\tilde{x}_{n,0} < \tilde{x}_{n,1} < \dots < \tilde{x}_{n,n+1}$ as the family x_1, \dots, x_n, a, b ordered increasingly. According to this definition $\tilde{x}_{n,0} = a$ and $\tilde{x}_{n,n+1} = b$. As a convention we denote $\tilde{x}_{0,0} := a$ and $\tilde{x}_{0,1} := b$. For $n \geq 2$, one of the following situations may occur:

- $\mathcal{Y}_n = \{b\}$ and $\mathcal{J}_n = \{[\tilde{x}_{n,1}, b]\}$
- $\mathcal{Y}_n = \{a\}$ and $\mathcal{J}_n = \{[a, \tilde{x}_{n,n}]\}$

– $\mathcal{Y}_n = \{a, b\}$ and for some $j \in \{2, \dots, n\}$, $\mathcal{J}_n = \{[a, \tilde{x}_{n,j-1}], [\tilde{x}_{n,j}, b]\}$

In any case $(a, b) \setminus \left(\bigcup_{J \in \mathcal{J}_n} J\right)$ is an interval of form $(\tilde{x}_{n,j-1}, \tilde{x}_{n,j})$.

We set $\{J\}_0 = \emptyset$. Let $n \geq 1$. There is a $j \in \{1, \dots, n\}$ such that $x_n \in (\tilde{x}_{n-1,j-1}, \tilde{x}_{n-1,j})$. Conditionally on $(a, b) \setminus \left(\bigcup_{J \in \mathcal{J}_{n-1}} J\right) = (\tilde{x}_{n-1,j-1}, \tilde{x}_{n-1,j})$, the point $B_{T_n}^{(x_n)}$ equals $\tilde{x}_{n-1,j-1}$ with probability $\frac{\tilde{x}_{n-1,j} - x_n}{\tilde{x}_{n-1,j} - \tilde{x}_{n-1,j-1}}$ and $\tilde{x}_{n-1,j}$ with probability $\frac{x_n - \tilde{x}_{n-1,j-1}}{\tilde{x}_{n-1,j} - \tilde{x}_{n-1,j-1}}$. By induction we get that

$$\mathbb{P}\left((a, b) \setminus \left(\bigcup_{J \in \mathcal{J}_n} J\right) = (\tilde{x}_{n,j-1}, \tilde{x}_{n,j})\right) = \frac{\tilde{x}_{n,j} - \tilde{x}_{n,j-1}}{b - a}$$

Hence

$$\mathbb{P}(\mathcal{Y}_\infty = \{a\}) \leq \lim_{n \rightarrow +\infty} \mathbb{P}\left((a, b) \setminus \left(\bigcup_{J \in \mathcal{J}_n} J\right) = (\tilde{x}_{n,0}, \tilde{x}_{n,1})\right) = \lim_{n \rightarrow +\infty} \frac{\tilde{x}_{n,1} - \tilde{x}_{n,0}}{b - a} = 0$$

and similarly $\mathbb{P}(\mathcal{Y}_\infty = \{b\}) = 0$. Thus $\mathcal{Y}_\infty = \{a, b\}$. Almost surely for n large enough \mathcal{J}_n will be of form $\{[a, \tilde{x}_{n,j-1}], [\tilde{x}_{n,j}, b]\}$ for a random $j \in \{2, \dots, n\}$. We denote by $p_{n,1}^+$ respectively $p_{n,2}^-$ the random values of $\tilde{x}_{n,j-1}$ respectively $\tilde{x}_{n,j}$. Almost surely, neither of the non-decreasing sequence $(p_{n,1}^+)_n$ or non-increasing sequence of $(p_{n,2}^-)_n$ is stationary. This fact follows from the same argument according to which \mathcal{Y}_∞ is not reduced to one point. Moreover $p_{n,2}^- - p_{n,1}^+$, bounded by $\sup_{2 \leq j \leq n} (\tilde{x}_{n,j} - \tilde{x}_{n,j-1})$, converges to 0. It follows that a.s. \mathcal{Z}_∞ is reduced to one point, the common limit of $p_{n,1}^+$ and $p_{n,2}^-$. Finally if $\tilde{a} < \tilde{b}$ are two values taken by the sequence $(x_n)_{n \geq 1}$ then

$$\mathbb{P}(\mathcal{Z}_\infty \subseteq (\tilde{a}, \tilde{b})) = \frac{\tilde{b} - \tilde{a}}{b - a}$$

It follows that the unique point in \mathcal{Z}_∞ is distributed uniformly on (a, b) . \square

We consider now the case of the Brownian motion on \mathbb{R} with a non-zero Radon killing measure κ . $G(x, y) = u_\uparrow(x \wedge y)u_\downarrow(x \vee y)$ will be the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - \kappa$. The law of $(\mathcal{Y}_n, \mathcal{J}_n)$ may be expressed explicitly. Let Q_n be the cardinal of \mathcal{Y}_n . Let $Y_{n,1}, Y_{n,2}, \dots, Y_{n,Q(n)}$ be the points in \mathcal{Y}_n ordered in the increasing sense. Denote by $[p_{n,1}^-, p_{n,1}^+], [p_{n,2}^-, p_{n,2}^+], \dots, [p_{n,Q(n)}^-, p_{n,Q(n)}^+]$ the intervals in \mathcal{J}_n ordered in the increasing sense. For all $q \in \{1, \dots, Q_n\}$, $Y_{n,q} \in [p_{n,q}^-, p_{n,q}^+]$. It happens with positive probability that for some q , $p_{n,q}^- = p_{n,q}^+$ if one of the starting points x_1, \dots, x_n is an atom of κ . To compute recursively the joint law of above random variables we use the following facts: Given a killed Brownian path $(B_t^{(x)})_{0 \leq t < \zeta}$ starting from x , the distribution of $B_{\zeta^-}^{(x)}$ is $G(x, y)\kappa(dy)$ (see section 2.2). Given $a < x$, let T_a be the first time $B^{(x)}$ hits a . Then

$$\mathbb{P}_x(T_a \leq \zeta) = \frac{u_\downarrow(x)}{u_\downarrow(a)} = \frac{G(x, a)}{G(a, a)}$$

On the event $T_a > \zeta$, the distribution of $B_{\zeta^-}^{(x)}$ is:

$$(G(x, y) - \mathbb{P}_x(T_a \leq \zeta)G(a, y))1_{y>a}\kappa(dy) = \left(G(x, y) - \frac{G(x, a)G(a, y)}{G(a, a)} \right) 1_{y>a}\kappa(dy)$$

More generally, if $a < x < b$ and $\tilde{\zeta}$ is the first time $B^{(x)}$ gets either killed by the killing measure κ or hits a or b then

- The probability that $B_{\tilde{\zeta}^-}^{(x)} = a$ is:

$$\frac{u_\downarrow(a)u_\uparrow(x) - u_\downarrow(x)u_\uparrow(a)}{u_\downarrow(a)u_\uparrow(b) - u_\downarrow(b)u_\uparrow(a)} = \frac{\det \begin{pmatrix} G(x, b) & G(a, b) \\ G(a, x) & G(a, a) \end{pmatrix}}{\det \begin{pmatrix} G(b, b) & G(a, b) \\ G(a, b) & G(a, a) \end{pmatrix}}$$

- The probability that $B_{\tilde{\zeta}^-}^{(x)} = b$ is:

$$\frac{u_\downarrow(x)u_\uparrow(b) - u_\downarrow(b)u_\uparrow(x)}{u_\downarrow(a)u_\uparrow(b) - u_\downarrow(b)u_\uparrow(a)} = \frac{\det \begin{pmatrix} G(a, x) & G(a, b) \\ G(x, b) & G(b, b) \end{pmatrix}}{\det \begin{pmatrix} G(a, a) & G(a, b) \\ G(a, b) & G(b, b) \end{pmatrix}}$$

- The distribution of $B_{\tilde{\zeta}^-}^{(x)}$ on (a, b) is:

$$\frac{\det \begin{pmatrix} G(x, y) & G(a, y) & G(y, b) \\ G(a, x) & G(a, a) & G(a, b) \\ G(x, b) & G(a, b) & G(b, b) \end{pmatrix}}{\det \begin{pmatrix} G(a, a) & G(a, b) \\ G(a, b) & G(b, b) \end{pmatrix}} 1_{a<y<b}\kappa(dy)$$

Above expressions give the law of $(\mathcal{Y}_1, \mathcal{J}_1)$ and the law of $(\mathcal{Y}_{n+1}, \mathcal{J}_{n+1})$ conditionally on $(\mathcal{Y}_n, \mathcal{J}_n)$. By induction one can derive the law of $(\mathcal{Y}_n, \mathcal{J}_n)$. We will express it using a single identity involving a determinant. However this single identity may correspond to different configurations: We will divide the set of indices $\{1, \dots, Q_n\}$ in three categories E_n^- , E_n^+ and $E_n^{-,+}$ where for $q \in E_n^-$, $Y_{n,q} = p_{n,q}^-$, for $q \in E_n^+$, $Y_{n,q} = p_{n,q}^+$ and for $q \in E_n^{-,+}$, $p_{n,q}^- < Y_{n,q} < p_{n,q}^+$. For instance on the figure 2.a, $Q_5 = 3$, $E_5^- = \{3\}$, $E_5^+ = \{1\}$ and $E_5^{-,+} = \{2\}$.

Proposition 6.3. — Let $q \in \{1, \dots, n\}$. Let $(E_n^-, E_n^+, E_n^{-,+})$ be a partition of $\{1, \dots, q\}$:

$$\{1, \dots, q\} = E_n^- \amalg E_n^+ \amalg E_n^{-,+}$$

Let x^- be an increasing function from $E_n^- \amalg E_n^{-,+}$ to $\{x_1, \dots, x_n\}$ and x^+ an increasing function from $E_n^+ \amalg E_n^{-,+}$ to $\{x_1, \dots, x_n\}$. We assume that the sets $x^-(E_n^- \amalg E_n^{-,+})$ and $x^+(E_n^+ \amalg E_n^{-,+})$ are disjoint, that for every $i \in E_n^{-,+}$ $x^-(i) < x^+(i)$ and that for every $i \in E_n^- \amalg E_n^{-,+}$ and $j \in E_n^+ \amalg E_n^{-,+}$ such that $i \neq j$, $(x^+(j) - x^-(i))$ has the

same sign as $(j - i)$. Let $(\Delta_i)_{1 \leq i \leq n}$ be a family of disjoint bounded intervals each of which may be open, closed or semi-open such that for every $i < j$, $\max \Delta_i < \min \Delta_j$, that for every i , $\min \Delta_i \geq x^-(i)$ if $i \in E_n^- \amalg E_n^{-,+}$, $\max \Delta_i \leq x^+(i)$ if $i \in E_n^+ \amalg E_n^{-,+}$, and that for all i

$$x^-(i-1), x^+(i-1) < \min \Delta_i, \max \Delta_i < x^-(i+1), x^+(i+1)$$

where in the previous inequalities one should only consider the terms that are defined. Let $p_i^-(y_i)$ and $p_i^+(y_i)$ be the functions defined by: $p_i^-(y_i) = x^-(i)$ if $i \in E_n^- \amalg E_n^{-,+}$ and y_i otherwise. $p_i^+(y_i) = x^+(i)$ if $i \in E_n^+ \amalg E_n^{-,+}$ and y_i otherwise. Then

(6.3.1)

$$\begin{aligned} \mathbb{P}(Q_n = q, \forall i \in E_n^-, p_{n,i}^- = x^-(i), p_{n,i}^+ = Y_{n,i} \forall i \in E_n^+, p_{n,i}^- = x^+(i), p_{n,i}^+ = Y_{n,i}, \\ \forall i \in E_n^{-,+}, p_{n,i}^- = x^-(i), p_{n,i}^+ = x^+(i), \forall r \in \{1, \dots, q\}, Y_{n,r} \in \Delta_r) = \\ \int_{y_1 \in \Delta_1} \dots \int_{y_q \in \Delta_q} \det(G(p_i^-(y_i), p_j^+(y_j)))_{1 \leq i, j \leq q} \prod_{1 \leq r \leq q} \kappa(dy_i) \end{aligned}$$

$\det(G(p_i^-(y_i), p_j^+(y_j)))_{1 \leq i, j \leq q}$ may be rewritten as a simpler product:

$$(6.3.2) \quad G(p_1^-(y_1), p_1^+(y_1)) \prod_{1 \leq r \leq q-1} \left(G(p_{r+1}^-(y_{r+1}), p_{r+1}^+(y_{r+1})) \right. \\ \left. - \frac{G(p_r^-(y_r), p_{r+1}^+(y_{r+1}))G(p_r^+(y_r), p_{r+1}^-(y_{r+1}))}{G(p_r^-(y_r), p_r^+(y_r))} \right)$$

If σ is a permutation of $\{1, \dots, n\}$, then $(\mathcal{Y}_n(x_{\sigma(1)}, \dots, x_{\sigma(n)}), \mathcal{J}_n(x_{\sigma(1)}, \dots, x_{\sigma(n)}))$ has the same law as $(\mathcal{Y}_n(x_1, \dots, x_n), \mathcal{J}_n(x_1, \dots, x_n))$. Moreover, for any $n' > n$ and any permutation σ of $\{n+1, \dots, n'\}$, the law of $(\mathcal{Y}_{n'}(x_1, \dots, x_n, x_{\sigma(n+1)}, \dots, x_{\sigma(n')}), \mathcal{J}_{n'}(x_1, \dots, x_n, x_{\sigma(n+1)}, \dots, x_{\sigma(n')}))$ conditionally on $(\mathcal{Y}_n(x_1, \dots, x_n), \mathcal{J}_n(x_1, \dots, x_n))$ is the same as the law of $(\mathcal{Y}_{n'}(x_1, \dots, x_n, x_{n+1}, \dots, x_{n'}), \mathcal{J}_{n'}(x_1, \dots, x_n, x_{n+1}, \dots, x_{n'}))$ conditionally on $(\mathcal{Y}_n(x_1, \dots, x_n), \mathcal{J}_n(x_1, \dots, x_n))$.

Proof. — We will only give the sketch of a short proof. First let's check that the determinant $\det(G(p_i^-(y_i), p_j^+(y_j)))_{1 \leq i, j \leq q}$ may be indeed expressed as a product (6.3.2). We use the fact that for any $a < b < \tilde{a} < \tilde{b} \in \mathbb{R}$:

$$G(a, \tilde{b})G(b, \tilde{a}) = G(a, \tilde{a})G(b, \tilde{b}) = u_\uparrow(a)u_\uparrow(b)u_\downarrow(\tilde{a})u_\downarrow(\tilde{b})$$

By subtracting from the last line in the matrix $(G(p_i^-(y_i), p_j^+(y_j)))_{1 \leq i, j \leq q}$, which is $(G(p_q^-(y_q), p_j^+(y_j)))_{1 \leq j \leq q}$, the second to last line $(G(p_{q-1}^-(y_{q-1}), p_j^+(y_j)))_{1 \leq j \leq q}$ multiplied by $\frac{G(p_{q-1}^-(y_{q-1}), p_q^+(y_q))}{G(p_{q-1}^-(y_{q-1}), p_{q-1}^+(y_{q-1}))}$ we get zero for all coefficient on the last line,

except the diagonal one. Thus $\det (G(p_i^-(y_i), p_j^+(y_j)))_{1 \leq i, j \leq q}$ equals

$$\det(G(p_i^-(y_i), p_j^+(y_j)))_{1 \leq i, j \leq q-1} \times \left(G(p_q^-(y_q), p_q^+(y_q)) - \frac{G(p_{q-1}^-(y_{q-1}), p_q^+(y_q))G(p_{q-1}^+(y_{q-1}), p_q^-(y_q))}{G(p_{q-1}^-(y_{q-1}), p_{q-1}^+(y_{q-1}))} \right)$$

By induction we get (6.3.2).

Next step is to check that $(\mathcal{Y}_n(x_1, \dots, x_{n-2}, x_{n-1}, x_n), \mathcal{J}_n(x_1, \dots, x_{n-2}, x_{n-1}, x_n))$ and $(\mathcal{Y}_n(x_1, \dots, x_{n-2}, x_n, x_{n-1}), \mathcal{J}_n(x_1, \dots, x_{n-2}, x_n, x_{n-1}))$ have the same law conditionally on $(\mathcal{Y}_{n-2}(x_1, \dots, x_{n-2}), \mathcal{J}_{n-2}(x_1, \dots, x_{n-2}))$. This can be done using the explicit expressions for the conditional destitution of $B_{T_{n-1}^-}^{(x_{n-1})}$, $B_{T_n^-}^{(x_n)}$, $B_{T_{n-1}^-}^{(x_n)}$ and $B_{T_n^-}^{(x_{n-1})}$. This invariance by transposition of the two last starting points implies in turn all the invariances by permutation stated in the proposition.

From the invariance by permutation follows that one only needs to prove (6.3.1) in case $x_1 < x_2 < \dots < x_n$. In this case one can prove (6.3.1) by induction on n using the expression (6.3.2) for $\det (G(p_i^-(y_i), p_j^+(y_j)))_{1 \leq i, j \leq q}$. \square

The fact that the law of the tree obtained after n steps of Wilson's algorithm is invariant under permutations of the starting points (x_1, \dots, x_n) is something that is also satisfied in case of random walks on a true finite graph. The product (6.3.2) can be further rewritten as

$$(6.3.3) \quad u_\uparrow(p_1^-(y_1))u_\downarrow(p_q^+(y_q)) \prod_{1 \leq r \leq q-1} (u_\downarrow(p_r^+(y_r))u_\uparrow(p_{r+1}^-(y_{r+1})) - u_\uparrow(p_r^+(y_r))u_\downarrow(p_{r+1}^-(y_{r+1})))$$

Next we will show that \mathcal{Y}_∞ and \mathcal{Z}_∞ are a.s. discrete.

Lemma 6.4. — For all $n \geq 2$ and $q \in \{2, \dots, n\}$:

$$\mathbb{P}(\mathcal{Y}_\infty \cap (p_{n,q-1}^-, p_{n,q}^+) = \emptyset | p_{n,q-1}^-, p_{n,q}^+, Q_n \geq q) = \frac{2(p_{n,q}^+ - p_{n,q-1}^-)}{u_\downarrow(p_{n,q-1}^-)u_\uparrow(p_{n,q}^+) - u_\uparrow(p_{n,q-1}^-)u_\downarrow(p_{n,q}^+)}$$

Proof. — Let n and q be fixed. For $n' > n$, let

$$N(n') := \sharp(\{x_{n+1}, \dots, x_{n'}\} \cap (p_{n,q-1}^-, p_{n,q}^+))$$

and $\tilde{x}_{n',1} < \tilde{x}_{n',2} < \dots < \tilde{x}_{n',N(n')}$ the points of $\{x_{n+1}, \dots, x_{n'}\} \cap (p_{n,q-1}^-, p_{n,q}^+)$ ordered increasingly. Conventionally we define $\tilde{x}_{n',0} := p_{n,q-1}^-$ and $\tilde{x}_{n',N(n')+1} := p_{n,q}^+$. The condition $\mathcal{Y}_{n'} \cap (p_{n,q-1}^-, p_{n,q}^+) = \emptyset$ is satisfied if and only if for some $i \in \{1, 2, \dots, N(n') + 1\}$, necessarily unique, the following holds:

$$[p_{n,q-1}^-, \tilde{x}_{n',i-1}] \subseteq \bigcup_{J \in \mathcal{J}_m} J \text{ and } [\tilde{x}_{n',i}, p_{n,q}^+] \subseteq \bigcup_{J \in \mathcal{J}_{n'}} J$$

Thus

$$\begin{aligned} \mathbb{P}(\mathcal{Y}_{n'} \cap (p_{n,q-1}^-, p_{n,q}^+) = \emptyset | p_{n,q-1}^-, p_{n,q}^+, Q_n \geq q) = \\ \sum_{i=1}^{N(n')+1} \mathbb{P}\left([p_{n,q-1}^-, \tilde{x}_{n',i-1}] \subseteq \bigcup_{J \in \mathcal{J}_m} J, [\tilde{x}_{n',i}, p_{n,q}^+] \subseteq \bigcup_{J \in \mathcal{J}_{n'}} J \mid p_{n,q-1}^-, p_{n,q}^+, Q_n \geq q\right) \end{aligned}$$

Let $T_{n',i}$ be the first time $B^{(\tilde{x}_{n',i})}$ hits either $p_{n,q-1}^-$ or $p_{n,q}^+$ or gets killed by the killing measure κ . For $i \in \{1, 2, \dots, N(n') + 1\}$ let $T_{n',i,\tilde{x}_{n',i-1}}$ be the first time $B^{(\tilde{x}_{n',i})}$ hits $\tilde{x}_{n',i-1}$. Since the law of $(\mathcal{Y}_{n'}, \mathcal{J}_{n'})$ conditionally on $(\mathcal{Y}_n, \mathcal{J}_n)$ is invariant by permutation of points in $(x_{n+1}, \dots, x_{n'})$, we get that

$$\begin{aligned} \mathbb{P}\left([p_{n,q-1}^-, \tilde{x}_{n',i-1}] \subseteq \bigcup_{J \in \mathcal{J}_m} J, [\tilde{x}_{n',i}, p_{n,q}^+] \subseteq \bigcup_{J \in \mathcal{J}_{n'}} J \mid p_{n,q-1}^-, p_{n,q}^+, Q_n \geq q\right) = \\ \mathbb{P}\left(B_{T_{n',i-1}^-}^{(\tilde{x}_{n',i-1})} = p_{n,q-1}^-, B_{T_{n',i}^-}^{(\tilde{x}_{n',i})} = p_{n,q}^+, T_{n',i} < T_{n',i,\tilde{x}_{n',i-1}} \mid p_{n,q-1}^-, p_{n,q}^+, Q_n \geq q\right) = \\ \frac{u_\downarrow(\tilde{x}_{n',i-1})u_\uparrow(\tilde{x}_{n',i}) - u_\uparrow(\tilde{x}_{n',i-1})u_\downarrow(\tilde{x}_{n',i})}{u_\downarrow(p_{n,q-1}^-)u_\uparrow(p_{n,q}^+) - u_\uparrow(p_{n,q-1}^-)u_\downarrow(p_{n,q}^+)} \end{aligned}$$

It follows that

$$\begin{aligned} \mathbb{P}(\mathcal{Y}_{n'} \cap (p_{n,q-1}^-, p_{n,q}^+) = \emptyset | p_{n,q-1}^-, p_{n,q}^+, Q_n \geq q) = \\ \sum_{i=1}^{N(n')+1} \frac{u_\downarrow(\tilde{x}_{n',i-1})u_\uparrow(\tilde{x}_{n',i}) - u_\uparrow(\tilde{x}_{n',i-1})u_\downarrow(\tilde{x}_{n',i})}{u_\downarrow(p_{n,q-1}^-)u_\uparrow(p_{n,q}^+) - u_\uparrow(p_{n,q-1}^-)u_\downarrow(p_{n,q}^+)} \end{aligned}$$

If $\tilde{x}_{n',i-1}$ is close to $\tilde{x}_{n',i}$ then

$$\begin{aligned} u_\downarrow(\tilde{x}_{n',i-1})u_\uparrow(\tilde{x}_{n',i}) - u_\uparrow(\tilde{x}_{n',i-1})u_\downarrow(\tilde{x}_{n',i}) \\ = W(u_\downarrow, u_\uparrow)(\tilde{x}_{n',i-1})(\tilde{x}_{n',i} - \tilde{x}_{n',i-1}) + o(\tilde{x}_{n',i} - \tilde{x}_{n',i-1}) \\ = 2(\tilde{x}_{n',i} - \tilde{x}_{n',i-1}) + o(\tilde{x}_{n',i} - \tilde{x}_{n',i-1}) \end{aligned}$$

The sequence $(x_{n'})_{n' \geq n+1}$ is dense in $(p_{n,q-1}^-, p_{n,q}^+)$. Thus

$$\begin{aligned} \lim_{n' \rightarrow +\infty} \mathbb{P}(\mathcal{Y}_{n'} \cap (p_{n,q-1}^-, p_{n,q}^+) = \emptyset | p_{n,q-1}^-, p_{n,q}^+, Q_n \geq q) = \\ \frac{2(p_{n,q}^+ - p_{n,q-1}^-)}{u_\downarrow(p_{n,q-1}^-)u_\uparrow(p_{n,q}^+) - u_\uparrow(p_{n,q-1}^-)u_\downarrow(p_{n,q}^+)} \end{aligned}$$

□

Proposition 6.5. — *Let $a < b \in \mathbb{R}$. Then for all $n \geq 1$*

$$(6.3.4) \quad \mathbb{E}[\#\mathcal{Y}_n \cap [a, b]] \leq \int_{[a,b]} G(x, x) \kappa(dx)$$

It follows that a.s. for all $a < b \in \mathbb{R}$, $\mathcal{Y}_\infty \cap [a, b]$ is finite.

Proof. — Let $\tilde{a} < \tilde{b} \in [a, b]$ where \tilde{a} is close to \tilde{b} . We will first show that for all $n \geq 1$

$$(6.3.5) \quad \mathbb{P}\left(\mathcal{Y}_n \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) \leq \int_{[\tilde{a}, \tilde{b}]} G(x, x) \kappa(dx) + o(\tilde{b} - \tilde{a})$$

where $o(\tilde{b} - \tilde{a})$ is uniform over \tilde{a} and \tilde{b} close to each other in $[a, b]$. Then we will deduce (6.3.4) by partitioning the interval $[a, b]$ in small subintervals $[\tilde{a}, \tilde{b}]$ and approximating the expected number of points in $[\tilde{a}, \tilde{b}]$ by the probability of presence of one point. Let $n \geq 1$. Then

$$\mathbb{P}\left(\mathcal{Y}_n(x_1, \dots, x_n) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) \leq \mathbb{P}\left(\mathcal{Y}_{n+2}(x_1, \dots, x_n, \tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right)$$

Since the law of \mathcal{Y}_{n+2} is invariant by permutation of the starting points:

$$\mathbb{P}\left(\mathcal{Y}_{n+2}(x_1, \dots, x_n, \tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) = \mathbb{P}\left(\mathcal{Y}_{n+2}(\tilde{a}, \tilde{b}, x_1, \dots, x_n) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right)$$

But

$$(6.3.6) \quad \mathbb{P}\left(\mathcal{Y}_{n+2}(\tilde{a}, \tilde{b}, x_1, \dots, x_n) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) = \mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) \\ + \mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] = \emptyset, \mathcal{Y}_{n+2}(\tilde{a}, \tilde{b}, x_1, \dots, x_n) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right)$$

We start Wilson's algorithm by launching first $B^{(\tilde{a})}$ starting from \tilde{a} followed by $B^{(\tilde{b})}$ starting \tilde{b} . Then

$$\mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) = \mathbb{P}\left(B_{T_1}^{(\tilde{a})} \in [\tilde{a}, \tilde{b}]\right) + \mathbb{P}\left(B_{T_1}^{(\tilde{a})} \notin [\tilde{a}, \tilde{b}], B_{T_1}^{(\tilde{a})} \leq \tilde{a}, B_{T_2}^{(\tilde{b})} \in [\tilde{a}, \tilde{b}]\right)$$

Applying proposition 6.3 we get that

$$\mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) = \\ \int_{x \in [\tilde{a}, \tilde{b}]} \left(G(\tilde{a}, x) + \int_{y \leq \tilde{a}} (G(y, \tilde{a})G(x, \tilde{b}) - G(y, \tilde{b})G(\tilde{a}, x)) \kappa(dy) \right) \kappa(dx)$$

For $x \in \mathbb{R}$, let $T_{1,x}$ be the first time $B^{(\tilde{a})}$ hits x . Then

$$G(\tilde{a}, x) + \int_{y \leq \tilde{a}} (G(y, \tilde{a})G(x, \tilde{b}) - G(y, \tilde{b})G(\tilde{a}, x)) \kappa(dy) = \\ G(x, x) \left(\mathbb{P}(T_1 \geq T_{1,x}) + \frac{G(x, \tilde{b})}{G(x, x)} \mathbb{P}(T_1 < T_{1,x}, B_{T_1}^{(\tilde{a})} \leq \tilde{a}) \right) \leq G(x, x)$$

Thus

$$(6.3.7) \quad \mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) \leq \int_{[\tilde{a}, \tilde{b}]} G(x, x) \kappa(dx)$$

Further

$$\mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] = \emptyset, \mathcal{Y}_{n+2}(\tilde{a}, \tilde{b}, x_1, \dots, x_n) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) \leq \\ \mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] = \emptyset, \mathcal{Y}_\infty(\tilde{a}, \tilde{b}, (x_j)_{j \geq 1}) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right)$$

Applying lemma 6.4 and proposition 6.3 we get that

$$\begin{aligned}
& \mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] = \emptyset, \mathcal{Y}_\infty(\tilde{a}, \tilde{b}, (x_j)_{j \geq 1}) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) = \\
& \mathbb{P}\left(\mathcal{Y}_\infty(\tilde{a}, \tilde{b}, (x_j)_{j \geq 1}) \cap [\tilde{a}, \tilde{b}] \neq \emptyset \mid \mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] = \emptyset\right) \times \mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] = \emptyset\right) \\
& = \left(1 - \frac{2(\tilde{b} - \tilde{a})}{u_\downarrow(\tilde{a})u_\uparrow(\tilde{b}) - u_\uparrow(\tilde{a})u_\downarrow(\tilde{b})}\right) \\
& \quad \times \int_{y \leq a} \int_{z \geq b} \mathbf{1}_{y, z \notin [\tilde{a}, \tilde{b}]} \det \begin{pmatrix} G(y, a) & G(y, z) \\ G(a, b) & G(b, z) \end{pmatrix} \kappa(dy) \kappa(dz) \\
& \leq (u_\downarrow(\tilde{a})u_\uparrow(\tilde{b}) - u_\uparrow(\tilde{a})u_\downarrow(\tilde{b}) - 2(\tilde{b} - \tilde{a})) \int_{y \leq b} u_\uparrow(y) \kappa(dy) \int_{z \geq a} u_\downarrow(z) \kappa(dz)
\end{aligned}$$

But

$$u_\downarrow(\tilde{a})u_\uparrow(\tilde{b}) - u_\uparrow(\tilde{a})u_\downarrow(\tilde{b}) - 2(\tilde{b} - \tilde{a}) = o(\tilde{b} - \tilde{a})$$

Thus

$$(6.3.8) \quad \mathbb{P}\left(\mathcal{Y}_2(\tilde{a}, \tilde{b}) \cap [\tilde{a}, \tilde{b}] = \emptyset, \mathcal{Y}_{n+2}(\tilde{a}, \tilde{b}, x_1, \dots, x_n) \cap [\tilde{a}, \tilde{b}] \neq \emptyset\right) = o(\tilde{b} - \tilde{a})$$

Combining (6.3.6), (6.3.7) and (6.3.8) we get (6.3.5).

Now for $j \in \mathbb{N}^*$ and $i \in \{1, \dots, 2^j\}$ consider the intervals $\Delta_{i,j}$ defined by

$$\Delta_{i,j} = \begin{cases} [a + (i-1)2^{-j}(b-a), a + i2^{-j}(b-a)] & \text{if } i \leq 2^j - 1 \\ [a + (1-2^{-j})(b-a), b] & \text{if } i = 2^j \end{cases}$$

Then $\mathbb{E}[\#\mathcal{Y}_n \cap [a, b)]$ is the increasing limit of $\sum_{i=1}^{2^j} \mathbb{P}(\mathcal{Y}_n \cap \Delta_{i,j} \neq \emptyset)$. But

$$\sum_{i=1}^{2^j} \mathbb{P}(\mathcal{Y}_n \cap \Delta_{i,j} \neq \emptyset) \leq \sum_{i=1}^{2^j} \int_{\Delta_{i,j}} G(x, x) \kappa(dx) + 2^j o(2^{-j})$$

(6.3.4) follows. Since (6.3.4) holds for all n , it also holds at the limit when n tends to $+\infty$. This implies that $\mathcal{Y}_\infty \cap [a, b]$ is a.s. finite. \square

Proposition 6.6. — *Almost surely all the intervals in \mathcal{J}_∞ are open.*

Proof. — We need only to show that for any $n \geq 1$ and $q \in \{1, \dots, n\}$

$$(6.3.9) \quad \mathbb{P}(Q_n \geq q, \forall n' \geq n, \min(t_{n,n'}([p_{n,q}^-, p_{n,q}^+])) = p_{n,q}^-) = 0$$

and

$$\mathbb{P}(Q_n \geq q, \forall n' \geq n, \max(t_{n,n'}([p_{n,q}^-, p_{n,q}^+])) = p_{q,n}^+) = 0$$

Let n and q be fixed. We will show (6.3.9). We will also assume that $q \geq 2$. The proof is similar if $q = 1$. We need to show that a.s. the following conditional probability converges to 0:

$$\lim_{n' \rightarrow +\infty} \mathbb{P}(\min(t_{n,n'}([p_{n,q}^-, p_{n,q}^+])) = p_{n,q}^- \mid \mathcal{Y}_n, \mathcal{J}_n, Q_n \geq q) = 0$$

We recall that for $n'' \geq n+1$, $B^{(x_{n''})}$ is a Brownian motion starting from $x_{n''}$ and it is independent from $(\mathcal{Y}_n, \mathcal{J}_n)$. Let $T_{n'', p_{q,n}^-}$ be the first time it hits $p_{q,n}^-$ and $\tilde{T}_{n''}$ the first time it either hits $\bigcup_{J \in \mathcal{J}_n} J$ or gets killed by the killing measure κ . Since the law of $(\mathcal{Y}_{n'}, \mathcal{J}_{n'})$ conditionally on $(\mathcal{Y}_n, \mathcal{J}_n)$ is invariant by permutation of points in $(x_{n+1}, \dots, x_{n'})$, we get that

$$\begin{aligned} \mathbb{P}(\min(\iota_{n,n'}([p_{n,q}^-, p_{n,q}^+])) &= p_{n,q}^- | (\mathcal{Y}_n, \mathcal{J}_n), Q_n \geq q) \\ &\leq \inf_{n+1 \leq n'' \leq n'} 1 - 1_{p_{n,q-1}^+ < x_{n''} < p_{n,q}^-} \mathbb{P}(\tilde{T}_{n''} = T_{n'', p_{q,n}^-} | p_{n,q-1}^+, p_{n,q}^-, Q_n \geq q) \end{aligned}$$

But $\mathbb{P}(\tilde{T}_{n''} = T_{n'', p_{q,n}^-} | p_{n,q-1}^+, p_{n,q}^-)$ is close to 1 if $x_{n''}$ is close enough to $p_{n,q}^-$. There is always a subsequence of $(x_{n''})_{n'' \geq n+1}$ made of points in $(p_{n,q-1}^+, p_{n,q}^-)$ which converges to $p_{n,q}^-$. It follows that

$$\inf_{n'' \geq n+1} 1 - 1_{p_{n,q-1}^+ < x_{n''} < p_{n,q}^-} \mathbb{P}(\tilde{T}_{n''} = T_{n'', p_{q,n}^-} | p_{n,q-1}^+, p_{n,q}^-, Q_n \geq q) = 0$$

which concludes the proof. \square

From proposition 6.6 follows that \mathcal{Z}_∞ is closed. Moreover it does not contain any of the points of the sequence $(x_n)_{n \geq 1}$. Since the sequence $(x_n)_{n \geq 1}$ is everywhere dense, the connected components of \mathcal{Z}_∞ are single points. One can see that

- If $y < \tilde{y}$ are two consecutive points in \mathcal{Y}_∞ then $\sharp(\mathcal{Z}_\infty \cap (y, \tilde{y})) = 1$.
- If \mathcal{Y}_∞ is bounded from below and $y = \min \mathcal{Y}_\infty$ then $\mathcal{Z}_\infty \cap (-\infty, y] = \emptyset$.
- If \mathcal{Y}_∞ is bounded from above and $y = \max \mathcal{Y}_\infty$ then $\mathcal{Z}_\infty \cap [y, +\infty) = \emptyset$.

See figure 2.b. The set \mathcal{Z}_∞ may be empty, which for instance happens almost surely if κ is a Dirac measure. For $n \geq 1$ we define

$$\mathcal{Z}_n := \left\{ \frac{p_{n,q-1}^- + p_{n,q}^+}{2} \mid 2 \leq q \leq Q_n \right\}$$

We will write $\mathcal{Z}_n(x_1, \dots, x_n)$ and $\mathcal{Z}_\infty((x_n)_{n \geq 1})$ whenever we need to emphasize the dependence on the starting points.

Proposition 6.7. — *The law of $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ does not depend on the starting points $(x_n)_{n \geq 1}$.*

Proof. — Let $(\tilde{x}_n)_{n \geq 1}$ be another sequence of pairwise disjoint points in \mathbb{R} . We will show that the sequence $(\mathcal{Y}_{2n}(x_1, \dots, x_n, \tilde{x}_1, \dots, \tilde{x}_n), \mathcal{Z}_{2n}(x_1, \dots, x_n, \tilde{x}_1, \dots, \tilde{x}_n))$ converges in law to $(\mathcal{Y}_\infty((x_n)_{n \geq 1}), \mathcal{Z}_\infty((x_n)_{n \geq 1}))$ and that $(\mathcal{Y}_{2n}(\tilde{x}_1, \dots, \tilde{x}_n, x_1, \dots, x_n), \mathcal{Z}_{2n}(\tilde{x}_1, \dots, \tilde{x}_n, x_1, \dots, x_n))$ converges to $(\mathcal{Y}_\infty((\tilde{x}_n)_{n \geq 1}), \mathcal{Z}_\infty((\tilde{x}_n)_{n \geq 1}))$. Since the two couples of point processes $(\mathcal{Y}_{2n}(x_1, \dots, x_n, \tilde{x}_1, \dots, \tilde{x}_n), \mathcal{Z}_{2n}(x_1, \dots, x_n, \tilde{x}_1, \dots, \tilde{x}_n))$ and $(\mathcal{Y}_{2n}(\tilde{x}_1, \dots, \tilde{x}_n, x_1, \dots, x_n), \mathcal{Z}_{2n}(\tilde{x}_1, \dots, \tilde{x}_n, x_1, \dots, x_n))$ have the same law, this will finish the proof.

For the convergence in law we will use the topology of uniform convergence on compact sets of collections of points in \mathbb{R} . It can be defined using the following

metric: Let d_H be the Hausdorff metric on compact subsets of \mathbb{R} . One may use the metric d_{PP} on point processes:

$$d_{PP}(\mathcal{X}, \tilde{\mathcal{X}}) := d_H(\tan^{-1}(\mathcal{X}) \cup \{-1, 1\}, \tan^{-1}(\tilde{\mathcal{X}}) \cup \{-1, 1\})$$

In order to simplify the notations we will write:

$$(\mathcal{Y}_n, \mathcal{Z}_n) := (\mathcal{Y}_n(x_1, \dots, x_n), \mathcal{Z}_n(x_1, \dots, x_n))$$

$$(\mathcal{Y}_\infty, \mathcal{Z}_\infty) := (\mathcal{Y}_\infty((x_n)_{n \geq 1}), \mathcal{Z}_\infty((x_n)_{n \geq 1}))$$

$$(\tilde{\mathcal{Y}}_{2n}, \tilde{\mathcal{Z}}_{2n}) := (\mathcal{Y}_{2n}(x_1, \dots, x_n, \tilde{x}_1, \dots, \tilde{x}_n), \mathcal{Z}_{2n}(x_1, \dots, x_n, \tilde{x}_1, \dots, \tilde{x}_n))$$

We can construct $((\mathcal{Y}_n, \mathcal{Z}_n))_{n \geq 1}$, $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and $((\tilde{\mathcal{Y}}_{2n}, \tilde{\mathcal{Z}}_{2n}))_{n \geq 1}$ on the same probability space using independent Brownian motions starting from the points in $(x_n)_{n \geq 1}$ and $(\tilde{x}_n)_{n \geq 1}$ and killed by the measure κ . We construct the sequence $((\mathcal{Y}_n, \mathcal{Z}_n))_{n \geq 1}$ using the Wilson's algorithm described in introduction. This way $\mathcal{Y}_n \subseteq \mathcal{Y}_{n+1}$ and $\mathcal{Y}_\infty = \bigcup_{n \geq 1} \mathcal{Y}_n$. In order to construct $\tilde{\mathcal{Y}}_{2n}$, we first construct \mathcal{Y}_n and then continue the Wilson's algorithm using the Brownian motions starting from $\tilde{x}_1, \dots, \tilde{x}_n$. This way $\mathcal{Y}_n \subseteq \tilde{\mathcal{Y}}_{2n}$ but not necessarily $\tilde{\mathcal{Y}}_{2n} \subseteq \tilde{\mathcal{Y}}_{2(n+1)}$.

Let $C > 0$ and $\varepsilon \in (0, \frac{C}{2})$. Let $\delta \in (0, 1)$, δ small. There is $N \in \mathbb{N}^*$ such that

$$\mathbb{P}(\mathcal{Y}_N \cap [-C, C] = \mathcal{Y}_\infty \cap [-C, C]) \geq 1 - \delta$$

There is $\varepsilon' \in (0, \varepsilon)$ such that for all $a < b \in [-C, C]$ satisfying $b - a \leq \varepsilon'$ the following holds:

$$1 - \frac{2(b-a)}{u_\downarrow(a)u_\uparrow(b) - u_\uparrow(a)u_\downarrow(b)} \leq \frac{\delta}{N}$$

There is $N' \geq N$ such that with probability $1 - 2\delta$ the following two conditions hold:

$$(6.3.10) \quad \mathcal{Y}_{N'} \cap [-C, C] = \mathcal{Y}_\infty \cap [-C, C]$$

$$(6.3.11) \quad \text{Leb}([-C, C] \setminus \bigcup_{J \in \mathcal{J}_{N'}} J) \leq \varepsilon'$$

We define the following two random variables:

$$K^- := \min_{J \in \mathcal{J}_{N'}, J \subseteq [-C, C]} (\min J) \quad K^+ := \max_{J \in \mathcal{J}_{N'}, J \subseteq [-C, C]} (\max J)$$

If (6.3.11) holds, then $[-\frac{C}{2}, \frac{C}{2}] \subseteq [K^-, K^+]$. If (6.3.10) and (6.3.11) hold then for $n \geq N'$, $[K^-, K^+] \setminus \bigcup_{J \in \mathcal{J}_n} J$ is made of at most N intervals, each of length at most ε' . Consider the following condition on $\tilde{\mathcal{Y}}_{2n}$:

$$(6.3.12) \quad \tilde{\mathcal{Y}}_{2n} \cap [K^-, K^+] = \mathcal{Y}_n \cap [K^-, K^+]$$

Applying lemma 6.4 we get that for all $n \geq N'$

$$\mathbb{P}(\tilde{\mathcal{Y}}_{2n} \text{ satisfies (6.3.12)} \mid (6.3.10) \text{ and (6.3.11) hold}) \geq 1 - \delta$$

This implies that for all $n \geq N'$

$$\mathbb{P}(\tilde{\mathcal{Y}}_{2n} \text{ satisfies (6.3.12), and (6.3.10) and (6.3.11) hold.}) \geq 1 - 3\delta$$

Let $n \geq N'$. On the event when (6.3.10) and (6.3.11) hold and $\tilde{\mathcal{Y}}_{2n}$ satisfies (6.3.12), which happens with probability at least $1 - 3\delta$, the following is true:

$$\begin{aligned} & - \tilde{\mathcal{Y}}_{2n} \cap [K^-, K^+] = \mathcal{Y}_\infty \cap [K^-, K^+] \\ & - d_H(\tilde{\mathcal{Z}}_{2n} \cap [K^-, K^+], \mathcal{Z}_\infty \cap [K^-, K^+]) \leq \varepsilon \end{aligned}$$

In particular with probability at least $1 - 3\delta$

$$\begin{aligned} & - d_{PP}(\tilde{\mathcal{Y}}_{2n}, \mathcal{Y}_\infty) \leq 1 - \tan^{-1}\left(\frac{C}{2}\right) \\ & - d_H(\tilde{\mathcal{Z}}_{2n}, \mathcal{Z}_\infty) \leq \varepsilon + (1 - \tan^{-1}\left(\frac{C}{2}\right)) \end{aligned}$$

Since C is arbitrary large and ε and δ are arbitrary small, this implies that $(\tilde{\mathcal{Y}}_{2n}, \tilde{\mathcal{Z}}_{2n})$ converges in law as $n \rightarrow +\infty$ to $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$. \square

Next we identify the law of \mathcal{Y}_∞ as a determinantal fermionic point process. For generalities on this processes see [11], chapter 4, and [28].

Proposition 6.8. — *Let $n \geq 1$ and $a_1 < b_1 < a_2 < b_2 < \dots < a_n < b_n \in \mathbb{R}$. Then*

$$(6.3.13) \quad \mathbb{E} \left[\prod_{r=1}^n \#(\mathcal{Y}_\infty \cap [a_r, b_r]) \right] = \int_{[a_1, b_1]} \dots \int_{[a_n, b_n]} \det(G(y_i, y_j))_{1 \leq i, j \leq n} \prod_{r=1}^n \kappa(dy_r)$$

In other words \mathcal{Y}_∞ is a determinantal point process on \mathbb{R} with reference measure κ and determinantal kernel G .

Proof. — Consider points $\tilde{a}_r < \tilde{b}_r \in [a_r, b_r]$ for $r \in \{1, \dots, n\}$. We will show that

$$(6.3.14) \quad \mathbb{P} \left(\forall r \in \{1, \dots, n\}, \mathcal{Y}_\infty \cap [\tilde{a}_r, \tilde{b}_r] \neq \emptyset \right) =$$

$$\int_{[\tilde{a}_1, \tilde{b}_1]} \dots \int_{[\tilde{a}_n, \tilde{b}_n]} \det(G(y_i, y_j))_{1 \leq i, j \leq n} \prod_{r=1}^n \kappa(dy_r)$$

$$+ \left(\sum_{r=1}^n O(\tilde{b}_r - \tilde{a}_r) \right) \times \prod_{r=1}^n \kappa([\tilde{a}_r, \tilde{b}_r]) + \sum_{\substack{E \subseteq \{1, \dots, n\} \\ E \neq \emptyset}} \prod_{r \in E} o(\tilde{b}_r - \tilde{a}_r) \prod_{r \notin E} \kappa([\tilde{a}_r, \tilde{b}_r])$$

where the quantities $O(\tilde{b}_r - \tilde{a}_r)$ and $o(\tilde{b}_r - \tilde{a}_r)$ are uniform over $\tilde{a}_r < \tilde{b}_r \in [a_r, b_r]$, \tilde{a}_r close to \tilde{b}_r . From (6.3.14) one deduces (6.3.13) by splitting the intervals $[a_r, b_r]$ in small subintervals and approximating the number of points in $\mathcal{Y}_\infty \cap [a_r, b_r]$ by the number of subintervals of $[a_r, b_r]$ that contain a point in \mathcal{Y}_∞ .

As the law of \mathcal{Y}_∞ does not depend on the choice of everywhere dense sequence of starting points, we will assume that the first $2n$ starting points in Wilson's algorithm are in order $\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_n, \tilde{b}_n$. We will show that for all non-empty subsets E of

$\{1, \dots, n\}$

$$(6.3.15) \quad \mathbb{P} \left(\forall r \in E, \mathcal{Y}_{2n} \cap [\tilde{a}_r, \tilde{b}_r] = \emptyset, \mathcal{Y}_\infty \cap [\tilde{a}_r, \tilde{b}_r] \neq \emptyset, \forall r \notin E, \mathcal{Y}_{2n} \cap [\tilde{a}_r, \tilde{b}_r] \neq \emptyset \right) \\ = \prod_{r \in E} o(\tilde{b}_r - \tilde{a}_r) \prod_{r \notin E} \kappa([\tilde{a}_r, \tilde{b}_r])$$

Further we will show that for any $r_0 \in \{1, \dots, n\}$

(6.3.16)

$$\mathbb{P}(\forall r \in \{1, \dots, n\}, \mathcal{Y}_{2n} \cap [\tilde{a}_r, \tilde{b}_r] \neq \emptyset, [\tilde{a}_{r_0}, \tilde{b}_{r_0}] \not\subseteq \bigcup_{J \in \mathcal{J}_{2n}} J) = O(\tilde{b}_{r_0} - \tilde{a}_{r_0}) \prod_{r=1}^n \kappa([\tilde{a}_r, \tilde{b}_r])$$

If for all $r \in \{1, \dots, n\}$, $\mathcal{Y}_{2n} \cap [\tilde{a}_r, \tilde{b}_r] \neq \emptyset$ and $[\tilde{a}_r, \tilde{b}_r] \subseteq \bigcup_{J \in \mathcal{J}_{2n}} J$ then necessarily $Q_{2n} = n$ and $\mathcal{J}_{2n} = \{[\tilde{a}_r, \tilde{b}_r] | 1 \leq r \leq n\}$. We will use the fact that according to (6.3.1)

$$(6.3.17) \quad \mathbb{P} \left(Q_{2n} = n, \mathcal{J}_{2n} = \{[\tilde{a}_r, \tilde{b}_r] | 1 \leq r \leq n\} \right) \\ = \int_{[\tilde{a}_1, \tilde{b}_1]} \dots \int_{[\tilde{a}_n, \tilde{b}_n]} \det \left(G^{\tilde{a}_i, \tilde{b}_j} \right)_{1 \leq i, j \leq n} \prod_{r=1}^n \kappa(dy_r) \\ = \int_{[\tilde{a}_1, \tilde{b}_1]} \dots \int_{[\tilde{a}_n, \tilde{b}_n]} \det (G(y_i, y_j))_{1 \leq i, j \leq n} \prod_{r=1}^n \kappa(dy_r) + \left(\sum_{r=1}^n O(\tilde{b}_r - \tilde{a}_r) \right) \times \prod_{r=1}^n \kappa([\tilde{a}_r, \tilde{b}_r])$$

Let's show (6.3.15). A closed expression of the probability in (6.3.15) can be computed using (6.3.1) and lemma 6.4. Since many different configurations (different values of Q_{2n} and configurations of \mathcal{J}_{2n}) contribute to the probability in (6.3.15), we won't give the closed expression and only give the estimates. Let E be a non-empty subset of $\{1, \dots, n\}$. If $r \notin E$, then the condition $\mathcal{Y}_{2n} \cap [\tilde{a}_r, \tilde{b}_r] \neq \emptyset$ contributes by a factor $O(\kappa([\tilde{a}_r, \tilde{b}_r]))$ to the probability in (6.3.15). If $r \in E$, then the two conditions $\mathcal{Y}_{2n} \cap [\tilde{a}_r, \tilde{b}_r] = \emptyset$ and $\mathcal{Y}_\infty \cap [\tilde{a}_r, \tilde{b}_r] \neq \emptyset$ imply that $(\tilde{a}_r, \tilde{b}_r) \cap \bigcup_{J \in \mathcal{J}_{2n}} J = \emptyset$. According to the identity (6.3.3), the condition $(\tilde{a}_r, \tilde{b}_r) \cap \bigcup_{J \in \mathcal{J}_{2n}} J = \emptyset$ contributes to the probability in (6.3.15) by a factor

$$O(u_\downarrow(\tilde{a}_r)u_\uparrow(\tilde{b}_r) - u_\uparrow(\tilde{a}_r)u_\downarrow(\tilde{b}_r)) = O(\tilde{b}_r - \tilde{a}_r)$$

According to the lemma 6.4, the additional condition $\mathcal{Y}_\infty \cap [\tilde{a}_r, \tilde{b}_r] \neq \emptyset$ contributes to the probability in (6.3.15) by a factor

$$1 - \frac{2(\tilde{b}_r - \tilde{a}_r)}{u_\downarrow(\tilde{a}_r)u_\uparrow(\tilde{b}_r) - u_\uparrow(\tilde{a}_r)u_\downarrow(\tilde{b}_r)} = o(1)$$

(6.3.15) follows.

We deal now with (6.3.16). As in the previous case, the condition that for all $r \in \{1, \dots, n\}$, $\mathcal{Y}_{2n} \cap [\tilde{a}_r, \tilde{b}_r] \neq \emptyset$ contributes by a factor $O\left(\prod_{r=1}^n \kappa([\tilde{a}_r, \tilde{b}_r])\right)$ to the probability in (6.3.16). The condition $[\tilde{a}_{r_0}, \tilde{b}_{r_0}] \not\subseteq \bigcup_{J \in \mathcal{J}_{2n}} J$ implies that there is

$i \in \{2, \dots, Q_{2n}\}$ such that $\tilde{a}_{r_0} < p_{2n,i-1}^+ < p_{2n,i}^- < \tilde{b}_{r_0}$. As previously, this contributes by a factor $O(\tilde{b}_{r_0} - \tilde{a}_{r_0})$ to the probability. Combining (6.3.15), (6.3.16) and (6.3.17) yields (6.3.14). \square

Let \mathfrak{G}_κ be the following operator defined for functions in $\mathbb{L}^2(d\kappa)$ with compact support:

$$(\mathfrak{G}_\kappa f)(x) := \int_{\mathbb{R}} G(x, y) f(y) \kappa(dy)$$

A standard condition for a determinantal point process with kernel G relative to the measure κ to be well defined is \mathfrak{G}_κ to be positive semi-definite, contracting and locally trace class. We explain why this is true. Let f be a compactly supported $\mathbb{L}^2(d\kappa)$ function. Then the weak second derivative of $\mathfrak{G}_\kappa f$ is

$$d\left(\frac{d(\mathfrak{G}_\kappa f)}{dx}\right) = 2(\mathfrak{G}_\kappa f - f)d\kappa$$

$\mathfrak{G}_\kappa f$ and $\frac{d(\mathfrak{G}_\kappa f)}{dx}$ are square-integrable and

$$(6.3.18) \quad \begin{aligned} \int_{\mathbb{R}} (\mathfrak{G}_\kappa f) f d\kappa &= \int_{\mathbb{R}} (\mathfrak{G}_\kappa f)^2 d\kappa + \frac{1}{2} \int_{\mathbb{R}} (\mathfrak{G}_\kappa f) d\left(\frac{d(\mathfrak{G}_\kappa f)}{dx}\right) \\ &= \int_{\mathbb{R}} (\mathfrak{G}_\kappa f)^2 d\kappa + \frac{1}{2} \int_{\mathbb{R}} \left(\frac{d(\mathfrak{G}_\kappa f)}{dx}\right)^2 dx \end{aligned}$$

Identity (6.3.18) shows that \mathfrak{G}_κ is positive semi-definite. It also shows that $\int_{\mathbb{R}} (\mathfrak{G}_\kappa f)^2 d\kappa \leq \int_{\mathbb{R}} (\mathfrak{G}_\kappa f) f d\kappa$, which implies that \mathfrak{G}_κ is contracting and hence can be continuously extended to a contraction of the whole space $\mathbb{L}^2(d\kappa)$. \mathfrak{G}_κ is locally trace class because it is positive semi-definite and its functional kernel is continuous (see theorem 2.12 in [27], chapter 2).

Next we give a criterion for \mathcal{Y}_∞ to be finite or just to be finite in the neighbourhood of either $+\infty$ or $-\infty$.

Proposition 6.9. — *If $\int_{(0,+\infty)} x\kappa(dx) < +\infty$ then almost surely $\#\mathcal{Y}_\infty \cap (0, +\infty)$ is finite. Moreover*

$$(6.3.19) \quad \mathbb{E}[\#\mathcal{Y}_\infty \cap (0, +\infty)] = \int_{(0,+\infty)} G(x, x) \kappa(dx) < +\infty$$

If $\int_{(0,+\infty)} x\kappa(dx) = +\infty$ then almost surely $\#\mathcal{Y}_\infty \cap (0, +\infty) = +\infty$. In general, for all $a \in \mathbb{R}$

$$(6.3.20) \quad \mathbb{P}(\mathcal{Y}_\infty \cap (a, +\infty) = \emptyset) = u_\downarrow(+\infty) \int_{(-\infty, a]} u_\uparrow(x) \kappa(dx)$$

Similarly, if $\int_{\mathbb{R}} |x| \kappa(dx) < +\infty$ then a.s. $\#\mathcal{Y}_\infty$ is finite and

$$\mathbb{E}[\#\mathcal{Y}_\infty] = \int_{\mathbb{R}} G(x, x) \kappa(dx) < +\infty$$

If $\int_{\mathbb{R}} |x| \kappa(dx) = +\infty$ then a.s. $\#\mathcal{Y}_\infty = +\infty$.

Proof. — We need only to deal with the finiteness of $\sharp(\mathcal{Y}_\infty \cap (0, +\infty))$. If $\int_{(0, +\infty)} x k(dx) < +\infty$ then (6.3.19) holds according to 2.3 and hence $\sharp(\mathcal{Y}_\infty \cap (0, +\infty))$ is finite a.s.

We will prove (6.3.20). If $\int_{(0, +\infty)} x k(dx) = +\infty$ then according 2.3 $u_\downarrow(+\infty) > 0$ and thus $\sharp(\mathcal{Y}_\infty \cap (0, +\infty)) = +\infty$ a.s. Let $a < b \in \mathbb{R}$. We assume that the two first starting points in Wilson's algorithm are a and b . Then

(6.3.21)

$$\begin{aligned} \mathbb{P}(\mathcal{Y}_\infty \cap (a, b] = \emptyset) &= \mathbb{P}(B_{T_1}^{(a)} > b) + \mathbb{P}(B_{T_1}^{(a)} \leq a, B_{T_2}^{(b)} = a) \\ &= \mathbb{P}(B_{\zeta_1}^{(a)} > b) + \mathbb{P}(B_{\zeta_1}^{(a)} \leq a) \times \mathbb{P}(B^{(b)} \text{ hits } a \text{ before time } \zeta_2) \\ &= \int_{(b, +\infty)} G(a, x) \kappa(dx) + \left(\int_{(-\infty, a]} G(a, x) \kappa(dx) \right) \times \frac{u_\downarrow(b)}{u_\downarrow(a)} \\ &= \int_{(b, +\infty)} G(a, x) \kappa(dx) + u_\downarrow(b) \int_{(-\infty, a]} u_\uparrow(x) \kappa(dx) \end{aligned}$$

Letting b go to $+\infty$ in (6.3.21) gives (6.3.20). \square

Next we will show that \mathcal{Z}_∞ is a determinantal point process with kernel \mathcal{K} relatively to the Lebesgue measure where

$$\begin{aligned} \mathcal{K}(y, z) &:= -\frac{1}{2} \frac{du_\uparrow}{dx}((y \wedge z)^+) \frac{u_\downarrow}{dz}((y \vee z)^-) \\ &= 2 \int_{(-\infty, y \wedge z]} u_\uparrow(x) \kappa(dx) \times \int_{[y \vee z, +\infty)} u_\downarrow(x) \kappa(dx) \end{aligned}$$

Proposition 6.10. — Let $n \geq 1$ and $a_1 < b_1 < a_2 < b_2 < \dots < a_n < b_n \in \mathbb{R}$. Then

$$(6.3.22) \quad \mathbb{E} \left[\prod_{r=1}^n \sharp(\mathcal{Z}_\infty \cap (a_r, b_r)) \right] = \int_{(a_1, b_1)} \dots \int_{(a_n, b_n)} \det(\mathcal{K}(z_i, z_j))_{1 \leq i, j \leq n} \prod_{r=1}^n dz_r$$

If for $r \in \{1, 2, \dots, n\}$, $\kappa(\{a_r\}) = \kappa(\{b_r\}) = 0$ then

(6.3.23)

$$\mathbb{P}(\forall r \in \{1, 2, \dots, n\}, \sharp(\mathcal{Z}_\infty \cap (a_r, b_r)) = 1) = \det(\mathcal{K}(a_i, b_j))_{1 \leq i, j \leq n} \times \prod_{r=1}^n (b_r - a_r)$$

Proof. — We will only prove (6.3.23). (6.3.22) can be deduced from (6.3.23) by dividing the intervals (a_r, b_r) in small subintervals and approximating the expected number of points in these subintervals by the probability to have one single point per subinterval. Observe that if the measure κ has atoms then \mathcal{K} is not continuous. Yet $z \mapsto \frac{du_\uparrow}{dx}(z^+)$ is right-continuous and $z \mapsto \frac{du_\downarrow}{dx}(z^-)$ is left-continuous. So the approximation can still be done.

Consider the Wilson's algorithm where the $2n$ first starting points are in order $a_1, b_1, a_2, b_2, \dots, a_n, b_n$. Then

$$(6.3.24) \quad \mathbb{P}(\forall r \in \{1, 2, \dots, n\}, \#(\mathcal{Z}_\infty \cap (a_r, b_r)) = 1) = \\ \mathbb{P}\left(\forall r \in \{1, 2, \dots, n\}, (a_r, b_r) \subseteq \mathbb{R} \setminus \bigcup_{J \in \mathcal{J}_{2n}} J, (a_r, b_r) \cap \mathcal{Y}_\infty = \emptyset\right)$$

Applying lemma 6.4 we get that (6.3.24) equals

$$(6.3.25) \quad \mathbb{P}\left(\forall r \in \{1, 2, \dots, n\}, (a_r, b_r) \subseteq \mathbb{R} \setminus \bigcup_{J \in \mathcal{J}_{2n}} J\right) \times \prod_{r=1}^n \frac{2(b_r - a_r)}{u_\downarrow(a_r)u_\uparrow(b_r) - u_\uparrow(a_r)u_\downarrow(b_r)}$$

Further

$$(6.3.26) \quad \mathbb{P}\left(\forall r \in \{1, 2, \dots, n\}, (a_r, b_r) \subseteq \mathbb{R} \setminus \bigcup_{J \in \mathcal{J}_{2n}} J\right) = \\ \mathbb{P}\left(B_{T_1^-}^{(a_1)} \leq a_1, B_{T_{2n}^-}^{(b_n)} \geq b_n, \forall r \in \{1, \dots, n-1\}, b_r \leq B_{T_{2r}^-}^{(b_r)} \leq B_{T_{2r+1}^-}^{(a_{r+1})} \leq a_{r+1}\right)$$

Applying (6.3.1) and (6.3.3) we get that (6.3.26) equals

$$(6.3.27) \quad \prod_{r=1}^n (u_\downarrow(a_r)u_\uparrow(b_r) - u_\uparrow(a_r)u_\downarrow(b_r)) \times \int_{(-\infty, a_1]} u_\uparrow(y_1)\kappa(dy_1) \times \int_{[b_n, +\infty)} u_\downarrow(z_n)\kappa(dy_n) \\ \times \prod_{r=1}^{n-1} \left(\kappa([b_r, a_{r+1}]) + \int_{b_r \leq y_r < \tilde{y}_r \leq a_{r+1}} (u_\downarrow(y_r)u_\uparrow(\tilde{y}_r) - u_\uparrow(y_r)u_\downarrow(\tilde{y}_r))\kappa(dy_r)\kappa(d\tilde{y}_r) \right)$$

But

$$(6.3.28) \quad \int_{b_r \leq y_r < \tilde{y}_r \leq a_{r+1}} u_\downarrow(y_r)u_\uparrow(\tilde{y}_r)\kappa(dy_r)\kappa(d\tilde{y}_r) \\ = \frac{1}{2} \int_{b_r \leq y_r \leq a_{r+1}} u_\downarrow(y_r) \left(\frac{du_\uparrow}{dx}(a_{r+1}) - \frac{du_\uparrow}{dx}(y_r^+) \right) \kappa(dy_r) \\ = \frac{1}{4} \left(\frac{du_\downarrow}{dx}(a_{r+1}) - \frac{du_\downarrow}{dx}(b_r) \right) \frac{du_\uparrow}{dx}(a_{r+1}) - \frac{1}{2} \int_{b_r \leq y_r \leq a_{r+1}} u_\downarrow(y_r) \frac{du_\uparrow}{dx}(y_r^+) \kappa(dy_r)$$

and

$$(6.3.29) \quad - \int_{b_r \leq y_r < \tilde{y}_r \leq a_{r+1}} u_\uparrow(y_r)u_\downarrow(\tilde{y}_r)\kappa(dy_r)\kappa(d\tilde{y}_r) \\ = -\frac{1}{2} \int_{b_r \leq y_r \leq a_{r+1}} u_\uparrow(y_r) \left(\frac{du_\downarrow}{dx}(a_{r+1}) - \frac{du_\downarrow}{dx}(y_r^+) \right) \kappa(dy_r) \\ = -\frac{1}{4} \left(\frac{du_\uparrow}{dx}(a_{r+1}) - \frac{du_\uparrow}{dx}(b_r) \right) \frac{du_\downarrow}{dx}(a_{r+1}) + \frac{1}{2} \int_{b_r \leq y_r \leq a_{r+1}} u_\uparrow(y_r) \frac{du_\downarrow}{dx}(y_r) \kappa(dy_r)$$

Combining (6.3.28) and (6.3.29) we get that

$$\begin{aligned}
& \int_{b_r \leq y_r < \tilde{y}_r \leq a_{r+1}} (u_\downarrow(y_r)u_\uparrow(\tilde{y}_r) - u_\uparrow(y_r)u_\downarrow(\tilde{y}_r)) \kappa(dy_r) \kappa(d\tilde{y}_r) \\
&= \frac{1}{4} \left(\frac{du_\uparrow}{dx}(b_r) \frac{du_\downarrow}{dx}(a_{r+1}) - \frac{du_\downarrow}{dx}(b_r) \frac{du_\uparrow}{dx}(a_{r+1}) \right) \\
&\quad - \frac{1}{2} \int_{b_r \leq y_r \leq a_{r+1}} \left(u_\downarrow(y_r) \frac{du_\uparrow}{dx}(y_r^+) - u_\uparrow(y_r) \frac{du_\downarrow}{dx}(y_r^+) \right) \kappa(dy_r) \\
&= \frac{1}{4} \left(\frac{du_\uparrow}{dx}(b_r) \frac{du_\downarrow}{dx}(a_{r+1}) - \frac{du_\downarrow}{dx}(b_r) \frac{du_\uparrow}{dx}(a_{r+1}) \right) - \kappa([b_r, a_{r+1}])
\end{aligned}$$

It follows that (6.3.27) equals

$$\begin{aligned}
(6.3.30) \quad & \prod_{r=1}^n (u_\downarrow(a_r)u_\uparrow(b_r) - u_\uparrow(a_r)u_\downarrow(b_r)) \times \left(-\frac{1}{4} \frac{du_\uparrow}{dx}(a_1) \frac{du_\downarrow}{dx}(b_n) \right) \\
& \times \prod_{r=1}^{n-1} \left(\frac{1}{4} \left(\frac{du_\uparrow}{dx}(b_r) \frac{du_\downarrow}{dx}(a_{r+1}) - \frac{du_\downarrow}{dx}(b_r) \frac{du_\uparrow}{dx}(a_{r+1}) \right) \right) \\
& = \frac{1}{2^n} \prod_{r=1}^n (u_\downarrow(a_r)u_\uparrow(b_r) - u_\uparrow(a_r)u_\downarrow(b_r)) \times \det(\mathcal{K}(a_i, b_j))_{1 \leq i, j \leq n}
\end{aligned}$$

(6.3.25) together with (6.3.30) gives (6.3.23). \square

To see that the operator induced by the kernel \mathcal{K} on $\mathbb{L}^2(L\text{eb})$ is positive semi-definite, one can check that for any \mathbb{L}^2 function f with compact support

$$\int_{\mathbb{R}^2} f(y)\mathcal{K}(y, z)f(z)dydz = \int_{\mathbb{R}^2} G(\tilde{y}, \tilde{z}) \left(\int_{\tilde{y}}^{\tilde{z}} f(x)dx \right)^2 \kappa(d\tilde{y})\kappa(d\tilde{z})$$

To see that \mathcal{K} induces a contraction one can check that for any \mathcal{C}^1 function f with compact support

$$\int_{\mathbb{R}^2} f(y)\mathcal{K}(y, z)f(z)dydz = \int_{\mathbb{R}} f(x)^2 dx - \frac{1}{2} \int_{\mathbb{R}^2} \frac{df}{dx}(\tilde{y})G(\tilde{y}, \tilde{z}) \frac{df}{dx}(\tilde{z})d\tilde{y}d\tilde{z}$$

and that $\int_{\mathbb{R}^2} \frac{df}{dx}(\tilde{y})G(\tilde{y}, \tilde{z}) \frac{df}{dx}(\tilde{z})d\tilde{y}d\tilde{z} \geq 0$.

The determinantal kernels G and \mathcal{K} both satisfy the following relation: for any $x \leq y \leq z \in \mathbb{R}$

$$(6.3.31) \quad G(x, y)G(y, z) = G(x, z)G(y, y) \quad \mathcal{K}(x, y)\mathcal{K}(y, z) = \mathcal{K}(x, z)\mathcal{K}(y, y)$$

For $x \in \mathbb{R}$ and $y, z > x$, we define

$$(6.3.32) \quad G^{(x \times)}(y, z) := G(y, z) - \frac{G(x, y)G(x, z)}{G(x, x)} \quad \mathcal{K}^{(x \triangleright)}(y, z) := \mathcal{K}(y, z) - \frac{\mathcal{K}(x, y)\mathcal{K}(x, z)}{\mathcal{K}(x, x)}$$

Relation (6.3.31) ensures that $\det(G(y_i, y_j))_{1 \leq i, j \leq n}$ and $\det(\mathcal{K}(z_i, z_j))_{1 \leq i, j \leq n}$ can be factorised as follows: If $y_1 < y_2 < \dots < y_n$ then

$$(6.3.33) \quad \det(G(y_i, y_j))_{1 \leq i, j \leq n} = G(y_1, y_1) \prod_{r=2}^n G^{(y_{r-1} \times)}(y_r, y_r)$$

If $z_1 < z_2 < \dots < z_n$ then

$$(6.3.34) \quad \det(\mathcal{K}(z_i, z_j))_{1 \leq i, j \leq n} = \mathcal{K}(z_1, z_1) \prod_{r=2}^n \mathcal{K}^{(z_{r-1} \triangleright)}(z_r, z_r)$$

The relations (6.3.31) or equivalently the factorisations (6.3.33) and (6.3.34) imply that the spacings between consecutive points of \mathcal{Y}_∞ respectively \mathcal{Z}_∞ are independent, that is to say conditionally on \mathcal{Y}_∞ having a point at y_0 , the position of the next higher point y is independent on $\mathcal{Y}_\infty \cap (-\infty, y_0)$, and similarly for \mathcal{Z}_∞ ([28], section 2.4). Conditionally on $y_0 \in \mathcal{Y}_\infty$ the distribution of its higher neighbour in \mathcal{Y}_∞ is of the form $f_G(y_0, y)\kappa(dy)$. Similarly denote $f_{\mathcal{K}}(z_0, z)dz$ the distribution between two consecutive points in \mathcal{Z}_∞ conditionally on z_0 be the lowest one. Following relations relate $G^{(y_0 \times)}(y, y)$ respectively $\mathcal{K}^{(z_0 \triangleright)}(z, z)$ to f_G respectively $f_{\mathcal{K}}$:

$$\begin{aligned} G^{(y_0 \times)}(y, y) &= f_G(y_0, y) \\ &+ \sum_{j \geq 2} \int_{y_0 < y_1 < \dots < y_{j-1} < y} f_G(y_0, y_1) f_G(y_1, y_2) \dots f_G(y_{j-1}, y) \kappa(dy_1) \dots \kappa(dy_{j-1}) \end{aligned}$$

$$\begin{aligned} \mathcal{K}^{(z_0 \triangleright)}(z, z) &= f_{\mathcal{K}}(z_0, z) \\ &+ \sum_{j \geq 2} \int_{z_0 < z_1 < \dots < z_{j-1} < z} f_{\mathcal{K}}(z_0, z_1) f_{\mathcal{K}}(z_1, z_2) \dots f_{\mathcal{K}}(z_{j-1}, z) dz_1 \dots dz_{j-1} \end{aligned}$$

If $\int_{(0, +\infty)} x k(dx) < +\infty$, i.e. $\mathcal{Y}_\infty \cap (0, +\infty)$ a.s. finite, then $\int_{(y_0, +\infty)} f_G(y_0, y)\kappa(dy) < 1$ and $\int_{z_0}^{+\infty} f_{\mathcal{K}}(z_0, z)dz < 1$.

Given a couple of interwoven point processes $(\mathcal{Y}, \mathcal{Z})$ on \mathbb{R} such that between any two consecutive point in \mathcal{Y} lies one single point of \mathcal{Z} and such that for any J bounded subinterval of \mathbb{R} $\mathcal{Y} \cap J$ satisfies the constraint

$$\mathbb{E}[\sharp(\mathcal{Y} \cap J)] < +\infty$$

the joint distribution of $(\mathcal{Y}, \mathcal{Z})$ can be fully described using the family of measures $(M_n(\mathcal{Y}, \mathcal{Z}))_{n \geq 0}$ defined by

$$\int_{\mathbb{R}} f(y_0) M_0(\mathcal{Y}, \mathcal{Z})(dy_0) = \mathbb{E} \left[\sum_{y_0 \in \mathcal{Y}} f(y_0) \right]$$

$$\begin{aligned} & \int_{y_0 < z_1 < y_1 < \dots < z_n < y_n} f(y_0, z_1, y_1, \dots, z_n, y_n) M_n(\mathcal{Y}, \mathcal{Z})(dy_0, dz_1, dy_1, \dots, dz_n, dy_n) \\ &= \mathbb{E} \left[\sum_{\substack{y_0, \dots, y_n \\ n+1 \text{ consecutive points in } \mathcal{Y} \\ z_1, \dots, z_n \in \mathcal{Z} \\ y_0 < z_1 < y_1 < \dots < z_n < y_n}} f(y_0, z_1, y_1, \dots, z_n, y_n) \right] \end{aligned}$$

$M_n(\mathcal{Y}, \mathcal{Z})(dy_0, dz_1, dy_1, \dots, dz_n, dy_n)$ is the infinitesimal probability for y_0, y_1, \dots, y_n being $n+1$ consecutive points in \mathcal{Y} and z_1, \dots, z_n being the n points in \mathcal{Z} separating them. In case of $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$, $M_0(\mathcal{Y}_\infty, \mathcal{Z}_\infty)(dy_0) = G(y_0, y_0)\kappa(dy_0)$.

Proposition 6.11. — For $n \geq 1$

(6.3.35)

$$\begin{aligned} M_n(\mathcal{Y}_\infty, \mathcal{Z}_\infty)(dy_0, dz_1, \dots, dz_n, dy_n) &= 2^n u_\uparrow(y_0) u_\downarrow(y_n) \kappa(dy_0) dz_1 \dots dz_n \kappa(dy_n) \\ &= 2^n G(y_0, y_n) \kappa(dy_0) dz_1 \dots dz_n \kappa(dy_n) \end{aligned}$$

Moreover

$$f_G(y_0, y) = 2(y - y_0) \frac{u_\downarrow(y)}{u_\downarrow(y_0)} \quad \kappa(dy) \text{ — almost everywhere}$$

$$f_\kappa(z_0, z) = 2\kappa((z_0, z)) \left(\frac{du_\downarrow}{dx}(z_0) \right)^{-1} \frac{du_\downarrow}{dx}(z) \quad dz \text{ — almost everywhere}$$

The distribution on \mathcal{Z}_∞ conditionally on \mathcal{Y}_∞ is the following: given two consecutive points $y_1 < y_2$ in \mathcal{Y}_∞ , then the point of \mathcal{Z}_∞ lying between them is distributed uniformly on (y_1, y_2) and independently on the behaviour of \mathcal{Z}_∞ on $(-\infty, y_1) \cup (y_2, +\infty)$. The distribution on \mathcal{Y}_∞ conditionally on \mathcal{Z}_∞ is the following: given two consecutive points $z_1 < z_2$ in \mathcal{Z}_∞ , then the point of \mathcal{Y}_∞ lying between them is distributed on (z_1, z_2) according the measure $1_{z_1 < y < z_2} \frac{\kappa(dy)}{\kappa((z_1, z_2))}$ and independently on the behaviour of \mathcal{Y}_∞ on $(-\infty, z_1) \cup (z_2, +\infty)$. If $\int_{(-\infty, 0)} |x| \kappa(dx) < +\infty$, then $\min \mathcal{Y}_\infty$ is distributed conditionally on \mathcal{Z}_∞ according to the measure $1_{y < \min \mathcal{Z}_\infty} \frac{\kappa(dy)}{\kappa((-\infty, \min \mathcal{Z}_\infty))}$ and it is independent on the behaviour of \mathcal{Y}_∞ on $(-\infty, \min \mathcal{Z}_\infty)$. Similarly for the distribution of $\max \mathcal{Y}_\infty$ conditionally on $\max \mathcal{Z}_\infty$ if $\int_{(0, +\infty)} x \kappa(dx) < +\infty$.

Proof. — Let $a_0 < b_0 < \tilde{a}_1 < \tilde{b}_1 < a_1 < b_1 < \dots < \tilde{a}_n < \tilde{b}_n < a_n < b_n \in \mathbb{R}$. Let $\mathcal{C}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, a_1, b_1, \dots, \tilde{a}_n, \tilde{b}_n, a_n, b_n)$ corresponding to the following conditions:

- $\mathcal{Y}_\infty \cap [a_0, b_0] \neq \emptyset$, $\mathcal{Y}_\infty \cap [a_n, b_n] \neq \emptyset$
- $\forall r \in \{1, \dots, n\}$, $\#(\mathcal{Y}_\infty \cap [a_r, b_r]) = 1$
- $\forall r \in \{1, \dots, n\}$, $\#(\mathcal{Z}_\infty \cap (\tilde{a}_r, \tilde{b}_r)) = 1$
- $\forall r \in \{0, \dots, n-1\}$, $(\mathcal{Y}_\infty \cup \mathcal{Z}_\infty) \cap (b_r, \tilde{a}_{r+1}) = \emptyset$, $(\mathcal{Y}_\infty \cup \mathcal{Z}_\infty) \cap [\tilde{b}_r, a_{r+1}) = \emptyset$

We will compute the probability of $\mathcal{C}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, a_1, b_1, \dots, \tilde{a}_n, \tilde{b}_n, a_n, b_n)$. Consider that we execute the Wilson's algorithm where the $2n$ first starting points are $\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_n, \tilde{b}_n$. The only configurations that contribute to the studied event are

those where $B_{T_1^-}^{(\tilde{a}_1)} \in [a_0, b_0]$, $B_{T_{2^n}^-}^{(\tilde{b}_n)} \in [a_n, b_n]$ and for $r \in \{1, \dots, n-1\}$, $B_{T_{2^r}^-}^{(\tilde{b}_r)} = B_{T_{2^{r+1}}^-}^{(\tilde{a}_{r+1})} \in [a_{r+1}, b_{r+1}]$. We further need that for $r \in \{1, \dots, n\}$, $\mathcal{Y}_\infty \cap (\tilde{a}_r, \tilde{b}_r) = \emptyset$. Thus applying (6.3.1), (6.3.3) and lemma 6.4 we get the probability of the event $\mathcal{C}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, a_1, b_1, \dots, \tilde{a}_n, \tilde{b}_n, a_n, b_n)$ equals

$$\int_{[a_0, b_0]} u_\uparrow(y_0) \kappa(dy_0) \times \int_{[a_n, b_n]} u_\downarrow(y_n) \kappa(dy_n) \times \prod_{r=1}^{n-1} \kappa([a_r, b_r]) \times \prod_{r=1}^n 2(\tilde{b}_r - \tilde{a}_r)$$

The above probability also equals $M_n(\mathcal{Y}_\infty, \mathcal{Z}_\infty)([a_0, b_0] \times [\tilde{a}_1, \tilde{b}_1] \times [a_1, b_1] \times \dots \times [\tilde{a}_n, \tilde{b}_n] \times [a_n, b_n])$ and gives the expression of (6.3.35). To get the expressions of f_G and $f_{\mathcal{K}}$ just observe that

$$G(y_0, y_0) f_G(y_0, y) \kappa(dy_0) \kappa(dy) = M_1([y_0, y_0 + dy_0] \times (y_0, y) \times [y, y + dy])$$

$$\mathcal{K}(z_0, z_0) f_{\mathcal{K}}(z_0, z) dz_0 dz = M_3((-\infty, z_0) \times [z_0, z_0 + dz_0] \times (z_0, z) \times [z, z + dz] \times (z, +\infty))$$

Expression (6.3.35) gives also the law of \mathcal{Z}_∞ conditionally on \mathcal{Y}_∞ and the law of \mathcal{Y}_∞ conditionally on \mathcal{Z}_∞ , except for the possible extremal points of \mathcal{Y}_∞ . Let's deal with the distribution of $\max \mathcal{Y}_\infty$ conditionally on $\max \mathcal{Z}_\infty$ in case $\int_{(0, +\infty)} x \kappa(dx) < +\infty$. Again according to (6.3.35), conditionally on $z_0 \in \mathcal{Z}_\infty$, the distribution of $\min \mathcal{Y}_\infty \cap (z_0, +\infty)$ is proportional to $1_{y > z_0} u_\downarrow(y) \kappa(dy)$. To obtain the distribution $\max \mathcal{Y}_\infty$ conditionally on $\max \mathcal{Z}_\infty$, one must weight $u_\downarrow(y)$ by $1 - \int_{\tilde{y} > y} f_G(y, \tilde{y}) \kappa(d\tilde{y})$, i.e. the probability of not having any point in \mathcal{Y}_∞ consecutive to y . But

$$\begin{aligned} \int_{\tilde{y} > y} f_G(y, \tilde{y}) \kappa(d\tilde{y}) &= 2 \int_{\tilde{y} > y} (\tilde{y} - y) \frac{u_\downarrow(\tilde{y})}{u_\downarrow(y)} \kappa(d\tilde{y}) \\ &= \lim_{\tilde{y} \rightarrow +\infty} \frac{\tilde{y} - y}{u_\downarrow(y)} \frac{du_\downarrow}{dx}(\tilde{y}^+) - \frac{1}{u_\downarrow(y)} \int_{\tilde{y} > y} \frac{du_\downarrow}{dx}(\tilde{y}^+) d\tilde{y} \end{aligned}$$

But

$$(\tilde{y} - y) \frac{du_\downarrow}{dx}(\tilde{y}^+) = (\tilde{y} - y) \int_{(\tilde{y}, +\infty)} 2u_\downarrow(x) \kappa(dx) \leq 2 \int_{(\tilde{y}, +\infty)} (x - y) u_\downarrow(x) \kappa(dx) \rightarrow 0$$

It follows that:

$$\int_{\tilde{y} > y} f_G(y, \tilde{y}) \kappa(d\tilde{y}) = -\frac{1}{u_\downarrow(y)} \int_{\tilde{y} > y} \frac{du_\downarrow}{dx}(\tilde{y}^+) d\tilde{y} = 1 - \frac{u_\downarrow(+\infty)}{u_\downarrow(y)}$$

Thus $1_{y > z_0} u_\downarrow(y) (1 - \int_{\tilde{y} > y} f_G(y, \tilde{y}) \kappa(d\tilde{y})) \kappa(dy)$ is simply proportional to $1_{y > z_0} \kappa(dy)$. \square

Proposition 6.12. — In case $\int_{\mathbb{R}} |x| \kappa(dx) < +\infty$

$$\mathbb{P}(\#\mathcal{Y}_\infty = 1) = u_\uparrow(-\infty) u_\downarrow(+\infty) \kappa(\mathbb{R})$$

Conditionally on $\#\mathcal{Y}_\infty = 1$ the unique point in \mathcal{Y}_∞ is distributed according $\frac{\kappa(dy)}{\kappa(\mathbb{R})}$.

Proof. — The distribution of the unique point y_0 of \mathcal{Y}_∞ on the event $\#\mathcal{Y}_\infty = 1$ is given by the following sieve identity:

$$\begin{aligned} & \left(G(y_0, y_0) - \int_{y_{-1} < y_0} G(y_{-1}, y_{-1}) f_G(y_{-1}, y_0) \kappa(dy_{-1}) \right. \\ & \quad - \int_{y_1 > y_0} G(y_0, y_0) f_G(y_0, y_1) \kappa(dy_1) \\ & \quad \left. + \int_{y_{-1} < y_0} \int_{y_1 > y_0} G(y_{-1}, y_{-1}) f_G(y_{-1}, y_0) f_G(y_0, y_1) \kappa(dy_{-1}) \kappa(dy_1) \right) \kappa(dy_0) \end{aligned}$$

It is the infinitesimal probability of \mathcal{Y}_∞ having a point at y_0 minus the probability of having a point at y_0 and an other lower, minus the probability of having a point at y_0 and an other higher, plus the probability of having a point at y_0 surrounded by two neighbours on both sides. The identity can be further factorized as

$$\begin{aligned} & \left(u_\uparrow(y_0) - 2 \int_{y_{-1} < y_0} (y_0 - y_{-1}) u_\uparrow(y_{-1}) \kappa(dy_{-1}) \right) \\ & \quad \times \left(u_\downarrow(y_0) - 2 \int_{y_1 > y_0} (y_1 - y_0) u_\downarrow(y_1) \kappa(dy_1) \right) \times \kappa(dy_0) \end{aligned}$$

According to the calculation done in the proof of proposition 6.11 this the above equals $u_\uparrow(-\infty)u_\downarrow(+\infty)\kappa(dy_0)$. \square

Now let's describe $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ in two particular cases. If the killing rate is uniform, that is $\kappa(dy) = cdy$ where c is constant, then

$$cf_G(x_0, x) = f_\kappa(x_0, x) = 2c(x - x_0)e^{-\sqrt{2c}(x-x_0)}$$

Both the spacings of \mathcal{Y}_∞ and \mathcal{Z}_∞ are i.i.d. gamma-2 variables with mean $\sqrt{\frac{2}{c}}$. Actually the union $\mathcal{Y}_\infty \cup \mathcal{Z}_\infty$ is a Poisson point process with intensity $\sqrt{2c}dx$. If the killing measure is of form $\kappa = c \sum_{j \in \mathbb{Z}} \delta_j$ where c is constant, then again the spacings between consecutive points in \mathcal{Y}_∞ are i.i.d random variables, this time integer valued. Let N_2 be a random variable with same distribution as this spacings. For any $j \in \mathbb{N}$

$$\mathbb{P}(N_2 = j) = 2cj(1 + \sqrt{2c})^{-j}$$

N_2 can be written as $N_2 = N_1 + \tilde{N}_1 - 1$ where N_1 and \tilde{N}_1 are two independent geometric variables of parameter $(1 + \sqrt{2c})^{-1}$. Actually, if $y_0 < y$ are two consecutive points in \mathcal{Y}_∞ and z the point of \mathcal{Z}_∞ lying between them, then conditionally on y_0 , $(\lfloor z \rfloor - y_0, y - \lfloor z \rfloor)$ has the same law as $(N_1 - 1, \tilde{N}_1)$. Moreover $\{\lfloor z \rfloor | z \in \mathcal{Z}_\infty\}$ has the same law as \mathcal{Y}_∞ .

6.4. Determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$: general case

Let I be an open subinterval of \mathbb{R} and L be the generator of a transient diffusion on I of form $L = \frac{1}{m(x)} \frac{d}{dx} \left(\frac{1}{w(x)} \frac{d}{dx} \right) - \kappa$ with zero Dirichlet boundary conditions on

∂I with sample path denoted $(X_t)_{0 \leq t < \zeta}$. We will describe, without proof, the law of $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ in this generic case. It can be derived in the same way as it was done in the previous section. Let G be the Green's function of L relatively to the measure $m(y)dy$, factorisable as $G(x, y) = u_\uparrow(x \wedge y)u_\downarrow(x \vee y)$.

Proposition 6.13. — \mathcal{Y}_∞ and \mathcal{Z}_∞ are a.s. discrete point processes. Let ∂I be the boundary of I in $\mathbb{R} \cup \{-\infty, +\infty\}$. Almost surely

$$\mathcal{Y}_\infty \cap \partial I = \{y \in \partial I | \mathbb{P}(X_{\zeta^-} = y) > 0\}$$

If $\kappa \neq 0$, the points in $\mathcal{Y}_\infty \cap I$ are a determinantal point process with determinantal kernel $G(x, y)$ relatively the reference measure $m(y)\kappa(dy)$. \mathcal{Z}_∞ is a determinantal point process on I with determinantal kernel

$$\frac{du_\uparrow}{dx}((y \wedge z)^+) \frac{du_\downarrow}{dx}((y \vee z)^-)$$

relatively to the reference measure $\frac{dz}{w(z)}$. Given two consecutive points $y_1 < y_2$ in \mathcal{Y}_∞ , then the point of \mathcal{Z}_∞ lying between them is distributed according to the measure $1_{y_1 < z < y_2} \frac{w(z)dz}{\int_{(y_1, y_2)} w(a)da}$ and independently on the behaviour of \mathcal{Z}_∞ on $(-\infty, y_1) \cup (y_2, +\infty)$. Given two consecutive points $z_1 < z_2$ in \mathcal{Z}_∞ , then the point of \mathcal{Y}_∞ lying between them is distributed on (z_1, z_2) according the measure $1_{z_1 < y < z_2} \frac{m(y)\kappa(dy)}{\int_{(z_1, z_2)} m(q)\kappa(dq)}$ and independently on the behaviour of \mathcal{Y}_∞ on $(-\infty, z_1) \cup (z_2, +\infty)$.

CHAPTER 7

MONOTONE COUPLINGS FOR THE POINT PROCESSES $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$

7.1. Conditioning

In this chapter we will deal with monotone coupling for the determinantal point processes \mathcal{Y}_∞ and \mathcal{Z}_∞ introduced in chapter 6. We will restrict to the Brownian case. Consider two different killing measures κ and $\tilde{\kappa}$ on \mathbb{R} , with $\kappa \leq \tilde{\kappa}$, and the couples of determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ respectively $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ corresponding to the Brownian motion on \mathbb{R} with killing measure κ respectively $\tilde{\kappa}$. We will show that one can couple $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ on the same probability space such that $\mathcal{Z}_\infty \subseteq \tilde{\mathcal{Z}}_\infty$ and $\tilde{\mathcal{Y}}_\infty \subseteq \mathcal{Y}_\infty \cup \text{Supp}(\tilde{\kappa} - \kappa)$. Moreover if κ and $\tilde{\kappa}$ are proportional we may also have $\mathcal{Y}_\infty \subseteq \tilde{\mathcal{Y}}_\infty$. We will provide an explicit construction for these couplings in the section 7.2.

In the section 7.1 we will prove conditioning results for $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$: what is obtained if \mathcal{Y}_∞ or \mathcal{Z}_∞ is conditioned by either containing a point at a given location or not containing any points in a given interval. These results will be used in the next section. The conditional law we will obtain are analogous to those of the Uniform Spanning Tree on a finite undirected connected graph: Let \mathbb{G} be such a graph, E the set of its edges, C a weight function on E and Υ the corresponding Uniform Spanning Tree on \mathbb{G} . Let E_1 and E_2 be two disjoint subsets of E such that E_1 contains no cycles and such that erasing the edges in E_2 does not disconnect \mathbb{G} . The law of Υ conditioned by $E_1 \subseteq \Upsilon$ and $\Upsilon \cap E_2 = \emptyset$ can be described as follows: Let \mathbb{G}' be the graph obtained from \mathbb{G} through erasing the edges in E_2 and contracting (i.e. identifying the two end vertices) the edges in E_1 . The edges of \mathbb{G}' are in one to one correspondence with $E \setminus E_2$. If we keep the same weight function C on these edges and take Υ' an Uniform Spanning Tree on \mathbb{G}' , then $\Upsilon' \cup E_1$ has the same law as Υ conditioned by $E_1 \subseteq \Upsilon$ and $\Upsilon \cap E_2 = \emptyset$ (see proposition 4.2 in [1]).

Let κ be a Radon measure on \mathbb{R} and $G(x, y) = u_\uparrow(x \wedge y)u_\downarrow(x \vee y)$ the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - \kappa$. First we will restrict the Brownian motion with killing measure κ to a half-line by adding either a killing or a reflecting boundary point and describe

what is obtained if we apply the Wilson's algorithm to it. This is related to some of the conditional laws we are interested in. Diffusions with reflection were not discussed so far.

For $x_0 < y$ let

$$u_{\uparrow}^{(x_0 \times)}(y) := u_{\uparrow}(y) - \frac{u_{\uparrow}(x_0)}{u_{\downarrow}(x_0)} u_{\uparrow}(y)$$

and for $x_0 < y, z$ let

$$\begin{aligned} G^{(x_0 \times)}(y, z) &:= u_{\uparrow}^{(x_0 \times)}(y \wedge z) u_{\downarrow}(y \vee z) \\ \mathcal{K}^{(x_0 \times)}(y, z) &:= -\frac{1}{2} \frac{du_{\uparrow}^{(x_0 \times)}}{dx}((y \wedge z)^+) \frac{du_{\downarrow}}{dx}((y \vee z)^-) \end{aligned}$$

$G^{(x_0 \times)}$ was already introduced in (6.3.32). For $y < x_0$ let

$$u_{\downarrow}^{(\times x_0)}(y) := u_{\downarrow}(y) - \frac{u_{\downarrow}(x_0)}{u_{\uparrow}(x_0)} u_{\uparrow}(y)$$

and for $y, z < x_0$ let

$$\begin{aligned} G^{(\times x_0)}(y, z) &:= u_{\uparrow}(y \wedge z) u_{\downarrow}^{(\times x_0)}(y \vee z) \\ \mathcal{K}^{(\times x_0)}(y, z) &:= -\frac{1}{2} \frac{du_{\uparrow}}{dx}((y \wedge z)^+) \frac{du_{\downarrow}^{(\times x_0)}}{dx}((y \vee z)^-) \end{aligned}$$

$G^{(x_0 \times)}$ respectively $G^{(\times x_0)}$ is the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - \kappa$ restricted to the interval $(x_0, +\infty)$ respectively $(-\infty, x_0)$ with zero Dirichlet boundary condition at x_0 .

Let $x_0 \in \mathbb{R}$ such that $\kappa(\{x_0\}) = 0$. For $x_0 < y$ let

$$u_{\uparrow}^{(x_0 \triangleright)}(y) := u_{\uparrow}(y) + \left(\frac{du_{\downarrow}}{dx}(x_0) \right)^{-1} \frac{du_{\uparrow}}{dx}(x_0) u_{\downarrow}(y)$$

and for $y, z < x_0$ let

$$\begin{aligned} G^{(x_0 \triangleright)}(y, z) &:= u_{\uparrow}^{(x_0 \triangleright)}(y \wedge z) u_{\downarrow}(y \vee z) \\ \mathcal{K}^{(x_0 \triangleright)}(y, z) &:= -\frac{1}{2} \frac{du_{\uparrow}^{(x_0 \triangleright)}}{dx}((y \wedge z)^+) \frac{du_{\downarrow}}{dx}((y \vee z)^-) \end{aligned}$$

$\mathcal{K}^{(x_0 \triangleright)}$ was already introduced in (6.3.32). For $y < x_0$ let

$$u_{\downarrow}^{(\triangleleft x_0)}(y) := u_{\downarrow}(y) + \left(\frac{du_{\uparrow}}{dx}(x_0) \right)^{-1} \frac{du_{\downarrow}}{dx}(x_0) u_{\uparrow}(y)$$

and for $y, z < x_0$ let

$$\begin{aligned} G^{(\triangleleft x_0)}(y, z) &:= u_{\uparrow}(y \wedge z) u_{\downarrow}^{(\triangleleft x_0)}(y \vee z) \\ \mathcal{K}^{(\triangleleft x_0)}(y, z) &:= -\frac{1}{2} \frac{du_{\uparrow}}{dx}((y \wedge z)^+) \frac{du_{\downarrow}^{(\triangleleft x_0)}}{dx}((y \vee z)^-) \end{aligned}$$

$G^{(x_0^\triangleright)}$ respectively $G^{(\triangleleft x_0)}$ is the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - \kappa$ restricted to the interval $[x_0, +\infty)$ respectively $(-\infty, x_0]$ with zero Neumann boundary condition at x_0 . Equivalently $G^{(x_0^\triangleright)}$ respectively $G^{(\triangleleft x_0)}$ is the restriction to $[x_0, +\infty)$ respectively $(-\infty, x_0]$ of the Green's function on \mathbb{R} of $\frac{1}{2} \frac{d^2}{dx^2} - 1_{[x_0, +\infty)} \kappa$ respectively $\frac{1}{2} \frac{d^2}{dx^2} - 1_{(-\infty, x_0]} \kappa$.

Consider now $x_0 \in \mathbb{R}$ and $(x_n)_{n \geq 1}$ a dense sequence of pairwise disjoint points in $(x_0, +\infty)$. We consider the Wilson's algorithm applied to the Brownian motion on $(x_0, +\infty)$ with killing measure κ and killing boundary x_0 , where $(x_n)_{n \geq 0}$ is the sequence of starting points. Let $\mathcal{Y}_\infty^{(x_0^\times)}$ and $\mathcal{Z}_\infty^{(x_0^\times)}$ be the interwoven point processes in $[x_0, +\infty)$ obtained as result. See figure 3.a for an illustration of the first four steps of Wilson's algorithm and of $(\mathcal{Y}_\infty^{(x_0^\times)}, \mathcal{Z}_\infty^{(x_0^\times)})$. According to proposition 6.13, $x_0 \in \mathcal{Y}_\infty^{(x_0^\times)}$ a.s., $\mathcal{Y}_\infty^{(x_0^\times)} \cap (x_0, +\infty)$ is a determinantal point process with determinantal kernel $G^{(x_0^\times)}$ relatively to the measure $1_{(x_0, +\infty)} \kappa$ and $\mathcal{Z}_\infty^{(x_0^\times)}$ is a determinantal point process with kernel $\mathcal{K}^{(x_0^\times)}$ relatively to the measure $1_{z > x_0} dz$. The distribution of the $2n$ closest to x_0 points in $(\mathcal{Y}_\infty^{(x_0^\times)} \cap (x_0, +\infty)) \cup \mathcal{Z}_\infty^{(x_0^\times)}$, the odd-numbered belonging to $\mathcal{Y}_\infty^{(x_0^\times)} \cap (x_0, +\infty)$ and the even-numbered to $\mathcal{Z}_\infty^{(x_0^\times)}$, is given by the measure

$$M_n^{(x_0^\times)}(\mathcal{Y}_\infty^{(x_0^\times)}, \mathcal{Z}_\infty^{(x_0^\times)})(dz_1, dy_1, \dots, dz_n, dy_n) := 2^n \frac{u_\downarrow(y_n)}{u_\downarrow(x_0)} dz_1 \kappa(dy_1) \dots dz_n \kappa(dy_n)$$

Its total mass equals $\mathbb{P}(\#\mathcal{Y}_\infty^{(x_0^\times)} \geq n + 1)$. If the Wilson's algorithm is applied to the Brownian motion on $(-\infty, x_0)$, killed at x_0 and with killing measure κ , and $(\mathcal{Y}_\infty^{(\times x_0)}, \mathcal{Z}_\infty^{(\times x_0)})$ are the point processes returned by the algorithm, then the distribution of the $2n$ closest to x_0 points in $(\mathcal{Y}_\infty^{(\times x_0)} \cap (-\infty, x_0)) \cup \mathcal{Z}_\infty^{(\times x_0)}$ is given by the measure

$$M_n^{(\times x_0)}(\mathcal{Y}_\infty^{(\times x_0)}, \mathcal{Z}_\infty^{(\times x_0)})(dz_{-1}, dy_{-1}, \dots, dz_{-n}, dy_{-n}) := 2^n \frac{u_\uparrow(y_{-n})}{u_\uparrow(x_0)} dz_{-1} \kappa(dy_{-1}) \dots dz_{-n} \kappa(dy_{-n})$$

Let now $x_0 \in \mathbb{R}$ such that $\kappa(\{x_0\}) = 0$. If we replace the Brownian motion on $(x_0, +\infty)$ killed in x_0 by a Brownian motion on $[x_0, +\infty)$ reflected in x_0 , and keep the killing measure κ , we get another pair $(\mathcal{Y}_\infty^{(x_0^\blacktriangleright)}, \mathcal{Z}_\infty^{(x_0^\blacktriangleright)})$ of interwoven point processes on $[x_0, +\infty)$. The pair $(\mathcal{Y}_\infty^{(x_0^\blacktriangleright)}, \mathcal{Z}_\infty^{(x_0^\blacktriangleright)})$ can be also obtain through applying Wilson's algorithm to a Brownian motion on \mathbb{R} with the killing measure $1_{(x_0, +\infty)} \kappa$. See figure 3.b for an illustration of $(\mathcal{Y}_\infty^{(x_0^\blacktriangleright)}, \mathcal{Z}_\infty^{(x_0^\blacktriangleright)})$. Observe the difference with figure 3.a at the third step of Wilson's algorithm. $\mathcal{Y}_\infty^{(x_0^\blacktriangleright)}$ is a determinantal point process with determinantal kernel $G^{(x_0^\blacktriangleright)}$ relatively to the measure $1_{(x_0, +\infty)} \kappa$. $\mathcal{Z}_\infty^{(x_0^\blacktriangleright)}$ is a determinantal point process with kernel $\mathcal{K}^{(x_0^\blacktriangleright)}$ relatively to the measure $1_{z > x_0} dz$. The distribution of the $2n - 1$ closest to x_0 points in $\mathcal{Y}_\infty^{(x_0^\blacktriangleright)} \cup \mathcal{Z}_\infty^{(x_0^\blacktriangleright)}$, the odd-numbered

belonging to $\mathcal{Z}_\infty^{(x_0 \triangleright)}$ and the even-numbered to $\mathcal{Y}_\infty^{(x_0 \triangleright)}$, is given by the measure

$$M_n^{(x_0 \triangleright)}(\mathcal{Y}_\infty^{(x_0 \triangleright)}, \mathcal{Z}_\infty^{(x_0 \triangleright)})(dy_1, dz_1, \dots, dz_{n-1}, dy_n) := -2^n \left(\frac{du_\downarrow}{dx}(x_0) \right)^{-1} u_\downarrow(y_n) \kappa(dy_1) dz_1 \dots dz_{n-1} \kappa(dy_n)$$

If the Wilson's algorithm is applied to the Brownian motion on $(-\infty, x_0]$, reflected at x_0 and with killing measure κ , and $(\mathcal{Y}_\infty^{(\triangleleft x_0)}, \mathcal{Z}_\infty^{(\triangleleft x_0)})$ are the point processes returned by the algorithm, then the distribution of the $2n - 1$ closest to x_0 points in $\mathcal{Y}_\infty^{(\triangleleft x_0)} \cup \mathcal{Z}_\infty^{(\triangleleft x_0)}$ is given by the measure

$$M_n^{(\triangleleft x_0)}(\mathcal{Y}_\infty^{(\triangleleft x_0)}, \mathcal{Z}_\infty^{(\triangleleft x_0)})(dy_{-1}, dz_{-1}, \dots, dz_{-n+1}, dy_{-n}) := 2^n \left(\frac{du_\uparrow}{dx}(x_0) \right)^{-1} u_\uparrow(y_{-n}) \kappa(dy_{-1}) dz_{-1} \dots dz_{-n+1} \kappa(dy_{-n})$$




Fig. 3.a - Illustration of the first four steps of Wilson's algorithm in case of killing at x_0 and of $(\mathcal{Y}_\infty^{(x_0 \times)}, \mathcal{Z}_\infty^{(x_0 \times)})$: x-dots represent the points of $\mathcal{Y}_n^{(x_0 \times)}$, diamonds the points of $\mathcal{Z}_n^{(x_0 \times)}$ and thick lines the intervals in $\mathcal{J}_n^{(x_0 \times)}$.




Fig. 3.b - Illustration of the first four steps of Wilson's algorithm in case of reflection at x_0 and of $(\mathcal{Y}_\infty^{(x_0 \triangleright)}, \mathcal{Z}_\infty^{(x_0 \triangleright)})$: x-dots represent the points of $\mathcal{Y}_n^{(x_0 \triangleright)}$, diamonds the points of $\mathcal{Z}_n^{(x_0 \triangleright)}$ and thick lines the intervals in $\mathcal{J}_n^{(x_0 \triangleright)}$.

Let \mathcal{Y}_∞ and \mathcal{Z}_∞ be the determinantal point processes associated to the Brownian motion on \mathbb{R} with killing measure κ . Let $n, n' \in \mathbb{N}^*$. The following two factorizations hold:

$$\begin{aligned} M_{n+n'}(\mathcal{Y}_\infty, \mathcal{Z}_\infty)(dy_{-n'}, dz_{-n'}, \dots, dy_{-1}, dz_{-1}, dy_0, dz_1, dy_1, \dots, dz_n, dy_n) = \\ M_{n'}^{(\times y_0)}(\mathcal{Y}_\infty^{(\times y_0)}, \mathcal{Z}_\infty^{(\times y_0)})(dz_{-1}, dy_{-1}, \dots, dz_{-n'}, dy_{-n'}) \times G(y_0, y_0)\kappa(dy_0) \\ \times M_n^{(y_0 \times)}(\mathcal{Y}_\infty^{(y_0 \times)}, \mathcal{Z}_\infty^{(y_0 \times)})(dz_1, dy_1, \dots, dz_n, dy_n) \\ M_{n+n'-1}(\mathcal{Y}_\infty, \mathcal{Z}_\infty)(dy_{-n'}, dz_{-n'+1}, \dots, dz_{-1}, dy_{-1}, dz_0, dy_1, dz_1, \dots, dz_{n-1}, dy_n) = \\ M_{n'}^{(\triangleleft z_0)}(\mathcal{Y}_\infty^{(\triangleleft z_0)}, \mathcal{Z}_\infty^{(\triangleleft z_0)})(dy_{-1}, dz_{-1}, \dots, dz_{-n'+1}, dy_{-n'}) \times \mathcal{K}(z_0, z_0)dz_0 \\ \times M_n^{(z_0 \triangleright)}(\mathcal{Y}_\infty^{(z_0 \triangleright)}, \mathcal{Z}_\infty^{(z_0 \triangleright)})(dy_1, dz_1, \dots, dz_{n-1}, dy_n) \end{aligned}$$

The above factorisations imply the following:

Property 7.1. — *Let $\varepsilon > 0$ and let F_1 and F_2 be two measurable non-negative functionals on couples of point processes on \mathbb{R} and f a measurable non-negative function on \mathbb{R} . Then*

$$\begin{aligned} \mathbb{E} \left[\sum_{y_0 \in \mathcal{Y}_\infty} f(y_0) F_1(\mathcal{Y}_\infty \cap (-\infty, y_0], \mathcal{Z}_\infty \cap (-\infty, y_0]) F_2(\mathcal{Y}_\infty \cap [y_0, +\infty), \mathcal{Z}_\infty \cap [y_0, +\infty)) \right] \\ = \int_{\mathbb{R}} f(y_0) G(y_0, y_0) \mathbb{E}[F_1(\mathcal{Y}_\infty^{(\times y_0)}, \mathcal{Z}_\infty^{(\times y_0)})] \mathbb{E}[F_2(\mathcal{Y}_\infty^{(y_0 \times)}, \mathcal{Z}_\infty^{(y_0 \times)})] \kappa(dy_0) \end{aligned}$$

and

$$\begin{aligned} \mathbb{E} \left[\sum_{z_0 \in \mathcal{Z}_\infty} f(z_0) F_1(\mathcal{Y}_\infty \cap (-\infty, z_0], \mathcal{Z}_\infty \cap (-\infty, z_0]) F_2(\mathcal{Y}_\infty \cap [z_0, +\infty), \mathcal{Z}_\infty \cap [z_0, +\infty)) \right] \\ = \int_{\mathbb{R}} f(z_0) \mathcal{K}(z_0, z_0) \mathbb{E}[F_1(\mathcal{Y}_\infty^{(\triangleleft z_0)}, \mathcal{Z}_\infty^{(\triangleleft z_0)})] \mathbb{E}[F_2(\mathcal{Y}_\infty^{(z_0 \triangleright)}, \mathcal{Z}_\infty^{(z_0 \triangleright)})] dz_0 \end{aligned}$$

If $y_0 \in \text{Supp}(\kappa)$, then conditionally on $y_0 \in \mathcal{Y}_\infty$, $(\mathcal{Y}_\infty \cap (-\infty, y_0], \mathcal{Z}_\infty \cap (-\infty, y_0])$ and $(\mathcal{Y}_\infty \cap [y_0, +\infty), \mathcal{Z}_\infty \cap [y_0, +\infty))$ are independent, $(\mathcal{Y}_\infty \cap (-\infty, y_0], \mathcal{Z}_\infty \cap (-\infty, y_0])$ has the same law as $(\mathcal{Y}_\infty^{(\times y_0)}, \mathcal{Z}_\infty^{(\times y_0)})$ and $(\mathcal{Y}_\infty \cap [y_0, +\infty), \mathcal{Z}_\infty \cap [y_0, +\infty))$ has the same law as $(\mathcal{Y}_\infty^{(y_0 \times)}, \mathcal{Z}_\infty^{(y_0 \times)})$.

If $\kappa((-\infty, z_0)) > 0$, $\kappa((z_0, +\infty)) > 0$ and $\kappa(\{z_0\}) = 0$, then conditionally on $z_0 \in \mathcal{Z}_\infty$, $(\mathcal{Y}_\infty \cap (-\infty, z_0], \mathcal{Z}_\infty \cap (-\infty, z_0])$ and $(\mathcal{Y}_\infty \cap [z_0, +\infty), \mathcal{Z}_\infty \cap [z_0, +\infty))$ are independent, $(\mathcal{Y}_\infty \cap (-\infty, z_0], \mathcal{Z}_\infty \cap (-\infty, z_0])$ has the same law as $(\mathcal{Y}_\infty^{(\triangleleft z_0)}, \mathcal{Z}_\infty^{(\triangleleft z_0)})$ and $(\mathcal{Y}_\infty \cap [z_0, +\infty), \mathcal{Z}_\infty \cap [z_0, +\infty))$ has the same law as $(\mathcal{Y}_\infty^{(z_0 \triangleright)}, \mathcal{Z}_\infty^{(z_0 \triangleright)})$.

Let $y_0 \in \mathbb{R}$ and $c > 0$. We will denote by $(\mathcal{Y}_\infty^{(y_0)}, \mathcal{Z}_\infty^{(y_0)})$ the pair of interwoven determinantal point processes corresponding to the killing measure $\kappa + c\delta_{y_0}$, conditioned on $\mathcal{Y}_\infty^{(y_0)}$ containing y_0 . The law of $(\mathcal{Y}_\infty^{(y_0)}, \mathcal{Z}_\infty^{(y_0)})$ does not depend on the value of c according to the property 7.1. $(\mathcal{Y}_\infty^{(y_0)} \cap (y_0, +\infty), \mathcal{Z}_\infty^{(y_0)} \cap (y_0, +\infty))$ and

$(\mathcal{Y}_\infty^{(y_0)} \cap (-\infty, y_0), \mathcal{Z}_\infty^{(y_0)} \cap (-\infty, y_0))$ are independent. The distribution of the $2n$ closest to y_0 points in $(\mathcal{Y}_\infty^{(y_0)} \cup \mathcal{Z}_\infty^{(y_0)}) \cap (y_0, +\infty)$, on the event $\sharp(\mathcal{Y}_\infty^{(y_0)} \cap (y_0, +\infty)) \geq n$, is

$$(7.1.1) \quad \mathbb{1}_{y_0 < z_1 < y_1 < \dots < z_n < y_n} 2^n \frac{u_\downarrow(y_n)}{u_\downarrow(y_0)} dz_1 \kappa(dy_1) \dots dz_n \kappa(dy_n)$$

The distribution of the $2n$ closest to y_0 points in $(\mathcal{Y}_\infty^{(y_0)} \cup \mathcal{Z}_\infty^{(y_0)}) \cap (-\infty, y_0)$ is

$$(7.1.2) \quad \mathbb{1}_{y_0 > z_{-1} > y_{-1} > \dots > z_{-n} > y_{-n}} 2^n \frac{u_\uparrow(y_{-n})}{u_\uparrow(y_0)} dz_{-1} \kappa(dy_{-1}) \dots dz_{-n} \kappa(dy_{-n})$$

Let $a < b \in \mathbb{R}$. Next we will describe what happens if we condition by $\mathcal{Z}_\infty \cap [a, b] = \emptyset$. This condition implies in particular that $\sharp(\mathcal{Y}_\infty \cap [a, b]) \leq 1$. Let $\widehat{\mathbb{R}}$ be the quotient space where in \mathbb{R} we identify to one point all the points lying in $[a, b]$. $\widehat{\mathbb{R}}$ is homeomorphic to \mathbb{R} . Let $\hat{\pi}$ be the projection from \mathbb{R} to $\widehat{\mathbb{R}}$. Let θ be the class of $[a, b]$ in $\widehat{\mathbb{R}}$. We define on $\widehat{\mathbb{R}}$ the metric $d_{\widehat{\mathbb{R}}}$:

- If $x < y < a$ or $b < x < y$ then $d_{\widehat{\mathbb{R}}}(\hat{\pi}(x), \hat{\pi}(y)) = y - x$.
- If $x < a$ and $y > b$ then $d_{\widehat{\mathbb{R}}}(\hat{\pi}(x), \hat{\pi}(y)) = (y - x) - (b - a)$.
- If $x < a$ then $d_{\widehat{\mathbb{R}}}(\hat{\pi}(x), \theta) = a - x$.
- If $x > b$ then $d_{\widehat{\mathbb{R}}}(\hat{\pi}(x), \theta) = x - b$.

$\widehat{\mathbb{R}}$ endowed with $d_{\widehat{\mathbb{R}}}$ is isometric to \mathbb{R} . So we can define a standard Brownian motion on $\widehat{\mathbb{R}}$. Let $\hat{\kappa}$ be the measure κ pushed forward by $\hat{\pi}$ on $\widehat{\mathbb{R}}$. In particular $\hat{\kappa}(\{\theta\}) = \kappa([a, b])$. Let $(\widehat{\mathcal{Y}}_\infty, \widehat{\mathcal{Z}}_\infty)$ be the pair of interwoven determinantal point processes on $\widehat{\mathbb{R}}$ obtained by applying the Wilson's algorithm to the Brownian motion on $\widehat{\mathbb{R}}$ with killing measure $\hat{\kappa}$.

Proposition 7.2. — *Conditionally on $\mathcal{Z}_\infty \cap [a, b] = \emptyset$, $(\hat{\pi}(\mathcal{Y}_\infty), \hat{\pi}(\mathcal{Z}_\infty))$ has the same distribution as $(\widehat{\mathcal{Y}}_\infty, \widehat{\mathcal{Z}}_\infty)$. Moreover on the event $\mathcal{Y}_\infty \cap [a, b] \neq \emptyset$, the unique point in $\mathcal{Y}_\infty \cap [a, b]$ is distributed according the probability measure $\frac{\mathbb{1}_{a < y < b} \kappa(dy)}{\kappa([a, b])}$.*

Proof. — First we compute $\mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset)$. We consider that a and b are the first two starting points in the Wilson's algorithm. Then

$$\begin{aligned} \mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset) &= \mathbb{P}(B_{T_1^-}^{(a)} > b) + \mathbb{P}(B_{T_1^-}^{(a)} < a, B_{T_2^-}^{(b)} = a) + \mathbb{P}(B_{T_1^-}^{(a)} = B_{T_2^-}^{(b)} \in [a, b]) \\ &= \frac{1}{2} \frac{du_\uparrow}{dx}(a^-) u_\downarrow(b) - \frac{1}{2} u_\uparrow(a) \frac{du_\downarrow}{dx}(b^+) + u_\uparrow(a) u_\downarrow(b) \kappa([a, b]) \end{aligned}$$

Next we determine the Green's function \widehat{G} of $\frac{1}{2} \frac{d^2}{dx^2} - \hat{\kappa}$ on $\widehat{\mathbb{R}}$. Let \hat{u}_\uparrow and \hat{u}_\downarrow be two solutions on $\widehat{\mathbb{R}}$ to

$$\frac{1}{2} \frac{d\hat{u}}{dx} - \hat{u} \hat{\kappa} = 0$$

with the initial conditions $\hat{u}_\uparrow(\theta) = u_\uparrow(a)$, $\frac{d\hat{u}_\uparrow}{dx}(\theta^-) = \frac{du_\uparrow}{dx}(a^-)$, $\hat{u}_\downarrow(\theta) = u_\downarrow(b)$ and $\frac{d\hat{u}_\downarrow}{dx}(\theta^+) = \frac{du_\downarrow}{dx}(b^+)$. Then for $x \leq a$, $\hat{u}_\uparrow(\hat{\pi}(x)) = u_\uparrow(x)$ and for $x \geq b$, $\hat{u}_\downarrow(\hat{\pi}(x)) =$

$u_\downarrow(x)$. \hat{u}_\uparrow and \hat{u}_\downarrow are positive, \hat{u}_\uparrow is non-decreasing and \hat{u}_\downarrow non-increasing. Moreover:

$$\frac{d\hat{u}_\uparrow}{dx}(\theta^+) = \frac{d\hat{u}_\uparrow}{dx}(\theta^-) + 2\hat{u}_\uparrow(\theta)\hat{\kappa}(\{\theta\}) = \frac{du_\uparrow}{dx}(a^-) + 2u_\uparrow(a)\kappa([a, b])$$

The Wronskian of \hat{u}_\downarrow and \hat{u}_\uparrow equals

$$\begin{aligned} W(\hat{u}_\downarrow, \hat{u}_\uparrow) &= \hat{u}_\downarrow(\theta) \frac{d\hat{u}_\uparrow}{dx}(\theta^+) - \hat{u}_\uparrow(\theta) \frac{d\hat{u}_\downarrow}{dx}(\theta^+) \\ &= \frac{du_\uparrow}{dx}(a^-)u_\downarrow(b) - u_\uparrow(a) \frac{du_\downarrow}{dx}(b^+) + 2u_\uparrow(a)u_\downarrow(b)\kappa([a, b]) \\ &= 2\mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset) \end{aligned}$$

Thus \widehat{G} equals

$$\widehat{G}(\tilde{x}, \tilde{y}) = \frac{\hat{u}_\uparrow(\tilde{x} \wedge \tilde{y})\hat{u}_\downarrow(\tilde{x} \vee \tilde{y})}{\mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset)}$$

In particular if $x \leq a$ and $y \geq b$ then

$$(7.1.3) \quad \widehat{G}(\hat{\pi}(x), \hat{\pi}(y)) = \frac{u_\uparrow(x)u_\downarrow(y)}{\mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset)} = \frac{G(x, y)}{\mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset)}$$

To prove the equality in law, we need to consider the probabilities of the events $\mathcal{C}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, a_1, b_1, \dots, \tilde{a}_n, \tilde{b}_n, a_n, b_n)$ where $n \geq 1$ and $a_0 < b_0 < \tilde{a}_1 < \tilde{b}_1 < a_1 < b_1 < \dots < \tilde{a}_n < \tilde{b}_n < a_n < b_n \in \mathbb{R}$, corresponding to following conditions:

- $\mathcal{Y}_\infty \cap [a_0, b_0] \neq \emptyset$, $\mathcal{Y}_\infty \cap [a_n, b_n] \neq \emptyset$
- $\forall r \in \{1, \dots, n\}$, $\#\mathcal{Y}_\infty \cap [a_r, b_r] = 1$
- $\forall r \in \{1, \dots, n\}$, $\#\mathcal{Z}_\infty \cap (\tilde{a}_r, \tilde{b}_r) = 1$
- $\forall r \in \{0, \dots, n-1\}$, $(\mathcal{Y}_\infty \cup \mathcal{Z}_\infty) \cap (b_r, \tilde{a}_r) = \emptyset$, $(\mathcal{Y}_\infty \cup \mathcal{Z}_\infty) \cap [\tilde{b}_r, a_{r+1}] = \emptyset$

We will also assume that either all of the $[a_r, b_r]$ do not intersect $[a, b]$ or one of the $[a_r, b_r]$ is contained in $[a, b]$ and the other do not intersect $[a, b]$. The probabilities of such events determine the joint law of $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ on the event $\#\mathcal{Y}_\infty \geq 2$, $\mathcal{Z}_\infty \cap [a, b] = \emptyset$. We will denote $\widehat{\mathcal{C}}_n(\cdot)$ the analogously defined events where we replace $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ by $(\widehat{\mathcal{Y}}_\infty, \widehat{\mathcal{Z}}_\infty)$. We do not need to deal with the event $\#\mathcal{Y}_\infty = 1$ because then $\mathcal{Z}_\infty = \emptyset$.

We first consider the case of $[a, b] \cap (\bigcup_{r=0}^n [a_r, b_r]) = \emptyset$. If there is $r_0 \in \{0, n-1\}$ such that $b_{r_0} < a$ and $b < a_{r_0+1}$ then

$$\begin{aligned} &\mathbb{P}\left(\mathcal{C}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, a_1, b_1, \dots, \tilde{a}_n, \tilde{b}_n, a_n, b_n), \mathcal{Z}_\infty \cap [a, b] = \emptyset\right) \\ &= \int_{[a_0, b_0]} u_\uparrow(y_0)\kappa(dy_0) \times \int_{[a_n, b_n]} u_\downarrow(y_n)\kappa(dy_n) \times \prod_{r=1}^{n-1} \kappa([a_r, b_r]) \\ &\quad \times \prod_{r \neq r_0} 2(\tilde{b}_r - \tilde{a}_r) \times 2Leb([\tilde{a}_{r_0}, \tilde{b}_{r_0}] \setminus [a, b]) \end{aligned}$$

Using (7.1.3) we get that the above equals

$$\mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset) \times \mathbb{P}\left(\widehat{\mathcal{C}}_n(\hat{\pi}(a_0), \hat{\pi}(b_0), \hat{\pi}(\tilde{a}_1), \hat{\pi}(\tilde{b}_1), \dots, \hat{\pi}(a_n), \hat{\pi}(b_n))\right)$$

If $b < a_0$, then we consider a Wilson's algorithm where the $2(n+1)$ first starting points are $\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_n, \tilde{b}_n, a, b$. The conditions $\mathcal{C}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, a_1, b_1, \dots, \tilde{a}_n, \tilde{b}_n, a_n, b_n)$ and $\mathcal{Z}_\infty \cap [a, b] = \emptyset$ are satisfied if and only if the following is true:

- $B_{T_1^-}^{(\tilde{a}_1)} \in [a_0, b_0]$, $B_{T_{2n}^-}^{(\tilde{b}_n)} \in [a_n, b_n]$, for all $r \in \{1, \dots, n-1\}$, $B_{T_{2r}^-}^{(\tilde{b}_r)} = B_{T_{2r+1}^-}^{(\tilde{a}_{r+1})} \in [a_r, b_r]$ and for all $r \in \{1, \dots, n\}$, $\mathcal{Y}_\infty \cap (\tilde{a}_r, \tilde{b}_r) = \emptyset$.
- Either $B_{T_{2n+1}^-}^{(a)} \in (b, B_{T_1^-}^{(\tilde{a}_1)})$ or $B_{T_{2n+2}^-}^{(b)} < a$ or $B_{T_{2n+1}^-}^{(a)} = B_{T_{2n+2}^-}^{(b)} \in [a, b]$.

Then

$$\begin{aligned} & \mathbb{P}\left(\mathcal{C}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, a_1, b_1, \dots, \tilde{a}_n, \tilde{b}_n, a_n, b_n), \mathcal{Z}_\infty \cap [a, b] = \emptyset\right) \\ &= \int_{[a_n, b_n]} u_\downarrow(y_n) \kappa(dy_n) \times \prod_{r=1}^{n-1} \kappa([a_r, b_r]) \times \prod_{r=1}^n 2(\tilde{b}_r - \tilde{a}_r) \\ & \quad \times \left(u_\uparrow(a) \int_{b < y < y_0, y_0 \in [a_0, b_0]} (u_\downarrow(y) u_\uparrow(y_0) - u_\uparrow(y) u_\downarrow(y_0)) \kappa(dy) \kappa(dy_0) + u_\uparrow(a) \kappa([a_0, b_0]) \right) \\ & \quad + \left(\int_{y_{-1} < a} u_\uparrow(y_{-1}) \kappa(dy_{-1}) + u_\uparrow(a) \kappa([a_0, b_0]) \right) \\ & \quad \times \int_{[a_0, b_0]} (u_\downarrow(b) u_\uparrow(y_0) - u_\uparrow(b) u_\downarrow(y_0)) \kappa(dy_0) \\ &= \int_{[a_n, b_n]} u_\downarrow(y_n) \kappa(dy_n) \times \prod_{r=1}^{n-1} \kappa([a_r, b_r]) \times \prod_{r=1}^n 2(\tilde{b}_r - \tilde{a}_r) \\ & \quad \times \left(\frac{1}{2} u_\uparrow(a) \int_{[a_0, b_0]} \left(\frac{du_\downarrow}{dx}(b^+) u_\uparrow(y_0) - \frac{du_\uparrow}{dx}(b^+) u_\downarrow(y_0) \right) \kappa(dy_0) \right. \\ & \quad \left. + \left(\frac{1}{2} \frac{du_\uparrow}{dx}(a^-) + u_\uparrow(a) \kappa([a_0, b_0]) \right) \int_{[a_0, b_0]} (u_\downarrow(b) u_\uparrow(y_0) - u_\uparrow(b) u_\downarrow(y_0)) \kappa(dy_0) \right) \end{aligned}$$

But for $y_0 \geq b$

$$\begin{aligned} \hat{u}_\uparrow(\hat{\pi}(y_0)) &= \frac{1}{2} u_\uparrow(a) \left(\frac{du_\downarrow}{dx}(b^+) u_\uparrow(y_0) - \frac{du_\uparrow}{dx}(b^+) u_\downarrow(y_0) \right) \\ & \quad + \left(\frac{1}{2} \frac{du_\uparrow}{dx}(a^-) + u_\uparrow(a) \kappa([a_0, b_0]) \right) (u_\downarrow(b) u_\uparrow(y_0) - u_\uparrow(b) u_\downarrow(y_0)) \end{aligned}$$

Indeed one can check the initial conditions $\hat{u}_\uparrow(\hat{\pi}(b)) = u_\uparrow(a)$ and $\frac{d\hat{u}_\uparrow}{dx}(\hat{\pi}(b)^+) = \frac{du_\uparrow}{dx}(a^-) + 2u_\uparrow(a) \kappa([a_0, b_0])$. It follows that

$$\begin{aligned} & \mathbb{P}\left(\mathcal{C}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, \dots, a_n, b_n), \mathcal{Z}_\infty \cap [a, b] = \emptyset\right) \\ &= \int_{[\hat{\pi}(a_0), \hat{\pi}(b_0)]} \hat{u}_\uparrow(\tilde{y}_0) \kappa(d\tilde{y}_0) \times \int_{[\hat{\pi}(a_n), \hat{\pi}(b_n)]} \hat{u}_\downarrow(\tilde{y}_n) \kappa(d\tilde{y}_n) \times \prod_{r=1}^{n-1} \kappa([a_r, b_r]) \times \prod_{r=1}^n 2(\tilde{b}_r - \tilde{a}_r) \\ &= \mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset) \times \mathbb{P}\left(\widehat{\mathcal{C}}_n(\hat{\pi}(a_0), \hat{\pi}(b_0), \hat{\pi}(\tilde{a}_1), \hat{\pi}(\tilde{b}_1), \dots, \hat{\pi}(a_n), \hat{\pi}(b_n))\right) \end{aligned}$$

Similar holds if $b_n < a$.

Now we consider the case when there is $r_0 \in \{0, \dots, n\}$ such that $[a_{r_0}, b_{r_0}] \subseteq [a, b]$ and $[a, b] \cap (\bigcup_{r \neq r_0} [a_r, b_r]) = \emptyset$. If $1 \leq r_0 \leq n-1$ then

$$\begin{aligned} & \mathbb{P} \left(\mathcal{E}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, \dots, a_n, b_n), \mathcal{Z}_\infty \cap [a, b] = \emptyset \right) \\ &= \int_{[a_0, b_0]} u_\uparrow(y_0) \kappa(dy_0) \times \int_{[a_n, b_n]} u_\downarrow(y_n) \kappa(dy_n) \times \prod_{r=1}^{n-1} \kappa([a_r, b_r]) \\ & \quad \times \prod_{r \neq r_0, r_0+1} 2(\tilde{b}_r - \tilde{a}_r) \times 2Leb([\tilde{a}_{r_0}, \tilde{b}_{r_0}] \setminus [a, b]) \times 2Leb([\tilde{a}_{r_0+1}, \tilde{b}_{r_0+1}] \setminus [a, b]) \\ &= \frac{\kappa([a_{r_0}, b_{r_0}])}{\kappa([a, b])} \times \mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset) \times \mathbb{P} \left(\widehat{\mathcal{E}}_n(\hat{\pi}(a_0), \hat{\pi}(b_0), \hat{\pi}(\tilde{a}_1), \hat{\pi}(\tilde{b}_1), \dots, \hat{\pi}(a_n), \hat{\pi}(b_n)) \right) \end{aligned}$$

Moreover $\hat{\pi}(a_{r_0}) = \hat{\pi}(b_{r_0}) = \theta$. If $r_0 = 0$ then

$$\begin{aligned} & \mathbb{P} \left(\mathcal{E}_n(a_0, b_0, \tilde{a}_1, \tilde{b}_1, \dots, a_n, b_n), \mathcal{Z}_\infty \cap [a, b] = \emptyset \right) \\ &= u_\uparrow(a) \kappa([a_0, b_0]) \times \int_{[a_n, b_n]} u_\downarrow(y_n) \kappa(dy_n) \\ & \quad \times \prod_{r=1}^{n-1} \kappa([a_r, b_r]) \times \prod_{r=2}^n 2(\tilde{b}_r - \tilde{a}_r) \times 2Leb([\tilde{a}_1, \tilde{b}_1] \setminus [a, b]) \\ &= \frac{\kappa([a_0, b_0])}{\kappa([a, b])} \times \mathbb{P}(\mathcal{Z}_\infty \cap [a, b] = \emptyset) \times \mathbb{P} \left(\widehat{\mathcal{E}}_n(\hat{\pi}(a_0), \hat{\pi}(b_0), \hat{\pi}(\tilde{a}_1), \hat{\pi}(\tilde{b}_1), \dots, \hat{\pi}(a_n), \hat{\pi}(b_n)) \right) \end{aligned}$$

and $\hat{\pi}(a_0) = \hat{\pi}(b_0) = \theta$. We have a similar expression if $r_0 = n$. \square

Next we deal with the condition of the determinantal point process \mathcal{Y}_∞ not charging a given subinterval of \mathbb{R} . We will consider the following more general situation: Let κ and $\tilde{\kappa}$ be two different killing measures on \mathbb{R} , with $\kappa \leq \tilde{\kappa}$, and the couples of determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ respectively $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ corresponding to the Brownian motion on \mathbb{R} with killing measure κ respectively $\tilde{\kappa}$. Let \tilde{G} be the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - \tilde{\kappa}$, factorized as

$$\tilde{G}(x, y) = \tilde{u}_\uparrow(x \wedge y) \tilde{u}_\downarrow(x \vee y)$$

Let

$$\tilde{\mathcal{K}}(y, z) := -\frac{1}{2} \frac{d\tilde{u}_\uparrow}{dx}((y \wedge z)^+) \frac{d\tilde{u}_\downarrow}{dx}((y \vee z)^-)$$

We will assume that $\tilde{\kappa} - \kappa$ has a first moment, that is to say

$$\int_{\mathbb{R}} |x|(\tilde{\kappa}(dx) - \kappa(dx)) < +\infty$$

Let χ be the Radon-Nikodym derivative

$$\chi := \frac{d\kappa}{d\tilde{\kappa}}$$

By definition $0 \leq \chi \leq 1$. Let $\Delta\tilde{\mathcal{Y}}$ be the point process obtained from $\tilde{\mathcal{Y}}_\infty$ as follows: Given a point y in $\tilde{\mathcal{Y}}_\infty$ we chose to erase it with probability $\chi(y)$ and keep it with

probability $1 - \chi(y)$, each choice being independent from the other choices and the position of other points. It is immediate to check that $\Delta\tilde{\mathcal{Y}}$ is a determinantal point process with determinantal kernel $(\tilde{G}(x, y))_{x, y \in \mathbb{R}}$ relatively to the measure $(1 - \chi)\tilde{\kappa}$, that is to say the measure $\tilde{\kappa} - \kappa$. We will show that conditionally on $\Delta\tilde{\mathcal{Y}} = \emptyset$, $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ has the same law as $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$. In case $1 - \chi$ being the indicator function of a bounded subinterval of \mathbb{R} , this gives the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ conditioned on $\tilde{\mathcal{Y}}_\infty$ not charging this subinterval.

Lemma 7.3. — $\Delta\tilde{\mathcal{Y}}$ is a.s. finite. Let

$$v_{\kappa, \tilde{\kappa}}(y) := \left(\tilde{u}_\uparrow(y) - \int_{y_{-1} < y} \tilde{u}_\uparrow(y_{-1})(u_\downarrow(y_{-1})u_\uparrow(y) - u_\uparrow(y_{-1})u_\downarrow(y))(\tilde{\kappa} - \kappa)(dy_{-1}) \right) \\ \times \left(\tilde{u}_\downarrow(y) - \int_{y_1 > y} \tilde{u}_\downarrow(y_1)(u_\uparrow(y_1)u_\downarrow(y) - u_\downarrow(y_1)u_\uparrow(y))(\tilde{\kappa} - \kappa)(dy_1) \right)$$

Then

$$\mathbb{P}(\#\Delta\tilde{\mathcal{Y}} = 1) = \int_{\mathbb{R}} v_{\kappa, \tilde{\kappa}}(y)(\tilde{\kappa} - \kappa)(dy)$$

The distribution of the unique point in $\Delta\tilde{\mathcal{Y}}$ conditionally on $\#\Delta\tilde{\mathcal{Y}} = 1$ is

$$\frac{v_{\kappa, \tilde{\kappa}}(y)(\tilde{\kappa} - \kappa)(dy)}{\mathbb{P}(\#\Delta\tilde{\mathcal{Y}} = 1)}$$

Furthermore

$$\mathbb{P}(\#\Delta\tilde{\mathcal{Y}} \geq 2) \leq \frac{1}{2} \left(\int_{\mathbb{R}} \tilde{G}(y, y)(\tilde{\kappa}(dy) - \kappa(dy)) \right)^2$$

and $\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset) > 0$.

Proof. — First let us check that $\int_{\mathbb{R}} \tilde{G}(y, y)(\tilde{\kappa}(dy) - \kappa(dy)) < +\infty$. Since $\tilde{\kappa} - \kappa$ has a first moment, we need only to show that $\tilde{G}(y, y)$ grows sub-linearly in the neighbourhood of $-\infty$ and $+\infty$. Let $a < b \in \mathbb{R}$ such that $\tilde{\kappa}((a, b)) > 0$. Let $\tilde{G}_{a, b}$ be the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - 1_{(a, b)}\tilde{\kappa}$. Then $\tilde{G}_{a, b}(y, y)$ is affine on $(-\infty, a)$ and on $(b, +\infty)$. Moreover $\tilde{G}(y, y) \leq \tilde{G}_{a, b}(y, y)$. Thus we get

$$\mathbb{E}[\#\Delta\tilde{\mathcal{Y}}] = \int_{\mathbb{R}} \tilde{G}(y, y)(\tilde{\kappa}(dy) - \kappa(dy)) < +\infty$$

In particular $\Delta\tilde{\mathcal{Y}}$ is a.s. finite.

To bound $\mathbb{P}(\#\Delta\tilde{\mathcal{Y}} \geq 2)$ we use the following:

$$\mathbb{P}(\#\Delta\tilde{\mathcal{Y}} \geq 2) \leq \frac{1}{2} \mathbb{E}[\#\Delta\tilde{\mathcal{Y}}(\#\Delta\tilde{\mathcal{Y}} - 1)] \\ = \frac{1}{2} \int_{\mathbb{R}^2} (\tilde{G}(x, x)\tilde{G}(y, y) - \tilde{G}(x, y)^2)(\tilde{\kappa}(dx) - \kappa(dx))(\tilde{\kappa}(dy) - \kappa(dy)) \\ \leq \frac{1}{2} \left(\int_{\mathbb{R}} \tilde{G}(y, y)(\tilde{\kappa}(dy) - \kappa(dy)) \right)^2$$

The expression of $\mathbb{E}[\#\Delta\tilde{\mathcal{Y}}(\#\Delta\tilde{\mathcal{Y}} - 1)]$ that we used is general for determinantal point processes.

Let's prove now that $\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset) > 0$. $\Delta\tilde{\mathcal{Y}}$ is determinantal point process associated to a trace-class self-adjoint positive semi-definite contraction operator on $\mathbb{L}^2(d\tilde{\kappa} - d\kappa)$. $\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset) > 0$ if and only if all the eigenvalues of the operator are strictly less than 1 (see theorem 4.5.3 in [11]). Let $f \in \mathbb{L}^2(\tilde{\kappa} - \kappa)$. Let

$$F(x) := \int_{\mathbb{R}} \tilde{G}(x, y) f(y) (\tilde{\kappa}(dy) - \kappa(dy))$$

F is continuous, dominated by

$$\tilde{G}(x, x)^{\frac{1}{2}} \left(\int_{\mathbb{R}} \tilde{G}(y, y) (\tilde{\kappa}(dy) - \kappa(dy)) \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}} f(y)^2 (\tilde{\kappa}(dy) - \kappa(dy)) \right)^{\frac{1}{2}}$$

and has left-side and right-side derivatives at every point. F satisfies the equation

$$-\frac{1}{2} \frac{d^2 F}{dx^2} + F\tilde{\kappa} = f(\tilde{\kappa} - \kappa)$$

Assume by absurd that $f = F(\tilde{\kappa} - \kappa)$ -almost everywhere. Then

$$\begin{aligned} \int_{\mathbb{R}} F(x)^2 (\tilde{\kappa}(dx) - \kappa(dx)) &= \int_{\mathbb{R}} f(x) F(x) (\tilde{\kappa}(dx) - \kappa(dx)) \\ &= \int_{\mathbb{R}} F(x)^2 \tilde{\kappa}(dx) + \frac{1}{2} \int_{\mathbb{R}} \frac{dF}{dx}(x)^2 dx \end{aligned}$$

Thus F is necessarily constant. But then this means that $(\tilde{\kappa} - \kappa)(\mathbb{R}) = \tilde{\kappa}(\mathbb{R})$, which is impossible because κ is non zero. Thus 1 is not an eigenvalue of the operator defining the determinantal process $\Delta\tilde{\mathcal{Y}}$ and thus $\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset) > 0$.

As for $\tilde{\mathcal{Y}}_{\infty}$, the spacing between consecutive points of $\Delta\tilde{\mathcal{Y}}$ are independent. By construction $\Delta\tilde{\mathcal{Y}} \subseteq \text{Supp}(\tilde{\kappa} - \kappa)$. Given $y_0 \in \text{Supp}(\tilde{\kappa} - \kappa)$, let

$$1_{y > y_0} f_{\Delta\tilde{\mathcal{Y}}}(y_0, y) (\tilde{\kappa}(dy) - \kappa(dy))$$

be the distribution of the lowest point in $\Delta\tilde{\mathcal{Y}} \cap (y_0, +\infty)$ conditionally on $y_0 \in \Delta\tilde{\mathcal{Y}}$. Since y_0 may be the maximum of $\Delta\tilde{\mathcal{Y}}$, $f_{\Delta\tilde{\mathcal{Y}}}(y_0, y) (\tilde{\kappa}(dy) - \kappa(dy)) < 1$. For y to be $\min \Delta\tilde{\mathcal{Y}} \cap (y_0, +\infty)$, y must belong to $\tilde{\mathcal{Y}}_{\infty}$, all points in $y' \in \tilde{\mathcal{Y}}_{\infty} \cap (y_0, y)$ must be erased (probability $\chi(y')$ for each), and y must be kept (probability $1 - \chi(y)$). For $y' > y_0$, let $f_{\tilde{G}}(y_0, y')$ be

$$f_{\tilde{G}}(y_0, y') = 2(y' - y_0) \frac{\tilde{u}_{\downarrow}(y')}{\tilde{u}_{\downarrow}(y_0)}$$

$1_{y' > y_0} f_{\tilde{G}}(y_0, y') \tilde{\kappa}(dy')$ is the distribution of $\min \tilde{\mathcal{Y}}_\infty \cap (y_0, +\infty)$ conditionally on $y_0 \in \tilde{\mathcal{Y}}_\infty$ (proposition 6.11). $f_{\Delta\tilde{\mathcal{Y}}}$ and $f_{\tilde{G}}$ are related as follows:

$$\begin{aligned} f_{\Delta\tilde{\mathcal{Y}}}(y_0, y) &= f_{\tilde{G}}(y_0, y) + \sum_{j \geq 2} \int_{y_0 < \dots < y_{j-1} < y} f_{\tilde{G}}(y_0, y_1) \dots f_{\tilde{G}}(y_{j-1}, y) \prod_{i=1}^{j-1} \chi(y_i) \kappa(dy_i) \\ &= \frac{\tilde{u}_\downarrow(y)}{\tilde{u}_\downarrow(y_0)} \left(2(y - y_0) + \sum_{j \geq 2} 2^j \int_{y_0 < \dots < y_{j-1} < y} (y_1 - y_0) \dots (y - y_{j-1}) \prod_{i=1}^{j-1} \kappa(dy_i) \right) \end{aligned}$$

But

$$\begin{aligned} &2(y - y_0) + \sum_{j \geq 2} 2^j \int_{y_0 < \dots < y_{j-1} < y} (y_1 - y_0) \dots (y - y_{j-1}) \prod_{i=1}^{j-1} \kappa(dy_i) \\ &= \frac{u_\downarrow(y_0)}{u_\downarrow(y)} \left(f_G(y_0, y) + \sum_{j \geq 2} \int_{y_0 < \dots < y_{j-1} < y} f_G(y_0, y_1) \dots f_G(y_{j-1}, y) \prod_{i=1}^{j-1} \kappa(dy_i) \right) \\ &= \frac{u_\downarrow(y_0)}{u_\downarrow(y)} \left(G(y, y) - \frac{G(y_0, y)^2}{G(y_0, y_0)} \right) = u_\downarrow(y_0) u_\uparrow(y) - u_\uparrow(y_0) u_\downarrow(y) \end{aligned}$$

(see section 6.3). It follows that

$$f_{\Delta\tilde{\mathcal{Y}}}(y_0, y) = \frac{\tilde{u}_\downarrow(y)}{\tilde{u}_\downarrow(y_0)} (u_\downarrow(y_0) u_\uparrow(y) - u_\uparrow(y_0) u_\downarrow(y))$$

In particular, if $y_0 < y_1 < \dots < y_n \in \mathbb{R}$, the infinitesimal probability that $\Delta\tilde{\mathcal{Y}}$ has a point at each of the locations y_i and no points in-between is

$$\begin{aligned} &\tilde{G}(y_0, y_0) f_{\Delta\tilde{\mathcal{Y}}}(y_0, y_1) \dots f_{\Delta\tilde{\mathcal{Y}}}(y_{n-1}, y_n) \prod_{i=0}^n (\tilde{\kappa}(dy_i) - \kappa(dy_i)) \\ &= \tilde{u}_\uparrow(y_0) \tilde{u}_\downarrow(y_n) \prod_{i=1}^n (u_\downarrow(y_i) u_\uparrow(y_{i-1}) - u_\uparrow(y_i) u_\downarrow(y_{i-1})) \prod_{i=0}^n (\tilde{k}(dy_i) - k(dy_i)) \end{aligned}$$

Thus the expression of $v_{\kappa, \tilde{\kappa}}(y)$ is a sieve identity obtained as follows: $v_{\kappa, \tilde{\kappa}}(y)(\tilde{\kappa} - \kappa)(dy)$ is the infinitesimal probability that $\Delta\tilde{\mathcal{Y}}$ contains a point at y , from which we subtract the infinitesimal probabilities to have a point at y at another below respectively above, and to which we add the infinitesimal probability to have a point at y and points both below and above y . \square

Next we deal with the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ conditionally on $\Delta\tilde{\mathcal{Y}} = 0$. Let $y_0 \in \text{Supp}(\tilde{\kappa} - \kappa)$. First we will compute the probability that $\Delta\tilde{\mathcal{Y}} \cap (y_0, +\infty) \neq \emptyset$ conditionally on $y_0 \in \tilde{\mathcal{Y}}_\infty$.

Lemma 7.4. — *There are positive constants c_1 and c_2 such that for all $x \in \mathbb{R}$*

$$(7.1.4) \quad \int_{y < x} (u_{\downarrow}(y)u_{\uparrow}(x) - u_{\uparrow}(y)u_{\downarrow}(x))\tilde{u}_{\uparrow}(y)(\tilde{\kappa}(dy) - \kappa(dy)) = \tilde{u}_{\uparrow}(x) - c_1 u_{\uparrow}(x)$$

$$(7.1.5) \quad \int_{y > x} (u_{\uparrow}(y)u_{\downarrow}(x) - u_{\downarrow}(y)u_{\uparrow}(x))\tilde{u}_{\downarrow}(y)(\tilde{\kappa}(dy) - \kappa(dy)) = \tilde{u}_{\downarrow}(x) - c_2 u_{\downarrow}(x)$$

In particular

$$v_{\kappa, \tilde{\kappa}}(y) = c_1 c_2 u_{\uparrow}(y)u_{\downarrow}(y)$$

Proof. — We will prove (7.1.5). The proof of (7.1.4) is similar. Let f be the function

$$f(x) := \tilde{u}_{\downarrow}(x) - \int_{y > x} (u_{\uparrow}(y)u_{\downarrow}(x) - u_{\downarrow}(y)u_{\uparrow}(x))\tilde{u}_{\downarrow}(y)(\tilde{\kappa}(dy) - \kappa(dy))$$

The derivative of f , defined everywhere except at most countably many points, is

$$\frac{df}{dx}(x) = \frac{d\tilde{u}_{\downarrow}}{dx}(x) - \int_{y > x} \left(u_{\uparrow}(y) \frac{du_{\downarrow}}{dx}(x) - u_{\downarrow}(y) \frac{du_{\uparrow}}{dx}(x) \right) \tilde{u}_{\downarrow}(y)(\tilde{\kappa}(dy) - \kappa(dy))$$

The weak second derivative of f is:

$$\begin{aligned} \frac{d^2 f}{dx^2}(x) &= \frac{d^2 \tilde{u}_{\downarrow}}{dx^2}(x) - \int_{y > x} \left(u_{\uparrow}(y) \frac{d^2 u_{\downarrow}}{dx^2}(x) - u_{\downarrow}(y) \frac{d^2 u_{\uparrow}}{dx^2}(x) \right) \tilde{u}_{\downarrow}(y)(\tilde{\kappa}(dy) - \kappa(dy)) \\ &\quad + \left(u_{\uparrow}(x) \frac{du_{\downarrow}}{dx}(x) - u_{\downarrow}(x) \frac{du_{\uparrow}}{dx}(x) \right) \tilde{u}_{\downarrow}(x)(\tilde{\kappa}(dx) - \kappa(dx)) \\ &= 2\tilde{u}_{\downarrow}(x)\tilde{\kappa}(dx) \\ &\quad - \int_{y > x} (u_{\uparrow}(y)u_{\downarrow}(x) - u_{\downarrow}(y)u_{\uparrow}(x))\tilde{u}_{\downarrow}(y)(\tilde{\kappa}(dy) - \kappa(dy)) \times \kappa(dx) \\ &\quad + 2\tilde{u}_{\downarrow}(x)(\tilde{\kappa}(dx) - \kappa(dx)) \\ &= 2\tilde{u}_{\downarrow}(x)\kappa(dx) \\ &\quad - \int_{y > x} (u_{\uparrow}(y)u_{\downarrow}(x) - u_{\downarrow}(y)u_{\uparrow}(x))\tilde{u}_{\downarrow}(y)(\tilde{\kappa}(dy) - \kappa(dy)) \times \kappa(dx) \\ &= 2f(x)\kappa(dx) \end{aligned}$$

Thus f satisfies the same differential equation as u_{\downarrow} . Moreover $|f|$ is dominated by

$$\tilde{u}_{\downarrow}(x) + u_{\downarrow}(x) \int_{y > x} G(y, y)(\tilde{\kappa}(dy) - \kappa(dy))$$

Thus f is bounded on the intervals of the type $(a, +\infty)$. It follows that there is a constant $c_2 \in \mathbb{R}$ such that $f \equiv c_2 u_{\downarrow}$. Thus we get the identity (7.1.5). Let's show that $c_2 > 0$. Let $x \in \text{Supp}(\tilde{\kappa})$. Then

$$\begin{aligned} 1 - \frac{1}{\tilde{u}_{\downarrow}(x)} \int_{y > x} (u_{\uparrow}(y)u_{\downarrow}(x) - u_{\downarrow}(y)u_{\uparrow}(x))\tilde{u}_{\downarrow}(y)(\tilde{\kappa}(dy) - \kappa(dy)) \\ = 1 - \int_{y > x} f_{\Delta \tilde{Y}}(x, y)(\tilde{\kappa}(dy) - \kappa(dy)) = \mathbb{P}(\Delta \tilde{Y} \cap (x, +\infty) = \emptyset | x \in \tilde{Y}_{\infty}) \end{aligned}$$

The above conditional probability is positive because according to the lemma 7.3, $\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset) > 0$. Thus f is positive and $c_2 > 0$. \square

Lemma 7.5. — *Conditionally on the event $\Delta\tilde{\mathcal{Y}} = \emptyset$, $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ has the same law as $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$.*

Proof. — It is enough to show that conditionally on $\Delta\tilde{\mathcal{Y}} = \emptyset$, $\tilde{\mathcal{Y}}_\infty$ has the same law as \mathcal{Y}_∞ . Indeed in both cases the points of $\tilde{\mathcal{Z}}_\infty$ respectively \mathcal{Z}_∞ are distributed independently and uniformly between any two consecutive points of $\tilde{\mathcal{Y}}_\infty$ respectively \mathcal{Y}_∞ . For $n \geq 1$ and $y_1 < \dots < y_n$, let $\rho_n(dy_1, \dots, dy_n)$ be the infinitesimal probability for $\tilde{\mathcal{Y}}_\infty$ having a point at each of the locations y_i and none in-between, conditionally on $\Delta\tilde{\mathcal{Y}} = \emptyset$. We need only to show that

$$(7.1.6) \quad \rho_n(dy_1, \dots, dy_n) = 2^{n-1} u_\uparrow(y_1) u_\downarrow(y_n) \prod_{i=2}^n (y_i - y_{i-1}) \prod_{i=1}^n \kappa(dy_i)$$

For $y_1 < \dots < y_n$ to be n consecutive points in $\tilde{\mathcal{Y}}_\infty$ and for $\Delta\tilde{\mathcal{Y}} = \emptyset$, we need $y_1 < \dots < y_n$ to be n consecutive points in $\tilde{\mathcal{Y}}_\infty$, to choose not to erase any of y_i (probability $\chi(y_i)$) and finally we need that $\Delta\tilde{\mathcal{Y}} \cap (-\infty, y_1) = \emptyset$ and $\Delta\tilde{\mathcal{Y}} \cap (y_n, +\infty) = \emptyset$. Thus

$$\begin{aligned} \rho_n(dy_1, \dots, dy_n) &= \frac{1}{\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset)} 2^{n-1} \tilde{u}_\uparrow(y_1) \tilde{u}_\downarrow(y_n) \\ &\times \left(1 - \frac{1}{\tilde{u}_\uparrow(y_1)} \int_{y < y_1} (u_\downarrow(y) u_\uparrow(y_1) - u_\uparrow(y) u_\downarrow(y_1)) \tilde{u}_\uparrow(y) (\tilde{\kappa}(dy) - \kappa(dy)) \right) \\ &\times \left(1 - \frac{1}{\tilde{u}_\downarrow(y_n)} \int_{y > y_n} (u_\uparrow(y) u_\downarrow(y_n) - u_\downarrow(y) u_\uparrow(y_n)) \tilde{u}_\downarrow(y) (\tilde{\kappa}(dy) - \kappa(dy)) \right) \\ &\times \prod_{i=2}^n (y_i - y_{i-1}) \prod_{i=1}^n \chi(y_i) \tilde{\kappa}(dy_i) \end{aligned}$$

Applying lemma 7.4 we get that

$$\rho_n(dy_1, \dots, dy_n) = \frac{c_1 c_2}{\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset)} 2^{n-1} u_\uparrow(y_1) u_\downarrow(y_n) \prod_{i=2}^n (y_i - y_{i-1}) \prod_{i=1}^n \kappa(dy_i)$$

Since the constant $\frac{c_1 c_2}{\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset)}$ does not depend on n , the previous equations implies that

$$\mathbb{P}(\tilde{\mathcal{Y}}_\infty \neq \emptyset | \Delta\tilde{\mathcal{Y}} = \emptyset) = \frac{c_1 c_2}{\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset)} \mathbb{P}(\mathcal{Y}_\infty \neq \emptyset)$$

But $\mathbb{P}(\tilde{\mathcal{Y}}_\infty \neq \emptyset | \Delta\tilde{\mathcal{Y}} = \emptyset) = \mathbb{P}(\mathcal{Y}_\infty \neq \emptyset) = 1$. Thus

$$\frac{c_1 c_2}{\mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset)} = 1$$

and 7.1.6 holds. \square

Corollary 7.6. — *Let $a < b \in \mathbb{R}$ such that $\tilde{\kappa}(\mathbb{R} \setminus [a, b]) > 0$. Conditionally on $\tilde{\mathcal{Y}}_\infty \cap [a, b] = \emptyset$, $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ has the same law as the pair of interwoven determinantal*

point processes obtained from the Wilson's algorithm applied to the Brownian motion with killing measure $1_{\mathbb{R} \setminus [a, b]} \kappa$.

Lemma 7.7. — *Conditionally on $\sharp \Delta \tilde{\mathcal{Y}} = 1$ and on the position of the unique point Y in $\Delta \tilde{\mathcal{Y}}$, $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ has the same law as $(\mathcal{Y}_\infty^{(Y)}, \mathcal{Z}_\infty^{(Y)})$.*

Proof. — It is enough to show that conditionally on $\sharp \Delta \tilde{\mathcal{Y}} = 1$ and on the position of the unique point Y in $\Delta \tilde{\mathcal{Y}}$, $\tilde{\mathcal{Y}}_\infty$ has the same law as $\mathcal{Y}_\infty^{(Y)}$. Indeed the points of $\tilde{\mathcal{Z}}_\infty$ respectively $\mathcal{Z}_\infty^{(Y)}$ are independently and uniformly distributed between any two consecutive points in $\tilde{\mathcal{Y}}_\infty$ respectively $\mathcal{Z}_\infty^{(Y)}$.

Let $n \geq 1$ and $i_0 \in \{1, \dots, n\}$. Let $y_1 < \dots < y_n \in \mathbb{R}$. The infinitesimal probability for y_1, \dots, y_n being n consecutive points in $\tilde{\mathcal{Y}}_\infty$ and $\Delta \tilde{\mathcal{Y}} = \{y_{i_0}\}$ is

$$\begin{aligned}
& 2^{n-1} \tilde{u}_\uparrow(y_1) \tilde{u}_\downarrow(y_n) \\
& \times \left(1 - \frac{1}{\tilde{u}_\uparrow(y_1)} \int_{y < y_1} (u_\downarrow(y) u_\uparrow(y_1) - u_\uparrow(y) u_\downarrow(y_1)) \tilde{u}_\uparrow(y) (\tilde{\kappa}(dy) - \kappa(dy)) \right) \\
& \times \left(1 - \frac{1}{\tilde{u}_\downarrow(y_n)} \int_{y > y_n} (u_\uparrow(y) u_\downarrow(y_n) - u_\downarrow(y) u_\uparrow(y_n)) \tilde{u}_\downarrow(y) (\tilde{\kappa}(dy) - \kappa(dy)) \right) \\
& \times \prod_{i=2}^n (y_i - y_{i-1}) \prod_{i \neq i_0} \kappa(dy_i) \times (\tilde{\kappa} - \kappa)(dy_{i_0}) \\
& = c_1 c_2 2^{n-1} u_\uparrow(y_1) u_\downarrow(y_n) \prod_{i=2}^n (y_i - y_{i-1}) \prod_{i \neq i_0} \kappa(dy_i) \times (\tilde{\kappa} - \kappa)(dy_{i_0}) \\
& = v_{\kappa, \tilde{\kappa}}(y_{i_0}) (\tilde{\kappa} - \kappa)(dy_{i_0}) \times 2^{i_0-1} \frac{u_\uparrow(y_1)}{u_\uparrow(y_{i_0})} \prod_{i=1}^{i_0-1} (y_{i+1} - y_i) \kappa(dy_i) \\
& \times 2^{n-i_0} \frac{u_\downarrow(y_n)}{u_\downarrow(y_{i_0})} \prod_{i=i_0+1}^n (y_i - y_{i-1}) \kappa(dy_i)
\end{aligned} \tag{7.1.7}$$

In 7.1.7 appears the infinitesimal probability for $\Delta \tilde{\mathcal{Y}} = \{y_{i_0}\}$ times the infinitesimal probability for y_1, \dots, y_n being n consecutive points in $\mathcal{Y}_\infty^{(y_0)}$ (compare with expressions 7.1.1 and 7.1.2). \square

7.2. Couplings

In this section we will prove the monotone coupling results for $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ stated at the beginning of section 7.1. The construction of the coupling will be explicit. However it will not appeal to Wilson's algorithm used to define $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$. First we will describe analogous monotone coupling results for Uniform Spanning Trees on finite graphs. In this case no explicit construction is known in general and the proof relies on Strassen's theorem and the conditions for stochastic domination between determinantal processes shown in [20].

Proposition 7.8. — *Let \mathbb{G} be a finite connected undirected graph with E its set of edges, and $(C(e))_{e \in E}$ a positive weight function on E . Let F be a subset of E . Let $(\tilde{C}(e))_{e \in E}$ be another weight function such $\tilde{C} \geq C$ and $\tilde{C} = C$ on $E \setminus F$. Let Υ be the Uniform Spanning Tree of \mathbb{G} corresponding to the weights C and $\tilde{\Upsilon}$ the Uniform Spanning Tree of \mathbb{G} corresponding to the weights \tilde{C} . There is a coupling of Υ and $\tilde{\Upsilon}$ such that*

$$(7.2.1) \quad \tilde{\Upsilon} \cap (E \setminus F) \subseteq \Upsilon \cap (E \setminus F)$$

In case F is made of all edges adjacent to a particular vertex x_0 , and \tilde{C} is proportional to C on F , then there is a coupling satisfying the additional condition

$$(7.2.2) \quad \Upsilon \cap F \subseteq \tilde{\Upsilon} \cap F$$

Proof. — It is enough to prove the first coupling ((7.2.1)) in case F is a single edge ($F = \{e\}$). Then by induction on $\sharp F$ the general result will follow. From definition of Uniform Spanning Trees is clear that $\mathbb{P}(e \in \Upsilon) \leq \mathbb{P}(e \in \tilde{\Upsilon})$. Moreover, Υ conditionally on $e \in \Upsilon$ respectively $e \notin \Upsilon$ has the same law as $\tilde{\Upsilon}$ conditionally on $e \in \tilde{\Upsilon}$ respectively $e \notin \tilde{\Upsilon}$. A possible coupling is the following: first we couple $1_{e \in \Upsilon}$ with $1_{e \in \tilde{\Upsilon}}$ in a way such that $1_{e \in \Upsilon} \leq 1_{e \in \tilde{\Upsilon}}$. In case $1_{e \in \Upsilon} = 1_{e \in \tilde{\Upsilon}} = 0$ respectively $1_{e \in \Upsilon} = 1_{e \in \tilde{\Upsilon}} = 1$ we sample for both Υ and $\tilde{\Upsilon}$ the same tree having the law of Υ conditioned by $e \notin \Upsilon$ respectively $e \in \Upsilon$. In case $1_{e \in \Upsilon} = 0$ and $1_{e \in \tilde{\Upsilon}} = 1$, we use the fact that on the edges in $E \setminus \{e\}$, the law of Υ conditioned by $e \in \Upsilon$ is stochastically dominated by the law of Υ conditioned by $e \notin \Upsilon$, which implies the existence of a monotone coupling by Strassen's theorem. See theorems 5.2, 5.3 and 5.5 in [20].

Now we consider the case of F made of all edges adjacent to a particular vertex x_0 , and \tilde{C} is proportional to C on F . Let $(\Upsilon, \tilde{\Upsilon})$ be a coupling satisfying (7.2.1). In general it does not satisfy (7.2.2). To deal with this issue we will re-sample the edges of Υ and $\tilde{\Upsilon}$ contained in F , that is to say sample Υ' having the same law as Υ , $\tilde{\Upsilon}'$ having the same law as $\tilde{\Upsilon}$, such that $\Upsilon' \cap (E \setminus F) = \Upsilon \cap (E \setminus F)$, $\tilde{\Upsilon}' \cap (E \setminus F) = \tilde{\Upsilon} \cap (E \setminus F)$ and such that $\Upsilon' \cap F \subseteq \tilde{\Upsilon}' \cap F$. Let $\mathcal{T}_1, \dots, \mathcal{T}_N$ be the connected components of $\Upsilon \cap (E \setminus F)$. (7.2.1) ensures that each connected component of $\tilde{\Upsilon}' \cap (E \setminus F)$ is contained in one of the \mathcal{T}_i . Let $\mathcal{T}_{1,1}, \dots, \mathcal{T}_{1,q_1}, \dots, \mathcal{T}_{N,1}, \dots, \mathcal{T}_{N,q_N}$ be the connected components of $\tilde{\Upsilon}' \cap (E \setminus F)$, where $\mathcal{T}_{i,j} \subseteq \mathcal{T}_i$. Conditionally on $\mathcal{T}_1, \dots, \mathcal{T}_N$, $\Upsilon \cap F$ has the following law: for each \mathcal{T}_i one chooses an edge connecting x_0 to \mathcal{T}_i with probability proportional to C , and independently from the edges of Υ that will connect x_0 to other $(\mathcal{T}_{i'})_{i' \neq i}$. Similarly for the law of $\tilde{\Upsilon}$ conditionally on $\mathcal{T}_{1,1}, \dots, \mathcal{T}_{1,q_1}, \dots, \mathcal{T}_{N,1}, \dots, \mathcal{T}_{N,q_N}$. To construct Υ' and $\tilde{\Upsilon}'$ we use the fact that \tilde{C} is proportional to C on F :

- We start with Υ and $\tilde{\Upsilon}$ satisfying (7.2.1).
- Then we remove from Υ and $\tilde{\Upsilon}$ the edges contained in F .
- For each $\mathcal{T}_{i,j}$, we add to $\tilde{\Upsilon}'$ an edge connecting x_0 to $\mathcal{T}_{i,j}$, chosen proportionally to its weight under C , each choice being independent from the others.

- For each $i \in \{1, \dots, N\}$, there are q_i edges in $\tilde{\Upsilon}'$ connecting x_0 to \mathcal{T}_i , one for each $(\mathcal{T}_{i,j})_{1 \leq j \leq q_i}$. In order to construct Υ' , we need to choose one out of q_i to keep and remove the others. We chose to keep the edge corresponding to $\mathcal{T}_{i,j}$ with probability proportional to:

$$\sum_{\substack{e \text{ connecting} \\ x_0 \text{ to } \mathcal{T}_{i,j}}} C(e)$$

The choice is done independently for each $i \in \{1, \dots, N\}$.

By construction $\Upsilon' \cap F \subseteq \tilde{\Upsilon}' \cap F$. \square

Consider now two different killing measures κ and $\tilde{\kappa}$ on \mathbb{R} , with $\kappa \leq \tilde{\kappa}$, and the couples of determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ respectively $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ corresponding to the Brownian motion on \mathbb{R} with killing measure κ respectively $\tilde{\kappa}$. We want to show that one can couple $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ on the same probability space such that $\mathcal{Z}_\infty \subseteq \tilde{\mathcal{Z}}_\infty$ and $\tilde{\mathcal{Y}}_\infty \subseteq \mathcal{Y}_\infty \cup \text{Supp}(\tilde{\kappa} - \kappa)$, and if κ and $\tilde{\kappa}$ are proportional also have $\mathcal{Y}_\infty \subseteq \tilde{\mathcal{Y}}_\infty$. The condition $\mathcal{Z}_\infty \subseteq \tilde{\mathcal{Z}}_\infty$ and $\tilde{\mathcal{Y}}_\infty \subseteq \mathcal{Y}_\infty \cup \text{Supp}(\tilde{\kappa} - \kappa)$ is analogous to (7.2.1). The condition $\mathcal{Y}_\infty \subseteq \tilde{\mathcal{Y}}_\infty$ is analogous to (7.2.2), where the cemetery \dagger plays the role of the distinguished vertex x_0 . We used the stochastic domination principle ([20]) for determinantal point process with determinantal kernel a projection operator. It ensures the existence of a monotone coupling but does not give one explicitly (see open questions [20]). However for $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ we will construct a whole family of rather explicit monotone couplings.

Let \tilde{G} be the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - \tilde{\kappa}$, factorized as

$$\tilde{G}(x, y) = \tilde{u}_\uparrow(x \wedge y) \tilde{u}_\downarrow(x \vee y)$$

Let

$$\tilde{\mathcal{K}}(y, z) := -\frac{1}{2} \frac{d\tilde{u}_\uparrow}{dx}((y \wedge z)^+) \frac{d\tilde{u}_\downarrow}{dx}((y \vee z)^-)$$

Let $\mathfrak{G}_{\tilde{\kappa}}$ be the operator on $\mathbb{L}^2(d\tilde{\kappa})$ defined on functions with compact support as follows:

$$(\mathfrak{G}_{\tilde{\kappa}} f)(x) := \int_{\mathbb{R}} \tilde{G}(x, y) f(y) \tilde{\kappa}(dy)$$

In case $\tilde{\kappa} = c\kappa$ where c is a constant, $c > 1$, we have the following resolvent identity, which follows from lemma 2.8:

$$(7.2.3) \quad \frac{1}{c} \mathfrak{G}_{c\kappa} \mathfrak{G}_\kappa = \frac{1}{c} \mathfrak{G}_\kappa \mathfrak{G}_{c\kappa} = \frac{1}{c-1} \left(\mathfrak{G}_\kappa - \frac{1}{c} \mathfrak{G}_{c\kappa} \right)$$

Next we prove that a simple necessary but not sufficient condition for monotone couplings to exist is satisfied. It won't be used in the sequel but we prefer to give a direct proof for it.

Proposition 7.9. — *For any $z_1, \dots, z_n \in \mathbb{R}$ such that $\tilde{\kappa}(\{z_i\}) = 0$*

$$(7.2.4) \quad \det(\tilde{\mathcal{K}}(z_i, z_j))_{1 \leq i, j \leq n} \geq \det(\mathcal{K}(z_i, z_j))_{1 \leq i, j \leq n}$$

If $\tilde{\kappa} = c\kappa$, $c > 1$, then for any $y_1, \dots, y_n \in \text{Supp}(\kappa)$

$$(7.2.5) \quad c^n \det(\tilde{G}(y_i, y_j))_{1 \leq i, j \leq n} \geq \det(G(y_i, y_j))_{1 \leq i, j \leq n}$$

Proof. — We will first show (7.2.4). To begin with we will show that for any $z_1 \in \mathbb{R}$, $\tilde{\mathcal{K}}(z_1, z_1) \geq \mathcal{K}(z_1, z_1)$. The Wronskian

$$W(u_\uparrow, \tilde{u}_\uparrow)(z) := u_\uparrow(z) \frac{d\tilde{u}_\uparrow}{dx}(z^+) - \tilde{u}_\uparrow(z) \frac{du_\uparrow}{dx}(z^+)$$

is non-negative. Indeed $W(u_\uparrow, \tilde{u}_\uparrow)(-\infty) = 0$ and

$$dW(u_\uparrow, \tilde{u}_\uparrow) = 2u_\uparrow \tilde{u}_\uparrow (d\tilde{\kappa} - d\kappa) \geq 0$$

Similarly the Wronskian

$$W(u_\downarrow, \tilde{u}_\downarrow)(z) := u_\downarrow(z) \frac{d\tilde{u}_\downarrow}{dx}(z^+) - \tilde{u}_\downarrow(z) \frac{du_\downarrow}{dx}(z^+)$$

is non-positive. Using the fact that

$$W(u_\downarrow, u_\uparrow) = W(\tilde{u}_\downarrow, \tilde{u}_\uparrow) \equiv 2$$

we get

$$\begin{aligned} \tilde{\mathcal{K}}(z_1, z_1) - \mathcal{K}(z_1, z_1) &= \frac{1}{2} \left(\frac{du_\uparrow}{dx}(z_1^+) \frac{du_\downarrow}{dx}(z_1^+) - \frac{d\tilde{u}_\uparrow}{dx}(z_1^+) \frac{d\tilde{u}_\downarrow}{dx}(z_1^+) \right) \\ &= \frac{1}{4} \left(\frac{du_\uparrow}{dx}(z_1^+) \frac{du_\downarrow}{dx}(z_1^+) W(\tilde{u}_\downarrow, \tilde{u}_\uparrow) - \frac{d\tilde{u}_\uparrow}{dx}(z_1^+) \frac{d\tilde{u}_\downarrow}{dx}(z_1^+) W(u_\downarrow, u_\uparrow) \right) \\ &= \frac{1}{4} \left(\frac{du_\downarrow}{dx}(z_1^+) \frac{d\tilde{u}_\downarrow}{dx}(z_1^+) W(u_\uparrow, \tilde{u}_\uparrow)(z_1) \right. \\ &\quad \left. - \frac{du_\uparrow}{dx}(z_1^+) \frac{d\tilde{u}_\uparrow}{dx}(z_1^+) W(u_\downarrow, \tilde{u}_\downarrow)(z_1) \right) \geq 0 \end{aligned}$$

To prove (7.2.4) in general, we will use the factorization (6.3.34). For $x_0 < z$, let

$$\tilde{u}_\uparrow^{(x_0 \triangleright)}(z) := \tilde{u}_\uparrow(z) + \left(\frac{d\tilde{u}_\downarrow}{dx}(x_0^-) \right)^{-1} \frac{d\tilde{u}_\uparrow}{dx}(x_0^-) \tilde{u}_\downarrow(z)$$

Factorization (6.3.34) ensures that we only need to prove that for $x_0 < z$ with $\kappa(\{x_0\}) = 0$:

$$-\frac{d\tilde{u}_\uparrow^{(x_0 \triangleright)}}{dx}(z^+) \frac{d\tilde{u}_\downarrow}{dx}(z) \geq -\frac{du_\uparrow^{(x_0 \triangleright)}}{dx}(z^+) \frac{du_\downarrow}{dx}(z)$$

First observe that the Wronskian

$$W(u_\uparrow^{(x_0 \triangleright)}, \tilde{u}_\uparrow^{(x_0 \triangleright)})(z) := u_\uparrow^{(x_0 \triangleright)}(z) \frac{d\tilde{u}_\uparrow^{(x_0 \triangleright)}}{dx}(z^+) - \tilde{u}_\uparrow^{(x_0 \triangleright)}(z) \frac{du_\uparrow^{(x_0 \triangleright)}}{dx}(z^+)$$

is non-negative on $[x_0, +\infty)$. Indeed $W(u_\uparrow^{(x_0 \triangleright)}, \tilde{u}_\uparrow^{(x_0 \triangleright)})(x) = 0$ and

$$dW(u_\uparrow^{(x_0 \triangleright)}, \tilde{u}_\uparrow^{(x_0 \triangleright)}) = 2u_\uparrow^{(x_0 \triangleright)}(z) \tilde{u}_\uparrow^{(x_0 \triangleright)}(z) (d\tilde{\kappa} - d\kappa) \geq 0$$

The sequel of the proof works as in the previous case.

Let's prove now (7.2.5). First we consider the case $n = 1$. From the resolvent identity (7.2.3) follows that

$$\mathfrak{G}_{c\kappa} - \mathfrak{G}_\kappa = (c - 1)(\mathfrak{G}_\kappa - \mathfrak{G}_{c\kappa}\mathfrak{G}_\kappa)$$

Since $\mathfrak{G}_{c\kappa}$ is contracting, this implies that $\mathfrak{G}_\kappa \leq \mathfrak{G}_{c\kappa}$, where the inequality stands for positive semi-definite operators on $\mathbb{L}^2(d\kappa)$. Let $y_1 \in \text{Supp}(\kappa)$. Then for any $\varepsilon > 0$

$$(7.2.6) \quad c \int_{(y_1-\varepsilon, y_1+\varepsilon)^2} \tilde{G}(x, y)\kappa(dx)\kappa(dy) \geq \int_{(y_1-\varepsilon, y_1+\varepsilon)^2} G(x, y)\kappa(dx)\kappa(dy)$$

Since $y_1 \in \text{Supp}(\kappa)$, both sides of (7.2.6) are positive. The continuity of G and \tilde{G} ensures that $c\tilde{G}(y_1, y_1) \geq G(y_1, y_1)$. In case of general n , we use the factorization (6.3.33). It is enough to prove that for any $x_0 < y$, $y \in \text{Supp}(\kappa)$

$$(7.2.7) \quad c\tilde{G}^{(x_0 \times)}(y, y) \geq G^{(x_0 \times)}(y, y)$$

where

$$\tilde{G}^{(x_0 \times)}(y, y) := \tilde{G}(y, y) - \frac{\tilde{G}(x_0, y)^2}{\tilde{G}(x_0, x_0)}$$

\tilde{G} is the restriction to $(x_0, +\infty)^2$ of the Green's function of $\frac{1}{2}\frac{d^2}{dx^2} - 1_{(x_0, +\infty)}\tilde{\kappa}$. Let $\mathfrak{G}_\kappa^{(x_0 \times)}$ and $\mathfrak{G}_{c\kappa}^{(x_0 \times)}$ be the operators on $\mathbb{L}^2(1_{(x_0, +\infty)}d\kappa)$ defined for functions f with compact support as

$$\begin{aligned} (\mathfrak{G}_\kappa^{(x_0 \times)} f)(x) &:= \int_{(x_0, +\infty)} G^{(x_0 \times)}(x, y)f(y)\kappa(dy) \\ (\mathfrak{G}_{c\kappa}^{(x_0 \times)} f)(x) &:= c \int_{(x_0, +\infty)} \tilde{G}^{(x_0 \times)}(x, y)f(y)\kappa(dy) \end{aligned}$$

$\mathfrak{G}_\kappa^{(x_0 \times)}$ and $\mathfrak{G}_{c\kappa}^{(x_0 \times)}$ are contractions and satisfy a resolvent identity similar to (7.2.3), which similarly implies (7.2.7). \square

The resolvent identity (7.2.3) implies that \mathfrak{G}_κ and $\mathfrak{G}_{c\kappa}$ commute and that $\mathfrak{G}_\kappa \leq \mathfrak{G}_{c\kappa}$. It was shown in case of determinantal point processes on discrete space that this a sufficient condition for a monotone coupling to exist. See theorem 7.1 in [20].

To construct the couplings we will give several procedures that take deterministic arguments, among which pairs of interwoven sets of points, and return pairs of interwoven random point processes. The first procedure we describe will be used as sub-procedure in subsequent procedures.

Procedure 7.10. — *Arguments:*

- a pair $(\mathcal{Y}, \mathcal{Z})$ of disjoint discrete sets of points in \mathbb{R} such that between any two points in \mathcal{Y} lies a single point in \mathcal{Z} and vice-versa, and such that $\inf \mathcal{Y} \cup \mathcal{Z} \in \mathcal{Y} \cup \{-\infty\}$, $\sup \mathcal{Y} \cup \mathcal{Z} \in \mathcal{Y} \cup \{+\infty\}$
- a positive Radon measure κ
- a point $y_0 \in \mathbb{R}$ such that $y_0 \notin \mathcal{Z}$

Procedure:

- (i) If $y_0 \notin \mathcal{Y}$, we define a random variable Z distributed as follows:
 - (i a) If there are $y' \in \mathcal{Y}$, $z' \in \mathcal{Z} \cup \{+\infty\}$, such that $y' < z'$, $y_0 \in (y', z')$ and $\mathcal{Y} \cap (y', z') = \mathcal{Z}_\infty \cap (y', z') = \emptyset$ then Z is distributed according to

$$\frac{1_{z \in (y', y_0)}}{u_\uparrow(y_0) - u_\uparrow(y')} \frac{du_\uparrow}{dx}(z) dz$$
 - (i b) If there are $y' \in \mathcal{Y}$, $z' \in \mathcal{Z} \cup \{-\infty\}$, such that $z' < y'$, $y_0 \in (z', y')$ and $\mathcal{Y} \cap (z', y') = \mathcal{Z} \cap (z', y') = \emptyset$ then Z is distributed according to

$$\frac{-1_{z \in (y_0, y')}}{u_\downarrow(y') - u_\downarrow(y_0)} \frac{du_\downarrow}{dx}(z) dz$$
- (ii) If there are $y' \in \mathcal{Y}$, $z' \in \mathcal{Z} \cup \{+\infty\}$, such that $y' < z'$, $y_0 \in (y', z')$ and $\mathcal{Y} \cap (y', z') = \mathcal{Z} \cap (y', z') = \emptyset$, then
 - (ii a) with probability $\frac{u_\uparrow(y')}{u_\uparrow(y_0)}$ we set

$$(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}}) = (\mathcal{Y} \cup \{y_0\} \setminus \{y'\}, \mathcal{Z})$$
 - (ii b) and with probability $1 - \frac{u_\uparrow(y')}{u_\uparrow(y_0)}$ we set

$$(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}}) = (\mathcal{Y} \cup \{y_0\}, \mathcal{Z} \cup \{Z\})$$
- (iii) If there are $y' \in \mathcal{Y}$, $z' \in \mathcal{Z} \cup \{-\infty\}$, such that $z' < y'$, $y_0 \in (z', y')$ and $\mathcal{Y} \cap (z', y') = \mathcal{Z} \cap (z', y') = \emptyset$, then
 - (iii a) with probability $\frac{u_\downarrow(y')}{u_\downarrow(y_0)}$ we set

$$(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}}) = (\mathcal{Y} \cup \{y_0\} \setminus \{y'\}, \mathcal{Z})$$
 - (iii b) and with probability $1 - \frac{u_\downarrow(y')}{u_\downarrow(y_0)}$ we set

$$(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}}) = (\mathcal{Y} \cup \{y_0\}, \mathcal{Z} \cup \{Z\})$$
- (iv) If $y_0 \in \mathcal{Y}$, we set $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}}) = (\mathcal{Y}, \mathcal{Z})$.

Return: $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$.

Lemma 7.11. — If procedure 7.10 is applied to the pair of interwoven determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ corresponding to the killing measure κ , then its result $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ has the same law as $(\mathcal{Y}_\infty^{(y_0)}, \mathcal{Z}_\infty^{(y_0)})$.

Proof. — By construction $y_0 \in \tilde{\mathcal{Y}}$. Let $\tilde{Z}_1 < \tilde{Y}_1 < \dots < \tilde{Z}_n < \tilde{Y}_n$ be the $2n$ closest points to y_0 in $(\tilde{\mathcal{Y}} \cup \tilde{\mathcal{Z}}) \cap (y_0, +\infty)$. On the event $\min(\mathcal{Y}_\infty \cup \mathcal{Z}_\infty) \cap (y_0, +\infty) \in \mathcal{Z}_\infty$ (point (ii) in procedure 7.10) their distribution is given by

$$(7.2.8) \quad 1_{y_0 < z_1 < y_1 < \dots < z_n < y_n} 2^n \left(\int_{(-\infty, y_0)} u_\uparrow(y') \kappa(dy') \right) u_\downarrow(y_n) dz_1 \kappa(dy_1) \dots dz_n \kappa(dy_n)$$

On the event $\min(\mathcal{Y}_\infty \cup \mathcal{Z}_\infty) \cap (y_0, +\infty) \in \mathcal{Y}_\infty$ (point (iii) in procedure 7.10), the distribution of $\min(\mathcal{Y}_\infty \cup \mathcal{Z}_\infty) \cap (y_0, +\infty)$ is (see proposition 6.11)

$$\begin{aligned} & \mathbf{1}_{y' > y_0} 2 \left(\int_{(-\infty, y_0)} u_\uparrow(y_{-1})(y_0 - y_{-1}) \kappa(dy_{-1}) \right) u_\downarrow(y') \kappa(dy') \\ & \quad + \mathbf{1}_{y' > y_0} \frac{u_\uparrow(+\infty)}{u_\uparrow(y')} G(y_0, y_0) \kappa(dy') \\ & = \mathbf{1}_{y' > y_0} (u_\uparrow(y_0) - u_\uparrow(+\infty)) u_\downarrow(y') \kappa(dy') + \mathbf{1}_{y' > y_0} u_\uparrow(+\infty) u_\downarrow(y') \kappa(dy') \\ & = \mathbf{1}_{y' > y_0} u_\uparrow(y_0) u_\downarrow(y') \kappa(dy') \end{aligned}$$

Thus on the event $\min(\mathcal{Y}_\infty \cup \mathcal{Z}_\infty) \cap (y_0, +\infty) \in \mathcal{Y}_\infty$ (point (iii) in procedure 7.10), the distribution of $(Z_1, Y_1, \dots, Z_n, Y_n)$ is

$$\begin{aligned} & \mathbf{1}_{y_0 < z_1 < \dots < y_n} \left(\int_{y_0 < y' < z_1} \frac{u_\downarrow(y')}{u_\downarrow(y_0)} u_\uparrow(y_0) u_\downarrow(y') 2^n \frac{u_\downarrow(y_n)}{u_\downarrow(y')} \kappa(dy') \right) dz_1 \kappa(dy_1) \dots dz_n \kappa(dy_n) \\ & \quad + \mathbf{1}_{y_0 < z_1 < \dots < y_n} \frac{-1}{u_\downarrow(y_0)} \frac{du_\downarrow}{dx}(z_1) u_\uparrow(y_0) u_\downarrow(y_1) 2^{n-1} \frac{u_\downarrow(y_n)}{u_\downarrow(y_1)} dz_1 \kappa(dy_1) \dots dz_n \kappa(dy_n) \end{aligned} \quad (7.2.9)$$

The term (7.2.9) corresponds to the case when a point is removed from \mathcal{Y}_∞ (case (iii a) in procedure 7.10) and (7.2.10) to the case when Z is added to \mathcal{Z}_∞ (case (iii b) in procedure 7.10). The sum of the densities that appear in (7.2.8), (7.2.9) and (7.2.10) is

$$\begin{aligned} & 2^n \left(\int_{(-\infty, y_0)} u_\uparrow(y') \kappa(dy') \right) u_\downarrow(y_n) + \left(\int_{y_0 < y' < z_1} \frac{u_\downarrow(y')}{u_\downarrow(y_0)} u_\uparrow(y_0) u_\downarrow(y') 2^n \frac{u_\downarrow(y_n)}{u_\downarrow(y')} \kappa(dy') \right) \\ & \quad + \frac{-1}{u_\downarrow(y_0)} \frac{du_\downarrow}{dx}(z_1) u_\uparrow(y_0) u_\downarrow(y_1) 2^{n-1} \frac{u_\downarrow(y_n)}{u_\downarrow(y_1)} \\ & = 2^{n-1} \frac{du_\uparrow}{dx}(y_0) u_\downarrow(y_n) + \frac{2^{n-1}}{u_\downarrow(y_0)} \left(\frac{du_\downarrow}{dx}(z_1) - \frac{du_\downarrow}{dx}(y_0^+) \right) u_\uparrow(y_0) u_\downarrow(y_n) \\ & \quad + \frac{-2^{n-1}}{u_\downarrow(y_0)} \frac{du_\downarrow}{dx}(z_1) u_\uparrow(y_0) u_\downarrow(y_n) \\ & = 2^{n-1} u_\downarrow(y_n) \left(\frac{du_\uparrow}{dx}(y_0) + \frac{-1}{u_\downarrow(y_0)} \frac{du_\downarrow}{dx}(y_0^+) u_\uparrow(y_0) \right) = 2^n \frac{u_\downarrow(y_n)}{u_\downarrow(y_0)} \end{aligned}$$

So we obtain the density which appears in (7.1.1).

It remains to prove that $(\tilde{\mathcal{Y}} \cap (y_0, +\infty), \tilde{\mathcal{Z}} \cap (y_0, +\infty))$ and $(\tilde{\mathcal{Y}} \cap (-\infty, y_0), \tilde{\mathcal{Z}} \cap (-\infty, y_0))$ are independent. Let $Z_{-1} > Y_{-1} > \dots > Z_{-n'} > Y_{-n'}$ be the n' closest points to y_0 in $(\tilde{\mathcal{Y}} \cup \tilde{\mathcal{Z}}) \cap (-\infty, y_0)$. The distribution of the family of points $(Z_{-1}, Y_{-1}, \dots, Z_{-n'}, Y_{-n'}, Z_1, Y_1, \dots, Z_n, Y_n)$ on the event $\#\{\tilde{\mathcal{Y}} \cap (-\infty, y_0)\} \geq n, \#\{\tilde{\mathcal{Y}} \cap (y_0, +\infty)\} \geq n'$ is

$$\mathbf{1}_{y_0 < y' < z_1} 2^{n+n'} u_\uparrow(y_{-n'}) u_\downarrow(y_n) \frac{u_\downarrow(y')}{u_\downarrow(y_0)} \kappa(dy') - \frac{2^{n+n'-1} u_\uparrow(y_{-n'}) u_\downarrow(y_n)}{u_\downarrow(y_0)} \frac{du_\downarrow}{dx}(z_1)$$

$$(7.2.12) \\ + \int_{z_{-1} < y' < y_0} 2^{n+n'} u_\uparrow(y_{-n'}) u_\downarrow(y_n) \frac{u_\uparrow(y')}{u_\uparrow(y_0)} \kappa(dy') + \frac{2^{n+n'-1} u_\uparrow(y_{-n'}) u_\downarrow(y_n)}{u_\uparrow(y_0)} \frac{du_\uparrow}{dx}(z_{-1}) \\ \times 1_{y_{-n'} < z_{-n'} < \dots < z_{-1} < y_0 < z_1 < \dots < z_n < y_n} \kappa(dy_{-n'}) dz_{-n'} \dots dz_{-1} dz_1 \dots dz_n \kappa(dy_n)$$

The term (7.2.11) corresponds to point (iii) in procedure 7.10 and (7.2.12) to point (ii) in procedure 7.10. One can check that the sum of the densities equals

$$2^{n+n'} \frac{u_\uparrow(y_{-n'})}{u_\uparrow(y_0)} \frac{u_\downarrow(y_n)}{u_\downarrow(y_0)}$$

Thus $(\tilde{\mathcal{Y}} \cap (y_0, +\infty), \tilde{\mathcal{Z}} \cap (y_0, +\infty))$ and $(\tilde{\mathcal{Y}} \cap (-\infty, y_0), \tilde{\mathcal{Z}} \cap (-\infty, y_0))$ are independent. \square

Lemma 7.12. — *We consider the subspace of triples $((\mathcal{Y}, \mathcal{Z}), \kappa, y_0)$ consisting of a pair of discrete sets of points $(\mathcal{Y}, \mathcal{Z})$, a Radon measure κ and a point $y_0 \in \mathbb{R}$, and which satisfies the restrictions on the arguments of procedure 7.10. We assume this subspace endowed with the product topology obtained from the topology of uniform convergence on compact subsets for the pairs $(\mathcal{Y}, \mathcal{Z})$, the vague topology for the measures κ and standard order topology on \mathbb{R} . If $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ is the pair of point processes obtained by applying procedure 7.10 to the arguments $((\mathcal{Y}, \mathcal{Z}), \kappa, y_0)$, then its law depends continuously on $((\mathcal{Y}, \mathcal{Z}), \kappa, y_0)$.*

Proof. — From lemma 2.4 it follows that the cumulative distribution function of Z (point (i) in procedure 7.10) depends uniformly continuously on $((\mathcal{Y}, \mathcal{Z}), \kappa, y_0)$ in the neighbourhood of triples where $y_0 \notin \mathcal{Y}$. Moreover the probabilities to make either the choice (ii a) or the choice (ii b), as well as to make either the choice (iii a) or the choice (iii b), depend continuously on $((\mathcal{Y}, \mathcal{Z}), \kappa, y_0)$. Thus the law of $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ depends continuously on $((\mathcal{Y}, \mathcal{Z}), \kappa, y_0)$ in the neighbourhood of triples where $y_0 \notin \mathcal{Y}$. Moreover in the neighbourhood of triples where $y_0 \in \mathcal{Y}$, with high probability, converging to 1, $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}}) = (\mathcal{Y}, \mathcal{Z})$. Thus the law of $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ is continuous also at these triples. \square

First we will describe a coupling in case when $\tilde{\kappa}$ and κ differ by an atom: $\tilde{\kappa} = \kappa + c\delta_{y_0}$. We construct the coupling as follows:

Procedure 7.13. — *Arguments:*

- a pair $(\mathcal{Y}, \mathcal{Z})$ of disjoint discrete sets of points in \mathbb{R} such that between any two points in \mathcal{Y} lies a single point in \mathcal{Z} and vice-versa, and such that $\inf \mathcal{Y} \cup \mathcal{Z} \in \mathcal{Y} \cup \{-\infty\}$, $\sup \mathcal{Y} \cup \mathcal{Z} \in \mathcal{Y} \cup \{+\infty\}$
- two positive Radon measures κ and $\tilde{\kappa}$ where $\tilde{\kappa}$ is of form $\tilde{\kappa} = \kappa + c\delta_{y_0}$ and $y_0 \notin \mathcal{Z}$.

Procedure:

- (i) Let β be a Bernoulli r.v. of parameter $c\tilde{G}(y_0, y_0)$.
- (ii) If $\beta = 0$ we set $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}}) = (\mathcal{Y}, \mathcal{Z})$.

- (iii) If $\beta = 1$, we apply the procedure 7.10 to the arguments $(\mathcal{Y}, \mathcal{Z})$, κ and y_0 and set $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ to be its result.

Return: $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$.

$(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ constructed this way satisfies the following: between any two consecutive points in $\tilde{\mathcal{Y}}$ lies a single point in $\tilde{\mathcal{Z}}$ and between any two consecutive points in $\tilde{\mathcal{Z}}$ lies a point in $\tilde{\mathcal{Y}}$. By construction $\mathcal{Z} \subseteq \tilde{\mathcal{Z}}$ and $\tilde{\mathcal{Y}} \subseteq \mathcal{Y} \cup \{y_0\}$.

Proposition 7.14. — *If procedure 7.13 is applied to the pair of interwoven determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ corresponding to the measure κ , then the returned pair of point processes $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ has the law of the interwoven determinantal point processes $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ corresponding to $\tilde{\kappa} = \kappa + c\delta_{y_0}$.*

Proof. — Observe that a.s. $y_0 \notin \mathcal{Z}_\infty$. First we deal with the case $\kappa(\{y_0\}) = 0$. Then almost surely $y_0 \notin \mathcal{Y}_\infty$ and $y_0 \in \tilde{\mathcal{Y}}$ if and only if $\beta = 1$. But

$$\mathbb{P}(\beta = 1) = \mathbb{P}(y_0 \in \tilde{\mathcal{Y}}) = c\tilde{G}(y_0, y_0)$$

According to corollary 7.6, conditionally on $y_0 \notin \tilde{\mathcal{Y}}$, $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ has the same law as $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$, that is to say the same law as $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ conditionally on $\beta = 0$. According to lemma 7.11, conditionally on $\beta = 1$, $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ follows the same law as $(\tilde{\mathcal{Y}} \cap (-\infty, y_0), \tilde{\mathcal{Z}} \cap (-\infty, y_0))$, which is also the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ conditioned on $y_0 \in \tilde{\mathcal{Y}}_\infty$.

We deal now with the case $\kappa(\{y_0\}) > 0$.

$$\begin{aligned} \mathbb{P}(y_0 \in \tilde{\mathcal{Y}}_\infty) &= \tilde{\kappa}(\{y_0\})\tilde{G}(y_0, y_0) \\ \mathbb{P}(y_0 \in \tilde{\mathcal{Y}}) &= \mathbb{P}(\beta = 1) + \mathbb{P}(\beta = 0, y_0 \in \mathcal{Y}_\infty) \\ &= c\tilde{G}(y_0, y_0) + (1 - c\tilde{G}(y_0, y_0))\kappa(\{y_0\})G(y_0, y_0) \end{aligned}$$

But G and \tilde{G} satisfy the resolvent identity (see lemma 2.8):

$$\tilde{G}(y_0, y_0)\kappa(\{y_0\})G(y_0, y_0) = \frac{\kappa(\{y_0\})}{\tilde{\kappa}(\{y_0\}) - \kappa(\{y_0\})}(G(y_0, y_0) - \tilde{G}(y_0, y_0))$$

It follows that $\mathbb{P}(y_0 \in \tilde{\mathcal{Y}}) = \mathbb{P}(y_0 \in \tilde{\mathcal{Y}}_\infty)$. Let $\tilde{\kappa} := \kappa - \kappa(\{y_0\})\delta_{y_0}$ and $(\check{\mathcal{Y}}_\infty, \check{\mathcal{Z}}_\infty)$ be the interwoven determinantal point processes corresponding to $\tilde{\kappa}$. Let $\tilde{\kappa}' := \tilde{\kappa} - \kappa(\{y_0\})\delta_{y_0}$ and $(\check{\mathcal{Y}}'_\infty, \check{\mathcal{Z}}'_\infty)$ be the interwoven determinantal processes corresponding to $\tilde{\kappa}'$. According to corollary 7.6, $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ conditioned by $y_0 \notin \tilde{\mathcal{Y}}$ has the same law as $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ conditioned by $y_0 \notin \mathcal{Y}_\infty$, which is the same law as $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ conditioned by $y_0 \notin \tilde{\mathcal{Y}}_\infty$, and it is the law of $(\check{\mathcal{Y}}_\infty, \check{\mathcal{Z}}_\infty)$. For $y_0 \in \tilde{\mathcal{Y}}$ there are two possibilities: either $y_0 \in \mathcal{Y}_\infty$ or $y_0 \notin \mathcal{Y}_\infty$ and $\beta = 1$. In the first case, it follows from proposition 7.1 that $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ conditioned on $y_0 \in \mathcal{Y}_\infty$ has the same law as $(\check{\mathcal{Y}}_\infty, \check{\mathcal{Z}}_\infty)$ conditioned on $y_0 \in \check{\mathcal{Y}}_\infty$. In the second case $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ conditioned on $y_0 \notin \mathcal{Y}_\infty$ has the same law as $(\check{\mathcal{Y}}_\infty, \check{\mathcal{Z}}_\infty)$. This brings us back to the situation $\kappa(\{y_0\}) = 0$. According to what was proved earlier, conditionally on $y_0 \notin \mathcal{Y}_\infty$ and $\beta = 1$, $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ has the same

law as $(\tilde{\mathcal{Y}}'_\infty, \tilde{\mathcal{Z}}'_\infty)$ conditioned on $y_0 \in \tilde{\mathcal{Y}}'_\infty$. But this is the same law as for $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ conditioned on $y_0 \in \tilde{\mathcal{Y}}_\infty$. So again, $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ has the same law as $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$. \square

Next we consider the more general case where the measure $\tilde{\kappa} - \kappa$ has a first moment:

$$\int_{\mathbb{R}} |x|(\tilde{\kappa}(dx) - \kappa(dx)) < +\infty$$

First we describe a procedure that does not give a coupling between $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ but allows to approach it.

Procedure 7.15. — *Arguments:*

- a pair $(\mathcal{Y}, \mathcal{Z})$ of disjoint discrete sets of points in \mathbb{R} such that between any two points in \mathcal{Y} lies a single point in \mathcal{Z} and vice-versa, and such that $\inf \mathcal{Y} \cup \mathcal{Z} \in \mathcal{Y} \cup \{-\infty\}$, $\sup \mathcal{Y} \cup \mathcal{Z} \in \mathcal{Y} \cup \{+\infty\}$
- two positive Radon measures $\kappa, \tilde{\kappa}$ such that $\kappa \leq \tilde{\kappa}$ and $\int_{\mathbb{R}} |x|(\tilde{\kappa}(dx) - \kappa(dx)) < +\infty$ and $(\tilde{\kappa} - \kappa)(\mathcal{Z}) = 0$.

Procedure:

- (i) Let β be a Bernoulli r.v. of parameter

$$\int_{\mathbb{R}} v_{\kappa, \tilde{\kappa}}(y)(\tilde{\kappa} - \kappa)(dy)$$

(see notations of proposition 7.3)

- (ii) Let Y be a real r.v. independent from β distributed according to

$$\frac{v_{\kappa, \tilde{\kappa}}(y)(\tilde{\kappa} - \kappa)(dy)}{\mathbb{P}(\beta = 1)}$$

- (iii) If $\beta = 0$ we set $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}}) = (\mathcal{Y}, \mathcal{Z})$.
- (iv) If $\beta = 1$, we apply the procedure 7.10 to the arguments $(\mathcal{Y}, \mathcal{Z})$, κ and Y and set $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ to be its result.

Return: $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$.

Observe that in case $\tilde{\kappa}$ and κ differ only by an atom, procedure 7.15 is the same as procedure 7.13.

Lemma 7.16. — *Let $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ respectively $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ be the pair of interwoven determinantal point processes corresponding to the killing measure κ respectively $\tilde{\kappa}$. We assume that the procedure 7.15 is applied to $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and that $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ is the returned pair of point processes. Then the total variation distance between the law of $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ and the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ is less or equal to $\left(\int_{\mathbb{R}} \tilde{G}(y, y)(\tilde{\kappa}(dy) - \kappa(dy)) \right)^2$.*

Proof. — Let $\Delta\tilde{\mathcal{Y}}$ be the determinantal point process defined in section 7.1 (see lemma 7.3). According to lemma 7.5, the law of $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ conditionally on $\beta = 0$ is the same as the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ conditionally on $\Delta\tilde{\mathcal{Y}} = \emptyset$. From lemmas 7.11 and 7.7 follows

that the law of $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ conditionally on $\beta = 1$ is the same as the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ conditionally on $\sharp\Delta\tilde{\mathcal{Y}} = 1$. Moreover $\mathbb{P}(\beta = 1) = \mathbb{P}(\sharp\Delta\tilde{\mathcal{Y}} = 1)$. However

$$\mathbb{P}(\beta = 0) = \mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset) + \mathbb{P}(\sharp\Delta\tilde{\mathcal{Y}} \geq 2) \geq \mathbb{P}(\Delta\tilde{\mathcal{Y}} = \emptyset)$$

It follows that the total variation distance between the law of $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ and the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ is less or equal to $2\mathbb{P}(\sharp\Delta\tilde{\mathcal{Y}} \geq 2)$, which according lemma 7.3 is less or equal to $\left(\int_{\mathbb{R}} \tilde{G}(y, y)(\tilde{\kappa}(dy) - \kappa(dy))\right)^2$. \square

Corollary 7.17. — *Let $\kappa_0 \leq \kappa_1 \leq \dots \leq \kappa_j$ be positive Radon measures such that $\int_{\mathbb{R}} |x|(\kappa_j(dx) - \kappa_0(dx)) < +\infty$. Let G_i be the Green's function of $\frac{1}{2}\frac{d^2}{dx^2} - \kappa_i$ and $(\mathcal{Y}_\infty^{(i)}, \mathcal{Z}_\infty^{(i)})$ the pair of interwoven determinantal point processes corresponding to κ_i . Let $((\mathcal{Y}^{(i)}, \mathcal{Z}^{(i)}))_{0 \leq i \leq j}$ be the sequence of pairs of interwoven point processes defined as follows: $(\mathcal{Y}^{(0)}, \mathcal{Z}^{(0)}) := (\mathcal{Y}_\infty^{(0)}, \mathcal{Z}_\infty^{(0)})$; given $(\mathcal{Y}^{(i-1)}, \mathcal{Z}^{(i-1)})$, $(\mathcal{Y}^{(i)}, \mathcal{Z}^{(i)})$ is obtained by applying procedure 7.15 to the arguments $(\mathcal{Y}^{(i-1)}, \mathcal{Z}^{(i-1)})$, κ_{i-1} and κ_i . Then the total variation distance between the law of $(\mathcal{Y}^{(j)}, \mathcal{Z}^{(j)})$ and the law of $(\mathcal{Y}_\infty^{(j)}, \mathcal{Z}_\infty^{(j)})$ is less or equal to*

$$\sum_{i=1}^j \left(\int_{\mathbb{R}} G_{i-1}(y, y)(\kappa_i(dy) - \kappa_{i-1}(dy)) \right)^2$$

Proof. — Let $(\mathcal{Y}'^{(i)}, \mathcal{Z}'^{(i)})$ be the pair of point processes obtained by applying procedure 7.15 to the arguments $(\mathcal{Y}_\infty^{(i-1)}, \mathcal{Z}_\infty^{(i-1)})$, κ_{i-1} and κ_i . According to lemma 7.16, the total variation distance between the law of $(\mathcal{Y}'^{(i)}, \mathcal{Z}'^{(i)})$ and the law of $(\mathcal{Y}_\infty^{(i)}, \mathcal{Z}_\infty^{(i)})$ is less or equal to $\left(\int_{\mathbb{R}} G_{i-1}(y, y)(\kappa_i(dy) - \kappa_{i-1}(dy))\right)^2$. We denote by d_q the total variation distance between the law of $(\mathcal{Y}^{(q)}, \mathcal{Z}^{(q)})$ and the law of $(\mathcal{Y}_\infty^{(q)}, \mathcal{Z}_\infty^{(q)})$. The total variation distance between the law of $(\mathcal{Y}'^{(i)}, \mathcal{Z}'^{(i)})$ and the law of $(\mathcal{Y}^{(i)}, \mathcal{Z}^{(i)})$ is less or equal to d_{i-1} . It follows that

$$d_i \leq d_{i-1} + \left(\int_{\mathbb{R}} G_{i-1}(y, y)(\kappa_i(dy) - \kappa_{i-1}(dy)) \right)^2$$

and thus

$$d_j \leq \sum_{i=1}^j \left(\int_{\mathbb{R}} G_{i-1}(y, y)(\kappa_i(dy) - \kappa_{i-1}(dy)) \right)^2$$

\square

Next we give a true monotone coupling between $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$. We still consider that $\kappa \leq \tilde{\kappa}$ and that $\int_{\mathbb{R}} |x|(\tilde{\kappa}(dx) - \kappa(dx)) < +\infty$. To construct the coupling we will use a continuous monotonic increasing path in the space of measures, $(\kappa_q)_{0 \leq q \leq 1}$, joining κ to $\tilde{\kappa}$ ($\kappa_0 = \kappa$, $\kappa_1 = \tilde{\kappa}$). Such a path is defined as follows: Let Λ be a positive Radon measure on $\mathbb{R} \times [0, 1]$ satisfying the following constraints:

- For any $q \in [0, 1]$, $\Lambda(\mathbb{R} \times \{q\}) = 0$
- For any A Borel subset of \mathbb{R} , $\Lambda(A \times [0, 1]) = \tilde{\kappa}(A)$

For $q \in [0, 1]$, we define κ_q as the measure on \mathbb{R} satisfying, for any A Borel subset of \mathbb{R}

$$\kappa_q(A) = \kappa_0(A) + \Lambda(A \times [0, q])$$

For any $q \leq q' \in [0, 1]$, $\kappa_q \leq \kappa_{q'}$. Moreover the map $q \mapsto \kappa_q$ is continuous for the vague topology. In the sequel we will denote G_q the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - \kappa_q$ (for $x \leq y$, $G_q(x, y) = u_{q, \uparrow}(x)u_{q, \downarrow}(y)$) and use the measure $G_q(y, y)\Lambda(dy, dq)$, which is finite.

Procedure 7.18. — *Arguments:*

- a pair $(\mathcal{Y}, \mathcal{Z})$ of disjoint discrete sets of points in \mathbb{R} such that between any two points in \mathcal{Y} lies a single point in \mathcal{Z} and vice-versa, and such that $\inf \mathcal{Y} \cup \mathcal{Z} \in \mathcal{Y} \cup \{-\infty\}$, $\sup \mathcal{Y} \cup \mathcal{Z} \in \mathcal{Y} \cup \{+\infty\}$
- two positive Radon measures $\kappa, \tilde{\kappa}$ such that $\kappa \leq \tilde{\kappa}$ and $\int_{\mathbb{R}} |x|(\tilde{\kappa}(dx) - \kappa(dx)) < +\infty$ and $(\tilde{\kappa} - \kappa)(\mathcal{Z}) = 0$.
- a continuous monotonic increasing path in the space of measures, $(\kappa_q)_{0 \leq q \leq 1}$, joining κ to $\tilde{\kappa}$, obtained by integrating the Radon measure Λ on $\mathbb{R} \times [0, 1]$.

Procedure:

- (i) First sample a Poisson point process of intensity $G_q(y, y)\Lambda(dy, dq)$ on $\mathbb{R} \times [0, 1]$: $((Y_j, q_j))_{1 \leq j \leq N}$, the points being ordered in the increasing sense of q_j .
- (ii) Then construct recursively the sequence $((\mathcal{Y}^{(j)}, \mathcal{Z}^{(j)}))_{0 \leq j \leq N}$ of pairs of interwoven point processes as follows: $(\mathcal{Y}^{(0)}, \mathcal{Z}^{(0)})$ is set to be $(\mathcal{Y}, \mathcal{Z})$. $(\mathcal{Y}^{(j)}, \mathcal{Z}^{(j)})$ is obtained by applying procedure 7.10 to the arguments $(\mathcal{Y}^{(j-1)}, \mathcal{Z}^{(j-1)})$, κ_{q_j} and Y_j .
- (iii) $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ is set to be $(\mathcal{Y}^{(N)}, \mathcal{Z}^{(N)})$

Return: $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$.

The condition $(\tilde{\kappa} - \kappa)(\mathcal{Z}) = 0$ ensures that a.s., none of $\mathcal{Y}^{(j)}$ lies in \mathcal{Z} . By construction $\mathcal{Z} \subseteq \tilde{\mathcal{Z}}$ and $\tilde{\mathcal{Y}} \subseteq \mathcal{Y} \cup \text{Supp}(\tilde{\kappa} - \kappa)$. $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ differs from $(\mathcal{Y}, \mathcal{Z})$ only by a finite number of points. The law of $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ depends only on the "geometrical path" $(\kappa_q)_{0 \leq q \leq 1}$ and not on its parametrization: if θ is an increasing homomorphism from $[0, 1]$ to itself, then procedure 7.18 applied the path $(\kappa_{\theta(q)})_{0 \leq q \leq 1}$ returns the same result (in law). Below an illustration of procedure 7.18:




Fig. 4 - Illustration of procedure 7.18: On the left are represented $(\mathcal{Y}, \mathcal{Z})$ and the Poisson process $((Y_j, q_j))_{1 \leq j \leq N}$. On the right are represented the successive $((\mathcal{Y}^{(j)}, \mathcal{Z}^{(j)}))_{0 \leq j \leq N}$. x-dots represent the points of $\mathcal{Y}^{(j)}$ and diamonds the points of $\mathcal{Z}^{(j)}$.

Proposition 7.19. — *Let $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ respectively $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ be the couple of interwoven determinantal point processes corresponding to the killing measure κ respectively $\tilde{\kappa}$. We assume that the procedure 7.18 is applied to $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and that $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ is the returned couple of point processes. Then $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ has the same law as $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$.*

Proof. — Observe that a.s. $(\tilde{\kappa} - \kappa)(\mathcal{Z}_\infty) = 0$. Let $n \in \mathbb{N}^*$. We define the family $((\mathcal{Y}^{(j,n)}, \mathcal{Z}^{(j,n)}))_{0 \leq j \leq n}$ of interwoven point processes as follows: $(\mathcal{Y}^{(0,n)}, \mathcal{Z}^{(0,n)})$ equals $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$. Given $(\mathcal{Y}^{(j-1,n)}, \mathcal{Z}^{(j-1,n)})$, $(\mathcal{Y}^{(j,n)}, \mathcal{Z}^{(j,n)})$ is obtained by applying procedure 7.15 to the arguments $(\mathcal{Y}^{(j-1,n)}, \mathcal{Z}^{(j-1,n)})$, $\kappa_{\frac{j-1}{n}}$ and $\kappa_{\frac{j}{n}}$. We will show that as n tends to infinity, the law of $(\mathcal{Y}^{(n,n)}, \mathcal{Z}^{(n,n)})$ converges in total variation to the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ and converges weakly to the law of $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$, which will imply that $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$ and $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ have the same law.

Applying corollary 7.17, we get that the total variation distance between the law of $(\mathcal{Y}^{(n,n)}, \mathcal{Z}^{(n,n)})$ and the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ is bounded by

$$\begin{aligned} & \sum_{j=1}^n \left(\int_{\mathbb{R}} G_{\frac{j-1}{n}}(y, y) (\kappa_{\frac{j}{n}}(dy) - \kappa_{\frac{j-1}{n}}(dy)) \right)^2 \\ & \leq \sup_{x \in \mathbb{R}} \left(\frac{G_0(x, x)}{1 + |x|} \right)^2 \sum_{j=1}^n \left(\int_{\mathbb{R}} (1 + |y|) (\kappa_{\frac{j}{n}}(dy) - \kappa_{\frac{j-1}{n}}(dy)) \right)^2 \\ & \leq \sup_{x \in \mathbb{R}} \left(\frac{G_0(x, x)}{1 + |x|} \right)^2 \int_{\mathbb{R}} (1 + |y|) (\tilde{\kappa}(dy) - \kappa(dy)) \\ & \quad \times \sup_{1 \leq j \leq n} \int_{\mathbb{R}} (1 + |y|) (\kappa_{\frac{j}{n}}(dy) - \kappa_{\frac{j-1}{n}}(dy)) \end{aligned}$$

The continuity of the path $(\kappa_q)_{0 \leq q \leq 1}$ ensures that

$$\lim_{n \rightarrow +\infty} \sup_{1 \leq j \leq n} \int_{\mathbb{R}} (1 + |y|) (\kappa_{\frac{j}{n}}(dy) - \kappa_{\frac{j-1}{n}}(dy)) = 0$$

and hence the total variation distance between the law of $(\mathcal{Y}^{(n,n)}, \mathcal{Z}^{(n,n)})$ and the law of $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ converges to 0 as n tends to infinity.

We define a random finite set E_n of points in $\mathbb{R} \times \{\frac{1}{n}, \frac{2}{n}, \dots, \frac{n}{n}\}$ as follows: Let $(\beta_{1,n}, \beta_{2,n}, \dots, \beta_{n,n})$ be a family of independent Bernoulli variables, $\beta_{i,n}$ being of parameter

$$\int_{\mathbb{R}} v_{\kappa_{\frac{i-1}{n}}, \kappa_{\frac{i}{n}}}(y) (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy)$$

Whenever $\beta_{i,n} = 1$, we add to E_n a point $(Y_{i,n}, \frac{i-1}{n})$ to E_n where $Y_{i,n}$ is a r.v. distributed according the measure

$$\frac{1}{\mathbb{P}(\beta_{i,n} = 1)} v_{\kappa_{\frac{i-1}{n}}, \kappa_{\frac{i}{n}}}(y) (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy)$$

The $(Y_{i,n}, \frac{i-1}{n})$ are assumed to be independent and independent from the family $(\beta_{1,n}, \beta_{2,n}, \dots, \beta_{n,n})$. The pair $(\mathcal{Y}^{(n,n)}, \mathcal{Z}^{(n,n)})$ is sampled as follows: starting from $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$, independent from E_n , we apply successively, for i ranging from 1 to n , the procedure 7.10 with the arguments $\kappa_{\frac{i-1}{n}}$ and $Y_{i,n}$ whenever $\beta_{i,n} = 1$. At the end we get $(\mathcal{Y}^{(n,n)}, \mathcal{Z}^{(n,n)})$. According to lemma 2.4, the law of the pair of point processes returned by procedure 7.10 depends continuously on the arguments. So to prove that $(\mathcal{Y}^{(n,n)}, \mathcal{Z}^{(n,n)})$ converges in law to $(\tilde{\mathcal{Y}}, \tilde{\mathcal{Z}})$, we only need to show that the random set of point E_n converges in law to the Poisson point process $((Y_j, q_j))_{1 \leq j \leq N}$ used in

procedure 7.18. All of the functions $v_{\kappa_{\frac{i-1}{n}}, \kappa_{\frac{i}{n}}}(y)$ are dominated by $G_0(y, y)$. Moreover

$$\begin{aligned}
& |v_{\kappa_{\frac{i-1}{n}}, \kappa_{\frac{i}{n}}}(y) - G_{\frac{i}{n}}(y, y)| \\
& \leq u_{\frac{i}{n}, \downarrow}(y) \int_{y_{-1} < y} u_{\frac{i}{n}, \uparrow}(y_{-1}) (u_{\frac{i-1}{n}, \downarrow}(y_{-1}) u_{\frac{i-1}{n}, \uparrow}(y) - u_{\frac{i-1}{n}, \uparrow}(y_{-1}) u_{\frac{i-1}{n}, \downarrow}(y)) \\
& \quad \times (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy_{-1}) \\
& + u_{\frac{i}{n}, \uparrow}(y) \int_{y_1 > y} u_{\frac{i}{n}, \downarrow}(y_1) (u_{\frac{i-1}{n}, \uparrow}(y_1) u_{\frac{i-1}{n}, \downarrow}(y) - u_{\frac{i-1}{n}, \downarrow}(y_1) u_{\frac{i-1}{n}, \uparrow}(y)) \\
& \quad \times (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy_1) \\
& + \int_{y_{-1} < y} u_{\frac{i}{n}, \uparrow}(y_{-1}) (u_{\frac{i-1}{n}, \downarrow}(y_{-1}) u_{\frac{i-1}{n}, \uparrow}(y) - u_{\frac{i-1}{n}, \uparrow}(y_{-1}) u_{\frac{i-1}{n}, \downarrow}(y)) \\
& \quad \times (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy_{-1}) \\
& \times \int_{y_1 > y} u_{\frac{i}{n}, \downarrow}(y_1) (u_{\frac{i-1}{n}, \uparrow}(y_1) u_{\frac{i-1}{n}, \downarrow}(y) - u_{\frac{i-1}{n}, \downarrow}(y_1) u_{\frac{i-1}{n}, \uparrow}(y)) \\
& \quad \times (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy_1) \\
& \leq G_0(y, y) \int_{y_{-1} < y} G_0(y_{-1}, y_{-1}) (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy_{-1}) \\
& + G_0(y, y) \int_{y_1 > y} G_0(y_1, y_1) (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy_1) \\
& + G_0(y, y) \int_{y_{-1} < y} G_0(y_{-1}, y_{-1}) (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy_{-1}) \\
& \quad \times \int_{y_1 > y} G_0(y_1, y_1) (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy_1)
\end{aligned}$$

Thus given any bounded interval J

$$\lim_{n \rightarrow +\infty} \sup_{1 \leq i \leq n} \sup_{y \in J} |v_{\kappa_{\frac{i-1}{n}}, \kappa_{\frac{i}{n}}}(y) - G_{\frac{i}{n}}(y, y)| = 0$$

It follows that

$$\lim_{n \rightarrow +\infty} \sup_{1 \leq i \leq n} \mathbb{P}(\beta_{i,n} = 1) = 0$$

and the measure

$$\sum_{i=1}^n v_{\kappa_{\frac{i-1}{n}}, \kappa_{\frac{i}{n}}}(y) (\kappa_{\frac{i}{n}} - \kappa_{\frac{i-1}{n}})(dy) \otimes \delta_{\frac{i}{n}}(dq)$$

converges weakly to $G_q(y, y)\Lambda(dy, dq)$, which is the intensity of the Poisson point process $((Y_j, q_j))_{1 \leq j \leq N}$. Thus the random sets E_n are compound Bernoulli approximations of the Poisson point process $((Y_j, q_j))_{1 \leq j \leq N}$ and converge in law to the latter. \square

Given a continuous monotonic increasing path $(\kappa_q)_{0 \leq q \leq 1}$ in the space of Radon measures and a pair of interwoven determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ corresponding to κ_0 , used as argument, procedure 7.18 yields non-homogeneous Markov q -parametrized process in the space of interwoven pairs of discrete sets of points whose one-dimensional marginal at any value q_0 of the parameter is the pair of interwoven determinantal point processes corresponding to the killing measure κ_{q_0} . This corresponds to sampling only the partial Poisson point process of intensity $\mathbb{1}_{0 \leq q \leq q_0} G_q(y, y) \Lambda(dy, dq)$ and successively applying procedure 7.10 for each of its points. In general, multidimensional marginals corresponding to $q_1 < \dots < q_n$ depend not only on $\kappa_{q_1}, \dots, \kappa_{q_n}$ but on the whole path $(\kappa_q)_{q_1 \leq q \leq q_n}$. For instance consider two different paths $(\kappa_q)_{0 \leq q \leq 1}$ and $(\hat{\kappa}_q)_{0 \leq q \leq 1}$ where

$$\begin{aligned} - \kappa_0 &= \hat{\kappa}_0 = \delta_{-\frac{1}{2}} + \delta_{\frac{1}{2}} \\ - \kappa_1 &= \hat{\kappa}_1 = \delta_{-\frac{3}{2}} + \delta_{-\frac{1}{2}} + \delta_{\frac{1}{2}} + \delta_{\frac{3}{2}} \\ - \kappa_q &= 2q\delta_{-\frac{1}{2}} + \delta_{-\frac{1}{2}} + \delta_{\frac{1}{2}} \text{ for } q \in [0, \frac{1}{2}] \text{ and } \kappa_q = \delta_{-\frac{1}{2}} + \delta_{-\frac{1}{2}} + \delta_{\frac{1}{2}} + (2q-1)\delta_{\frac{3}{2}} \\ &\quad \text{for } q \in [\frac{1}{2}, 1] \\ - \hat{\kappa}_q &= \delta_{-\frac{1}{2}} + \delta_{\frac{1}{2}} + 2q\delta_{\frac{1}{2}} \text{ for } q \in [0, \frac{1}{2}] \text{ and } \hat{\kappa}_q = (2q-1)\delta_{-\frac{1}{2}} + \delta_{-\frac{1}{2}} + \delta_{\frac{1}{2}} + \delta_{\frac{3}{2}} \\ &\quad \text{for } q \in [\frac{1}{2}, 1] \end{aligned}$$

Let $G_q(x, y) = u_{q,\uparrow}(x \wedge y) u_{q,\downarrow}(x \vee y)$ be the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - \kappa_q$ and $\hat{G}_q(x, y) = \hat{u}_{q,\uparrow}(x \wedge y) \hat{u}_{q,\downarrow}(x \vee y)$ the Green's function of $\frac{1}{2} \frac{d^2}{dx^2} - \hat{\kappa}_q$. Let $((\mathcal{Y}_\infty, \mathcal{Z}_\infty), (\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty))$ be the coupling between the point process corresponding to κ_0 respectively κ_1 induced by the path $(\kappa_q)_{0 \leq q \leq 1}$ and $((\mathcal{Y}_\infty, \mathcal{Z}_\infty), (\hat{\mathcal{Y}}_\infty, \hat{\mathcal{Z}}_\infty))$ the coupling induced by the path $(\hat{\kappa}_q)_{0 \leq q \leq 1}$. Then

$$\begin{aligned} \mathbb{P}(\mathcal{Y}_\infty = \left\{ -\frac{1}{2}, \frac{1}{2} \right\}, \tilde{\mathcal{Y}}_\infty = \left\{ -\frac{3}{2}, \frac{1}{2}, \frac{3}{2} \right\}) \\ = \mathbb{P}(\mathcal{Y}_\infty = \left\{ -\frac{1}{2}, \frac{1}{2} \right\}) \times G_{\frac{1}{2}}\left(-\frac{3}{2}\right) \frac{u_{\frac{1}{2},\downarrow}\left(-\frac{1}{2}\right)}{u_{\frac{1}{2},\downarrow}\left(-\frac{3}{2}\right)} G_1\left(\frac{3}{2}\right) \left(1 - \frac{u_{1,\uparrow}\left(\frac{1}{2}\right)}{u_{1,\uparrow}\left(\frac{3}{2}\right)}\right) \end{aligned}$$

$$\begin{aligned} \mathbb{P}(\mathcal{Y}_\infty = \left\{ -\frac{1}{2}, \frac{1}{2} \right\}, \hat{\mathcal{Y}}_\infty = \left\{ -\frac{3}{2}, \frac{1}{2}, \frac{3}{2} \right\}) \\ = \mathbb{P}(\mathcal{Y}_\infty = \left\{ -\frac{1}{2}, \frac{1}{2} \right\}) \times \hat{G}_{\frac{1}{2}}\left(\frac{3}{2}\right) \left(1 - \frac{\hat{u}_{\frac{1}{2},\uparrow}\left(\frac{1}{2}\right)}{\hat{u}_{\frac{1}{2},\uparrow}\left(\frac{3}{2}\right)}\right) \hat{G}_1\left(-\frac{3}{2}\right) \frac{\hat{u}_{1,\downarrow}\left(-\frac{1}{2}\right)}{\hat{u}_{1,\downarrow}\left(-\frac{3}{2}\right)} \end{aligned}$$

But

$$\hat{G}_{\frac{1}{2}}\left(\frac{3}{2}\right) = G_{\frac{1}{2}}\left(-\frac{3}{2}\right) \quad \hat{G}_1\left(-\frac{3}{2}\right) = G_1\left(\frac{3}{2}\right)$$

and

$$\frac{\hat{u}_{\frac{1}{2},\uparrow}\left(\frac{1}{2}\right)}{\hat{u}_{\frac{1}{2},\uparrow}\left(\frac{3}{2}\right)} = \frac{u_{\frac{1}{2},\downarrow}\left(-\frac{1}{2}\right)}{u_{\frac{1}{2},\downarrow}\left(-\frac{3}{2}\right)} \quad \frac{\hat{u}_{1,\downarrow}\left(-\frac{1}{2}\right)}{\hat{u}_{1,\downarrow}\left(-\frac{3}{2}\right)} = \frac{u_{1,\uparrow}\left(\frac{1}{2}\right)}{u_{1,\uparrow}\left(\frac{3}{2}\right)}$$

Thus

$$\frac{\mathbb{P}\left(\mathcal{Y}_\infty = \left\{-\frac{1}{2}, \frac{1}{2}\right\}, \tilde{\mathcal{Y}}_\infty = \left\{-\frac{3}{2}, \frac{1}{2}, \frac{3}{2}\right\}\right)}{\mathbb{P}\left(\mathcal{Y}_\infty = \left\{-\frac{1}{2}, \frac{1}{2}\right\}, \hat{\mathcal{Y}}_\infty = \left\{-\frac{3}{2}, \frac{1}{2}, \frac{3}{2}\right\}\right)} = \frac{\frac{u_{\frac{1}{2}, \downarrow}\left(-\frac{1}{2}\right)}{u_{\frac{1}{2}, \downarrow}\left(-\frac{3}{2}\right)}}{1 - \frac{u_{\frac{1}{2}, \downarrow}\left(-\frac{1}{2}\right)}{u_{\frac{1}{2}, \downarrow}\left(-\frac{3}{2}\right)}} \times \frac{1 - \frac{u_{1, \uparrow}\left(\frac{1}{2}\right)}{u_{1, \uparrow}\left(\frac{3}{2}\right)}}{\frac{u_{1, \uparrow}\left(\frac{1}{2}\right)}{u_{1, \uparrow}\left(\frac{3}{2}\right)}}$$

But

$$\frac{u_{\frac{1}{2}, \downarrow}\left(-\frac{1}{2}\right)}{u_{\frac{1}{2}, \downarrow}\left(-\frac{3}{2}\right)} = \frac{3}{11} \frac{u_{1, \uparrow}\left(\frac{1}{2}\right)}{u_{1, \uparrow}\left(\frac{3}{2}\right)} = \frac{11}{41}$$

Thus

$$\frac{\mathbb{P}\left(\mathcal{Y}_\infty = \left\{-\frac{1}{2}, \frac{1}{2}\right\}, \tilde{\mathcal{Y}}_\infty = \left\{-\frac{3}{2}, \frac{1}{2}, \frac{3}{2}\right\}\right)}{\mathbb{P}\left(\mathcal{Y}_\infty = \left\{-\frac{1}{2}, \frac{1}{2}\right\}, \hat{\mathcal{Y}}_\infty = \left\{-\frac{3}{2}, \frac{1}{2}, \frac{3}{2}\right\}\right)} = \frac{45}{44} \neq 1$$

The two couplings are different.

If $\tilde{\kappa} - \kappa$ does not have a first moment we can still construct a coupling between $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ as follows: Consider a continuous monotonic increasing path $(\kappa_q)_{0 \leq q \leq 1}$ joining κ to $\tilde{\kappa}$ satisfying the constraint

$$\forall q \in [0, 1), \int_{\mathbb{R}} |x|(\kappa_q(dx) - \kappa_0(dx)) < +\infty$$

Given $q_0 \in (0, 1)$, one can apply procedure 7.18 to the arguments $(\mathcal{Y}_\infty, \mathcal{Z}_\infty), \kappa, \kappa_{q_0}$ and the partial path $(\kappa_q)_{0 \leq q \leq q_0}$. As result we get a two interwoven determinantal point processes corresponding to the killing measure κ_{q_0} . At the limit as q_0 tends to 1 we get something that has the same law as $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$.

Next we prove the existence of stronger couplings in case $\tilde{\kappa} = c\kappa$ where $c > 1$ is a constant.

Proposition 7.20. — *If $\tilde{\kappa} = c\kappa$ with $c > 1$ then there is a coupling between $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ such that $\mathcal{Z}_\infty \subseteq \tilde{\mathcal{Z}}_\infty$ and $\mathcal{Y}_\infty \subseteq \tilde{\mathcal{Y}}_\infty$.*

Proof. — Consider a coupling between $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$ and $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$ given by procedure 7.18, possible extended to the case where κ does not have a first moment. Then $\mathcal{Z}_\infty \subseteq \tilde{\mathcal{Z}}_\infty$ but in general $\mathcal{Y}_\infty \not\subseteq \tilde{\mathcal{Y}}_\infty$. So we will sample other point processes \mathcal{Y}'_∞ and $\tilde{\mathcal{Y}}'_\infty$ that conditionally on \mathcal{Z}_∞ respectively $\tilde{\mathcal{Z}}_\infty$ have the same law as \mathcal{Y}_∞ respectively $\tilde{\mathcal{Y}}_\infty$, and such that $\mathcal{Y}'_\infty \subseteq \tilde{\mathcal{Y}}'_\infty$. For each connected component \tilde{J} of $\mathbb{R} \setminus \tilde{\mathcal{Z}}_\infty$ we sample a point $\tilde{Y}_{\tilde{J}}$ according the measure $\frac{1_{y \in \tilde{J}} \tilde{\kappa}(dy)}{\tilde{\kappa}(\tilde{J})}$. We assume that conditionally on $\tilde{\mathcal{Z}}_\infty$, all the $\tilde{Y}_{\tilde{J}}$ are independent from \mathcal{Z}_∞ and independent one from another. We set

$$\tilde{\mathcal{Y}}'_\infty := \{\tilde{Y}_{\tilde{J}} | \tilde{J} \text{ connected component of } \mathbb{R} \setminus \tilde{\mathcal{Z}}_\infty\}$$

Then $(\tilde{\mathcal{Y}}'_\infty, \tilde{\mathcal{Z}}_\infty)$ has the same law as $(\tilde{\mathcal{Y}}_\infty, \tilde{\mathcal{Z}}_\infty)$. Let be J a connected component of $\mathbb{R} \setminus \mathcal{Z}_\infty$ and $\tilde{J}_1, \dots, \tilde{J}_{N_J}$ the connected components of $J \setminus \tilde{\mathcal{Z}}_\infty$. On J we define the

r.v. Y_J as follows: Y_J takes value in $\tilde{Y}_{\tilde{J}_n}$ and

$$\mathbb{P}(Y_J = \tilde{Y}_{\tilde{J}_n} | J, \tilde{J}_1, \dots, \tilde{J}_{N_J}) = \frac{\kappa(\tilde{J}_n)}{\kappa(J)}$$

We set

$$\mathcal{Y}'_\infty := \{Y_J | J \text{ connected component of } \mathbb{R} \setminus \mathcal{Z}_\infty\}$$

By construction $\mathcal{Y}'_\infty \subseteq \tilde{\mathcal{Y}}'_\infty$. Moreover the proportionality of κ and $\tilde{\kappa}$ ensures that $(\mathcal{Y}'_\infty, \mathcal{Z}_\infty)$ has the same law as $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$. \square

BIBLIOGRAPHY

- [1] I. BENJAMINI, R. LYONS, Y. PERES & O. SCHRAMM – “Unifrom spanning forests”, *The Annals of Probability* **29** (2001), no. 1, p. 1–65.
- [2] J. BERTOIN & J. PITMAN – “Two coalescents derived from the ranges of stable subordinators”, *Electronic Journal of Probability* **5** (1999), no. 7.
- [3] P. BIANE – “Relations entre pont et excursion du mouvement brownien réel.”, *Annales de l’institut Henri Poincaré* **22** (1986), no. 1, p. 1–7.
- [4] G. BIRKHOFF & G. C. ROTA – *Ordinary differential equations*, 4th ed., John Wiley and Sons, 1989.
- [5] L. BREIMAN – *Probability*, Classics in applied mathematics, vol. 7, SIAM, 1992.
- [6] L. CHAUMONT & G. U. BRAVO – “Markovian bridges : weak continuity and pathwise construction”, *The Annals of Probability* **39** (2011), no. 2, p. 609–647.
- [7] E. B. DYNKIN – “Gaussian and non-gaussian random fields associated with markov processes”, *Journal of Functional Analysis* **55** (1984), p. 344–376.
- [8] ———, “Local times and quantum fields”, in *Seminar on Stochastic Processes, Gainesville 1983*, Progress in Probability and Statistics, vol. 7, Birkhauser, 1984, p. 69–84.
- [9] ———, “Polynomials of the occupation field and related random fields”, *Journal of Functional Analysis* **58** (1984), p. 20–52.
- [10] P. FITZSIMMONS & J. ROSEN – “Markovian loop soups: permanental processes and isomorphism theorems”, arXiv:1211.5163, Nov. 2012.
- [11] J. B. HOUGH, M. KRISHNAPUR, Y. PERES & B. VIRAG – *Zeros of gaussian analytic functions and determinantal point processes*, University Lecture Series, vol. 51, American Mathematical Society, 2009.

- [12] K. ITÔ & H. P. MCKEAN – *Diffusion processes and their sample paths*, Grundlehren der mathematischen Wissenschaften, vol. 125, Springer, 1974.
- [13] Y. L. JAN – *Markov paths, loops and fields*, Lecture Notes in Mathematics, École d’Été de Probabilités de Saint-Flour, vol. 2026, Springer, 2011.
- [14] Y. L. JAN & S. LEMAIRE – “Markovian loop clusters on graphs”, To be published in *Illinois Journal of Mathematics*.
- [15] Y. L. JAN, M. MARCUS & J. ROSEN – “Permanental fields, loop soups and continuous additive functionals”, arXiv:1209.1804, Sept. 2012.
- [16] I. KARATZAS & S. E. SHREVE – *Brownian motion and stochastic calculus*, 2nd ed., Graduate Texts in Mathematics, vol. 113, Springer, 2010.
- [17] K. KAWAZU & S. WATANABE – “Branching processes with immigration and related limit theorems”, *Theory of Probability and its Applications* **16** (1971), no. 1, p. 36–54.
- [18] G. F. LAWLER, O. SCHRAMM & W. WERNER – “Conformal restriction: the chordal case”, *Journal of American Mathematical Society* **16** (2003), no. 4, p. 917–955.
- [19] G. F. LAWLER & W. WERNER – “The brownian loop-soup”, *Probability Theory and Related Fields* **128** (2004), p. 565–588.
- [20] R. LYONS – “Determinantal probability measures”, *Publications Mathématiques de l’IHÉS* **98** (2003), p. 167–212.
- [21] H. P. MCKEAN – “Elementary solutions for certain parabolic partial differential equations”, *Transactions of the American Mathematical Society* **82** (1956), no. 2, p. 519–548.
- [22] J. P. P. FITZSIMMONS & M. YOR – “Markovian bridges: Construction, palm interpretation, and splicing”, in *Seminar on Stochastic Processes 1992* (Boston), Birkhäuser, 1993, p. 101–134.
- [23] J. PITMAN & M. YOR – “Decomposition at the maximum for excursions and bridges of one-dimensional diffusions”, in *Ito’s Stochastic Calculus and Probability Theory*, Springer, 1996, p. 293–310.
- [24] D. REVUZ & M. YOR – *Continuous martingales and brownian motion*, 3rd ed., Grundlehren der mathematischen Wissenschaften, vol. 293, Springer, 1999.
- [25] P. SALMINEN, P. VALLOIS & M. YOR – “On the excursion theory for linear diffusions”, *Japanese Journal of Mathematics* **2** (2007), no. 1, p. 97–127.
- [26] S. SHEFFIELD & W. WERNER – “Conformal loop ensembles: the markovian characterization and the loop-soup construction”, *Annals of Mathematics* **176** (2012), no. 3, p. 1827–1917.

- [27] B. SIMON – *Trace ideals and their applications*, 2nd ed., Mathematical Surveys and Monographs, vol. 120, American Mathematical Society, 2005.
- [28] A. SOSHNIKOV – “Determinantal random point fields”, *Uspekhi Matematicheskikh Nauk* **55** (2000), no. 5, p. 107–160.
- [29] G. TESCHL – *Ordinary differential equations and dynamical systems*, Graduate Studies in Mathematics, vol. 140, American Mathematical Society, 2012.
- [30] W. VERVAAT – “A relation between brownian bridge and brownian excursion”, *The Annals of Probability* **7** (1979), no. 1, p. 143–149.
- [31] D. B. WILSON – “Generating random spanning trees more quickly than the cover time”, in *Proceedings of the Twenty-Eighth Annual ACM Symposium on the Theory of Computing*, Association for Computing Machinery, 1996, p. 296–303.
- [32] D. ZHAN – “Loop-erasure of planar brownian motion”, *Communications in Mathematical Physics* **303** (2012), no. 3, p. 709–720.