



HAL
open science

Every totally real algebraic integer is a tree eigenvalue

Justin Salez

► **To cite this version:**

| Justin Salez. Every totally real algebraic integer is a tree eigenvalue. 2013. hal-00789806v1

HAL Id: hal-00789806

<https://hal.science/hal-00789806v1>

Preprint submitted on 18 Feb 2013 (v1), last revised 4 Sep 2014 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Every totally real algebraic integer is a tree eigenvalue

Justin Salez

Université Paris Diderot & LPMA

Abstract

Every graph eigenvalue is in particular a totally real algebraic integer, i.e. a zero of some real-rooted monic polynomial with integer coefficients. Conversely, the fact that every such number occurs as an eigenvalue of some finite graph is a remarkable result, conjectured forty years ago by Hoffman, and proved twenty years later by Bass, Estes and Guralnick. This note provides an independent, elementary proof of a stronger statement, namely that the graph may always be chosen to be a tree.

Keywords: adjacency matrix, tree, eigenvalue, totally real algebraic integer
2010 MSC: 05C05, 05C25, 05C31, 05C50

1. Introduction

By definition, the eigenvalues of a finite undirected graph $G = (V, E)$ are the roots of the characteristic polynomial $\Phi_G(X) := \det(A - XI)$, where $A = (A_{ij})_{i,j \in V}$ is the adjacency matrix of G :

$$A_{ij} = \begin{cases} 1 & \text{if } \{i, j\} \in E; \\ 0 & \text{otherwise.} \end{cases} .$$

Graph eigenvalues capture a considerable amount of structural information, and their study constitutes the very essence of spectral graph theory [1, 2].

It follows from their definition that graph eigenvalues belong to the ring of **totally real algebraic integers**, i.e. zeros of real-rooted monic polynomials with integer coefficients. Conversely, the remarkable fact that every totally real algebraic integer is an eigenvalue of some finite graph is a remarkable result, conjectured forty years ago by Hoffman [3], and established twenty

URL: <http://www.proba.jussieu.fr/~salez/> (Justin Salez)

years later by Bass, Estes and Guralnick [4]. The purpose of this note is to give an elementary and self-contained proof of a stronger statement, namely that the graph may be chosen to be a tree.

Theorem 1. *Every totally real algebraic integer occurs as the eigenvalue of some finite tree.*

Given two trees S and T with respective eigenvalues $\lambda_1, \dots, \lambda_m$ and μ_1, \dots, μ_n (counted with multiplicities), it is easy to construct a new tree whose eigenvalues include $\lambda_1, \dots, \lambda_m, \mu_1, \dots, \mu_n$ (with multiplicities) : simply root S and T at arbitrary vertices, then duplicate each rooted tree, and connect the four resulting roots to a common additional vertex. Consequently, we obtain the following Corollary.

Corollary 1. *Any finite multi-set of totally real algebraic integers is contained in the spectrum of some finite tree.*

Trees undoubtedly play a special and important role in many aspects of graph theory. We therefore believe that the strengthening provided by Theorem 1 may be of independent interest, beyond the fact that it provides a considerably simpler proof of Hoffman's conjecture.

2. Reformulation of the problem and outline of the proof

Let $T = (V, E)$ be a finite tree, and $o \in V$ an arbitrary vertex. Our starting point is the following well-known and elementary recursion for the characteristic polynomial of T (see e.g. [2, Proposition 5.1.1]) :

$$-\Phi_T(X) = X\Phi_{T \setminus o}(X) + \sum_{i \sim o} \Phi_{T \setminus i \setminus o}(X). \quad (1)$$

Note that $T \setminus o$ consists of $\deg(o)$ disjoint subtrees T_i , one for each neighbor $i \sim o$, and hence that $\Phi_{T \setminus o}(X) = \prod_{i \sim o} \Phi_{T_i}(X)$. Dividing (1) by $X\Phi_{T \setminus o}(X)$, one obtains that the rational function

$$\mathfrak{f}_{(T,o)}(X) := 1 + \frac{\Phi_T(X)}{X\Phi_{T \setminus o}(X)}$$

satisfies the following recursion :

$$\mathfrak{f}_{(T,o)}(X) = \frac{1}{X^2} \sum_{i \sim o} \frac{1}{1 - \mathfrak{f}_{(T_i,i)}(X)},$$

the sum being interpreted as 0 when empty (i.e. when T is reduced to o). It follows that a number $\lambda \in \mathbb{C} \setminus \{0\}$ is a tree eigenvalue if and only if $1 \in \mathcal{F}(\lambda)$, where $\mathcal{F} = \mathcal{F}(\lambda)$ is the smallest subset of \mathbb{C} satisfying for every $d \in \mathbb{N}$

$$(\alpha_1, \dots, \alpha_d) \in \mathcal{F} \implies \frac{1}{\lambda^2} \sum_{i=1}^d \frac{1}{1 - \alpha_i} \in \mathcal{F}.$$

Equivalently, $\mathcal{F} = \mathcal{F}(\lambda)$ is the smallest subset of \mathbb{C} satisfying :

$$\begin{aligned} \text{(ZERO)} \quad & 0 \in \mathcal{F}; \\ \text{(MAP)} \quad & \alpha \in \mathcal{F} \implies \frac{1}{\lambda^2(1 - \alpha)} \in \mathcal{F}; \\ \text{(ADD)} \quad & \alpha, \beta \in \mathcal{F} \implies \alpha + \beta \in \mathcal{F}. \end{aligned}$$

Remark 1. *Strictly speaking, one should work in the extended complex plane $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ or restrict the property (MAP) to the case $\alpha \neq 1$ to avoid divisions by zero. We choose to ignore this issue, as it would unnecessarily lengthen the proof. Our goal being precisely to show that $1 \in \mathcal{F}$, the skeptical reader may simply assume that $1 \notin \mathcal{F}$ throughout the paper, and obtain a contradiction in the end.*

For example, \mathcal{F} contains $\frac{1}{\lambda^2 - k}$ for every $k \in \mathbb{N}$, as proved by the following chain of implications :

$$0 \in \mathcal{F} \xrightarrow{\text{(MAP)}} \frac{1}{\lambda^2} \in \mathcal{F} \xrightarrow{\text{(ADD)}} \frac{k}{\lambda^2} \in \mathcal{F} \xrightarrow{\text{(MAP)}} \frac{1}{\lambda^2 - k} \in \mathcal{F}. \quad (2)$$

We will in fact determine \mathcal{F} explicitly, for an arbitrary totally real algebraic number $\lambda \neq 0$. Specifically, we will successively prove the following facts.

- I. If λ is a totally real algebraic number then \mathcal{F} contains a positive integer.
- II. If \mathcal{F} contains a positive integer, then \mathcal{F} is stable under negation.
- III. If \mathcal{F} is stable under negation, then

$$\mathcal{F} = \left\{ \frac{P(\lambda^2)}{Q(\lambda^2)} : P, Q \in \mathbb{Z}[X], Q \text{ monic, } \deg(P) < \deg(Q) \right\}.$$

In particular, if λ is a totally real algebraic integer, then \mathcal{F} is the field $\mathbb{Q}(\lambda^2)$ which contains 1, and Theorem 1 is proved. Facts I, II and III will be established respectively in Section 4, 5 and 6. In Section 3 below, we recall some basic properties of algebraic numbers that will be useful in the sequel.

3. Algebraic preliminaries

A number $\zeta \in \mathbb{C}$ is *algebraic* if it is a zero of some polynomial with rational coefficients. All such polynomials are then multiple of a unique monic polynomial $P \in \mathbb{Q}[X]$, called the *minimal polynomial* of ζ . The *degree* of ζ is the degree of P . The algebraic number ζ is called

- *totally real* if all the roots of P are real ;
- *totally positive* if all the roots of P are non-negative.

The following Lemma gathers the basic properties of algebraic numbers that will be used in the sequel. These are well-known (see e.g. [5]) and follow directly from the fact that if $P(X) = \prod_{i=1}^n (X - \alpha_i)$ and $Q(X) = \prod_{j=1}^m (X - \beta_j)$ have rational coefficients, then so do the polynomials

$$\begin{aligned} \prod_{i=1}^n \prod_{j=1}^m (X - \alpha_i - \beta_j), & \quad \prod_{i=1}^n \left(X - \frac{1}{\alpha_i} \right), & \quad \prod_{i=1}^n \prod_{j=1}^m (X - \alpha_i \beta_j) \\ \prod_{i=1}^n (X + \alpha_i), & \quad \prod_{i=1}^n (X^2 - \alpha_i^2), & \quad \prod_{i=1}^n (X^2 - \alpha_i). \end{aligned}$$

Lemma 1 (Elementary algebraic properties).

- The totally real algebraic numbers form a subfield of \mathbb{C} .
- The set of totally positive algebraic numbers is stable under $+$, \times , \div .
- If α is totally real and if $\beta \neq 0$ is totally positive, then $\alpha + k\beta$ is totally positive for all sufficiently large $k \in \mathbb{N}$.
- The squares of totally real algebraic numbers are the totally positive algebraic numbers.

We shall also use twice the following elementary result.

Lemma 2. Let ζ be an algebraic number of degree n , and let $q_1 > \dots > q_n$ be rational numbers. Then, there exists $m_1, \dots, m_n \in \mathbb{Z}$ such that

$$\frac{m_1}{\zeta - q_1} + \dots + \frac{m_n}{\zeta - q_n} \in \mathbb{N}^*. \quad (3)$$

Moreover, for every $1 \leq i \leq n$, m_i has the same sign as $(-1)^i P(q_i)$, where $P \in \mathbb{Q}[X]$ is the minimal polynomial of ζ .

Proof. Since $q_1 > \dots > q_n$ are pairwise distinct, the n polynomials

$$Q_i(X) := \prod_{j \neq i} (X - q_j) \quad (1 \leq i \leq n)$$

form a basis (over \mathbb{Q}) of the vector space $V = \{R \in \mathbb{Q}[X] : \deg(R) < n\}$. In other words, every $R \in \mathbb{Q}[X]$ with $\deg(R) < n$ can be written uniquely as

$$R(X) = \theta_1 Q_1(X) + \dots + \theta_n Q_n(X), \quad (4)$$

with $\theta_1, \dots, \theta_n$ in \mathbb{Q} . Evaluating this identity at $X = q_i$ yields

$$\theta_i = \frac{R(q_i)}{\prod_{j \neq i} (q_i - q_j)}. \quad (5)$$

In particular, one may take $R(X) = \prod_{i=1}^n (X - q_i) - P(X)$ (note that $\deg R < n$ since P is monic with degree n) and evaluate (4) at $X = \zeta$ to get :

$$\frac{\theta_1}{\zeta - q_1} + \dots + \frac{\theta_n}{\zeta - q_n} = 1,$$

It follows from (5) that θ_i must have the same sign as $(-1)^i P(q_i)$, and multiplying this identity by a large enough positive integer yields the lemma. \square

4. If λ is totally real, then \mathcal{F} contains a positive integer

Proof. We first apply Lemma 2 to $\zeta = \lambda^2$ with $q_i = i, 1 \leq i \leq n$. From (2) and the stability of \mathcal{F} under internal addition, it follows that the sum appearing in (3) is the difference of two elements in \mathcal{F} . In other words, we have found $\Delta \in \mathbb{N}^*$ and an element α with the following property:

$$\alpha \in \mathcal{F} \text{ and } \alpha - \Delta \in \mathcal{F}. \quad (6)$$

Since λ is a totally real algebraic number, it follows from part (a) of Lemma 1 and the minimality of \mathcal{F} that \mathcal{F} is included in the set of totally real algebraic numbers. In particular, α is a totally real algebraic number. In fact we may assume without loss of generality that $1 - \alpha$ is totally positive, because $\alpha' = \alpha + \beta$ will also satisfy (6) for any $\beta \in \mathcal{F}$, and choosing $\beta = \frac{k}{\lambda^2 - m}$ with $k, m \in \mathbb{N}$ large enough eventually makes $1 - \alpha'$ totally positive, by parts (b) and (c) of Lemma 1. Thus, parts (b) and (d) now guarantee that the number

$$\zeta' := \lambda^2(1 - \alpha)$$

is totally positive. Moreover, for $i, j \in \mathbb{N}$, we observe that

$$\begin{aligned} \left\{ \alpha, \alpha - \Delta, \frac{1}{\lambda^2} \right\} \subseteq \mathcal{F} &\stackrel{\text{(ADD)}}{\implies} (j(\Delta - 1) + 1)\alpha + j(\alpha - \Delta) + \frac{i}{\lambda^2} \in \mathcal{F} \\ &\stackrel{\text{(MAP)}}{\implies} \frac{1}{(\Delta j + 1)\zeta' - i} \in \mathcal{F} \\ &\stackrel{\text{(ADD)}}{\implies} \frac{1}{\zeta' - \frac{i}{\Delta j + 1}} \in \mathcal{F}, \end{aligned}$$

which shows that

$$\frac{1}{\zeta' - q} \in \mathcal{F} \quad \text{for every } q \in \mathcal{Q} := \left\{ \frac{i}{\Delta j + 1} : i, j \in \mathbb{N} \right\}. \quad (7)$$

Finally, let $P \in \mathbb{Q}[X]$ be the minimal polynomial of ζ' , which has $n := \deg(P)$ pairwise distinct non-negative zeros. Since \mathcal{Q} is dense in $[0, \infty)$, one can find $q_1 > \dots > q_n$ in \mathcal{Q} that interleave those zeros, in the sense that $P(q_i)$ has sign $(-1)^i$ for every $1 \leq i \leq n$. A second application of Lemma 2 now provides us with non-negative integers m_1, \dots, m_n such that

$$\frac{m_1}{\zeta' - q_1} + \dots + \frac{m_n}{\zeta' - q_n} \in \mathbb{N}^*.$$

Together with (7), this identity proves that \mathcal{F} contains a positive integer. \square

5. If \mathcal{F} contains a positive integer then \mathcal{F} is stable under negation

Proof. \mathcal{F} satisfies (ZERO) and (ADD), and hence so does the subset

$$\mathcal{F}^* := \mathcal{F} \cap (-\mathcal{F}).$$

We will now use the assumption to prove that \mathcal{F}^* also satisfies (MAP). This will imply the desired equality $\mathcal{F}^* = \mathcal{F}$, by minimality of \mathcal{F} . By assumption, \mathcal{F} contains a positive integer n , allowing us to write :

$$\begin{aligned} \alpha \in \mathcal{F}^* &\implies -\alpha \in \mathcal{F} \\ &\stackrel{\text{(ADD)}}{\implies} -(n-1)\alpha \in \mathcal{F} \\ &\stackrel{\text{(ADD)}}{\implies} -(n-1)\alpha + n \in \mathcal{F} \\ &\stackrel{\text{(MAP)}}{\implies} \frac{-1}{\lambda^2(1-\alpha)(n-1)} \in \mathcal{F} \\ &\stackrel{\text{(ADD)}}{\implies} \frac{-1}{\lambda^2(1-\alpha)} \in \mathcal{F}. \end{aligned}$$

Strictly speaking, the above argument is valid only if $n \neq 1$. But if $n = 1$, one may use the following alternative argument :

$$\alpha \in \mathcal{F}^* \implies -\alpha \in \mathcal{F} \xrightarrow{\text{(ADD)}} -\alpha + 2 \in \mathcal{F} \xrightarrow{\text{(MAP)}} \frac{-1}{\lambda^2(1-\alpha)} \in \mathcal{F}.$$

On the other hands, we trivially have

$$\alpha \in \mathcal{F}^* \implies \alpha \in \mathcal{F} \xrightarrow{\text{(MAP)}} \frac{1}{\lambda^2(1-\alpha)} \in \mathcal{F}.$$

Those two facts together precisely mean that \mathcal{F}^* satisfies (MAP). \square

6. If \mathcal{F} is stable under negation, then

$$\mathcal{F} = \left\{ \frac{P(\lambda^2)}{Q(\lambda^2)} : P, Q \in \mathbb{Z}[X], Q \text{ monic, } \deg(P) < \deg(Q) \right\}.$$

Proof. The inclusion \subseteq is immediate, since the set on the right-hand side satisfies (ZERO), (ADD) and (MAP). Conversely, we will now show that \mathcal{F} contains $\frac{P(\lambda^2)}{Q(\lambda^2)}$ for every $P, Q \in \mathbb{Z}[X]$ with Q monic and $\deg(P) < \deg(Q)$. The proof is by induction over $n = \deg Q$. The case $n = 0$ is simply the fact that $0 \in \mathcal{F}$. Now, assume that the claim holds for some $n \in \mathbb{N}$, and let

$$P(X) = p_n X^n + \cdots + p_0 \quad \text{and} \quad Q(X) = X^{n+1} + q_n X^n + \cdots + q_0,$$

with $p_0, \dots, p_n, q_0, \dots, q_n \in \mathbb{Z}$. Since $(\mathcal{F}, +)$ is a group by assumption, it is in fact enough to consider the following two special cases :

- Case 1 : $P(X) = X^n$.
- Case 2 : $P(X)$ is monic of degree n with $P(0) = 1$.

For the first case, observe that $\frac{1}{1+\lambda^{2n}} \in \mathcal{F}$ by our induction hypothesis and therefore,

$$\frac{1}{\lambda^{2n+2}} = \frac{1}{\lambda^2 \left(1 - \frac{1}{1+\lambda^{2n}}\right)} - \frac{1}{\lambda^2} \in \mathcal{F}.$$

Since \mathcal{F} also contains $\frac{1}{\lambda^2}, \dots, \frac{1}{\lambda^{2n}}$ by our induction hypothesis, one sees that

$$\frac{\lambda^{2n}}{Q(\lambda^2)} = \frac{1}{\lambda^2 \left(1 + \frac{q_n}{\lambda^2} + \cdots + \frac{q_0}{\lambda^{2n+2}}\right)} \in \mathcal{F},$$

as desired. In the second case, observe that

$$R(X) := P(X) - \frac{Q(X) - Q(0)P(X)}{X}$$

is an element of $\mathbb{Z}[X]$ with degree less than n , so our induction hypothesis guarantees that $\frac{R(\lambda^2)}{P(\lambda^2)} \in \mathcal{F}$, and therefore

$$\frac{P(\lambda^2)}{Q(\lambda^2)} = \frac{1}{\lambda^2 \left(1 - \frac{R(\lambda^2)}{P(\lambda^2)} + \frac{Q(0)}{\lambda^2}\right)} \in \mathcal{F}.$$

□

7. Acknowledgment

The author warmly thanks Louis-Hadrien Robert for helpful discussions on the problem.

References

- [1] D. M. Cvetković, M. Doob, H. Sachs, Spectra of graphs, 3rd Edition, Johann Ambrosius Barth, Heidelberg, 1995, theory and applications.
- [2] A. E. Brouwer, W. H. Haemers, Spectra of graphs, Universitext, Springer, New York, 2012. doi:10.1007/978-1-4614-1939-6.
URL <http://dx.doi.org/10.1007/978-1-4614-1939-6>
- [3] A. J. Hoffman, Eigenvalues of graphs, in: Studies in graph theory, Part II, Math. Assoc. Amer., Washington, D. C., 1975, pp. 225–245. Studies in Math., Vol. 12.
- [4] H. Bass, D. R. Estes, R. M. Guralnick, Eigenvalues of symmetric matrices and graphs, J. Algebra 168 (2) (1994) 536–567. doi:10.1006/jabr.1994.1244.
URL <http://dx.doi.org/10.1006/jabr.1994.1244>
- [5] S. Lang, Algebra, 3rd Edition, Vol. 211 of Graduate Texts in Mathematics, Springer-Verlag, New York, 2002. doi:10.1007/978-1-4613-0041-0.
URL <http://dx.doi.org/10.1007/978-1-4613-0041-0>