



Dust properties in comet 67P/Churyumov-Gerasimenko's coma during its 2008-2009 apparition from polarization observations

Edith Hadamcik, Anny Chantal Levasseur-Regourd, Jérémie Lasue, A. K. Sen, Rajiv Gupta

► To cite this version:

Edith Hadamcik, Anny Chantal Levasseur-Regourd, Jérémie Lasue, A. K. Sen, Rajiv Gupta. Dust properties in comet 67P/Churyumov-Gerasimenko's coma during its 2008-2009 apparition from polarization observations. European Planetary Science Congress 2010, Sep 2010, Rome, Italy. EPSC Abstracts, 5, pp. EPSC2010-879, 2010. hal-00793260

HAL Id: hal-00793260

<https://hal.science/hal-00793260>

Submitted on 11 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Dust properties in comet 67P/Churyumov-Gerasimenko's coma during its 2008-2009 apparition from polarization observations

E. Hadamcik (1) A.C. Levasseur-Regourd (1) J. Lasue (2), A.K. Sen (3) and R. Gupta (4)
(1) UPMC Univ. Paris 06, Université Versailles Saint Quentin, LATMOS-IPSL, 78280 Guyancourt, France
(2), LPI, Houston, USA, (3), Assam Univ., Silchar, India, (4), IUCAA, Pune, India. (edith.hadamcik@latmos.ipsl.fr),

Abstract

Comet 67P/Churyumov-Gerasimenko is the main target of the Rosetta mission. Remote observations from India and France, of the scattered light (and mainly its linear polarization) by the dust allowed us to conclude on different physical properties pre- and post-perihelion. Large dust particles are likely ejected from the surface pre-perihelion and small grains eventually in fluffy aggregates post-perihelion possibly originating from the subsurface.

1. Introduction

The ESA/Rosetta spacecraft will encounter comet 67P/Churyumov-Gerasimenko in 2014. The last observation period to prepare the mission was in 2008-2009. Remote observations of the light scattered by the dust in the coma allow to tentatively determine the physical properties of the particles.

The intensity and linear polarization evolution were followed in the coma from about 2 months pre-perihelion to about 2 months post-perihelion.

2. Observations

In a French-Indian collaboration, two telescopes were used with polarimetric imaging capabilities [1,2]. A 0.80 m telescope at observatoire de Haute-Provence

(OHP) in France and a 2 m telescope at Girawali Observatory in India. The observations periods are 25-27 December, 2008, 30 April and 1 May, 2009 in India. 17-19 March, 2009 in France. The phase angle ranged from 35.8° to 28.8°.

3. Results

For the two first periods the coma is asymmetric with very steep intensity radial decrease (down to -1.5) mainly in the solar direction in agreement with previous observations in 1982-83 and 1995-96. Two months after perihelion the coma is about isotropic with a radial decrease close to the nominal value -1. Before perihelion and two months after the aperture polarization values are comparable to polarization values measured on other comets at such phase angles. The sharp radial decrease in intensity and the feature in the tailward direction without any difference in polarization in the coma before perihelion suggest the presence of large dark particles. The post-perihelion increase in intensity (about 2 magnitude) and in polarization (1 per cent) is significantly above the error bars and suggests the ejection of dust particles with different physical properties. More polarized structures than in the surrounding coma are noticed on the polarization maps with evolution of the structures in 24 h suggesting small micron or sub-micron-sized grains eventually included in fluffy aggregates. An

important seasonal effect suggests that the different grains originate from different hemispheres of the nucleus.




Figure 1: Intensity images of the 3 periods : 1st column in negative, 2nd column treated images to enhance the structures.




Figure 2: Polarization maps about 3 weeks post-perihelion. FOV 40000 km.

4. Discussion

As suggested in the present work, [3] from their observations conclude on the presence of large particles pre-perihelion. [4] modeled the dust environment of the comet and found similar

conclusions. They suggest a strong seasonal effect due to the obliquity of the spin axis with ejection of small grains after perihelion.

From the study of thermal evolution of the nucleus as a function of its shape and of the obliquity of the axis, [5] discuss the relations between the nucleus activity and the internal structure. Differences in dust mantling depend on the inclination of the axis with a strong influence on the local surface and sub-surface activity and a North-South dichotomy of the nucleus.

Polarization and intensity variations in the coma of 67P/Churyumov-Gerasimenko are reminiscent of those noticed for some comets such as 9P/Tempel 1 and 81P/Wild 2.

5. Summary and Conclusions

The presence of rather large particles can be suggested before and just after perihelion and the ejection of smaller grains eventually in fluffy aggregates post-perihelion. An important seasonal effect related to the obliquity of the comet suggests that the different grains originate from the surface or subsurface and from different hemispheres of the nucleus.

Acknowledgements

We gratefully acknowledge ESA for the narrow-band filters at IGO. OHP and IUCAA for allocation of observation time, PNP for funding the French observations.

References

- [1] Levasseur-Regourd, A.C., Hadamcik, E., Sen, A.K., Gupta, R., Lasue, J., Recent polarimetric observations of comet 67P/Churyumov-Gerasimenko, In: Icy bodies in the solar system, IAU 263, Cambridge University Press, in press, 2010.
- [2] Hadamcik, E., Sen, A.K., Levasseur-Regourd, A.C., Gupta, R., Lasue, J., Polarimetric observations of comet 67P/Churyumov-Gerasimenko during its 2008-2009 apparition, A&A, (in press), 2010.
- [3] Tozzi, G.P., Boehnhardt, H., et al., Observations of 67P/Churyumov-Gerasimenko, BAAS, DPS 41, 2009.
- [4] Fulle, M., & the Giada team. Rosetta Science Working Team, December 2009.
- [5] De Sanctis, M.C., Lasue, J., Capria, M.T., et al., Shape and obliquity effects on the thermal evolution of the Rosetta target 67P/Churyumov-Gerasimenko cometary nuclei. Icarus 207, 341-358, 2010