



**HAL**  
open science

## Large Deviations for Quasi-Arithmetically Self-Normalized Random Variables

Jean-Marie Aubry, Marguerite Zani

► **To cite this version:**

Jean-Marie Aubry, Marguerite Zani. Large Deviations for Quasi-Arithmetically Self-Normalized Random Variables. ESAIM: Probability and Statistics, 2013, 17 (1), pp.1–12. hal-00796315

**HAL Id: hal-00796315**

**<https://hal.science/hal-00796315>**

Submitted on 4 Mar 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## LARGE DEVIATIONS FOR QUASI-ARITHMETICALLY SELF-NORMALIZED RANDOM VARIABLES

JEAN-MARIE AUBRY<sup>1</sup> AND MARGUERITE ZANI<sup>2</sup>

**Abstract.** We introduce a family of convex (concave) functions called sup (inf) of powers, which are used as generator functions for a special type of quasi-arithmetic means. Using these means we generalize the large deviation result that was obtained in the homogeneous case by Shao [14] on self-normalized statistics. Furthermore, in the homogenous case, we derive the Bahadur exact slope for tests using self-normalized statistics.

**Résumé.** Nous introduisons une famille de fonctions convexes (concaves) appelées sup (inf) de puissances, qui sont employées comme fonctions génératrices pour un type spécial de moyennes quasi-arithmétiques. À l'aide de ces moyennes, nous généralisons le résultat que Shao [14] a obtenu dans le cas homogène sur les statistiques autonormalisées. De plus, dans le cas homogène, nous obtenons la pente exacte de Bahadur pour les tests utilisant les statistiques autonormalisées.

**1991 Mathematics Subject Classification.** 60F10, 62F05.

The dates will be set by the publisher.

### INTRODUCTION

The use of self-normalization in statistics dates back to Student's  $t$ -distribution

$$T_n := \sqrt{n} \frac{\overline{X}_n - m}{S_n}$$

where  $\overline{X}_n := \frac{1}{n} \sum_{i=1}^n X_i$  is the empirical mean and  $S_n^2 := \frac{1}{n-1} \sum_{i=1}^n (X_i - \overline{X}_n)^2$  is the empirical unbiased variance of a Gaussian sample  $X_1, \dots, X_n$  of mean  $m$  and unknown variance. This statistic  $T_n$  is widely used to test  $m = m_0$  against alternative and it is well known that  $T_n$  has normal limit distribution when  $n \rightarrow \infty$  (even when the variables  $X_i$  are i.i.d. non Gaussian). Shao [14] proved the associated large deviations results for i.i.d. random variables  $X_i$  that are self-normalized by homogenous means, i.e. an asymptotic estimate for

$$\mathbb{P} \left( \frac{\overline{X}_n}{M_{p,n}} \geq x \right)^{\frac{1}{n}} \tag{1}$$

---

*Keywords and phrases:* large deviations, self-normalised statistics, Bahadur exact slope

<sup>1</sup> jmaubry@math.cnrs.fr

<sup>2</sup> zani@univ-paris12.fr

Laboratoire d'Analyse et Mathématiques Appliquées (CNRS UMR 8050), Université Paris-Est Créteil, 61 av. du Général de Gaulle, 94010 Créteil Cedex, France.

with  $M_{p,n} := \left(\frac{1}{n} \sum_{i=1}^n |X_i|^p\right)^{\frac{1}{p}}$ ,  $p > 1$  and minimal hypotheses on the common law of  $X_i$ . There have been several further papers on this subject by Dembo and Shao [6, 7], Lai and Shao [10], Shao [15] and a book by de la Pena, Lai and Shao [5]. For related results see also Tchirina [16].

The difficulty here is, of course, that the self-normalized random variables  $X_i/M_{p,n}$  are far from independent and the result is indeed quite different from the classical Cramér-Chernoff large deviations [2]; in particular, no exponential moment hypothesis on  $X_i$  (actually, no moment hypothesis at all) is necessary. It is worth noting that Shao's applications, notably to Student's  $t$ -distribution, make only use of the case  $p = 2$ , i.e. the *quadratic* mean self-normalization; besides, the results for other values of  $p > 1$  (the so-called *Hölder* or *power* means) present no extra arduousness.

There has however been a renewed interest in recent litterature for *quasi-arithmetic* means, where the power function  $x \mapsto x^p$  is replaced by a more general strictly monotonic and continuous *generator function*  $\phi : [0, \infty) \rightarrow [0, \infty)$ . More precisely, by a famous result of Kolmogorov [9] and Nagumo [12], any reasonable notion of mean can be written as

$$M_{\phi,n} := \phi^{-1} \left( \frac{1}{n} \sum_{i=1}^n \phi(X_i) \right)$$

for such a function  $\phi$ . Modern applications of quasi-arithmetic means in a probabilistic framework range from Ordered Weighted Averaging operators, introduced in decision theory by Yager [17], to statistics, because of their links with Archimedean copulas [3, 4] and covariance functions [13].

Another potential application is the following. Suppose that the i.i.d. variables  $X_i$  follow an unknown law  $\mu(\theta, c)$ , where  $c$  is the parameter of interest. In cases where  $\theta$  is simply a scale parameter (meaning that there exists a function  $\nu$  such that the law of  $X_i/\nu(\theta)$  does not depend on  $\theta$ ), self-normalization by any power mean yields a scale-free statistic that can be used to test for  $c$ . Large-deviation type results can then be translated in terms of Bahadur asymptotic efficiency, see § 3. In more general cases, one can try and find  $\phi$  such that  $\frac{E(X)}{\phi^{-1}(E(\phi(X)))} = f(c)$  and does not depend on  $\theta$ . This seems to be a difficult problem, but ad-hoc examples (starting from  $\phi$ ) can be constructed to show at least that the class of laws  $\mu(\theta, c)$  for which it can be solved is not empty. Suppose that for  $\theta > 0$  and  $\sqrt{3}/2 < c < 1$ ,  $\mu(\theta, c)$  is the uniform law on  $[\theta, \theta + \lambda(\theta, c)]$  where

$$\lambda(\theta, c) = \frac{6\theta - 3c^2(1 + 2\theta) + c \left( 3 + \sqrt{3(1-c)(3 + 4\theta^2 + c(-3 + 4\theta + 4\theta^2))} \right)}{4c^2 - 3}$$

Then for  $\phi(x) = x + x^2$  one has indeed  $\frac{E(X)}{\phi^{-1}(E(\phi(X)))} = c$  for all  $\theta$  and the statistic

$$T_n := \frac{\overline{X_n}}{M_{\phi,n}} \quad (2)$$

can be used to test for  $c$ .

So the natural question arises, whether Shao's asymptotic results on (1) can be extended using a more general quasi-arithmetic mean  $M_{\phi,n}$ . In this paper, we show large deviations for statistics of type (2). The main drawback and technical difficulty is that now  $M_{\phi,n}$  is no longer homogenous (indeed the only homogeneous quasi-arithmetic means are the power means and the geometric mean, see [8, p. 68]). For this reason, we need to restrict ourselves to a subclass of quasi-arithmetic means where the function  $\phi$  enjoys some extra properties, studied in § 1. The main theorem on self-normalized large deviations is then stated and proved in § 2. In § 3, we use these results to determine Bahadur exact slope for self-normalized statistics.

## 1. INFIMUM AND SUPREMUM OF POWERS

We shall apply Shao's method in [14] to a more general class of self-normalized large deviations, where the power function  $x \mapsto x^p$  used as a generator function for the normalizing mean is replaced by a more general

convex function  $\phi$  or, equivalently,  $x \mapsto x^{\frac{1}{p}}$  is replaced by the concave function  $\psi := \phi^{-1}$ . The first step of our approach consists in writing  $\psi$  as an infimum of concave power functions, which can be done only in certain cases. Alternatively, the condition can be expressed in terms of  $\phi$  written as a supremum of convex power functions.

### 1.1. Characterization of admissible generator functions

In the following,  $\partial\omega(x)$  denotes the superdifferential (resp. subdifferential) at  $x$  of the concave (resp. convex) function  $\omega$ ;  $a \vee b$  denotes the maximum of  $a$  and  $b$ .

**Proposition 1.1.** *For  $\psi : [0, \infty) \rightarrow [0, \infty)$ , the following are equivalent.*

(i) *There exists  $\mu : [1, \infty) \rightarrow (0, \infty)$  such that for all  $x > 0$ ,*

$$\psi(x) = \inf_{p \geq 1} \mu(p)x^{\frac{1}{p}} \quad (3)$$

(ii) *The function  $\omega : x \mapsto \log(\psi(\exp(x)))$  is proper closed concave on  $\mathbb{R}$  and  $\partial\omega(x) \subset [0, 1]$  for all  $x \in \mathbb{R}$ .*

*Proof.* Let us start by assuming (i). Then, writing  $\alpha := \frac{1}{p}$ ,  $y := \log(x)$  and  $\vartheta(\alpha) := -\log(\mu(\frac{1}{\alpha}))$ , we turn (3) into

$$\omega(y) = \inf_{0 < \alpha \leq 1} \alpha y - \vartheta(\alpha) \quad (4)$$

So  $\omega$  is the concave conjugate of  $\tilde{\vartheta}$ , the concave hull of  $\vartheta$  restricted to  $(0, 1]$ ; the closure of  $\tilde{\vartheta}$  has support  $\subset [0, 1]$  hence the conclusion.

Conversely, suppose that (ii) holds. Let  $\vartheta$  be the concave conjugate of  $\omega$ , it is a proper closed concave function with support  $\subset [0, 1]$  because of the hypothesis on the superdifferential of  $\omega$ . Inverting the Fenchel-Legendre transformation, we can write  $\omega(y) = \inf_{0 \leq \alpha \leq 1} \alpha y - \vartheta(\alpha)$ , which by closedness is equivalent to (4), hence (i) with  $\mu(p) := \exp(-\vartheta(\frac{1}{p}))$ . □

Remarks: (i) clearly implies that  $\psi$  itself is nondecreasing, concave and that  $\psi(0) = 0$  but this is not enough (take  $\psi(x) = x + \sqrt{x}$  for instance). The example  $\mu(p) = 1$  for all  $p \geq 1$  shows however that  $\psi$  is not necessarily (strictly) increasing. Possible infinite values for  $\mu$  mean that the infimum on  $p$  is actually taken on the support  $S_\mu := [1, \infty) \setminus \mu^{-1}(\{\infty\})$ .

When a function  $\psi$  satisfies the hypothesis of Proposition 1.1, we shall say that it is an *infimum of powers* with weight  $\mu$ . Note that we can (and will) always assume, without loss of generality, that  $\mu(p) = \exp(-\vartheta(\frac{1}{p}))$  for a proper closed concave function  $\vartheta$ ; in particular  $\mu$  is continuous on its support. We shall say in that case that  $\mu$  is a *regular weight* and remark that  $\inf_{p \geq 1} \mu(p) > 0$ . If furthermore  $\psi$  is differentiable, then  $\mu(p)$  can be parametrized by the contact point  $s > 0$ ,  $p(s) = \frac{\psi(s)}{s\psi'(s)}$  and  $\mu(p(s)) = \psi(s)s^{-\frac{1}{p(s)}}$  (see examples below).

**Proposition 1.2.** *For  $\phi : [0, \infty) \rightarrow [0, \infty]$ , the following are equivalent.*

(i) *There exists  $\nu : [1, \infty) \rightarrow [0, \infty)$  such that for all  $x > 0$ ,*

$$\phi(x) = \sup_{p \geq 1} \nu(p)x^p \quad (5)$$

(ii) *The function  $\omega : x \mapsto \log(\phi(\exp(x)))$  is proper closed convex on  $\mathbb{R}$  and  $\partial\omega(x) \subset [1, \infty)$  for all  $x \in \mathbb{R}$  where  $\phi(x) < \infty$ .*

The proof is very similar to that of Proposition 1.1 and is omitted (note that  $\phi$  can take infinite values).

Examples: some inf of powers with parametrization.

$\psi(x)$	$\omega(x)$	$p(s)$	$\mu(p(s))$
$\log(1+x)$	$\log \log(1+e^x)$	$\frac{(1+s)\log(1+s)}{s}$	$s^{\frac{s}{(1+s)\log(1+s)}} \log(1+s)$
$\sqrt{1+x}-1$	$\log(\sqrt{1+e^x}-1)$	$\frac{2(1+s-\sqrt{1+s})}{s}$	$s^{\frac{s}{2(-1-s+\sqrt{1+s})}} (\sqrt{1+s}-1)$
$1-e^{-x}$	$\log(1-e^{-e^x})$	$\frac{e^s-1}{s}$	$(1-e^{-s})s^{\frac{s}{e^s-1}}$

## 1.2. Bijectivity

Let us give the necessary and sufficient conditions under which an infimum (resp. supremum) of powers is bijective, in which case its converse is a supremum (resp. infimum) of powers.

**Proposition 1.3.** *Let  $\psi$  be an inf of powers with regular weight  $\mu$ . Then the following are equivalent.*

- (i)  $\lim_{p \rightarrow \infty} \mu(p) = \infty$ ;
- (ii)  $\psi$  is unbounded;
- (iii)  $\psi$  is a bijection :  $[0, \infty) \rightarrow [0, \infty)$  and  $\psi^{-1}(y) = \sup_{p \geq 1} \mu(p)^{-p} y^p$ .

*Proof.* Suppose (i), fix any  $C < \infty$  and pick  $p_0$  such that  $\mu(p) \geq C$  when  $p \geq p_0$ . Then choose  $x := 1 \vee \left(\frac{C}{\inf_{p > 1} \mu(p)}\right)^{p_0}$ . We thus have  $\inf_{p \geq p_0} \mu(p)x^{\frac{1}{p}} \geq C$  and  $\inf_{1 < p \leq p_0} \mu(p)x^{\frac{1}{p}} \geq (\inf_{p > 1} \mu(p))x^{\frac{1}{p_0}} \geq C$  so  $\psi(x) \geq C$ . Conversely, since  $\mu(p) = \exp(-\vartheta(\frac{1}{p}))$  with  $\vartheta$  concave, if  $\mu$  does not have infinite limit at  $\infty$  then it has to be bounded. In that case  $\psi$  is also bounded (by the same constant).

Being concave and taking value 0 at 0,  $\psi$  unbounded implies that it is strictly increasing and surjective on  $[0, \infty)$ , hence it is a bijection. Furthermore we have  $x \geq \psi^{-1}(y) \iff y \leq \psi(x) \iff \forall p > 1, y \leq \mu(p)x^{\frac{1}{p}} \iff \forall p > 1, x \geq \mu(p)^{-p} y^p \iff x \geq \sup_{p > 1} \mu(p)^{-p} y^p$ . The converse is trivial.  $\square$

**Proposition 1.4.** *Let  $\phi$  be a sup of powers with regular weight  $\nu$ . Then the following are equivalent.*

- (i)  $\lim_{p \rightarrow \infty} \nu(p)^{\frac{1}{p}} = 0$ ;
- (ii)  $\phi$  is finitely-valued;
- (iii)  $\phi$  is a bijection :  $[0, \infty) \rightarrow [0, \infty)$  and  $\phi^{-1}(x) = \inf_{p \geq 1} \nu(p)^{-\frac{1}{p}} x^{\frac{1}{p}}$ .

The proof is very similar to that of Proposition 1.3 and is omitted.

## 1.3. Operations

We shall now see that the class of functions satisfying the inf (resp. sup) of powers property is ‘‘large’’, as it is under minor conditions stable by powers, composition, and products. We give the proofs only in the inf case.

**Proposition 1.5.** *If  $\psi$  is an inf (resp. sup) of powers, then so is  $\psi^\alpha : x \mapsto \psi(x)^\alpha$  for  $0 < \alpha \leq 1$  (resp. for  $\alpha \geq 1$ ).*

*Proof.* Take  $\mu^\alpha(p) := \mu(\alpha p)^\alpha$  if  $p > \frac{1}{\alpha}$ ,  $\mu^\alpha(p) := \infty$  else.  $\square$

**Proposition 1.6.** *If  $\psi_1$  and  $\psi_2$  are inf (resp. sup) of powers, then so is  $\psi_1 \circ \psi_2$ .*

*Proof.* Using Proposition 1.5,

$$\begin{aligned}\psi_1(\psi_2(x)) &= \inf_{p_1 > 1} \mu_1(p_1) \psi_2(x)^{\frac{1}{p_1}} \\ &= \inf_{p_2 > p_1 > 1} \mu_1(p_1) \mu_2\left(\frac{p_2}{p_1}\right)^{\frac{1}{p_1}} x^{\frac{1}{p_2}} \\ &= \inf_{p_2 > 1} \underbrace{\inf_{p_1 \in (1, p_2)} \mu_1(p_1) \mu_2\left(\frac{p_2}{p_1}\right)^{\frac{1}{p_1}}}_{=:\mu(p_2)} x^{\frac{1}{p_2}}\end{aligned}$$

□

**Proposition 1.7.** *If  $\psi_1$  and  $\psi_2$  are inf of powers with  $p_1 \leq \frac{1}{\alpha_1} \Rightarrow \mu_1(p_1) = \infty$  and  $p_2 \leq \frac{1}{\alpha_2} \Rightarrow \mu_2(p_2) = \infty$  for some  $\alpha_1 + \alpha_2 \leq 1$ , then  $\psi_1\psi_2$  is also an inf of powers. If  $\phi_1$  and  $\phi_2$  are sup of powers (no extra condition), then so is  $\phi_1\phi_2$ .*

*Proof.* We can write

$$\begin{aligned}\psi_1(x)\psi_2(x) &= \inf_{p_1 > \frac{1}{\alpha_1}} \inf_{p_2 > \frac{1}{\alpha_2}} \mu_1(p_1) \mu_2(p_2) x^{\frac{1}{p_1} + \frac{1}{p_2}} \\ &= \inf_{p > 1} \underbrace{\inf_{\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p}} \mu_1(p_1) \mu_2(p_2)}_{=:\mu(p)} x^{\frac{1}{p}}\end{aligned}$$

□

## 2. SELF-NORMALIZED LARGE DEVIATIONS

Let  $\psi$  be a bijective inf of powers with weight  $\mu$  (cf. Proposition 1.3) and let  $\phi := \psi^{-1}$  extended to  $\mathbb{R}$  by even symmetry ( $\phi(x) := \phi(|x|)$ ). Let  $(X_i)_{i \geq 1}$  be a sequence of independent random variables having the same law as an arbitrary  $X \neq 0$ . Their empirical mean

$$\overline{X}_n := \frac{1}{n} \sum_{i=1}^n X_i$$

will be normalized by the quasi-arithmetic mean

$$M_{\phi, n} := \psi\left(\frac{1}{n} \sum_{i=1}^n \phi(X_i)\right)$$

**Theorem 1.** *Suppose that there exist  $1 < \pi_0 < \pi_1 < \infty$  such that  $\mu(p) < \infty$  only if  $p \in S_\mu := [\pi_0, \pi_1]$ .*

*Then for any  $x > \begin{cases} \frac{0 \vee \mathbb{E}X}{\psi(\mathbb{E}\phi(X))} & \text{if } E(|X|) < \infty \\ 0 & \text{if } E(|X|) = \infty \end{cases}$ ,*

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\frac{\overline{X}_n}{M_{\phi, n}} \geq x\right)^{\frac{1}{n}} = \sup_{c > 0, p \in S_\mu} \inf_{t \geq 0} \mathbb{E} \exp\left(t\left(cX - x \frac{\mu(p)}{p} (\phi(X) + (p-1)c^{\frac{p}{p-1}})\right)\right)$$

*Proof.* For short we shall always write  $q := \frac{p}{p-1}$ , whenever  $p > 1$  is considered. We repeatedly use the fact that  $x \mapsto \frac{x^p}{p}$  and  $y \mapsto \frac{y^q}{q}$  are convex conjugate functions and that for any  $x, y \geq 0$  we have  $xy = \inf_{b>0} \frac{(b^{-1}x)^p}{p} + \frac{(by)^q}{q}$ .

Since  $\psi$  is an inf of powers, we obtain for  $x \geq 0$

$$\begin{aligned}
\mathbb{P}\left(\frac{\overline{X_n}}{M_{\phi,n}} \geq x\right) &= \mathbb{P}\left(\sum_{i=1}^n X_i \geq \inf_{p \in S_\mu} x\mu(p)n^{\frac{1}{q}} \left(\sum_{i=1}^n \phi(X_i)\right)^{\frac{1}{p}}\right) \\
&= \mathbb{P}\left(\sum_{i=1}^n X_i \geq \inf_{b>0, p \in S_\mu} x\mu(p) \left(\frac{b^{-p}}{p} \sum_{i=1}^n \phi(X_i) + n\frac{b^q}{q}\right)\right) \\
&= \mathbb{P}\left(\sup_{b>0, p \in S_\mu} \sum_{i=1}^n \left(X_i - x\mu(p) \left(\frac{b^{-p}}{p} \phi(X_i) + \frac{b^q}{q}\right)\right) \geq 0\right) \\
&= \mathbb{P}\left(\sup_{c>0, p \in S_\mu} \sum_{i=1}^n \left(cX_i - x\frac{\mu(p)}{p}(\phi(X_i) + (p-1)c^q)\right) \geq 0\right) \tag{6}
\end{aligned}$$

Then we can finish along the general lines of [14], with a significant complication due to the non-homogeneity of  $\phi$ . We prove the lower bound in § 2.1 and the upper bound in § 2.2.  $\square$

Remark: the hypothesis on the support of  $\mu$  excludes some interesting examples. It could be replaced by a decay hypothesis on the tail distribution of  $X$ ; this aspect will be studied in a subsequent paper.

## 2.1. Lower bound

Suppose first  $E(|X|) < \infty$ . Since we look for one-sided large deviations in (6) we need to have

$$\mathbb{E}\left(cX - x\frac{\mu(p)}{p}(\phi(X) + (p-1)c^q)\right) < 0 \tag{7}$$

Observe that, by Young's inequality, with  $\beta(p) := \frac{\mu(p)^{1-p}}{p} \left(\frac{0 \vee \mathbb{E}X}{x}\right)^p$  we have for all values of  $c > 0$  and  $p \in S_\mu$

$$c\frac{0 \vee \mathbb{E}X}{x} - \mu(p)\frac{c^q}{q} - \beta(p) \leq 0 \tag{8}$$

while the hypothesis on  $x$  combined to (3) implies  $x > \frac{0 \vee \mathbb{E}X}{\mu(p)(\mathbb{E}\phi(X))^{\frac{1}{p}}}$ , hence

$$\beta(p) - \frac{\mu(p)}{p}\mathbb{E}\phi(X) < 0 \tag{9}$$

Adding up (8) and (9) yields precisely (7) and we obtain

$$\liminf_{n \rightarrow \infty} \mathbb{P}\left(\frac{\overline{X_n}}{M_{\phi,n}} \geq x\right)^{\frac{1}{n}} \geq \inf_{t \geq 0} \mathbb{E} \exp\left(t\left(cX - x\frac{\mu(p)}{p}(\phi(X) + (p-1)c^q)\right)\right) \tag{10}$$

If  $E(|X|) = \infty$  and  $x > 0$  then the right-hand side of (10) is zero, so the inequality is trivial.

Remark: so far we used no assumption on  $\phi$ , except that it takes non-negative values on the range of  $X$ . When  $\phi = \psi^{-1}$ , this function is convex and Jensen's inequality implies that the lower bound on  $x$  is  $\leq 1$ .

## 2.2. Upper bound

Now we use the fact that  $\phi := \psi^{-1}$ . Let  $K > 3$  be fixed and let  $[\pi_0', \pi_1'] := S'_\mu := \left\{ p \in S_\mu : \frac{\mu(p)}{p} \leq C \right\}$ , where  $C$  is determined by the following (proof in Appendix A):

**Lemma 2.1.** *There exists  $C < \infty$  such that  $\inf_{t \geq 0} \mathbb{E} \exp(t(KX - xC\phi(X))) \leq \frac{1}{K}$ .*

We deduce from (6) that  $\mathbb{P}\left(\frac{\overline{X_n}}{M_{\phi,n}} \geq x\right) \leq I_1 + I_2 + I_3$ , with

$$\begin{aligned} I_1 &:= \mathbb{P}\left(\sup_{c > K, p \in S_\mu} \sum_{i=1}^n \left(cX_i - x \frac{\mu(p)}{p} (\phi(X_i) + (p-1)c^q)\right) \geq 0\right) \\ I_2 &:= \mathbb{P}\left(\sup_{0 < c \leq K, p \in S'_\mu} \sum_{i=1}^n \left(cX_i - x \frac{\mu(p)}{p} (\phi(X_i) + (p-1)c^q)\right) \geq 0\right) \\ I_3 &:= \mathbb{P}\left(\sup_{0 < c \leq K, p \in S_\mu \setminus S'_\mu} \sum_{i=1}^n \left(cX_i - x \frac{\mu(p)}{p} (\phi(X_i) + (p-1)c^q)\right) \geq 0\right) \end{aligned}$$

We start with  $I_1$ . Let  $K'(p) := \frac{xK^{q-1}\mu(p)}{2q}$ : the restriction  $c > K$  implies that  $K'(p) < x \frac{\mu(p)}{2p} (p-1)c^{q-1}$ , so

$$\begin{aligned} &\sup_{c > K, p \in S_\mu} \sum_{i=1}^n \left(cX_i - x \frac{\mu(p)}{p} (\phi(X_i) + (p-1)c^q)\right) \geq 0 \\ \Rightarrow &\sup_{c > K, p \in S_\mu} \sum_{i=1}^n \left(|X_i| \mathbf{1}_{|X_i| > K'(p)} + K'(p) - x \frac{\mu(p)}{p} \left(\frac{\phi(X_i)}{c} + (p-1)c^{q-1}\right)\right) \geq 0 \\ \Rightarrow &\sup_{p \in S_\mu} \sum_{i=1}^n |X_i| \mathbf{1}_{|X_i| > K'(p)} - x \frac{\mu(p)}{p} \inf_{c > K} \sum_{i=1}^n \left(\frac{\phi(X_i)}{c} + \frac{1}{2}(p-1)c^{q-1}\right) \geq 0 \\ \Rightarrow &\sup_{p \in S_\mu} \sum_{i=1}^n |X_i| \mathbf{1}_{|X_i| > K'(p)} - x \mu(p) \inf_{b > K^{\frac{1}{p}}} \left(\frac{1}{p} \left(\frac{(\sum_{i=1}^n \phi(X_i))^{\frac{1}{p}}}{b}\right)^p + \frac{1}{q} \left(\left(\frac{n}{2}\right)^{\frac{1}{q}} b\right)^q\right) \geq 0 \end{aligned}$$

by Young's inequality,

$$\Rightarrow \sup_{p \in S_\mu} \sum_{i=1}^n |X_i| \mathbf{1}_{|X_i| > K'(p)} - x \mu(p) \left(\sum_{i=1}^n \phi(X_i)\right)^{\frac{1}{p}} \left(\frac{n}{2}\right)^{\frac{1}{q}} \geq 0$$

by Proposition 1.3,

$$\Rightarrow \sup_{p \in S_\mu} \sum_{i=1}^n |X_i| \mathbf{1}_{|X_i| > K'(p)} - x \left(\sum_{i=1}^n |X_i|^p\right)^{\frac{1}{p}} \left(\frac{n}{2}\right)^{\frac{1}{q}} \geq 0$$

and using Hölder's inequality,

$$\Rightarrow \sup_{p \in S_\mu} \sum_{i=1}^n \mathbf{1}_{|X_i| > K'(p)} - \frac{n}{2} x^q \geq 0$$

Finally we obtain that

$$\begin{aligned} I_1 &\leq \mathbb{P}\left(\sup_{p \in S_\mu} \sum_{i=1}^n \left(\mathbf{1}_{|X_i| > K'(p)} - \frac{x^q}{2}\right) \geq 0\right) \\ &\leq \mathbb{P}\left(\sum_{i=1}^n \left(\mathbf{1}_{|X_i| > \inf_{p \in S_\mu} K'(p)} - \tau\right) \geq 0\right) \end{aligned}$$

with  $\tau := \min\left(\frac{x^{\frac{\pi_0}{2}}}{2}, 1\right) > 0$ . Applying the Chernoff bound on binomial random variables, we deduce that

$$\limsup_{n \rightarrow \infty} I_1^{\frac{1}{n}} \leq \left(\frac{e}{\tau} \mathbb{P}\left(|X| > \inf_{p \in S_\mu} K'(p)\right)\right)^\tau$$

Since  $\inf_{p \in S_\mu} \mu(p) > 0$ , when  $K \rightarrow \infty$  the quantity  $\inf_{p \in S_\mu} K'(p) \rightarrow \infty$  and therefore  $\limsup_{n \rightarrow \infty} I_1^{\frac{1}{n}} \rightarrow 0$ .

Let us now bound  $I_2$ . We discretize  $c \in [0, K]$  and  $p \in S'_\mu$  with a step  $\Delta$  to be chosen later (Lemma 2.2). Let  $T_l$  denote the interval  $[\pi_0' + l\Delta, \pi_0' + (l+1)\Delta]$  and let  $p_l$  realize  $\inf_{p \in T_l} \frac{\mu(p)}{p}$ .

$$\begin{aligned} I_2 &\leq \sum_{j=1}^{\frac{K}{\Delta}} \sum_{l=0}^{\frac{\pi_1' - \pi_0'}{\Delta} - 1} \mathbb{P}\left(\sum_{i=1}^n \left(j\Delta X_i - x \frac{\mu(p_l)}{p_l} \left(\Phi(X_i) + \inf_{p \in T_l} (p-1)((j-1)\Delta)^q\right)\right) \geq 0\right) \\ &\leq \sum_{j=1}^{\frac{K}{\Delta}} \sum_{l=0}^{\frac{\pi_1' - \pi_0'}{\Delta} - 1} \inf_{t \geq 0} \left(\mathbb{E} \exp\left(t \left(j\Delta X - x \frac{\mu(p_l)}{p_l} \left(\Phi(X) + \inf_{p \in T_l} (p-1)((j-1)\Delta)^q\right)\right)\right)\right)^n \end{aligned}$$

By our choice of  $C$  in Lemma 2.1, for any  $0 \leq l < \frac{\pi_1' - \pi_0'}{\Delta}$  we have  $\frac{\mu(p_l)}{p_l} \leq C$ . Let  $Y_{j,l}$  be standard Gaussian independent random variables,  $\delta > 0$  and

$$\xi_{j,l} := \delta Y_{j,l} + j\Delta X - x \frac{\mu(p_l)}{p_l} \left(\Phi(X) + (p_l - 1)(j\Delta)^{\frac{p_l}{p_l-1}}\right)$$

**Lemma 2.2.** *There exists  $\Delta > 0$  such that for all  $j, l$ , the infimum over  $t \geq 0$  of  $\exp(t\xi_{j,l})$  is reached for  $t_{j,l} \in \left[0, \frac{1}{\Delta K^{\frac{\pi_0}{\pi_0-1} + 1}}\right]$ .*

Let us now define  $D(j, l) := (p_l - 1)(j\Delta)^{\frac{p_l}{p_l-1}} - \inf_{p \in T_l} (p-1)((j-1)\Delta)^q$ .

**Lemma 2.3.** *There exists a constant  $C' < \infty$  (depending only on  $\pi_0$  and  $\pi_1$ ) such that  $\max_{j,l} D(j, l) \leq C' K^{\frac{\pi_0}{\pi_0-1}} \log(K)\Delta$ .*

Using Lemma 2.2,

$$\begin{aligned} I_2 &\leq \sum_{j=1}^{\frac{K}{\Delta}} \sum_{l=0}^{\frac{\pi_1 - \pi_0}{\Delta} - 1} \exp\left(n x t_{j,l} \frac{\mu(p_l)}{p_l} D(j, l)\right) \left(\mathbb{E} \exp(t_{j,l} \xi_{j,l})\right)^n \\ &\leq \frac{K(\pi_1 - \pi_0)}{\Delta^2} \exp\left(n \frac{x C}{\Delta K^{\frac{\pi_0}{\pi_0-1} + 1}} \max_{j,l} D(j, l)\right) \\ &\quad \times \sup_{c > 0, p \in S'_\mu} \inf_{t \geq 0} \left(e^{\frac{(t\delta)^2}{2}} \mathbb{E} \exp\left(t \left(cX - x \frac{\mu(p)}{p} \left(\Phi(X) + (p-1)c^q\right)\right)\right)\right)^n \end{aligned}$$

and from Lemma 2.3 we deduce that

$$\limsup_{n \rightarrow \infty} I_2^{\frac{1}{n}} \leq \exp \left( x C'' \frac{\log(K)}{K} \right) \sup_{c > 0, p \in S'_\mu} \inf_{t \geq 0} e^{\frac{(t\delta)^2}{2}} \mathbb{E} \exp \left( t \left( cX - x \frac{\mu(p)}{p} (\Phi(X) + (p-1)c^q) \right) \right)$$

The bound on  $I_3$  is simpler: since  $p \in S_\mu \setminus S'_\mu$ ,

$$I_3 \leq \mathbb{P} \left( \sum_{i=1}^n KX_i - xC\phi(X_1) \right) \leq \inf_{t \geq 0} (\mathbb{E} \exp(t(KX - xC\phi(X))))^n \leq \left( \frac{1}{K} \right)^n$$

by Lemma 2.1, hence  $\limsup_{n \rightarrow \infty} I_3^{\frac{1}{n}} \rightarrow 0$  when  $K \rightarrow \infty$ .

Putting all together, the upper bound follows by letting  $K \rightarrow \infty$ : for any  $\delta > 0$  we have

$$\limsup_{n \rightarrow \infty} \mathbb{P} \left( \frac{\overline{X}_n}{M_{\phi,n}} \geq x \right)^{\frac{1}{n}} \leq \sup_{c > 0, p \in S_\mu} \inf_{t \geq 0} e^{\frac{(t\delta)^2}{2}} \mathbb{E} \exp \left( t \left( cX - x \frac{\mu(p)}{p} (\Phi(X) + (p-1)c^q) \right) \right)$$

Combining this with the following completes the proof of Theorem 1.

**Lemma 2.4** (straightforward adaptation from [14, Lemma 2.1]). *For any random variable  $X$  we have*

$$\begin{aligned} \limsup_{\delta \searrow 0} \inf_{b \geq 0} \inf_{t \geq 0} e^{\frac{(t\delta)^2}{2}} \mathbb{E} \exp \left( t \left( cX - x \frac{\mu(p)}{p} (\Phi(X) + (p-1)c^q) \right) \right) \\ = \sup_{b \geq 0} \inf_{t \geq 0} \mathbb{E} \exp \left( t \left( cX - x \frac{\mu(p)}{p} (\Phi(X) + (p-1)c^q) \right) \right) \end{aligned}$$

for  $x > \frac{0 \vee \mathbb{E}X}{\psi(\mathbb{E}\phi(X))}$ . Moreover, the convergence is uniform in  $x \in [\alpha, 1]$  for any  $\frac{0 \vee \mathbb{E}X}{\psi(\mathbb{E}\phi(X))} < \alpha < 1$ .

### 3. AN APPLICATION TO BAHADUR EXACT SLOPE

#### 3.1. Bahadur results

Let us recall here some basic facts about Bahadur exact slopes of test statistics. For a reference, see [1] and [11]. Consider a sample  $X_1, \dots, X_n$  having common law  $\mu_\theta$  depending on a parameter  $\theta \in \Theta$ . To test  $(\mathcal{H}_0) : \theta \in \Theta_0$  against the alternative  $(\mathcal{H}_1) : \theta \in \Theta_1 := \Theta \setminus \Theta_0$ , we use a test statistic  $S_n$ , large values of  $S_n$  rejecting the null hypothesis. The  $p$ -value of this test is by definition  $H_n(S_n)$ , where

$$H_n(t) := \sup_{\theta \in \Theta_0} P_\theta(S_n \geq t)$$

The Bahadur exact slope  $c(\theta)$  of  $S_n$  is then given by the following relation

$$c(\theta) = -2 \liminf_{n \rightarrow \infty} \frac{1}{n} \log(H_n(S_n)) \quad (11)$$

Quantitatively, for  $\theta \in \Theta_1$ , the larger  $c(\theta)$  is, the faster  $S_n$  rejects  $H_0$ .

A theorem of Bahadur (Theorem 7.2. in [1]) gives the following characterization of  $c(\theta)$ : suppose that  $\lim_n n^{-1/2} S_n =: b(\theta)$  for any  $\theta \in \Theta_1$ , and that  $\lim_n n^{-1} \log(H_n(n^{1/2}t)) =: -F(t)$  under any  $\theta \in \Theta_0$ . If  $F$  is continuous on an interval  $I$  containing  $b(\Theta_1)$ , then  $c(\theta)$  is given by:

$$c(\theta) = 2F(b(\theta)) \quad (12)$$

### 3.2. The homogeneous case

For the seek of simplicity, we assume now that we are in the homogeneous case, i.e.  $M_{\phi,n} = M_{p,n} := (\frac{1}{n} \sum_{i=1}^n |X_i|^p)^{\frac{1}{p}}$ , with  $p > 1$ . Proofs can be adapted to the general case, adding suitable assumptions on  $\phi$ .

We want to test whether the sample  $X_1, \dots, X_n$  has exponential distribution  $\mathcal{E}(\beta)$ , where  $\beta > 0$  is unknown (null hypothesis  $H_0$ ), against the alternative that it has gamma distribution  $\gamma(\theta, \beta)$  for some  $\theta > 1$  and  $\beta > 0$  (alternative hypothesis  $H_1$ ). To achieve this we use the statistic

$$T_n = \frac{\overline{X_n}}{M_{p,n}}$$

Under  $H_0$ ,  $T_n$  is scale-free, therefore we can assume  $\beta = 1$ .

From Theorem 1.1. of Shao [14],

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log P(T_n \geq x) = -I(x)$$

where

$$I(x) = -\log \left( \sup_{c>0} \inf_{t \geq 0} \mathbb{E} \exp \left( t \left( cX - x \left( \frac{X^p}{p} + \frac{c^q}{q} \right) \right) \right) \right) \quad (13)$$

for any  $x > \frac{E(X)}{(E(X^p))^{1/p}} = \Gamma(p+1)^{-\frac{1}{p}}$ . In order to apply Bahadur results, it remains to study the continuity of  $I$  on its domain. This is the object of the following lemma:

**Lemma 3.1.** *The rate function  $I$  is continuous on  $(\Gamma(p+1)^{-\frac{1}{p}}, 1)$ .*

*Proof of Lemma 3.1.* Set for any  $(t, c, x) \in [0, \infty) \times (0, \infty) \times (T_0, +\infty)$

$$H(t, c, x) = \mathbb{E} \exp \left( t \left( cX - x \left( \frac{X^p}{p} + \frac{c^q}{q} \right) \right) \right)$$

Denote by

$$\Phi(u) = cu - x \left( \frac{u^p}{p} + \frac{c^q}{q} \right)$$

The function  $\Phi$  is convex on  $\mathbb{R}^+$  and reached its maximum at  $u_0 = (\frac{c}{x})^{1/p}$ . Moreover  $\Phi(u_0) > 0$  if and only if  $x < 1$ . Since  $X$  has full support on  $\mathbb{R}^+$ ,

$$P \left( cX - x \left( \frac{X^p}{p} + \frac{c^q}{q} \right) > 0 \right) > 0$$

From Lemma 1 and Lemma 3 of [2], we know that for any  $(c, x)$ ,  $\inf_{t \geq 0} H(t, c, x)$  is reached at some unique and finite  $t = t(c, x) > 0$ . The function  $t(c, x)$  is given by the equation

$$\frac{\partial}{\partial t} H(t, c, x) = 0$$

more precisely,  $\frac{\partial}{\partial t} H$  being  $\mathcal{C}^1$  on its domain, by the Implicit Function Theorem  $t$  is  $\mathcal{C}^1$  at each point  $(c, x)$ . Thus the same can be said of

$$\underline{H}(c, x) = \inf_{t \geq 0} H(t, c, x) = H(t(c, x), c, x)$$

Since  $H \rightarrow 0$  when  $c \rightarrow \infty$ , we also know that  $\sup_{c>0} H$  is reached, therefore  $\sup_{c>0} \underline{H}(c, x)$  is reached for some  $c = c(x)$  and this function is continuous on  $(\Gamma(p+1)^{-\frac{1}{p}}, 1)$ . So is  $I(x) = -\log(\underline{H}(c(x), x))$ .  $\square$

Suppose that we want to test  $H_0$  against the alternative that the sample has  $\Gamma(\theta, \beta)$  distribution for  $\theta > 1$  and  $\beta > 0$ . Under the alternative,

$$T_n \rightarrow T(\theta, \beta) = \theta \left( \frac{\Gamma(\theta)}{\Gamma(\theta + p)} \right)^{1/p}$$

For  $\theta > 1$ ,  $T(\theta, \beta) > \Gamma(p+1)^{-\frac{1}{p}}$  and the Bahadur exact slope is therefore

$$c(\theta) = -\log \left( \sup_{c>0} \inf_{t \geq 0} \mathbb{E} \exp \left( t \left( cX - \theta \left( \frac{\Gamma(\theta)}{\Gamma(\theta + p)} \right)^{1/p} \left( \frac{X^p}{p} + \frac{c^q}{q} \right) \right) \right) \right)$$

## APPENDIX A. TECHNICAL LEMMATA

*Proof of Lemma 2.1.* Take any  $p > 1$  such that  $\mu(p) < \infty$ . Then  $\phi(z) \geq \mu(p)^{-p} z^p$  and in particular  $Kz - x\phi(z) \rightarrow -\infty$  when  $z \rightarrow \infty$ , so  $\mathbb{E} \exp(KX - x\phi(X)) < \infty$ . Observe also that for any  $z > 0$ ,  $Kz - xC\phi(z) \rightarrow -\infty$  when  $C \rightarrow \infty$ . By Lebesgue's dominated convergence theorem, it follows that  $\mathbb{E} \exp(KX - xC\phi(X)) \rightarrow 0$  when  $C \rightarrow \infty$ .  $\square$

*Proof of Lemma 2.2.* It is easily checked that  $\xi_{j,k}$  satisfies the same conditions as  $\xi_j$  in [14], in particular, thanks to the hypothesis on  $x$ ,  $-\infty \leq \mathbb{E}\xi_{j,l} < 0$ . Therefore we know that there exists  $t_{j,l} \geq 0$  such that

$$\begin{aligned} 1 &\geq \inf_{t \geq 0} \mathbb{E} \exp(t\xi_{j,l}) = \mathbb{E} \exp(t_{j,l}\xi_{j,l}) \\ &\geq \mathbb{E} \exp \left( t_{j,l} \left( \delta Y_{j,l} - x \frac{\mu(p_l)}{p_l} \left( \Phi(X_{j,l}) + (p_l - 1)(j\Delta)^{\frac{p_l}{p_l-1}} \right) \right) \right) \end{aligned}$$

choosing  $B$  large enough so  $\mathbb{P}(X \geq B) \geq \frac{1}{e}$ ,

$$\geq \exp \left( \frac{(t_{j,l}\delta)^2}{2} - 1 - t_{j,l}xC \left( \phi(B) + (\pi_1 - 1)K^{\frac{\pi_0}{\pi_0-1}} \right) \right) =: \exp(R(t_{j,l}))$$

It follows that  $t_{j,l}$  has to be smaller than  $t^+$ , the largest root of  $R$ , so the lemma is proved with  $\Delta := \frac{1}{t^+ K^{\frac{\pi_0}{\pi_0-1} + 1}}$ .  $\square$

*Proof of Lemma 2.3.* We have

$$\begin{aligned} D(j, l) &= \underbrace{(p_l - 1)(j\Delta)^{\frac{p_l}{p_l-1}} - (p_l - 1)((j-1)\Delta)^{\frac{p_l}{p_l-1}}}_{D_1} \\ &\quad + \underbrace{(p_l - 1)((j-1)\Delta)^{\frac{p_l}{p_l-1}} - \inf_{p \in T_l} (p-1)((j-1)\Delta)^q}_{D_2} \end{aligned}$$

with

$$D_1 \leq (p_l - 1) \sup_{c \in [j\Delta, (j-1)\Delta]} \frac{\partial}{\partial c} c^{\frac{p_l}{p_l-1}} \Delta \leq \pi_1 K^{\frac{1}{\pi_0-1}} \Delta$$

and

$$D_2 \leq \sup_{p \in T_l} \left| \frac{\partial}{\partial p} (p-1)((j-1)\Delta)^q \right| \Delta \leq K^{\frac{\pi_0}{\pi_0-1}} \log(K) \left( 1 + \frac{1}{(\pi_0 - 1)^2} \right) \Delta$$

$\square$

## REFERENCES

- [1] BAHADUR, R.R. Some limit theorems in statistics CBMS-NSF Regional Conference Series in Applied Mathematics, 1971 Philadelphia, Pa.: SIAM, Society for Industrial and Applied Mathematics
- [2] CHERNOFF, H. A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. *Ann. Math. Statist.* 23, 493–507 (1952).
- [3] CUVELIER, E., AND NOIRHOMME-FRAITURE, M. An approach to stochastic process using quasi-arithmetic means. In *Recent advances in stochastic modeling and data analysis*. World Scientific, 2007, pp. 2–9.
- [4] CUVELIER, E., AND NOIRHOMME-FRAITURE, M. Parametric families of probability distributions for functional data using quasi-arithmetic means with Archimedean generators. In *Functional and operatorial statistics*, Contrib. Statist. Springer, 2008, pp. 127–133.
- [5] DE LA PEÑA, V. H., LAI, T. L., AND SHAO, Q.-M. *Self-normalized processes*. Probability and its Applications (New York). Springer-Verlag, Berlin, 2009. Limit theory and statistical applications.
- [6] DEMBO, A., AND SHAO, Q.-M. Self-normalized large deviations in vector spaces. In *High dimensional probability (Oberwolfach, 1996)*, vol. 43 of *Progr. Probab.* Birkhäuser, Basel, 1998, pp. 27–32.
- [7] DEMBO, A., AND SHAO, Q.-M. Large and moderate deviations for Hotelling’s  $T^2$ -statistic. *Electron. Comm. Probab.* 11 (2006), 149–159 (electronic).
- [8] HARDY, G. H., LITTLEWOOD, J. E., AND PÓLYA, G. *Inequalities*. Cambridge, at the University Press, 1952. 2d ed.
- [9] KOLMOGOROFF, A. Sur la notion de la moyenne. *Rendiconti Accad. d. L. Roma (6)* 12 (1930), 388–391.
- [10] LAI, T. L., AND SHAO, Q.-M. Self-normalized limit theorems in probability and statistics. In *Asymptotic theory in probability and statistics with applications*, vol. 2 of *Adv. Lect. Math. (ALM)*. Int. Press, Somerville, MA, 2008, pp. 3–43.
- [11] NIKITIN, Y. Asymptotic efficiency of non parametric tests. Cambridge, at the University Press, 1995.
- [12] NAGUMO, M. Über eine Klasse der Mittelwerte. *Japanese Journ. of Math.* 7 (1930), 71–79.
- [13] PORCU, E., MATEU, J., AND CHRISTAKOS, G. Quasi-arithmetic means of covariance functions with potential applications to space-time data. *Journal of Multivariate Analysis* 100, 8 (2009), 1830 – 1844.
- [14] SHAO, Q.-M. Self-normalized large deviations. *Ann. Probab.* 25, 1 (1997), 285–328.
- [15] SHAO, Q.-M. A note on the self-normalized large deviation. *Chinese J. Appl. Probab. Statist.* 22, 4 (2006), 358–362.
- [16] TCHIRINA, A. V. Large deviations for a class of scale-free statistics under the gamma distribution. *J. Math. Sciences* 128, 1 (2005), 2640–2655.
- [17] YAGER, R. R. On ordered weighted averaging aggregation operators in multicriteria decisionmaking. *IEEE Trans. Systems Man Cybernet.* 18, 1 (1988), 183–190.