



HAL
open science

Lyapunov spectrum of a relativistic stochastic flow in the Poincaré group.

Camille Tardif

► **To cite this version:**

Camille Tardif. Lyapunov spectrum of a relativistic stochastic flow in the Poincaré group.. 2013.
hal-00798519

HAL Id: hal-00798519

<https://hal.science/hal-00798519>

Preprint submitted on 8 Mar 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lyapunov spectrum of a relativistic stochastic flow in the Poincaré group.

Camille Tardif *

March 8, 2013

Abstract

We determine the Lyapunov spectrum and stable manifolds of some stochastic flows on the Poincaré group associated to Dudley's relativistic processes.

Key words: Relativistic processes. Lévy processes in Lie groups. Poincaré group. Lyapunov spectrum. Hyperbolic dynamics.

AMS Subject Classification: 37H15, 60G51, 83A05

Contents

1	Introduction	1
2	Dudley processes and their lift in the Poincaré group	3
2.1	Minkowski space-time and Poincaré group	3
2.2	Dudley processes	4
3	Asymptotic random variables	6
3.1	Iwasawa decomposition in G	7
3.2	Asymptotic behavior of the G -component	8
3.3	Asymptotic behavior of the $\mathbb{R}^{1,d}$ -component	13
3.4	Geometric description of the convergence	14
4	Lyapunov spectrum and stable manifolds	15
4.1	Lyapunov spectrum	15
4.2	Stable manifolds	17
4.3	Projection on $\mathbb{H}^d \times \mathbb{R}^{1,d}$ of the stable manifolds	21

1 Introduction

In 1966 Dudley [6] defined a class of relativistic processes with Lorentzian-covariant dynamics in the framework of special relativity. Such a process ξ_t with values in Minkowski space-time $\mathbb{R}^{1,d}$, is differentiable and has velocity smaller than the speed of light. So it can be parametrized by its proper time, which amounts to impose to the velocity $\dot{\xi}_t$ to be an element of the unit pseudo sphere \mathbb{H}^d of $\mathbb{R}^{1,d}$. The restriction to the tangent space of \mathbb{H}^d of Minkowski ambient pseudo-metric turns \mathbb{H}^d into a Riemannian manifold of constant negative curvature. The invariance of the process $(\xi_t, \dot{\xi}_t)$ by the natural action of the set of Lorentz transforms on $\mathbb{H}^d \times \mathbb{R}^{1,d}$ imposes to the laws of ξ_t to be invariant by the action of the isometries of \mathbb{H}^d . Among this class of

*camille.tardif@uni.lu Université du Luxembourg.

relativistic processes, there is essentially only one which is continuous. It corresponds to the case where $\dot{\xi}_t$ is a Riemannian Brownian motion in the hyperbolic space and in this case $(\dot{\xi}_t, \xi_t)$ is called *Dudley diffusion*. Forty years after this seminal work, Franchi and Le Jan [8] extended Dudley diffusion to the framework of any Lorentz manifold. They defined relativistic processes with Lorentzian-covariant dynamics on generic Lorentzian manifolds by rolling without slipping a Dudley diffusion on the unit tangent space. They studied the asymptotic behavior of such diffusion in the Schwarzschild space-time. Bailleul [3] succeeded to compute the Poisson boundary of Dudley diffusion in Minkowski space-time and showed that it coincides with the causal boundary of $\mathbb{R}^{1,d}$. The asymptotic behavior of relativistic diffusions was investigated in other non flat Lorentzian manifolds ([1], [7]) with the aim of describing how the asymptotic behavior of the diffusion reflects the asymptotic geometry of the manifold.

In this work we ask a new question concerning these processes dealing with the asymptotic behaviour of some stochastic flow associated to it. As Brownian motion on a Riemannian manifold, the relativistic diffusion [8] is obtained by projecting a diffusion process with values in the orthonormal frame bundle, solution of a stochastic differential equation. This SDE generates a stochastic flow which, in our Lorentzian framework, consists in a stochastic perturbation of the geodesic flow. Existence and computation, for example, of the Lyapunov spectrum and stable manifolds of these flows may be investigated in the same way as it was done by Carverhill and Elworthy [5] for the canonical stochastic flow in the Riemannian framework. The main difficulty to study the flow of relativistic processes comes from the fact that the orthonormal frame bundle of a Lorentz manifold is never compact. Nevertheless in this article we provide a study of the asymptotic dynamics of the stochastic flow generated by Dudley processes in the Minkowski space-time (without restricting ourselves to the diffusion case). Precisely, in this framework, the orthonormal frame bundle is identified with the Poincaré group $\tilde{G} := PSO(1, d) \times \mathbb{R}^{1,d}$ and denoting ϕ_t the left invariant stochastic flow associated to one of Dudley's processes in \tilde{G} we obtain the description of the Lyapunov spectrum and the stable manifolds of φ_t . Precisely we obtain the following two results.

Theorem 1 (Lyapunov spectrum). *There exist a constant $\alpha > 0$ and two asymptotic random Lie sub-algebras $V_\infty^- \subset V_\infty^0$ of $\text{Lie}(\tilde{G})$ such that for some norm $\|\cdot\|$ on $\text{Lie}(\tilde{G})$ and $\tilde{X} \in \text{Lie}(\tilde{G})$ we have for almost every trajectory*

$$\frac{1}{t} \log \|d\varphi_t(\text{Id})(\tilde{X})\|_{\varphi_t(\text{Id})} \xrightarrow{t \rightarrow +\infty} \begin{cases} \alpha & \text{if } \tilde{X} \in \text{Lie}(\tilde{G}) \setminus V_\infty^0 \\ 0 & \text{if } \tilde{X} \in V_\infty^0 \setminus V_\infty^- \\ -\alpha & \text{if } \tilde{X} \in V_\infty^- \setminus \{0\} \end{cases}$$

Theorem 2 (Stable manifolds). *Denote by $\mathcal{V}_\infty^- := \exp(V_\infty^-)$ and d the distance associated to a left invariant and $\text{Ad}(SO(d))$ -invariant Riemannian metric in \tilde{G} . Then for any two distinct points \tilde{g}' and \tilde{g} in \tilde{G} we have*

- If $\tilde{g}' \in \tilde{g}\mathcal{V}_\infty^-$ then

$$\frac{1}{t} \log d(\varphi_t(\tilde{g}), \varphi_t(\tilde{g}')) \xrightarrow{t \rightarrow +\infty} -\alpha.$$

- If $\tilde{g}' \notin \tilde{g}\mathcal{V}_\infty^-$ then

$$\liminf_{t \rightarrow \infty} d(\varphi_t(\tilde{g}), \varphi_t(\tilde{g}')) > 0.$$

We begin by constructing, in section 2, Dudley processes as projections of left Lévy processes on the Poincaré group \tilde{G} , identified with the orthonormal frame bundle of the Minkowski space-time. These Lévy processes are solutions of stochastic integral equations and induce a left invariant stochastic flow φ_t in \tilde{G} . In section 3 we find the asymptotic behavior of Dudley processes and exhibit the asymptotic random variables $(\theta_\infty, \lambda_\infty) \in \mathbb{S}^{d-1} \times \mathbb{R}_+^*$. Finally in section 4 we prove Theorems 1 and 2 and explicit the projection of the stable manifold in $\mathbb{H}^d \times \mathbb{R}^{1,d}$ by showing that it corresponds to a skew product of a horosphere by a line.

Note that stochastic flows generated by Lévy processes on semi-simple Lie groups were intensively studied by Liao ([12], [11], [13]). But his results cannot be used directly here since our Lévy processes lie in the Poincaré group which is not semi-simple. Moreover in our work we

suppose only that the Levy measure is integrable at infinity whereas Liao [13] request the entire integrability of it.

Our work is also strongly inspired by the work of Bailleul and Raugi [4] where the authors used Raugi's methods [14] to find the Poisson boundary of Dudley diffusion.

2 Dudley processes and their lift in the Poincaré group

We present in this section the geometrical framework of special relativity and define a natural class of relativistic Markov processes with Lorentzian-covariant dynamics introduced by Dudley in [6]. They are obtained by projecting left Lévy processes with values in the Poincaré group and are described by two parameters: a diffusion coefficient $\sigma \in \mathbb{R}$ and a jump intensity Lévy measure ν on \mathbb{R}_+^* .

2.1 Minkowski space-time and Poincaré group

The Minkowski space-time $\mathbb{R}^{1,d}$ is $\mathbb{R} \times \mathbb{R}^d$ endowed with the Lorentz quadratic form q defined by

$$\forall \xi = (\xi^0, \xi^1, \dots, \xi^d) \in \mathbb{R} \times \mathbb{R}^d, \quad q(\xi) = (\xi^0)^2 - (\xi^1)^2 - \dots - (\xi^d)^2.$$

We denote by $\vec{\xi} := (\xi^1, \dots, \xi^d)^t$ the space component of ξ .

Set

$$Q = \text{Diag}(1, -1, \dots, -1)$$

the matrix of q in the canonical basis (e_0, e_1, \dots, e_d) . Time orientation is given by the constant vector field e_0 and some $\xi \in \mathbb{R}^{1,d}$ is said to be future oriented when $q(\xi, e_0) > 0$. A path γ_s in $\mathbb{R}^{1,d}$ is said to be *time-like* when it is differentiable almost everywhere and $q(\dot{\gamma}_s) > 0$ and $q(\dot{\gamma}_s, e_0) > 0$. The Poincaré group is the group of affine q -isometries which preserve orientation and time-orientation. It is the semi-direct product connected group

$$\tilde{G} := PSO(1, d) \ltimes \mathbb{R}^{1,d}$$

where $G := PSO(1, d)$ denotes the group of linear q -isometries which preserve orientation and time-orientation. An element $\tilde{g} = (g, \xi) \in \tilde{G}$ is made up of its linear part $g \in G$ and its translation part ξ . We identify G with the sub-group of \tilde{G} which fixes 0. By this way, we identify $\mathbb{R}^{1,d}$ with the homogeneous space \tilde{G}/G . The identity element of \tilde{G} and G is denoted by Id (thus for us $\text{Id} = (\text{Id}, 0)$). At $\tilde{g} = (g, \xi) \in \tilde{G}$ we associate the affine frame $((g(e_0), g(e_1), \dots, g(e_d)); \xi)$ of $\mathbb{R}^{1,d}$ and \tilde{G} is identified with the bundle of q -orthonormal, oriented and time-oriented, frames over $\mathbb{R}^{1,d}$. We denote by

$$\begin{aligned} \tilde{\pi} : \quad \tilde{G} &\longrightarrow \mathbb{R}^{1,d} \\ \tilde{g} = (g, \xi) &\longmapsto \xi \end{aligned}$$

the projection associated to this trivial fibration and $G = \tilde{\pi}^{-1}\{0\}$. The canonical basis being fixed we identify G with the matrix group

$$G = \{g \in \text{SL}(\mathbb{R}^{d+1}), gQg^t = Q, q(g(e_0), e_0) > 0\},$$

and its Lie algebra is

$$\begin{aligned} \text{Lie}(G) &= \{X \in \mathcal{M}_{d+1}(\mathbb{R}), XQ - QX^t = 0\} \\ &= \left\{ \begin{pmatrix} 0 & b^t \\ b & C \end{pmatrix}, b \in \mathbb{R}^d, C \in \mathcal{M}_d(\mathbb{R}) \text{ s.t } C = -C^t \right\}. \end{aligned}$$

We have $\text{Lie}(\tilde{G}) = \text{Lie}(G) \times \mathbb{R}^{1,d}$ and for $\tilde{X} = (X, x), \tilde{Y} = (Y, y) \in \text{Lie}(\tilde{G})$

$$[\tilde{X}, \tilde{Y}] = ([X, Y], Xy - Yx).$$

We identify $\text{Lie}(G)$ with $\text{Ker}(d_{\text{Id}}\tilde{\pi})$ and its elements are vertical for the fibration $\tilde{\pi}$. We set

$$\begin{aligned} V_i &:= e_0 e_i^t + e_i e_0^t \quad i = 1, \dots, d \\ V_{ij} &:= [V_i, V_j] = e_i e_j^t - e_j e_i^t \quad j > i. \end{aligned}$$

Moreover we set

$$H_0 := (0, e_0) \in \text{Lie}(\tilde{G})$$

which is horizontal for the fibration $\tilde{\pi}$.

Notation For $\tilde{X} \in \text{Lie}(\tilde{G})$ we denote by \tilde{X}^t the left invariant vector field in \tilde{G} associated. Denote by K the subgroup of G made of the rotations of \mathbb{R}^d . We have

$$K := \left\{ \begin{pmatrix} 1 & 0 \\ 0 & k \end{pmatrix}, k \in SO(d) \right\},$$

and K is also the stabilizer of e_0 under the action of G on $\mathbb{R}^{1,d}$. The homogeneous space G/K can be identified with the orbit of e_0 under the action of G which is the unit pseudo sphere $\mathbb{H}^d := \{\xi \in \mathbb{R}^{1,d}, q(\xi) = 1, \xi^0 > 0\}$ and is a Riemannian manifold of constant negative curvature when its tangent space is endowed with the restriction of q on it.

For $r \in \mathbb{R}^+$ and $\theta \in \mathbb{S}^{d-1} \subset \mathbb{R}^d$, define

$$S(r, \theta) := \exp \left(r \sum_{i=1}^d \theta^i V_i \right) = \begin{pmatrix} \cosh(r) & \sinh(r)\theta^t \\ \sinh(r)\theta & \text{Id} + (\cosh(r) - 1)\theta\theta^t \end{pmatrix}.$$

Each $g \in G$ can be decomposed in *polar form* $g = S(r(g), \theta(g))R$ where $R \in K$.

2.2 Dudley processes

In this paragraph we define the relativistic processes introduced by Dudley in [6]. These processes enjoy two natural properties:

- they are \tilde{G} -invariant i.e their dynamics are invariant by a change of q -orthonormal frame
- their trajectories in $\mathbb{R}^{1,d}$ are time-like: they are almost everywhere differentiable, and the tangents vectors are time-like and time oriented.

First remark that no Markov processes with values in $\mathbb{R}^{1,d}$ is \tilde{G} -invariant. Indeed, the law at some time $t > 0$ of such process starting at 0 would be a G -invariant probability measure in $\mathbb{R}^{1,d}$ which is necessary trivial by the following lemma.

Lemma 1. *The only G -invariant probability measure in $\mathbb{R}^{1,d}$ is the Dirac measure at 0.*

Proof. Let μ be a G -invariant probability measure in $\mathbb{R}^{1,d}$. First suppose that the support of μ is not contained in the q -orthogonal hyperplane of some light-like line $\{u(e_0 + \hat{\theta}), u \in \mathbb{R}\}$ ($\hat{\theta} := \sum_{i=1}^d \hat{\theta}^i e_i \in \mathbb{S}^{d-1}$). So there exist a compact set C in the complement of this hyperplane such that $\mu(C) > 0$. For $g = S(r, \hat{\theta})$, $r > 0$ and $\xi = (\xi^0, \vec{\xi}) \in \mathbb{R}^{1,d}$, denoting $\|\cdot\|$ the Euclidean norm in $\mathbb{R}^{1,d}$ we have

$$\begin{aligned} \|g(\xi)\|^2 &= 2q(g(\xi), e_0)^2 - q(g(\xi))^2 = 2q(\xi, g^{-1}(e_0))^2 - q(\xi)^2 = 2 \left(\cosh(r)\xi^0 - \sinh(r)\hat{\theta} \cdot \vec{\xi} \right)^2 - q(\xi)^2 \\ &= 2 \left(\cosh(r)(\xi^0 - \hat{\theta} \cdot \vec{\xi}) + e^{-r}\hat{\theta} \cdot \vec{\xi} \right)^2 - q(\xi)^2 \\ &= 2 \left(\cosh(r)q(\xi, e_0 + \hat{\theta}) + e^{-r}\hat{\theta} \cdot \vec{\xi} \right)^2 - q(\xi)^2. \end{aligned}$$

Since C is a compact set in the complement of the q -orthogonal hyperplane to $\{u(e_0 + \hat{\theta}), u \in \mathbb{R}\}$ then $\inf_{\xi \in C} |q(\xi, e_0 + \hat{\theta})| > 0$ and thus r can be chosen such that $\inf_{\xi \in C} \|g(\xi)\|$ is arbitrary large. Thus g can be chosen such that C and $g(C)$ are disjoint. Furthermore $q(g(\xi), e_0 + \hat{\theta}) = q(\xi, g^{-1}(e_0 + \hat{\theta})) = e^r q(\xi, e_0 + \hat{\theta})$ and thus $g(C)$ belongs to the complement of the hyperplane q -orthogonal to $e_0 + \hat{\theta}$. By iteration we can find a sequence $g_k = S(r_k, \hat{\theta})$, $k \in \mathbb{N}$ such that the compact sets $g_k(C)$

are pairwise disjoint. Thus we obtain a contradiction writing $1 \geq \mu(\cup_k g_k(C)) = \sum_k \mu(g_k(C)) = \sum_k \mu(C) = +\infty$.

Now if the support of μ is contained in some hyperplan tangent to the light-cone and is not restricted to 0, we can find a compact set C in this hyperplan with $0 \notin C$ and $\mu(C) > 0$ and we can choose a rotation $R \in K$ such that $R(C)$ is not in the hyperplan. Thus $\mu(C) = \mu(R(C)) = 0$ and it gives a contradiction. So we proved that μ is necessary the Dirac measure in $\{0\}$. \square

Thus $\mathbb{R}^{1,d} = \tilde{G}/G$ cannot be the space of states of some non-trivial \tilde{G} -invariant Markov process. But \tilde{G} -homogeneous spaces of the form \tilde{G}/\tilde{K} where \tilde{K} is a compact subgroup of \tilde{G} have some \tilde{K} -invariant probability measure and may be endowed with some \tilde{G} -invariant Markov processes (see [13] or [9]). The smallest spaces of states we can consider correspond to the maximal compact sub-group of \tilde{G} . Thus it is natural to consider the space of states $\tilde{G}/K \simeq \mathbb{H}^d \times \mathbb{R}^{1,d}$. The group K is seen as the subgroup of \tilde{G} which stabilize 0 and e_0 under the action of \tilde{G} on $\mathbb{R}^{1,d}$.

We denote by $\pi : \tilde{G} \mapsto \mathbb{H}^d \times \mathbb{R}^{1,d} \simeq \tilde{G}/K$ the canonical projection

$$\forall \tilde{g} = (g, \xi) \in \tilde{G} \quad \pi(\tilde{g}) = (g(e_0), \xi).$$

The following Proposition exhibit all the relativistic processes in $\mathbb{H}^d \times \mathbb{R}^{1,d}$. It is essentially an application of a result of Liao (Theorem 2.1 and 2.2 p 42 in [13]).

Proposition 1. *The Markov processes on $\mathbb{H}^d \times \mathbb{R}^{1,d}$, starting at (ζ_0, ξ_0) , which are \tilde{G} -invariant and whose trajectories are time-like are of the form (ζ_s, ξ_s) where ζ_s is a G -invariant Markov process on \mathbb{H}^d and $\xi_t = \xi_0 + a \int_0^t \zeta_s ds$; a being some positive constant. For such a process there exist $\sigma > 0$ and a measure ν on \mathbb{R}^+ satisfying*

$$\int_0^{+\infty} \min(1, r^2) \nu(dr) < +\infty,$$

such that $(\zeta_t, \xi_t) = \pi(\tilde{g}_t)$ in law where \tilde{g}_t is a left Levy process on \tilde{G} starting at \tilde{g}_0 s.t $(\zeta_0, \xi_0) = \pi(\tilde{g}_0)$ of which generator $\tilde{\mathcal{L}}$ is defined by

$$\begin{aligned} \forall f \in C^2(\tilde{G}) \quad \tilde{\mathcal{L}}f(\tilde{g}) &= aH_0^l f(\tilde{g}) + \frac{\sigma^2}{2} \sum_{i=1}^d (V_i^l)^2 f(\tilde{g}) \\ &+ \int_0^{+\infty} \int_{\mathbb{S}^d} \left(f(\tilde{g}S(r, \theta)) - f(\tilde{g}) - r \mathbf{1}_{r \in [0,1]} \sum_{i=1}^d \theta^i V_i^l f(\tilde{g}) \right) \nu(dr) d\theta. \end{aligned}$$

Definition 1 (Dudley processes). When $a = 1$ then $\dot{\xi}_t = \zeta_t$ and ξ_t is parametrized by its *proper time*, i.e $q(\dot{\xi}_t) = 1$. When moreover σ or ν is non trival we call (ξ_t, ζ_t) a Dudley process and we consider exclusively these processes in the sequel. When $\nu = 0$ (ξ_t, ζ_t) is continuous and is called *Dudley diffusion*.

Remark 1. *The process ξ_t is differentiable and $\dot{\xi}_t$ is càdlàg.*

Proof of Proposition 1. Let (ζ_t, ξ_t) a Markov (Feller) process on $\mathbb{H}^d \times \mathbb{R}^{1,d}$ starting at (ζ_0, ξ_0) , which is \tilde{G} -invariant and whose trajectories are time-like. By choosing \tilde{g}_0 such that $(\zeta_0, \xi_0) = \pi(\tilde{g}_0)$ and considering the Markov process $\tilde{g}_0^{-1}(\zeta_t, \xi_t)$ it remains to prove the proposition in the case where $(\zeta_0, \xi_0) = (e_0, 0)$.

Set

$$H_i := (0, e_i) \in \text{Lie}(\tilde{G}), \quad i = 1, \dots, d.$$

The family $\{H_0, H_i, V_i, V_{ij}\}_{i < j \in \{1, \dots, d\}}$ form an orthonormal basis of $\text{Lie}(\tilde{G})$ for an $Ad(K)$ -invariant inner product on $\text{Lie}(\tilde{G})$ and $\text{Ker } d_{\text{Id}}\pi = \{V_{ij}\}_{i < j \in \{1, \dots, d\}}$.

We set

$$X_0 := H_0 \quad X_i := H_i \quad X_{d+i} := V_i \quad i = 1, \dots, d.$$

An $\tilde{h} = (h, \xi) \in \tilde{G}$ can be decomposed in $\tilde{h} = \exp\left(\sum_{i=0}^d x^i(\tilde{h})X_i\right) \exp\left(\sum_{i=d+1}^{2d} x^i(\tilde{h})X_i\right) R$ where $R \in K$, $x^0(\tilde{h}) = \xi^0$ and for $i = 1, \dots, d$ $x^i(\tilde{h}) = \xi^i$ and $x^{d+i}(\tilde{h}) = r(h)\theta^i(h)$. By Theorem 2.1 and

2.2 p 42 of [13], (ζ_t, ξ_t) coincide in law with $\pi(\tilde{g}_t)$ where \tilde{g}_t is a left Levy process in \tilde{G} , starting at Id, which is K -right invariant and generated by

$$\tilde{\mathcal{L}}f(\tilde{g}) = \frac{1}{2} \sum_{i,j=0}^{2d} a_{ij} X_i^l X_j^l f(\tilde{g}) + \sum_{i=0}^{2d} b^i X_i^l f(\tilde{g}) + \int_{\tilde{G}} \left(f(\tilde{g}\tilde{h}) - f(\tilde{g}) - \sum_{i=0}^{2d} \mathbf{1}_{r(h) \leq 1} x^i(\tilde{h}) X_i f(\tilde{g}) \right) \tilde{\Pi}(d\tilde{h}).$$

The matrice $A := (a_{ij})$ is a positive symmetric, $(b^i)_i \in \mathbb{R}^{2d+1}$ and $\tilde{\Pi}$ is a Levy measure invariant by K -conjugation in \tilde{G} . The right K -invariance of \tilde{g}_t ensures that for all $k \in K \simeq SO(d)$, $\text{diag}(1, k, k) \text{Adiag}(1, k, k)^{-1} = A$. Thus, A is necessarily of the form

$$\begin{pmatrix} \hat{\sigma} & 0 & 0 \\ 0 & \tilde{\sigma} \text{Id} & \sigma' \text{Id} \\ 0 & \sigma' \text{Id} & \sigma \text{Id} \end{pmatrix},$$

where $\hat{\sigma}, \tilde{\sigma} \geq 0$ and $\tilde{\sigma} \geq (\sigma')^2$. Moreover, using again K -invariance it comes $b^i = 0$ for $i = 1, \dots, 2d$ and we set $a := b^0$. The trajectories of \tilde{g}_t projected in $\mathbb{R}^{1,d}$ need to be differentiable so the jump measure Π is supported on G and $\hat{\sigma}$ and $\tilde{\sigma}$ are necessarily null. Since the trajectories are time-oriented we have $a > 0$. The push forward of $\tilde{\Pi}$ by π is supported on $G/K \simeq \mathbb{H}^d$ and is K -invariant. Thus Π can be chosen of the form

$$\forall f \in C_0(G) \quad \Pi f = \int_0^{+\infty} \int_{\mathbb{S}^{d-1}} f(S(r, \theta)) \nu(dr) d\theta$$

where ν is a Levy measure on \mathbb{R}_+^* (i.e satisfying $\int \min(1, r^2) \nu(dr) < +\infty$). \square

Denote by g_t the G -component of \tilde{g}_t . Thus $g_t(e_0) = \dot{\xi}_t$ and $\xi_t = \int_0^t g_s(e_0) ds$. By definition g_t is a G -valued left Levy process, K -right invariant, generated by \mathcal{L} defined by

$$\forall f \in C^2(G) \quad \mathcal{L}f(g) = \frac{\sigma^2}{2} \sum_{i=1}^d (V_i^l)^2 f(g) + \int_0^{+\infty} \int_{\mathbb{S}^d} \left(f(\tilde{g}S(r, \theta)) - f(\tilde{g}) - r \mathbf{1}_{r \in [0,1]} \sum_{i=1}^d \theta^i V_i^l f(\tilde{g}) \right) \nu(dr) d\theta.$$

Denote by Π the Levy measure supported on G defined by

$$\Pi f = \int_0^{+\infty} \int_{\mathbb{S}^{d-1}} f(S(r, \theta)) \nu(dr) d\theta.$$

Define $U_0 := \{g \in G, r(g) \leq 1\}$ which is a K invariant neighborhood of Id in G . For $f \in C^2(G)$ we have the following Itô formula (see [2]) for g_t

$$\begin{aligned} f(g_t) &= f(\text{Id}) + \sigma \sum_{i=1}^d \int_0^t V_i^l f(g_{s-}) dB_s^i + \frac{\sigma^2}{2} \int_0^t \sum_{i=1}^d (V_i^l)^2 f(g_{s-}) ds + \int_0^t \int_{U_0} (f(g_{s-h}) - f(g_{s-})) \tilde{N}(ds, dh) \\ &+ \int_0^t \int_{U_0} \left(f(g_{s-h}) - f(g_{s-}) - r(h) \sum_{i=1}^d \theta^i(h) V_i^l f(g_{s-}) \right) ds \Pi(dh) \\ &+ \int_0^t \int_{(U_0)^c} (f(g_{s-h}) - f(g_{s-})) N(ds, dh), \end{aligned} \quad (1)$$

where B_t is a Brownian motion of \mathbb{R}^d , N is a Poisson random measure on $\mathbb{R} \times G$ of intensity measure $dt \otimes \Pi$ and $\tilde{N}(ds, dh) := N(ds, dh) - ds \Pi(dh)$ is the compensated random measure associated.

3 Asymptotic random variables

In this section we determine the asymptotic behavior of $\pi(\tilde{g}_t) = (g_t(e_0), \xi_t)$ under an integrability condition on the jump intensity measure ν (Ass.1). Writing $g_t = n_t a_t k_t$ in some Iwasawa decomposition of G we first prove (Prop.2), applying the Itô formula (1) and the law of large number, that the abelian term $a_t = \exp(\alpha_t V_1)$ is *positively contracting*, $\frac{\alpha_t}{t}$ converges almost surely to a

positive constant α depending explicitly on σ and ν . Next we prove (Prop.4) that the nilpotent term n_t converges almost surely to an asymptotic random variable n_∞ and this convergence is exponentially fast with rate α . Then we investigate (Prop.5) the asymptotic behavior of ξ_t in $\mathbb{R}^{1,d}$. Geometrically, seen in the projective space, the \mathbb{H}^d -valued process $g_t(e_0)$ converges to a limit angle $\theta_\infty \in \partial\mathbb{H}^d \simeq \mathbb{S}^{d-1}$ of which n_∞ is a stereographic projection. Moreover, the process ξ_t is asymptotic to some affine hyperplane q -orthogonal to θ_∞ of which position is fixed by another asymptotic random variable $\lambda_\infty \in \mathbb{R}_+^*$. Figure 1 sum up the asymptotic results.




Figure 1: Asymptotic behavior of a Dudley diffusion

3.1 Iwasawa decomposition in G

Although a polar decomposition of G was used to introduce g_t (defining the K invariant measure Π), Iwasawa decomposition seems to be more adapted to describe its asymptotic dynamics. Introduce briefly this decomposition.

The maximal abelian subalgebra contained in $\text{Vect}\{V_1, \dots, V_d\}$, which is the orthogonal subspace of $\text{Lie}(G)$ of \mathcal{K} for the Killing form, is of dimension one. Let choose $\mathcal{A} := \text{Vect}\{V_1\}$ one of them. The linear endomorphism $\text{ad}(V_1)$ of $\text{Lie}(G)$ is diagonalisable with eigenvalues $-1, 0$ and 1 . Set

$$\mathcal{N} = \{X \in \text{Lie}(G), \text{ad}(V_1)X = -X\} \quad \overline{\mathcal{N}} = \{X \in \text{Lie}(G), \text{ad}(V_1)X = X\}$$

the eigenspace corresponding respectively to the eigenvalue -1 and 1 . Explicitly

$$\mathcal{N} = \text{Vect}\{V_i - V_{1i}, \quad i = 2, \dots, d\} \quad \text{and} \quad \overline{\mathcal{N}} = \text{Vect}\{V_i + V_{1i}, \quad i = 2, \dots, d\}.$$

The eigenspace corresponding to 0 is $\mathcal{A} \oplus \mathcal{M}$ where \mathcal{M} is the sub algebra of elements of \mathcal{K} which commute with the elements of \mathcal{A} . Explicitly

$$\mathcal{M} = \text{Vect}\{V_{ij}, \quad i, j = 2 \dots d\} = \left\{ \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & C \end{pmatrix}, \quad C \in \text{so}(d-1) \right\}.$$

The subspace \mathcal{N} is a nilpotent Lie algebra (even abelian since $[\mathcal{N}, \mathcal{N}] = 0$). The corresponding Iwasawa decomposition of $\text{Lie}(G)$ is

$$\text{Lie}(G) = \mathcal{N} \oplus \mathcal{A} \oplus \mathcal{K}.$$

For $X \in \text{Lie}(G)$ we denote by $\{X\}_{\mathcal{N}}$ (resp. $\{X\}_{\mathcal{A}}$ and $\{X\}_{\mathcal{K}}$) its projection in \mathcal{N} (resp. \mathcal{A} and \mathcal{K}) thus $X = \{X\}_{\mathcal{N}} + \{X\}_{\mathcal{A}} + \{X\}_{\mathcal{K}}$. Denoting by $A := \exp(\mathcal{A})$, $N := \exp(\mathcal{N})$ the subgroup corresponding we obtain the corresponding Iwasawa decompositions of G

$$G = NAK.$$

Moreover, the mapping from $N \times A \times K$ to G which maps (n, a, k) to nak is an analytic diffeomorphism. For $g \in G$ we denote by $g = (g)_N(g)_A(g)_K$ its decomposition in Iwasawa coordinates. To simplify notations set $n_t := (g_t)_N$, $a_t := (g_t)_A$ and $k_t = (g_t)_K$, thus $g_t = n_t a_t k_t$.

Note that we have other Iwasawa decompositions like $\text{Lie}(G) = \overline{N} \oplus \mathcal{A} \oplus \mathcal{K}$ (with $G = \overline{N}AK$).

Iwasawa and polar coordinates.

For $g \in G$ written in polar form $g = \exp\left(r \sum_{i=1}^d \theta^i (V_i - V_{1i})\right) R$ where $R \in K$, we have $q(e_0, g(e_0)) = \cosh(r)$.

Now denoting by $b \in \mathbb{R}^{d-2}$ and $u \in \mathbb{R}$ such that $(g)_N = \exp\left(\sum_{i=2}^d b^i (V_i - V_{1i})\right)$ and $(g)_A = \exp(uV_1)$ we can compute explicitly $q(e_0, g(e_0))$ in terms of b and u and we obtain

$$\cosh(r) = \left(1 + \frac{\|b\|^2}{2}\right) \cosh(u) + \frac{\|b\|^2}{2} \sinh(u). \quad (2)$$

Moreover u can be expressed in term of θ and r via

$$e^u = \cosh(r) + \theta^1 \sinh(r) \quad (3)$$

3.2 Asymptotic behavior of the G -component

The aim of this section is to show that n_t converges almost surely to an asymptotic N -valued random variable n_∞ and that the convergence is exponentially fast. This result, stated in Proposition 4, appears to be a consequence of the contracting property of a_t . For this we need the following integrability condition on ν . The group G is semi simple and the tools used in this section are very closed from those of Liao [13]. Nevertheless we present a self-contained proof in our specific framework and our results are established under a weaker assumption than the ones of [13] (see remark 3). Namely we suppose that the following integrability condition is satisfied.

Assumption 1.

$$\int_1^{+\infty} r \nu(dr) < +\infty.$$

The following proposition computes explicitly the linear drift of a_t which appears to be positive. The proof is essentially a consequence of the law of large number.

Proposition 2. *Let denote by α_t the \mathbb{R} -valued process such that $a_t = \exp(\alpha_t V_1)$. Then the following convergence holds almost surely*

$$\frac{\alpha_t}{t} \xrightarrow{t \rightarrow \infty} \alpha > 0.$$

The positive constant α is

$$\alpha := \frac{d-1}{2} \sigma^2 + \int_0^{+\infty} \frac{r \cosh(r) - \sinh(r)}{\sinh(r)} \nu(dr).$$

Proof. First define $\log : A \rightarrow \mathcal{A} \simeq \mathbb{R}$ $\exp(uV_1) \mapsto uV_1$ and apply Itô formula (1) to the smooth map $f : g \mapsto \log(g)_A$. Remark that for $g, h \in G$ $(gh)_A = (g)_A ((g)_K h)_A$ and

$$f(g_s h) - f(g_{s-}) = \log(k_{s-} g)_A.$$

Moreover

$$\begin{aligned} V_i^l f(g_{s-}) &= \frac{d}{dt} \log(g_{s-} e^{tV_i})_A \Big|_{t=0} = \frac{d}{dt} \log(e^{tAd(k_{s-})V_i})_A \Big|_{t=0} = \{Ad(k_{s-})V_i\}_{\mathcal{A}} \\ (V_i^l)^2 f(g_{s-}) &= \frac{d}{dt} \{Ad(g_{s-} e^{tV_i})_K V_i\}_{\mathcal{A}} \Big|_{t=0} = \frac{d}{dt} \{Ad(e^{tAd(k_{s-})V_i})_K Ad(k_{s-})V_i\}_{\mathcal{A}} \Big|_{t=0} \\ &= \{[\{Ad(k_{s-})V_i\}_{\mathcal{K}}, Ad(k_{s-})V_i]\}_{\mathcal{A}}. \end{aligned}$$

For $k \in K$ we have $Ad(k)V_i = \sum_{j=1}^d k_{ij} V_j$ where $k = (k_{ij}) \in K \simeq SO(d)$. Then we compute

$$\sum_{i=1}^d \{[\{Ad(k_{s-})V_i\}_{\mathcal{K}}, Ad(k_{s-})V_i]\}_{\mathcal{A}} = \sum_{i=1}^d \sum_{j=1}^d \sum_{l=1}^d (k_{s-})_{ij} (k_{s-})_{il} \{[\{V_j\}_{\mathcal{K}}, V_l]\}_{\mathcal{A}},$$

since $\{V_j\}_{\mathcal{K}} = V_{1j}$ (and 0 for $j = 1$) and $[V_{1j}, V_l] = V_1 \delta_{jl}$ we get

$$\sum_{i=1}^d \{[\{Ad(k_{s-})V_i\}_{\mathcal{K}}, Ad(k_{s-})V_i]\}_{\mathcal{A}} = \sum_{j=2}^d \sum_{i=1}^d ((k_{s-})_{ij})^2 V_1 = (d-1)V_1.$$

Thus Itô formula (1) can be written

$$\begin{aligned} \log(a_t) &= M_t + \frac{d-1}{2} \sigma^2 t + \int_0^t \int_{U_0^c} \log(k_{s-}h)_A N(ds, dh) \\ &\quad + \int_0^t \int_{U_0} \left(\log(k_{s-}h)_A - r(h) \sum_{i=1}^d \theta^i(h) \{Ad(k_{s-})V_i\}_{\mathcal{A}} \right) ds \Pi(dh), \end{aligned}$$

where $M_t := \sigma \sum_{i=1}^d \int_0^t \{Ad(k_{s-})V_i\}_{\mathcal{A}} dB_s^i + \int_0^t \int_{U_0} (k_{s-}h)_A \tilde{N}(ds, dh)$ is a martingale. Its bracket is $\sigma^2 \int_0^t \sum_{i=1}^d \|\{Ad(k_{s-})V_i\}_{\mathcal{A}}\|^2 ds + \int_0^t \int_{U_0} \|\log(k_{s-}h)_A\|^2 ds \Pi(dh) = t \left(\sigma^2 + \int_{U_0} \|\log(h)_A\|^2 \Pi(dh) \right)$ and thus we obtain that almost surely

$$\frac{M_t}{t} \xrightarrow[t \rightarrow +\infty]{} 0.$$

Moreover, making the change of variable $h' = k_{s-} h k_{s-}^{-1}$ we obtain using the K -invariant by conjugation of Π

$$\begin{aligned} &\int_0^t \int_{U_0} \left(\log(k_{s-}h)_A - r(h) \sum_{i=1}^d \theta^i(h) \{Ad(k_{s-})V_i\}_{\mathcal{A}} \right) ds \Pi(dh) \\ &= \int_0^t \int_{U_0} \log(h')_A - r(h') \sum_{i=1}^d \theta^i(k_{s-}^{-1}h') \{Ad(k_{s-})V_i\}_{\mathcal{A}} \Pi(dh') ds \\ &= \int_0^t \int_{U_0} \log(h')_A - r(h') \sum_{i=1}^d \sum_{j=1}^d (k_{s-})_{ij} \theta^j(h') (k_{s-})_{i1} V_1 \Pi(dh') ds \\ &= \int_0^t \int_{U_0} \log(h')_A - r(h') \theta^1(h') V_1 \Pi(dh') ds \\ &= t \int_{\mathbb{S}^{d-1}} \int_0^1 \log(S(r, \theta))_A - r \theta^1 V_1 \nu(dr) d\theta \end{aligned}$$

By (3) we have $\log(S(r, \theta))_A = \log(\cosh(r) + \theta^1 \sinh(r)) V_1$ and the previous term equals

$$\begin{aligned} t \int_{\mathbb{S}^{d-1}} \int_0^1 \log(\cosh(r) + \theta^1 \sinh(r)) - r \theta^1 \nu(dr) d\theta V_1 &= t \int_0^1 \left(\int_{-1}^1 \log(\cosh(r) + u \sinh(r)) - ru \frac{du}{2} \right) \nu(dr) V_1 \\ &= t \int_0^1 \frac{r \cosh(r) - \sinh(r)}{\sinh(r)} \nu(dr) \end{aligned}$$

It remains to consider the asymptotic behavior of the stochastic integral $\int_0^t \int_{U_0^c} \log(k_s - h) {}_A N(ds, dh)$. We know that there exist T_i jump times of a Poisson process of intensity measure $\Pi(U_0^c)$ and random variable h_n i.i.d of law $\Pi|_{U_0^c}/\Pi(U_0^c)$ and independent of $(T_i)_i$ such that

$$\int_0^t \int_{U_0^c} \log(k_s - h) {}_A N(ds, dh) = \sum_{n, T_n \leq t} \log(k_{T_n^-} - h_n) {}_A.$$

Moreover, by invariance of Π under conjugation by K we check easily that the random variables $h'_n := Ad(k_{T_n^-})h_n$ are i.i.d of common law $\Pi|_{U_0^c}/\Pi(U_0^c)$ and we have

$$\int_0^t \int_{U_0^c} \log(k_s - h) {}_A N(ds, dh) = \sum_{n=1}^{N_t} \log(h'_n) {}_A,$$

where N_t is a Poisson process of intensity measure $\Pi(U_0^c)$ and independent of $(h'_n)_n$. Moreover since $\int_1^{+\infty} r\nu(dr) < +\infty$ then $\log(h'_n) {}_A$ is integrable and the law of large number ensures that

$$\frac{1}{t} \int_0^t \int_{U_0^c} \log(k_s - h) {}_A N(ds, dh) \xrightarrow{t \rightarrow +\infty} \Pi(U_0^c) \mathbb{E}[\log(h'_n) {}_A].$$

The proof is ended by checking that

$$\mathbb{E}[\log(h'_n) {}_A] = \frac{1}{\Pi(U_0^c)} \int_1^{+\infty} \frac{r \cosh(r) - \sinh(r)}{\sinh(r)} \nu(dr) V_1.$$

□

Remark 2. When $\int_1^{+\infty} r\nu(dr) = +\infty$ we obtain $\mathbb{E}[|\log(h'_n) {}_A|] = +\infty$. Nevertheless

$$\begin{aligned} \mathbb{E}[-\min(\log(h'_n) {}_A, 0)] &= \int_1^{+\infty} \int_{-1}^1 -\min(\log(\cosh(r + \theta^1 \sinh(r))), 0) \frac{d\theta^1}{2} \nu(dr) \\ &= \int_1^{+\infty} \int_{e^{-r}}^1 -\log(v) \frac{dv}{1 \sinh(r)} \nu(dr) \\ &= \int_1^{+\infty} \frac{1}{2 \sinh(r)} (1 - e^{-r}(r+1)) \nu(dr) < +\infty. \end{aligned}$$

Now, applying a generalized law of large numbers we deduce that almost surely

$$\frac{1}{t} \int_0^t \int_{U_0^c} \log(k_s - h) {}_A N(ds, dh) \xrightarrow{t \rightarrow +\infty} +\infty,$$

and so

$$\frac{\alpha_t}{t} \xrightarrow{t \rightarrow +\infty} +\infty.$$

The following proposition establishes that g_t is bounded in expectation on a finite time interval. This result is used to prove the convergence of n_t in the next Proposition.

Proposition 3. Fix $T > 0$. Then

$$\mathbb{E} \left[\sup_{t \in [0, T]} r(g_t) \right] < +\infty.$$

Proof. We cannot directly apply Itô formula to $g \mapsto r(g)$ since it is not regular at Id . But we can find a smooth function \tilde{r} such that $\tilde{r} \geq r$ on U_0 and $\tilde{r} = r$ on U_0^c . For such a function we have

$$\tilde{r}(g_t) = \tilde{r}(Id) + \hat{m}_t + \tilde{m}_t + I_t + J_t + \int_0^t \int_{U_0^c} (\tilde{r}(g_s - h) - \tilde{r}(g_{s-})) N(ds, dh). \quad (4)$$

Where $\hat{m}_t := \sigma \int_0^t V_i^l \tilde{r}(g_{s-}) dB_s^i$ and $\tilde{m}_t := \int_0^t \int_{U_0} (\tilde{r}(g_{s-h}) - \tilde{r}(g_{s-})) \tilde{N}(ds, dh)$ are martingales, $I_t := \sigma^2 \int_0^t \sum_{i=1}^d (V_i^l)^2 \tilde{r}(g_{s-}) ds$ and

$$J_t := \int_0^t \int_{r \in [0,1]} \int_{\theta \in \mathbb{S}^{d-1}} \left(\tilde{r}(g_{s-S(r,\theta)}) - \tilde{r}(g_{s-}) - r \sum_{i=1}^d \theta^i V_i^l \tilde{r}(g_{s-}) \right) d\theta \nu(dr) ds$$

are processes with finite variation.

To prove the proposition we will bound the supremum on $[0, T]$ of each of these five terms by means of $\|X^l \tilde{r}\|_\infty$ and $\|(X^l)^2 \tilde{r}\|_\infty$ for some $X \in \text{Vect}\{V_i, i = 1, \dots, d\}$. Thus we need the following lemma.

Lemma 2. *For $X \in \mathcal{P} = \text{Vect}\{V_i, i = 1, \dots, d\}$ we have $\|X^l \tilde{r}\|_\infty < +\infty$ and $\|(X^l)^2 \tilde{r}\|_\infty < +\infty$.*

Proof of the lemma. U_0 being a compact set it suffices to show that the supremum is finite on U_0^c . Since $\tilde{r} = r$ on U_0^c it remains to prove that $\sup_{g \in U_0^c} |X^l r(g)| < +\infty$ and $\sup_{g \in U_0^c} |(X^l)^2 r(g)| < +\infty$. The polar decomposition of g can be written $g = \tilde{k} \exp(r(g)V_1)k$ for some $\tilde{k}, k \in K$. Setting $x \in \mathbb{R}^d$ such that $X = \sum_i x^i V_i$ we have

$$r(g \exp(sX)) = r \left(\exp(r(g)V_1) \exp \left(s \sum_{i=1}^d (kx)^i V_i \right) \right).$$

Let $\theta \in \mathbb{S}^{d-1}$ be such that $(kx)^i = \|x\| \theta^i$. Then we compute explicitly, for $g \in U_0^c$:

$$\begin{aligned} r \left(\exp(r(g)V_1) \exp \left(s \sum_{i=1}^d (kx)^i V_i \right) \right) &= (\cosh)^{-1} (\cosh(r(g)) \cosh(s\|x\|) + \theta^1 \sinh(r(g)) \sinh(s\|x\|)) \\ &= r(g) + s\|x\|\theta^1 + \frac{s^2}{2} \|x\|^2 \frac{\cosh(r(g))}{\sinh(r(g))} (1 - (\theta^1)^2) + O(s^3). \end{aligned}$$

Then it comes that

$$X^l r(g) = \left. \frac{d}{ds} r(g \exp(sX)) \right|_{s=0} = \|x\| \theta^1,$$

and thus $\sup_{g \in U_0^c} |X^l r(g)| \leq \|x\|$.

Moreover

$$(X^l)^2 r(g) = \left. \frac{d^2}{ds^2} r(g \exp(sX)) \right|_{s=0} = \|x\|^2 \frac{\cosh(r(g))}{\sinh(r(g))} (1 - (\theta^1)^2)$$

and so $\sup_{g \in U_0^c} |(X^l)^2 r(g)| \leq 2\|x\|^2$. □

Return to the proof of Proposition 3. By Doob's norm inequalities (see [10]) we obtain

$$\mathbb{E} \left[\sup_{t \in [0, T]} |\hat{m}_t|^2 \right] \leq 4\mathbb{E} [(\hat{m}_T)^2] \leq 4\sigma^2 \mathbb{E} \left[\int_0^T \sum_{i=1}^d |V_i^l \tilde{r}(g_{s-})|^2 ds \right] \leq 4\sigma^2 T d \max_i \|V_i^l \tilde{r}\|_\infty^2 < +\infty.$$

and

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} |\tilde{m}_t|^2 \right] &\leq 4\mathbb{E} [(\tilde{m}_T)^2] \leq 4\mathbb{E} \left[\int_0^T \int_{r \in [0,1]} \int_{\theta \in \mathbb{S}^{d-1}} |\tilde{r}(g_{s-S(r,\theta)}) - \tilde{r}(g_{s-})|^2 \nu(dr) d\theta ds \right] \\ &\leq 8dT \int_0^1 r^2 \nu(dr) \max_i \|V_i^l \tilde{r}\|_\infty^2 < +\infty. \end{aligned}$$

We have also

$$\mathbb{E} \left[\sup_{t \in [0, T]} |I_t| \right] \leq \sigma^2 T d \max_i \|(V_i^l)^2 \tilde{r}\|_\infty < +\infty.$$

Moreover applying a Taylor inequality to $u \in [0, 1] \mapsto \tilde{r}(g_s - S(ur, \theta))$ it comes

$$\left| \tilde{r}(g_s - S(r, \theta)) - \tilde{r}(g_{s-}) - r \sum_{i=1}^d \theta^i V_i^l \tilde{r}(g_{s-}) \right| \leq \frac{1}{2} r^2 \sup_{\theta \in \mathbb{S}^{d-1}} \left\| \left(\sum_i \theta^i V_i^l \right)^2 \tilde{r} \right\|_\infty < +\infty$$

and thus

$$\mathbb{E} \left[\sup_{t \in [0, T]} |J_t| \right] \leq \frac{T}{2} \left(\int_0^1 r^2 \nu(dr) \right) \sup_{\theta \in \mathbb{S}^{d-1}} \left\| \left(\sum_i \theta^i V_i^l \right)^2 \tilde{r} \right\|_\infty < +\infty.$$

Finally, to bound the supremum of the last term of (4) we need the assumption 1. Indeed, since $|\tilde{r}(g_s - S(r, \theta)) - \tilde{r}(g_{s-})| \leq r \sup_{\theta \in \mathbb{S}^{d-1}} \left\| \left(\sum_i \theta^i V_i^l \right) \tilde{r} \right\|_\infty$ we obtain

$$\mathbb{E} \left[\sup_{t \in [0, T]} \left| \int_0^t \int_{U_0^c} \tilde{r}(g_{s-h}) - \tilde{r}(g_{s-}) ds \Pi(dh) \right| \right] \leq T d \left(\int_1^{+\infty} r \nu(dr) \right) \max_i \|V_i^l \tilde{r}\|_\infty < +\infty.$$

□

We can now state the main result of this section, the convergence of n_t to some asymptotic random variable n_∞ and the speed of convergence.

Proposition 4. *Denote by $b_t = (b_t^i)_{i=2, \dots, d}$ the \mathbb{R}^{d-1} -valued process such that*

$$n_t = \exp \left(\sum_{i=2}^d b_t^{i-1} (V_i - V_{1i}) \right).$$

Then b_t converges almost surely to b_∞ exponentially fast with rate α , i.e

$$\limsup_{t \rightarrow +\infty} \frac{1}{t} \log \|b_t - b_\infty\| \leq -\alpha. \quad (5)$$

As a consequence n_t converges almost-surely to $n_\infty := \exp \left(\sum_{i=2}^d b_\infty^{i-1} (V_i - V_{1i}) \right)$ and defining $h_t := e^{-t\alpha V_1} n_\infty^{-1} g_t$ we obtain

$$\lim_{t \rightarrow +\infty} \frac{1}{t} r(h_t) = 0 \quad \text{a.s.} \quad (6)$$

Proof. Denoting by $[t]$ the integer part of t we decompose $b_t = \sum_{j=1}^{[t]} (b_j - b_{j-1}) + b_t - b_{[t]}$. To prove that b_t converge and (5) holds it is sufficient to verify that

$$\limsup_{j \rightarrow \infty} \frac{1}{j} \log \sup_{s \in [0, 1]} \|b_j - b_{j+s}\| \leq -\alpha \quad \text{a.s.} \quad (7)$$

For $j \in \mathbb{N}$ and $s \in [0, 1]$ we have $g_j^{-1} g_{j+s} = k_j^{-1} a_j^{-1} n_j^{-1} n_{j+s} a_{j+s} k_{j+s}$ and thus

$$n_j^{-1} n_{j+s} = a_j (g_j^{-1} g_{j+s} k_j)_N a_j^{-1}. \quad (8)$$

Denote by $\tilde{b}_{j,s} = (\tilde{b}_{j,s}^i)_{i=2, \dots, d} \in \mathbb{R}^{d-1}$ such that $(g_j^{-1} g_{j+s} k_j)_N = \exp \left(\sum_{i=2}^d \tilde{b}_{j,s}^i (V_i - V_{1i}) \right)$. Then, (8) implies that

$$b_{j+s} - b_j = e^{-\alpha j} \tilde{b}_{j,s}. \quad (9)$$

Since g_t is a Levy process, $(\sup_{s \in [0, 1]} r(g_j^{-1} g_{j+s}))_{j \in \mathbb{N}}$ are i.i.d random variables. Moreover, by Proposition 3 their common expectation $\mathbb{E}[\sup_{s \in [0, 1]} r(g_s)]$ is finite. Thus, applying the law of large number it comes

$$\frac{1}{j} \sup_{s \in [0, 1]} r(g_j^{-1} g_{j+s}) \xrightarrow{j \rightarrow \infty} 0 \quad \text{a.s.} \quad (10)$$

Since $r((g_j^{-1}g_{j+s})_N) \leq 2r(g_j^{-1}g_{j+s})$ and, by (2), $\|\tilde{b}_{j,s}\|^2 = 2\left(\cosh\left(r(g_j^{-1}g_{j+s}k_j)_N\right) - 1\right)$ we deduce from (10) that for $\varepsilon > 0$ there exists j_0 such that for $j > j_0$

$$\|\tilde{b}_{j,s}\| \leq e^{\varepsilon j}. \quad (11)$$

Since, by Proposition 2, $\alpha_j = j\alpha + o(j)$ thus (7) follows from (9) and (11). To finish the proof of the proposition we need to check (6). We have

$$r(h_t) = r(e^{-\alpha t V_1} n_\infty^{-1} n_t a_t) \leq r(e^{-\alpha t V_1} a_t) + r(a_t^{-1} n_\infty^{-1} n_t a_t),$$

and $r(e^{-\alpha t V_1} a_t) = |\alpha_t - \alpha t| = o(t)$ and we obtain from (2) (since $a_t^{-1} n_\infty^{-1} n_t a_t \in N$),

$$r(a_t^{-1} n_\infty^{-1} n_t a_t) = \cosh^{-1}\left(1 + e^{2\alpha t} \frac{\|b_t - b_\infty\|^2}{2}\right).$$

So by (5) we have also $r(a_t^{-1} n_\infty^{-1} n_t a_t) = o(t)$ and (6) holds. \square

Remark 3. • *Without integrability condition, we can nevertheless show that g_t satisfy the irreducibility and contraction conditions of [13] and deduce that α_t converges almost surely to $+\infty$ and n_t converges to n_∞ . Nevertheless, by remark 2, we obtain in the case where $\int_1^{+\infty} r\nu(dr) = +\infty$, that almost surely $\frac{\alpha_t}{t}$ converges to $+\infty$.*

- *In [13] the author uses a stronger hypothesis to prove the rate of convergence of a Lévy process in a semi-simple group. It corresponds in our case to assume that $\int_0^{+\infty} r\nu(dr) < +\infty$.*

3.3 Asymptotic behavior of the $\mathbb{R}^{1,d}$ -component

The linear endomorphism $\xi \mapsto \exp(V_1)\xi$ is diagonalisable with eigenvalues $-1, 0, +1$. Denote by U^-, U^0 and U^+ the respective eigenspaces. Explicitly $U^- = \text{Vect}\{e_0 - e_1\}$, $U^0 = \text{Vect}\{e_2, \dots, e_d\}$ and $U^+ = \text{Vect}\{e_0 + e_1\}$. For $\xi \in \mathbb{R}^{1,d}$ we denote by $(\xi)^-, (\xi)^0$ and $(\xi)^+$ its projection on each eigenspace. Explicitly we obtain

$$\begin{aligned} (\xi)^- &= -\frac{1}{2}q(\xi, e_0 + e_1)(e_0 - e_1), & (\xi)^+ &= \frac{1}{2}q(\xi, e_0 - e_1)(e_0 + e_1) \\ (\xi)^0 &= \sum_{i=2}^d q(\xi, e_i)e_i. \end{aligned}$$

Recall that by definition, $\xi_t = \int_0^t g_s(e_0)ds$. The following proposition gives the asymptotic behavior of ξ_t .

Proposition 5. *There exists an asymptotic random variable $\lambda_\infty > 0$ such that*

$$(n_\infty^{-1}\xi_t)^- \xrightarrow[t \rightarrow +\infty]{} \lambda_\infty(e_0 - e_1),$$

and moreover

$$\limsup_{t \rightarrow +\infty} \frac{1}{t} \log \|(n_\infty^{-1}\xi_t)^- - \lambda_\infty(e_0 - e_1)\| \leq -\alpha. \quad (12)$$

We also have

$$\limsup_{t \rightarrow +\infty} \frac{1}{t} \log \|(n_\infty^{-1}\xi_t)^0\| \leq 0, \quad \text{and,} \quad \limsup_{t \rightarrow +\infty} \frac{1}{t} \log \|(n_\infty^{-1}\xi_t)^+\| \leq \alpha. \quad (13)$$

Proof. We have

$$-\frac{1}{2}q(n_\infty^{-1}\xi_t, e_0 + e_1) = \int_0^t -\frac{1}{2}q(n_\infty^{-1}n_s a_s(e_0), e_0 + e_1)ds \quad (14)$$

and the integrand can be written

$$\begin{aligned} -\frac{1}{2}q(n_\infty^{-1}n_s a_s(e_0), e_0 + e_1) &= -\frac{1}{2}q(e^{-\alpha s V_1} n_\infty^{-1} n_s a_s(e_0), e^{-\alpha s V_1}(e_0 + e_1)) \\ &= -\frac{1}{2}e^{-\alpha s} q(h_s(e_0), (e_0 + e_1)), \end{aligned}$$

where $h_s = e^{-\alpha s} n_\infty^{-1} g_s$ as defined in Proposition 4.

Denote by $(\tilde{r}_s, \theta_s) \in \mathbb{R}^+ \times \mathbb{S}^{d-1}$ the polar decomposition of $h_s(e_0) \in \mathbb{H}^d$. So $\tilde{r} = r(h_s)$ and

$$-\frac{1}{2}e^{-\alpha s} q(h_s(e_0), (e_0 + e_1)) = \frac{1}{2}e^{-\alpha s} \left(\cosh(\tilde{r}_s) - \tilde{\theta}_s^1 \sinh(\tilde{r}_s) \right) \in \frac{1}{2}[e^{-(\alpha s + \tilde{r}_s)}, e^{-(\alpha s - \tilde{r}_s)}].$$

Proposition 4 ensures that $\tilde{r}_s = o(s)$ a.s, so fixing $\varepsilon > 0$ arbitrary small we can find $s_0 > 0$ such that for all $s > s_0$ the integrand of (14) is positive and bounded by $e^{-(\alpha - \varepsilon)s}$. This ensures the convergence of $(n_\infty^{-1} \xi_t)^-$ to $\lambda_\infty(e_0 - e_1)$ with $\lambda_\infty > 0$. Moreover for $t > s_0$

$$\left| -\frac{1}{2}q(n_\infty^{-1} \xi_t, e_0 + e_1) - \lambda_\infty \right| \leq \int_t^{+\infty} e^{-(\alpha - \varepsilon)s} ds = \frac{1}{\alpha - \varepsilon} e^{-(\alpha - \varepsilon)t},$$

which prove (12).

Now

$$(n_\infty^{-1} \xi_t)^0 = \sum_{i=2}^d \int_0^t q(n_\infty^{-1} n_s a_s(e_0), e_i) ds e_i,$$

and for $i = 2, \dots, d$ we have $q(n_\infty^{-1} n_s a_s(e_0), e_i) = q(e^{-\alpha s V_1} n_\infty^{-1} n_s a_s(e_0), e_i) = \tilde{\theta}_s^i \sinh(\tilde{r}_s)$ so $|q(n_\infty^{-1} n_s a_s(e_0), e_i)| \leq e^{\tilde{r}_s}$ and this ensures that $\limsup_{t \rightarrow +\infty} \frac{1}{t} \log \|(n_\infty^{-1} \xi_t)^0\| \leq 0$.

Moreover,

$$(n_\infty^{-1} \xi_t)^+ = \left(\int_0^t \frac{1}{2} q(n_\infty^{-1} n_s a_s(e_0), e_0 - e_1) ds \right) (e_0 - e_1)$$

and

$$\begin{aligned} \frac{1}{2}q(n_\infty^{-1} n_s a_s(e_0), e_0 - e_1) &= \frac{1}{2}e^{\alpha s} q(h_t(e_0), e_0 - e_1) \\ &= \frac{1}{2}e^{\alpha s} \left(\cosh(\tilde{r}_s) - \tilde{\theta}_s^1 \sinh(\tilde{r}_s) \right) \in \frac{1}{2}[e^{\alpha s - \tilde{r}_s}, e^{\alpha s + \tilde{r}_s}]. \end{aligned}$$

So

$$\|(n_\infty^{-1} \xi_t)^+\| \leq \|(n_\infty^{-1} \xi_{s_0})^+\| + \int_{s_0}^t e^{(\alpha + \varepsilon)s} ds,$$

and thus $\limsup_{t \rightarrow +\infty} \frac{1}{t} \log \|(n_\infty^{-1} \xi_t)^+\| \leq \alpha$. \square

3.4 Geometric description of the convergence

Denote by $\mathbf{p} : \mathbb{R}^{1,d} \setminus \{0\} \rightarrow \mathbb{P}^d \mathbb{R}$ the projection onto the projective space of dimension d . The hyperboloid \mathbb{H}^d is mapped onto the interior of a projective ball and its boundary $(\partial \mathbb{H}^d \simeq \mathbb{S}^{d-1})$ is the image of the q -isotropy cone

$$\partial \mathbb{H}^d := \mathbf{p}(\{\xi, q(\xi) = 0\} \setminus \{0\}).$$

From the relation

$$q(\dot{\xi}_t, n_t(e_0 + e_1)) = q(g_t(e_0), n_t(e_0 + e_1)) = q(e_0, a_t^{-1}(e_0 + e_1)) = e^{-\alpha t} \xrightarrow{t \rightarrow \infty} 0,$$

we deduce that all limit points of $\mathbf{p}(\dot{\xi}_t)$ are q -orthogonal to $\theta_\infty := \mathbf{p}(n_\infty(e_0 + e_1))$. Since the only point of $\overline{\mathbf{p}(\mathbb{H}^d)}$ which is q -orthogonal to θ_∞ is θ_∞ itself it comes that $\mathbf{p}(\dot{\xi}_t)$ converges to θ_∞ in $\mathbb{P}^d \mathbb{R}$. Now, identifying $\mathbb{P}^d \mathbb{R}$ with its affine chart $\{\xi, \xi^0 = 1\}$ we can consider that $\theta_\infty \in \mathbb{S}^d$. From (2)

we deduce that $r_t \rightarrow +\infty$ and since $\mathbf{p}(\dot{\xi}_t) = \mathbf{p}(e_0 + \theta_t \frac{\sinh(r_t)}{\cosh(r_t)})$ it comes that θ_t converges to θ_∞ in \mathbb{S}^d . The two asymptotic random variables θ_∞ and n_∞ are linked by

$$\mathbf{p}(e_0 + \theta_\infty) = \mathbf{p}(n_\infty(e_0 + e_1))$$

or more explicitly, $b_\infty \in \mathbb{R}^{d-1}$ (defined by $n_\infty = \exp\left(\sum_{i=2}^d b_\infty^{i-1}(V_1 - V_{1i})\right)$) is the stereographic projection of θ_∞

$$\theta_\infty = \frac{1}{1 + \|b_\infty\|^2} \begin{pmatrix} 1 - \|b_\infty\|^2 \\ 2b_\infty \end{pmatrix}.$$

Concerning the asymptotic behavior of ξ_t , Proposition 5 ensures that $q(\xi_t, n_\infty(e_0 + e_1))$ converges to λ_∞ . Thus geometrically ξ_t is asymptotic to an affine hyperplan which is q -orthogonal to $n_\infty(e_0 + e_1)$ (or $e_0 + \theta_\infty$) and passing by $\lambda_\infty(e_0 - e_1)$.

4 Lyapunov spectrum and stable manifolds

4.1 Lyapunov spectrum

The Levy process \tilde{g}_t , with values in \tilde{G} and starting at some \tilde{g} , can be obtained by solving the following left invariant stochastic integro-differential equation in \tilde{G}

$$\begin{aligned} \forall f \in C^2(\tilde{G}), \quad f(\tilde{g}_t) = f(\tilde{g}) + \sigma \sum_{i=1}^d \int_0^t V_i^l f(\tilde{g}_{s-}) \circ dB_s^i + \int_0^t H_0^l(\tilde{g}_{s-}) ds + \int_0^t \int_{U_0} (f(\tilde{g}_{s-h}) - f(\tilde{g}_{s-})) \tilde{N}(ds, dh) \\ + \int_0^t \int_{U_0} \left(f(\tilde{g}_{s-h}) - f(\tilde{g}_{s-}) - r(h) \sum_{i=1}^d \theta^i(h) V_i^l f(\tilde{g}_{s-}) \right) ds \Pi(dh) \quad (15) \\ + \int_0^t \int_{(U_0)^c} (f(\tilde{g}_{s-h}) - f(\tilde{g}_{s-})) N(ds, dh). \end{aligned}$$

This stochastic differential equation induces a stochastic flow φ_t in \tilde{G} which maps \tilde{g} on the solution at time t and starting at \tilde{g} of (15). By left invariance φ_t is also defined by

$$\begin{aligned} \varphi_t : \tilde{G} &\longrightarrow \tilde{G} \\ \tilde{g} &\longmapsto \tilde{g} \tilde{g}_t, \end{aligned}$$

where \tilde{g}_t is starting at Id.

Denote by $\|\cdot\|$ any norm on $\text{Lie}(\tilde{G})$ and by $\|\cdot\|_{\tilde{g}}$ the left invariant (Finsler) metric associated in \tilde{G} on $T_{\tilde{g}}\tilde{G}$. For $v \in T_{\tilde{g}}\tilde{G}$ we aim to investigate the asymptotic exponential rate of growth or decay of $\|d\varphi_t(\tilde{g})(v)\|_{\varphi_t(\tilde{g})}$. Denote by $L_{\tilde{g}}$ the left translation by \tilde{g} in \tilde{G} . By left invariance of the flow, $\|d\varphi_t(\tilde{g})(v)\|_{\varphi_t(\tilde{g})} = \|d\varphi_t(\text{Id})(\tilde{X})\|_{\varphi_t(\text{Id})}$ where $\tilde{X} := (dL_{\tilde{g}})^{-1}(v) \in T_{\text{Id}}\tilde{G} = \text{Lie}(\tilde{G})$. For $\tilde{g} = (g, \xi) \in \tilde{G}$ and $\tilde{X}, \tilde{Y} \in \text{Lie}(\tilde{G})$ it comes

$$\text{Ad}(\tilde{g})(\tilde{X}) = (\text{Ad}(g)(X), gx - \text{Ad}(g)(X)\xi) \quad (16)$$

$$\text{ad}(\tilde{Y})(\tilde{X}) = (\text{ad}(Y)(X), Yx - Xy). \quad (17)$$

The endomorphism $\tilde{X} \mapsto \text{ad}(V_1, 0)(\tilde{X})$ is diagonalisable on $\text{Lie}(\tilde{G})$. Its eigenvalues are $-1, 0, 1$ and we denote by \tilde{U}^- , \tilde{U}^0 and \tilde{U}^+ the eigenspaces associated.

We can check that

$$\tilde{X} \in \tilde{U}^+ \iff X \in \bar{\mathcal{N}} \text{ and } x \in U^+$$

$$\tilde{X} \in \tilde{U}^0 \iff X \in \mathcal{A} \oplus \mathcal{M} \text{ and } x \in U^0$$

$$\tilde{X} \in \tilde{U}^- \iff X \in \mathcal{N} \text{ and } x \in U^-.$$

Set

$$\tilde{g}_\infty := (n_\infty, \lambda_\infty(e_0 - e_1)) \in \tilde{G}$$

and $V_\infty^- := \text{Ad}(\tilde{g}_\infty)(\tilde{U}^+)$, $V_\infty^0 := \text{Ad}(\tilde{g}_\infty)(\tilde{U}^0 + \tilde{U}^+)$.

We denote by

$$\tilde{h}_t := (e^{-t\alpha V_1}, 0) \tilde{g}_\infty^{-1} \tilde{g}_t,$$

where we recall that $\tilde{g}_t := \varphi_t(\text{Id})$ is starting at Id .

Theorem 1. *Let $\tilde{X} \in \text{Lie}(\tilde{G})$. For almost every trajectory*

$$\frac{1}{t} \log \|d\varphi_t(\text{Id})(\tilde{X})\|_{\varphi_t(\text{Id})} \xrightarrow{t \rightarrow +\infty} \begin{cases} \alpha & \text{if } \tilde{X} \in \text{Lie}(\tilde{G}) \setminus V_\infty^0 \\ 0 & \text{if } \tilde{X} \in V_\infty^0 \setminus V_\infty^- \\ -\alpha & \text{if } \tilde{X} \in V_\infty^- \setminus \{0\} \end{cases}$$

Proof. By left invariance of $\|\cdot\|$, $\|d\varphi_t(\text{Id})(\tilde{X})\|_{\varphi_t(\text{Id})} = \|\text{Ad}(\tilde{g}_t^{-1})(\tilde{X})\|$. We set, for $\tilde{g} \in \tilde{G}$

$$\|\text{Ad}(\tilde{g})\| := \sup_{\tilde{X} \neq 0} \frac{\|\text{Ad}(\tilde{g})(\tilde{X})\|}{\|\tilde{X}\|}.$$

Let $\tilde{X} \in \text{Lie}(\tilde{G})$. Writting $\tilde{g}_t^{-1} = (\tilde{h}_t)^{-1} (e^{-t\alpha V_1}, 0) \tilde{g}_\infty^{-1}$ we deduce that

$$\begin{aligned} \frac{\|\text{Ad}((e^{-t\alpha V_1}, 0) \tilde{g}_\infty^{-1})(\tilde{X})\|}{\|\text{Ad}(\tilde{h}_t)\|} &\leq \|\text{Ad}(\tilde{g}_t^{-1})(\tilde{X})\| \\ &\leq \|\text{Ad}(\tilde{h}_t)^{-1}\| \|\text{Ad}((e^{-t\alpha V_1}, 0) \tilde{g}_\infty^{-1})(\tilde{X})\|. \end{aligned} \quad (18)$$

Suppose for the moment that

$$\limsup_{t \rightarrow +\infty} \frac{1}{t} \log \|\text{Ad}(\tilde{h}_t)^{-1}\| \leq 0 \quad (19)$$

$$\text{and } \limsup_{t \rightarrow +\infty} \frac{1}{t} \log \|\text{Ad}(\tilde{h}_t)\| \leq 0. \quad (20)$$

Then we deduce from (18) that $\frac{1}{t} \log \|\text{Ad}(\tilde{g}_t^{-1})(\tilde{X})\|$ and $\frac{1}{t} \log \|\text{Ad}((e^{-t\alpha V_1}, 0) \tilde{g}_\infty^{-1})(\tilde{X})\|$ have the same limit when t goes to ∞ . The linear isomorphism $\text{Ad}(e^{-t\alpha V_1}, 0)$ is diagonalisable with eigenvalues $e^{-\alpha t}$, 1 and $e^{\alpha t}$ associated respectively to the eigenspaces \tilde{U}^+ , \tilde{U}^0 and \tilde{U}^- . Decomposing $\text{Ad}(\tilde{g}_\infty)^{-1}(\tilde{X})$ in the direct sum $\tilde{U}^- \oplus \tilde{U}^0 \oplus \tilde{U}^+$ and using a Euclidean norm $\|\cdot\|$ on $\text{Lie}(\tilde{G})$ for which this decomposition is orthogonal, we deduce easily the theorem (note that the convergence is independant of the chosen norm).

Thus it remains to prove (19) and (20). We have

$$\begin{aligned} \tilde{h}_t &= (e^{-t\alpha V_1}, 0) \tilde{g}_\infty^{-1} \tilde{g}_t = (e^{-t\alpha V_1}, 0) (n_\infty^{-1}, -\lambda_\infty(e_0 - e_1)) (n_t a_t k_t, \xi_t) \\ &= (h_t, e^{-t\alpha V_1} (n_\infty^{-1} \xi_t) - \lambda_\infty e^{t\alpha} (e_0 - e_1)) \\ &= (\text{Id}, e^{-t\alpha V_1} (n_\infty^{-1} \xi_t) - \lambda_\infty e^{t\alpha} (e_0 - e_1)) (h_t, 0) \end{aligned}$$

Let $\varepsilon > 0$. By Proposition 4 we can find $t_0 > 0$ such that $\forall t > t_0$ $r(h_t) \leq \varepsilon t$ and by Proposition 5 we have

$$\begin{aligned} \|e^{-t\alpha V_1} (n_\infty^{-1} \xi_t) - \lambda_\infty e^{t\alpha} (e_0 - e_1)\| &\leq e^{\alpha t} \|(n_\infty^{-1} \xi_t)_- - \lambda_\infty (e_0 - e_1)\| + \|(n_\infty^{-1} \xi_t)_0\| + e^{-\alpha t} \|(n_\infty^{-1} \xi_t)_+\| \\ &\leq e^{\varepsilon t} \end{aligned}$$

Now using the following Lemma 3 we deduce easily (19) and (20). \square

Lemma 3. *There exist positive constants α, β, γ such that for $g \in G$ and $\xi \in \mathbb{R}^{1,d}$*

$$\begin{aligned} \|\text{Ad}(g, 0)\| &\leq \alpha e^{r(g)} \\ \|\text{Ad}(\text{Id}, \xi)\| &\leq \beta \|\xi\| + \gamma. \end{aligned}$$

Proof. All norms are equivalent and it suffices to check the inequalities for some particular norms. Let choose the following $SO(d)$ -invariant euclidean norm on $\text{Lie}(\tilde{G})$

$$\|(X, x)\| := \sqrt{\text{Tr}(X^t X) + x^t x}.$$

We obtain easily

$$\|\text{Ad}(g, 0)\| = e^{r(g)}.$$

Taking now $\|(X, x)\| := \sqrt{\text{Tr}(X^t X) + x^t x}$ we get

$$\begin{aligned} \|\text{Ad}(\text{Id}, \xi)(X, x)\| &= \sqrt{\text{Tr}(X^t X) + \|x - X\xi\|^2} \leq \|(X, x)\| + \|X\xi\| \\ &\leq \|(X, x)\| + \sqrt{\text{Tr}(X^t X)} \max_i |\xi^i| \leq \|(X, x)\| \left(1 + \max_i |\xi^i|\right). \end{aligned}$$

Thus $\|\text{Ad}(\text{Id}, \xi)\| \leq 1 + \alpha\|\xi\|$ for a constant $\alpha > 0$ independant of ξ . □

4.2 Stable manifolds

First, remark that V_∞^- and V_∞^0 are Lie sub-algebras of $\text{Lie}(\tilde{G})$. Denote by

$$\mathcal{V}_\infty^- := \exp(V_\infty^-), \quad \text{and} \quad \mathcal{V}_\infty^0 := \exp(V_\infty^0)$$

the closed subgroup of \tilde{G} associated.

Fix now a euclidean norm $\|\cdot\|$ on $\text{Lie}(\tilde{G})$ which is $\text{Ad}(K)$ -invariant. Such a norm is of the form

$$\left\| \left(\begin{pmatrix} 0 & b^t \\ b & C \end{pmatrix}, x \right) \right\| := \sqrt{\kappa^2 b^t b + \beta^2 \text{Tr}(C^t C) + \gamma^2 \bar{x}^t \bar{x} + \delta^2 (x^0)^2},$$

for some positive constants κ, β, γ and δ . We denote by d the distance in \tilde{G} associated to the left invariant Riemannian metric induced by $\|\cdot\|$. To simplify notations, we denote by $d(g, h)$ the distance between $(g, 0)$ and $(h, 0)$ for $g, h \in G$.

The following result shows that the stable manifold associated to φ_t is $\varphi_0 \mathcal{V}_\infty^-$.

Theorem 2. *Let \tilde{g} and \tilde{g}' two distinct points in \tilde{G} .*

- *If $\tilde{g}' \in \tilde{g} \mathcal{V}_\infty^-$ then*

$$\frac{1}{t} \log d(\varphi_t(\tilde{g}), \varphi_t(\tilde{g}')) \xrightarrow{t \rightarrow +\infty} -\alpha.$$

- *If $\tilde{g}' \notin \tilde{g} \mathcal{V}_\infty^-$ then*

$$\liminf_{t \rightarrow \infty} d(\varphi_t(\tilde{g}), \varphi_t(\tilde{g}')) > 0.$$

The properties of d we need in the proof of Theorem 2 are sum up in the following proposition

Proposition 6. *i) Left invariance*

$$\forall \tilde{g}, \tilde{h} \in \tilde{G}, \quad d(\tilde{g}, \tilde{g}\tilde{h}) = d(\text{Id}, \tilde{h}).$$

Thus $d(\text{Id}, \tilde{g}^{-1}) = d(\text{Id}, \tilde{g})$ and triangularity inequality writes:

$$\forall \tilde{g}, \tilde{h} \in \tilde{G}, \quad d(\text{Id}, \tilde{g}\tilde{h}) \leq d(\text{Id}, \tilde{g}) + d(\text{Id}, \tilde{h})$$

ii) K -right invariance

$$\forall \tilde{g} \in \tilde{G} \text{ and } k \in K, \quad d((k, 0), \tilde{g}(k, 0)) = d(\text{Id}, \tilde{g})$$

iii) For $\tilde{X} \in \text{Lie}(\tilde{G})$

$$d(\text{Id}, \exp(\tilde{X})) \leq \|\tilde{X}\|.$$

iv) There exists a neighborhood \mathcal{O} of 0 in $\text{Lie}(\tilde{G})$ and a constant $C > 0$ such that

$$\forall \tilde{X} \in \mathcal{O}, \quad C \|\tilde{X}\| \leq d(\text{Id}, \exp(\tilde{X}))$$

v) $\forall (g, \xi) \in \tilde{G}$

$$d(\text{Id}, g) \leq d(\text{Id}, (g, \xi))$$

vi) For $g = S(r, \theta)R$ and $g' = S(r', \theta')R'$ we have

$$\frac{\kappa}{\sqrt{\kappa^2 + 2\beta^2}} d(S(r, \theta), S(r', \theta')) \leq d(g, g')$$

vii) For all $r \geq 0$ and $\theta \in \mathbb{S}^{d-1}$

$$d(\text{Id}, S(r, \theta)) = \kappa r$$

Proof of Proposition 6. i) and ii). The left and K -right invariance follows from the definition of the metric as being a left invariant Riemannian metric on \tilde{G} defined from an $\text{Ad}(K)$ -invariant inner product on $\text{Lie}(\tilde{G})$. Inequality iii) is obtained remarking that the length of the path $t \in [0, 1] \mapsto \exp(t\tilde{X})$ is equal to $\|\tilde{X}\|$.

iv). Denote by $\widehat{\exp} : \text{Lie}(\tilde{G}) \rightarrow \tilde{G}$ be the exponential map at Id induced by the metric $\|\cdot\|$ in \tilde{G} : for $\tilde{X} \in \text{Lie}(\tilde{G})$, $\widehat{\exp}(\tilde{X}) = \gamma_{\tilde{X}}(1)$ where $t \in [0, 1] \mapsto \gamma_{\tilde{X}}(t)$ is the geodesic starting from Id in the direction \tilde{X} . The differential at 0 of $\widehat{\exp}$ is known to be identity and there exists a sufficient small neighborhood \mathcal{O}' of 0 in $\text{Lie}(\tilde{G})$ such that:

$$\forall \tilde{X} \in \mathcal{O}', \quad \|\tilde{X}\| = d(I, \widehat{\exp}(\tilde{X})). \quad (*)$$

Furthermore, the map $\widehat{\exp}^{-1} \circ \exp$ can be defined in a neighborhood of 0 and its differential at 0 is the identity: $\widehat{\exp}^{-1} \circ \exp(\tilde{X}) = \tilde{X} + o(\|\tilde{X}\|)$. So we can find \mathcal{O} neighborhood of 0 and $C > 0$ such that for all $\tilde{X} \in \mathcal{O}$, $C \|\tilde{X}\| \leq \|\widehat{\exp}^{-1} \circ \exp(\tilde{X})\| \leq \frac{1}{C} \|\tilde{X}\|$. Taking \mathcal{O} small enough so that $\widehat{\exp}^{-1} \circ \exp(\mathcal{O}) \subset \mathcal{O}'$, we can apply (*) to $\widehat{\exp}^{-1} \circ \exp(\tilde{X})$, thus yielding $\|\widehat{\exp}^{-1} \circ \exp(\tilde{X})\| = d(I, \exp(\tilde{X}))$ for every $\tilde{X} \in \mathcal{O}$.

v). Each path $s \in [0, 1] \mapsto (g_s, \xi_s)$ joining Id to (g, ξ) is of length $\int_0^1 \|(g_s^{-1}\dot{g}_s, g_s^{-1}\dot{\xi}_s)\| ds$ which is greater than $\int_0^1 \|(g_s^{-1}\dot{g}_s, 0)\| ds$ corresponding to the path $s \in [0, 1] \mapsto (g_s, 0)$ joining Id to $(g, 0)$.

vi). Consider a path $s \in [0, 1] \mapsto S(r_s, \theta_s)R_s$ joining g to g' . We compute, using dot notation for $\frac{d}{ds}$

$$R_s^{-1} S(r_s, \theta_s)^{-1} \frac{d}{ds} (S(r_s, \theta_s) R_s) = \begin{pmatrix} 0 & \dot{r}_s \theta_s^t + \sinh(r_s) \dot{\theta}_s^t \\ \dot{r}_s \theta_s + \sinh(r_s) \dot{\theta}_s & (\cosh(r_s) - 1) (\dot{\theta}_s \theta_s^t - \theta_s \dot{\theta}_s^t) + R_s^{-1} \dot{R}_s \end{pmatrix}.$$

Its length $l := \int_0^1 \|R_t^{-1} S(r_t, \theta_t)^{-1} \frac{d}{dt} S(r_t, \theta_t) R_t\| dt$ is larger than

$$\int_0^1 \kappa \|\dot{r}_s \theta_s + \sinh(r_s) \dot{\theta}_s\| ds = \int_0^1 \kappa \sqrt{(\dot{r}_s)^2 + \sinh(r_s)^2 \|\dot{\theta}_s\|^2} ds.$$

Moreover, the path $s \mapsto S(r_s, \theta_s)$ which join $S(r, \theta)$ to $S(r', \theta')$ is of length

$$\int_0^1 \sqrt{\kappa^2 \left((\dot{r}_s)^2 + \sinh(r_s)^2 \|\dot{\theta}_s\|^2 \right) + \beta^2 \left(2(\cosh(r) - 1)^2 \|\dot{\theta}_s\|^2 \right)} ds \quad (21)$$

$$\leq \frac{\sqrt{\kappa^2 + 2\beta^2}}{\kappa} \int_0^1 \kappa \sqrt{(\dot{r}_s)^2 + \sinh(r_s)^2 \|\dot{\theta}_s\|^2} ds. \quad (22)$$

Thus

$$d(S(r, \theta), S(r', \theta')) \leq \frac{\sqrt{\kappa^2 + 2\beta^2}}{\kappa} l$$

and taking the infimum over all the path joining g to g' we obtain vi). \square

Proof of Theorem 2. Let $\tilde{Y} \in \text{Lie}(\tilde{G}) \setminus \{0\}$ be such that $\exp(\tilde{Y}) = \tilde{g}^{-1}\tilde{g}'$. Then

$$d(\varphi_t(\tilde{g}), \varphi_t(\tilde{g}')) = d\left(\text{Id}, \tilde{g}_t^{-1} \exp(\tilde{Y}) \tilde{g}_t\right) = d\left(\text{Id}, \exp\left(\text{Ad}(\tilde{g}_t^{-1})(\tilde{Y})\right)\right).$$

By Theorem 1, if $\tilde{Y} \in \text{Ad}(g_\infty)(\tilde{U}^+)$ (i.e. if $g' \in g\mathcal{V}_\infty^-$) then $\|\text{Ad}(g_t^{-1})(\tilde{Y})\|$ converge to 0 exponentially fast with rate α and so for large t it evolves in \mathcal{O} . Thus, using *iii*) and *iv*) of Proposition 6 we obtain the first point of the theorem as a direct consequence of Theorem 1.

Set $\tilde{X} := \text{Ad}(\tilde{g}_\infty^{-1})(\tilde{Y})$, thus

$$\tilde{Y} = \text{Ad}(\tilde{g}_\infty)\left(\left(\tilde{X}\right)^+ + \left(\tilde{X}\right)^0 + \left(\tilde{X}\right)^-\right),$$

and write

$$\begin{aligned} \left(\tilde{X}\right)^+ &= (X^+, x^+), \text{ where } X^+ \in \bar{\mathcal{N}} \text{ and } x^+ \in U^+ \\ \left(\tilde{X}\right)^0 &= (X^0, x^0), \text{ where } X^0 \in \mathcal{A} \oplus \mathcal{M} \text{ and } x^0 \in U^0 \\ \left(\tilde{X}\right)^- &= (X^-, x^-), \text{ where } X^- \in \mathcal{N} \text{ and } x^- \in U^-. \end{aligned}$$

Now suppose that $\tilde{g}' \notin \tilde{g}\mathcal{V}_\infty^-$ which is equivalent to $\left(\tilde{X}\right)^0 \neq 0$ or $\left(\tilde{X}\right)^- \neq 0$.

Suppose first that $\left(\tilde{X}\right)^- \neq 0$. Thus $\tilde{Y} \in \text{Lie}(\tilde{G}) \setminus V_\infty^0$ and by Theorem 1 $\|\text{Ad}(\tilde{g}_t^{-1})\tilde{Y}\|$ converges to $+\infty$ exponentially fast. Now suppose by contradiction that $\liminf_{t \rightarrow +\infty} d\left(\text{Id}, \exp\left(\text{Ad}(\tilde{g}_t^{-1})\tilde{Y}\right)\right) = 0$. Then we can find a s_t such that $d\left(\text{Id}, \exp\left(\text{Ad}(\tilde{g}_{s_t}^{-1})(\tilde{Y})\right)\right)$ converges to 0 and for large t $\text{Ad}(\tilde{g}_{s_t}^{-1})(\tilde{Y})$ lies in \mathcal{O} . The inequality *iv*) of Proposition 6 give us the contradiction and we have proved the second point of the Theorem if $\left(\tilde{X}\right)^- \neq 0$.

So we can suppose $\left(\tilde{X}\right)^- = 0$ and $\left(\tilde{X}\right)^0 \neq 0$.

First case: $X^0 \neq 0$. By *v*) of Proposition 6, $d(\text{Id}, g_t^{-1} \exp(Y) g_t) \leq d\left(\text{Id}, \tilde{g}_t^{-1} \exp(\tilde{Y}) \tilde{g}_t\right)$ and it remains to prove that $\liminf_{t \rightarrow \infty} d(\text{Id}, g_t^{-1} \exp(Y) g_t)$ is positive. But $Y = \text{Ad}(n_\infty)(X)$ and $X = X^+ + X^0 \in \bar{\mathcal{N}} \oplus \mathcal{A} \oplus \mathcal{M} \setminus \bar{\mathcal{N}}$. Consider an Iwasawa decomposition of $\exp(X)$ in G

$$\exp(X) = \bar{n} a m, \quad n \in \bar{\mathcal{N}}, \quad a \in A, \quad \text{and } m \in M,$$

and $am \neq \text{Id}$. Since $\forall g, h \quad d(\text{Id}, gh) \leq d(\text{Id}, g) + d(\text{Id}, h)$ we get

$$\begin{aligned} d(\text{Id}, g_t^{-1} \exp(Y) g_t) &= d(\text{Id}, h_t^{-1} e^{-t\alpha V_1} \bar{n} a m e^{t\alpha V_1} h_t^{-1}) \\ &\geq d(\text{Id}, h_t^{-1} e^{-t\alpha V_1} a m e^{t\alpha V_1} h_t) - d(\text{Id}, h_t^{-1} e^{-t\alpha V_1} \bar{n} e^{t\alpha V_1} h_t). \end{aligned}$$

Writting $\bar{n} = \exp(Z)$, $Z \in \bar{\mathcal{N}}$ and $d(\text{Id}, h_t^{-1} e^{-t\alpha V_1} \bar{n} e^{t\alpha V_1} h_t)$ is dominated by $e^{-t\alpha + r(h_t)} \|Z\|$ (by Lemma 3 and *iii*) of Proposition 6) and converges exponentially fast to zero (recall that by Proposition 4 $r(h_t) = o(t)$ a.s.). Thus it remains to prove that $\liminf d(\text{Id}, h_t^{-1} e^{-t\alpha V_1} a m e^{t\alpha V_1} h_t) > 0$ to finish the proof in the first case. This is ensured by the following Lemma.

Lemma 4. *Let $a \in A$ and $m \in M$ s.t $am \neq \text{Id}$. Then $\exists C > 0$, $\forall g \in G$, $d(\text{Id}, g^{-1} a m g) > C$.*

Proof of lemma 4. Consider the polar decomposition $g = S(r, \theta)R$. Suppose first that $a = \text{Id}$ and

$m \neq \text{Id}$. Then we get

$$\begin{aligned}
d(\text{Id}, g^{-1}mg) &= d(\text{Id}, S(r, -\theta)mS(r, \theta)) = d(S(r, \theta), mS(r, \theta)) = d(S(r, \theta), S(r, m\theta)m) \\
&\geq \frac{\kappa}{\sqrt{\kappa^2 + 2\beta^2}} d(S(r, \theta), S(r, m\theta)) \quad \text{by } vi) \text{ of Proposition 6} \\
&\geq \frac{\kappa}{\sqrt{\kappa^2 + 2\beta^2}} (d(S(r, \theta), S(r, \theta)m^{-1}) - d(S(r, \theta)m^{-1}, S(r, m\theta))) \\
&= \frac{\kappa}{\sqrt{\kappa^2 + 2\beta^2}} (d(\text{Id}, m^{-1}) - d(S(r, \theta), mS(r, \theta))) \\
&= \frac{\kappa}{\sqrt{\kappa^2 + 2\beta^2}} (d(\text{Id}, m) - d(\text{Id}, g^{-1}mg))
\end{aligned}$$

Thus $d(\text{Id}, g^{-1}mg) \geq \frac{\kappa}{\kappa + \sqrt{\kappa^2 + 2\beta^2}} d(\text{Id}, m) > 0$.

Suppose now $a \neq \text{Id}$. Let $u \neq 0$ such that $a = \exp(uV_1)$, then an explicit computation gives:

$$\cosh r(g^{-1}amg) = \cosh(u) (\cosh(r)^2 - ((m\theta)^1)^2 \sinh(r)^2) - \sinh(r)^2 \sum_{i=2}^d \theta^i (m\theta)^i \quad (23)$$

$$= \cosh(u) + \left(\cosh(u)(1 - ((m\theta)^1)^2) - \sum_{i=2}^d \theta^i (m\theta)^i \right) \sinh(r)^2 \quad (24)$$

$$\geq \cosh(u) + (1 - \theta^t(m\theta)) \sinh(r)^2 \quad \text{we used } (m\theta)^1 = \theta^1 \quad (25)$$

$$\geq \cosh(u). \quad (26)$$

Then by *vi*) and *vii*) of Proposition 6 it comes

$$d(\text{Id}, g^{-1}amg) \geq \frac{\kappa}{\sqrt{\kappa^2 + 2\beta^2}} \kappa u > 0.$$

□

Return to the proof of Theorem 2.

Second case: $\mathbf{X}^0 = \mathbf{0}$ but $\mathbf{x}^0 \neq \mathbf{0}$. So $\tilde{X} = (X^+, x^+ + x^0)$ and explicitly

$$\exp(\tilde{X}) = (\exp(X^+), x^0 + x^+ + \frac{X^+x^0}{2}) = (\text{Id}, \xi)(\exp(X^+), 0),$$

where we have set $\xi := x^0 + x^+ + \frac{X^+x^0}{2}$.

Thus

$$\begin{aligned}
\tilde{g}_t^{-1} \exp(\tilde{Y}) \tilde{g}_t &= \tilde{h}_t^{-1} (e^{-t\alpha V_1}, 0) \exp(\tilde{X}) (e^{t\alpha V_1}, 0) \tilde{h}_t \\
&= (\text{Id}, h_t^{-1} e^{-t\alpha V_1} \xi) (\exp(\text{Ad}(h_t^{-1} e^{-t\alpha V_1}) X), 0),
\end{aligned}$$

and

$$d(\text{Id}, \tilde{g}_t^{-1} \exp(\tilde{Y}) \tilde{g}_t) \geq d(\text{Id}, (\text{Id}, h_t^{-1} e^{-t\alpha V_1} \xi)) - d(\text{Id}, (\exp(\text{Ad}(h_t^{-1} e^{-t\alpha V_1}) X), 0)).$$

As done previously in the first case, $d(\text{Id}, (\exp(\text{Ad}(h_t^{-1} e^{-t\alpha V_1}) X), 0))$ converges exponentially fast to 0 and it remains to prove that

$$\liminf_{t \rightarrow \infty} d(\text{Id}, (\text{Id}, h_t^{-1} e^{-t\alpha V_1} \xi)) > 0. \quad (27)$$

Suppose by contradiction that we can find s_t such that $d(\text{Id}, (\text{Id}, h_{s_t}^{-1} e^{-s_t \alpha V_1}(\xi)))$ converges to 0. By *iv*) of Proposition 6 for large t

$$d(\text{Id}, (\text{Id}, h_{s_t}^{-1} e^{-s_t \alpha V_1}(\xi))) \geq \|h_{s_t}^{-1} e^{-s_t \alpha V_1}(\xi)\|$$

Since $\frac{X+x^0}{2} \in U^+$ we obtain directly that $q(\xi) = q(x^0)$ which is negative since x^0 is supposed to be non zero. But

$$\begin{aligned} \|h_{s_t}^{-1} e^{-s_t \alpha V_1}(\xi)\|^2 &= \gamma^2 \left(\sum_{i=1}^d q(h_{s_t}^{-1} e^{-s_t \alpha V_1}(\xi), e_i)^2 \right) + \delta^2 q(h_{s_t}^{-1} e^{-s_t \alpha V_1}(\xi), e_0)^2 \\ &\geq \min(\gamma, \delta)^2 \left(\sum_{i=0}^d q(h_{s_t}^{-1} e^{-s_t \alpha V_1}(\xi), e_i)^2 \right) \\ &= \min(\gamma, \delta)^2 \left(2q(h_{s_t}^{-1} e^{-s_t \alpha V_1}(\xi), e_0)^2 - q(h_{s_t}^{-1} e^{-s_t \alpha V_1}(\xi))^2 \right) \geq -\min(\gamma, \delta)^2 q(x^0) > 0. \end{aligned}$$

□

4.3 Projection on $\mathbb{H}^d \times \mathbb{R}^{1,d}$ of the stable manifolds

We explicit here the projection of \mathcal{V}_∞^- on $\mathbb{H}^d \times \mathbb{R}^{1,d}$. Recall that by definition an element of \mathcal{V}_∞^- is of the form $\tilde{g}_\infty \exp(X, x) \tilde{g}_\infty^{-1}$ where $(X, x) \in \bar{\mathcal{N}} \times U^+$. We deduce, since in this case $\exp(X, x) = (\exp(X), x)$, that an element of $\pi(\mathcal{V}_\infty^-)$ is of the form

$$(n_\infty \exp(X) n_\infty^{-1}(e_0), u n_\infty(e_0 + e_1) + \lambda_\infty (\text{Id} - n_\infty \exp(X) n_\infty^{-1})(e_0 - e_1)), \quad (28)$$

where X lies in $\bar{\mathcal{N}}$ and $u \in \mathbb{R}$.

Since $\exp(X)(e_0 + e_1) = e_0 + e_1$ for $X \in \bar{\mathcal{N}}$ we obtain

$$q(n_\infty \exp(X) n_\infty^{-1}(e_0), n_\infty(e_0 + e_1)) = q(n_\infty^{-1} e_0, e_0 + e_1) = q(e_0, n_\infty(e_0 + e_1))$$

and thus when X describes $\bar{\mathcal{N}}$ then $n_\infty \exp(X) n_\infty^{-1}(e_0)$ draws the intersection between \mathbb{H}^d and the affine hyperplan passing by e_0 and q -orthogonal to $n_\infty(e_0 + e_1)$. This submanifold of $\mathbb{R}^{1,d}$ is a paraboloid of codimension 2 and is mapped by \mathbf{p} (the projection onto the projective space) on a sphere tangent at $\partial \mathbb{H}^d$ in θ_∞ and passing by $\mathbf{p}(e_0)$. It is called the *horosphere* tangent at θ_∞ and passing by e_0 and is denoted by \mathcal{H}_∞ .

Moreover, since

$$q(n_\infty \exp(X) n_\infty^{-1}(e_0 - e_1), n_\infty(e_0 + e_1)) = q(e_0 - e_1, e_0 + e_1) = 0,$$

we get that when X describes \mathcal{N} then $n_\infty \exp(X) n_\infty^{-1}(e_0 - e_1)$ describes the intersection between the light cone $\{\xi, q(\xi) = 0\}$ and the hyperplan passing by $e_0 - e_1$ and q -orthogonal to $n_\infty(e_0 + e_1)$. Thus, when X describes \mathcal{N} then $(\text{Id} - n_\infty \exp(X) n_\infty^{-1})(e_0 - e_1)$ draws a paraboloid \mathcal{P}_∞ in the hyperplan q -orthogonal to $n_\infty(e_0 + e_1)$. For each $\dot{\xi}$ in the horosphere \mathcal{H}_∞ corresponds a unique $X_{\dot{\xi}} \in \mathcal{N}$ such that $\dot{\xi} = n_\infty \exp(X_{\dot{\xi}}) n_\infty^{-1}(e_0)$ and the one-to-one function $\psi : \dot{\xi} \mapsto (\text{Id} - n_\infty \exp(X_{\dot{\xi}}) n_\infty^{-1})(e_0 - e_1)$ maps \mathcal{H}_∞ on \mathcal{P}_∞ .

Then by (28), we obtain the following one-to-one map

$$\begin{aligned} \mathcal{H}_\infty \times \langle n_\infty(e_0 + e_1) \rangle &\longrightarrow \pi(\mathcal{V}_\infty^-) \\ (\dot{\xi}, \xi) &\longmapsto (\dot{\xi}, \xi + \lambda_\infty \psi(\dot{\xi})) \end{aligned}$$

and $\pi(\mathcal{V}_\infty^-)$ is a skew product of the line $\langle n_\infty(e_0 + e_1) \rangle$ with the horosphere \mathcal{H}_∞ .




Figure 2: $\pi(\mathcal{V}_\infty^-)$ is a skew-product of a horosphere with a line

References

- [1] J. Angst. *Etude de diffusions à valeurs dans des variétés lorentziennes*. PhD thesis, Université de Strasbourg, 2009.
- [2] David Applebaum and Hiroshi Kunita. Lévy flows on manifolds and Lévy processes on Lie groups. *J. Math. Kyoto Univ.*, 33(4):1103–1123, 1993.
- [3] Ismael Bailleul. Poisson boundary of a relativistic diffusion. *Probab. Theory Related Fields*, 141(1-2):283–329, 2008.
- [4] Ismael Bailleul and Albert Raugi. Where does randomness lead in spacetime? *ESAIM Probab. Stat.*, 14:16–52, 2010.
- [5] A. P. Carverhill and K. D. Elworthy. Lyapunov exponents for a stochastic analogue of the geodesic flow. *Trans. Amer. Math. Soc.*, 295(1):85–105, 1986.
- [6] R. M. Dudley. Lorentz-invariant Markov processes in relativistic phase space. *Ark. Mat.*, 6:241–268, 1966.
- [7] Jacques Franchi. Relativistic diffusion in Gödel’s universe. *Comm. Math. Phys.*, 290(2):523–555, 2009.
- [8] Jacques Franchi and Yves Le Jan. Relativistic diffusions and Schwarzschild geometry. *Comm. Pure Appl. Math.*, 60(2):187–251, 2007.
- [9] G. A. Hunt. Semi-groups of measures on Lie groups. *Trans. Amer. Math. Soc.*, 81:264–293, 1956.
- [10] Olav Kallenberg. *Foundations of modern probability*. Probability and its Applications (New York). Springer-Verlag, New York, second edition, 2002.
- [11] Ming Liao. Stable manifolds for stochastic flows induced by Lévy processes on Lie groups. *Proc. London Math. Soc. (3)*, 83(2):493–512, 2001.
- [12] Ming Liao. Dynamical properties of Lévy processes in Lie groups. *Stoch. Dyn.*, 2(1):1–23, 2002.
- [13] Ming Liao. *Lévy processes in Lie groups*, volume 162 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2004.
- [14] A. Raugi. Fonctions harmoniques sur les groupes localement compacts à base dénombrable. *Bull. Soc. Math. France Mém.*, 54:5–118, 1977.