



**HAL**  
open science

# Invariant Hilbert schemes and desingularizations of symplectic reductions for classical groups

Ronan Terpereau

► **To cite this version:**

Ronan Terpereau. Invariant Hilbert schemes and desingularizations of symplectic reductions for classical groups. 2013. hal-00799369v1

**HAL Id: hal-00799369**

**<https://hal.science/hal-00799369v1>**

Preprint submitted on 12 Mar 2013 (v1), last revised 21 Dec 2013 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# INVARIANT HILBERT SCHEMES AND DESINGULARIZATIONS OF SYMPLECTIC REDUCTIONS FOR CLASSICAL GROUPS

RONAN TERPEREAU

ABSTRACT. Let  $G \subset GL(V)$  be a classical group acting on the symplectic vector space  $W = V^{\oplus n'} \oplus V^{*\oplus n'}$ , and  $\mu : W \rightarrow Lie(G)^*$  the corresponding moment map. In this article, we use the theory of invariant Hilbert schemes to construct canonical desingularizations of the irreducible components of the symplectic reduction  $\mu^{-1}(0)//G$  for classes of examples where  $G = GL(V)$ ,  $O(V)$ , or  $Sp(V)$ . It is known that such components are isomorphic to closures of nilpotent orbits in simple Lie algebras, and we determine all the cases where the Hilbert-Chow morphism is a symplectic desingularization.

## CONTENTS

1. Introduction and statement of the main results	1
2. Generalities on symplectic varieties and closures of nilpotent orbits	4
3. Case of $GL_n$	6
4. Case of $Sp_n$ ( $n$ even)	14
References	20

## 1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

We work over the field of complex numbers  $\mathbb{C}$ . Let  $V$  and  $V'$  be two vector spaces of dimensions  $n$  and  $n'$  respectively, and let

$$(1) \quad W := \text{Hom}(V', V) \times \text{Hom}(V, V'),$$

which is naturally equipped with an action of  $GL(V') \times GL(V)$ :

$$\forall (g', g) \in GL(V') \times GL(V), \forall (u_1, u_2) \in W, (g', g) \cdot (u_1, u_2) := (g \circ u_1 \circ g'^{-1}, g' \circ u_2 \circ g^{-1}).$$

The  $GL(V)$ -module  $W$  is *symplectic*, that is, equipped with a  $GL(V)$ -invariant non-degenerate skew-symmetric bilinear form  $\Omega$  defined by:

$$(2) \quad \forall (u_1, u_2), (u'_1, u'_2) \in W, \Omega((u_1, u_2), (u'_1, u'_2)) := \text{tr}(u'_1 \circ u_2) - \text{tr}(u_1 \circ u'_2),$$

where  $\text{tr}(\cdot)$  denotes the trace. The symplectic  $GL(V)$ -module  $W$  has a  $GL(V)$ -equivariant *moment map* defined by:

$$(3) \quad \mu_{GL(V)} : \begin{array}{ccc} W & \rightarrow & \mathfrak{gl}(V)^* \\ (u_1, u_2) & \mapsto & (h \mapsto \text{tr}(u_2 \circ h \circ u_1)) \end{array}$$

where  $\mathfrak{gl}(V)$  denotes the Lie algebra of  $GL(V)$ . If  $G \subset GL(V)$  is any algebraic subgroup, and  $\mathfrak{g}$  is the Lie algebra of  $G$ , then the moment map  $\mu_G : W \rightarrow \mathfrak{g}^*$  for the action of  $G$  on  $W$  is obtained by composing  $\mu_{GL(V)}$  with the linear projection  $\mathfrak{gl}(V)^* \rightarrow \mathfrak{g}^*$ . To simplify the notation, we will denote  $\mu$  instead of  $\mu_G$ .

The map  $\mu$  is  $G$ -equivariant, and thus the set-theoretic fiber  $\mu^{-1}(0)$  is a  $G$ -stable closed subset of  $W$ . From now on, we suppose that  $G$  is reductive, and we denote  $W//G$  the *symplectic reduction* of  $W$  by  $G$ , that is, the categorical quotient  $\mu^{-1}(0)//G := \text{Spec}(\mathbb{C}[\mu^{-1}(0)]^G)$ . Let us mention that if  $G'$  is an algebraic subgroup of the  $G$ -automorphism group  $\text{Aut}^G(\mu^{-1}(0))$ , then  $G'$  acts on  $W//G$  and the quotient morphism  $\mu^{-1}(0) \rightarrow W//G$  is  $G'$ -equivariant. **The aim of this article** is to construct canonical desingularizations of the irreducible components of  $W//G$  for classes of examples where  $G = GL(V)$ ,  $O(V)$ , or  $Sp(V)$ . In these three cases, it is known that the irreducible components of  $W//G$  are closures of nilpotent orbits in simple Lie algebras of type A, C and D respectively. The geometry of such closures has been extensively studied ([KP79, KP81, KP82, Fu03a, Fu03b, Fu06b, Nam06]). In particular, the normalizations of such closures are *symplectic varieties* (as defined by Beauville in [Bea00]) whose symplectic desingularizations are the so-called *Springer desingularizations*, obtained by collapsing the cotangent bundle over some Grassmannians (see Section 2 for details).

Before stating our results, we need to recall briefly the definition of the *invariant Hilbert scheme*, constructed by Alexeev and Brion (see [AB05, Bri] for more details), in the particular case we are interested. Let  $G \subset GL(V)$  be a reductive algebraic subgroup,  $\text{Irr}(G)$  the set of isomorphism classes of irreducible representations of  $G$ , and  $h : \text{Irr}(G) \rightarrow \mathbb{N}$  a *Hilbert function*. If  $X \subset W$  a  $G$ -stable closed subscheme, then the invariant Hilbert scheme  $\text{Hilb}_h^G(X)$  is the moduli space that parametrizes the  $G$ -stable closed subschemes  $Z$  of  $X$  such that

$$\mathbb{C}[Z] \cong \bigoplus_{M \in \text{Irr}(G)} M^{\oplus h(M)}$$

as a  $G$ -module. We now suppose that  $X$  is reduced and that  $X//G$  is irreducible. If  $h = h_0$  is the Hilbert function of the general fiber of the quotient morphism  $\nu : X \rightarrow X//G$ , then there exists a projective morphism

$$\gamma : \text{Hilb}_{h_0}^G(X) \rightarrow X//G,$$

called the *Hilbert-Chow morphism*, that sends a closed subscheme  $Z \subset X$  to the point  $Z//G \subset X//G$ . If  $G' \subset \text{Aut}^G(X)$  is any algebraic subgroup, then it is known that  $G'$  acts on  $X//G$  and on  $\text{Hilb}_{h_0}^G(X)$  such that  $\nu$  and  $\gamma$  are  $G'$ -equivariant. The Hilbert-Chow morphism induces an isomorphism over the flat locus  $U \subset X//G$  of  $\nu$ . The *main component* of  $\text{Hilb}_{h_0}^G(X)$  is the irreducible component defined by

$$\text{Hilb}_{h_0}^G(X)^{\text{main}} := \overline{\gamma^{-1}(U)}.$$

Then the restriction  $\gamma : \text{Hilb}_{h_0}^G(X)^{\text{main}} \rightarrow X//G$  is a projective birational morphism, and thus  $\gamma$  is a candidate for a canonical desingularization of  $X//G$ .

We can now state the main result of this paper:

**Theorem A.** [Sections 3.3 and 4.3] *With the above notation, let  $G = GL(V)$ ,  $O(V)$  or  $Sp(V)$ , then  $W//G$  is*

- (1) *irreducible if  $G = GL(V)$ ,  $O(V)$ , or  $Sp(V)$  with  $(n' > n$  or  $n'$  odd);*
- (2) *the union of two isomorphic irreducible components if  $G = Sp(V)$  with  $(n' \leq n$  and  $n'$  even).*

*Let  $X$  denote  $\mu^{-1}(0)$  in Case (1) resp. the preimage of one of the two irreducible components of  $W//G$  by the quotient morphism  $\mu^{-1}(0) \rightarrow W//G$  in Case (2). Then*

the Hilbert-Chow morphism  $\gamma : \text{Hilb}_{h_0}^G(X)^{\text{main}} \rightarrow X//G$  is a Springer desingularization (and the unique one) if and only if

- $G = GL(V)$ ,  $n \geq n' - 1$ , and  $n'$  is even; or
- $G = O(V)$ , and  $n \geq 2n' - 1$ ; or
- $G = Sp(V)$ ,  $n$  and  $n'$  are even, and  $n \geq 2n' - 2$ .

Moreover,  $\gamma$  is still a desingularization (that strictly dominates the Springer desingularizations when they exist) if

- $G = GL(V)$  and  $((n = 1$  and  $n' \geq 3)$  or  $(n = 2$  and  $n' \geq 4))$ ; or
- $G = O(V)$  and  $(n = 1 < n'$  or  $n = 2 \leq n')$ ; or
- $G = Sp(V)$  and  $(n = 2 < n'$  or  $n = 4 \leq n')$ .

With the notation of Theorem A, we have no idea whether the Hilbert-Chow morphism  $\gamma : \text{Hilb}_{h_0}^G(X)^{\text{main}} \rightarrow X//G$  is always a desingularization or not. Let us mention that the case  $G = Sp(V)$  with  $(n = 2$  and  $n' = 3)$  was already handled by Becker in [Bec11]. In this case,  $W///G$  is a closure of a nilpotent orbit that admits two Springer desingularizations, and Becker showed that  $\gamma : \text{Hilb}_{h_0}^G(\mu^{-1}(0)) \rightarrow W///G$  is a desingularization that dominates them both. To obtain this result, Becker first used the existence of natural morphisms from the invariant Hilbert scheme to Grassmannians to identify  $\text{Hilb}_{h_0}^G(\mu^{-1}(0))^{\text{main}}$  with the total space of a homogeneous line bundle over a Grassmannian, and then she showed that  $\text{Hilb}_{h_0}^G(\mu^{-1}(0)) = \text{Hilb}_{h_0}^G(\mu^{-1}(0))^{\text{main}}$  by computing the tangent space of  $\text{Hilb}_{h_0}^G(\mu^{-1}(0))$  at every point of  $\text{Hilb}_{h_0}^G(\mu^{-1}(0))^{\text{main}}$ . In the present article, our most important theoretical result is a *reduction principle* that generalizes the first step of Becker's method.

**Reduction Principle.** [Propositions 3.8 and 4.8] *We take the notation of Theorem A, and we make the extra assumption that  $n' \geq 2n$  or  $(n' < 2n$  and  $n'$  is even) when  $G = GL(V)$ , and  $n' \geq n$  or  $(n' < n$  and  $n', n$  are even) when  $G = Sp(V)$ .*

*Then the main component  $\text{Hilb}_{h_0}^G(X)^{\text{main}}$  is the total space of a homogeneous fiber bundle over a flag variety whose fiber is isomorphic to  $\text{Hilb}_{h_{W'}}^G(W')^{\text{main}}$ , where  $W' = V^{\oplus N} \oplus V^{*\oplus N}$  with  $N := h_0(V^*)$ , and  $h_{W'}$  is the Hilbert function of the general fiber of the quotient morphism  $W' \rightarrow W'//G$ .*

A similar reduction principle (but in a different setting) was already obtained and used in [Tera]. Once we have performed the reduction step, we use [Tera, §1, Theorem] to identify the fiber  $\text{Hilb}_{h_{W'}}^G(W')^{\text{main}}$ . Unfortunately, the results that we get in [Tera] are very partial and that is why we are able to determine  $\text{Hilb}_{h_0}^G(X)^{\text{main}}$  only in the cases given by Theorem A. The last step to prove Theorem A is then to compare the Hilbert-Chow morphism  $\gamma : \text{Hilb}_{h_0}^G(X)^{\text{main}} \rightarrow X//G$  with the Springer desingularizations of  $X//G$ . We thus have a general method to determine  $\text{Hilb}_{h_0}^G(X)^{\text{main}}$  that will certainly be helpful to determine further examples when  $G = GL(V)$ ,  $O(V)$  or  $Sp(V)$ . In addition, the author thinks that the same method can be used to treat the case of  $G = SL(V)$ . However, when  $G = SO(V)$ , we do not know how to obtain a reduction principle as above.

Let now  $G \subset GL(V)$  be any reductive algebraic subgroup,  $X \subset \mu^{-1}(0)$  a maximal  $G$ -stable closed subset such that  $X//G$  is an irreducible component of  $W///G$ , and  $h_0$  the Hilbert function of the general fiber of the quotient morphism  $X \rightarrow X//G$ . Then it is generally a difficult problem to determine whether the main component

$\text{Hilb}_{h_0}^G(X)^{\text{main}}$  coincides with the whole invariant Hilbert scheme  $\text{Hilb}_{h_0}^G(X)$  or not. In this direction, we obtain

**Proposition B.** [Propositions 3.18 and 4.10] *With the notation of Theorem A, if  $G = GL(V)$  and  $n' \geq 2n$ , then the invariant Hilbert scheme  $\text{Hilb}_{h_0}^G(X)$  has at least two irreducible components (and exactly two when  $n = 1$ ). On the other hand, if  $G = O(V)$  or  $Sp(V)$ , and  $n' \geq n = 2$ , then  $\text{Hilb}_{h_0}^G(X)$  is irreducible.*

In Section 2, we recall some basic facts about symplectic varieties and closures of nilpotent orbits in simple Lie algebras. The case of  $GL(V)$  is treated in Section 3, and the case of  $Sp(V)$  is treated in Section 4. The case of  $O(V)$  is quite similar to the case of  $GL(V)$ , and details can be found in the thesis [Terb] from which this article is extracted: the Reduction Principle is [Terb, Proposition 3.4.10], Theorem A is [Terb, Corollaire 3.4.11 and Proposition A.2.5], and Proposition B is [Terb, Proposition 3.4.17].

**Acknowledgments:** I am deeply thankful to Michel Brion for proposing this subject to me, for a lot of helpful discussions, and for his patience. I thank Tanja Becker for exchange of knowledge on invariant Hilbert schemes by e-mail and during her stay in Grenoble in October 2010. I also thank Bart Van Steirteghem for helpful discussions during his stay in Grenoble in Summer 2011.

## 2. GENERALITIES ON SYMPLECTIC VARIETIES AND CLOSURES OF NILPOTENT ORBITS

**2.1. Symplectic varieties and symplectic desingularizations.** Let us first recall the definitions of symplectic variety and symplectic desingularization (see [Bea00] or the survey [Fu06a] for more details). Let  $Y$  be a normal variety whose regular part  $Y_{\text{reg}}$  admits a *symplectic form*  $\Omega$  (that is,  $\Omega$  is a holomorphic 2-form which is closed and non-degenerate at every point of  $Y_{\text{reg}}$ ) such that, for any desingularization  $f : \tilde{Y} \rightarrow Y$ , the 2-form  $f^*(\Omega)$  extends to a 2-form on the whole  $\tilde{Y}$ , then we say that  $Y$  is a *symplectic variety*. Moreover, if  $f : \tilde{Y} \rightarrow Y$  is a desingularization such that  $f^*(\Omega)$  extends to a symplectic form on  $\tilde{Y}$ , then we say that  $f$  is a *symplectic desingularization* of  $Y$ . It must be underlined that symplectic varieties do not always admit symplectic desingularizations, and when they do, there may be several of them.

Let  $W = \text{Hom}(V', V) \times \text{Hom}(V, V')$ , and  $G \subset GL(V)$  a reductive algebraic subgroup. We consider the symplectic reduction  $W///G$  as in the introduction. The following conjecture motivates the study (and the name!) of  $W///G$ :

**Conjecture 2.1** (Kaledin, Lehn, Sorger). With the above notation, the irreducible components  $Y_1, \dots, Y_r$  of  $W///G$  are symplectic varieties. Moreover, if every  $Y_i$  admits a symplectic desingularization, then the quotient  $\text{Hom}(V', V)//G$  is smooth.

When  $G$  is a finite group, Conjecture 2.1 was proved by Kaledin and Verbitsky, but the general case remains open. Let us mention that Becker showed in [Bec09] that the converse of the second part of Conjecture 2.1 holds for  $G = Sp(V)$  with  $n = 2$ . In our setting, that is when  $G = GL(V)$ ,  $O(V)$  or  $Sp(V)$ , one easily checks that Conjecture 2.1 holds (see [Terb, §A.2] for details).

**2.2. Closures of nilpotent orbits.** We now recall some basic facts concerning the closures of nilpotent orbits (see [CM93, Fu03a] for more details). Let  $\mathfrak{g}'$  be a semi-simple Lie algebra, and  $G'$  the adjoint group of  $\mathfrak{g}'$ . The following (well-known) result is due to Kostant, Kirillov, Souriau, and Panyushev:

**Theorem 2.2.** *The normalization of the closure of an adjoint orbit in a semi-simple Lie algebra is a symplectic variety.*

When  $\mathfrak{g}'$  is a simple Lie algebra of type A, C or D, one can associate partitions to nilpotent orbits. More precisely:

- If  $\mathfrak{g}' = \mathfrak{sl}_{n'}$ , then there exists a one-to-one correspondence between the partitions  $\mathbf{d} = (d_1 \geq \dots \geq d_k)$  of  $n'$  and the nilpotent orbits  $\mathcal{O}_{[d_1, \dots, d_k]}$  of  $\mathfrak{g}'$  ([CM93, §3.1]).
- If  $\mathfrak{g}' = \mathfrak{sp}_{2n'}$ , then there exists a one-to-one correspondence between the partitions  $\mathbf{d} = (d_1 \geq \dots \geq d_k)$  of  $2n'$ , in which every odd  $d_i$  occurs with even multiplicity, and the nilpotent orbits  $\mathcal{O}_{[d_1, \dots, d_k]}$  of  $\mathfrak{g}'$  ([CM93, §5.1]).
- If  $\mathfrak{g}' = \mathfrak{so}_{2n'}$ , then there exists a correspondence (which is not bijective anymore) between some partitions of  $2n'$  and the nilpotent orbits of  $\mathfrak{g}'$  ([CM93, §5.1]). Let  $\mathbf{d} = (d_1 \geq \dots \geq d_k)$  be a partition of  $2n'$  such that each even  $d_i$  occurs with even multiplicity. If at least one of the  $d_i$  is odd, then one associates to  $\mathbf{d}$  a unique nilpotent orbit  $\mathcal{O}_{[d_1, \dots, d_k]}$ ; else, one associates to  $\mathbf{d}$  two distinct nilpotent orbits  $\mathcal{O}_{[d_1, \dots, d_k]}^I$  and  $\mathcal{O}_{[d_1, \dots, d_k]}^{II}$ .

If  $\mathfrak{g}'$  is of type A, then  $\overline{\mathcal{O}_{\mathbf{d}}}$  is always normal ([KP79]). In types B, C and D, the geometry of  $\overline{\mathcal{O}_{\mathbf{d}}}$  was studied in [KP82]; in particular, if  $\mathfrak{g}'$  is of type C and  $d_1 + d_2 \leq 4$  resp. of type D and  $d_1 \leq 2$ , then  $\overline{\mathcal{O}_{\mathbf{d}}}$  is normal. From now on, we only consider partitions  $\mathbf{d}$  such that each  $d_i \leq 2$ . By Theorem 2.2, the variety  $\overline{\mathcal{O}_{\mathbf{d}}}$  is symplectic, and we are going to describe its symplectic desingularizations (see [Fu03a, Fu06b, FN04] for details).

Let  $f : Z \rightarrow \overline{\mathcal{O}_{\mathbf{d}}}$  be a symplectic desingularization. Then, by [Fu03a, Proposition 3.1], the group  $G'$  acts naturally on  $Z$  in such a way that  $f$  is  $G'$ -equivariant. One says that  $f$  is a *Springer desingularization* if there exists a parabolic subgroup  $P \subset G'$  and a  $G'$ -equivariant isomorphism between  $Z$  and the total space of the cotangent bundle over  $G'/P$ , denoted  $\mathcal{T}^*(G'/P)$ . Then, under this isomorphism, the map  $f$  becomes

$$\mathcal{T}^*(G'/P) \cong G' \times^P \mathfrak{u} \longrightarrow \mathfrak{g}', \quad (g', x) \longmapsto Ad(g').x,$$

where  $\mathfrak{u}$  is the nilradical of the Lie algebra of  $P$ , and  $G' \times^P \mathfrak{u} := (G' \times \mathfrak{u}) / \equiv$  with  $(g', u) \equiv (g' \circ p^{-1}, Ad(p).u)$ .

**Theorem 2.3.** ([Fu03a, Theorem 3.3]) *With the above notation, if  $f : Z \rightarrow \overline{\mathcal{O}_{\mathbf{d}}}$  is a symplectic desingularization, then  $f$  is a Springer desingularization.*

When  $\mathfrak{g}'$  is of type A, C or D, the Springer desingularizations of  $\overline{\mathcal{O}_{\mathbf{d}}}$  are known (up to isomorphism):

- Let  $\mathfrak{g}' = \mathfrak{sl}_{n'}$  and  $\mathbf{d} = [2^N, 1^{n'-2N}]$ . We denote  $\text{Gr}(p, \mathbb{C}^{n'})$  the Grassmannian of  $p$ -dimensional subspaces of  $\mathbb{C}^{n'}$ , and  $\mathcal{T}_1^*$  resp.  $\mathcal{T}_2^*$ , the cotangent bundle over  $\text{Gr}(N, \mathbb{C}^{n'})$  resp. over  $\text{Gr}(n' - N, \mathbb{C}^{n'})$ . By [Fu06b, §2], if  $N < \frac{n'}{2}$ , then  $\mathcal{T}_1^*$  and  $\mathcal{T}_2^*$  are the two Springer desingularizations of  $\overline{\mathcal{O}_{\mathbf{d}}}$ ; else,  $\mathcal{T}_1^* = \mathcal{T}_2^*$  is the unique Springer desingularization of  $\overline{\mathcal{O}_{\mathbf{d}}}$ .

- Let  $\mathfrak{g}' = \mathfrak{sp}_{2n'}$  and  $\mathbf{d} = [2^N, 1^{2(n'-N)}]$ . Then  $\overline{\mathcal{O}_{\mathbf{d}}}$  admits a Springer desingularization if and only if  $N = n'$  ([Fu03a, Proposition 3.19]). We denote  $\text{IGr}(p, \mathbb{C}^{2n'})$  the Grassmannian of isotropic  $p$ -dimensional subspaces of  $\mathbb{C}^{2n'}$ , and  $\mathcal{T}^*$  the cotangent bundle over  $\text{IGr}(n', \mathbb{C}^{2n'})$ . By [FN04, Proposition 3.5], if  $N = n'$ , then  $\mathcal{T}^*$  is the unique Springer desingularization of  $\overline{\mathcal{O}_{\mathbf{d}}}$ .
- Let  $\mathfrak{g}' = \mathfrak{so}_{2n'}$  and  $\mathbf{d} = [2^N, 1^{2(n'-N)}]$  (with  $N$  even). We recall that if  $d_1 = \dots = d_k = 2$ , then one associates to  $\mathbf{d}$  two distinct nilpotent orbits  $\mathcal{O}_{\mathbf{d}}^I$  and  $\mathcal{O}_{\mathbf{d}}^{II}$ . By [Fu03a, Proposition 3.20], the variety  $\overline{\mathcal{O}_{\mathbf{d}}}$  admits a Springer desingularization if and only if  $N \in \{n' - 1, n'\}$ . We denote  $\text{OGr}(p, \mathbb{C}^{2n'})$  the Grassmannian of isotropic  $p$ -dimensional subspaces of  $\mathbb{C}^{2n'}$ . The Grassmannian  $\text{OGr}(p, \mathbb{C}^{2n'})$  is irreducible except if  $p = n'$ , in which case  $\text{OGr}(n', \mathbb{C}^{2n'}) = \text{OG}^I \cup \text{OG}^{II}$  is the union of two irreducible components (exchanged by the natural action of the orthogonal group  $O(\mathbb{C}^{2n'})$ ). We denote  $\mathcal{T}_I^*$  resp.  $\mathcal{T}_{II}^*$ , the cotangent bundle over  $\text{OG}^I$ , resp. over  $\text{OG}^{II}$ . If  $N = n' - 1$ , then  $\mathcal{T}_I^*$  and  $\mathcal{T}_{II}^*$  are the two Springer desingularizations of  $\overline{\mathcal{O}_{[2^{n'-1}, 1^2]}}$  by [Fu06b, §2]. If  $N = n'$ , then  $\mathcal{T}_I^*$  resp.  $\mathcal{T}_{II}^*$ , is the unique Springer desingularization of  $\overline{\mathcal{O}_{[2^{n'}]}}^I$  resp. of  $\overline{\mathcal{O}_{[2^{n'}]}}^{II}$ , by [FN04, Proposition 3.5].

*Remark 2.4.* We will see in Sections 3 and 4 that the irreducible components of the symplectic reduction  $W//G$ , when  $G = GL(V)$  or  $Sp(V)$ , are closures of nilpotent orbits  $\mathcal{O}_{\mathbf{d}}$  in simple Lie algebras of type A and C respectively, where every  $d_i \leq 2$ . If  $G = O(V)$ , then it is shown in [Terb, §3.4] that  $W//G$  is the closure of the nilpotent orbit  $\mathcal{O}_{[2^N, 1^{2(n'-N)}]}$  in type D, where  $N := \min(n', n)$ .

### 3. CASE OF $GL_n$

In this section, we take  $G = GL(V)$  and  $G' = GL(V')$ , both acting on  $W = \text{Hom}(V', V) \times \text{Hom}(V, V')$  as in the introduction. We denote  $\mathfrak{g}$  resp.  $\mathfrak{g}'$ , the Lie algebra of  $G$  resp. of  $G'$ , and  $N := \min(E(\frac{n'}{2}), n)$ , where  $n := \dim(V)$ ,  $n' := \dim(V')$ , and  $E(\cdot)$  is the lower integer part.

**3.1. The quotient morphism.** The two main results of this section are Proposition 3.3, which describes the symplectic reduction  $W//G := \mu^{-1}(0)//G$ , and Corollary 3.6, which gives the Hilbert function  $h_0$  of the general fiber of the quotient morphism  $\nu: \mu^{-1}(0) \rightarrow W//G$ .

By (3), the zero fiber of the moment map  $\mu: W \rightarrow \mathfrak{g}^*$  is

$$\mu^{-1}(0) = \{(u_1, u_2) \in W \mid u_1 \circ u_2 = 0\},$$

which is a  $G' \times G$ -stable closed subset of  $W$ . Let us determine the irreducible components of  $\mu^{-1}(0)$  as well as their dimensions. Let  $m \in \{0, \dots, n'\}$ ; we define the closed subset

$$(4) \quad X_m := \left\{ (u_1, u_2) \in W \mid \begin{array}{l} \text{Im}(u_2) \subset \text{Ker}(u_1); \\ \text{rk}(u_2) \leq \min(n, m); \\ \dim(\text{Ker}(u_1)) \geq \max(n' - n, m). \end{array} \right\} \subset \mu^{-1}(0),$$

and we consider the diagram

$$\begin{array}{ccc}
 Z_m := \{(u_1, u_2, L) \in W \times \text{Gr}(m, V') \mid \text{Im}(u_2) \subset L \subset \text{Ker}(u_1)\} & & \\
 p_1 \downarrow & \searrow p_2 & \\
 X_m & & \text{Gr}(m, V')
 \end{array}$$

where the  $p_i$  are the natural projections. We fix  $L_0 \in \text{Gr}(m, V')$ ; the second projection equips  $Z_m$  with a structure of homogeneous vector bundle over  $\text{Gr}(m, V')$  whose fiber over  $L_0$  is isomorphic to  $F_m := \text{Hom}(V'/L_0, V) \times \text{Hom}(V, L_0)$ . Hence,  $Z_m$  is a smooth variety of dimension  $m(n' - m) + n'n$ .

**Proposition 3.1.** *The irreducible components of  $\mu^{-1}(0)$  are*

$$\begin{cases} X_0, \dots, X_{n'} & \text{if } n' \leq n; \\ X_{n'-n}, \dots, X_n & \text{if } n < n' < 2n; \\ X_n & \text{if } n' \geq 2n; \end{cases}$$

where  $X_m$  is defined by (4).

*Proof.* We have

$$\mu^{-1}(0) = \{(u_1, u_2) \in W \mid \text{Im}(u_2) \subset \text{Ker}(u_1)\} = \bigcup_{i=0}^{n'} X_i.$$

Furthermore, for every  $m \in \{0, \dots, n'\}$ , the morphism  $p_1$  is surjective and  $Z_m$  is irreducible, hence  $X_m$  is irreducible.

If  $n' \geq 2n$ , then

$$\begin{cases} X_0 \subset \dots \subset X_n; \\ X_n = \dots = X_{n'-n}; \\ X_{n'-n} \supset \dots \supset X_{n'}; \end{cases}$$

and thus  $\mu^{-1}(0) = X_n$ .

If  $n' < 2n$ , then

$$\begin{cases} X_0 \subset \dots \subset X_{\max(0, n'-n)}; \\ X_{\min(n', n)} \supset \dots \supset X_{n'}; \end{cases}$$

and one easily checks that there is no other inclusion relation between the  $X_m$ .  $\square$

**Corollary 3.2.** *The dimension of  $\mu^{-1}(0)$  is*

$$\dim(\mu^{-1}(0)) = \begin{cases} nn' + \frac{1}{4}n'^2 & \text{if } n' < 2n \text{ and } n' \text{ is even;} \\ nn' + \frac{1}{4}(n'^2 - 1) & \text{if } n' < 2n \text{ and } n' \text{ is odd;} \\ 2nn' - n^2 & \text{if } n' \geq 2n. \end{cases}$$

*Proof.* By Proposition 3.1, it is sufficient to compute the dimension of  $X_m$  for some  $m$ . If  $m \leq n$  or  $m \geq n' - n$ , one may check that the map  $p_1 : Z_m \rightarrow X_m$  is birational, and thus  $P(m) := \dim(X_m) = \dim(Z_m) = m(n' - m) + n'n$ . It remains simply to study the variations of the polynomial  $P$  to obtain the result.  $\square$



Let  $l \in \{0, \dots, N\}$ . We fix a basis  $\mathcal{B}$  of  $V$  resp.  $\mathcal{B}'$  of  $V'$ , and we introduce some notation that we will use in the proofs of Proposition 3.3 and Lemma 3.5:

$$(5) \quad \bullet (u_1^l, u_2^l) := \left( \begin{bmatrix} 0_{l, n'-l} & I_l \\ 0_{n-l, n'-l} & 0_{n-l, l} \end{bmatrix}, \begin{bmatrix} I_l & 0_{l, n-l} \\ 0_{n'-l, l} & 0_{n'-l, n-l} \end{bmatrix} \right) \in W;$$

$$(6) \quad \bullet f_l := \begin{bmatrix} 0_{l, n'-l} & I_l \\ 0_{n'-l, n'-l} & 0_{n'-l, l} \end{bmatrix} \in \mathfrak{g}'.$$

**Proposition 3.3.** *Let  $\mathcal{O}_{\mathbf{d}} \subset \mathfrak{g}' \cong \mathfrak{gl}_{n'}$  denote the nilpotent orbit associated to a partition  $\mathbf{d}$  of  $n'$ . Then the symplectic reduction of  $W$  by  $G$  is  $W///G = \overline{\mathcal{O}_{[2N, 1^{n'-2N}]}}$ .*

*Proof.* If  $f \in W///G$ , then there exists  $(u_1, u_2) \in \mu^{-1}(0)$  such that  $f = u_2 \circ u_1$ , and thus  $f \circ f = (u_2 \circ u_1) \circ (u_2 \circ u_1) = u_2 \circ (u_1 \circ u_2) \circ u_1 = 0$ , whence the inclusion " $\subset$ ". Now, let  $f \in \overline{\mathcal{O}_{[2N, 1^{n'-2N}]}}$ . Up to conjugation by an element of  $G'$ , we can suppose that  $f = f_l$  for some  $l \leq N$ , where  $f_l$  is defined by (6). But then  $u_2^l \circ u_1^l = f_l$  and  $u_1^l \circ u_2^l = 0$ , where  $u_1^l$  and  $u_2^l$  are defined by (5), and thus  $f \in W///G$ .  $\square$

**Corollary 3.4.** *The symplectic reduction  $W///G \subset \mathfrak{g}'$  decomposes into  $N+1$  orbits for the adjoint action of  $G'$ :*

$$U_i := \mathcal{O}_{[2^i, 1^{n'-2i}]}, \quad \text{for } i = 0, \dots, N.$$

The closures of the nilpotent orbits  $U_i$  are nested in the following way:

$$\{0\} = \overline{U_0} \subset \dots \subset \overline{U_N} = W///G.$$

Hence,  $W///G$  is a symplectic variety (see Section 2), of dimension  $2N(n' - N)$  ([CM93, Corollary 6.1.4]), and whose singular locus is  $\overline{U_{N-1}}$  ([KP81, §3.2]).

By Corollary 3.2, the dimension of the general fiber of the quotient morphism  $\nu$  is

$$(7) \quad \begin{cases} nn' - \frac{1}{4}n'^2 & \text{if } n' < 2n \text{ and } n' \text{ is even;} \\ nn' - \frac{1}{4}(n'^2 - 1) & \text{if } n' < 2n \text{ and } n' \text{ is odd;} \\ n^2 & \text{if } n' \geq 2n. \end{cases}$$

If  $n' < 2n$ , then  $N = E(\frac{n'}{2})$ , and we denote

$$(8) \quad H := \left\{ \begin{bmatrix} M & 0_{n-N, N} \\ 0_{N, n-N} & I_N \end{bmatrix}, M \in GL_{n-N} \right\} \cong GL_{n-N},$$

which is a reductive algebraic subgroup of  $G \cong GL_n$ .

**Proposition 3.5.** *The general fiber of the quotient morphism  $\nu : \mu^{-1}(0) \rightarrow W///G$  is isomorphic to*

$$\begin{cases} G & \text{if } n' \geq 2n; \\ G/H & \text{if } n' < 2n \text{ and } n' \text{ is even;} \end{cases}$$

where  $H \subset G$  is the subgroup defined by (8).

*Proof.* We first suppose that  $n' < 2n$  and  $n'$  is even (that is,  $N = \frac{n'}{2}$ ). With the notation (5), and by [SB00, §I.6.2.5, Theorem 10], we have the equivalence

$$G.(u_1^N, u_2^N) \text{ is closed in } \mu^{-1}(0) \Leftrightarrow C_G(H).(u_1^N, u_2^N) \text{ is closed in } \mu^{-1}(0).$$

We have  $C_G(H) = \left\{ \begin{bmatrix} M & 0 \\ 0 & \lambda I_{n-N} \end{bmatrix}, M \in GL_N, \lambda \in \mathbb{G}_m \right\}$ , where  $\mathbb{G}_m$  denotes the multiplicative group. Hence

$$C_G(H) \cdot (u_1^N, u_2^N) = \left\{ \left( \begin{bmatrix} 0 & M \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} M^{-1} & 0 \\ 0 & 0 \end{bmatrix} \right), M \in GL_N \right\} \subset \mu^{-1}(0)$$

is a closed subset, and thus  $G \cdot (u_1^N, u_2^N)$  is the unique closed orbit contained in the fiber  $\nu^{-1}(f_N)$ , where  $f_N$  is defined by (6). One may check that  $\text{Stab}_G((u_1^N, u_2^N)) = H$ . Furthermore,  $\dim(G/H) = N(2n - N)$ , which is also the dimension of the general fiber of  $\nu$  by (7), and thus  $\nu^{-1}(f_N) \cong G/H$ .

We now suppose that  $n' \geq 2n$  (that is,  $N = n$ ). One may check that  $\text{Stab}_G((u_1^{n'}, u_2^{n'})) = Id$ , and thus the fiber  $\nu^{-1}(f_n)$  contains a unique closed orbit isomorphic to  $G$ . But  $\dim(G) = n^2$  is the dimension of the general fiber of  $\nu$  by (7), hence  $\nu^{-1}(f_n) \cong G$ .  $\square$

If  $n' < 2n$  and  $n'$  is odd, then the situation is more complicated (except the case  $n' = 1$  which is trivial) because the general fiber of the quotient morphism  $\nu$  is reducible. From now on, we will only consider the cases where  $n' \geq 2n$  or ( $n' < 2n$  and  $n'$  is even).

**Corollary 3.6.** *The Hilbert function  $h_0$  of the general fiber of the quotient morphism  $\nu: \mu^{-1}(0) \rightarrow W//G$  is given by:*

$$\forall M \in \text{Irr}(G), h_0(M) = \begin{cases} \dim(M) & \text{if } n' \geq 2n; \\ \dim(M^H) & \text{if } n' < 2n \text{ and } n' \text{ is even;} \end{cases}$$

where  $H \subset G$  is the subgroup defined by (8).

**3.2. The reduction principle for the main component.** In this section we prove the reduction principle when  $G = GL(V)$  (Proposition 3.8).

The subset  $\mu^{-1}(0) \subset W$  being  $G' \times G$ -stable, it follows from [Bri, Lemma 3.3] that the invariant Hilbert scheme

$$\mathcal{H} := \text{Hilb}_{h_0}^G(\mu^{-1}(0))$$

is a  $G'$ -stable closed subscheme of  $\text{Hilb}_{h_0}^G(W)$ . The scheme  $\text{Hilb}_{h_0}^G(W)$  was studied in [Tera]; let us recall

**Proposition 3.7.** ([Tera, §4.4]) *Let  $h_0$  be the Hilbert function given by Corollary 3.6, and  $G' = GL(V')$  acting naturally on  $\text{Gr}(n' - h_0(V), V'^*) \times \text{Gr}(n' - h_0(V^*), V')$ . Then there exists a  $G'$ -equivariant morphism*

$$\begin{array}{ccc} \rho : \text{Hilb}_{h_0}^G(W) & \rightarrow & \text{Gr}(n' - h_0(V), V'^*) \times \text{Gr}(n' - h_0(V^*), V') \\ Z & \mapsto & (\text{Ker}(f_Z^1), \text{Ker}(f_Z^2)) \end{array}$$

where  $f_Z^1: V'^* \cong \text{Mor}^G(W, V) \rightarrow \text{Mor}^G(Z, V)$  and  $f_Z^2: V' \cong \text{Mor}^G(W, V^*) \rightarrow \text{Mor}^G(Z, V^*)$  are the restriction maps.

By Corollary 3.6, we have  $h_0(V) = h_0(V^*) = N$ . We identify  $\text{Gr}(n' - N, V'^*)$  with  $\text{Gr}(N, V')$ , and we denote

$$A_i := \{(L_1, L_2) \in \text{Gr}(N, V') \times \text{Gr}(n' - N, V') \mid \dim(L_1 \cap L_2) = N - i\}, \text{ for } i = 0, \dots, N.$$

The  $A_i$  are the  $N + 1$  orbits for the action of  $G'$  on  $\text{Gr}(N, V') \times \text{Gr}(n' - N, V')$ , and

$$A_0 = \overline{A_0} \subset \overline{A_1} \subset \dots \subset \overline{A_N} = \text{Gr}(N, V') \times \text{Gr}(n' - N, V').$$

In particular,  $A_N$  is the unique open orbit and

$$(9) \quad A_0 = \mathcal{F}_{N, n'-N} := \{(L_1, L_2) \in \text{Gr}(N, V') \times \text{Gr}(n' - N, V') \mid L_1 \subset L_2\},$$

which is a partial flag variety, is the unique closed orbit. Let

- $a_0 := (L_1, L_2) \in A_0$ , and  $P$  the parabolic subgroup of  $G'$  stabilizing  $a_0$ ;
- $W' := \{(u_1, u_2) \in W \mid L_2 \subset \text{Ker}(u_1) \text{ and } \text{Im}(u_2) \subset L_1\}$ , which is a  $P \times G'$ -module contained in  $\mu^{-1}(0)$ ;
- $\mathcal{H}' := \text{Hilb}_{h_0}^G(W')$ , and  $\mathcal{H}'^{\text{main}}$  its main component.

Let us note that, if  $n' \geq 2n$  or ( $n' < 2n$  and  $n'$  is even), then  $h_0$  coincides with the Hilbert function of the general fiber of the quotient morphism  $W' \rightarrow W'//G$  by [Tera, Proposition 4.13]; in particular,  $\mathcal{H}'^{\text{main}}$  is well-defined. We are going to show

**Proposition 3.8.** *If  $n' \geq 2n$  or ( $n' < 2n$  and  $n'$  is even), and with the above notation, we have a  $G'$ -equivariant isomorphism*

$$\mathcal{H}^{\text{main}} \cong G' \times^P \mathcal{H}'^{\text{main}}.$$

First of all, we need

**Lemma 3.9.** *If  $n' \geq 2n$  or ( $n' < 2n$  and  $n'$  is even), then the morphism  $\rho$  of Proposition 3.7 sends  $\mathcal{H}^{\text{main}}$  onto  $A_0$ , the  $G'$ -variety defined by (9).*

*Proof.* As the quotient morphism  $\nu : \mu^{-1}(0) \rightarrow W'//G$  is flat over the open orbit  $U_N$ , the restriction of the Hilbert-Chow morphism  $\gamma$  to  $\gamma^{-1}(U_N)$  is an isomorphism. We fix  $f_N \in U_N$ , and we denote  $Q := \text{Stab}_{G'}(f_N)$ , and  $Z_N$  the unique point of  $\mathcal{H}$  such that  $\gamma(Z_N) = f_N$ . As  $\gamma$  is  $G'$ -equivariant,  $Z_N$  is  $Q$ -stable. In addition,  $\rho$  is also  $G'$ -equivariant, hence  $\rho(Z_N)$  is a fixed point for the action of  $Q$ . But one may check that  $\text{Gr}(N, V') \times \text{Gr}(n' - N, V')$  has a unique fixed point for  $Q$ , which is contained in  $A_0$ . Then, as  $A_0$  is  $G'$ -stable, we have  $\rho(Z) \in A_0$ , for every  $Z \in \gamma^{-1}(U_N)$ . Hence,  $\rho^{-1}(A_0)$  is a closed subset of  $\text{Hilb}_{h_0}^G(W)$  containing  $\gamma^{-1}(U_N)$ , and the result follows.  $\square$

The restriction  $\rho|_{\mathcal{H}^{\text{main}}} : \mathcal{H}^{\text{main}} \rightarrow A_0$  is  $G'$ -equivariant, hence  $\mathcal{H}^{\text{main}}$  is the total space of a  $G'$ -homogeneous fiber bundle over  $A_0$ . Let  $F$  be the scheme-theoretic fiber of  $\rho|_{\mathcal{H}^{\text{main}}}$  over  $a_0$ . The action of  $P$  on  $\mathcal{H}^{\text{main}}$ , induced by the action of  $G'$ , stabilizes  $F$ , and we have a  $G'$ -equivariant isomorphism

$$(10) \quad \mathcal{H}^{\text{main}} \cong G' \times^P F.$$

Hence, to show Proposition 3.8, we have to determine  $F$  as a  $P$ -scheme. We start by considering  $F'$ , the scheme-theoretic fiber of the restriction  $\rho|_{\mathcal{H}} : \mathcal{H} \rightarrow \text{Gr}(N, V') \times \text{Gr}(n' - N, V')$  over  $a_0$ , as a  $P$ -scheme. The proof of the next lemma is analogous to the proof of [Tera, Lemma 3.7].

**Lemma 3.10.** *With the above notation, we have a  $P$ -equivariant isomorphism*

$$F' \cong \mathcal{H}',$$

where  $P$  acts on  $\mathcal{H}'$  via its action on  $W'$ .

As  $\mathcal{H}^{\text{main}}$  is a variety of dimension  $2N(n' - N)$ , we deduce that  $F$  is a variety of dimension  $N^2$ . By Lemma 3.10, the fiber  $F$  is isomorphic to a subvariety of  $\mathcal{H}'^{\text{main}}$ , but  $\dim(\mathcal{H}'^{\text{main}}) = N^2$ , and thus we have a  $P$ -equivariant isomorphism

$$(11) \quad F \cong \mathcal{H}'^{\text{main}},$$

and Proposition 3.8 follows.

*Remark 3.11.* The scheme  $\mathcal{H}'$  is  $P$ -stable and identifies with a closed subscheme of  $\mathcal{H}$ , hence we have an inclusion of  $G'$ -schemes  $G' \times^P \mathcal{H}' \subset \mathcal{H}$ .

**3.3. Proof of Theorem A for  $GL(V)$ .** First, we recall the following result:

**Theorem 3.12.** [Tera, §1, Theorem] *Let  $G = GL(V)$ ,  $W = \text{Hom}(V', V) \times \text{Hom}(V, V')$ , and  $h_W$  the Hilbert function of the general fiber of the quotient morphism  $W \rightarrow W//G$ . We denote  $n := \dim(V)$ ,  $n' := \dim(V')$ , and  $Y_0$  the blow-up of  $W//G = \text{End}(V')^{\leq n}$  at 0. In the following cases, the invariant Hilbert scheme  $\mathcal{H}' := \text{Hilb}_{h_W}^G(W)$  is a smooth variety and the Hilbert-Chow morphism is the succession of blow-up described as follows:*

- if  $n \geq 2n' - 1$ , then  $\mathcal{H}' \cong W//G = \text{End}(V')$ ;
- if  $n' > n = 1$  or  $n' = n = 2$ , then  $\mathcal{H}' \cong Y_0$ ;
- if  $n' > n = 2$ , then  $\mathcal{H}'$  is isomorphic to the blow-up of  $Y_0$  along the strict transform of  $\text{End}(V')^{\leq 1}$ .

Let us now consider the following diagram

$$(12) \quad \begin{array}{ccc} & \mathcal{F}_{N, n'-N} & \\ p_1 \swarrow & & \searrow p_2 \\ \text{Gr}(N, V') & & \text{Gr}(n'-N, V') \end{array}$$

where  $\mathcal{F}_{N, n'-N}$  is defined by (9),  $p_1$  and  $p_2$  being the natural projections. We denote  $\underline{V}'$  the constant vector bundle over  $\mathcal{F}_{N, n'-N}$  with fiber  $V'$ , and  $T_1$  resp.  $T_2$ , the pull-back of the tautological bundle over  $\text{Gr}(N, V')$  by  $p_1$ , resp. over  $\text{Gr}(n'-N, V')$  by  $p_2$ . In particular, if  $N = \frac{n'}{2}$ , then  $\mathcal{F}_{N, n'-N} = \text{Gr}(N, V')$  and  $T := T_1 = T_2$  is the tautological bundle over  $\text{Gr}(N, V')$ .

We deduce from Proposition 3.8 and Theorem 3.12 that we have the following  $G'$ -equivariant isomorphisms

$$(13) \quad \mathcal{H}^{\text{main}} \cong \begin{cases} \text{Hom}(\underline{V}'/T, T) & \text{if } n \geq n' - 1 \text{ and } n' \text{ is even;} \\ \text{Hom}(\underline{V}'/T_2, T_1) & \text{if } n = 1 \text{ and } n' \geq 3; \\ \text{Bl}_0(\text{Hom}(\underline{V}'/T_2, T_1)) & \text{if } n = 2 \text{ and } n' \geq 4; \end{cases}$$

where  $\text{Bl}_0(\cdot)$  denotes the blow-up along the zero section. In all these cases,  $\mathcal{H}^{\text{main}}$  is smooth, and thus the Hilbert-Chow morphism  $\gamma : \mathcal{H}^{\text{main}} \rightarrow W//G$  is a (distinguished) desingularization.

On the other hand, we saw in Section 2 that the Springer desingularizations of  $W//G$  are the cotangent bundles  $\mathcal{T}_1^* := \mathcal{T}^* \text{Gr}(N, V')$  and  $\mathcal{T}_2^* := \mathcal{T}^* \text{Gr}(n'-N, V') \cong \mathcal{T}^* \text{Gr}(N, V'^*)$ . We then distinguish between two cases:

- (1) If  $N < \frac{n'}{2}$ , then let us show by contradiction that  $\gamma : \mathcal{H}^{\text{main}} \rightarrow W//G$  can not be a Springer desingularization. First, we consider the isomorphism of  $G' \times G$ -modules

$$W \rightarrow \text{Hom}(V'^*, V^*) \times \text{Hom}(V^*, V'^*), \quad (u_1, u_2) \mapsto (u_2^*, u_1^*),$$

where  $u_\bullet^*$  denotes the pull-back of  $u_\bullet$ . Denoting  $\mathcal{H}_0 := \text{Hilb}_{h_0}^G(\mu'^{-1}(0))$ , where  $\mu'$  is the moment map for the natural action of  $G$  on  $\text{Hom}(V'^*, V^*) \times \text{Hom}(V^*, V'^*)$ , we have an isomorphism of  $G'$ -varieties  $\mathcal{H}^{\text{main}} \cong \mathcal{H}_0^{\text{main}}$ . Now if we suppose that (say)  $\mathcal{H}^{\text{main}} \cong \mathcal{T}_1^*$ , then we get that  $\mathcal{H}_0^{\text{main}} \cong \mathcal{T}_2^*$ , and thus  $\mathcal{T}_1^* \cong \mathcal{T}_2^*$  as a  $G'$ -variety, which is absurd.

However, one easily checks that if  $n \in \{1, 2\}$  and  $n' \geq 2n + 1$ , then  $\gamma :$

$\mathcal{H}^{\text{main}} \rightarrow W//G$  dominates the two Springer desingularizations  $\mathcal{T}_1^*$  and  $\mathcal{T}_2^*$  (see [Terb, §A.2.2] for details).

- (2) If  $N = \frac{n'}{2}$ , then  $\mathcal{T}^* := \mathcal{T}_1^* = \mathcal{T}_2^*$  is the unique Springer desingularization of  $W//G$ . Let us show that  $\gamma: \mathcal{H}^{\text{main}} \rightarrow W//G$  is a Springer desingularization if and only if  $n \geq n' - 1$ . The implication " $\Leftarrow$ " is given by (13) since  $\mathcal{T}^* \cong \text{Hom}(\underline{V}'/T, T)$ . The other implication is given by:

**Lemma 3.13.** *If  $N = \frac{n'}{2}$  and the Hilbert-Chow morphism  $\gamma: \mathcal{H}^{\text{main}} \rightarrow W//G$  is a Springer desingularization, then  $n \geq n' - 1$ .*

*Proof.* We suppose that  $\gamma: \mathcal{H}^{\text{main}} \rightarrow W//G$  is a Springer desingularization, that is,  $\mathcal{H}^{\text{main}} \cong \mathcal{T}^*$  as a  $G'$ -variety. We fix  $L \in \text{Gr}(N, V')$ , and we define  $P \subset G'$ ,  $W'$ , and  $\mathcal{H}'^{\text{main}}$  as in Section 3.2. We have  $\mathcal{T}^* \cong G' \times^P \text{Hom}(V'/L, L)$ , and it follows from (10) and (11) that  $\mathcal{H}^{\text{main}} \cong G' \times^P \mathcal{H}'^{\text{main}}$ . Hence,  $\mathcal{H}'^{\text{main}} \cong \text{Hom}(V'/L, L)$  as a  $P$ -variety. We denote  $\gamma': \mathcal{H}'^{\text{main}} \rightarrow W'//G$  the restriction of the Hilbert-Chow morphism. As  $\gamma'$  is projective and birational, and  $W'//G = \text{Hom}(V'/L, L)$  is smooth, the Zariski's Main Theorem implies that  $\gamma'$  is an isomorphism. It follows that the quotient morphism  $\nu': W' \rightarrow W'//G$  is flat, and thus  $n \geq 2N - 1$  by [Tera, Corollary 4.12].  $\square$

In addition, if  $n' = 4$  and  $n = 2$ , then by (13) we have  $\mathcal{H}^{\text{main}} \cong \text{Bl}_0(\mathcal{T}^*)$ , and thus  $\gamma$  dominates the unique Springer desingularization of  $W//G$ .

**3.4. Reducibility of the invariant Hilbert scheme.** The aim of this section is to show Proposition B for  $GL(V)$ , that is, to show

**Proposition 3.14.** *If  $n' \geq 2n$ , then the invariant Hilbert scheme  $\mathcal{H}$  is reducible.*

We suppose that  $n' \geq 2n$ , then  $N = n$ . We fix

$$(14) \quad a_n = (L'_1, L'_2) \in A_n$$

a point of the open  $G'$ -orbit of  $\text{Gr}(n, V') \times \text{Gr}(n' - n, V')$ , and we consider

$$\begin{aligned} W'' &:= \{(u_1, u_2) \in W \mid L'_2 \subset \text{Ker}(u_1) \text{ and } \text{Im}(u_2) \subset L'_1\} \\ &\cong \text{Hom}(V'/L'_2, V) \times \text{Hom}(V, L'_1), \end{aligned}$$

which is a  $G$ -submodule of  $W$ . As  $V' = L'_1 \oplus L'_2$ , we have a natural identification  $W'' \cong \text{Hom}(L'_1, V) \times \text{Hom}(V, L'_1)$  as a  $G$ -module. We saw at the beginning of Section 1 that the  $G$ -module  $W''$  is symplectic and has a  $G$ -equivariant moment map that we denote  $\mu': W'' \rightarrow \mathfrak{g}^*$ . The proof of the next lemma is analogous to the proof of [Tera, Lemma 3.7].

**Lemma 3.15.** *We suppose that  $n' \geq 2n$ , and let  $\rho: \mathcal{H} \rightarrow \text{Gr}(n, V') \times \text{Gr}(n' - n, V')$  be the morphism of Proposition 3.7. The scheme-theoretic fiber  $F''$  of  $\rho$  over the point  $a_n$ , defined by (14), is isomorphic to the invariant Hilbert scheme  $\text{Hilb}_{h_0}^G(\mu'^{-1}(0))$ , where  $h_0$  is the Hilbert function defined by  $h_0(M) = \dim(M)$ , for every  $M \in \text{Irr}(G)$ , and  $\mu': W'' \rightarrow \mathfrak{g}^*$  is the moment map defined above.*

*Remark 3.16.* The Hilbert function  $h_0$  of Lemma 3.15 does not generally coincide with the Hilbert function of the general fiber of the quotient morphism  $\mu'^{-1}(0) \rightarrow \mu'^{-1}(0)//G$ .

By Lemma 3.9, the morphism  $\rho : \text{Hilb}_{h_0}^G(W) \rightarrow \text{Gr}(n, V') \times \text{Gr}(n' - n, V')$  of Proposition 3.7 sends  $\mathcal{H}^{\text{main}}$  onto  $A_0$ . Hence, to show Proposition 3.14, it is enough, by Lemma 3.15, to show that  $\text{Hilb}_{h_0}^G(\mu'^{-1}(0))$  is non-empty. We denote  $V'' := L'_1$ , and we equip  $W'' \cong \text{Hom}(V'', V) \times \text{Hom}(V, V'')$  with the natural action of  $G'' := GL(V'')$ . Then we have

$$(15) \quad \begin{aligned} \mathbb{C}[W'']_2 \cong & (S^2(V'') \otimes S^2(V^*)) \oplus (S^2(V''^*) \otimes S^2(V)) \\ & \oplus (\Lambda^2(V'') \otimes \Lambda^2(V^*)) \oplus (\Lambda^2(V''^*) \otimes \Lambda^2(V)) \\ & \oplus ((sl(V'') \oplus M_0) \otimes (sl(V) \oplus V_0)) \text{ as a } G'' \times G\text{-module,} \end{aligned}$$

where  $M_0$  resp.  $V_0$ , is the trivial  $G''$ -module resp. the trivial  $G$ -module, and  $sl(V'') := \{f \in \text{End}(V'') \mid \text{tr}(f) = 0\}$ .

We denote  $I_0$  the ideal of  $\mathbb{C}[W'']$  generated by  $(sl(V'') \otimes V_0) \oplus (M_0 \otimes V_0) \oplus (M_0 \otimes sl(V)) \subset \mathbb{C}[W'']_2$ . The ideal  $I_0$  is homogeneous,  $G'' \times G$ -stable, and contains the ideal generated by the homogeneous  $G''$ -invariants of positive degree of  $\mathbb{C}[W'']$ . In particular,  $I_0$  identifies with an ideal of  $\mathbb{C}[\mu'^{-1}(0)]$ .

**Proposition 3.17.** *Let  $I_0 \subset \mathbb{C}[W'']$  be the ideal defined above, then  $I_0$  is a point of the invariant Hilbert scheme  $\text{Hilb}_{h_0}^G(\mu'^{-1}(0))$  defined in Lemma 3.15.*

*Proof.* We have to check that the ideal  $I_0$  has the good Hilbert function, that is,

$$\mathbb{C}[W'']/I_0 \cong \bigoplus_{M \in \text{Irr}(G)} M^{\oplus \dim(M)}$$

as a  $G$ -module. To do that, we are going to adapt the method used by Kraft and Schwarz to prove [KS, Theorem 9.1].

We denote  $R := V''^* \otimes V$ , which is an irreducible  $G'' \times G$ -submodule of  $W''^* \cong R \oplus R^*$ . Then  $R$  and  $R^*$  are orthogonal modulo  $I_0$ , which means that the image of the  $G'' \times G$ -submodule  $R \otimes R^* \subset \mathbb{C}[W'']_2$  in  $\mathbb{C}[W'']/I_0$  is isomorphic to the highest weight component of  $R \otimes R^*$  (that is,  $sl(V'') \otimes sl(V)$ ). Then, by [Bri85, Lemme 4.1], any irreducible  $G'' \times G$ -submodule of  $\mathbb{C}[R]$  is orthogonal to any irreducible  $G'' \times G$ -submodule of  $\mathbb{C}[R^*]$ , and thus the natural morphism

$$\phi : \mathbb{C}[R]^{U'' \times U} \otimes \mathbb{C}[R^*]^{U'' \times U} \rightarrow (\mathbb{C}[W'']/I_0)^{U'' \times U}$$

is surjective, where  $U''$  resp.  $U$ , denotes the unipotent radical of a Borel subgroup  $B'' \subset G''$  resp.  $B \subset G$ . Furthermore, if  $T'' \subset B''$  resp.  $T \subset B$ , is a maximal torus, then  $\phi$  is  $T'' \times T$ -equivariant.

Now by [Pro07, §13.5.1] we have the following isomorphisms of  $T'' \times T$ -algebras  $\mathbb{C}[R]^{U'' \times U} \cong \mathbb{C}[x_1, \dots, x_n]$ , where  $x_i \in \Lambda^i V'' \otimes \Lambda^i V^*$  is a highest weight vector, and  $\mathbb{C}[R^*]^{U'' \times U} \cong \mathbb{C}[y_1, \dots, y_n]$ , where  $y_j \in \Lambda^j V''^* \otimes \Lambda^j V$  is a highest weight vector. Hence, we have an exact sequence

$$0 \rightarrow K_0 \rightarrow \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] \rightarrow (\mathbb{C}[W'']/I_0)^{U'' \times U} \rightarrow 0,$$

where  $K_0$  is the kernel of  $\phi$ . One may check that the ideal  $K_0$  is generated by the products  $x_r y_s$  with  $r + s > n$  (see [KS, §9, Proof of Theorem 9.1(1)]).

We denote  $\Lambda = \langle \epsilon_1, \dots, \epsilon_n \rangle$  the weight lattice of the linear group  $GL_n$ , and  $\Lambda_+ \subset \Lambda$  the subset of dominant weights, that is, weights of the form  $r_1 \epsilon_1 + \dots + r_n \epsilon_n$ , with  $r_1 \geq \dots \geq r_n$ . If  $\lambda \in \Lambda_+$ , then we denote  $S^\lambda(\mathbb{C}^n)$  the irreducible  $GL_n$ -module of highest weight  $\lambda$ . We fix  $\lambda = k_1 \epsilon_1 + \dots + k_t \epsilon_t - k_{t+1} \epsilon_{t+1} - \dots - k_n \epsilon_n \in \Lambda_+$ , where each

$k_i$  is a nonnegative integer. One easily checks that the weight of the monomial

$$x_{n-t}^{k_{t+1}} x_{n-t-1}^{k_{t+2}-k_{t+1}} x_{n-t-2}^{k_{t+3}-k_{t+2}} \dots x_1^{k_n-k_{n-1}} y_t^{k_t} y_{t-1}^{k_{t-1}-k_t} y_{t-2}^{k_{t-2}-k_{t-1}} \dots y_1^{k_1-k_2}$$

for the action of  $T'' \times T$  is  $(\lambda^*, \lambda)$ , where  $\lambda^*$  denotes the highest weight of the  $GL_n$ -module  $S^\lambda(\mathbb{C}^{n^*})$ , and that  $\lambda$  uniquely determines this monomial. We get that the isotypic component of the  $G$ -module  $S^\lambda(V)$  in  $\mathbb{C}[W'']/I_0$  is the  $G'' \times G$ -module  $S^\lambda(V''^*) \otimes S^\lambda(V)$ . As  $\dim(V) = \dim(V'') = n$ , we have  $\dim(S^\lambda(V)) = \dim(S^\lambda(V''^*))$  for every  $\lambda \in \Lambda^+$ . In other words, each irreducible  $G$ -module  $M$  occurs in  $\mathbb{C}[W'']/I_0$  with multiplicity  $\dim(M)$ .  $\square$

By Proposition 3.17, the scheme  $\text{Hilb}_{h_0}^G(\mu'^{-1}(0))$  is non-empty, and thus  $\mathcal{H}$  has an irreducible component, different from  $\mathcal{H}^{\text{main}}$ , of dimension greater or equal to  $\dim(A_n) = 2n(n' - n)$ , which implies Proposition 3.14.

**3.5. Study of the case  $n = 1$ .** In all this Section, we suppose that  $n = 1$  and  $n' \geq 2$  (the case  $n' = 1$  being trivial). Then  $G = \mathbb{G}_m$  is the multiplicative group,  $W//G = \overline{\mathcal{O}_{[2,1^{n'-2}]}} \subset \mathfrak{g}'$ , and the morphism of Proposition 3.7 is  $\rho : \text{Hilb}_{h_0}^G(W) \rightarrow \mathbb{P}(V') \times \mathbb{P}(V'^*)$ . The Segre embedding gives a  $G'$ -equivariant isomorphism  $\mathbb{P}(V') \times \mathbb{P}(V'^*) \cong \mathbb{P}(\mathfrak{g}'^{\leq 1})$ , where  $\mathfrak{g}'^{\leq 1} := \{f \in \mathfrak{g}' \mid \text{rk}(f) \leq 1\}$ , and thus we can consider  $\rho' : \text{Hilb}_{h_0}^G(W) \rightarrow \mathbb{P}(\mathfrak{g}'^{\leq 1})$ , the morphism induced by  $\rho$ .

We saw in Section 3.3 that  $\mathcal{H}^{\text{main}}$  is a smooth variety, and in Section 3.4 that  $\mathcal{H}$  is always reducible. In this section, we determine the irreducible components of  $\mathcal{H}$  when  $n = 1$ .

**Proposition 3.18.** *We equip  $\mathcal{H}$  with its reduced structure. If  $n' > n = 1$  and with the above notation, we have a  $G'$ -equivariant isomorphism*

$$\mathcal{H} \cong \left\{ (f, L) \in \overline{\mathcal{O}_{[2,1^{n'-2}]}} \times \mathbb{P}(\mathfrak{g}'^{\leq 1}) \mid f \in L \right\}.$$

In particular,  $\mathcal{H}$  is the union of two smooth irreducible components of dimension  $2n' - 2$  defined by:

- $C_1 := \left\{ (f, L) \in \overline{\mathcal{O}_{[2,1^{n'-2}]}} \times \mathbb{P}(\overline{\mathcal{O}_{[2,1^{n'-2}]}}) \mid f \in L \right\} = \mathcal{H}^{\text{main}}$ , and the Hilbert-Chow morphism  $\gamma : \mathcal{H}^{\text{main}} \rightarrow \overline{\mathcal{O}_{[2,1^{n'-2}]}}$  is the blow-up of  $\overline{\mathcal{O}_{[2,1^{n'-2}]}}$  at 0;
- $C_2 := \left\{ (0, L) \in \overline{\mathcal{O}_{[2,1^{n'-2}]}} \times \mathbb{P}(\mathfrak{g}'^{\leq 1}) \right\} \cong \mathbb{P}(\mathfrak{g}'^{\leq 1})$ .

*Proof.* By [Tera, §1, Theorem], we have a  $G'$ -equivariant isomorphism

$$\gamma \times \rho' : \text{Hilb}_{h_0}^G(W) \rightarrow \left\{ (f, L) \in \mathfrak{g}'^{\leq 1} \times \mathbb{P}(\mathfrak{g}'^{\leq 1}) \mid f \in L \right\}.$$

Then, by [Bri, Lemma 3.3], we have a  $G'$ -equivariant closed embedding

$$\gamma \times \rho' : \mathcal{H} \hookrightarrow \mathcal{Y} := \left\{ (f, L) \in \overline{\mathcal{O}_{[2,1^{n'-2}]}} \times \mathbb{P}(\mathfrak{g}'^{\leq 1}) \mid f \in L \right\}.$$

One may check that  $\mathcal{Y}$  is the union of the two closed subset  $C_1$  and  $C_2$ , both of dimension  $2n' - 2$ . The morphism  $\gamma \times \rho'$  sends  $\mathcal{H}^{\text{main}}$  into  $C_1$ ; the varieties  $\mathcal{H}^{\text{main}}$  and  $C_1$  have the same dimension, hence  $\gamma \times \rho' : \mathcal{H}^{\text{main}} \rightarrow C_1$  is an isomorphism. On the other hand, we saw in Section 3.4 that  $\mathcal{H}$  admits a second irreducible component, denoted  $\mathcal{H}_2$ , of dimension at least  $2n' - 2$ , which is the dimension of  $C_2$ , and thus  $\gamma \times \rho'$  is an isomorphism between  $\mathcal{H}_2$  and  $C_2$ .  $\square$

*Remark 3.19.* One may check that the component  $C_2$  of Proposition 3.18 is formed by the homogeneous ideals of  $\mathbb{C}[\mu^{-1}(0)]$ .

When  $n' \geq 2n \geq 4$ , great dimensional irreducible components (other than  $\mathcal{H}^{\text{main}}$ ) may appear. For instance, if  $n = 2$  and  $n' \geq 4$ , then one may check that the irreducible component formed by the homogeneous ideals of  $\mathbb{C}[\mu^{-1}(0)]$  is of dimension  $4n' - 5$ , whereas the main component  $\mathcal{H}^{\text{main}}$  is of dimension  $4n' - 8$ . In addition, we showed in Section 3.4 that  $\mathcal{H}$  has at least two components, but  $\mathcal{H}$  may have much more components.

#### 4. CASE OF $Sp_n$ ( $n$ EVEN)

In this section, we denote  $E := V' \oplus V'^*$ , and we take  $G = Sp(V)$  and  $G' = SO(E)$ . We identify  $W = \text{Hom}(V', V) \times \text{Hom}(V, V') \cong \text{Hom}(E, V)$ , equipped with the natural action of  $G' \times G$ . We denote  $\mathfrak{g}$  resp.  $\mathfrak{g}'$ , the Lie algebra of  $G$  resp. of  $G'$ , and we recall that we denote  $n := \dim(V)$  (which is even), and  $n' := \dim(V')$ .

**4.1. The quotient morphism.** The main results of this section are Proposition 4.3, which describes the irreducible components of the symplectic reduction  $W///G := \mu^{-1}(0)//G$ , and Corollary 4.6, which gives the Hilbert function of the general fiber of the quotient morphism for each irreducible component of  $W///G$ . Contrary to the case of  $GL(V)$  studied in Section 3, we will see that  $W///G$  is reducible when  $n' \leq n$  and  $n'$  is even.

As  $G$  resp.  $G'$ , preserves a non-degenerate bilinear form on  $V$  resp. on  $E$ , we have a canonical isomorphism  $V \cong V^*$  resp.  $E \cong E^*$ . If  $w \in \text{Hom}(E, V)$ , we denote  ${}^t w \in \text{Hom}(V^*, E^*) \cong \text{Hom}(V, E)$  the transpose of  $w$ . Then, by [Bec09, Proposition 3.1], the zero fiber of the moment map  $\mu : W \rightarrow \mathfrak{g}^*$  is

$$\mu^{-1}(0) = \{w \in W \mid w \circ {}^t w = 0\},$$

which is a  $G' \times G$ -stable closed subset of  $W$ .

*Remark 4.1.* One may check that the biggest subgroup of  $GL(E)$  that stabilizes  $\mu^{-1}(0)$  in  $W$  is the orthogonal group  $O(E)$ . However, for practical reasons (study of the nilpotent orbits of  $\mathfrak{g}'$ ), we prefer to consider the action of  $G' = SO(E)$ .

The proof of the next proposition is analogous to the one of Proposition 3.1 and Corollary 3.2.

**Proposition 4.2.** *The zero fiber of the moment map  $\mu : W \rightarrow \mathfrak{g}^*$  is*

- a variety of dimension  $2n'n - \frac{1}{2}n(n+1)$  if  $n' > n$ ;
- the union of two varieties of dimension  $n'n + \frac{1}{2}n'(n'-1)$  if  $n' \leq n$ .

The following result was proved by Becker ([Bec09, Proposition 3.6]):

**Proposition 4.3.** *Let  $\mathbf{d}$  be a partition of  $2n'$ , and  $\mathcal{O}_{\mathbf{d}}$  resp.  $\mathcal{O}_{\mathbf{d}}^I$  and  $\mathcal{O}_{\mathbf{d}}^{II}$ , denote the nilpotent orbit(s) of  $\mathfrak{g}' \cong \mathfrak{so}_{2n'}$  associated to  $\mathbf{d}$ . Then the symplectic reduction of  $W$  by  $G$  is*

$$W///G = \begin{cases} \overline{\mathcal{O}_{[2^n, 1^{2(n'-n)]}}} & \text{if } n' > n; \\ \overline{\mathcal{O}_{[2^{n'-1}, 1^2]}} & \text{if } n' < n \text{ and } n' \text{ is odd}; \\ \overline{\mathcal{O}_{[2^{n'}]}^I \cup \mathcal{O}_{[2^{n'}]}^{II}} & \text{if } n' \leq n \text{ and } n' \text{ is even.} \end{cases}$$

**Corollary 4.4.** *The orbits for the adjoint action of  $G'$  on  $W///G$  are*

- $U_i := \mathcal{O}_{[2^i, 1^{2(n'-i)]}}$ , for  $i = 0, 2, \dots, n$ , if  $n' > n$ ;
- $U_i := \mathcal{O}_{[2^i, 1^{2(n'-i)]}}$ , for  $i = 0, 2, \dots, n'-1$ , if  $n' < n$  and  $n'$  is odd;



- $U_i := \mathcal{O}_{[2^i, 12(n'-i)]}$ , for  $i = 0, 2, \dots, n' - 2$ , and  $U_{n'}^I := \mathcal{O}_{[2^{n'}]}^I$ ,  $U_{n'}^{II} := \mathcal{O}_{[2^{n'}]}^{II}$ , if  $n' \leq n$  and  $n'$  is even.

The closures of the nilpotent orbits  $U_i$  are nested in the following way:

$$\begin{cases} \{0\} = \overline{U_0} \subset \overline{U_2} \subset \dots \subset \overline{U_n} & \text{if } n' > n; \\ \{0\} = \overline{U_0} \subset \overline{U_2} \subset \dots \subset \overline{U_{n'-1}} & \text{if } n' < n \text{ and } n' \text{ is odd}; \\ \{0\} = \overline{U_0} \subset \overline{U_2} \subset \dots \subset \overline{U_{n'-2}} = \overline{U_{n'}^I} \cap \overline{U_{n'}^{II}} & \text{if } n' \leq n \text{ and } n' \text{ is even.} \end{cases}$$

If ( $n' > n$  or  $n'$  is odd) resp. if ( $n' \leq n$  and  $n'$  is even), then the symplectic reduction  $W///G$  is the closure of a nilpotent orbit resp. is the union of two closures of nilpotent orbits, and thus the irreducible components of  $W///G$  are symplectic varieties (see Section 2). If  $n' > n$ , then  $W///G$  is of dimension  $2n'n - n(n+1)$  ([CM93, Corollary 6.1.4]), and its singular locus is  $\overline{U_{n-2}}$  ([KP82, Theorem 2]). On the other hand, if  $n' \leq n$ , then each irreducible component of  $W///G$  is of dimension  $n'(n'-1)$ , and the singular locus of  $W///G$  is  $\overline{U_{n'-2}}$  resp.  $\overline{U_{n'-3}}$ , when  $n'$  is even resp. when  $n'$  is odd.

We are now interested by the Hilbert function of the general fiber of the quotient morphism for each irreducible component of  $W///G$ . We will distinguish between the following cases:

- If  $n' > n$ , then  $W///G$  is irreducible, and we denote  $h_0$  the Hilbert function of the general fiber of the quotient morphism  $\nu : \mu^{-1}(0) \rightarrow W///G$ . By Proposition 4.2, the dimension of the general fiber of  $\nu$  is  $\frac{1}{2}n(n+1)$ .
- If  $n' \leq n$  and  $n'$  is even, then by Proposition 4.2, the zero fiber  $\mu^{-1}(0)$  is the union of two irreducible components that we denote  $X_I$  and  $X_{II}$ . Let  $\nu_I : X_I \rightarrow Y_I$  and  $\nu_{II} : X_{II} \rightarrow Y_{II}$  be the quotient morphisms. Up to the exchange of  $X_I$  and  $X_{II}$ , we can suppose that  $Y_I = \overline{U_{n'}^I}$  and  $Y_{II} = \overline{U_{n'}^{II}}$ . The orthogonal group  $O(E)$  acts transitively on  $U_{n'}^I \cup U_{n'}^{II}$ , hence the general fibers of  $\nu_I$  and  $\nu_{II}$  are isomorphic. In particular, the general fibers of  $\nu_I$  and  $\nu_{II}$  have the same Hilbert function, denoted  $h_0$ , and the same dimension, which is  $n'n - \frac{1}{2}n'(n'-1)$ .
- If  $n' < n$  and  $n'$  is odd, then  $W///G$  is irreducible, and we denote  $h_0$  the Hilbert function of the general fiber of the quotient morphism  $\nu : \mu^{-1}(0) \rightarrow W///G$ . The general fiber being reducible, determining  $h_0$  is more complicated than in the previous cases (except the case  $n' = 1$  which is trivial). From now on, we will always exclude the case  $n' < n$  and  $n'$  is odd.

If  $n' < n$  and  $n'$  is even, then we denote

$$(16) \quad H := \left\{ \begin{bmatrix} M & 0_{n-n', n'} \\ 0_{n', n-n'} & I_{n'} \end{bmatrix}, M \in Sp_{n-n'} \right\} \cong Sp_{n-n'},$$

which is a reductive algebraic subgroup of  $G \cong Sp_n$ . The proof of the next proposition is analogous to the proof of Proposition 3.5:

**Proposition 4.5.** *If  $n' > n$ , then the general fiber of the quotient morphism  $\nu : \mu^{-1}(0) \rightarrow W///G$  is isomorphic to  $G$ .*

*If  $n' = n$ , then the general fibers of the quotient morphisms  $\nu_I : X_I \rightarrow Y_I$  and  $\nu_{II} : X_{II} \rightarrow Y_{II}$  are isomorphic to  $G$ .*

*If  $n' < n$  and  $n'$  is even, then the general fibers of  $\nu_I$  and  $\nu_{II}$  are isomorphic to  $G/H$ , where  $H \subset G$  is the subgroup defined by (16).*

**Corollary 4.6.** *The Hilbert function  $h_0$  defined above is given by:*

$$\forall M \in \text{Irr}(G), h_0(M) = \begin{cases} \dim(M) & \text{if } n' \geq n; \\ \dim(M^H) & \text{if } n' < n \text{ and } n' \text{ is even;} \end{cases}$$

where  $H \subset G$  is the subgroup defined by (16).

**4.2. The reduction principle for the main component.** In this section, we give the guidelines to prove the reduction principle when  $G = Sp(V)$  (Proposition 4.8). The strategy is the same as for  $GL(V)$  (see Section 3.2), but as the symplectic reduction  $W///G$  is reducible when  $n' \leq n$  and  $n'$  is even, it seems necessary to give some details.

As  $\mu^{-1}(0)$  is a  $G' \times G$ -stable closed subset of  $W$ , it follows from [Bri, Lemma 3.3] that the invariant Hilbert scheme

$$\mathcal{H} := \text{Hilb}_{h_0}^G(\mu^{-1}(0))$$

is a  $G'$ -stable closed subscheme of  $\text{Hilb}_{h_0}^G(W)$ . As we aim at constructing canonical desingularizations of the irreducible components of  $W///G$ , we consider the two  $G'$ -stable closed subschemes  $\mathcal{H}_I := \text{Hilb}_{h_0}^G(X_I)$  and  $\mathcal{H}_{II} := \text{Hilb}_{h_0}^G(X_{II})$  instead of  $\mathcal{H}$  when  $n' \leq n$  and  $n'$  is even. Let us note that if we fix  $g'_0 \in O(E) \backslash SO(E)$  and make  $G'$  act on  $X_{II}$  by  $(g'_0 g' g'_0^{-1}) \cdot x$  for every  $g' \in G'$  and every  $x \in X_{II}$ , then  $\phi: X_I \rightarrow X_{II}$ ,  $x \mapsto g'_0 \cdot x$  is a  $G' \times G$ -equivariant isomorphism, and thus  $\mathcal{H}_I \cong \mathcal{H}_{II}$  as a  $G'$ -scheme. We denote  $\mathcal{H}_I^{\text{main}}$  resp.  $\mathcal{H}_{II}^{\text{main}}$ , the main component of  $\mathcal{H}_I$  resp. of  $\mathcal{H}_{II}$ . We always have the (set-theoretic) inclusion  $\mathcal{H}_I \cup \mathcal{H}_{II} \subset \mathcal{H}$ , but this may not be an equality. If  $n' > n$ , then  $W///G$  is irreducible, and we denote  $\mathcal{H}^{\text{main}}$  the main component of  $\mathcal{H}$ .

The scheme  $\text{Hilb}_{h_0}^G(W)$  was studied in [Terb]. In particular, we obtained

**Proposition 4.7.** ([Terb, §1.5.1]) *Let  $h_0$  be the Hilbert function given by Corollary 4.6, and  $G' = SO(E)$  acting naturally on  $\text{Gr}(2n' - h_0(V^*), E)$ . Then there exists a  $G'$ -equivariant morphism*

$$\begin{array}{ccc} \rho & : & \text{Hilb}_{h_0}^G(W) \rightarrow \text{Gr}(2n' - h_0(V^*), E) \\ & & Z \mapsto \text{Ker}(f_Z) \end{array}$$

where  $f_Z: E \cong \text{Mor}^G(W, V^*) \rightarrow \text{Mor}^G(Z, V^*)$  is the restriction map.

We identify  $\text{Gr}(2n' - h_0(V^*), E)$  with  $\text{Gr}(h_0(V^*), E^*)$ . By Corollary 4.6, if  $n' > n$  or  $(n' \leq n$  and  $n'$  is even), then  $h_0(V^*) = N := \min(n', n)$ . We denote  $q$  the non-degenerate quadratic form on  $E$  preserved by  $O(E)$ . We have a canonical isomorphism  $E \cong E^*$ , and thus  $q$  identifies with a non-degenerate quadratic form on  $E^*$ . For  $i = 0, \dots, N$ , we denote

$$A_i := \{L \in \text{Gr}(N, E^*) \mid q|_L \text{ is of rank } i\}.$$

If  $n' > n$ , then the  $A_i$  are the  $n+1$  orbits for the action of  $G'$  on  $\text{Gr}(n, E^*)$ . However, if  $n' \leq n$ , then the  $A_i$  are  $G'$ -orbits for  $i = 1, \dots, n'$ , but the isotropic Grassmannian  $A_0 = \text{OGr}(n', E^*)$  is the union of two  $G'$ -orbits, denoted  $\text{OGr}^I$  and  $\text{OGr}^{II}$ , which are exchanged by the action of any element of  $O(E) \backslash SO(E)$ . In any case, we have

$$\text{OGr}(N, E^*) = \overline{A_0} \subset \overline{A_1} \subset \dots \subset \overline{A_N} = \text{Gr}(N, E^*).$$

Let us now fix some notation:

- $L_0 \in A_0$ , and  $P$  the parabolic subgroup of  $G'$  stabilizing  $L_0$ ;

- $W' := \text{Hom}(E/L_0^\perp, V)$ , which identifies with a  $P \times G$ -module contained in  $\mu^{-1}(0)$ ;
- $\mathcal{H}' := \text{Hilb}_{h_0}^G(W')$ , and  $\mathcal{H}'^{\text{main}}$  its main component.

It must be underlined that, if  $n' > n$  or ( $n' \leq n$  and  $n'$  is even), then the Hilbert function of the general fiber of the quotient morphism  $W' \rightarrow W'//G$  coincides with the Hilbert function  $h_0$  of Corollary 4.6 (in particular,  $\mathcal{H}'^{\text{main}}$  is well defined).

Proceeding as for Lemma 3.9, one may check that, if  $n' > n$  resp. if ( $n' \leq n$  and  $n'$  is even), then the morphism  $\rho$  of Proposition 4.7 sends  $\mathcal{H}^{\text{main}}$  resp.  $\mathcal{H}_I^{\text{main}}$  and  $\mathcal{H}_{II}^{\text{main}}$ , onto  $A_0$ . More precisely, if  $n' \leq n$  and  $n'$  is even, then  $\rho$  sends  $\mathcal{H}_I^{\text{main}}$  onto one of the irreducible component of  $A_0$ , and  $\mathcal{H}_{II}^{\text{main}}$  onto the other component. Up to the exchange of these two components, we can suppose that  $\rho$  sends  $\mathcal{H}_I^{\text{main}}$  onto  $\text{OGr}^I$ , and  $\mathcal{H}_{II}^{\text{main}}$  onto  $\text{OGr}^{II}$ .

It follows that the restriction of  $\rho$  equips  $\mathcal{H}^{\text{main}}$  resp.  $\mathcal{H}_I^{\text{main}}$ , resp.  $\mathcal{H}_{II}^{\text{main}}$ , with a structure of a  $G'$ -homogeneous fiber bundle over  $A_0$  resp. over  $\text{OGr}^I$ , resp. over  $\text{OGr}^{II}$ . Hence, it is enough to determine the fiber  $F_0$  over  $L_0$  to determine  $\mathcal{H}^{\text{main}}$  resp.  $\mathcal{H}_I^{\text{main}}$ , resp.  $\mathcal{H}_{II}^{\text{main}}$ . Proceeding as in Section 3.2, we obtain that  $F_0$  is isomorphic to  $\mathcal{H}'^{\text{main}}$  as a  $P$ -scheme. We deduce

**Proposition 4.8.** *With the above notation, we have the following  $G'$ -equivariant isomorphisms:*

- If  $n' > n$ , then

$$\mathcal{H}^{\text{main}} \cong G' \times^P \mathcal{H}'^{\text{main}}.$$

- If ( $n' \leq n$  and  $n'$  is even) and  $L_0 \in \text{OGr}^I$  resp.  $L_0 \in \text{OGr}^{II}$ , then

$$\mathcal{H}_I^{\text{main}} \cong G' \times^P \mathcal{H}'^{\text{main}} \quad \text{resp.} \quad \mathcal{H}_{II}^{\text{main}} \cong G' \times^P \mathcal{H}'^{\text{main}}.$$

**4.3. Proof of Theorem A for  $\text{Sp}(\mathbf{V})$ .** Let us start by recalling

**Theorem 4.9.** [Tera, §1, Theorem] *Let  $G = \text{Sp}(V)$ ,  $W = \text{Hom}(E, V)$ , and  $h_W$  the Hilbert function of the general fiber of the quotient morphism  $W \rightarrow W//G$ . We denote  $n := \dim(V)$ ,  $e := \dim(E)$ , and  $Y_0$  the blow-up of  $W//G = \Lambda^2(E^*)^{\leq n} := \{Q \in \Lambda^2(E^*) \mid \text{rk}(Q) \leq n\}$  at 0. In the following cases, the invariant Hilbert scheme  $\mathcal{H}' := \text{Hilb}_{h_W}^G(W)$  is a smooth variety, and the Hilbert-Chow morphism is the succession of blow-up described as follows:*

- if  $n \geq 2e - 2$ , then  $\mathcal{H}' \cong W//G = \Lambda^2(E^*)$ ;
- if  $e > n = 2$  or  $e = n = 4$ , then  $\mathcal{H}' \cong Y_0$ ;
- if  $e > n = 4$ , then  $\mathcal{H}'$  is isomorphic to the blow-up of  $Y_0$  along the strict transform of  $\Lambda^2(E^*)^{\leq 2}$ .

If  $n' > n$ , then we denote  $T$  the tautological bundle over  $A_0 = \text{OGr}(n, E^*)$ . If  $n' \leq n$  and  $n'$  is even, then we denote  $T_I$  resp.  $T_{II}$ , the tautological bundle over  $\text{OGr}^I$  resp. over  $\text{OGr}^{II}$ . We deduce from Proposition 4.8 and Theorem 4.9 that we have the following  $G'$ -equivariant isomorphisms

$$(17) \quad \mathcal{H}^{\text{main}} \cong \begin{cases} \Lambda^2(T) & \text{if } n' > n = 2; \\ \text{Bl}_0(\Lambda^2(T)) & \text{if } n' > n = 4; \end{cases}$$

$$(18) \quad \mathcal{H}_\bullet^{\text{main}} \cong \begin{cases} \Lambda^2(T_\bullet) & \text{if } n \geq 2n' - 2 \text{ and } n' \text{ is even;} \\ \text{Bl}_0(\Lambda^2(T_\bullet)) & \text{if } n' = n = 4; \end{cases}$$

where  $\bullet$  stands for  $I$  or  $II$ , and  $\text{Bl}_0(\cdot)$  denotes the blow-up along the zero section. In all these cases, the main component of the invariant Hilbert scheme is smooth,

and thus the Hilbert-Chow morphism  $\gamma: \mathcal{H}^{\text{main}} \rightarrow W///G$  resp.  $\gamma: \mathcal{H}_{\bullet}^{\text{main}} \rightarrow Y_{\bullet}$ , is a (distinguished) desingularization.

It remains to compare  $\gamma$  with the Springer desingularizations (when they exist) of the irreducible components of  $W///G$ . We saw in Section 2 that the irreducible components of  $W///G$  have Springer desingularizations if and only if  $n' \leq n+1$ . We then distinguish between the following cases:

- (1) If  $n' \leq n+1$  and  $n'$  is odd, then  $W///G$  admits two Springer desingularizations, which are given by the cotangent bundles  $\mathcal{T}_I^*$  and  $\mathcal{T}_{II}^*$  over  $\text{OGr}^I$  and  $\text{OGr}^{II}$  respectively. The natural action of the orthogonal group  $O(E)$  on  $\text{OGr}(n', E^*)$  induces an action on the cotangent bundle  $\mathcal{T}^*\text{OGr}(n', E^*)$  that exchanges  $\mathcal{T}_I^*$  and  $\mathcal{T}_{II}^*$ . On the other hand, it follows from Remark 4.1, that the group  $O(E)$  stabilizes  $\mathcal{H}^{\text{main}}$ , and thus  $\gamma: \mathcal{H}^{\text{main}} \rightarrow W///G$  can not be a Springer desingularization. However, if  $n \in \{2, 4\}$  and  $n' = n+1$ , then one may show that  $\gamma$  dominates the two Springer desingularizations of  $W///G$  (see [Bec11, Introduction] for the case  $n=2$ , the case  $n=4$  being analogous).
- (2) If  $n' \leq n$  and  $n'$  is even, then  $Y_{\bullet}$  has a unique Springer desingularization, which is given by the cotangent bundle  $\mathcal{T}_{\bullet}^* \cong \Lambda^2(T_{\bullet})$  over  $\text{OGr}_{\bullet}$ . Proceeding as we did for  $GL(V)$  in Section 3.3, one may show that  $\gamma: \mathcal{H}_{\bullet}^{\text{main}} \rightarrow Y_{\bullet}$  is the Springer desingularization if and only if  $n \geq 2n' - 2$ . In addition, if  $n' = n = 4$ , then by (18) we have  $\mathcal{H}_{\bullet}^{\text{main}} \cong \text{Bl}_0(\mathcal{T}_{\bullet}^*)$ , and thus  $\gamma$  dominates the unique Springer desingularization of  $Y_{\bullet}$ .

**4.4. Study of the case  $n=2$ .** In this section, we suppose that  $n' \geq n=2$  (the case  $n'=1$  being trivial). Then  $G \cong Sp_2$ , and the morphism of Proposition 4.7 is  $\rho: \text{Hilb}_{h_0}^G(W) \rightarrow \text{Gr}(2, E^*)$ . Denoting  $\mathfrak{g}'^{\leq 2} := \{f \in \mathfrak{g}' \mid \text{rk}(f) \leq 2\}$ , we have a  $G'$ -equivariant isomorphism

$$(19) \quad \mathbb{P}(\mathfrak{g}'^{\leq 2}) \cong \text{Gr}(2, E^*),$$

and thus we can consider  $\rho': \text{Hilb}_{h_0}^G(W) \rightarrow \mathbb{P}(\mathfrak{g}'^{\leq 2})$ , the morphism induced by  $\rho$ . By Proposition 4.3, we have

$$W///G = \begin{cases} \overline{\mathcal{O}_{[2^2, 1^{2n'-4}]}} & \text{if } n' \geq 3; \\ \overline{\mathcal{O}_{[2^2]}^I \cup \mathcal{O}_{[2^2]}^{II}} & \text{if } n' = 2. \end{cases}$$

We will show that if  $n' \geq 3$  resp. if  $n' = 2$ , then  $\mathcal{H}$  resp.  $\mathcal{H}_{\bullet}$  (where  $\bullet$  stands for  $I$  or  $II$ ), coincides with its main component. In particular, the geometric properties of the invariant Hilbert scheme for  $G = Sp(V)$  seem to be quite different from the case of  $G = GL(V)$  studied in Section 3. Let us recall that the case ( $n' = 3$  and  $n = 2$ ) was treated by Becker in [Bec11]; she showed that  $\mathcal{H}$  coincides with  $\mathcal{H}^{\text{main}}$  as a scheme, and is the total space of a line bundle over  $\text{OGr}(2, E^*)$ .

**Proposition 4.10.** *We equip all the invariant Hilbert schemes with their reduced structures. If  $n' > n = 2$ , then  $\mathcal{H} = \mathcal{H}^{\text{main}}$  is a smooth variety isomorphic to*

$$\text{Bl}_0(\overline{\mathcal{O}_{[2^2, 1^{2n'-4}]}}) := \{(f, L) \in \overline{\mathcal{O}_{[2^2, 1^{2n'-4}]}} \times \mathbb{P}(\overline{\mathcal{O}_{[2^2, 1^{2n'-4}]}}) \mid f \in L\},$$

and the Hilbert-Chow morphism  $\gamma: \mathcal{H} \rightarrow W///G$  is the blow-up of  $\overline{\mathcal{O}_{[2^2, 1^{2n'-4}]}}$  at 0. If  $n' = n = 2$ , then  $\text{Hilb}_{h_0}^G(\mu^{-1}(0)) = \mathcal{H}_I \cup \mathcal{H}_{II}$  is the union of two smooth varieties isomorphic to  $\text{Bl}_0(\overline{\mathcal{O}_{[2^2]}^I})$  and  $\text{Bl}_0(\overline{\mathcal{O}_{[2^2]}^{II}})$  respectively, and the intersection  $\mathcal{H}_I \cap \mathcal{H}_{II}$

is formed by the homogeneous ideals of  $\mathbb{C}[\mu^{-1}(0)]$ . Moreover, the Hilbert-Chow morphism  $\gamma: \mathcal{H}_I \rightarrow \overline{\mathcal{O}}_{[2^2]}^I$  resp.  $\gamma: \mathcal{H}_{II} \rightarrow \overline{\mathcal{O}}_{[2^2]}^{II}$ , is the blow-up of  $\overline{\mathcal{O}}_{[2^2]}^{II}$  resp. of  $\overline{\mathcal{O}}_{[2^2]}^I$ , at 0.

*Proof.* The proofs for the cases  $n' = 2$  and  $n' \geq 3$  are quite similar, and thus we will only consider the case  $n' \geq 3$  (which is simpler in terms of notation!). It follows from [Bri, Lemma 3.3] and [Tera, §1, Theorem] that we have a closed embedding

$$\gamma \times \rho': \mathcal{H} \hookrightarrow \mathcal{Y} := \left\{ (f, L) \in \overline{\mathcal{O}}_{[2^2, 1^{2n'-4}]} \times \mathbb{P}(\mathfrak{g}'^{\leq 2}) \mid f \in L \right\}.$$

One may check that  $\mathcal{Y}$  is the union of the two subvarieties  $C_1$  and  $C_2$  defined by:

- $C_1 := \text{Bl}_0(\overline{\mathcal{O}}_{[2^2, 1^{2n'-4}]})$ ;
- $C_2 := \left\{ (0, L) \in \overline{\mathcal{O}}_{[2^2, 1^{2n'-4}]} \times \mathbb{P}(\mathfrak{g}'^{\leq 2}) \right\} \cong \mathbb{P}(\mathfrak{g}'^{\leq 2})$ .

The subvarieties  $C_1$  and  $C_2$  are of dimension  $4n' - 6$  and  $4n' - 4$  respectively. The morphism  $\gamma \times \rho'$  sends  $\mathcal{H}^{\text{main}}$  into  $C_1$ ; the varieties  $\mathcal{H}^{\text{main}}$  and  $C_1$  have the same dimension, hence  $\gamma \times \rho': \mathcal{H}^{\text{main}} \rightarrow C_1$  is an isomorphism.

Now it follows from [Terb, Proposition 3.3.13] that the component  $C_2$  identifies with the closed subset of  $\text{Hilb}_{h_0}^G(W)$  formed by the homogeneous ideals of  $\mathbb{C}[W]$ . Let us describe this correspondance. If  $L \in C_2 \cong \mathbb{P}(\mathfrak{g}'^{\leq 2})$ , then we define the ideal  $I_L \subset \mathbb{C}[W]$  generated by the homogeneous  $G$ -invariants of positive degree of  $\mathbb{C}[W]$ , and by the  $G$ -module  $L^\perp \otimes V \subset \mathbb{C}[W]_1 \cong E \otimes V$ , where  $L$  is identified with a 2-dimensional subspace of  $E^*$  with (19). Let us show that  $I_L$  is a point of  $\mathcal{H}$  if and only if  $L \in \text{OGr}(2, E^*)$ ; the result will follow since  $\mathbb{P}(\overline{\mathcal{O}}_{[2^2, 1^{2n'-4}]})$  identifies with  $\text{OGr}(2, E^*)$  via the isomorphism (19), and that  $\left\{ (0, L) \in \overline{\mathcal{O}}_{[2^2, 1^{2n'-4}]} \times \mathbb{P}(\overline{\mathcal{O}}_{[2^2, 1^{2n'-4}]}) \right\}$  is a subvariety of  $C_1$ .

We denote  $W' := \text{Hom}(E/L^\perp, V)$ , then

$$\mathbb{C}[W']_2 \cong (S^2(E/L^\perp) \otimes S^2(V)) \oplus (\Lambda^2(E/L^\perp) \otimes \Lambda^2(V))$$

as a  $G$ -module. Let  $I'_L$  be the ideal of  $\mathbb{C}[W']$  generated by  $\Lambda^2(E/L^\perp) \otimes \Lambda^2(V) \subset \mathbb{C}[W']_2$ , then one may check that

$$\mathbb{C}[W']/I'_L \cong \mathbb{C}[W]/I_L \cong \bigoplus_{M \in \text{Irr}(G)} M^{\oplus \dim(M)}$$

as a  $G$ -module. Hence

$$\begin{aligned} I_L \in \mathcal{H} &\Leftrightarrow I_L \cap \mathbb{C}[W]_2 \supset E_0 \otimes S^2(V), \text{ where } E_0 \text{ is the trivial representation of } G'; \\ &\Leftrightarrow q_L = 0, \text{ where } q \text{ is the quadratic form preserved by } G'; \\ &\Leftrightarrow L \in \text{OGr}(2, E^*). \end{aligned}$$

□

*Remark 4.11.* In the proof of Proposition 4.10, we show that if  $n' > n = 2$ , then the homogeneous ideals of  $\mathcal{H}$  are contained in  $\mathcal{H}^{\text{main}}$ . Using analogous arguments, one may check that this statement is true more generally when  $n' > n \geq 2$ .

## REFERENCES

- [AB05] V. Alexeev and M. Brion. Moduli of affine schemes with reductive group action. *J. Algebraic Geom.*, 14:83–117, 2005.
- [Bea00] A. Beauville. Symplectic singularities. *Invent. Math.*, 139:no.3, 541–549, 2000.
- [Bec09] T. Becker. On the existence of symplectic resolutions of symplectic reductions. *Mathematische Zeitschrift*, 265:343–363, 2009.
- [Bec11] T. Becker. An example of an  $SL_2$ -Hilbert scheme with multiplicities. *Transform. Groups*, 16:no. 4, 915–938, 2011.
- [Bri] M. Brion. Invariant Hilbert schemes. arXiv: 1102.0198, to appear in "*Handbook of Moduli*".
- [Bri85] M. Brion. Représentations exceptionnelles des groupes semi-simples. *Ann. Sci. Ecole Norm. Sup.*, 2:345–387, 1985.
- [CM93] D. Collingwood and W. McGovern. *Nilpotent orbits in semisimple Lie algebras*. Van Nostrand Reinhold Mathematics Series, vol. 296. Van Nostrand Reinhold Co., New York, 1993.
- [FN04] B. Fu and Y. Namikawa. Uniqueness of crepant resolutions and symplectic singularities. *Ann. de l'institut Fourier*, 54:1–19, 2004.
- [Fu03a] B. Fu. Symplectic resolutions for nilpotent orbits. *Invent. Math.*, 151:167–186, 2003.
- [Fu03b] B. Fu. Symplectic resolutions for nilpotent orbits (II). *C. R. Acad. Sci. Paris*, 337:277–281, 2003.
- [Fu06a] B. Fu. A survey on symplectic singularities and resolutions. *Ann. Math. Blaise Pascal*, 13:209–236, 2006.
- [Fu06b] B. Fu. Symplectic resolutions for nilpotent orbits (III). *C. R. Acad. Sci. Paris*, 342:585–588, 2006.
- [KP79] H. Kraft and C. Procesi. Closures of conjugacy classes of matrices are normal. *Invent. Math.*, 53:227–247, 1979.
- [KP81] H. Kraft and C. Procesi. Minimal singularities in  $GL_n$ . *Invent. Math.*, 62:503–515, 1981.
- [KP82] H. Kraft and C. Procesi. On the geometry of conjugacy classes in classical groups. *Comment. Math. Helvetici*, 57:539–602, 1982.
- [KS] H. Kraft and G. W. Schwarz. Representations with a reduced null cone. arXiv: 1112.3634, to appear in *Progress in Mathematics* (Birkhäuser), a volume in honor of Nolan Wallach.
- [Nam06] Y. Namikawa. Birational geometry of symplectic resolutions of nilpotent orbits. *Moduli spaces and arithmetic geometry*, 45, Math. Soc. Japan, Tokyo:75–116, Adv. Stud. Pure Math., 2006.
- [Pro07] C. Procesi. *Lie Groups, an Approach through Invariants and Representations*. Universitext. Springer, New York, 2007.
- [SB00] G. W. Schwarz and M. Brion. *Théorie des invariants et géométrie des variétés quotients*. Travaux en cours, vol. 61. Hermann, Paris, 2000.
- [Tera] R. Terpereau. Invariant Hilbert schemes and desingularizations of quotients by classical groups. arXiv: 1301.4020.
- [Terb] R. Terpereau. Schémas de Hilbert invariants et théorie classique des invariants (Ph.D. thesis). arXiv: 1211.1472.

Université Grenoble I, Institut Fourier,  
 UMR 5582 CNRS-UJF, BP 74,  
 38402 St. Martin d'Hères Cédex, FRANCE  
*E-mail address:* ronan.terpereau@ujf-grenoble.fr