



Mercury in Bay Bolete (*Xerocomus badius*): bioconcentration by fungus and assessment of element intake by humans eating fruiting bodies

Jerzy Falandysz, Anna Katarzyna Kojta, Grażyna Jarzyńska, Malgorzata Drewnowska, Anna Dryżalowska, Daria Wydmańska, Izabela Kowalewska, Anna Wacko, Monika Szlosowska, Kannan Kurunthachalam, et al.

► To cite this version:

Jerzy Falandysz, Anna Katarzyna Kojta, Grażyna Jarzyńska, Malgorzata Drewnowska, Anna Dryżalowska, et al.. Mercury in Bay Bolete (*Xerocomus badius*): bioconcentration by fungus and assessment of element intake by humans eating fruiting bodies. Food Additives and Contaminants, 2012, pp.1. <10.1080/19440049.2012.662702>. <hal-00801034>

HAL Id: hal-00801034

<https://hal.science/hal-00801034v1>

Submitted on 15 Mar 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



**Mercury in Bay Bolete (*Xerocomus badius*):
bioconcentration by fungus
and assessment of element intake by humans eating
fruiting bodies**

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2011-508.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	22-Jan-2012
Complete List of Authors:	Falandysz, Jerzy; University of Gdańsk, Institute of Environmental Sciences & Public Health Kojta, Anna; University of Gdańsk, Chemistry; University of Gdańsk, Institute of Environmental Sciences & Public Health Jarzyńska, Grażyna; University of Gdańsk, Institute of Environmental Sciences & Public Health Drewnowska, Małgorzata; University of Gdańsk, Institute of Environmental Sciences & Public Health Dryżałowska, Anna; University of Gdańsk, Institute of Environmental Sciences & Public Health Wydmańska, Daria; University of Gdańsk, Institute of Environmental Sciences & Public Health Kowalewska, Izabela; University of Gdańsk, Institute of Environmental Sciences & Public Health Wacko, Anna; University of Gdańsk, Institute of Environmental Sciences & Public Health Szłosowska, Monika; University of Gdańsk, Institute of Environmental Sciences & Public Health Kurunthachalam, Kannan; Wadsworth Center, NYS Department of Health Szefer, Piotr; Medical University of Gdańsk, Department of Food Sciences
Methods/Techniques:	Metals analysis - AAS, Reference materials, Statistical analysis, Risk assessment
Additives/Contaminants:	Heavy metals - mercury
Food Types:	Mushrooms
Abstract:	Concentrations of Hg were determined in 221 specimens of Bay Bolete and in 221 samples of corresponding forest topsoil layer (0-10 cm) collected from 18 sites across geographically and industrially diverse regions of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>Poland in 2000-2008. Mercury concentrations in caps and stipes of Bay Bolete were strongly correlated ($p < 0.0001$), whereas the relationship between the Hg concentrations in soil and mushrooms varied depending on the sampling location. The Bay Bolete showed a lower bioconcentration potential of Hg at sites with elevated soil Hg concentrations and a higher at sites with lower Hg concentrations in soil. In a view of Hg content, the consumption of Bay Boletes (caps or a whole mushrooms) at the regions surveyed at least at a rate up to 2.5 kg per capita weekly in a mushrooming season is safe and will not result in exceeding of currently allowable Hg intake doses.</p>

SCHOLARONE™
Manuscripts

**Mercury in Bay Bolete (*Xerocomus badius*): bioconcentration by fungus
and assessment of element intake by humans eating fruiting bodies**

J. Falandysz^a, A.K. Kojta^{a*}, G. Jarzyńska^a, M. Drewnowska^a, A. Dryżałowska^a, D. Wydmańska^a,
I. Kowalewska^a, A. Wacko^a, M. Szlosowska^a, K. Kannan^b and P. Szefer^c

^aResearch Group of Environmental Chemistry, Ecotoxicology & Food Toxicology, Institute of
Environmental Sciences & Public Health, University of Gdańsk,
18 Sobieskiego Str., PL 80-952 Gdańsk, Poland

^bWadsworth Center, New York State Department of Health, and Department of Environmental
Health Sciences, School of Public Health, State University of New York at Albany,
Empire State Plaza, Albany, NY 12201-0509, USA

^cDepartment of Food Sciences, Medical University of Gdańsk, 107 Hallera Ave., PL 80-416
Gdańsk, Poland

* Corresponding author. Email: annakojta@gmail.com

Abstract

Concentrations of mercury (Hg) were determined in 221 specimens of Bay Bolete and in 221 samples of corresponding forest topsoil layer (0-10 cm) collected from 18 sites across geographically and industrially diverse regions of Poland in 2000-2008. Mercury concentrations in caps and stipes of Bay Bolete were strongly correlated ($p < 0.0001$), whereas the relationship between the Hg concentrations in soil and mushrooms varied depending on the sampling location. The Bay Bolete showed a lower bioconcentration potential of Hg at sites with elevated soil Hg concentrations and a higher at sites with lower Hg concentrations in soil. In a view of Hg content, the consumption of Bay Boletes (caps or a whole mushrooms) at the regions surveyed at least at a rate up to 2.5 kg *per capita* weekly in a mushrooming season is safe and will not result in exceeding of currently allowable Hg intake doses.

Keywords: food contamination; heavy metals; mushrooms; nutrition; mercury.

Introduction

Contamination of food resources with Hg from environmental releases of this hazardous metal is a continuous threat to food safety (Olivero et al., 2002). In order to protect human exposure from Hg found in the food supply, the Joint FAO/WHO Expert Committee on Food Additives established for total Hg a new provisional tolerable weekly intake (PTWI) of 0.004 mg kg⁻¹ body weight in 2010 (JECFA, 2010). A few earlier studies have shown that wild mushrooms can accumulate mercury from the soil to a considerable amount (Alonso et al., 2000; Chudzyński et al., 2009 and 2011; Falandysz et al., 2001a, 2002a-b, 2003a-d and 2004; Melgar et al., 2009; Stijve and Cardinale, 1974; Zimmermannova et al., 2001). The Bay Bolete mushroom, due to its size and seasonal abundance, is by volume, one of the dominant species of mushrooms harvested

in wild each year in Poland. For example, 1705 tons of mushrooms have been harvested from the wild by commercial harvesters in Poland in 2000, and it was 5914 tons in 2008, of which 3275 tons were solely of the genera *Xerocomus* (*X. badius*, *X. subtomentosus*, *X. chrysenteron*) (GUS, 2009). These quantities do not account for the amount harvested annually by numerous non-commercial collectors in Poland, where wild mushroom picking is a traditional activity.

Some species of higher fungi (macromycetes) are known for their ability to accumulate metallic elements including hazardous heavy metals such as Hg, Cd or Pb (Brzostowski et al., 2009 and 2011a-b; Chudzyński and Falandysz, 2008; Doğan et al., 2006; Falandysz et al., 2001b, 2007a-c, 2008a-b and 2011; Frankowska et al., 2010; Gast et al., 1988; Gucia et al., 2011a-b; Jarzyńska et al., 2011; Kojta et al., 2011; Malinowska et al., 2004; Melgar et al., 1998; Stijve and Besson, 1976). Concentrations of trace elements in underlying soil or other substratum and bioavailability are among the factors that can affect the concentrations of metals in fruiting bodies (carpophores) of fungi (Bargagli and Baldi, 1984; Borovička et al., 2010; Falandysz et al., 1994; Řanda and Kučera, 2004; Stijve et al., 2004).

Nevertheless, in nature, it is difficult to establish quantitative relationship between the concentrations of metallic element and metalloid content in mushrooms and the underlying substratum (Gast et al., 1988; Falandysz et al., 2011). If mushrooms and topsoil were collected concurrently from both contaminated and background (free of local pollution sources) areas, it is possible to elucidate the relationship soil metal concentrations and residue levels in mushrooms (Barcan et al., 1998; Falandysz et al., 2002c; Carvalho et al., 2005; Stijve and Roschnik, 1974).

In background areas, Cd concentrations in fruiting bodies of the King Bolete (*Boletus edulis*) fungus increased with Cd concentrations in topsoil (Falandysz et al., 2011). A positive association was found between soil methyl mercury concentrations and the concentrations in fruiting bodies of fungi, although the number of evidences is very limited (Fischer et al., 1995). A

1
2
3 67 similar relationship was also observed for total Hg in Parasol Mushroom (*Macrolepiota procera*)
4
5 68 (Falandysz and Chwir, 1997).
6
7

8 69 In this study, we determined total Hg concentrations in Bay Bolete (*X. badius*), and in
9
10 70 forest topsoil layer below the fruit bodies, at eighteen sites distantly distributed across Poland.
11
12 71 We examined bioconcentration potential of Hg in Bay Bolete and daily intake rates of Hg from
13
14 72 mushroom consumption. Bay Bolete is a mycorrhizal fungus associated with coniferous
15
16 73 woodland and mixed woodland. The symbiotic trees are supplied by fungus with portion of
17
18 74 minerals absorbed by mycelium. Nevertheless, plants are very poor in Hg while mushrooms are
19
20 75 relatively abundant in Hg (Cibulka et al., 1996).
21
22
23
24

25 76 **Materials and methods**
26
27

28 77 Bay Bolete, *Xerocomus badius* (Fr.) Kühn, is a mycorrhizal mushroom and is a common
29
30 78 species found in mixed woodlands of several European countries. The mushroom and topsoil
31
32 79 samples (0-10 cm layer; ca. 100 g) were collected from eighteen, geographically diverse
33
34 80 regions of Poland during 2000-2008 (Figure 1; Table 1). Mushrooms were cleaned up from
35
36 81 adhering plant material and soil particles with a plastic knife and brush and, if necessary, the
37
38 82 bottom part of stipe was cut away. Next, the fruiting bodies were initially air-dried for 48 h, and
39
40 83 then were oven-dried at 65 °C to a constant weight. Dried mushrooms were crushed and ground
41
42 84 in a ceramic mortar to fine powder and kept in a sealed polyethylene bags. Soil substrate
43
44 85 samples, after the removal of visible organisms, stones, sticks, and leaves, were air-dried for
45
46 86 approximately 10 weeks and then sieved through a 2 mm sieve (Falandysz, 2002; Falandysz et
47
48 87 al., 2003a).
49
50
51
52
53

54 88 Analysis of total Hg content in fruiting bodies and soil was conducted using a direct
55
56 89 sample thermal decomposition coupled with gold wool trap, desorption and cold-vapor atomic
57
58
59
60

absorption spectroscopy (CV–AAS; Mercury analyzer type MA–2000, Nippon Instruments Corporation, Takatsuki, Japan) determination (Jarzyńska and Falandysz, 2011a). The procedure for Hg determination was validated through analysis of several certified reference materials including: CS-M-1 (dried fruiting bodies of mushroom, Cow Bolete *Suillus bovinus*); CTA-OTL-1 (tobacco leaves) and INCT-TL-1 (tea leaves) produced by the Institute of Nuclear Chemistry and Technology in Warsaw, Poland. The declared total Hg contents of the certified reference materials were: $0.174 \pm 0.018 \mu\text{g g}^{-1}$ dw for CS-M-1 (our measurements were $0.179 \pm 0.008 \mu\text{g g}^{-1}$ dw; $n = 3$); $0.043 \mu\text{g g}^{-1}$ dw for CTA-OTL-1 (our measurements were $0.046 \pm 0.002 \mu\text{g g}^{-1}$ dw; $n = 3$); and $0.005 \pm 0.001 \mu\text{g g}^{-1}$ dw for INCT-TL-1 (our measurements were $0.005 \pm 0.000 \mu\text{g g}^{-1}$ dw; $n = 3$). In addition, with every set of 10 mushroom or soil samples analyzed, one procedural blank was included; no contamination or interference was found in blanks (Jarzyńska and Falandysz, 2011a).

Data were evaluated statistically using the computer software Statistica version 8.0. Mercury concentrations in mushrooms and soils have no Gaussian distribution. Data transformation, which aimed to obtain their log-normal distribution, was unsuccessful. Moreover, data variances were heterogeneous (Barlett test). Consequently, statistical analyses were performed with nonparametric tests. A nonparametric Tukey-type multiple comparisons (Nemenyi test) following nonparametric analysis of variance (Kruskall-Wallis test) were applied to indicate diversity of Hg content in soil substratum, and caps and stipes or whole fruiting bodies of specimens from the certain sampling sites surveyed. Further examination of differences in Hg concentrations between the caps (also stipes and soil) from certain localizations was made using Mann–Whitney U test.

Results and discussion

Hg in fruiting bodies

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Concentrations of total Hg in caps and stipes of Bay Bolete were high in specimens collected from Złotoryja (site 12 Z), and these concentrations differed significantly from samples collected at other locations (Figure 3). The arithmetic mean concentrations of Hg in Bay Bolete's cap varied between 0.051 ± 0.030 and $1.0 \pm 1.1 \mu\text{g g}^{-1}$ dw, depending on the location. The corresponding concentrations of Hg in stipes of Bay Bolete at these sites were between 0.027 ± 0.018 and $0.68 \pm 0.86 \mu\text{g Hg g}^{-1}$ dw. The whole fruiting bodies collected from Kłodzka Dale (14 KD) contained $0.26 \pm 0.06 \mu\text{g Hg g}^{-1}$ dw. A total Hg concentration of $1.0 \mu\text{g g}^{-1}$ dw ($0.1 \mu\text{g g}^{-1}$ wet weight) exceeded in two specimens from Włoszczowa (15 W) and two from Złotoryja (12 Z) (Table 1).

Mercury concentrations in Bay Bolete at sites that are considered as background (unpolluted) areas varied between 0.051 ± 0.030 and $0.32 \pm 0.17 \mu\text{g g}^{-1}$ dw in caps and between 0.027 ± 0.018 and $0.23 \pm 0.12 \mu\text{g g}^{-1}$ dw in stipes (Table 1). These concentrations are relatively small and comparable to a range of values between 0.26 ± 0.17 and $0.50 \pm 0.10 \mu\text{g g}^{-1}$ dw in caps, and between 0.089 ± 0.026 and $0.16 \pm 0.07 \mu\text{g g}^{-1}$ dw in stipes, reported for the mushroom Larch Bolete (*Suillus grevillei*) as well as between 0.095 ± 0.082 and $0.28 \pm 0.07 \mu\text{g g}^{-1}$ dw in caps and between 0.045 ± 0.018 and $0.13 \pm 0.02 \mu\text{g g}^{-1}$ dw in stipes reported for Slippery Jack (*Suillus luteus*) in Poland (Chudzyński et al., 2009 and 2011). The fungus, Bay Bolete, can be considered as a weak accumulator of Hg in fruiting bodies in comparison with King Bolete (*Boletus edulis*) which contained between 1.2 ± 1.4 and $7.6 \pm 3.1 \mu\text{g Hg g}^{-1}$ dw in caps or with Parasol Mushroom (*Macrolepiota procera*) which contained between 1.1 ± 1.0 and $8.4 \pm 7.4 \mu\text{g Hg g}^{-1}$ dw in caps (Falandysz and Gucia, 2008; Falandysz et al., 2007b and 2008a).

The caps of Bay Bolete, on average, contained higher concentrations of Hg than the stipes (Table 1). The median value for caps to stipe Hg concentration quotient (Q_{CS}), calculated for 221 specimens collected across Poland, was 1.7, and the arithmetic mean was 1.8 ± 0.8 (0.1-6.6).

1
2
3 138 Nevertheless, the total Hg content in caps of Bay Bolete was significantly correlated with the
4
5 139 content in stipes ($r = 0.95$; $p < 0.0001$; Figure 2).
6
7

8 140 ***Hg content in topsoil and its relation to concentrations in mushroom***
9

10 141 Soil Hg content is an important factor that affects the concentrations of this element in terrestrial
11
12 142 plants and edible mushrooms. A concentration of Hg of $1.0 \mu\text{g g}^{-1}$ dw has been suggested as a
13
14 143 reference value for agricultural soil (Rundgren et al., 1992). No such value has been reported for
15
16 144 forest soils in Poland. The median Hg concentrations in topsoil at most sampling sites in this
17
18 145 study were below $0.1 \mu\text{g g}^{-1}$ (Table 1). In our earlier surveys, we found that the median values of
19
20 146 total Hg concentrations in forest soils in central and northern Europe were below $0.1 \mu\text{g g}^{-1}$
21
22 147 (Chudzyński et al., 2009 and 2011; Falandysz and Bielawski, 2001 and 2007; Falandysz and
23
24 148 Brzostowski, 2007; Falandysz and Chwir, 1997; Falandysz et al., 2001a, 2002a-c, 2003a-d and
25
26 149 2004). It is known that the forest upper layer of soil that is rich in organic matter can easily
27
28 150 absorb Hg deposited from the atmosphere and from weathering of parent bedrocks (Suchara and
29
30 151 Sucharová, 2002). Nevertheless, the content of organic matter in the top 0-10 cm forest soil layer
31
32 152 vary greatly depending on the region, bedrock, vegetation, and age of the forest. The Critical
33
34 153 Limit (CL) proposed for Hg(II) in soil is $0.13 \mu\text{g g}^{-1}$ (when soil properties are taken into account,
35
36 154 the value ranges between 0.04 and $1.2 \mu\text{g g}^{-1}$). The soil organic matter (SOM) normalized CL is
37
38 155 $3.3 \mu\text{g Hg g}^{-1}$ of SOM (varies between 0.03 and $3.3 \mu\text{g g}^{-1}$). This CL value is greater than the
39
40 156 value of $0.5 \mu\text{g g}^{-1}$ SOM, used to estimate Hg effects in European soil ecosystems (Tipping et al.,
41
42 157 2010). The total Hg content in forest topsoil (0-10 cm layer) in this study, and in our earlier
43
44 158 studies across Poland, was well below the recently proposed CL value for Hg(II) of $0.13 \mu\text{g g}^{-1}$.
45
46
47
48
49
50
51
52
53

54 159 In earlier two studies, the concentrations of Hg in forest soils were found not dependent
55
56 160 on soil type and texture or were elevated in soils of loose sands and slightly clayey relatively rich
57
58
59
60

in humus type (Falandysz et al., 2002b and 2003a). It is worth to mention that no relation with Hg of fruiting bodies and organic matter content or pH of the soils could be observed for fungi in a several studies (Falandysz and Bielawski, 2007; Falandysz and Brzostowski, 2007). No such relation could be observed also between soils organic matter content or pH and metals such as Cd and Pb sequestered in fruiting bodies of several fungi (Gast et al., 1998), as well as of Ag, Al, Ba, Ca, Cd, Co, Cr, Cu, Fe, Hg, K, Mg, Mn, Na, Ni, Sr, Pb, Rb and Zn by Pioson Pax, *Paxillus involutus*, mushroom (Brzostowski et al., 2011a), while uptake and sequestration in fruiting bodies of metallic elements essential to fungi (K, Na, Ca, Mg, Zn, Cu) is largely regulated by species. In other words, either organic matter content or pH of soils was found as not important factors that can determine the trace metals bioavailability to fungi. Nevertheless, in the complex process of minerals gaining and absorption by mycelium, and in case of mycorrhizal fungi theirs further transfer to symbiotic plant (tree) and also transfer and sequestering in fruiting bodies there are an unknown underlying variables.

The soils Hg concentrations increase moving from the north to the south of Poland (Figure 3). This spatial trend can be related to the degree of industrialization and urbanization of the areas surveyed. Dolnośląskie Voivodeship (the sites nos. 11–14; Fig, 1) is a highly industrialized and urbanized region in Southern Poland and differences in soils bedrock composition between the regions. Furthermore, industrial activities are predominant in Świętokrzyskie Voivodeship (the sites nos. 15 and 16; Southern Poland), where soil Hg concentrations varied considerably (Figs. 1 and 3, Table 1). Of the 18 locations surveyed in this study, 17 locations exhibited no clear relationship between Hg concentration in Bay Bolete and corresponding soil Hg content. At Karpacz (13 K), a tendency of increasing content of Hg in caps/stipes with increasing soil Hg content was found.

A potential of minerals uptake by fungus from the soil and further sequestration in fruiting bodies is assessed using the common concept of bioconcentration factor (BCF) or otherwise transfer factor or enrichment factor. They are simply the mushroom's constituent concentration divided by the soil (or other substratum media) mineral constituent concentration. The value of BCF for a given trace element is a measure of its site-specific bioavailability for fungus.

Elevated concentration of Hg in topsoil and in parallel a good bioavailability are the reasons for great concentrations of Hg in fruiting bodies of Bay Bolete at the Złotoryja site (Table 1; site 12 Z). However, for the two highly contaminated specimens from the Włoszczowa site (15 W; containing 2.7 and 1.4 $\mu\text{g Hg g}^{-1}$ dw in caps and 0.89 and 0.87 $\mu\text{g Hg g}^{-1}$ dw in stipes), the driving force for elevated concentrations seems to be great Hg bioavailability (BCF was 80 and 16 for caps and 26 and 10 for stipes).

Hg content of topsoil from the Karpacz site was elevated (compared to the other locations surveyed), although Hg concentration in Bay Bolete in that site was low. This can be attributed to low bioavailability of Hg at this site. Content of methylmercury (MeHg) and its proportion to inorganic Hg in soils can impact amount of total Hg sequestered in fruiting bodies (Fischer et al., 1995). At the Włoszczowa site (15 W), relatively elevated Hg content was found (0.28 and 0.17 $\mu\text{g g}^{-1}$ dw) in topsoil, while concentrations in caps (0.081 and 0.073 $\mu\text{g g}^{-1}$ dw) and in stipes (0.017 and 0.040 $\mu\text{g g}^{-1}$ dw), were low (BCF between 0.3 and 0.4 for caps and 0.06 and 0.2 for stipes). Thus, the concentrations of Hg accumulated in fruiting bodies varied widely ($p < 0.05$; Mann-Whitney U test) among the locations surveyed (Table 1).

Bioconcentration potential

Our results indicate that Bay Bolete can bioconcentrate Hg from the soil. Mercury was found in Bay Bolete, even in unpolluted areas where the BCF (bioconcentration factor) in caps was up to

1
2
3 208 11±6 (Table 1). Bioconcentration factor or transfer factor (TF) is a quotient of Hg concentration
4
5 209 in caps or stipes to soil Hg concentration. BCF value below 1 suggests exclusion or bio-exclusion
6
7
8 210 whereas the value above 1 suggests bioconcentration. The values of BCF of Hg in Bay Bolete
9
10 211 decreased with increasing soil Hg concentration (Figure 4).
11

12 212 ***Hg intake rates***
13
14

15 213 Both cap and stipe of Bay Bolete are edible and accumulation of Hg in both parts of this
16
17 214 mushroom is a human health concern. In a well-grown fruiting body of this fungus, cap is much
18
19
20 215 larger in mass than stipe, while in young specimens, the proportion is approximately equal, and in
21
22 216 much younger specimens, stipe is larger than cap. Nevertheless, there is no published data on Hg
23
24 217 content in caps, stipes or whole fruiting bodies of Bay Bolete collected at different growth stages.
25
26
27 218 The mushroom collectors harvest forest mushrooms daily and individuals, who are late during a
28
29 219 mushroom collection day, collect even young fruiting bodies. There are no limits regarding
30
31 220 allowable size or allowable daily harvest for mushroom collection or a mushroom species that
32
33 221 could be collected by an individual in Poland. For the assessment of daily intake rate of Hg from
34
35 222 Bay Bolete, we assumed that the fruiting bodies were collected from mature specimens.
36
37

38 223 Although mushrooms do not constitute a significant portion of human diet, they are
39
40 224 popular food items worldwide especially during their seasonal abundance. The data regarding
41
42 225 intake rates of wild-grown edible mushrooms at local, regional, national, or international scale
43
44 226 are highly limited. For example, as discussed by Zhang et al. (2010), consumption rate of wild-
45
46 227 grown mushrooms in Sweden was 1 kg *per capita* annually and Common Chanterelle
47
48 228 (*Cantharellus cibarius*) is the most commonly consumed mushroom in that country. In rural
49
50 229 areas of Sweden, as well as in rural areas of many other European countries, wild-mushroom
51
52
53 230 picking is a common practice and the intake rates *per capita* are expected to be greater than 1 kg
54
55 231 annually. In Czech Republic, 72 % of families take part in wild-mushroom picking, with an
56
57
58
59
60

average yield of 7 kg fresh weight *per familia* annually, while annual intake rate by some individuals can be up to 10 kg *per capita* fresh weight. In Sichuan Province in China, the annual wild mushroom consumption rate exceeded 20-24 kg *per capita* (Zhang et al., 2010).

In Poland, a meal of up to 300 g of fresh mushroom is considered as the common intake rate in a single dish *per capita* during a week period among mushroom dish fanciers or individuals with low income rural household. A single dish of up to 500 g of fruiting bodies (fresh weight) collected in wild has been used as daily intake rate of mushrooms *per capita*. (Zhang et al., 2010).

Mushrooms, depending on the species, are cooked in several ways. Any mushroom, wild or cultivated, should not be eaten fresh. Blanching (boiling in water for 10 minutes) of fresh fruiting bodies is a common procedure. Blanched mushrooms could be further baked, used as soup ingredients, pickled, or frozen. Dried mushrooms could be grounded (powdered) and used as a condiment. Dried mushrooms can be soaked in water and added (together with or without broth) to certain traditional dishes and further cooked without loss of mineral constituents. Soaking of fresh fruiting bodies of Bay Bolete in salt (2% of table salt per fresh mushroom weight) water at 20 °C for 5-15 min resulted in 5-10 % loss of total Hg content, while boiling for 15 min resulted in 15 % loss of this element (Svoboda et al., 2001).

It is not known if both organic and inorganic forms of Hg are leached during blanching or other type of processing. A 15 % loss of total Hg contained in fruiting bodies during blanching of Bay Bolete is considered to be low (Svoboda et al., 2001). Based on the large variation in total Hg content in Bay Bolete (Table 1), the 15% loss rate from food processing step can be ignored in the assessment of Hg intake rates by consumers.

For the assessment of potential risks from intake of Hg in the fruiting bodies of Bay Bolete, a reference dose (RfD; 0.0003 mg kg⁻¹ bm daily) and a Provisional Tolerable Weekly

1
2
3 256 Intake values (PTWI; 0.005 mg kg⁻¹ bw) have been used (JECFA, 1978, US EPA, 1987). The
4
5 257 PTWI of 5 µg kg⁻¹ body weight for total Hg established by the JECFA in 1978 was revised in
6
7
8 258 2010 (JECFA, 2010). The JECFA established a PTWI of 0.004 mg kg⁻¹ bw for inorganic Hg in
9
10 259 2010, based on the assumption that the predominant form of Hg in foods, other than fish and
11
12 260 shellfish, is inorganic Hg. A meal made with 300-500 g of fresh caps of Bay Bolete collected
13
14 261 from the Złotoryja site can result in Hg intake of 0.00025-0.0004 mg kg⁻¹ bw (assuming a 60 kg
15
16 262 body weight), based on the median Hg concentration of 0.058 µg g⁻¹ wet weight. If the intake is
17
18 263 calculated from the arithmetic mean of 0.10 µg Hg g⁻¹ wet weight (assuming 90 % water content
19
20 264 in caps), the intake rate will be 0.00043-0.0007 mg kg⁻¹ bw.
21
22
23

24 265 Wild mushrooms are more frequently and at higher rate eaten in a mushrooming season in
25
26 266 late summer and early fall, while usually much less in other periods of the year. In a view of Hg
27
28 267 content, the consumption of Bay Boletes (caps or a whole mushrooms) at the regions surveyed at
29
30 268 least at a rate up to 2.5 kg *per capita* weekly in a mushrooming season is safe and will not result
31
32 269 in exceeding of currently allowable Hg intake doses.
33
34
35

36 270 Bay Bolete collected from the Sierakowice site (an unpolluted area) contained the greatest
37
38 271 Hg concentration (Table 1). Consumption of fresh caps of Bay Bolete collected at this site at a
39
40 272 rate up to 7.5 kg should not result in Hg intake values exceeding the PTWI limit for a person
41
42 273 weighing 60 kg, assuming that no Hg from other foods is ingested. Forest soil at the Złotoryja
43
44 274 and Karpacz sites contained elevated concentrations of Hg, and Bay Bolete at the Złotoryja site
45
46 275 contained high levels of Hg (Figure 3; Table 1). Consumption of up to 2.4 kg of fresh caps of Bay
47
48 276 Bolete from Złotoryja should not exceed the PTWI limit, if no Hg from other foods is ingested.
49
50
51

52 277 Selenium in mushroom can provide protective role against Hg (Jarzyńska and Falandysz,
53
54 278 2011b). It is presumed that co-occurrence of selenium and MeHg or total Hg in stoichiometric
55
56 279 ratio could prevent the toxic effects of Hg in foods (Yoneda and Suzuki, 1997; Stijve and
57
58
59
60

Cardinale, 1974; Stijve and Besson, 1976). A review of published data on selenium content in a limited number of Bay Bolete samples collected in Europe indicated that selenium content in fruiting bodies usually varies around $0.20 \mu\text{g g}^{-1}$ dw (Falandysz, 2008), which is comparable to Hg content found in Bay Bolete in this study (Table 1).

The PTWI for MeHg is $0.0016 \text{ mg kg}^{-1} \text{ bw}$ (JECFA, 2007). We did not analyze MeHg content in Bay Bolete in this study. Fischer et al. (1995) reported that 0.70 to 1.1 % of total Hg content in Bay Bolete is in the form of MeHg in a highly contaminated area ($15\text{--}35 \mu\text{g Hg g}^{-1}$ dw). Pilz et al. (2011) in a recent study reported on MeHg content of five species of fungi: an unknown fungus, *Cordoncella* mushroom (King Bolete?), *Pleurotus djamer*, *P. citrinopileatus* and *P. ostreatus*, where this compound varied a somehow between the species but largely varied depending on the extraction method used, i.e. by ultrasonic (from 3 to 38 % of total Hg) or microwave assisted extraction (from 15 to 41 % of total Hg). Data available on MeHg in wild mushrooms are sparse and controversial. As reviewed recently, reported MeHg content of King Bolete (*Boletus edulis*) varied highly (from 0.6 to 39 % of total Hg) between the authors (Falandysz, 2010).

Conclusions

Hg concentrations in pooled caps of Bay Bolete collected from unpolluted areas in Poland were below $1.0 \mu\text{g Hg g}^{-1}$ dw. Mercury concentrations in caps and stipes of Bay Bolete were strongly correlated ($p < 0.0001$). The relationship between the concentrations of Hg in soil and in mushrooms varied depending on the location. The bioconcentration potential of Hg in Bay Bolete at sites with elevated soil Hg concentrations was lower than that in soils with low Hg content. In a view of Hg content, the consumption of Bay Boletes (caps or a whole mushrooms) at the regions surveyed at least at a rate up to $2.5 \text{ kg per capita}$ weekly in a mushrooming season is safe and will not result in exceeding of currently allowable Hg intake doses.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

304

305 **Acknowledgement**

306 Technical assistance by Joanna Gozdek, Aleksandra Jabłońska, Ewa Łukaszewicz, Dominika Romińska,
307 Daniel Siwicki and Aleksandra Stefańska is acknowledged. This study was supported by the Ministry of
308 Science and Higher Education under the grant no. DS-8130-4-0092-12. This article is a part of a Ph. D.
309 thesis by A.K. Kojta

310

311 **References**

312 Alonso J, Salgado MJ, Garcíá, MÁ, Melgar MJ. 2000. Accumulation of mercury in edible
313 macrofungi: Influence of some factors. Arch. Environ. Contam. Toxicol. 44: 18-162.

314 Barcan V, Kovnatsky EF, Smetannikova MS. 1998. Absorption of heavy metals in wild berries
315 and edible mushrooms in area affected by smelter emissions. Water Air Soil Poll. 103: 173-
316 195.

317 Bargagli R, Baldi F. 1984. Mercury and methyl mercury in higher fungi and their relation with
318 substrata in a cinnabar mining area. Chemosphere. 13: 1059-1071.

319 Borovička J, Kotrba P, Gryndler M, Mihaljevič M, Řanda Z. Rohovec J, Cajthaml T, Stijve T,
320 Dunn CE. 2010. Bioaccumulation of silver in ectomycorrhizal and saprobic macrofungi from
321 pristine and polluted areas. Sci. Total Environ. 408: 2733-2744.

322

323 Brzostowski A, Bielawski L, Orlikowska A, Plichta S, Falandysz J. 2009. Instrumental analysis
324 of metals profile in Poison Pax (*Paxillus involutus*) collected at two sites in Bory Tucholskie.
325 Chem. Anal. (Warsaw). 54: 907-919.

- 326 Brzostowski A, Jarzyńska G, Falandysz J, Zhang D. 2011a. Bioconcentration potential of
327 metallic elements by Poison Pax (*Paxillus involutus*) mushroom. J. Environ. Sci. Health Part
328 A. 46: 378-393.
- 329 Brzostowski A, Jarzyńska G, Kojta AK, Wydmańska D, Falandysz J. 2011b. Variations in metal
330 levels accumulated in Poison Pax (*Paxillus involutus*) mushroom collected at one site over four
331 years. J. Environ. Sci. Health Part A. 46: 581-588.
- 332 Carvalho MI, Pimentel AC, Fernandes B. 2005. Study of heavy metals in wild edible mushrooms
333 under different pollution conditions by X-ray fluorescence spectrometry. Anal. Sci. 21: 747-
334 750.
- 335 Chudzyński K, Bielawski L, Falandysz J. 2009. Mercury bio-concentration potential of Larch
336 Bolete, *Suillus grevillei*, mushroom. Bull. Environ. Contam. Toxicol. 83: 275-279.
- 337 Chudzyński K, Falandysz J. 2008. Multivariate analysis of elements content of Larch Bolete
338 (*Suillus grevillei*) mushroom. Chemosphere. 73: 1230-1239.
- 339 Chudzyński K, Jarzyńska G, Stefańska A, Falandysz J. 2011. Mercury content and bio-
340 concentration potential of Slippery Jack, *Suillus luteus*, mushroom. Food Chem. 125: 986-990.
- 341 Cibulka J, Šišák L, Pulkrab K, Miholová D, Száková J, Fučíková A, Slámová A, Stéhulová J,
342 Barláková S. 1996. Cadmium, lead, mercury and caesium levels in wild mushrooms and forest
343 berries from different localities of Czech Republic. Sci. Agaricult. Bohem. 27: 113-129.
- 344 Doğan HH, Sanda MA, Uyanöz R, Oztürk C, Çetin Ü. 2006. Contents of metals in some wild
345 mushrooms: its impact in human health. Biol. Trace Elem. Res. 110: 79-94.
- 346 Falandysz J. 2002. Mercury in mushrooms and soil of the Tarnobrzaska Plain, south-eastern
347 Poland. J. Environ. Sci. Health Part A. 37: 343-352.
- 348 Falandysz J. 2008. Selenium in edible mushrooms. J. Environ. Sci. Health Part C. 26: 256-299.

- 349 Falandysz J. 2010. Progress in Mycology. Jodhpur: Scientific Publishers. Mercury in certain
350 mushrooms species in Poland; p. 349-383.
- 351 Falandysz J, Bielawski L. 2001. Mercury content of wild edible mushrooms collected near the
352 town of Augustow. Pol. J. Environ. Stud. 10: 67-71.
- 353 Falandysz J, Bielawski L. 2007. Mercury and its bioconcentration factors in Brown Birch Scaber
354 Stalk (*Leccinum scabrum*) from various sites in Poland. Food Chem. 105: 635-640.
- 355 Falandysz J, Bielawski L, Kannan K, Gucia M, Lipka K, Brzostowski A. 2002a. Mercury in wild
356 mushrooms and underlying soil substrate from the great lakes land in Poland. J. Environ.
357 Monit. 4: 473-476.
- 358 Falandysz J, Bona H, Danisiewicz D. 1994. Silver uptake by *Agaricus bisporus* from an
359 artificially enriched substrate. Zeitschr. Lebensm. Unters. Forsch. 199: 225-228.
- 360 Falandysz J, Brzostowski A. 2007. Mercury and its bioconcentration factors in Poison Pax
361 (*Paxillus involutus*) from various sites in Poland. J. Environ. Sci. Health Part A. 42: 1095-
362 1100.
- 363 Falandysz J, Brzostowski A, Kawano M, Kannan K, Puzyn T, Lipka K. 2003a. Concentrations of
364 mercury in wild growing higher fungi and underlying substrate near Lake Wdzydze, Poland.
365 Water Air Soil Pollut. 148: 127-137.
- 366 Falandysz J, Chwir A. 1997. The concentrations and bioconcentration factors of mercury in
367 mushrooms from Mierzeja Wislana sand-bar, Northern Poland. Sci. Total Environ. 203: 221-
368 228.
- 369 Falandysz J, Frankowska A, Jarzyńska G, Dryżałowska A, Kojta AK, Zhang D. 2011. Survey on
370 composition and bioconcentration potential of 12 metallic elements in King Bolete (*Boletus*
371 *edulis*) mushroom that emerged at 11 spatially distant sites. J. Environ. Sci. Health Part B. 46:
372 231-246.

- 373 Falandysz J, Frankowska A, Mazur A. 2007a. Mercury and its bioconcentration factors in King
374 Bolete (*Boletus edulis*). J. Environ. Sci. Health Part A. 42: 2089-2095.
- 375 Falandysz J, Gucia M. 2008. Bioconcentration factors of mercury by parasol mushroom
376 (*Macrolepiota procera*). Environ. Geochem. Health. 30: 121-125.
- 377 Falandysz J, Gucia M, Brzostowski A, Kawano M, Bielawski L, Frankowska A, Wyrzykowska
378 B. 2003b. Content and bioconcentration of mercury in mushrooms from northern Poland. Food
379 Addit. Contam., 20: 247-253.
- 380 Falandysz J, Gucia M, Frankowska A, Kawano M, Skwarzec B. 2001a. Total mercury in wild
381 mushrooms and underlying soil substrate from the city of Umeå and its surroundings, Sweden.
382 Bull. Environ. Contam. Toxicol. 67: 767-770.
- 383 Falandysz J, Gucia M, Mazur A. 2007b. Content and bioconcentration factors of mercury by
384 Parasol Mushroom *Macrolepiota procera*. J. Environ. Sci. Health Part B. 42: 735-740.
- 385 Falandysz J, Gucia M, Skwarzec B, Frankowska A, Klawikowska K (2002b) Total mercury in
386 mushrooms and underlying soil from the Borecka Forest, Northeastern Poland. Arch. Environ.
387 Contam. Toxicol. 42: 145-154.
- 388 Falandysz J, Jędrusiak A, Lipka K, Kannan K, Kawano M, Gucia M, Brzostowski A, Dadej M.
389 2004. Mercury in wild mushrooms and underlying soil substrate from Koszalin, North-central
390 Poland. Chemosphere. 54: 461-466.
- 391 Falandysz J, Kawano M, Świeczkowski A, Brzostowski A, Dadej M. 2003c. Total mercury in
392 wild-grown higher mushrooms and underlying soil from Wdzydze Landscape Park, Northern
393 Poland. Food Chem. 81: 21-26.
- 394 Falandysz J, Kunito T, Kubota R, Bielawski L, Frankowska A, Falandysz JJ, Tanabe S. 2008a.
395 Multivariate characterization of elements accumulated in King Bolete *Boletus edulis*

- 396 mushroom at lowland and high mountain regions. J. Environ. Sci. Health Part A. 43: 1692-
397 1699.
- 398 Falandysz J, Kunito T, Kubota R, Brzostowski A, Mazur A, Falandysz JJ, Tanabe S. 2007c.
399 Selected elements of Poison Pax *Paxillus involutus*. J. Environ. Sci. Health Part A. 42: 1161-
400 1169.
- 401 Falandysz J, Kunito T, Kubota R, Gucia M, Mazur A, Falandysz JJ, Tanabe S. 2008b. Some
402 mineral constituents of Parasol Mushroom *Macrolepiota procera*. J. Environ. Sci. Health Part
403 B. 43: 187-192.
- 404 Falandysz J, Kunito T, Kubota R, Lipka, K.; Mazur, A.; Falandysz, J.J.; Tanabe, S. 2007d.
405 Selected elements in Fly Agaric *Amanita muscaria*. J. Environ. Sci. Health Part A. 42: 1615-
406 1623.
- 407 Falandysz J, Lipka K, Gucia M, Kawano M, Strumnik K, Kannan K. 2002c. Accumulation
408 factors of mercury in mushrooms from Zaborski Landscape Park, Poland. Environ. Intern. 28:
409 421-427.
- 410 Falandysz J, Lipka K, Kawano M, Brzostowski A, Dadej M, Jędrusiak A, Puzyn T. 2003d.
411 Mercury content and its bioconcentration factors in wild mushrooms at Łukta and Morąg,
412 Northeastern Poland. J. Agric. Food Chem. 51: 2832-2836.
- 413 Falandysz, J.; Lipka, K.; Mazur, A. 2007e. Mercury and its bioconcentration factors in Fly Agaric
414 (*Amanita muscaria*) from spatially distant sites in Poland. J. Environ. Sci. Health Part A. 42:
415 1625-1630.
- 416 Falandysz J, Szymczyk K, Ichihashi H, Bielawski L, Gucia M, Frankowska A, Yamasaki S.
417 2001b. ICP/MS and ICP/AES elemental analysis (38 elements) of edible wild mushrooms
418 growing in Poland. Food Addit. Contam. 18: 503-513.

- 419 Fischer RG, Rapsomanikis S, Andreae MO, Baldini F. 1995. Bioaccumulation of methylmercury
420 and transformation of inorganic mercury by macrofungi. Environ. Sci. Technol. 29: 993-999.
- 421 Frankowska A, Ziółkowska J, Bielawski L, Falandysz J. 2010. Profile and bioconcentration of
422 minerals by King Bolete (*Boletes edulis*) from the Płocka Dale in Poland. Food Add. Contam.
423 Part B. 3: 1-6.
- 424 Gast CH, Jansen E, Bierling J, Haanstra L. 1988. Heavy metals in mushrooms and their
425 relationship with soil characteristics. Chemosphere. 17: 789-799.
- 426 Gucia M, Jarzyńska G, Kojta AK, Falandysz J. 2012a. Temporal variability in mineral substances
427 of Parasol Mushroom (*Macrolepiota procera*) collected in the same sites. J. Environ. Sci.
428 Health Part B. 47: in press.
- 429 Gucia M, Kojta AK, Jarzyńska G, Rafał E, Roszak M, Osiej I, Falandysz J. 2012b. Multivariate
430 analysis of mineral constituents of edible Parasol Mushroom (*Macrolepiota procera*) and soils
431 beneath fruiting bodies collected from Northern Poland. Environ. Sci. Pollut. Res. 18: doi.
432 10.1007/s11356-011-0574-5.
- 433 GUS: Informator i opracowania statystyczne, Leśnictwo 2009 [Internet]. 2009. Warszawa:
434 Główny Urząd Statystyczny, ISSN 1230-574x; [cited 2010]. Available from:
435 <http://www.stat.gov.pl>.
- 436 Jarzyńska G, Falandysz J. 2011a. The determination of mercury in mushrooms by CV-AAS and
437 ICP-AES techniques. J. Environ. Sci. Health Part A. 46: 569-573.
- 438 Jarzyńska G, Falandysz J. 2011b. Selenium and 17 other largely essential and toxic metals in
439 muscle and organ meats of Red Deer (*Cervus elaphus*) - Consequences to human health.
440 Environ. Intern. 37: 882-888.

- 441 Jarzyńska G, Gucia M, Kojta AK, Rezulak K, Falandysz J. 2011. Profile of trace elements in
442 Parasol Mushroom (*Macrolepiota procera*) from Tucholskie Forest. J. Environ. Sci. Health
443 Part B. 46: 741-751.
- 444 JECFA. 1978. Evaluation of certain food additives and contaminants. Twenty-second report of
445 the Joint FAO/WHO Expert Committee on Food Additives, WHO Technical Report Series
446 631.
- 447 JECFA. 2007. Evaluation of certain food additives and contaminants. Sixty-seventh report of the
448 Joint FAO/WHO Expert Committee on Food Additives, WHO Technical Report Series 940.
- 449 JECFA. 2010. Joint FAO/WHO Expert Committee on Food Additives. Seventy-second meeting.
450 Rome, 16–25 February 2010. Summary and Conclusions. JECFA/72/SC. Food and Agriculture
451 Organization of the United Nations and World Health Organization. 16th March 2010.
- 452 Kojta AK, Gucia M, Jarzyńska G, Lewandowska M, Zakrzewska A, Falandysz J, Zhang D.:
453 2011. Phosphorous and metallic elements in Parasol Mushroom (*Macrolepiota procera*) and
454 soil from the Augustowska Forest and Elk regions in north-eastern Poland. Fresen. Environ.
455 Bull. 20: 3044-3052.
- 456 Malinowska E, Szefer P, Falandysz J. 2004. Metals bioaccumulation by bay bolete, *Xerocomus*
457 *badius*, from selected sites in Poland. Food Chem. 84: 405-416.
- 458 Melgar MJ, Alonso J, Garcíá MÁ. 2009. Mercury in edible mushrooms and soil.
459 Bioconcentration factors and toxicological risk. Sci. Total Environ. 407: 5328-5334.
- 460 Melgar MJ, Alonso J, Pérez-López M, Garcia MÁ. 1998. Influence of some factors in toxicity
461 and accumulation of cadmium from edible wild macrofungi in NW Spain. J. Environ. Sci.
462 Health Part B. 33: 439-455.
- 463 Olivero J, Johnson B, Arguello E, 2002. Human exposure to mercury in San Jorge river basin,
464 Colombia (South America). Sci. Total Environ. 289: 41-47.

- 465 Pilz C, Santos CMM, Nunes NAC, Müller EI, Flores ÉMM, Dressler VI. 2011. Speciation of Hg
466 in edible mushroom. Paper presented at: International Symposium Trace Elements in Food.
467 TEF-4; 19-22 June 2011; King's College, Aberdeen, Scotland..
- 468 Rundgren S, Rühling Å, Schluter K, Tyler G. 1992. Mercury in Soil - distribution, speciation and
469 biological effects. A review of the literature and comments on critical concentrations. Nordic
470 Council of Ministers, Copenhagen.
- 471 Řanda Z, Kučera J. 2004. Trace elements in higher fungi (mushrooms) determined by activation
472 analysis. J. Radioanal. Nuclear Chem. 259: 99-107.
- 473 Stijve T, Besson R. 1976. Mercury, cadmium, lead and selenium content of mushroom species
474 belonging to the genus *Agaricus*. Chemosphere. 7: 151-158.
- 475 Stijve T, Cardinale E. 1974. Selenium and mercury content of some edible mushrooms. Trav.
476 chim. alim. d'hyg. 65: 476-478.
- 477 Stijve T, Goessler W, Dupuy G. 2004. Influence of soil particles on concentrations of aluminium,
478 iron, calcium and other metals in mushrooms. Deutsch. Lebensm-Rundsch. 100: 10-13.
- 479 Stijve T, Roschnik R. 1974. Mercury and methyl mercury content of different species of fungi.
480 Trav. chim. alim. d'hyg. 65: 209-220.
- 481 Suchara I, Sucharová J. 2002. Distribution of sulphur and heavy metals in forest floor humus of
482 the Czech Republic. Water Air Soil Pollut. 136: 289-316.
- 483 Svoboda L, Kalač P, Špička J, Janoušková D. 2001. Leaching of cadmium, lead and mercury
484 from fresh and differently preserved edible mushrooms, *Xerocomus badius*, during soaking and
485 boiling. Food Chem. 79: 41-45.
- 486 Tipping E, Lofts S, Hooper H, Frey B, Spurgeon D, Svendsen C. 2010. Critical Limits for Hg(II)
487 in soils, derived from chronic toxicity data. Environ. Pollut. 158: 2465-2471.

1
2
3 488 US EPA. 1987. *United States Environmental Protection Agency Peer Review Workshop on*
4
5 489 *Mercury Issues*. Summary Report. Environmental Criteria and Assessment Office. Cincinnati,
6
7 490 OH: U.S. EPA. October 26-27.
8
9
10 491 Yoneda S, Suzuki KT. 1997. Detoxification of mercury by selenium by binding of equimolar Hg-
11
12 492 Se complex to a specific plasma protein. *Toxicol. Appl. Pharmacol.* 143: 274-280.
13
14 493 Zhang D, Frankowska A, Jarzyńska G, Kojta AK, Drewnowska M, Wydmańska D, Bielawski L,
15
16 494 Wang J, Falandysz J. 2010. Metals of King Bolete (*Boletus edulis*) collected at the same site
17
18 495 over two years. *Afr. J. Agric. Res.* 5: 3050-3055.
19
20 496 Zimmermannova K, Svoboda L, Kalač P. 2001. Mercury, cadmium, lead and copper contents in
21
22 497 fruiting bodies of selected edible mushrooms in contaminated Middle Spiš region *Ekologia*
23
24 498 (Bratislava). 20: 440-446.
25
26
27
28
29
30
31

32 500 **Figure Captions**

33
34 501 Figure 1. Locations of the sampling sites of Bay Bolete in Poland. Abbreviations: The communes
35
36 502 of Sierakowice (1) and Lipusz in Pomorskie Voivodeship (2); Augustowska Forest in Podlaskie
37
38 503 Voivodeship (3); Olsztynek (4) Szczytno in Warmia and Mazury Voivodeship (5); Commune of
39
40 504 Olszewo-Borki in Mazowieckie Voivodeship (6); Notecka Forest (7), Porążyn (8), Rogalin (9)
41
42 505 Turek in Wielkopolska Voivodeship (10); Lower Silesia Forest (11), Złotoryja (12), Karpacz (13)
43
44 506 and Kłodzka Dale in Dolnośląskie Voivodeship (14); Włoszczowa (15) and Starachowickie
45
46 507 Forest in Świętokrzyskie Voivodeship (16); Poniatowa in Lubelskie Voivodeship (17);
47
48 508 Chochołowska Valley in Małopolska Voivodeship (18).
49
50
51
52 509
53
54
55 510 Figure 2. Plot of Hg concentrations in caps against its concentrations in stipes of 221 fruit bodies
56
57 511 of Bay Bolete collected from Poland ($y = 0.0231 + 1.4949 \cdot x$; $r = 0.95$; $r^2 = 0.89$; $p < 0.0001$).
58
59
60

512

513 Figure 3. Median values of Hg concentration in caps or in whole specimens of Bay Bolete and in

514 soil substrate collected under mushrooms: the sampling sites are displayed starting from the north

515 to the south of Poland. Abbreviations: **CS** (1) Mojusz in Commune of Sierakowice and **CL** (2)

516 Commune of Lipusz in Pomorskie Voivodeship; **AF** (3) Augustowska Forest in Podlaskie

517 Voivodeship; **O** (4) Olsztynek and **S** (5) Szczytno in Warmia and Mazury Voivodeship; **O-B** (6)

518 Commune of Olszewo-Borki in Mazowieckie Voivodeship; **NF** (7) Notecka Forest, **P** (8)

519 Porążyn, **R** (9) Rogalin and **T** (10) Turek in Wielkopolska Voivodeship; **LSF** (11) Lower Silesia

520 Forest, **Z** (12) Złotoryja, **K** (13) Karpacz and **KD** (14) Kłodzka Dale in Dolnośląskie

521 Voivodeship; **W** (15) Włoszczowa and **SF** (16) Lipie in Starachowickie Forest in Świętokrzyskie

522 Voivodeship; **P-LV** (17) Poniatowa in Lubelskie Voivodeship; **CH** (18) Chochołowska Valley in


523 Małopolska Voivodeship.

1 Table 1. Mean, standard deviation, range and median values of total mercury concentration in fruiting bodies of mushroom and soil,
2 and cap to stipe concentration quotient and bioconcentration factor (BCF) values for mercury.


Site, year and number of samples	Hg ($\mu\text{g g}^{-1}$ dw)			$Q_{C/S}$	BCF	
	Cap	Stipe	Soil		Cap	Stipe
1 (CS) Mojusz, Commune of Sierakowice, Pomorskie Voivodeship, 2007 $n = 12$	0.32±0.17 (0.13-0.65) 0.29	0.23±0.12 (0.081-0.43) 0.21	0.11±0.09 (0.024-0.26) 0.053	1.4±0.3 (0.95-2.1) 1.4	5.4±5.4 (0.86-17) 3.3	4.1±4.1 (0.42-13) 2.4
2 (CL) Commune Lipusz, Pomorskie Voivodeship, 2006 $n = 15$	0.14±0.09 (0.080-0.41) 0.10	0.070±0.02 (0.039-0.13) 0.063	0.046±0.033 (0.022-0.14) 0.037	2.0±1.3 (1.3-6.6) 1.6	3.9±2.5 (0.60-8.4) 3.1	2.0±1.3 (0.45-5.1) 1.6
3 (AF) Augustowska Forest, Podlaskie Voivodeship, 2007 $n = 15$	0.099±0.033 (0.062-0.20) 0.091	0.067±0.020 (0.042-0.12) 0.060	0.075±0.046 (0.020-0.19) 0.046	1.5±0.1 (1.3-1.7) 1.5	2.0±1.4 (0.54-4.5) 1.5	1.3±0.8 (0.37-2.8) 1.2
4 (O) Olsztynek, Warmia and Mazury Voivodeship, 2002 $n = 15$	0.080±0.020 (0.052-0.13) 0.077	0.044±0.013 (0.031-0.083) 0.040	0.051±0.043 (0.013-0.20) 0.041	1.9±0.4 (1.1-2.5) 1.9	2.2±1.3 (0.29-5.1) 2.1	1.2±0.7 (0.21-2.9) 1.1
5 (S) Szczytno, Warmia and Mazury Voivodeship, 2002 $n = 15$	0.061±0.013 (0.041-0.093) 0.062	0.040±0.009 (0.029-0.058) 0.039	0.040±0.010 (0.022-0.054) 0.042	1.6±0.2 (1.3-2.0) 1.6	1.6±0.6 (1.0-2.8) 1.4	1.1±0.4 (0.65-2.0) 0.97
6 (O-B) Commune of Olszewo – Borki, Mazowieckie Voivodeship, 2007 $n = 15$	0.20±0.09 (0.047-0.39) 0.18	0.12±0.06 (0.025-0.22) 0.12	0.026±0.022 (0.0063-0.079) 0.022	1.7±0.6 (1.2-3.8) 1.6	11±6 (1.6-22) 13	7.2±5.2 (1.0-19) 4.4
7 (NF) Notecka Forest, Wielkopolska Voivodeship, 2008 $n = 15$	0.057±0.028 (0.028-0.13) 0.055	0.048±0.023 (0.007-0.091) 0.043	0.038±0.027 (0.009-0.072) 0.028	1.5±1.0 (0.68-4.4) 0.98	2.0±1.3 (0.46-5.2) 1.7	1.9±1.5 (0.18-5.5) 1.9
8 (P) Porażyn, Wielkopolska Voivodeship, 2008 (15)	0.052±0.025 (0.011-0.092) 0.057	0.031±0.017 (0.009-0.073) 0.032	0.073±0.030 (0.010-0.12) 0.077	1.7±0.5 (0.31-2.8) 1.8	1.1±1.3 (0.16-5.4) 0.78	0.64±0.71 (0.074-3.0) 0.45
9 (R) Rogalin, Wielkopolska Voivodeship,	0.092±0.050 (0.019-0.19)	0.042±0.021 (0.013-0.088)	0.10±0.08 (0.010-0.27)	2.2±0.7 (0.67-3.2)	2.3±2.8 (0.12-8.8)	1.0±1.3 (0.091-4.1)

2008 $n = 14$	0.088	0.041	0.098	2.3	1.2	0.50
10 (T) Turek, Wielkopolska Voivodeship, 2001-2002 $n = 11$	0.051±0.030 (0.016-0.11)	0.027±0.018 (0.0055-0.057)	0.048±0.006 (0.038-0.057)	2.3±1.4 (1.1-5.0)	1.1±0.6 (0.40-2.0)	0.58±0.38 (0.12-1.1)
11 (LSF) Lower Silesia Forest, Dolnośląskie Voivodeship, 2008 $n = 11$	0.041 0.21±0.21 (0.071-0.78)	0.027 0.095±0.095 (0.026-0.37)	0.046 0.15±0.10 (0.044-0.41)	1.7 2.3±0.9 (1.3-4.2)	0.76 1.7±1.7 (0.44-5.0)	0.47 0.72±0.62 (0.22-2.3)
12 (Z) Złotoryja, Dolnośląskie Voivodeship, 2008 $n = 9$	0.13 1.0±1.1 (0.38-3.7)	0.079 0.68±0.86 (0.21-2.8)	0.12 0.37±0.16 (0.18-0.71)	2.0 1.0±0.7 (0.13-2.3)	1.0 3.2±3.1 (0.82-10)	0.49 2.0±2.4 (0.43-7.8)
13 (K) Karpacz, Dolnośląskie Voivodeship, 2008 $n = 15$	0.58 0.25±0.14 (0.087-0.53)	0.30 0.13±0.06 (0.059-0.24)	0.36 0.50±0.20 (0.16-0.77)	1.0 2.1±0.9 (0.39-4.5)	1.5 0.55±0.31 (0.19-1.2)	1.0 0.28±0.13 (0.12-0.49)
14 (KD) Kłodzka Dale, Sudety Mountains, Dolnośląskie Voivodeship, 2001 $n = 5^{**}$	0.19 0.26±0.06 (0.18-0.32)	0.092 0.15±0.01 ^{***} (0.13-0.17)	0.46 0.14	2.0 ND	0.49 1.3±0.5 (1.3-2.4)	0.28 1.8
15 (W) Włoszowa, Świętokrzyskie Voivodeship, 2007 $n = 15$	0.36±0.73 (0.064-2.7)	0.17±0.31 (0.017-0.88)	0.068±0.073 (0.011-0.28)	2.7±1.0 (1.6-4.8)	9.3±20.0 (0.29-80)	3.8±7.2 (0.059-26)
16 (SF) Lipie, Starachowickie Forest, Świętokrzyskie Voivodeship, 2000 $n = 15$	0.098 0.11±0.03 (0.073-0.20)	0.042 0.084±0.040 (0.038-0.17)	0.047 0.14±0.05 (0.074-0.21)	2.5 1.5±0.4 (0.55-2.1)	2.3 0.87±0.39 (0.34-1.7)	1.1 0.68±0.49 (0.18-2.2)
17 (P-LV) Poniatowa, Lubelskie Voivodeship, 2008 $n = 15$	0.11 0.16±0.04 (0.12-0.2)	0.075 0.082±0.028 (0.062-0.13)	0.15	1.6 2.0±0.5 (1.7-2.1)	0.75	0.51
18 (CH) Chochołowska Dale, Małopolska Voivodeship, Tatra Mountains, 2000 $n = 9$	0.14 0.24±0.10 (0.10-0.42)	0.070 0.15±0.01 (0.13-0.17)	ND	1.7 1.5±0.4 (0.84-2.4)	ND	ND
	0.24	0.14	ND	1.5	ND	ND


Notes: * Number of samples and number of specimens (in parentheses); ND, not determined; ** a whole fruiting bodies; *** 15 samples



30x26mm (600 x 600 DPI)



30x22mm (600 x 600 DPI)



30x22mm (600 x 600 DPI)