



**HAL**  
open science

# ON THE INVERSE MAPPING CLASS MONOIDS

Rym Karoui, Vladimir Vershinin

► **To cite this version:**

Rym Karoui, Vladimir Vershinin. ON THE INVERSE MAPPING CLASS MONOIDS. *Journal of Knot Theory and Its Ramifications*, 2010, 19 (2), pp.163-179. 10.1142/S0218216510007814. hal-00803326

**HAL Id: hal-00803326**

**<https://hal.science/hal-00803326v1>**

Submitted on 21 Mar 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON THE INVERSE MAPPING CLASS MONOIDS

R. KAROUI AND V. V. VERSHININ

ABSTRACT. Braid groups and mapping class groups have many features in common. Similarly to the notion of inverse braid monoid inverse mapping class monoid is defined. It concerns surfaces with punctures, but among given  $n$  punctures several can be omitted. This corresponds to braids where the number of strings is not fixed. In the paper we give the analogue of the Dehn-Nilsen-Baer theorem, propose a presentation of the inverse mapping class monoid for a punctured sphere and study the word problem. This shows that certain properties and objects based on mapping class groups may be extended to the inverse mapping class monoids. We also give an analogues of Artin presentation with two generators.

## CONTENTS

1. Introduction	1
2. Inverse braid monoids and inverse mapping class monoids	2
3. Properties of inverse mapping class monoids	6
4. Inverse mapping class monoids for punctured sphere	8
5. The word problem	11
6. Examples	13
7. Acknowledgments	13
References	13

## 1. INTRODUCTION

Mapping class group is an important object in Topology, Complex Analysis, Algebraic Geometry and other domains. It is a lucky case when the method of Algebraic Topology works perfectly well, the application of the functor of fundamental group completely solves the topological problem: group of isotopy classes of homeomorphisms is described in terms of automorphisms of the fundamental group of the corresponding surface, as states the Dehn-Nilsen-Baer theorem, see [11], for example.

Let  $S_{g,b,n}$  be an oriented surface of the genus  $g$  with  $b$  boundary components and a set  $Q_n$  of  $n$  fixed points. Consider the group  $\text{Homeo}(S_{g,b,n})$  of orientation preserving self-homeomorphisms of  $S_{g,b,n}$  which fix pointwise the boundary (if it exists) and map the set  $Q_n$  into itself. Orientation reversing homeomorphisms also possible to consider, see [4], for example, but for the simplicity of exposition we restrict ourselves to orientation preserving case. Let  $\text{Homeo}^0(S_{g,b,n})$  be the normal subgroup of self-homeomorphisms of  $S_{g,b,n}$  which are isotopic to identity. Then the *mapping class group*  $\mathcal{M}_{g,b,n}$  is defined as a quotient group

---

2000 *Mathematics Subject Classification.* Primary 20F38; Secondary 20F36, 57M.

*Key words and phrases.* Inverse mapping class monoid, mapping class group, braid, inverse braid monoid, presentation.

The last author is partially supported by the Russian-Indian Grant 08-01-91300.




FIGURE 2.1

$$\mathcal{M}_{g,b,n} = \text{Homeo}(S_{g,b,n}) / \text{Homeo}^0(S_{g,b,n})$$

These groups are connected closely with braid groups. In [16] W. Magnus interpreted the braid group as the mapping class group of a punctured disc with the fixed boundary. Braid groups have a variety of generalizations, see [25], for example. One of generalizations is the *inverse braid monoid*  $IB_n$  constructed by D. Easdown and T. G. Lavers [5].

The notion of *inverse semigroup* was introduced by V. V. Wagner in 1952 [27]. By definition it means that for any element  $a$  of a semigroup (monoid)  $M$  there exists a unique element  $b$  (which is called *inverse*) such that

$$a = aba$$

and

$$b = bab.$$

The typical example of an inverse monoid is a monoid of partial (defined on a subset) injections of a set. For a finite set this gives us the notion of a *symmetric inverse monoid*  $I_n$  which generalizes and includes the classical symmetric group  $\Sigma_n$ . A presentation of symmetric inverse monoid was obtained by L. M. Popova [22], see also formulas (2.1), (2.3-2.4) below.

## 2. INVERSE BRAID MONOIDS AND INVERSE MAPPING CLASS MONOIDS

Inverse braid monoid arises from a very natural operation on braids: deleting several strands. By the application of this procedure to braids in  $Br_n$  we get *partial* braids [5]. Applying the standard procedure of multiplication of braids (concatenation) to partial braids we need to eliminate all not complete strands, as it is shown at Figure 2.1. If all strands are eliminated we get an *empty* braid. The set of all partial braids with this operation forms an inverse braid monoid  $IB_n$ .

Usually the braid group  $Br_n$  is given by the following Artin presentation [1]. It has the generators  $\sigma_i$ ,  $i = 1, \dots, n-1$ , and two types of relations:

$$(2.1) \quad \begin{cases} \sigma_i \sigma_j & = \sigma_j \sigma_i, & \text{if } |i-j| > 1, \\ \sigma_i \sigma_{i+1} \sigma_i & = \sigma_{i+1} \sigma_i \sigma_{i+1}. \end{cases}$$

There exist other presentations of the braid group. Let

$$\sigma = \sigma_1 \sigma_2 \dots \sigma_{n-1},$$




FIGURE 2.2

then the group  $Br_n$  is generated by  $\sigma_1$  and  $\sigma$  because

$$\sigma_{i+1} = \sigma^i \sigma_1 \sigma^{-i}, \quad i = 1, \dots, n-2.$$

The relations for the generators  $\sigma_1$  and  $\sigma$  are the following

$$(2.2) \quad \begin{cases} \sigma_1 \sigma^i \sigma_1 \sigma^{-i} &= \sigma^i \sigma_1 \sigma^{-i} \sigma_1 \text{ for } 2 \leq i \leq n/2, \\ \sigma^n &= (\sigma \sigma_1)^{n-1}. \end{cases}$$

The presentation (2.2) was given by Artin in the initial paper [1]. This presentation was also mentioned in the books by F. Klein [13] and by H. S. M. Coxeter and W. O. J. Moser [3].

Classical braid group  $Br_n$  can be defined also as the mapping class group of a disc  $D^2$  with  $n$  points deleted (or fixed) and with its boundary fixed, or as the subgroup of the automorphism group of a free group  $\text{Aut } F_n$ , generated by the following automorphisms:

$$\begin{cases} x_i &\mapsto x_{i+1}, \\ x_{i+1} &\mapsto x_{i+1}^{-1} x_i x_{i+1}, \\ x_j &\mapsto x_j, j \neq i, i+1. \end{cases}$$

Geometrically this action is depicted in Figure 2.2, where  $x_i$  correspond to the canonical loops on  $D^2$  which form the generators of the fundamental group of the punctured disc.

This topological interpretation of the braid group was continued in [26] for the inverse braid monoid. Let  $Q_n$  the set of  $n$  fixed points of a disc  $D^2$ . The fundamental group of  $D^2$  with these points deleted is isomorphic to  $F_n$ . Consider homeomorphisms of  $D^2$  onto a copy of the same disc with the condition that only  $k$  points of  $Q_n$ ,  $k \leq n$  (say  $i_1, \dots, i_k$ ) are mapped bijectively onto the  $k$  points (say  $j_1, \dots, j_k$ ) of the second copy of  $D^2$ ,  $k$  varies from 0 to  $n$  and  $n$  is fixed. Consider the isotopy classes of such homeomorphisms and denote the set of them by  $IM_n(D^2)$ . Evidently it is a monoid.

**Theorem 2.1.** [26] *The monoids  $IB_n$  and  $IM_n(D^2)$  are isomorphic.*

The following presentation for the inverse braid monoid was obtained in [5]. It has the generators  $\sigma_i, \sigma_i^{-1}$ ,  $i = 1, \dots, n-1$ ,  $\epsilon$ , and relations

$$(2.3) \quad \begin{cases} \sigma_i \sigma_i^{-1} = \sigma_i^{-1} \sigma_i = 1, \text{ for all } i, \\ \epsilon \sigma_i = \sigma_i \epsilon \text{ for } i \geq 2, \\ \epsilon \sigma_1 \epsilon = \sigma_1 \epsilon \sigma_1 \epsilon = \epsilon \sigma_1 \epsilon \sigma_1, \\ \epsilon = \epsilon^2 = \epsilon \sigma_1^2 = \sigma_1^2 \epsilon \end{cases}$$

and the braid relations (2.1).

Geometrically the generator  $\epsilon$  means that the first strand in the trivial braid is absent.

If we replace the first relation in (2.3) by the following set of relations

$$(2.4) \quad \sigma_i^2 = 1, \text{ for all } i,$$

and delete the superfluous relations

$$\epsilon = \epsilon\sigma_1^2 = \sigma_1^2\epsilon,$$

we get a presentation of the symmetric inverse monoid  $I_n$  [22]. We also can simply add the relations (2.4) if we do not worry about redundant relations. We get a canonical map [5]

$$(2.5) \quad \tau_n : IB_n \rightarrow I_n,$$

which is a natural extension of the corresponding map for the braid and symmetric groups.

More balanced relations for the inverse braid monoid were obtained in [9]. Let  $\epsilon_i$  denote the trivial braid with  $i$ th strand deleted, formally:

$$\begin{cases} \epsilon_1 &= \epsilon, \\ \epsilon_{i+1} &= \sigma_i^{\pm 1} \epsilon_i \sigma_i^{\pm 1}. \end{cases}$$

The generators are:  $\sigma_i, \sigma_i^{-1}, i = 1, \dots, n-1, \epsilon_i, i = 1, \dots, n$ , and relations are the following:

$$(2.6) \quad \begin{cases} \sigma_i \sigma_i^{-1} = \sigma_i^{-1} \sigma_i = 1, \text{ for all } i, \\ \epsilon_j \sigma_i = \sigma_i \epsilon_j, \text{ for } j \neq i, i+1, \\ \epsilon_i \sigma_i = \sigma_i \epsilon_{i+1}, \\ \epsilon_{i+1} \sigma_i = \sigma_i \epsilon_i, \\ \epsilon_i = \epsilon_i^2, \\ \epsilon_{i+1} \sigma_i^2 = \sigma_i^2 \epsilon_{i+1} = \epsilon_{i+1}, \\ \epsilon_i \epsilon_{i+1} \sigma_i = \sigma_i \epsilon_i \epsilon_{i+1} = \epsilon_i \epsilon_{i+1}, \end{cases}$$

plus the braid relations (2.1).

If we take the presentation (2.2) for the braid group we get a presentation of the inverse braid monoid with generators  $\sigma_1, \sigma_1^{-1}, \sigma, \sigma^{-1}, \epsilon$ , and relations [26]:

$$(2.7) \quad \begin{cases} \sigma_1 \sigma_1^{-1} = \sigma_1^{-1} \sigma_1 = 1, \\ \sigma \sigma^{-1} = \sigma^{-1} \sigma = 1, \\ \epsilon \sigma^i \sigma_1 \sigma^{-i} = \sigma^i \sigma_1 \sigma^{-i} \epsilon \text{ for } 1 \leq i \leq n-2, \\ \epsilon \sigma_1 \epsilon = \sigma_1 \epsilon \sigma_1 \epsilon = \epsilon \sigma_1 \epsilon \sigma_1, \\ \epsilon = \epsilon^2 = \epsilon \sigma_1^2 = \sigma_1^2 \epsilon, \end{cases}$$

plus (2.2).

Similar to braids the notion of *mapping class monoid* was introduced in [26]. The definition is as follows.

It is convenient to consider the surface  $S_{g,b,n}$ , sometimes with  $n$  points fixed, sometimes deleted. Let  $f$  be a homeomorphism of  $S_{g,b,n}$  which maps  $k$  points,  $k \leq n$ , from  $Q_n: \{i_1, \dots, i_k\}$  to  $k$  points  $\{j_1, \dots, j_k\}$  also from  $Q_n$ . The same way let  $h$  be a homeomorphism of  $S_{g,b,n}$  which maps  $l$  points,  $l \leq n$ , from  $Q_n$ , say  $\{s_1, \dots, s_l\}$  to  $l$  points  $\{t_1, \dots, t_l\}$  again from  $Q_n$ . Consider the intersection of the sets  $\{j_1, \dots, j_k\}$  and  $\{s_1, \dots, s_l\}$ , let it be the set of cardinality

$m$ , ( $m \leq k$ ), it may be empty. Then the composition of  $f$  and  $h$  maps  $m$  points of  $Q_n$  to  $m$  points (may be different) of  $Q_n$ . For example, suppose for the simplicity of notations that

$$\{j_1, \dots, j_k\} \cap \{s_1, \dots, s_l\} = \{j_1, \dots, j_m\} = \{s_1, \dots, s_m\},$$

then the composition  $h \circ f$  maps the points  $\{i_1, \dots, i_m\}$  to  $\{t_1, \dots, t_m\}$ . If  $m = 0$  then the composition has no relation to the set  $Q_n$ . Denote the set of isotopy classes of such maps by  $\mathcal{IM}_{g,b,n}$ . Composition defines a structure of monoid on  $\mathcal{IM}_{g,b,n}$ . It is evident that the monoid  $\mathcal{IM}_{g,b,n}$  is inverse, so we call it the *inverse mapping class monoid*. If  $g = 0$  and  $b = 1$  we get the inverse braid monoid. In the general case of  $\mathcal{IM}_{g,b,n}$  the role of the empty braid plays the mapping class group  $\mathcal{M}_{g,b}$  (without fixed points). Each element  $f \in \mathcal{IM}_{g,b,n}$  corresponds a (partial) bijection

$$\{i_1, \dots, i_k\} \rightarrow \{j_1, \dots, j_k\}.$$

This defines a canonical homomorphism to the symmetric inverse monoid:

$$(2.8) \quad \tau_{g,b,n} : \mathcal{IM}_{g,b,n} \rightarrow I_n.$$

We recall that a monoid  $M$  is *factorisable* if  $M = EG$  where  $E$  is a set of idempotents of  $M$  and  $G$  is a subgroup of  $M$ .

**Proposition 2.1.** [26] *The monoid  $\mathcal{IM}_{g,b,n}$  can be written in the form*

$$\mathcal{IM}_{g,b,n} = E\mathcal{M}_{g,b,n},$$

where  $E$  is a set of idempotents of  $\mathcal{IM}_{g,b,n}$  and  $\mathcal{M}_{g,b,n}$  is the corresponding mapping class group. So, this monoid is factorisable.

Similar to braids we introduce the idempotent elements  $\epsilon_i \in \mathcal{IM}_{g,b,n}$  as isotopy classes of identity map

$$Id : S_{g,b} \rightarrow S_{g,b},$$

where during isotopy all points of the set  $Q_n$  are fixed with the exception of the point with number  $i$ . The element  $\epsilon_1$  is denoted by  $\epsilon$ .

We call an element  $m$  of the mapping class monoid  $\mathcal{IM}_{g,b,n}$   *$i$ -Makanin* or  *$i$ -Brunnian* if it satisfies the equation:

$$(2.9) \quad \epsilon_i m = \epsilon_i.$$

Geometrically this means that if we fill the deleted point  $i$  at the surface, then a homeomorphism  $h$  lying in the class  $m$

$$h : S_{g,b} \rightarrow S_{g,b},$$

becomes isotopical to the identity map

$$Id : S_{g,b} \setminus \{1, \dots, \hat{i}, \dots, n\} \rightarrow S_{g,b} \setminus \{1, \dots, \hat{i}, \dots, n\}.$$

The condition (2.9) is equivalent to the condition

$$m\epsilon_{\tau(m)(i)} = \epsilon_{\tau(m)(i)},$$

where  $\tau$  is the canonical map to the symmetric inverse monoid (2.8). With the exception of  $\epsilon_i$  itself all such elements belong to the mapping class group  $\mathcal{M}_{g,b,n}$ . We denote the subgroup of  $i$ -Makanin elements of the mapping class group by  $A_i$ . The subgroups  $A_i$ ,  $i = 1, \dots, n$ , are conjugate. The intersection of all subgroups of  $i$ -Makanin elements is the of *Makanin* or *Brunnian* subgroup of the mapping class group

$$\text{Mak}_{g,b,n} = \bigcap_{i=1}^n A_i.$$

That is the same as  $m \in \text{Mak}_{g,b,n}$  if and only if the equation (2.9) holds for all  $i$ . Certain properties of *Makanin* subgroups of the mapping class groups are studied in the work Benson Farb, Christopher J. Leininger and Dan Margalit [7]

The purpose of this paper is to develop further the theory of inverse mapping class monoids, to demonstrate that canonical properties of mapping class groups often have there smooth continuation for the inverse mapping class monoid  $\mathcal{IM}_{g,b,n}$ .

### 3. PROPERTIES OF INVERSE MAPPING CLASS MONOIDS

Following the ideology of considering partial symmetries instead of global ones [15] we can define the monoid of partial automorphisms of a group as follows. For a group  $G$  consider a set of partial isomorphisms

$$f : H \rightarrow K,$$

where  $H, K$  are subgroups of  $G$ . The composition of  $f$  with the isomorphism

$$g : L \rightarrow M,$$

( $L, M$  are subgroups of  $G$ ) is a superposition of  $f$  and  $g$  which is defined on  $(K \cap L)f^{-1}$ :

$$fg : (K \cap L)f^{-1} \rightarrow (K \cap L)g,$$

The set of all such partial isomorphisms with this operation form a monoid which is evidently inverse and which we call the *inverse partial automorphism monoid of a group  $G$*  and denote by  $IPA(G)$ .

In the case when the group  $G$  is a finitely generated free group  $F_n$  the following submonoid  $EF_n$  of  $IPA(F_n)$  was defined in [26]. Let  $a$  be an element of the symmetric inverse monoid  $I_n$ ,  $a \in I_n$ , so  $a$  is a partial injection with the elements  $i_1, \dots, i_k$  as its domain of the definition. Denote by  $J_k = \{j_1, \dots, j_k\}$  the image of  $a$ . The monoid  $EF_n$  consists of isomorphisms

$$\langle x_{i_1}, \dots, x_{i_k} \rangle \rightarrow \langle x_{j_1}, \dots, x_{j_k} \rangle$$

expressed by

$$f_a : x_i \mapsto w_i^{-1} x_{a(i)} w_i,$$

if  $i$  is among  $i_1, \dots, i_k$  and not defined otherwise and  $w_i$  is a word on  $x_{j_1}, \dots, x_{j_k}$ . The composition of  $f_a$  and  $g_b$ ,  $a, b \in I_n$ , is defined for  $x_i$  belonging to the domain of  $a \circ b$ . We put  $x_{j_m} = 1$  in a word  $w_i$  if  $x_{j_m}$  does not belong to the domain of definition of  $g$ . We define a map  $\phi_n$  from  $IB_n$  to  $EF_n$  expanding the canonical inclusion

$$Br_n \rightarrow \text{Aut } F_n$$

by the condition that  $\phi_n(\epsilon)$  as a partial isomorphism of  $F_n$  is given by the formula

$$\phi_n(\epsilon)(x_i) = \begin{cases} x_i & \text{if } i \geq 2, \\ \text{undefined,} & \text{if } i = 1. \end{cases}$$

Using the presentation (2.3) we see that  $\phi_n$  is a well defined homomorphism of monoids

$$\phi_n : IB_n \rightarrow EF_n.$$

The following statement was proved in [26].

**Theorem 3.1.** *The homomorphism  $\phi_n$  is a monomorphism.*

As usual, we define the  $(g, b, n)$ -*surface group* as a group with the presentation

$$\pi_{g,b,n} = \langle a_1, c_1, \dots, a_g, c_g, v_1, \dots, v_b, u_1, \dots, u_n \mid \prod_{i=1}^n u_i \prod_{l=1}^b v_l \prod_{m=1}^g [a_m, c_m] \rangle .$$

It is the fundamental group of a surface of genus  $g$  with  $b$  components of a boundary and  $n$  punctures. The generators  $a_i, c_i$  correspond to the genus of a surface, the generators  $v_j$  correspond to boundary components and the generators  $u_k$  correspond to punctures.

In our construction of the inverse mapping class monoid  $\mathcal{IM}_{g,b,n}$  we are passing from maps which are fixing a set of  $n$  points to maps which are fixing the set with a smaller number of points. If we consider these points deleted it means we are filling the holes. On the level of fundamental groups this means passing to quotient group.

Let  $H$  be a quotient group of  $\pi_{g,b,n}$ , defined by the conditions

$$u_i = 1 \text{ for all } i \notin \{i_1, \dots, i_k\},$$

and let  $K$  be a quotient group of  $\pi_{g,b,n}$ , defined by the conditions

$$u_j = 1 \text{ for all } j \notin \{j_1, \dots, j_k\}.$$

Let  $t$  be an element of the symmetric inverse monoid  $I_n$ ,  $t \in I_n$ , so  $t$  is an injection with elements  $i_1, \dots, i_k$  constituting the domain of the definition of  $t$ . Let  $J_k = \{j_1, \dots, j_k\}$  be the image of  $t$ . Let  $w_i$  be a word on letters  $a_1, c_1, \dots, a_g, c_g, v_1, \dots, v_b, u_{j_1}, \dots, u_{j_k}$ . Let  $IAut \pi_{g,b,n}$  be the monoid consisting of isomorphisms

$$f_t : H \rightarrow K,$$

such that

$$\begin{cases} f_t(v_m) = v_m \text{ for } m = 1, \dots, b, \\ f_t(u_i) = w_i^{-1} u_{(i)t} w_i, \text{ if } i \text{ is among } i_1, \dots, i_k, \end{cases}$$

for all subgroups  $H$  and  $K$  of the type defined above, for all  $k = 0, 1, \dots, n$ . Here the index  $(i)t$  of the element  $u_{(i)t}$  means the image of the element  $i$  by the action of the injection  $t$ , so  $(i)t \in J_k$ .

The composition of  $f_t$  and  $g_s$ ,  $t, s \in I_n$ , is defined for  $u_i$  belonging to the domain of  $t \circ s$ . We put  $u_{j_m} = 1$  in a word  $w_i$  if  $u_{j_m}$  does not belong to the domain of definition of  $g_s$ .

By our construction to each element of  $f_t \in IAut \pi_{g,b,n}$  the element  $t \in I_n$  is associated. This gives a canonical homomorphism

$$(3.1) \quad \tau_{g,b,n} : IAut \pi_{g,b,n} \rightarrow I_n.$$

For the simplicity of exposition we restrict ourselves to the case of empty boundary which was described in the book of W. Magnus, A. Karrass and D. Solitar [18, Sec. 3.7], and we denote  $\pi_{g,0,n}$  by  $\pi_{g,n}$ . Let us define an equivalence relation in  $IAut \pi_{g,n}$ : for  $f_1$  and  $f_2$

$$f_1, f_2 : H \rightarrow K.$$

Such isomorphisms  $f_1$  and  $f_2$  are equivalent if they differ by an inner automorphism of the group  $H$ , their domain of definition, i.e. if there exists an element  $q \in H$  such that

$$f_1(q^{-1}xq) = f_2(x).$$

This equivalence is in fact a congruence on the monoid  $IAut \pi_{g,n}$  and we denote the corresponding quotient monoid by  $IOut \pi_{g,n}$ . This is similar to the classical case of groups when a



homomorphism is defined from a mapping class group of a surface to the group of outer automorphisms of the fundamental group of the surface. The homomorphism  $\tau_{g,b,n} : IAut \pi_{g,b,n} \rightarrow I_n$  of (3.1) factors through  $IOut \pi_{g,n}$  and we have a homomorphism

$$(3.2) \quad \tau_{g,b,n} : IOut \pi_{g,b,n} \rightarrow I_n.$$

Let us take an arbitrary element  $\eta$  of  $\mathcal{IM}_{g,n}$ . It is represented by a homeomorphism of a surface with  $k$  points deleted (among our  $n$  fixed points) onto another copy of the surface with (may be) different  $k$  points deleted

$$h : S_{g,n} \setminus \{i_1, \dots, i_k\} \rightarrow S_{g,n} \setminus \{j_1, \dots, j_k\}.$$

It defines the bijection  $\hat{h}$  between the conjugacy classes of  $\pi_1(S_{g,n} \setminus \{i_1, \dots, i_k\})$  and  $\pi_1(S_{g,n} \setminus \{j_1, \dots, j_k\})$ . We define  $\hat{h}$  as an algebraic image of  $\eta$  in  $IOut \pi_{g,n}$ . Formally a homomorphism of monoids

$$\psi_{g,n} : \mathcal{IM}_{g,n} \rightarrow IOut \pi_{g,n}.$$

is defined by the formula

$$\psi_{g,n}(\eta) = \hat{h}.$$

This homomorphism is compatible with homomorphisms  $\tau_{g,b,n}$  of (2.8) and (3.2).

**Theorem 3.2.** *The homomorphism  $\psi_n$  is an isomorphism of monoids.*

*Proof.* The monoid  $\mathcal{IM}_{g,n}$  as a set is a disjoint union of copies of mapping class groups  $\mathcal{M}_{g,k}$ , for  $k = 0, \dots, n$ , namely,  $C_n^k$  copies of  $\mathcal{M}_{g,k}$  for each  $k$ . Our constructions are done so that for each copy of  $\mathcal{M}_{g,k}$  the map  $\psi_{g,n}$  is a bijection because of the Dehn-Nilsen-Baer theorem.  $\square$

#### 4. INVERSE MAPPING CLASS MONOIDS FOR PUNCTURED SPHERE

The following presentation for the mapping class group of a punctured sphere  $\mathcal{M}_{0,n}$  was obtained by W. Magnus [16], see also [18]. Let  $\sigma_1, \dots, \sigma_{n-1}$ , denote the classes of homeomorphisms such that  $\sigma_i$  locally interchanges, say clockwise, the punctures with numbers  $i$  and  $i+1$ . Then the presentation has generators  $\sigma_1, \dots, \sigma_{n-1}$ , which satisfy the braid relations (2.1), the sphere relation

$$(4.1) \quad \sigma_1 \sigma_2 \dots \sigma_{n-2} \sigma_{n-1}^2 \sigma_{n-2} \dots \sigma_2 \sigma_1 = 1$$

and the following relation

$$(4.2) \quad (\sigma_1 \sigma_2 \dots \sigma_{n-2} \sigma_{n-1})^n = 1.$$

Let  $\Delta$  be the Garside's fundamental word in the braid group  $Br_n$  [8]. It can be in particular defined by the formula:

$$\Delta = \sigma_1 \dots \sigma_{n-1} \sigma_1 \dots \sigma_{n-2} \dots \sigma_1 \sigma_2 \sigma_1.$$

If we use Garside's notation  $\Pi_t \equiv \sigma_1 \dots \sigma_t$ , then  $\Delta \equiv \Pi_{n-1} \dots \Pi_1$ . If the generators  $\sigma_1, \sigma_2, \dots, \sigma_{n-2}, \sigma_{n-1}$ , are subject to the braid relations (2.1), then the condition (4.2) is equivalent to the following

$$\Delta^2 = 1.$$

If we consider the presentation of the braid group with two generators the sphere relation has the form

$$\sigma^n (\sigma_1^{-1} \sigma)^{1-n} = 1,$$

and the sphere mapping class relation is

$$\sigma^n = 1.$$

So, the mapping class group  $\mathcal{M}_{0,n}$  in two generators is described by the braid relations and the following two relations

$$(4.3) \quad \begin{cases} \sigma^n = 1, \\ (\sigma_1^{-1}\sigma)^{n-1} = 1. \end{cases}$$

**Theorem 4.1.** *We get a presentation of the inverse mapping class monoid for punctured sphere  $\mathcal{IM}_{0,n}$  if we take the generators  $\sigma_1, \sigma_2, \dots, \sigma_{n-2}, \sigma_{n-1}, \sigma_1^{-1}, \sigma_2^{-1}, \dots, \sigma_{n-2}^{-1}, \sigma_{n-1}^{-1}, \epsilon$  (or  $\epsilon_1, \epsilon_2, \dots, \epsilon_{n-1}, \epsilon_n$  instead of one  $\epsilon$ ) subject to the sphere braid relations (2.1), (4.1), the sphere mapping class relation (4.2) and the inverse braid relations (2.3) (or (2.6)).*

*Proof.* We use the fact that the mapping class group  $\mathcal{M}_{0,0,n}$  is a quotient group of the  $n$ -strand braid group for sphere and the ideas of the proof of the presentation (2.3) for the inverse braid monoid  $IB_n$  in the work [5].

Denote temporarily by  $P_n$  the monoid defined by the presentation by the generators  $\sigma_1, \sigma_2, \dots, \sigma_{n-2}, \sigma_{n-1}, \sigma_1^{-1}, \sigma_2^{-1}, \dots, \sigma_{n-2}^{-1}, \sigma_{n-1}^{-1}, \epsilon$  the sphere braid relations (2.1), (4.1), the sphere mapping class relation (4.2) and the inverse braid relations (2.3). To define a map

$$\Theta : P_n \rightarrow \mathcal{M}_{0,n}$$

we associate to each word in the alphabet on letters  $\sigma_1, \sigma_2, \dots, \sigma_{n-2}, \sigma_{n-1}, \sigma_1^{-1}, \sigma_2^{-1}, \dots, \sigma_{n-2}^{-1}, \sigma_{n-1}^{-1}, \epsilon$  the corresponding composition of (classes of) homeomorphisms. To juxtaposition of words there evidently corresponds the composition of (classes of) homeomorphisms. The fact that  $\Theta$  is well defined (respects the relations) for some relations are classical facts, for others it is evident, probably, with the exception of the third relation in (2.3). The equalities of this relation follow from the fact that all three given classes (containing identity) ignore the first two points.

Let  $\epsilon_{k+1,n}$  denote the the isotopy class of of a homeomorphism that fixes the first  $k$  points and does not care about the remaining  $n - k$  points. On the level of braids (on a sphere) it corresponds to the partial braid with the trivial first  $k$  strands and the absent rest  $n - k$  strands. The element  $\epsilon_{k+1,n}$  can be expressed using the generator  $\epsilon$  or the generators  $\epsilon_i$  as follows

$$(4.4) \quad \epsilon_{k+1,n} = \epsilon\sigma_{n-1} \dots \sigma_{k+1}\epsilon\sigma_{n-1} \dots \sigma_{k+2}\epsilon \dots \epsilon\sigma_{n-1}\sigma_{n-2}\epsilon\sigma_{n-1}\epsilon,$$

$$\epsilon_{k+1,n} = \epsilon_{k+1}\epsilon_{k+2} \dots \epsilon_n,$$

From our construction of the inverse mapping class monoid it follows that every element of  $\mathcal{IM}_{0,n}$  represented by a homeomorphism  $h$  of  $S_{g,b,n}$  which maps  $k$  points,  $k \leq n$ , from  $Q_n$ :  $\{i_1, \dots, i_k\}$  to  $k$  points  $\{j_1, \dots, j_k\}$  from  $Q_n$  can be expressed in the form

$$(4.5) \quad (\sigma_{i_1} \dots \sigma_1) \dots (\sigma_{i_k} \dots \sigma_k) \epsilon_{k+1,n} x \epsilon_{k+1,n} (\sigma_k \dots \sigma_{j_k}) \dots (\sigma_1 \dots \sigma_{j_1}),$$

$$k \in \{0, \dots, n\}, \quad x \in \mathcal{M}_{0,k}, \quad 0 \leq i_1 < \dots < i_k \leq n-1 \text{ and } 0 \leq j_1 < \dots < j_k \leq n-1.$$

Geometrically this means that we first send the points  $\{i_1, \dots, i_k\}$  to the points  $\{1, \dots, k\}$ , then apply a homeomorphism from the mapping class group  $\mathcal{M}_{0,k}$ , and then send the points  $\{1, \dots, k\}$  to the points  $\{j_1, \dots, j_k\}$ .

Note that in the formula (4.5) we can remove one of the  $\epsilon_{k+1,n}$ , but we shall use the form (4.5) because of convenience: two symbols  $\epsilon_{k+1,n}$  serve as markers to distinguish the elements of  $\mathcal{M}_{0,k}$ .

The element  $x$  belongs to the mapping class group  $\mathcal{M}_{0,k}$ , so it can be expressed as a word in the letters  $\sigma_1, \sigma_2, \dots, \sigma_{k-1}, \sigma_1^{-1}, \sigma_2^{-1}, \dots, \sigma_{k-1}^{-1}$ . This proves that homomorphism  $\Theta$  is onto.

Let us prove that  $\Theta$  is a monomorphism. Suppose that for two words  $W_1, W_2 \in P_n$  we have

$$\Theta(W_1) = \Theta(W_2).$$

It means that the corresponding homeomorphisms both map the set of points  $\{i_1, \dots, i_k\}$  onto the set of points  $\{j_1, \dots, j_k\}$  and they are isotopic in the class of homeomorphisms mapping  $\{i_1, \dots, i_k\}$  onto  $\{j_1, \dots, j_k\}$ . Using relations (2.1) and (2.3) the same way as in [5] transform the words  $W_1, W_2$  into the form (4.5)

$$\sigma(i_1, \dots, i_k; k) \epsilon_{k+1, n} x \epsilon_{k+1, n} \sigma(k; j_1, \dots, j_k).$$

Then the corresponding fragments  $\sigma(i_1, \dots, i_k; k)$  and  $\sigma(k; j_1, \dots, j_k; k)$  for  $W_1$  and  $W_2$  coincide. The elements  $x_1$  of  $W_1$  and  $x_2$  of  $W_2$  presented in the form 4.5 are the words in  $\sigma_1, \dots, \sigma_k$  and  $\sigma_1^{-1}, \dots, \sigma_k^{-1}$ , and they correspond after  $\Theta$  to isotopic homeomorphisms

$$h_1, h_2 : S_{0,k} \rightarrow S_{0,k}.$$

Hence  $x_1$  can be transformed into  $x_2$  using relations for the mapping class group  $\mathcal{M}_{0,k}$ . So, the words  $W_1$  and  $W_2$  represent the same element in  $P_n$ .  $\square$

**Proposition 4.1.** *There is a presentation of the inverse mapping class monoid for punctured sphere  $\mathcal{IM}_{0,n}$  with the generators  $\sigma_1, \sigma, \sigma_1^{-1}, \sigma^{-1}, \epsilon$  subject to the relations (2.2), (2.7) and (4.3).*  $\square$

The generators  $\epsilon_i$  commute with  $\Delta$  in the following way [26]:

$$\epsilon_i \Delta = \Delta \epsilon_{n+1-i}.$$

Let  $\mathcal{E}$  be the monoid generated by one idempotent generator  $\epsilon$ .

**Proposition 4.2.** *The abelianization of  $\mathcal{IM}_{0,n}$ ,  $n \geq 2$ , is an abelian monoid  $\mathcal{AM}$  generated (as an abelian monoid) by elements  $\epsilon$  and  $\bar{\alpha}$  subject to the following relations*

$$\begin{cases} 2\epsilon = \epsilon, \\ \epsilon + \bar{\alpha} = \epsilon \\ m\bar{\alpha} = 0, \text{ where } m = 2(n-1), \text{ if } n \text{ is even and } m = (n-1), \text{ if } n \text{ is odd.} \end{cases}$$

So,  $\mathcal{AM}$  is isomorphic to the quotient-monoid of  $\mathcal{E} \oplus \mathbb{Z}/m\mathbb{Z}$  by the relation  $\epsilon + \bar{1} = \epsilon$ . The canonical map of abelianization

$$a : \mathcal{IM}_{0,n} \rightarrow \mathcal{AM}$$

is given by the formula:

$$\begin{cases} a(\epsilon_i) = \epsilon, \\ a(\sigma_i) = \bar{\alpha}. \end{cases}$$

*Proof.* According to the presentation of the monoid  $\mathcal{IM}_{0,n}$ , given in Theorem 4.1 the homomorphism  $a$  is well defined. Now let  $f : \mathcal{IM}_{0,n} \rightarrow B$  be an arbitrary homomorphism from  $\mathcal{IM}_{0,n}$  to an abelian monoid  $B$ . Then the braid relations give that  $f(\sigma_i) = f(\sigma_j)$  for any  $i, j$ . Denote this element  $f(\sigma_i)$  by  $b$  and the image of  $\epsilon$  by  $\epsilon$ . Then the sphere braid relation and the sphere mapping class relation give that  $b^m = 1$ . Two last relations in (2.3) give that

$$\epsilon = \epsilon^2 = \epsilon b^2 = \epsilon b.$$

This defines a unique map

$$f_a : \mathcal{AM} \rightarrow B,$$

such that  $f = f_a a$  and so  $\mathcal{AM}$  is an abelianization of  $\mathcal{IM}_{0,n}$ .  $\square$

**Corollary 4.1.** *The monoid  $\mathcal{AM}$  as a set consists of the following elements*

$$\mathcal{AM} = \{\bar{0}, \epsilon, \bar{\alpha}, 2\bar{\alpha}, \dots, (m-1)\bar{\alpha}\}.$$

$\square$

**Proposition 4.3.** *The center of  $\mathcal{IM}_{0,n}$  for  $n \geq 1$  is isomorphic to  $\mathcal{E}$ :*

$$Z(\mathcal{IM}_{0,n}) = \{1\} \amalg \mathcal{M}_{0,0} \cong \mathcal{E},$$

*i.e. the only non-unit element of  $Z(\mathcal{IM}_{0,n})$  is the isotopy class of the identity map of a sphere such that isotopy does not respect punctures.*

*Proof.* If  $n = 1$ , then the monoid  $\mathcal{IM}_{0,1}$  consists of two elements, unit 1 and the idempotent  $\epsilon$  (Example 2 below) and so it is a commutative monoid isomorphic to  $\mathcal{E}$ . Let  $n \geq 2$ . Denote by  $\Phi$  the only element of  $\mathcal{M}_{0,0} \hookrightarrow \mathcal{IM}_{0,n}$ :

$$\Phi \in \mathcal{M}_{0,0} \hookrightarrow \mathcal{IM}_{0,n}.$$

For any element  $x$  of  $\mathcal{IM}_{0,n}$  we have

$$\Phi x = x\Phi = \Phi.$$

So,  $\Phi$  lies in the center. Let  $c$  be a non-unit element in the center different from  $\Phi$ . It does not belong to the mapping class group  $\mathcal{M}_{0,n}$ , because this group is centerless for  $n \geq 3$  [10] and for  $n = 2$  (Example 3 below)  $c$  can not be the generator  $\sigma_1$  of  $\mathcal{M}_{0,2} = \Sigma_2$ , since we have

$$\sigma_1 \epsilon_1 \neq \epsilon_1 \sigma_1.$$

So  $c$  is a homeomorphism of  $S^2$  which maps the points  $I_k = \{i_1, \dots, i_k\}$  to the points  $J_k = \{j_1, \dots, j_k\}$ ,  $0 < k < n$ . Take a (class of) homeomorphism  $x$  that maps one point in the complement of  $J_k$  to a point in  $I_k$  without restrictions to other points. Then  $cx$  is equal to  $\Phi$ , while  $xc$  is not.  $\square$

## 5. THE WORD PROBLEM

R. Gillette and J. Van Buskirk in [10] obtained an analogue of the Markov normal form [19] for the sphere braid groups  $Br_n(S^2)$  and for the punctured sphere mapping class group  $\mathcal{M}_{0,n}$ . This form can be obtained algorithmically, so, it gives a solution of the word problem. These ideas can be applied to the cases of inverse braid and mapping class monoids.

Let us recall the main points of the Gillette-Van Buskirk variant of Markov normal form. Define the elements  $s_{i,j}$ ,  $1 \leq i < j \leq m$ , of the classical braid group  $Br_m$  by the formula:

$$s_{i,j} = \sigma_{j-1} \dots \sigma_{i+1} \sigma_i^2 \sigma_{i+1}^{-1} \dots \sigma_{j-1}^{-1}.$$

These elements satisfy the following Burau relations:

$$(5.1) \quad \begin{cases} s_{i,j} s_{k,l} = s_{k,l} s_{i,j} \text{ for } i < j < k < l \text{ and } i < k < l < j, \\ s_{i,j} s_{i,k} s_{j,k} = s_{i,k} s_{j,k} s_{i,j} \text{ for } i < j < k, \\ s_{i,k} s_{j,k} s_{i,j} = s_{j,k} s_{i,j} s_{i,k} \text{ for } i < j < k, \\ s_{i,k} s_{j,k} s_{j,l} s_{j,k}^{-1} = s_{j,k} s_{j,l} s_{j,k}^{-1} s_{i,k} \text{ for } i < j < k < l. \end{cases}$$

The elements  $s_{i,j}$  with the relations (5.1) give a presentation of the pure braid group  $P_m$  [19].

Let us define the elements  $\sigma_{k,l}$ ,  $1 \leq k \leq l \leq m$  by the formulas

$$\begin{aligned}\sigma_{k,k} &= e, \\ \sigma_{k,l} &= \sigma_k^{-1} \dots \sigma_{l-1}^{-1}.\end{aligned}$$

We denote by the same symbols the images of the defined elements in the braid group of a sphere. Let  $w(x_1, \dots, x_m)$  be a word with possible entries of  $x_i^\delta$ , where  $x_i$  are some letters and  $\delta$  may be  $\pm 1$ .

**Theorem 5.1.** [10] *Every element of the group  $Br_n(S^2)$  can be uniquely written in the form*

$$\sigma_{i_n, n} \dots \sigma_{i_j, j} \dots \sigma_{i_2, 2} w_1(s_{1,2}, \dots, s_{1, n-1}) \dots w_j(s_{j, j+1}, \dots, s_{j, n-1}) \dots w_{n-3}(s_{n-3, n-2}, s_{n-2, n-1}) \Delta^{2\delta},$$

where elements  $s_{j, j+1}, \dots, s_{j, n-1}$  generate a free group and  $\delta$  may be 0 or 1. There exists an algorithm of obtaining the normal form for any word on letters  $\sigma_1, \dots, \sigma_{n-1}$ , which gives a solution of the word problem for  $Br_n(S^2)$ .

**Theorem 5.2.** [10] *Every element of the group  $\mathcal{M}_{0,n}$  can be uniquely written in the form*

$$\sigma_{i_n, n} \dots \sigma_{i_j, j} \dots \sigma_{i_2, 2} w_1(s_{1,2}, \dots, s_{1, n-1}) \dots w_j(s_{j, j+1}, \dots, s_{j, n-1}) \dots w_{n-3}(s_{n-3, n-2}, s_{n-2, n-1}),$$

where elements  $s_{j, j+1}, \dots, s_{j, n-1}$  generate a free group. There exists an algorithm of obtaining the normal form for any word on letters  $\sigma_1, \dots, \sigma_{n-1}$ , which gives a solution of the word problem for  $\mathcal{M}_{0,n}$ .

**Theorem 5.3.** *Every element  $w$  in  $IB_n(S^2)$  can be uniquely written in the form*

$$(5.2) \quad \sigma_{i_1} \dots \sigma_1 \dots \sigma_{i_k} \dots \sigma_k \epsilon_{k+1, n} x \epsilon_{k+1, n} \sigma_k \dots \sigma_{j_k} \dots \sigma_1 \dots \sigma_{j_1},$$

$$k \in \{0, \dots, n\}, 0 \leq i_1 < \dots < i_k \leq n-1, 0 \leq j_1 < \dots < j_k \leq n-1, \text{ and } x \in Br_k(S^2)$$

is written in the Gillette-Van Buskirk normal form for  $Br_k(S^2)$ . If  $w$  is expressed in terms of  $\sigma_1, \dots, \sigma_{n-1}, \sigma_1^{-1}, \dots, \sigma_{n-1}^{-1}, \epsilon$ , there is an effective algorithm for putting it in this normal form.

*Proof.* Given an element  $w$  of  $IB_n(S^2)$ , expressed as a word in the generators, the elements  $\sigma_{i_1}, \dots, \sigma_1, \dots, \sigma_{i_k}, \dots, \sigma_k$  and  $\sigma_k, \dots, \sigma_{j_k}, \dots, \sigma_1, \dots, \sigma_{j_1}$  of (5.2) are determined uniquely and algorithmically [5] by a given element of  $IB_n(S^2)$ . Geometrically they are determined by the fact which strands are absent and which ones are present in the braid  $w$ . The same way the element  $x \in Br_k(S^2)$  is determined as a word in the generators of  $Br_k(S^2)$ . We apply Gillette-Van Buskirk algorithm to it to put it into Gillette-Van Buskirk normal form which is also unique.  $\square$

**Theorem 5.4.** *Every element  $w$  in  $\mathcal{IM}_{0,n}$  can be uniquely written in the form*

$$\sigma_{i_1} \dots \sigma_1 \dots \sigma_{i_k} \dots \sigma_k \epsilon_{k+1, n} x \epsilon_{k+1, n} \sigma_k \dots \sigma_{j_k} \dots \sigma_1 \dots \sigma_{j_1},$$

$$k \in \{0, \dots, n\}, x \in \mathcal{M}_{0,k}, 0 \leq i_1 < \dots < i_k \leq n-1, 0 \leq j_1 < \dots < j_k \leq n-1,$$

where  $x$  is written in the Gillette-Van Buskirk normal form for  $\mathcal{M}_{0,k}$ . If  $w$  is expressed in terms of  $\sigma_1, \dots, \sigma_{n-1}, \sigma_1^{-1}, \dots, \sigma_{n-1}^{-1}, \epsilon$ , there is an effective algorithm for putting it in this normal form.

*Proof.* It is essentially the same as the proof of Theorem 5.3. Given an element  $w$  of  $\mathcal{IM}_{0,n}$ , expressed in its generators, the elements  $\sigma_{i_1}, \dots, \sigma_1, \dots, \sigma_{i_k}, \dots, \sigma_k$  and  $\sigma_k, \dots, \sigma_{j_k}, \dots, \sigma_1, \dots, \sigma_{j_1}$  are determined uniquely and algorithmically [5] by a given element of  $\mathcal{IM}_{0,n}$ . Geometrically they are determined by the fact which points participate in the description of a map of the class  $w$  and which ones are omitted. The same way the element  $x \in \mathcal{M}_{0,k}$  is determined as a word in generators of  $\mathcal{M}_{0,k}$ . We apply Gillette-Van Buskirk algorithm for  $\mathcal{M}_{0,k}$  to it to put it into Gillette-Van Buskirk normal form which is also unique.  $\square$

## 6. EXAMPLES

1. The monoid  $\mathcal{IM}_{0,0}$  consists of one element of the identical map of a sphere

$$Id : S_{0,0} \rightarrow S_{0,0},$$

2. The monoid  $\mathcal{IM}_{0,1}$  consists of two elements, say, unit 1 and the idempotent  $\epsilon$ .

3. The mapping class group  $\mathcal{M}_{0,2}$  is isomorphic to the symmetric group  $\Sigma_2$ , so the inverse mapping class monoid  $\mathcal{IM}_{0,2}$  is isomorphic to the symmetric inverse monoid  $I_2$ .

4. The mapping class group  $\mathcal{M}_{0,3}$  is isomorphic to the symmetric group  $\Sigma_3$ , so the inverse mapping class monoid  $\mathcal{IM}_{0,3}$  is isomorphic to the symmetric inverse monoid  $I_3$ .

5. The mapping class group of a once-punctured torus  $\mathcal{M}_{1,1}$  is isomorphic to the general linear group  $GL_2\mathbb{Z}$  (unimodular group  $SL_2\mathbb{Z}$  if we want to restrict ourselves to the orientation preserving homeomorphisms of a torus) as well as the mapping class group of a torus (without punctures) is isomorphic to  $GL_2\mathbb{Z}$  (again it is isomorphic to  $SL_2\mathbb{Z}$  if we want to restrict ourselves to the orientation preserving case). Then the inverse mapping class monoid  $\mathcal{IM}_{1,1}$  as a set is a disjoint union of two copies of  $GL_2\mathbb{Z}$ :

$$\mathcal{IM}_{1,1} = GL_2\mathbb{Z} \amalg GL_2\mathbb{Z}.$$

If we denote by  $\mathcal{IM}_{1,1}^+$  the inverse mapping class monoid of orientation preserving homeomorphisms then we have

$$\mathcal{IM}_{1,1}^+ = SL_2\mathbb{Z} \amalg SL_2\mathbb{Z}.$$

Let the classes of maps of the first copy of  $GL_2\mathbb{Z}$  respect the puncture and of the second copy are free. Then the multiplication in each copy is standard and multiplying a punctured element by a free one we get a standard product in  $GL_2\mathbb{Z}$  and put it in the free copy.

This example gives a good illustration for Theorem 3.2 if we recall the classical result of Nielsen that the group of outer automorphisms of the free group  $F_2$  (fundamental group of punctured torus) is isomorphic to  $GL_2\mathbb{Z}$ , see, for instance [18, Corollary N4, p.169].

## 7. ACKNOWLEDGMENTS

The authors are grateful to the referee for many useful remarks.

## REFERENCES

- [1] *E. Artin*, Theorie der Zöpfe. Abh. Math. Semin. Univ. Hamburg, 1925, v. 4, 47–72.
- [2] *J. A. Berrick, F. R. Cohen, Y. L. Wong and J. Wu*, Configurations, braids, and homotopy groups. J. Amer. Math. Soc. 19 (2006), no. 2, 265–326.
- [3] *H. S. M. Coxeter, W. O. J. Moser*, Generators and relations for discrete groups. 3rd ed. Ergebnisse der Mathematik und ihrer Grenzgebiete. Band 14. Berlin-Heidelberg-New York: Springer-Verlag. IX, 161 p. (1972).
- [4] *W. Dicks and E. Formanek*, Algebraic mapping-class groups of orientable surfaces with boundaries. pp. 57-115, in: Infinite groups: geometric, combinatorial and dynamical aspects. Progress in Mathematics 248, Birkhäuser Verlag, Basel, 2005.

- [5] *D. Easdown; T. G. Lavers*, The inverse braid monoid. *Adv. Math.* 186 (2004), no. 2, 438–455.
- [6] *E. Fadell and J. Van Buskirk*, The braid groups of  $E^2$  and  $S^2$ . *Duke Math. J.* 29 1962, 243–257.
- [7] *B. Farb, C. J. Leininger, D. Margalit* The lower central series and pseudo-Anosov dilatations. arXiv:math/0603675.
- [8] *F. A. Garside*, The braid group and other groups, *Quart. J. Math. Oxford Ser.* 1969, 20, 235–254.
- [9] *N. D. Gilbert*, Presentations of the inverse braid monoid. *J. Knot Theory Ramifications* 15 (2006), no. 5, 571–588.
- [10] *R. Gillette; J. Van Buskirk*, The word problem and consequences for the braid groups and mapping class groups of the 2-sphere. *Trans. Amer. Math. Soc.* 131. 1968. 277–296.
- [11] *N. V. Ivanov*, Mapping class groups, *Handbook of geometric topology*, 523–633, North-Holland, Amsterdam, 2002.
- [12] *D. L. Johnson*, Towards a characterization of smooth braids. *Math. Proc. Cambridge Philos. Soc.* 92 (1982), no. 3, 425–427.
- [13] *F. Klein*, *Vorlesungen über höhere Geometrie*. 3. Aufl., bearbeitet und herausgegeben von *W. Blaschke*. VIII + 405 S. Berlin, J. Springer (Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen Bd. 22) (1926).
- [14] Kourouka notebook: unsolved problems in group theory. Seventh edition. 1980. Akad. Nauk SSSR Sibirsk. Otdel., Inst. Mat., Novosibirsk, 1980. 115 pp. (Russian)
- [15] *M. V. Lawson*, *Inverse semigroups: the theory of partial symmetries*, World Scientific, Singapore, 1998.
- [16] *W. Magnus*, Über Automorphismen von Fundamentalgruppen berandeter Flächen. (German) *Math. Ann.* 109, 617-646 (1934).
- [17] *W. Magnus*, Braids and Riemann surfaces. *Comm. Pure Appl. Math.* 25 (1972) 151-161.
- [18] *W. Magnus; A. Karrass; D. Solitar*, *Combinatorial group theory. Presentations of groups in terms of generators and relations*. 2nd rev. ed. Dover Books on Advanced Mathematics. New York: Dover Publications, Inc. XII, 444 p. (1976).
- [19] *A. A. Markoff*, Foundations of the Algebraic Theory of Tresses, *Trudy Mat. Inst. Steklova*, No 16, 1945 (Russian, English summary).
- [20] *J. von Neumann*, On regular rings. *Proc. Natl. Acad. Sci. USA* 22, 707-713 (1936).
- [21] *M. Petrich*, *Inverse semigroups*. Pure and Applied Mathematics (New York). A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1984. x+674 pp.
- [22] *L. M. Popova*, Defining relations of a semigroup of partial endomorphisms of a finite linearly ordered set. (Russian) *Leningrad. Gos. Ped. Inst. Učen. Zap.* 238 1962 78–88.
- [23] *G. P. Scott*, Braid groups and the group of homeomorphisms of a surface. *Proc. Cambridge Philos. Soc.* 68, 1970, 605–617.
- [24] *V. V. Vershinin*, On presentations of generalizations of braids with few generators, *Fundam. Prikl. Mat.* Vol. 11, No 4, 2005. 23-32.
- [25] *V. V. Vershinin*, Braid groups, their Properties and Generalizations. *Handbook of Algebra*, vol. 4, Elsevier, Amsterdam a.o. 2006, p. 427-465.
- [26] *V. V. Vershinin*, On the inverse braid monoid. *Topology Appl.* 156 (2009) 1153–1166.
- [27] *V. V. Wagner*, Generalized groups. (Russian) *Doklady Akad. Nauk SSSR (N.S.)* 84, (1952). 1119–1122.
- [28] *O. Zariski*, On the Poincaré group of rational plane curves, *Am. J. Math.* 1936, 58, 607-619.

DÉPARTEMENT DES SCIENCES MATHÉMATIQUES, UNIVERSITÉ MONTPELLIER II, PLACE EUGÈNE BATAIL-  
LON, 34095 MONTPELLIER CEDEX 5, FRANCE

*E-mail address:* `Rym.Karoui@math.univ-montp2.fr`

DÉPARTEMENT DES SCIENCES MATHÉMATIQUES, UNIVERSITÉ MONTPELLIER II, PLACE EUGÈNE BATAIL-  
LON, 34095 MONTPELLIER CEDEX 5, FRANCE

*E-mail address:* `vershini@math.univ-montp2.fr`

SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK 630090, RUSSIA

*E-mail address:* `versh@math.nsc.ru`