



CO₂-SO₂ clathrate hydrate formation on early Mars

Eric Chassefière, Emmanuel Dartois, Jean-Michel Herri, Feng Tian, Frédéric Schmidt, Olivier Mousis, Azzedine Lakhli

► To cite this version:

Eric Chassefière, Emmanuel Dartois, Jean-Michel Herri, Feng Tian, Frédéric Schmidt, et al.. CO₂-SO₂ clathrate hydrate formation on early Mars. Icarus, 2013, 223 (2), pp.878-891. 10.1016/j.icarus.2013.01.001 . hal-00804822

HAL Id: hal-00804822

<https://hal.science/hal-00804822>

Submitted on 26 Mar 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CO₂-SO₂ clathrate hydrate formation on early Mars

Eric Chassefière^{a,b}, Emmanuel Dartois^c, Jean-Michel Herri^d, Feng Tian^e, Frédéric Schmidt^{a,b},
Olivier Mousis^{f,g}, Azzedine Lakhelif^f

^a Univ Paris-Sud, Laboratoire IDES, UMR8148, Université Paris-Sud, Bât. 504, Orsay, F-91405, France;

^b CNRS, Orsay, F-91405, France.

^c IAS, Université Paris-Sud, CNRS, France

^dCentre SPIN, ENS des Mines de Saint-Etienne, France

^eCenter for Earth System Sciences, Tsinghua University, Beijing, China

^fInstitut UTINAM, UMR 6213, Université de Franche-Comté & OSU THETA de Franche-Comté, France

^gUniversité de Toulouse; UPS-OMP; CNRS-INSU; IRAP; 14 Avenue Edouard Belin, 31400 Toulouse, France

Corresponding author

Eric Chassefière

IDES, Univ. Paris-Sud, CNRS

Tel : 33 1 69 15 67 48

Fax : 33 1 69 15 49 11

E-mail : eric.chassefiere@u-psud.fr

Second revised version submitted to Icarus

Abstract

It is generally agreed that a dense CO₂-dominant atmosphere was necessary in order to keep early Mars warm and wet. However, current models have not been able to produce surface temperature higher than the freezing point of water. Most sulfate minerals discovered on Mars are dated no earlier than the Hesperian, despite likely much stronger volcanic activities and more substantial release of sulfur-bearing gases into Martian atmosphere during the Noachian. Here we show, using a 1-D radiative-convective-photochemical model, that clathrate formation during the Noachian would have buffered the atmospheric CO₂ pressure of early Mars at ~2 bar and maintained a global average surface temperature ~230 K. Because clathrates trap SO₂ more favorably than CO₂, all volcanically outgassed sulfur would have been trapped in Noachian Mars cryosphere, preventing a significant formation of sulfate minerals during the Noachian and inhibiting carbonates from forming at the surface in acidic water resulting from the local melting of the SO₂-rich cryosphere. The massive formation of sulfate minerals at the surface of Mars during the Hesperian could be the consequence of a drop of the CO₂ pressure below a 2-bar threshold value at the late Noachian-Hesperian transition, which would have released sulfur gases into the atmosphere from both the Noachian sulfur-rich cryosphere and still active Tharsis volcanism. A lower value of the pressure threshold, down to ~0.5 bar, could have been sufficient to maintain middle and high latitude regions below the clathrate formation temperature during the Noachian and to make the trapping of SO₂ in clathrates efficient. Our hypothesis could allow to explain the formation of chaotic terrains and outflow channels, and the occurrence of episodic warm episodes facilitated by the release of SO₂ to the atmosphere. These episodes could explain the formation of valley networks and the degradation of impact craters, but remain to be confirmed by further modeling.

Key words : Early Mars, CO₂, SO₂, clathrates, sulfates, cryosphere, carbonates.

1. Introduction

Sulfur dioxide (SO_2) has been proposed as a possible greenhouse gas which worked together with CO_2 to raise the surface temperature of early Mars above the freezing point of water (Halevy et al., 2007; Johnson et al., 2008; Postawko and Kuhn, 1986), allowing liquid water to flow at the surface of the planet and carving the presently observed valley networks and outflow channels. However, the cooling from sulfur gases as a result of sulfate aerosol formation is an observational fact for the current atmosphere of Earth. On early Mars, the cooling effect of sulfate aerosols would also have counteracted efficiently the warming effect due to SO_2 greenhouse effect and sulfur outgassing. After a short period of warming, the presence of sulfate aerosols would have resulted in a colder surface, with a net cooling, instead of warming, of the planet (Tian et al., 2010). No mechanism to keep ancient Mars warm and wet through greenhouse effect has been generally accepted so far, suggesting a cold and wet early Mars. Much of the aqueous activity on Mars could have occurred in subsurface hydrothermal systems powered by magmatic or impact activity (Squyres and Kasting, 1994; Griffith and Shock, 1995; Segura et al., 2002; 2008), rather than at the surface of the planet.

Sulfur, under both oxidized (SO_2) and reduced (H_2S) forms, could have been released in large amount by volcanism during the Noachian. From several hundred millibar to around one bar of sulfur could have been outgassed along Martian history, most of which during the Noachian and the Hesperian (see e.g. Craddock and Greeley, 2009; Gaillard and Scaillet, 2009). The reconstructed evolution of the cumulated amount of released SO_2 since the early Noachian, shortly after accretion, is shown in Figure 1. It has been derived from both the CO_2 volcanic release rates calculated from a thermo-chemical evolution model of Mars (Grott et al., 2011) and a photogeological analysis of the Martian surface (Craddock and Greeley, 2009, referred to as CG09 in this paper). The ratio between SO_2 and CO_2 molar fractions in the

Fig.1

79 volcanic gas $f_{\text{SO}_2}/f_{\text{CO}_2}$ is typically in the range from 0.5 to ~ 1 (see Fig. 5 in Gaillard and
80 Scaillet, 2009). The evolutions of the CO_2 partial pressure (denoted by p_{CO_2} in the following)
81 for two values (0.01 and 1) of the surface fraction covered by hot upwellings (f_p) are plotted
82 on Figure 4b in Grott et al. (2011). By applying the ratio $f_{\text{SO}_2}/f_{\text{CO}_2}$ to Grott et al. curves, and
83 taking into account the difference of molar mass between CO_2 (44 g/mol) and SO_2 (64 g/mol),
84 we have plotted in Fig. 1 the corresponding evolutions of the cumulated amount of outgassed
85 SO_2 , expressed in pressure units, and the corresponding global equivalent layer (GEL)
86 thickness of the sulfate mineral assuming that all the released sulfur is involved in anhydrite
87 formation. The other plotted curve is obtained from a photogeological analysis of the Martian
88 surface yielding a cumulated lava extrusion volume of $0.7 \times 10^8 \text{ km}^3$ (CG09), by assuming
89 that basalts contain 7000 ppm sulfur by mass (an upper range according to Gaillard and
90 Scaillet, 2009), that is 10 times more than assumed in CG09. This curve would be ~ 3 times
91 lower assuming a lower range of 2500 ppm of S and 4 times higher assuming a lava extrusion
92 volume of $3 \times 10^8 \text{ km}^3$ (Gaillard and Scaillet, 2009). Both models and observations therefore
93 suggest a cumulated pressure of outgassed SO_2 of typically ~ 1 bar, corresponding to a ~ 40 m
94 thick sulfate GEL, within a factor of 3 both ways.

95
96 Interestingly, all sulfate minerals detected at the surface of Mars by OMEGA/Mars Express
97 have been emplaced during the Hesperian, a relatively late epoch in Martian history (Bibring
98 et al., 2006). Later observations of CRISM on MRO have revealed additional sulfate deposits,
99 a few of them under the form of interbedded phyllosilicate and sulfate layers in sediments of
100 Noachian craters (Terra Sirenum) in the Southern hemisphere (Murchie et al., 2009). These
101 intracrater deposits are relatively widespread, far from those mapped by OMEGA and at
102 higher elevation, excluding that they have been transported from equatorial regions (Wray et
103 al., 2011). Although a younger age for these sediments cannot be excluded, they are suspected
104 to have formed by evaporitic processes in an acidic context at the Noachian. Such formations

associating phyllosilicates and sulfates are also found in terrestrial acidic saline lake deposits (Baldrige et al., 2009). Most of sulfate deposits are observed in Hesperian terrains. They consist of several kinds of formations: (i) Hesperian layered sulfates under the form of extended deposits in Terra Meridiani, (ii) Interior Layered Deposits (ILD) throughout the Valles Marineris trough system consisting of massive mounds of layered material up to several kilometers thick, (iii) gypsum deposits in north polar dune field (Gaillard et al., 2012). Sulfur is also part of soil and dust at the global scale at an average level of ~6.8 % (King and McLennan, 2010). It has been suggested that some sulfates could have been formed by evaporitic processes in the Valles Marineris region prior to Tharsis formation, then redistributed by fluvial transport to Meridiani Planum, where they are now observed, during the elevation of the Tharsis uplift (Fan et al., 2008). If so, sulfates detected in Hesperian terrains could have been formed in the Noachian. The question of the origin of sulfates is debated, and at least some of the observed deposits could have formed in the late Noachian. In the present paper, we make the hypothesis that most of the sulfates observed in equatorial regions (layered deposits, ILD) have been formed at the Hesperian (not excluding a start at the late Noachian), and propose an explanation for the scarcity of sulfates in Noachian terrains, and their apparently massive deposition at the Hesperian.

The evolution of CO₂ on Mars is poorly constrained. Despite the discovery of carbonates in SNC meteorites and the recent orbital and in situ observations of carbonate rocks at some locations on the surface of Mars (see e.g. Niles et al., 2012), carbonates are not proved to be extensively present on Mars. The lack of global carbonate outcrops on Mars, which seems to contradict a massive presence of CO₂ in its early atmosphere, could be explained by either the action of sulfuric or sulfurous acids in large, standing bodies of water suppressing the formation of carbonates (Fairén et al., 2004; Halevy et al., 2007), or a rapid escape of early Martian CO₂ atmosphere (Tian et al. 2009, referred to as T09 in this paper), or both.

131

132 According to an early Mars upper atmosphere model (T09), the timescale for 1 bar of CO₂ to
133 be removed through thermal escape would have been ~1 Myr at 4.5 billion years ago (Ga) and
134 ~10 Myr at 4.1 Ga. Note that at first stages, during typically the first hundred million years,
135 carbon escape may have been slowed down by water hydrodynamic escape, which is not
136 taken into account in T09. If Mars was endowed with amounts of CO₂ similar to those found
137 on Venus and the Earth (~100 bar), and according to the model of T09, most of its initial CO₂
138 inventory would have been lost within the first 100 Myr after its formation. A mechanism that
139 can possibly help Mars to keep its CO₂ inventory is carbonate formation through weathering
140 of basalt (Pollack et al. 1987). The same authors proposed that a dense CO₂ atmosphere (1 to
141 5 bar) could have been maintained during the Noachian by constant recycling of CO₂ back
142 into the atmosphere through rapid burial and thermal decomposition of carbonate minerals.
143 They calculated a typical cycling time of CO₂ at 273 K of ~10 Myr, decreasing to ~1 Myr at
144 larger temperatures of 300 K for a CO₂ pressure of 1 bar, a temperature easily achievable in
145 the subsurface even if the surface was cold. If the formation of carbonate through weathering
146 has occurred at a rate comparable to that of carbon thermal escape, substantial amounts of
147 carbon could have been trapped in the format of carbonates and cycled through the
148 subsurface-hydrosphere-atmosphere system. If so, a significant atmosphere of CO₂ could have
149 survived until 4.1 Ga. The combination of a rapid formation of carbonate and a rapid thermal
150 escape of carbon on Noachian Mars could have decreased the atmospheric pressure of
151 Noachian Mars even more rapidly, resulting in shorter warm periods and longer, more
152 frequent cold periods on Noachian Mars. It is important to note that the timescales of
153 carbonate formation and their further destabilization are still poorly understood.

154

155 The present inventory of water on Mars is also poorly constrained. The total water content of
156 the two perennial polar caps corresponds to a GEL of 16 m depth (Smith et al., 2001), and the

ice deposits sequestered in the Dorsa Argentea Formation (DAF), near the south polar cap, could have represented a ~15 m thick GEL in the past (Head and Pratt, 2001). Nevertheless, only a fraction of the initial water could remain today in DAF reservoir, corresponding to a ~5-7.5 m thick GEL. Other reservoirs, expected to have been active during late Amazonian, could be present in tropical and mid-latitude regions (Head and Marchant, 2009). But they probably represent only a minor contribution to the global reservoir. The total inventory of the known reservoir, including near-surface stores that are distributed across middle to high latitudes, has been estimated to correspond to a 35 m thick GEL (Christensen, 2006). The mega-regolith capacity is large, with up to a ~500 m thick GEL potentially trapped in the cryosphere, and hypothetically several additional hundreds of meters GEL (up to ~500 m) of ground water surviving at depth below the cryosphere (Clifford et al., 2010). It has been suggested that most of ground ice has been lost by sublimation at low latitudes, and that only small amounts of groundwater would survive today (Grimm and Painter, 2009), with therefore less water in the megaregolith. A ~500 m thick GEL is generally assumed to be required to explain the formation of outflow channels (Carr, 1987), and most of this water could be trapped today as water ice, and possibly deep liquid water, in the subsurface, as well as in subsurface hydrated minerals. Based on an analysis of the present Mars' atmospheric D/H ratio, it has been suggested that a water GEL of up to ~300-400 m depth could have been stored in crustal serpentine since the late Noachian due to hydrothermalism triggered by magmatic activity (Chassefière and Leblanc, 2011b).

In the present paper, we suggest that the formation of CO₂-SO₂ clathrate hydrates (indifferently written “clathrate hydrates” or “clathrates” in this article) in a potentially CO₂-dominant (a few bar) early atmosphere of Mars should have significantly decreased the atmospheric SO₂ and CO₂ contents and should have caused subsequent sulfur enrichment in early Mars cryosphere. As shown hereafter, for a CO₂ pressure above 2 bar, all volcanically

released SO₂ and the fraction of atmospheric CO₂ in excess of 2 bar should have been converted to CO₂-SO₂ clathrates. In this way, the formation of sulfate particles in the atmosphere could have been inhibited and there would have been neither long term net cooling nor net warming from volcanic SO₂ eruptions. If atmospheric CO₂ pressure is between 1 and 2 bar, we show that a fraction of volcanically released SO₂ would have been converted to sulfate particles while the rest would have been trapped in clathrates. This mechanism would have resulted in a stabilization of the surface temperature at the clathrate equilibrium temperature after large volcanic events. At some time in the past, when the CO₂ pressure dropped below 1 bar, the formation of clathrates would have been totally inhibited and the sulfur injected by volcanism into the atmosphere would have been fully converted into atmospheric sulfate aerosols. Thus formation of sulfate minerals at the surface no earlier than at the Hesperian could be explained by a decrease in atmospheric CO₂ pressure below the 2 bar level occurring close to the late Noachian/ Hesperian transition.

In the following section, we describe the potential impact of CO₂-SO₂ clathrate formation on the sulfur content in the atmosphere and the surface temperature. Then, we present an extrapolation to low temperature of the thermodynamic properties of CO₂-SO₂ clathrates. New calculations of the surface temperature of Mars in presence of sulfate particles for different values of the CO₂ pressure are made, and consequences on the climate of early Mars examined. After an assessment of the consequences of the presence of SO₂ on the trapping of noble gases in clathrates, we discuss the possible implications of the present results and measurements to be done by future missions to improve our understanding of the impact of sulfur on early Mars' climate.

2. Formation of CO₂-SO₂ clathrate hydrate at high CO₂ pressure and SO₂ mixing ratio

209

210 The cooling effect of sulfate aerosols on the early Mars climate has been recently pointed out
 211 through a detailed 1-D photochemical/microphysical model of ancient Mars' atmosphere
 212 (Tian et al. 2010, referred to as T10 in this paper). Under certain CO₂ pressure, the surface
 213 temperature increases only slightly with enhanced level of atmospheric SO₂ at first because of
 214 the greenhouse contribution from SO₂. At atmospheric SO₂ concentration which could lead to
 215 meaningful surface warming, photochemistry processes induce the formation of sulfate
 216 aerosols which can reflect sunlight and cause surface cooling. The surface temperature for
 217 various SO₂ mixing ratios (T10) together with the equilibrium temperature of CO₂ clathrate
 218 hydrate (Mousis et al., 2012), and CO₂ ice, are plotted in Figure 2 as a function of the CO₂ Fig. 2
 219 pressure. At 3 bar atmospheric pressure and for a SO₂ mixing ratio of 10 ppmv, the amplitude
 220 of the cooling by sulfate aerosols is about 60 K (T10), resulting in a surface temperature low
 221 enough to allow the formation of CO₂ clathrate hydrates (and even CO₂ ice). For a lower SO₂
 222 mixing ratio of 1 ppmv the cooling is smaller, of the order of 20 K, but sufficient to fall in the
 223 domain of stability of CO₂ clathrates (but not of CO₂ ice).

224

225 First, in a regime without sulfur, CO₂ clathrates are stable for p_{CO2}>2 bar due to the cooling
 226 effect of the Rayleigh scattering (T10). As a consequence any CO₂ pressure in excess of 2 bar
 227 results in the formation of CO₂ clathrate hydrates until the atmospheric CO₂ pressure is
 228 reduced to ~2 bar, with a stabilization of p/T values close to 2 bar/230 K. Thus, the
 229 atmospheric CO₂ pressure could have been maintained at ~2 bar during most of the Noachian,
 230 if atmospheric escape is neglected. Because carbon escaped rapidly during the early
 231 Noachian, this regime is possible only if a few bar of carbon sequestered first in crustal
 232 carbonates was recycled to the atmosphere by volcanism during the Noachian, since the
 233 amount of carbon released from the mantle to the atmosphere during the late Noachian and at
 234 later epochs is rather modest, a few hundreds millibar according to existing models (Grott et

al., 2011). The warming effect of CO₂ clouds by reflection of the IR radiation emitted by the surface (Forget and Pierrehumbert, 1997) is not taken into account in this simulation. At p_{CO2}=2 bar, the presence of CO₂ clouds may result in a surface temperature increase by ~5 K (Colaprete and Toon, 2003), somewhat increasing the temperature and pressure thresholds for the formation of CO₂ clathrates. Also, 1-D modeling provides only global average values and the local conditions may differ significantly from the average case. Formation of clathrates is expected to be more efficient in cold regions (polar caps) than at middle or lower latitudes. The possibility that a part of the CO₂ inventory may be stored under the form of clathrates has no significant impact on the discussion made in the previous section relative to the role of thermal escape. Indeed, any later decrease of the atmospheric CO₂ content following thermal escape would have resulted in the destabilization of CO₂ clathrates and a transfer to the atmosphere of an amount of CO₂ equal to that removed to space. In this regime, the CO₂ is buffering pressure and temperature around 2 bar/230 K until the total volatile CO₂ reservoir (ice, clathrate, and the atmosphere combined) decreased below 2 bar.

Second, the sporadic addition of SO₂ in the atmosphere by volcanic events into the previous regime would have had the following effects: (i) initial increase of the surface temperature by direct greenhouse effect, (2) rapid decrease of the surface temperature by formation of sulfate aerosols, in some cases below the equilibrium temperature of CO₂ clathrates (cf Fig. 2). If the atmospheric SO₂, which is the source of sulfate aerosols, was trapped together with CO₂ in clathrates hydrates, the SO₂ atmospheric content would have been decreased, which would have slowed down or even inhibited the formation of sulfate aerosols and the subsequent cooling. Through this mechanism, the atmospheric mixing ratio of SO₂, following volcanic events, will stabilize close to the mixing ratio maintaining a surface temperature equal the CO₂-SO₂ clathrate equilibrium temperature. As an example, for p_{CO2} = 3 bar, this equilibrium mixing ratio is of the order of 0.1 ppm (Fig. 2). Any increase of the atmospheric SO₂ mixing

ratio above this equilibrium value would have resulted in aerosol formation, cooling of the surface, formation of clathrates, and the removal of the remaining fraction of SO₂ gas from the atmosphere to clathrates. The higher the SO₂/CO₂ mixing ratio in the clathrates with respect to the gas, the more efficient this mechanism. If SO₂ is enriched with respect to CO₂ in the clathrates, formation of CO₂-SO₂ clathrates results in a decrease of the atmospheric SO₂ mixing ratio. If SO₂ is depleted in the solid phase, clathration results in an enrichment of SO₂ in the atmosphere. As shown in the next section, SO₂ is enriched by a factor of ~100-500 in the clathrate phase at ~220 K. The condensation of only 1% (or less) of the atmospheric CO₂ in clathrates, with therefore no significant change in p,T conditions, would have resulted in the removal of most atmospheric SO₂ to the cryosphere, implying that the proposed mechanism is highly efficient. Thus, once the regime described in the last paragraph is reached at p_{CO2} = 2 bar / 230 K, any addition of atmospheric sulfur will be incorporated into CO₂-SO₂ clathrates.

The goal of the next section is to study the composition of CO₂-SO₂ clathrates as a function of gas composition, in particular the SO₂/CO₂ mixing ratio in the clathrate as a function of the gas mixing ratio to estimate the efficiency of CO₂-SO₂ clathrate formation in transferring SO₂ from the atmosphere to the cryosphere.

3. Thermodynamic modeling of CO₂-SO₂ clathrates in Martian atmospheric conditions

3.1 Thermodynamical approach

The van der Waals and Platteeuw (1959) model describes the equilibrium of hydrate phases by means of a convergence between a statistical thermodynamics approach implementing

Kihara parameters and a classical approach with a reference state parameters. In a recent publication (Herri and Chassefière, 2012), the Kihara parameters have been optimized to fit equilibrium data at low temperature concerning Martian atmospheric components: carbon dioxide (95.3%), nitrogen (2.7%), Argon (2%) and methane (<50ppb with an average at 15 ppb) (Owen et al., 1977; Mumma et al., 2009), using Handa and Tse (1986) reference state parameters. We observed that the stability of clathrate hydrate is firstly dependent on the partial pressure of carbon dioxide. However, the possible presence of SO₂ in the ancient Mars' atmosphere, a potential promoter for clathrate hydrate formation, could move the stability line for clathrate hydrate formation.

To test the influence of SO₂ we must implement in our models its Kihara parameters, retrieved from experimental data. Up to date, only few equilibrium data have been measured for sulfur dioxide. Above 0°C there are 36 equilibrium data from Van Berkum and Diepen (1979) in the range of temperature [285.2-293.84 K] and the range of pressure [0.253-392.330 MPa], and 2 equilibrium data at (280.2 K, 0.101 MPa) and (273.2 K, 0.0396 MPa) from Von Stackelberg (1949). Below 0°C Tamman and Krige (1925) measured the following experimental points: (261.15 K, 0.01695 MPa), (265.15 K, 0.01997 MPa), (269.15 K, 0.02597 MPa), (270.15 K, 0.02778 MPa), (271.15 K, 0.03051 MPa). Above 0°C, the data can not be easily used to constrain the Kihara parameters because it implies to describe the liquid phase in the presence of salts resulting from SO₂ dissolution (H₂SO₄, HSO₄⁻, HSO₄²⁻) and thus to model the speciation of the reaction as well as the activity coefficient of water. Below 0°C the situation is easier, dealing with ice, simplifying the equations. From a practical point of view, the restricted range of available temperature and pressure measurements hampers the possibility of retrieving only one set of Kihara parameters, but only interdependent sets of values. Our aim is to test within these sets of constrained values the consequence on the composition of (CO₂-SO₂) clathrate hydrate under Martian conditions.

The detailed description of the statistical-thermodynamic model and the strategy used for the retrieval of the SO₂ Kihara parameters required for the modeling are deferred to Appendix A. We, however, stress the importance of this aspect, defining the central parameters on which are based the scientific discussion outcomes in the following sections.

3.2 Extrapolation of experimental data to Martian temperatures

We explored the minimization in the full 3D parameter space (see appendix A) to deduce the minimum SO₂ enrichment during the formation of a mixed SO₂-CO₂ clathrate hydrate in contact with a CO₂ dominated atmosphere. Whatever the minimized set of Kihara parameters for SO₂, and exploring concentration ratios from 0.1 ppm to 1000 ppm of SO₂ in CO₂, the resulting mixed clathrate hydrate formed is always highly enriched in SO₂, by a factor above 60 with respect to gas phase, for all temperatures in the 170-240 K range (Fig.3).

Fig. 3

Falenty et al. (2011) performed experimental kinetics studies of CO₂ clathrate hydrate, monitoring the formation through the neutron diffraction pattern of hexagonal ice in the 185-195 K range. The data were extrapolated at low temperature through a model to compare with the actual polar frosts of Mars, at about 150 K, in order to show that the formation timescale for micron sized clathrates was longer than the typical seasonal variations regularly inducing the declathration. The nucleation rate depends not only on the temperature but on the ice grain sizes when nucleating pure clathrate hydrate from ice aerosol, as the nucleation proceeds in at least two steps: first surface nucleation followed by growth of the hydrate crystal. At 195 K, for a distribution of ice particles of about 1.5 micron of mean equivalent sphere radius, the nucleation proceeds in about one day in the laboratory. Extrapolating at the higher temperatures considered in Fig. 3 for the ancient Martian surface temperature, the nucleation

rate for hydrate formation of similar size ice grains would be higher by a factor of 3 at 200 K to 2 orders of magnitude at 230 K. Precise nucleation timescale with an SO₂-CO₂ clathrate is difficult to extrapolate only based on these experiments, and will depend on the particular mechanism by which the aerosols or surface clathrate kinetic formation proceed (heterogeneous nucleation, diffusion controlled,...), but if we take the high rate of hydrate nuclei growth, it seems reasonable to grow grains as large as 100 microns in hours to days.

4. Modeling of the effects of SO₂ and sulfate aerosols on surface temperature and consequences of CO₂-SO₂ clathrate formation on early Mars climate

For modeling the effects of SO₂ and sulfate aerosols on surface temperature we used the 1-D radiative-convective climate model and the 1-D photochemical model in T10, which allows us to investigate the combined climate effect of CO₂, SO₂, and sulfate aerosols. Most of the model details are described in T10 and will not be repeated here. SO₂ in the early Mars atmosphere could have originated from volcanic outgassing, which is simulated by assuming different surface upward fluxes of SO₂ and H₂S. The ratio between the two species is kept at unity, and previous sensitivity tests (T10) showed that modifying this ratio does not change the results of the photochemical and climate calculations significantly. In this work the relative solar luminosity is set to $S/S_0=0.75$. The atmospheric pressure is set to the following levels : 1.5, 1, and 0.5 bar, in addition to the level of 3 bar already studied in T10, and Figure 4 shows the relationship between surface temperature and atmospheric SO₂ concentration under different CO₂ pressures. As discussed in T10 the surface temperature increases with enhanced level of atmospheric SO₂ at first because of the greenhouse contribution from SO₂. Nevertheless, as atmospheric SO₂ concentration keeps increasing, photochemistry leads to the formation of sulfate aerosols which can reflect sunlight and cause surface cooling. The

Fig. 4

decrease of surface temperature with increasing atmospheric SO_2 is more dramatic under higher CO_2 pressure than under the lower CO_2 pressure because the sulfate aerosols fall slower in a denser atmosphere. It can be seen from Figure 4 that the maximum surface temperatures from combined greenhouse effect of CO_2 and SO_2 in early Martian atmosphere are 235, 235, 230, and 220 K under 3, 1.5, 1.0, and 0.5 bar CO_2 pressure respectively. Thus volcanic outgassing of SO_2 is not a solution to an early warm and wet Mars, consistent with the conclusion in T10.

For $p_{\text{CO}_2} = 3$ bar, the curve is everywhere below the CO_2 clathrate equilibrium temperature. Black triangles in Figure 4 correspond to the SO_2 mixing ratio for which the surface temperature is equal to the CO_2 clathrate equilibrium temperature. The same values are plotted as a function of p_{CO_2} in Figure 5. In the four cases ($p_{\text{CO}_2} = 0.5, 1, 1.5$ and 3 bar), we have indicated the lava volumes required to give rise to the corresponding SO_2 mixing ratio (Wilson and Head, 2002; Hanna and Phillips, 2006). For a CO_2 pressure larger than 2 bar that may be reached sporadically due to seasonal/orbital effect or also to the CO_2 outgassing from the crust (volcanism and disruption of carbonates), CO_2 clathrates are stable whatever the SO_2 mixing ratio in the atmosphere, resulting in the trapping of the fraction of atmospheric CO_2 in excess of 2 bar in the cryosphere under clathrate form. Any atmospheric sulfur released by volcanism is trapped in the cryosphere under the form of CO_2 - SO_2 clathrate. Occasionally an SO_2 mixing ratio of ~ 1 ppm may be reached since at this level, sulfur has no significant effect on the mean temperature. For higher values, the surface cooling down will result in CO_2 - SO_2 clathration. Note that an ample reservoir of H_2O is available in polar caps (a several 10-m thick global equivalent layer in present conditions, probably much more during the Noachian) to feed the formation of clathrates. As soon as an atmospheric H_2O molecule is consumed in clathrate formation, it is replaced by a molecule sublimated from the polar caps, due to the short dynamical time scale of the atmosphere (a few weeks). In terms of reservoirs, the required quantities of H_2O (and CO_2) for all S released by volcanism during the Noachian

Fig. 5

to be stored in clathrates is discussed later in this section. H_2S either is converted into SO_2 within the atmosphere in a typical time of a couple of weeks (Wong and Atreya, 2003), or is trapped directly in clathrates.

In the 0.5-2 bar p_{CO_2} range, the SO_2 level in the atmosphere following a volcanic eruption can reach typically 10 ppm or so, the excess being trapped in the cryosphere. According to the estimate of the characteristic growth time of a clathrate particle in the atmosphere made in Section 3 (a few hours to days for a 100 μm size particle), trapping occurs rapidly, simultaneously with the radiative cooling of the atmosphere due to sulfate particle formation, which occurs within a few months after SO_2 outgassing. After large volcanic events, the surface temperature is therefore stabilized at the CO_2 - SO_2 clathrate equilibrium temperature (~ 220 K, see Fig. 2) within a few months. Some of the SO_2 trapped in the cryosphere at previous times when $p_{\text{CO}_2} > 2$ bar (if it ever occurred) may similarly be released to the atmosphere. Also, the release of a few thousand cubic kilometers of lava may have happened frequently during Tharsis formation (Johnson et al., 2008, and references therein). During this period, the atmosphere could have been rich in SO_2 and sulfate particles (often at equilibrium condition described in Fig.5) and the SO_2 -rich cryosphere could have lost part of its SO_2 .

For a CO_2 pressure below 0.5 bar, there is no more formation of CO_2 - SO_2 clathrates for SO_2 mixing ratio < 200 ppm and all the released sulfur is converted to sulfates. A 200 ppm level of SO_2 in the atmosphere corresponds to the most important eruption supposed to have taken place on Mars, with a lava volume of 50 000 km^3 (Wilson and Head, 2002).

The potential consequences of CO_2 - SO_2 clathrate formation on early Mars climate are multiple. Let assume that the CO_2 pressure has been buffered at 2 bar during most of the Noachian. Whereas, during the time when $p_{\text{CO}_2} > 2$ bar, there is no SO_2 in the atmosphere and

a progressive storing of SO₂ in the cryosphere, the drop of the CO₂ pressure below 2 bar results in the appearance of SO₂ and sulfate aerosols in the atmosphere, due to both episodic volcanism and continuous release of SO₂ previously stored in the cryosphere, with the subsequent acidification and formation of sulfate minerals at the surface. The cryosphere is progressively depleted in SO₂. At this time, if a significant level of the SO₂ mixing ratio is maintained in the atmosphere, including during the periods between volcanic eruptions due to the continuous SO₂ release from the SO₂-rich cryosphere, the surface temperature is stabilized close to the CO₂ clathrate equilibrium temperature of ~220 K (210-230 K in the p_{CO2} range from 1-2 bar). Interestingly, the time when the CO₂ pressure dropped below 2 bar, resulting in sulfate precipitation at the surface, subsequent acidification of the surface and formation of sulfate minerals, could coincide with the age of sulfate minerals observed by OMEGA/ Mars Express (Bibring et al., 2006). If so, the CO₂ pressure had to be of the order of 2 bar at the late Noachian/ Hesperian transition.

The chronology of clays, sulfates and ferric oxides proposed by Bibring et al. (2006) is shown in Figure 6. We have scaled the CO₂ evolution profile calculated by Chassefière and Leblanc (2011a) by multiplying p_{CO2} by 2.5, in such a way to fit a pressure of ~2 bar at the end of the Noachian, just before sulfate minerals observed by OMEGA were emplaced. The estimated time profile of the crust production rate is also shown (Greeley and Schneid, 1991). The vertical grey band corresponds to the time interval when volcanic activity was significant according to Greeley and Schneid estimates. If the CO₂ pressure dropped below 2 bar at the end of the Noachian, the high volcanic activity during the first half of the Hesperian, when observed sulfate minerals formed, together with the release of SO₂ trapped in the cryosphere during Noachian volcanic events, should have resulted in the precipitation of large amounts of sulfate aerosols, with subsequent acidification of the surface and formation of sulfate minerals. The transformation of the volatile SO₂ from accessible near-surface clathrate reservoir, stored

Fig. 6

in the Noachian 2-bar regime, to sulfate sediment reservoir is not reversible since there is no efficient sediment recycling on Mars due to plate tectonics. However, Martian impacts may have contributed to recycling. The drop down of surface pressure has permitted the destabilization mechanism to be more and more efficient. The onset of sulfate mineral formation would have been triggered by the drop in CO₂ pressure below 2 bar. The end of sulfate mineral formation would be due to the sharp decrease of volcanic activity at the end of the Hesperian, and the total transformation from CO₂-SO₂ clathrate to sediment.

The origin of a possible CO₂ pressure drop at the end of the Noachian is unclear. Non-thermal escape is not expected to have removed more than ~10 mbar during the last 4 Gyr (Chassefière and Leblanc, 2011a). The sink for CO₂ is necessarily within the crust and formation of carbonates in subsurface hydrothermal systems may seem a good candidate (Griffith and Shock, 1995; Chassefière and Leblanc, 2011a). The reason for a stage of intense hydrothermalism at the early Hesperian could be the occurrence of a global scale episode of fissural volcanism between 3.8 and 3.6 Ga (Ody et al., 2012). The olivine-rich magma could have reached the surface through fractures and cracks in the basement rock caused by impacts during the late heavy bombardment (Ody et al., 2012). Such a context, with a highly fractured upper crust filled with both magma and liquid water from the melting cryosphere, may have favored the circulation of CO₂-rich hot waters in the subsurface and the deposition of carbonates in crustal cold water pools (Chassefière and Leblanc, 2011a). This episode could also be responsible for the trapping of large amounts of water in the crust by serpentinization (Chassefière and Leblanc, 2011b). Because in the same time, according to the massive sulfur precipitation scenario, the cryosphere would have lost most of its SO₂, less and less acidic conditions would have prevailed in the subsurface hydrosphere, making more and more efficient the precipitation of carbonates. These hypotheses remain to be confirmed by further studies.

470

471 Assuming (i) 2×10^{21} to 5×10^{21} g as the total sulfur degassed in the Noachian by the Tharsis
472 province (Gaillard et al. 2009), (ii) an equilibrium flux of S outgassing and sulfate
473 precipitation in the range from 2×10^9 to 2×10^{10} molecule $\text{cm}^{-2}\text{s}^{-1}$ for an atmospheric SO_2
474 mixing ratio in the range from 1 to 200 ppm (T10), we can estimate that the complete S
475 content from the clathrate reservoir is released in a time range from 40 to 1000 Myrs. The
476 cryosphere in the Noachian 2-bar regime has to be quite rich in SO_2 . Assuming for example a
477 SO_2/CO_2 ratio in the clathrate of 20%, the required water GEL thickness is in the range from
478 250 to 630 m, for a CO_2 global equivalent pressure in the range from 3 to 7.5 bar. A twice
479 lower ratio of 10% results in twice higher H_2O and CO_2 inventories (500-1300 m/ 6-15 bar).
480 A twice larger ratio of 40% results in twice lower H_2O and CO_2 inventories (125-315 m/ 1.5-
481 3.7 bar). These orders of magnitude estimates are reasonable for the global inventory of
482 Martian volatiles, as described in Section 1, implying a SO_2/CO_2 ratio in the clathrates of
483 several tens percents. If the sulfate is deposited in form of anhydrite (Gaillard et al. 2009) at
484 136 g.mol^{-1} , it will precipitate in a 20 to 50 m global layer.

485

486 Two majors geomorphological units are compatible with our scenario:

487

488 1) The presence of a thick Mars-wide sedimentary formation of sulfate, including in very
489 uncommon places at high topography, such the Interior Layer Deposit (ILD) at the top of
490 Valles Marineris. Eolian deposition was proposed from geomorphological arguments
491 (Michalski et al., 2011). For ILD in Valles Marineris, the formation as thick as 5 km (Nedell
492 et al., 1987) must be done in 400 Myrs, after the tectonic opening at 3.9 Ga and the formation
493 of the floor 3.5 Ga (Quantin et al., 2004). In our scenario, 20 m to 50 m GEL sulfur particle
494 could have precipitated directly from atmosphere in 40 to 1000 Myrs, in agreement with the
495 observation. The sedimentary formation of sulfate seems to be at very large scale (Bibring et

al., 2006) since it has been detected in different places on Mars such as Valles Marineris (Bibring et al., 2007, Le Deit et al, 2008), Meridiani Planum (Squyres et al., 2004), Aram Chaos (Massé et al., 2008) and Syrtis Major (Ehlmann et al., 2012).

2) The chaotic terrains, at equatorial region, could have been formed by disruption of the CO₂ clathrate in the past at the late Hesperian/Amazonian period. This interpretation is compatible with our scenario since the formation of clathrate must have been global during the Noachian, it should also affect equatorial region. The reason for such a disruption has been debated, including climatic change (Clifford and Parker, 2001), internal heat flux (Clifford and Parker, 2001), fracture propagation (Rodriguez et al., 2006), and seismic activity (Nummedal and Prior, 1981; Tanaka, 1999). We propose here that the disruption may be due to the pressure decrease below 2 bar. In our scenario, a GEL of CO₂-SO₂ clathrate ranging from 290 m to 720 m depth has been destabilized. Scaling this volume to the province of the chaos (1/10 surface of Mars) is compatible with the typical height loss in chaos ~3000 m. A schematic representation of the evolution of early Mars' under the effect of CO₂-SO₂ clathrate formation and destabilization is shown in Figure 7.

Fig.7

As previously stated, the surface temperature of ~230 K proposed to have prevailed during the Noachian is a global average value due to the use of a 1-D model. In the real Martian surface, there are places where the temperature is higher than 230 K, and clathrates don't form, for instance near the equator at low obliquity. They form only in the locations where the temperature reach 230 K, for instance at the pole at low obliquity. Due to the large variations of the obliquity of Mars on typical time periods of 10⁵-10⁶ yr (Laskar et al., 2002; Levrard et al., 2004), the colder regions of the planet regularly become warmer, and vice versa. We don't know if variations of orbital parameters similar to those calculated for the last ten million years occurred on early Mars. This is still unknown, especially under the scenario of giant

planet migration (Gomes et al., 2005; Morbidelli et al., 2007). Some chaotic variations of obliquity may have occurred at the Noachian (before giant planet migration) in specific dynamical conditions, but in other conditions obliquity may have remained stable (Brasser and Walsh, 2011). Making the hypothesis that early Mars obliquity changed in a similar fashion as it does today, a coming and going of ice deposits, including CO₂-SO₂ clathrates, between high latitude (at low obliquity) and low latitude (at high obliquity) regions through alternate sequences of sublimation and condensation should have occurred. On the recent Mars (last 10 Myr), the typical time scale for ice migration during a transition of the mean obliquity, like that which occurred 4 Myr ago, is ~1 Myr (Levrard et al., 2007). On ancient Mars, where water was more abundant and the CO₂ pressure higher, this time should have been larger. Assuming that the residence time of SO₂ in the atmosphere with respect to clathrate formation on the cold trap is 1 yr (see below), and that the equivalent of 1 bar of SO₂ is transferred in 1 Myr, the average SO₂ mixing ratio in a 2-bar atmosphere during an obliquity transition is 0.5 ppm. For an ice migration time of 10 Myr, and an amount of transferred SO₂ of 0.2 bar, the average SO₂ mixing ratio during obliquity transition is 10⁻² ppm. These values are small, with no expected effect on climate. An important question is to know if, during its transfer through the atmosphere, SO₂ is preserved from forming aerosols and being removed from the atmosphere-cryosphere system. The residence time of SO₂ in the atmosphere, before being removed by aerosol deposition, is 10²-10³ yr (Johnson et al., 2009, T10). The global mixing time of Mars atmosphere is ~0.5 yr (Krasnopolsky, 2005), of the same order as the Earth troposphere mixing time (~1 yr). The lifetime of a SO₂ molecule with respect to trapping in clathrates is therefore 2 to 3 orders of magnitude smaller than the time of SO₂ removal by aerosol deposition. SO₂ is therefore expected to survive ice migrations, making possible the preservation of a SO₂-rich cryosphere during the whole Noachian.

It must be emphasized that the threshold pressure value of 2 bar is obtained by using the 1-D model. The mean annual surface temperature is not uniform, decreasing from the equator to the poles. On Mars today, regions above 45° latitude in both hemispheres have an annual surface temperature lower by more than 10 K than the global average temperature (see e.g. Mellon et al., 2003). As shown on Figure 2, the difference between the mean global temperature and the CO₂ clathrate formation temperature is smaller than ≈10 K in the range 0.5-2 bar. By analogy with present Mars, it results that even with a CO₂ pressure of only 0.5 bar, regions above 45° latitude could have been below the clathrate formation temperature in annual average. If so, it may be thought that the cold trap formed by middle and high latitude regions may have been efficient enough to remove most of the SO₂ released to the atmosphere (including in equatorial regions) by volcanism. More accurate 2-D or 3-D models are required to precisely define the value of the pressure threshold, which could be smaller than 2 bar and even possibly 1 bar. Such a lower threshold value would be in better agreement with a strong carbon hydrodynamic escape ruling out a dense Martian atmosphere until the late Noachian (Tian et al., 2009), and the recent estimate of <1 bar obtained from the constraint provided by the ⁴⁰Ar/³⁶Ar ratios of trapped gases within Martian meteorite ALH 84001 (Cassata et al., 2012).

5. Influence of the presence of SO₂ on the trapping of argon, krypton and xenon in clathrates

Mousis et al. (2012) found recently that it was possible to account for the two orders of magnitude drop existing between the measured atmospheric abundances of non-radiogenic argon, krypton and xenon in Earth versus Mars (see Pepin, 1991) by invoking the trapping of these noble gases in clathrate deposits incorporated into the current Martian cryosphere. In their scenario, these authors estimated that masses of xenon, krypton and argon equivalent to

those found on Earth could be incorporated into clathrates if one assumes the simultaneous trapping of at least 2.3 bar of CO₂. This value is quite close to the maximum value of 2 bar imposed by the CO₂ clathrate saturation law, as previously explained, and is therefore compatible with constraints derived from the present work. Mousis et al. (2012) also considered the presence of atmospheric SO₂ on the clathrate composition. In order to quantify the influence of this species on the noble gas trapping in CO₂-dominated clathrate, they used approximate combination rules allowing them to retrieve a set of interaction potential parameters for SO₂-H₂O interactions. They then deduced that this species was a poor clathrate former in the presence of a CO₂-dominated gas and concluded that it has zero influence on the trapping efficiencies of other minor species, including noble gases.

The set of potential parameters determined for SO₂-H₂O interactions in the present work supersedes the exploratory one used by Mousis et al. (2012) and suggests that the mole fraction of SO₂ can be strongly enriched in CO₂-dominated clathrate compared to that existing in the gas phase. This implies that the question of the influence of this species on the noble gas trapping in CO₂-dominated clathrate remains open. Here we use a gas phase composition similar to the one defined by Mousis et al. (2012), in which the mole fraction of SO₂ is varied between 10⁻³, 1 and 10³ ppm, in order to investigate the influence of SO₂ on the trapping of other minor species. Abundances of CO₂, N₂, O₂, CO, Ar, Kr and Xe derive from Moroz (1998). In each of the three cases, the abundance of CO₂ is adjusted to allow the normalization to 1 of the sum of the mole fractions.

A set of SO₂-H₂O interaction parameters, which is representative of the range of solutions determined in Section 2 and Appendix A, has been selected ($a = 0.75 \text{ \AA}$, $\epsilon/K = 258.91 \text{ K}$ and $\sigma = 2.7 \text{ \AA}$). These parameters correspond to a mean enrichment factor of 100 of the SO₂ mole fraction in clathrate compared to the atmospheric one at a surface pressure of 1 bar,

irrespective of the considered SO_2 abundance (10^{-3} , 1 and 10^3 ppm). Figure 8 represents the clathrate composition computed in the three cases of the SO_2 abundance for atmospheric pressures up to 3 bar, allowing us to encompass all the plausible values discussed in the literature. At each pressure considered, the temperature used in our computations is the equilibrium temperature of CO_2 -dominated clathrate. The figure shows that the mole fractions of minor species are not altered when considering SO_2 atmospheric abundances ranging from 10^{-3} to 1 ppm. At a SO_2 atmospheric abundance of 10^3 ppm, the trapping efficiencies of Ar and Kr are decreased by a factor of 10 and that of Xe is almost unchanged.

Our calculations imply that the conclusions of Mousis et al. (2012) remain valid if one assumes that the SO_2 atmospheric abundance ranges between 10^{-3} and 1 ppm. However, for higher mixing ratios such as those estimated in the Noachian (see Sec. 4), the efficient clathration of SO_2 implies the trapping of up to several dozens of bar of atmospheric CO_2 if one wants to account for the losses of Ar and Kr via their clathration in the cryosphere. Because these values are implausible, it seems that the scenario proposing that the noble gas deficiency of the Martian atmosphere is due to their efficient sequestration in clathrates present in the cryosphere is not consistent with the one proposed here for SO_2 .

6. Discussion

We suggested in the present study that the formation of CO_2 - SO_2 clathrates at Noachian and Hesperian times could have played an important role in controlling and stabilizing the level of volcanic sulfur in the atmosphere, as well as the level of atmospheric CO_2 for earliest times. If the CO_2 pressure exceeded a threshold of $p_0 = 2$ bar (possibly less : 0.5-1 bar, which remains to be confirmed from more sophisticated 3D models) due to an efficient Noachian volcanism, the CO_2 in excess of p_0 (assumed to be ~ 2 bar for the discussion) could have been stored in

the cryosphere under the form of CO₂ clathrates. Indeed, due to the increasing albedo of the atmosphere through Rayleigh scattering for increasing CO₂ pressure above 1 bar, the surface temperature induced by a $p > \sim 2$ bar CO₂ atmosphere is smaller than the equilibrium temperature of clathrates, resulting in a saturation of CO₂ and its condensation under the form of clathrates. Such a clathrate buffer would have maintained the CO₂ pressure close to ~ 2 bar, and the surface temperature close to 230 K, resulting in a cold Mars at the Noachian. The cryosphere would have trapped all the sulfur released by volcanism under the form of sulfur-rich (enriched by a factor ~ 100 -500 with respect to the gas phase) CO₂-SO₂ clathrates.

Because the thermal conductivity of clathrates is low, smaller than that of water ice on the order of 5 to 6 times (Mellon, 1996), the formation of clathrate deposits should result in a higher thermal gradient, with an accumulation of heat beneath the clathrate layer. It has been suggested that, on present Mars, no more than a global equivalent atmospheric pressure of 112 mbar of CO₂, in a 4-km thick polar deposit, may be contained in the polar caps. For a larger amount of deposited CO₂, the warming at the base of the clathrate layer results in the liquefaction of CO₂, essentially precluding the stability of the system. On early Mars, the deposition of large amounts of CO₂ clathrates, assuming an atmospheric content larger than 2 bar, could have led to this kind of instability. But if clathrates were deposited on an extended fraction of the surface, much larger than the present caps, the maximum admissible value of the global equivalent atmospheric pressure of CO₂ trapped in clathrates could be much larger than 112 mbar. This phenomenon should have to be taken into account in any detailed modeling of massive clathrate deposition in the Noachian cryosphere.

For a surface pressure smaller than 2 bar and progressively decreasing, atmospheric sulfur is no longer trapped in clathrates, and an increasing fraction of the atmospheric SO₂, up to 10 ppm at $p_{\text{CO}_2}=1.5$ bar, and 30 ppm at $p_{\text{CO}_2}=1$ bar, can remain in the atmosphere. If the amount

of atmospheric SO₂ exceeds these thresholds, the cooling effect of sulfate aerosols results in a decrease of surface temperature below the clathrate equilibrium temperature and the condensation of the SO₂ in excess in CO₂-SO₂ clathrates. During this period, significant amount of SO₂ (and H₂S rapidly converted in SO₂ by atmospheric photochemistry) was converted into sulfate aerosols, which further settled down to the surface and possibly led to the formation of sulfate minerals. This sulfur may have been released, not only by volcanoes but also (and more continuously) by the SO₂-rich cryosphere formed when p_{CO2}>~2 bar. Through this mechanism, the cryosphere may have lost at this stage some of the volcanic SO₂ stored at earlier times, released back to the atmosphere. In this time range (2 to 1 bar p_{CO2} range), the surface temperature has been buffered, during and after episodes of sulfur release, at a temperature from 230 K (p_{CO2}=2 bar) to 210 K (p_{CO2}=1 bar). For p_{CO2}<1 bar, all the released SO₂ remains in the atmosphere, with no more trapping in clathrates.

We proposed the hypothesis that the formation of sulfate minerals, which have been observed from orbit and are formed during the Hesperian (Bibring et al., 2006), could have been triggered by a fall of early Mars atmosphere pressure below 2 bar and that this change occurred close to the late Noachian/Hesperian transition. This hypothesis requires that the total amount of CO₂ in both the atmosphere and the cryosphere during the Noachian has been greater than the content of a 2 bar atmosphere as a result of substantial volcanic outgassing and/or rapid carbonate formation, which might have been efficient enough to keep some CO₂ on early Mars despite rapid carbon escape. This constraint could be alleviated if further 3D-modeling shows that a pressure threshold value of 0.5-1 bar is sufficient to keep most of the released SO₂ trapped under clathrate form in the Noachian cryosphere. According to our hypothesis, Mars' cryosphere would have stored under the form of CO₂-SO₂ clathrates a significant amount of the sulfur outgassed during the whole Noachian (from the magma ocean crystallization to the 2-bar transition). At the transition between Noachian and Hesperian, this

reservoir would have been released back into the atmosphere through clathrate disruption that may be at the origin of the chaotic terrains. Then sulfur has precipitated in the form of aerosols, creating a large-scale sulfate deposit, in agreement with several in-situ and orbital observations. Today, the sulfate layer is not present everywhere but observed in some places such Valles Marineris (Bibring et al., 2007, Le Deit et al, 2008), Meridiani Planum (Squyres et al., 2004), Aram Chaos (Massé et al., 2008) and Syrtis Major (Ehlmann et al., 2012). First, the net aerosols precipitation flux in the ancient Mars should be controlled by local winds and topography in an analogous manner than snow precipitation (Forget et al., 2006). Second, the erosion may have removed significant part of the initial layer, especially because sulfate is highly transportable in soluble phase, such kieserite.

According to our scenario, the formation of carbonates should have been inhibited (at least in the superficial cryosphere) during most of the Noachian, although probably not at earliest times when the cryosphere was still SO₂-free, due to the acidification of water resulting from the local melting of the SO₂-rich cryosphere. Such fluid can precipitate sulfate phase locally in the subsurface but the cryosphere may be SO₂-rich enough (it can potentially store the complete SO₂ degassed from volcanism, see Sec. 4) implying unlikely conditions for carbonates precipitation. As previously mentioned it may be thought that during the massive release of SO₂ to the atmosphere after the end of the 2-bar stage at the Noachian-Hesperian transition, and the subsequent sequestration of sulfur in sediments, a less acidic cryosphere allowed the formation of carbonates, relatively deep in the subsurface due to cold and acidic surface conditions. Because the trapping of CO₂ in carbonates results in a decrease of the atmospheric pressure, and therefore a release of SO₂ to the atmosphere and its further sequestration in sediments, a positive feedback could have occurred. The removal of CO₂ in carbonates would have allowed SO₂ to be released and trapped in sulfate sediments, decreasing the acidity of the cryosphere and allowing more carbonates to be formed. Such a

runaway process could be one of the reasons for the sharp decrease of atmospheric CO₂ during the Hesperian, simultaneously with the formation of large sediment deposits, but remains a hypothesis to be further explored.

According to our scenario in the pre-Noachian and Noachian periods, the massive presence of sulfur in the cryosphere/atmosphere has kept low global temperature (near 230 K), in agreement with the results of clay minerals observations (Ehlmann et al., 2011). Under such conditions, clays may have been formed by alteration of the crust in the locally warmer region, where no SO₂-CO₂ clathrate can condense (equator during low obliquity). The low atmospheric SO₂ gas content may lead to acidic pH preventing carbonate from precipitating but permitting clay formation near the surface. More detailed modeling are required to discuss these implications.

Interestingly, the massive presence of sulfur in the pre-Noachian and Noachian cryosphere could have favored the occurrence of warm episodes, with liquid water at the surface and a global hydrological cycle. Such a warm episode, or series of warm episodes, with a typical duration of 10⁵-10⁷ years, could have allowed the formation of valley networks observed in the southern hemisphere (Hoke et al., 2011), and formed close to the late Noachian/Hesperian transition. Similar episodes seem to be required at the middle Noachian to explain the high degree of degradation of ancient craters (N. Mangold, personal communication, 2012). It has been suggested that SO₂, in addition to CO₂, could have played a major role in maintaining water in a liquid state at the surface of Mars for significant periods of time, with an ocean and a global hydrological cycle, and that such an episode could have been controlled by a well-developed SO₂ cycle (Halevy et al., 2007). The major objection in our scenario is that, starting from a cold state, any release of SO₂ to the atmosphere is followed by the formation of sulfate aerosols and a subsequent cooling of the atmosphere (T10). But any

catastrophic event such as a giant asteroid impact or a massive volcanic SO₂ release or a large-scale destabilization of methane clathrate from a highly magmatically active subsurface, might have been able to melt the superficial cryosphere and release huge amounts of SO₂ and H₂O (and possibly CO₂) into the atmosphere. Provided such a catastrophic event was able to trigger a global hydrological cycle, even for a short time of a few years or hundred years, the SO₂ (and CO₂) released at the same time from the melting cryosphere could have taken the relay and driven a somewhat longer period of warm and wet Mars, according to Halevy et al. scenario. Episodes of huge methane release from clathrates present on oceanic floors occurred during the history of Earth. At the Paleocene-Eocene transition, 55.5 Myr ago, massive dissociation of oceanic methane hydrate led to a warming of 5-6 °C during ~1000 yr (see e.g. Higgins and Schrag, 2006). The injection of CH₄ into early Mars atmosphere with a mixing ratio of a few percents, that is partial pressure of a few tens millibar, may have possibly raised the surface temperature above 0°C (Kasting, 1997), but further calculations by Tian (unpublished) don't confirm that CH₄, even at several percent level, is able to bring surface temperature close to the freezing point of water. Understanding if massive methane injections in the atmosphere of Mars, and/or giant impacts, have been able to initiate a SO₂ cycle-driven episode of warm and wet Mars (Halevy et al., 2007), breaking the background cold Noachian climate, is of prime interest to progress in our understanding of early Mars climate.

Whatever the precise triggering mechanisms of such possible warm episodes favored by the release of massive amounts of SO₂ together with CO₂ to the atmosphere may have been, our hypothesis may potentially explain the occurrence of outflow channels, valley networks and the degradation of impact craters. In Sec. 4, we suggested that the destabilization of clathrates when p_{CO2} dropped below 2 bar could be at the origin of chaotic terrains, and it may therefore as well explain the formation of outflow channels, provided sufficient amounts of water were present in the subsurface. Valley networks, formed at the late Noachian and which ended

relatively sharply at the beginning of the Hesperian (Fasset and Head, 2011), could be the result of warm episodes occurring when the CO₂ pressure was still close to 2 bar, and the temperature relatively “high” (230 K). Such episodes could explain the formation of some sulfate deposits during the Noachian. The postulated drop in pressure at the Noachian/Hesperian transition resulted in a general cooling of the planet from 230 K to 220 K (for p_{CO2}=1 bar), then 210 K (for p_{CO2}=0.5 bar) (see Fig. 2). Provided this drop was sharp enough, the cooling of the planet by ≈20 K during the Hesperian could have rapidly inhibited warm episodes possibly at the origin of valley networks. A recent study shows that the degradation of impact craters experienced a sharp transition at 3.7 Ga (Mangold et al., 2012). This transition between a period of substantial erosion due to a significant global fluvial activity (late Noachian) and a period of more regional and sporadic warming episodes (Hesperian) may be the result of the suggested drop in pressure, resulting in both a general cooling of the planet and episodes of massive SO₂-CO₂ release. The necessarily sporadic nature of clathrate destabilization during the Hesperian could be at the origin of the degradation of Hesperian craters, showing no relationship between age and degradation state (Mangold et al., 2012). In order to confirm these views, it will be necessary to build reliable scenarios of such warming episodes, possibly triggered by specific events (impacts, massive methane release, obliquity variations...) and favored by massive SO₂ release together with CO₂. If these events are intense enough, liquid water precipitation (rain) may incorporate atmospheric SO₂ so that the SO₂ aerosol deposition time could be shorter than in our arid model (SO₂ aerosols precipitation), implying a reduced cooling effect of aerosols, not sufficient to counteract the warming effect of SO₂ and other greenhouse gases. These extreme events provide also SO₂ precipitation before the Noachian-Hesperian boundary that could explain observed geological units. Also, clays may have been formed during such favorable climatic excursions (see e.g. Fan et al., 2008 ; Murchie et al., 2009 ; Wray et al., 2011).

Our results are based on extrapolations to low temperatures of the results obtained in Earth's laboratory conditions. In order to progress in our understanding of the role of the sulfur cycle in the early evolution of Mars, experimental measurements devoted to the measurements of CO₂-SO₂ clathrates thermodynamical and kinetic properties in Martian conditions would be necessary. Future space missions to Mars could also bring new results. Search for SO₂ plumes associated with outgassing from surviving buried cryospheric SO₂ clathrate-rich reservoirs by the Exomars TGO mission in 2016, and mapping of SO₂ (if detected), could help in characterizing a possible residual activity of sulfur in the Martian crust. SO₂ has never been detected in the Martian atmosphere, with a measured upper limit at 2 σ of 1 ppb (Krasnopolsky, 2005). Nevertheless, if some SO₂ has been trapped at depth in long life time clathrate layers, an episodic release due to tectonic or subsurface thermal events triggering the destabilization of clathrates cannot be ruled out. The dielectric constants of clathrates are too close to that of water ice to allow to them to be easily distinguished at radar wavelengths (Mousis et al., 2012), which makes subsurface clathrate layers hardly detectable from orbit. Buried reservoirs of SO₂ and H₂S clathrate-hydrates could be searched for through drilling from future landers on Mars. If some sulfur clathrate deposits survived close to some volcanoes, a concentric structure might be observed, like e.g. around the Hakon Mosby mud volcano in Marmara Sea where concentric layers of gas hydrates are observed, with a maximum concentration of clathrates in the sediment of 10-20% at about five hundred meters of the volcano (Fig. II.17 in Bourry, 2008). In a similar way, although in quite different conditions (subaerial volcanism, larger space and time scales), concentric deposits of CO₂-SO₂-H₂S clathrates could have been formed on Mars around volcanoes, with concentrations and composition expected to depend on the distance to the outgassing plume. The possibility for such deposits to have survived a long time is of course highly speculative, and we don't know at which depth they could have been preserved.

Acknowledgments : E. Chassefière, E. Dartois, A. Lakhlifi and F. Schmidt thank the interdisciplinary EPOV program of CNRS. O. Mousis acknowledges support from CNES. We feel indebted to the referees, M. Mischna and A. Fairén, for their constructive comments that allowed to significantly improve the quality of this paper. We thank L. Daumas for the drawing of figure 7.

References :

- Adisasmito, S., Frank, R.K., Sloan, E.D., 1991, J. Chem. Eng. Data, 36, 68-71.
- Baldrige, A. M., Hook, S. J., Crowley, J. K., Marion, G. M., Kargel, J. S., Michalski, J. L., Thomson, B. J., de SouzaFilho, C. R., Bridges, N. T., Brown, A. J., 2009. Contemporaneous deposition of phyllosilicates and sulfates: Using Australian acidic saline lake deposits to describe geochemical variability on Mars, Geophys. Res. Lett. 36, L19201, doi:10.1029/2009GL040069.
- Bibring, J.-P., Langevin, Y., Mustard, J. F., Poulet, F., Arvidson, R., Gendrin A., Gondet, B., Mangold, N., Pinet, P., Forget, F., 2006. Global Mineralogical and Aqueous Mars History Derived from OMEGA/Mars Express Data, Science 312, 5772, 400-404.
- Bibring, J.-P., Arvidson, R. E., Gendrin, A., Gondet, B., Langevin, Y., Le Mouélic, S., Mangold, N., Morris, R. V., Mustard, J. F., Poulet, F., Quantin, C., Sotin, C., 2007, Coupled Ferric Oxides and Sulfates on the Martian Surface, Science 317, 1206-1210.
- Bourry, C., 2008. Caractérisation physique et géochimique d'hydrates de gaz d'environnements géologiques différents, PhD thesis, University of Brest, France.
- Brasser, R., Walsh, K.J., 2011. Stability analysis of the Martian obliquity during the Noachian era, Icarus 213, 423–427.

831 Cassata, W.S., Shuster, D.L., Renne, P.R., Weiss, B.P., 2012. Trapped Ar isotopes in
832 meteorite ALH 84001 indicate Mars did not have a thick ancient atmosphere, *Icarus* 221,
833 461–465.

834 Chassefière, E., Leblanc, F., 2011a. Methane release and the carbon cycle on Mars. *Planet.*
835 *Space Sci.* 59, 207–217.

836 Chassefière, E., Leblanc, F., 2011b. Constraining methane release due to serpentinization by
837 the D/H ratio on Mars, *Earth Planet. Sci. Lett.* 310, 262–271.

838 Christensen, P., 2006. Water at the Poles and in Permafrost Regions of Mars,
839 *Elements* 2, 151–155.

840 Clifford, S. M., Parker, T. J., 2001, The Evolution of the Martian Hydrosphere: Implications
841 for the Fate of a Primordial Ocean and the Current State of the Northern Plains, *Icarus* 154,
842 40–79.

843 Clifford, S.M., Lasue, J., Heggy, E., Boisson, J., McGovern, P., Max, M.D., 2010. Depth of
844 the Martian cryosphere: Revised estimates and implications for the existence and detection
845 of subpermafrost groundwater, *J. Geophys. Res.* 115, E07001, doi:10.1029/2009JE003462.

846 Colaprete, A., Toon, O. B., 2003. Carbon dioxide clouds in an early dense Martian
847 atmosphere. *J. Geophys. Res.* 108, 5025, doi:10.1029/2002JE001967.

848 Craddock, R. A., Greeley, R., 2009. Minimum estimates of the amount and timing of gases
849 released into the martian atmosphere from volcanic eruptions. *Icarus*. 204, 512–526.

850 Danesh, A., 1998, *Phase Behaviour of Petroleum reservoirs Fluids*, Elsevier Science BV,
851 ISBN 0444821861.

852 Ehlmann, B.L., Mustard, J.F., Murchie, S.L., Bibring, J.-P., Meunier, A., Fraeman, A.A.,
853 Langevin, Y., 2011. Subsurface water and clay mineral formation during the early history
854 of Mars, *Nature* 479, 53, doi:10.1038/nature10582.

855 Ehlmann, B. L., Mustard, J. F., 2012, An in-situ record of major environmental transitions on
856 early Mars at Northeast Syrtis Major, *Geophys. Res. Lett.*, 39, L11202.

857 Fairén, A. G., et al., 2004. Inhibition of carbonate synthesis in acidic oceans on early Mars.
858 Nature. 431, 423-426.

859 Falabella, B.J., Vanpee, M., 1974. Experimental Determination of Gas Hydrate Equilibrium
860 below the Ice Point. Ind. Eng. Chem. Fundam.13, 228-231.

861 Falenty A., Genov G., Hansen Th. C., Kuhs W. F., Salamatina A. N., Falenty, 2011, Kinetics
862 of CO₂ Hydrate Formation from Water Frost at Low Temperatures: Experimental Results
863 and Theoretical Model. J. Phys. Chem. C 115, 4022–4032.

864 Fan, C., Schulze-Makuch, D., Fairén, A.G., Wolff, J.A., 2008. A new hypothesis for the
865 origin and redistribution of sulfates in the equatorial region of western Mars, Geophys.
866 Res. Lett. 35, L06201, doi:10.1029/2007GL033079.

867 Fasset, C. I., Head, J.W., 2011. Sequence and timing of conditions on early Mars, Icarus 211,
868 1204–1214.

869 Forget, F., Pierrehumbert, R. T., 1997. Warming Early Mars with Carbon Dioxide Clouds
870 That Scatter Infrared Radiation. Science. 278, 1273-1276.

871 Forget, F., Haberle, R. M., Montmessin, F., Levrard, B. & Head, J. W., 2006, Formation of
872 Glaciers on Mars by Atmospheric Precipitation at High Obliquity, Science, 311, 368-371

873 Gaillard, F., Scaillet B., 2009. The sulfur content of volcanic gases on Mars. Earth Planet. Sci.
874 Lett., 279, 1-2, 34-43.

875 Gaillard, F., Michalski, J., Berger, G., McLennan, S., Scaillet, B., 2012. Geochemical
876 reservoirs and timing of sulphur cycling on Mars, Space Sci. Rev., in press, doi
877 10.1007/s11214-012-9947-4.

878 Gomes, R., Levison, H., Tsiganis, K., Morbidelli, A., 2005, Origin of the cataclysmic Late
879 Heavy Bombardment period of the terrestrial planets, Nature 435, 466-469.

880 Griffith, L. L., Shock, E. L., 1995. A Geochemical Model for the Formation of Hydrothermal
881 Carbonates on Mars. Nature. 377, 406-408.

882 Greeley, R., Schneid, B.D., 1991. Magma generation on Mars - Amounts, rates, and
883 comparisons with earth, moon, and Venus, *Science* 254, 1991, 996-998.

884 Grott, M., et al., 2011. Volcanic outgassing of CO₂ and H₂O on Mars. *Earth and Planetary*
885 *Science Letters*. 308, 391-400.

886 Halevy, I., Zuber, M. T., Schrag, D. P., 2007. A Sulfur Dioxide Climate Feedback on Early
887 Mars, *Science* 318, 5858, 1903-1907.

888 Handa, Y.P. , Tse, J.S., 1986, Thermodynamic properties of empty lattices of structure I and
889 structure II clathrate hydrates, *J. Phys. Chem.* 23, 5917-5921.

890 Hanna, J. C., Phillips, R.J., 2006, Tectonic pressurization of aquifers in the formation of
891 Mangala and Athabasca valleys, Mars, *J. Geophys. Res.* 111, E03003,
892 doi:10.1029/2005JE002546.

893 Head, J. W., Pratt, S., 2001. Extensive Hesperian-aged south polar ice sheet on Mars:
894 Evidence for massive melting and retreat, and lateral flow and ponding of meltwater, *J.*
895 *Geophys. Res.* 106, E6, 12275-12300.

896 Head, J. W., Marchant, D. R., 2009. Inventory of Ice-related Deposits on Mars: Evidence for
897 Burial and Long-Term Sequestration of Ice in Non-Polar Regions and Implications for the
898 Water Budget and Climate Evolution, 40th Lunar and Planetary Science Conference,
899 (Lunar and Planetary Science XL), March 23-27, 2009 in The Woodlands, Texas, id.1356.

900 Herri, J.-M., Bouchemoua, A., Kwaterski, M., Fezoua, A., Ouabbas, Y., Cameirao, A., 2011.
901 Gas Hydrate Equilibria from CO₂-N₂ and CO₂-CH₄ gas mixtures – Experimental studies
902 and Thermodynamic Modelling. *Fluid Phase Equilibria* 301, 171-190.

903 Herri, J.M., Chassefière, E., 2012, Carbon dioxide, argon, nitrogen and methane clathrate
904 hydrates: thermodynamic modelling, investigation of their stability in Martian atmospheric
905 conditions and variability of methane trapping, *Planet. Space Sci.*, in press, doi
906 10.1016/j.pss.2012.07.028.

907 Higgins, J.A., Schrag, D.P., 2006. Beyond methane: Towards a theory for the Paleocene–

908 Eocene Thermal Maximum, *Earth Planet. Sci. Lett.* 245, 523-537.

909 Hoke, M. R. T., Hynek, B.M., Tucker, G.E., 2011, Formation timescales of large Martian
910 valley networks, *Earth Planet. Sci. Lett.* 312, 1-12.

911 Johnson, S. S., Mischna, M. A., Grove, T. L., Zuber, M. T., 2008. Sulfur-induced greenhouse
912 warming on early Mars, *J. Geophys. Res.* 113, E8, CiteID E08005.

913 Johnson, S.S., Pavlov, A.A., Mischna, M.A., 2009. Fate of SO₂ in the ancient Martian
914 atmosphere: Implications for transient greenhouse warming, *J. Geophys. Res.* 114, E11011,
915 doi:10.1029/2008JE003313.

916 Kasting, J.F., 1997. Warming early Earth and Mars, *Science* 276, 1213-1215.

917 King, P.P., McLennan, S.M., 2010. Sulfur on Mars, *Elements* 6, 107-112.

918 Krasnopolsky, V.A., 2005. A sensitive search for SO₂ in the martian atmosphere: Implications
919 for seepage and origin of methane, *Icarus* 178, 487–492.

920 Laskar, J., Levrard, B., Mustard, J. F. 2002. Orbital forcing of the martian polar layered
921 deposits. *Nature* 419, 375–377.

922 Le Deit, L., Le Mouélic, S., Bourgeois, O., Combe, J.-P., Mège, D., Sotin, C., Gendrin, A.,
923 Hauber, E., Mangold, N., Bibring, J.-P., 2008, Ferric oxides in East Candor Chasma,
924 Valles Marineris (Mars) inferred from analysis of OMEGA/Mars Express data:
925 Identification and geological interpretation», *J. Geophys. Res.* 113, Issue E7, CiteID
926 E07001.

927 Levrard, B., Forget, F., Montmessin, F., Laskar, J., 2004. Recent ice-rich deposits formed at
928 High latitude on Mars by sublimation of unstable equatorial ice during low obliquity,
929 *Nature* 431, Issue 7012, 1072-1075.

930 Mckoy, V., Sinagoglu, O. J., 1963. Theory of dissociation pressures of some gas hydrates. *J*
931 *Chem. Phys.* 38, 2946-2956.

932 Mangold, N., Adeli, S., Conway, S., Ansan, V., Langlais, B. 2012. A chronology of early
933 Mars climatic evolution from impact crater degradation, *J. Geophys. Res.* 117, E04003,

934 doi:10.1029/2011JE004005.

935 Massé, M., Le Mouélic, S., Bourgeois, O., Combe, J.-P., Le Deit, L., Sotin, C., Bibring, J.-P.,
936 Gondet, B., Langevin, Y., 2008, Mineralogical composition, structure, morphology, and
937 geological history of Aram Chaos crater fill on Mars derived from OMEGA Mars Express
938 data, *J. Geophys. Res.* 113, Issue E12, CiteID E12006.

939 Mellon, M.T., 1996. Limits on the CO₂ content of the Martian polar deposits, *Icarus* 124, 268-
940 279.

941 Mellon, M.T., Feldman, W.C., Prettyman, T.H., 2004. The presence and stability of groundice
942 in the southern hemisphere of Mars, *Icarus* 169, 324–340.

943 Michalski, J. & Niles, P. B., 2011, Origin of Martian Sulfates and Interior Layered Deposits
944 (ILDs) in the Valles Marineris by atmospherically driven processes, EPSC-DPS Joint
945 Meeting 2011, 1752.

946 Miller, S.L., Smythe, W.D., 1970. Carbon Dioxide Clathrate In The Martian Ice Cap, *Science*
947 170, 531-533.

948 Morbidelli, A., Tsiganis, K., Crida, A., Levison, H., Gomes, R., 2007, Dynamics of the giant
949 planets of the solar system in the gaseous protoplanetary disk and their relationship to the
950 current orbital architecture, *Astron. J.* 134:1790-1798.

951 Moroz, V. I., 1998. Chemical composition of the atmosphere of Mars. *Advances in Space*
952 Research 22, 449-457.

953 Mousis, O., Lunine, J.I., Chassefière, E., Montmessin, F., Lakhli, A., Picaud, S., Petit, J.-M.,
954 Cordier, D., 2012. Mars cryosphere: A potential reservoir for heavy noble gases?, *Icarus*
955 218, 80-87.

956 Mousis, O., Chassefière, E., Lasue, J., Chevrier, V., Elwood Madden, L.E., Lakhli, A.,
957 Lunine, J.I., Montmessin, F., Picaud, S., Schmidt, F., Swindle, T.D., 2012. Volatile
958 trapping in Martian clathrates, *Space Sci. Rev.*, in press, doi 10.1007/s11214-012-9942-9.

959 Mumma, M. J., Villanueva, G. L., Novak, R. E., Hewagama, T., Bonev, B. P., DiSanti, M. A.,
 960 Mandell, A. M., Smith, D. M., 2009. Strong Release of Methane on Mars in Northern
 961 Summer 2003, *Science* 323, 1041-1045.

962 Murchie, S.L., Mustard, J.F., Ehlmann, B.L., Milliken, R.E., Bishop, J.L., McKeown, N.K.,
 963 Noe Dobrea, E.Z., Seelos, F.P., Buczkowski, D.L., Wiseman, S.M., Arvidson, R.E., Wray,
 964 J.J., Swayze, G., Clark, R.N., Des Marais, D.J., McEwen, A.S., Bibring, J.P., 2009. A
 965 synthesis of Martian aqueous mineralogy after 1 Mars year of observations from the Mars
 966 Reconnaissance Orbiter, *J. Geophys. Res.* 114, E00D06, doi:10.1029/2009JE003342.

967 Niles, P.B., Catling, D.C., Berger, G., Chassefière, E., Ehlmann, B.L., Michalski, J.R.,
 968 Morris, R., Ruff, S.W., Sutter, B. 2012. Geochemistry of Carbonates on Mars :
 969 implications for climate history and nature of aqueous environments, *Space Sci. Rev.*, in
 970 press, doi 10.1007/s11214-012-9940-y.

971 Nummedal, D., Prior, D. B., 1981, Generation of Martian chaos and channels by debris flows,
 972 *Icarus* 45, 77–86.

973 Ody, A., Poulet, F., Bibring, J.-P., Loiseau, D., Carter, J., Gondet, B., Langevin, Y., 2012.
 974 Global investigation of olivine on Mars : insights into crust and mantle composition,
 975 submitted to *J. Geophys. Res.* (in revision).

976 Owen, T., K. Biemann, D.R. Rushneck, J.E. Biller, D.W. Howarth, LaFleur, A.L. 1977. The
 977 composition of the atmosphere at the surface of Mars. *J. Geophys. Res.* 82, 4635-4639.

978 Parrish, W.R., Prausnitz, J. M., 1972. Dissociation pressure of gas hydrates formed by gas
 979 mixtures. *Ind. Eng. Chem. Process Develop.*, 11, 26-35.

980 Pepin, R. O. 1991. On the origin and early evolution of terrestrial planet atmospheres and
 981 meteoritic volatiles. *Icarus* 92, 2-79.

982 Pollack, J. B., Kasting, J.F., Richardson, F.M., Poliakoff, K., 1987. The Case for a Wet,
 983 Warm Climate on Early Mars. *Icarus*. 71, 203-224.

984 Postawko, S. E., Kuhn, W. R., 1986. Effect of the greenhouse gases (CO₂, H₂O, SO₂) on
 985 Martian paleoclimate. *Journal of Geophysical Research (Proc. Lunar Planet. Sci. Conf.*
 986 *16th)*. 91, D431.

987 Quantin, C., Allemand, P., Mangold, N., Delacourt, C. 2004. Ages of Valles Marineris (Mars)
 988 landslides and implications for canyon history, *Icarus* 172, 2, 555- 572.

989 Rodriguez, J. A. P., Kargel, J., Crown, D. A., Bleamaster, L. F., Tanaka, K. L., Baker, V.,
 990 Miyamoto, H., Dohm, J. M., Sasaki, S., Komatsu, G., 2006. Headward growth of chasmata
 991 by volatile outbursts, collapse, and drainage: Evidence from Ganges chaos, Mars,
 992 *Geophys. Res. Lett.* 33, Issue 18, CiteID L18203.

993 Segura, T.L., Toon, O.B., Colaprete, A., Zahnle, K. 2002. Environmental Effects of Large
 994 Impacts on Mars, *Science* 298, Issue 5600, 1977-1980.

995 Segura, T.L., Toon, O.B., Colaprete, A., 2008. Modeling the environmental effects of
 996 moderate-sized impacts on Mars, *J. Geophys. Res.* 113, Issue E11, CiteID E11007.

997 Sloan, E.D., 1998. *Clathrate hydrates of natural gases*. 2nd Ed. Marcel Decker, New York.

998 Sloan, E.D., Koh, C.A., 2007. *Clathrate hydrates of natural gases*. 3rd Ed. CRC Press.

999 Smith, D. E, Zuber, M. T., Frey, H. V., Garvin, J. B., Head, J. W., Muhleman, D. O.,
 1000 Pettengill, G. H., Phillips, R. J., Solomon, S. C., Zwally, H. J., Banerdt, W. B., Duxbury,
 1001 T. C., Golombek, M. P., Lemoine, F. G., Neumann, G. A., Rowlands, D. D., Aharonson,
 1002 O., Ford, P. G., Ivanov, A. B., Johnson, C. L., McGovern, P. J., Abshire, J. B., Afzal, R. S.,
 1003 Sun, X., 2001. Mars Orbiter Laser Altimeter: Experiment summary after the first year of
 1004 global mapping of Mars, *J. Geophys. Res.* 106, E10, 23689-23722.

1005 Squyres, S. W., Kasting, J. F., 1994. Early Mars: How Warm and How Wet?
 1006 *Science* 265, Issue 5173, 744-749.

1007 Squyres, S. W., Grotzinger, J. P., Arvidson, R. E., Bell, J. F., Calvin, W., Christensen, P. R.,
 1008 Clark, B. C., Crisp, J. A., Farrand, W. H., Herkenhoff, K. E., Johnson, J. R., Klingelhöfer,
 1009 G., Knoll, A. H., McLennan, S. M., McSween, H. Y., Morris, R. V., Rice, J. W., Rieder,

1010 R., Soderblom, L. A., 2004, In Situ Evidence for an Ancient Aqueous Environment at
 1011 Meridiani Planum, Mars, *Science* 306, 1709-1714.
 1012 Tanaka, K. L., 1999, Debris-flow origin for the Simud/Tiu deposit on Mars, *J. Geophys. Res.*
 1013 104, 8637–8652.
 1014 Tian, F., Kasting, J.F., Solomon, S.C., 2009. Thermal escape of carbon from the early Martian
 1015 atmosphere. *Geophys. Res. Lett.* 36, Issue 2, CiteID L02205.
 1016 Tian, F., Claire, M.W., Haqq-Misra, J. D., Smith, M., Crisp, D. C., Catling, D., Zahnle, K.,
 1017 Kasting, J. F., 2010. Photochemical and climate consequences of sulfur outgassing on early
 1018 Mars. *Earth and Planetary Science Letters.* 295, 412-418.
 1019 Udachin, K.A. , Ratcliffe, C.I., Ripmeester, J.A. , 2002, Single Crystal Diffraction Studies of
 1020 Structures I. II and H Hydrates: Structure. Cage Occupancy and Composition, *Journal of*
 1021 *Supra Molecular Chemistry* 2, 405-408.
 1022 Van Berkum. J. G., Diepen. G. A. M., 1979, Phase equilibria in $\text{SO}_2 + \text{H}_2\text{O}$: the sulfur dioxide
 1023 gas hydrate. two liquid phases. and the gas phase in the temperature range 273 to 400 K
 1024 and at pressures up to 400 MPa, *J. Chem. Thermodyn.* 11., 317-334.
 1025 Von Stackelberg, M, 1949, Solid Gas Hydrates, *Die Naturwissenschaften* 36(12), 359-362.
 1026 Wilson, L., and J. W. Head III (2002), Tharsis-radial graben systems as the surface
 1027 manifestation of plume-related dike intrusion complexes: Models and implications, *J.*
 1028 *Geophys. Res.* 107(E8), 5057, doi:10.1029/2001JE001593.
 1029 Wong, A.-S., Atreya, S. K., Encrenaz, Th, 2003, Chemical markers of possible hot spots on
 1030 Mars, *J. Geophys. Res.* 110, E4, 5026, doi:10.1029/2002JE002003.
 1031 Wray, J.J., Murchie, S.L., Squyres, S.W., Seelos, F.P., Tornabene, L.L., 2011. Diverse
 1032 aqueous environments on ancient Mars revealed in the southern Highlands, *Geology* 37, 11,
 1033 1043–1046, doi: 10.1130/G30331A.1.

1034 Yasuda, K., Ohmura, R., 2008. Phase Equilibrium of Clathrate Hydrates Formed with
1035 Methane, Ethane, Propane or Carbon Dioxide at temperatures below the Freezing Point of
1036 Water. J. Chem. Eng. Data 53, 2182-2188
1037
1038

Appendix A

A.1 Description of the statistical-thermodynamic model

Three main clathrate hydrate structures have been identified: SI, SII and SH. They differ by their crystallographic structure in which water is organized in a three dimensional network. It liberates internal cavities of different polyhedral types called 5^{12} , $5^{12}6^2$, $5^{12}6^4$, $4^35^66^3$ and $5^{12}6^8$ (e^f describes a polyhedron: e is the number of edges of the face, and f is the number of faces with e edge). In Table A1 are described more precisely the SI and SII structures, the major ones to be formed due to the composition of Martian gases.

Table A1

In the case of clathrate hydrates in thermodynamic equilibrium, below the water freezing point, the equality of chemical potentials of water in the ice phase and in the hydrate phase can be written, by introducing a hypothetical reference state (phase β) corresponding to a hydrate with empty cavities:

$$\Delta\mu_w^{H-\beta} = \Delta\mu_w^{I-\beta}, \quad (1)$$

where $\Delta\mu_w^{H-\beta}$ and $\Delta\mu_w^{I-\beta}$ are the differences of the chemical potentials between water in hydrate (H) or ice (I) and water in the reference phase (β), respectively.

Modeling of $\Delta\mu_w^{H-\beta}$

$\Delta\mu_w^{H-\beta}$ is then determined from statistical thermodynamics whereas $\Delta\mu_w^{I-\beta}$ is determined by means of relations from classical thermodynamics that can be expressed as a function of the fugacity f_j of the gas j as:

$$\Delta\mu_w^{H-\beta} = RT \sum_i \nu_i \ln \left(1 - \sum_j \theta_j^i \right) = RT \sum_i \nu_i \ln \left(1 - \sum_j C_j^i f_j(T,P) \right) \quad (2)$$

where v_i is the number of cavities of type i , C_j^i is the Langmuir constant of component j in the cavity i describing the interaction potential between the encaged guest molecule and the surrounding water molecules, evaluated by assuming a spherically symmetrical cage and associated potential:

$$C_j^i = \frac{4\pi}{kT} \int_0^\infty \exp\left(-\frac{w(r)}{kT}\right) r^2 dr, \quad (3)$$

where w is the radial interaction potential between the guest molecule and the water molecules forming the structure. The interaction potential, when described by a Kihara model (Mckoy and Sinagolu, 1963), can be expressed as:

$$w(r) = 2z\varepsilon \left[\frac{\sigma^{12}}{\bar{R}^{11}r} \left(\delta^{10} + \frac{a}{\bar{R}} \delta^{11} \right) - \frac{\sigma^6}{\bar{R}^5 r} \left(\delta^4 + \frac{a}{\bar{R}} \delta^5 \right) \right] \quad (4)$$

$$\delta^N = \frac{1}{N} \left[\left(1 - r/\bar{R} - a/\bar{R} \right)^{-N} - \left(1 + r/\bar{R} - a/\bar{R} \right)^{-N} \right] \quad (5)$$

The gas parameters ε , σ and a are the Kihara parameters describing the potential we seek for and can be calculated from experimental data by fitting the model equations to corresponding hydrate equilibrium experimental data. The interaction potential takes into account the geometrical properties of the cavities (coordination number z and average mean radius \bar{R} , given in Table A1).

The equations that can be used to evaluate $\Delta\mu_w^{i-\beta}$, the difference in chemical potential between ice and the reference state $T_0 = 273.15$ K and the pressure $P_0 = 1$ bar are summarized in chapter 5 and in particular in Table 5.9 of Sloan and Koh (2008) and Table 4.3.1 of Herri et al (2011). The corresponding values of the reference state parameters used in the model are given in Table A2.

Table A2

Equilibrium :

A simultaneous minimization of the three Kihara parameters (σ , core radius and ϵ/k) on the clathrate hydrate model faces a high level of degeneracy, an issue amplified by the paucity of available equilibrium data in the sulfur dioxide case. In a rather well defined and measured system the degeneracy remains an issue (Herri et al., 2011). This is somehow already reflected in the distinct Kihara parameters deduced in the literature. Kihara parameters for pure substances can be evaluated from measurement of the viscosity, or can be calculated from the second virial coefficient, or from a combination of measurements (Tee et al, 1996). Also, they can be evaluated from Henry constants (Uno et al, 1975). The principle is to retrieve the Kihara parameters over experimental data with a limited number of pure substances. The authors propose a correlation to estimate the Kihara parameters for all kinds of gases. The optimization strategy we adopt to reduce the degeneracy is to set one of the parameters before to deduce the corresponding two others by minimization, as suggested in e.g. Mehta and Sloan (1996). They propose to fix the Kihara a value. Based on the parameterization by Tee et al (1966) and Uno et al (1975) for SO_2 , the core radius a value is estimated to lie close to $a=0.75 \text{ \AA}$, that we further set to explain the analysis performed.

Then, for a given set of Kihara parameters ϵ_j and σ_j , and a given temperature (resp. a given pressure), the calculated equilibrium pressure P_{calc} (resp. the calculated equilibrium temperature T_{calc}) corresponds to the value such as $\Delta\mu_w^{1-\beta} = \Delta\mu_w^{1-\beta}$. Then the calculated pressure and temperature are compared to the experimental ones P_{exp} and T_{exp} , and a deviation function can be defined as:

$$F(\epsilon_j, \sigma_j) = \sum_{l=1}^N \left| \frac{P_{calc}}{P_{exp}} - 1 \right| \rightarrow \min \text{ (resp. } F(\epsilon_j, \sigma_j) = \sum_{l=1}^N \left| \frac{T_{calc}}{T_{exp}} - 1 \right| \rightarrow \min \text{)} \quad (6)$$

The summation is performed over all N data of the set.

In the work of Herri and Chassefière (2012), the Kihara parameters for CO₂ clathrate hydrate were retrieved with such a scheme, with $\varepsilon/k=178.21\text{K}$, $a=0.6805\text{\AA}$ and $\sigma=2.873\text{\AA}$. By using the same method, the present work aims at retrieving the Kihara parameters for SO₂ hydrate.

A.2 Retrieving the SO₂ Kihara parameters

We focus on the value of $a=0.75\text{ \AA}$, but the minimization was performed over a broader range of a values. Then, the ε and σ can be optimized from experimental results. Using the Tamman and Krige (1925) data, the minimized ε and σ values dependencies are shown in Fig. A1, following a valley of minima all satisfying the available experimental data. Using another core radius value would give rise to another satisfying correlation. With the limited data available, covering only the range of temperature from 261.15 K to 271.15 K, corresponding pressure and temperature SO₂ clathrate hydrate equilibria at lower temperatures are shown in Fig. A2 (in this case the core radius was explored over a larger range). Nevertheless, taking into account the regression uncertainties, significant constraints can be set in the Martian context as shown in the next section.

Fig. A1

Fig. A2

1129 **Captions for tables :**

1130

1131 Table A1: Structure of SI and SII gas Hydrates

1132

1133 Table A2: Reference state parameters

1134

1135

Captions for figures :

Figure 1: Evolution of the cumulated amount of volcanically released sulfur in terms of SO₂ pressure (left vertical scale) and thickness of the corresponding sulfate mineral GEL (right vertical scale), assuming full conversion of sulfur into sulfate minerals. The “C&G” curve (dotted-dashed line) is obtained by multiplying by 10 the curve proposed in CG09 from a photogeological analysis of surface morphology (Greeley and Schneid, 1991). The two other curves are derived from CO₂ evolution curves proposed by Grott et al. (2011), as explained in the main text. These curves give the amount of CO₂ released from mantle carbon and don’t include a possible contribution of superficial carbon (e.g. carbonates) to the volcanic CO₂.

Figure 2: Mean surface temperature as a function of the CO₂ pressure for SO₂ mixing ratios of 0, 1, 10 and 100 ppm show the cooling effect of SO₂ aerosols at $p_{\text{CO}_2} = 3$ bar and mixing ratios of SO₂ of 1 ppm and 10 ppm. The equilibrium temperatures of CO₂ clathrate hydrate, and CO₂ ice, are plotted as a function of CO₂ pressure (from Mousis et al., 2012).

Figure 3: Ratio of concentration in mixed SO₂-CO₂ clathrate hydrate with respect to gas phase one as a function of temperature using the different constrained extrapolated SO₂ Kihara parameters. Initial SO₂ concentrations are 0.1 ppm (circles, blue curves) and 1000 ppm (triangles, red curves), covering most of the expected range for Martian conditions.

Figure 4: Surface temperature as a function the SO₂ mixing ratio for different values of the CO₂ partial pressure : 3 bar, 1.5 bar, 1 bar, 0.5 bar. The black triangles correspond to the SO₂ mixing ratio for which the temperature is equal to the CO₂ clathrate equilibrium temperature.

Figure 5: Equilibrium SO₂ mixing ratio as a function of CO₂ pressure. The lava volumes required to give rise to the corresponding SO₂ mixing ratio, calculated by using the same conversion factor as in Johnson et al. (2008), are indicated on the figure.

Figure 6: CO₂ evolution profile scaled on the Chassefière and Leblanc (2011a) profile to fit a 2 bar value at 3.8 Gyr b.p. (solid line). Blue rectangles with dotted lines and blue “average” dotted curve represent the evolution of the crust production rate derived from photogeological analysis (Greeley and Schneid, 1991). The chronology of clays, sulfates and ferric oxides proposed by Bibring et al. (2006) is shown at the top. The vertical grey band corresponds to the time interval when volcanic activity was significant according to Greeley and Schneid estimates.

Figure 7: Schematic representation of sulfur and CO₂ reservoir during the Noachian and Hesperian period. The total volatile surface/atmosphere CO₂ reservoir is decreasing due to thermal escape and possible formation of carbonates in the subsurface, from an initial budget ~100 bar. After magma ocean crystallization, CO₂ and water condense in form of clathrate (15 bar of CO₂ at maximum can be stored in form of clathrate when reacting with the total Martian water content of 1000 m GEL). After volcanic release of S, atmospheric SO₂ is systematically enriched in the condensating clathrate, triggered either by climatic changes or by global cooling due to aerosol formation. The CO₂ clathrate acts as an atmospheric buffer when the atmospheric pressure reaches 2 bar. After the complete consumption of CO₂ clathrate, the atmospheric pressure can drop below 2 bar and massive release of SO₂ from the clathrate induces the condensation of aerosols and precipitation as a Mars-wide sulfate layer (50 m represent the total SO₂ outgassed from Tharsis formation). This scenario represents the equilibrium state of surface/atmosphere but significant departure may have happened due to large impacts.

1187

1188 Figure 8: Mole fraction of volatiles encaged in clathrates calculated as a function of the
1189 surface pressure of CO₂ in the cases of, from top to bottom, SO₂ atmospheric abundances
1190 equal to 10⁻³, 1 and 10³ ppm.

1191

1192 Figure A1: Pearson's chi square test contour plot of the minimized ε/k versus σ for SO₂
1193 Kihara parameters, with core radius a fixed at 0.75Å. Contour levels are spaced in powers of
1194 3 of the best-regressed value. Pressure and temperature equilibrium data for SO₂ hydrate are
1195 taken from Tamman and Krige (1925)

1196

1197 Figure A2: Range of pressure and temperature extrapolated equilibrium for SO₂ clathrate
1198 hydrate using our model (small dots, colored range). The input Kihara parameters are
1199 obtained in the 3D $\{ \varepsilon/k, \sigma, a \}$ parameter space regression on experimental data by setting
1200 the core radius in the $a \in [0.55, 0.95]$ range, then regressing the degenerate Kihara parameters
1201 as shown in Fig. 3. Original data from Tamman and Krige (1925) (large circle) are displayed
1202 as well as water phase diagram limits.

1203

1204




Figure 1




Figure 2




Figure 3






Figure 4




Figure



Figure

6



Figure

7






Figure 8

$a=0.75\text{\AA}$



Figure

A1



Figure

A2