



HAL
open science

About the transcendentalty of some numbers

Jamel Ghannouchi

► **To cite this version:**

| Jamel Ghannouchi. About the transcendentalty of some numbers. 2013. hal-00808463v6

HAL Id: hal-00808463

<https://hal.science/hal-00808463v6>

Preprint submitted on 11 Apr 2013 (v6), last revised 8 May 2013 (v11)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

About the transcendentality of some numbers

Jamel Ghanouchi

Ecole Supérieure des Sciences et Techniques de Tunis

jamel.ghanouchi@topnet.tn

Abstract

Everyone knows that π , e or C_n , the Champernowne number are transcendental, but what about $\pi + e$, $\pi + C_n$ or $e + C_n$? In this paper, we show a method in the goal to know if they are.

The approach

A number is transcendental if it is not the root of a polynomial equation

$$a_n x^n + a_{n-1} x^{n-1} + \dots + a_0 = 0$$

where a_i are rational not all equal to zero, else it is algebraic. We know that π , e and C_n , the Champernowne number are transcendental, but we do not know anything about $e + \pi$, $\pi + C_n$ or $e + C_n$. Effectively, if A is transcendental : B transcendental and we do not know the nature of $A + B$ or $A - B$. But if B is algebraic, then $A + B$ and $A - B$ are transcendental. And if A and B are algebraics, their sum and their difference are algebraics. Let us try to solve this problem. Let

$$C_1 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 4 & 5 & 1 & 2 & 3 \\ 2 & 3 & 4 & 5 & 1 \\ 5 & 1 & 2 & 3 & 4 \\ 3 & 4 & 5 & 1 & 2 \end{pmatrix}$$

And let

$$C'_1 = \begin{pmatrix} 0 & 5 & 10 & 15 & 20 \\ 10 & 15 & 20 & 0 & 5 \\ 20 & 0 & 5 & 10 & 15 \\ 5 & 10 & 15 & 20 & 0 \\ 15 & 20 & 0 & 5 & 10 \end{pmatrix}$$

Thus $C_1 + C'_1$ is a magic square. It contains all the numbers from 1 to 25.

Let

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix}$$

And

$$C_2 = \begin{pmatrix} C_1 & C_1 + 25A & C_1 + 50A & C_1 + 75A & C_1 + 100A \\ C_1 + 75A & C_1 + 100A & C_1 & C_1 + 25A & C_1 + 50A \\ C_1 + 25A & C_1 + 50A & C_1 + 75A & C_1 + 100A & C_1 \\ C_1 + 100A & C_1 & C_1 + 25A & C_1 + 50A & C_1 + 75A \\ C_1 + 50A & C_1 + 75A & C_1 + 100A & C_1 & C_1 + 25A \end{pmatrix}$$

And

$$C'_2 = \begin{pmatrix} C'_1 & C'_1 & C'_1 & C'_1 & C'_1 \\ C'_1 & C'_1 & C'_1 & C'_1 & C'_1 \\ C'_1 & C'_1 & C'_1 & C'_1 & C'_1 \\ C'_1 & C'_1 & C'_1 & C'_1 & C'_1 \\ C'_1 & C'_1 & C'_1 & C'_1 & C'_1 \end{pmatrix}$$

The recurrence is

$$C_{i+1} = \begin{pmatrix} C_i & C_i + 5^i A & C_i + 2.5^i A & C_i + 3.5^i A & C_i + 4.5^i A \\ C_1 + 3.5^i A & C_1 + 4.5^i A & C_1 & C_1 + 5^i A & C_1 + 2.5^i A \\ C_1 + 5^i A & C_1 + 2.5^i A & C_1 + 3.5^i A & C_1 + 4.5^i A & C_1 \\ C_1 + 4.5^i A & C_1 & C_1 + 5^i A & C_1 + 2.5^i A & C_1 + 3.5^i A \\ C_1 + 2.5^i A & C_1 + 3.5^i A & C_1 + 4.5^i A & C_1 & C_1 + 5^i A \end{pmatrix}$$

And

$$C'_{i+1} = \begin{pmatrix} C'_i & C'_i & C'_i & C'_i & C'_i \\ C'_i & C'_i & C'_i & C'_i & C'_i \\ C'_i & C'_i & C'_i & C'_i & C'_i \\ C'_i & C'_i & C'_i & C'_i & C'_i \\ C'_i & C'_i & C'_i & C'_i & C'_i \end{pmatrix}$$

And $C_i + C'_i$ is a magic square containing all the numbers from 1 to

$$\begin{aligned} 5 + 20(5 + 5^2 + \dots + 5^{i-1}) + 20 &= 5 + 100(1 + 5 + 5^2 + \dots + 5^{i-2}) + 20 \\ &= 5 + 100\left(\frac{5^{i-1} - 1}{4}\right) + 20 = 5 + 25(5^{i-1} - 1) + 20 = 5^{i+1} \end{aligned}$$

The sum of C_{i+1} is $S_{i+1} = 5S_i + 10.5^i + 5S'_i$ ($S_1 = 65$, $S'_i = 50.5^{i-1}$). Hence

$$S_{i+1} = 5S_i + 60.5^i$$

$$5S_i = 5^2 S_{i-1} + 60.5^i$$

$$\begin{aligned}
& \dots \\
& 5^{i-1}S_2 = 5^i S_1 + 60.5^i \\
& 5^i S_1 = 5^i(65) = 13.5^{i+1} \\
& S_{i+1} = (60i + 65)5^i = (12i + 13)5^{i+1}
\end{aligned}$$

If M is the greatest number of the square, S the sum, the value of the integer $\pi.10^{p_i}$ that approaches π in the square (example between 1 and 25, $\pi.10^{p_i} = 3$ and between 1 and 125, it is 31). The number $\pi(p_i)$ depends of M and S . This square contains p_i first digit of π without the dot, we note it : $\pi.10^{p_i}$. It contains also $e.10^{p_i}$ and $C_n.10^{p_i}$ (and also $\pi + e$, πe , $\pi + C_n$, $\pi C_n \dots$). For $i + 1$, the greatest number in the square is $M = 5^{i+2}$ which has $E(\frac{(i+2)\log 5}{\log 10}) = i + 1$ ciphers and begins by 1, 2, 3 and then $p_i = i$, or by a number geater and then $p_i = i + 1$. The sum is $S_{i+1} = (12i + 13)5^{i+1}$ which has less than $p_i + 2$ ciphers. The sum $S_i.10^{-p_i}$ tends, in the infinity, to a number S which has less than three ciphers! But S_i is rational, thus S is rational and algebraic.

Main results

$$\left\{ \begin{array}{l}
S = 20 \\
A = S - \pi \\
B = S - e \\
C = S - C_n \\
A_1 = S - \pi - e \\
A_2 = S - \pi + e \\
A_3 = S - \pi e \\
A_4 = S - \pi^2
\end{array} \right.$$

We know that

$$S = \pi + A = e + B = \pi + e + A_1 = \pi - e + A_2 = \pi e + A_3$$

Definition A real number is composed if it is equal to $\pm p_1^{n_1} \dots p_i^{n_i}$ where p_j are integer prime numbers and n_j are rationals. We define other real prime numbers which can not be expressed like this : π , e , $\ln(2)$. Thus $\sqrt[q]{p} = p^{\frac{1}{q}}$ is composed.

Also $\sqrt[p]{p} + 1$ is prime, with p prime, hence $\sqrt[p]{p} - 1 = (p - 1)(\sqrt[p]{p} + 1)^{-1}$ is composed!

And π and e are primes in stead of π^{n_0} and e^{m_0} with $(n_0 - 1)(m_0 - 1) \neq 0$. We define the GCD of two numbers : If p_1 and p_2 are prime real numbers

$$p_1 \neq p_2 \Rightarrow GCD(p_1, p_2) = 1$$

$$n_1 n_2 < 0 \Rightarrow GCD(p_1^{n_1}, p_1^{n_2}) = 1$$

$$n_1 n_2 > 0 \Rightarrow GCD(p_1^{n_1}, p_1^{n_2}) = p_1^{\max(n_1, n_2) - \min(n_1, n_2)}$$

$$GCD(p_1^{n_1} p_2^{n_2} \dots p_i^{n_i}, p_1^{m'_1} p_2^{m'_2} \dots p_j^{m'_j}) = \prod_{i \neq j} GCD(p_i^{n_i}, p_j^{m'_j})$$

And if $x = p_1^{n_1} p_2^{n_2} \dots p_i^{n_i}$ and $y = p_{m_1}^{n_{m_1}} \dots p_{m_i}^{n_{m_i}}$ then y divides x if $1 < m_l < i$ for $1 \leq l \leq i$ and for $m_i = j$, $|n_{m_i}| < |j|$.

Theorem If T_1 and T_2 are transcendental prime real numbers, $T_1 T_2$ and $T_1 + T_2$ are transcendentals.

Proof of the theorem Let $T_1 = S - A$ and $T_2 = S - B$ (S algebraic) two real prime transcendental numbers. We have

$$nT_i^2 - mT_i + (-nT_i + m)T_i = 0$$

n is algebraic, $i \in \{1, 2\}$, as T_i is transcendental, it can not be solution of a polynomial equation, thus $(-nT_i + m)T_i$ is transcendental $\forall m, n$ algebraic. We have possibilities, for all (m, n)

1)

$T_1 T_2^n = T_A$ and $T_1 + mT_2^m$ are algebraics $\forall m, n$

2)

$T_1 T_2^n$ is algebraic and $T_1 + mT_2^m$ is transcendental, $\forall m, n$

3)

$T_1 T_2^n$ is transcendental and $T_1 + mT_2^m$ is algebraic, $\forall m, n$

4)

$T_1 T_2^n$ and $T_1 + mT_2^m$ are transcendentals, $\forall m, n$

Thus 1)

$m = n$ implies that $nT_1 T_2^n$ and $T_1 + nT_2^n$ are algebraics, but

$$T_1^2 - (T_1 + nT_2^n)T_2 + nT_1 T_2^n = 0$$

Is impossible means that they can not be algebraics simultaneously!

Else $m \neq n$ then $T_1 T_2^n = A_A$ and $T_1 + mT_2^m = A'_A$ are algebraics, then

$$T_1 T_2^n + mT_2^{m+n} = A_A + mT_2^{n+m} = A'_A T_2^n$$

And T_2 is solution of this algebraic equation : it impossible!

2)

$T_1 T_2^n - n = (T_1 - nT_2^{-n})T_2^n = T_A T_2^n = A_A$ is algebraic and $T_1 - nT_2^{-n} = T_A$ is transcendental.

And $T_1 T_2^{-n} + n = (T_1 + nT_2^n)T_2^{-n} = T'_A T_2^{-n} = A'_A$ is algebraic, with $T_1 + nT_2^n = T'_A$ transcendental

$T_A T'_A = A_A A'_A = (T_1 + nT_2^n)(T_1 - nT_2^{-n}) = T_1^2 - n^2 + nT_1 T_2^n - nT_1 T_2^{-n}$ means that T_1^2 is algebraic and it is impossible!

3)

$T_1T_2^n + n = (T_1 - nT_2^{-n})T_2^n = A_A T_2^n = T_A$ is transcendental and $T_1 - nT_2^{-n}$ is algebraic

$$2T_1T_2^{-n} + n = (2T_1 + nT_2^n)T_2^{-n} = A'_A T_2^{-n} = T'_A$$

$$\begin{aligned} A_A A'_A &= T_A T'_A = (T_1 T_2^n + n)(2T_1 T_2^{-n} + n) \\ &= 2T_1^2 + n^2 + nT_1(T_2 + T_2^{-n}) = n^2 + T_1(T_1 + nT_2^{-n} + T_1 + nT_2^n) \end{aligned}$$

And $n^2 + T_1 A''_A = A_A \cdot A'_A$

Where A''_A, A'_A, A_A are algebraics and it is impossible !

Thus $T_1 T_2^n$ and $T_1 + m T_2^m$ are transcendentals for all m, n .

Thus $T_1 T_2, T_1 T_2^{-1}, T_2 T_1^{-1}, T_1 - T_2$ and $T_1 + T_2$ are transcendentals ! In our hypothesis T_1 and T_2 are primes, we can not have $T_1 = u T_2^{-1}$, we can not have $T_1 + T_2 = A_A$ an algebraic number, as $T_1 T_2^m = T_2^m (A_A - T_2)$ is never algebraic !

Application of the theorem $T_1 = \pi$ and $T_2 = e$ are transcendental prime real numbers. We have $T_1 T_2 = e\pi$ and $T_1 + T_2 = e + \pi$, are transcendentals for example !

If π^m and e^n are the prime numbers, then $\pi^m e^n$ and $\pi^m + e^n$ are transcendental for all m, n .

Also, by the same way, we prove that $\pi^n, e^m, C_n^r, \pi^n e^m, C_n^r \pi^n$ and $C_n^r e^m$ are transcendentals for all n, m, r !

Conclusion

We did not presented like this, but we have given a method to find the nature of several numbers, we have shown the nature of $\pi^n e^m, \pi^n C_n^p, e^m C_n^p, \pi^n, e^m$ and $C_n^p \dots$

Références

- [1] Alan Baker, Transcendental number theory *Cambridge University Press*, (1975).