



**HAL**  
open science

## Complete Genome Sequence of the Piezophilic, Mesophilic, Sulfate-Reducing Bacterium *Desulfovibrio* *hydrothermalis* AM13(T.).

Boyang Ji, Gregory Gimenez, Valérie Barbe, Benoit Vacherie, Zoé Rouy,  
Amira Amrani, Marie-Laure Fardeau, Philippe Bertin, Didier Alazard, Sabine  
Leroy, et al.

### ► To cite this version:

Boyang Ji, Gregory Gimenez, Valérie Barbe, Benoit Vacherie, Zoé Rouy, et al.. Complete Genome Sequence of the Piezophilic, Mesophilic, Sulfate-Reducing Bacterium *Desulfovibrio hydrothermalis* AM13(T.). *Genome Announc*, 2013, 1 (1), epub ahead of print. 10.1128/genomeA.00226-12 . hal-00809071

**HAL Id: hal-00809071**

**<https://hal.science/hal-00809071>**

Submitted on 8 Apr 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Complete Genome Sequence of the Piezophilic, Mesophilic, Sulfate-Reducing Bacterium *Desulfovibrio hydrothermalis* AM13<sup>T</sup>

Boyang Ji,<sup>a</sup> Gregory Gimenez,<sup>b</sup> Valérie Barbe,<sup>c</sup> Benoît Vacherie,<sup>c</sup> Zoé Rouy,<sup>c</sup> Amira Amrani,<sup>d</sup> Marie-Laure Fardeau,<sup>d</sup> Philippe Bertin,<sup>e</sup> Didier Alazard,<sup>d</sup> Sabine Leroy,<sup>f</sup> Emmanuel Talla,<sup>a</sup> Bernard Ollivier,<sup>d</sup> Alain Dolla,<sup>a</sup> Nathalie Pradel<sup>d</sup>

Aix-Marseille Université, CNRS, Laboratoire de Chimie Bactérienne (LCB), UMR 7283, Marseille, France<sup>a</sup>; UMR 6236, Faculté de Médecine, Marseille, France<sup>b</sup>; CEA, Institut de Génétique/Genoscope, Laboratoire de Finition, Évry, France<sup>c</sup>; Aix-Marseille Université, Université du Sud Toulon-Var, CNRS, Institut National des Sciences de l'Univers (INSU), Institut de Recherche pour le Développement (IRD), Mediterranean Institute of Oceanography (MIO), UM110, Marseille, France<sup>d</sup>; UMR 7156, CNRS, Université Louis Pasteur, Strasbourg, France<sup>e</sup>; INRA, UR454 Microbiologie, Saint-Genès Champanelle, France<sup>f</sup>

***Desulfovibrio hydrothermalis* AM13<sup>T</sup> is a piezophilic, mesophilic, hydrogenotrophic sulfate-reducing bacterium collected from a deep-sea hydrothermal chimney on the East Pacific Rise (2,600 m depth, 13°N). We report the genome sequence of this bacterium, which includes a 3,702,934-bp chromosome and a circular plasmid of 5,328 bp.**

Received 18 December 2012 Accepted 31 December 2012 Published 21 February 2013

Citation Ji B, Gimenez G, Barbe V, Vacherie B, Rouy Z, Amrani A, Fardeau M-L, Bertin P, Alazard D, Leroy S, Talla E, Ollivier B, Dolla A, Pradel N. 2013. Complete genome sequence of the piezophilic, mesophilic, sulfate-reducing bacterium *Desulfovibrio hydrothermalis* AM13<sup>T</sup>. *Genome Announc.* 1(1):e00226-12. doi:10.1128/genomeA.00226-12.

Copyright © 2013 Ji et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Nathalie Pradel, [nathalie.pradel@ird.fr](mailto:nathalie.pradel@ird.fr).

Deep-sea sulfate-reducing microorganisms must be considered as key actors in Earth's sulfur cycles. Metagenomic studies have revealed that sulfate-reducing bacteria (SRB) are largely represented among the microorganisms that inhabit deep-sea environments (1–3). However, only a few SRB have been isolated from deep-sea environments (4–9). Among them, the *Desulfovibrio* genus is well-represented, with *Desulfovibrio profundus*, *Desulfovibrio piezophilus*, and *Desulfovibrio hydrothermalis* having been isolated previously (7–9). Here, we report the genome sequence of *D. hydrothermalis*, a piezo-mesophilic, hydrogenotrophic SRB isolated from a deep-sea hydrothermal chimney sample collected at a depth of 2,600 m (13°N on the East Pacific Rise) (7). It represents, with *D. piezophilus* (10), the second known genome of piezophilic bacteria in the genus *Desulfovibrio*, and thus gives the opportunity to specify the systems that are involved in the piezophilic lifestyles of *Desulfovibrio* species.

Genome sequencing was performed by the Genoscope (CEA, Évry, France), using 454-sequencing method (Roche) and Solexa technology. A mate-paired genomic library with 8-kb insert size was prepared according to the manufacturer's protocols (454 Life Sciences Corporation, Branford, CT). Approximately 19-fold coverage of GS FLX Titanium (<http://www.rocke.com/>) reads was assembled using Newbler. Gaps between contigs were closed with 1,038 PCR amplifications and the Consed editing software (<http://www.phrap.org/>). For the quality assessment, approximately 1,000-fold coverage of paired-end Illumina reads (36 bp) was mapped onto the whole-genome sequence.

The genome consists of two contigs of 284,269 bp and 3,418,665 bp, for a total of 3,702,934 bp, and of a circular plasmid of 5,328 bp. The average G+C contents for the chromosome and the plasmid DNA are 45.1% and 51.1%, respectively. A total of 3,543 coding DNA sequences (CDSs) were predicted, as well as 7 pseudogenes, 6 miscellaneous RNAs (misc-RNA), 5 disrupted rRNA operons, and 77 tRNA genes.

Phylogenetic analysis based on 16S rRNA shows that the clos-

est sequenced genomes to *D. hydrothermalis* are those of *Desulfovibrio salexigens* and *Desulfovibrio aespoensis*. Reciprocal best BLAST analysis indicated that *D. hydrothermalis* shares 2,824 and 2,069 orthologous proteins with *D. salexigens* and *D. aespoensis*, respectively, and 1,974 proteins are common to the three species.

Like other *Desulfovibrio* species, *D. hydrothermalis* possesses all the genes necessary for the sulfate-reduction pathway. Fatty acid membrane composition has been shown to play an important role in hydrostatic pressure adaptation (11). In *D. hydrothermalis*, all fatty acid biosynthesis pathways are present. Interestingly, three copies of the *fabF* gene (DESAMv2\_20244, DESAMv2\_21138, and DESAMv2\_21153), encoding a  $\beta$ -ketoacyl-acyl carrier protein (ACP) synthase II involved in monounsaturated fatty acid production and suggested to be involved in hydrostatic pressure adaptation (11), are found in the genome. Thus, the three copies of this gene might play a pivotal role in the adaptation strategy of *D. hydrothermalis* to hydrostatic pressure.

Comparative genomics, including the use of other *Desulfovibrio* species, will give insights into the adaptation of *Desulfovibrio* species to the piezophilic lifestyle that they encounter in the deep-sea biosphere.

**Nucleotide sequence accession numbers.** The final annotated genome of *D. hydrothermalis* is available in EMBL under the accession no. FO203522 (for the chromosome) and FO203523 (for the plasmid).

## ACKNOWLEDGMENTS

This work was supported by CNRS and IRD grants.

B.J. is a PhD student with a China Scholarship Council fellowship.

## REFERENCES

- Hu Y, Fu C, Yin Y, Cheng G, Lei F, Yang X, Li J, Ashforth EJ, Zhang L, Zhu B. 2010. Construction and preliminary analysis of a deep-sea sediment metagenomic fosmid library from Qiongdongnan basin, south China sea. *Mar. Biotechnol.* 12:719–727.
- Martin-Cuadrado AB, López-García P, Alba JC, Moreira D, Monticelli

- L, Strittmatter A, Gottschalk G, Rodríguez-Valera F. 2007. Metagenomics of the deep Mediterranean, a warm bathypelagic habitat. *PLoS One* 2:e914.
3. Stokke R, Roalkvam I, Lanzen A, Haflidason H, Steen IH. 2012. Integrated metagenomic and metaproteomic analyses of an ANME-1-dominated community in marine cold seep sediments. *Environ. Microbiol.* 14:1333–1346.
  4. Alain K, Postec A, Grinsard E, Lesongeur F, Prieur D, Godfroy A. 2010. *Thermodesulfatator atlanticus* sp. nov., a thermophilic, chemolithoautotrophic, sulfate-reducing bacterium isolated from a mid-Atlantic ridge hydrothermal vent. *Int. J. Syst. Evol. Microbiol.* 60:33–38.
  5. Audiffren C, Cayol JL, Joulain C, Casalot L, Thomas P, Garcia JL, Ollivier B. 2003. *Desulfonauticus submarinus* gen. nov., sp. nov., a novel sulfate-reducing bacterium isolated from a deep-sea hydrothermal vent. *Int. J. Syst. Evol. Microbiol.* 53:1585–1590.
  6. Jeanthon C, L'Haridon S, Cueff V, Banta A, Reysenbach AL, Prieur D. 2002. *Thermodesulfobacterium hydrogeniphilum* sp. nov., a thermophilic, chemolithoautotrophic, sulfate-reducing bacterium isolated from a deep-sea hydrothermal vent at Guaymas basin, and emendation of the genus *Thermodesulfobacterium*. *Int. J. Syst. Evol. Microbiol.* 52:765–772.
  7. Alazard D, Dukan S, Urios A, Verhé F, Bouabida N, Morel F, Thomas P, Garcia JL, Ollivier B. 2003. *Desulfovibrio hydrothermalis* sp. nov., a novel sulfate-reducing bacterium isolated from hydrothermal vents. *Int. J. Syst. Evol. Microbiol.* 53:173–178.
  8. Bale SJ, Goodman K, Rochelle PA, Marchesi JR, Fry JC, Weightman AJ, Parkes RJ. 1997. *Desulfovibrio profundus* sp. nov., a novel barophilic sulphate-reducing bacterium from deep sediment layers in the Japan Sea. *Int. J. Syst. Bacteriol.* 47:515–521.
  9. Khelaifa S, Fardeau ML, Pradel N, Aussignargues C, Garel M, Tamburini C, Cayol JL, Gaudron S, Gaill F, Ollivier B. 2011. *Desulfovibrio piezophilus*, sp. nov., a novel piezophilic sulfate-reducing bacterium isolated from wood falls in Mediterranean Sea. *Int. J. Syst. Evol. Microbiol.* 61:2706–2711.
  10. Pradel N, Ji BY, Gimenez G, Talla E, Lenoble P, Garel M, Tamburini C, Fourquet P, Lebrun R, Bertin P, Denis Y, Pophillat M, Barbe V, Ollivier B, Dolla A. The first genomic and proteomic characterization of a deep-sea sulfate reducer: insights into the piezophilic lifestyle of *Desulfovibrio piezophilus*. *PLoS One*, in press.
  11. Bartlett DH. 2002. Pressure effects on *in vivo* microbial processes. *Biochim. Biophys. Acta* 1595:367–381.