

Preperiodic dynatomic curves for $z \mapsto z^d + c$

Gao Yan

July 19, 2015

Abstract

The preperiodic dynatomic curve $\mathcal{X}_{n,p}$ is the closure in \mathbb{C}^2 of the set consisting of (c, z) such that z is a preperiodic point of the polynomial $z \mapsto z^d + c$ with preperiod n and period p ($n, p \geq 1$). We prove that each $\mathcal{X}_{n,p}$ has exactly $d - 1$ irreducible components, these components are all smooth and have pairwise transverse intersections at the singular points of $\mathcal{X}_{n,p}$. We also compute the genus of each component and the Galois group of the defining polynomial of $\mathcal{X}_{n,p}$.

Keywords and phrases: dynatomic curves, smooth, irreducible, Multibrot set, Galois group, complex dynamics.

AMS(2010) Subject Classification: 14H50, 37F45.

1 Introduction

Fix $d \geq 2$. For $c \in \mathbb{C}$, set $f_c(z) = z^d + c$. For $p \geq 1$, define

$$\check{\mathcal{X}}_{0,p} := \{(c, z) \in \mathbb{C}^2 \mid f_c^p(z) = z \text{ and for all } 0 < k < p, f_c^k(z) \neq z\}.$$

$$\mathcal{X}_{0,p} := \text{the closure of } \check{\mathcal{X}}_{0,p} \text{ in } \mathbb{C}^2.$$

It is known that all $\mathcal{X}_{0,p}$ are affine algebraic curves, called the *periodic dynatomic curves*. These curves have been the subject of several studies in algebraic and holomorphic dynamical systems. The known results for these curves mainly include the smoothness (Douady-Hubbard [DH1], Milnor [Mil1], Buff-Tan [BT]); irreducibility (Bousch [B], Buff-Tan [BT], Morton [Mo], Lau-Schleicher [LS], Schleicher [S]); the genus (Bousch [B]) and the associated Galois groups (Bousch [B], Morton [Mo], Lau-Schleicher [LS], Schleicher [S]).

In the present work, we study some topological and algebraic properties of *preperiodic dynatomic curves*.

Definition 1.1. For $n \geq 0$, $p \geq 1$, a point z is called a *p-periodic point* if $f_c^p(z) = z$ but $f_c^k(z) \neq z$ for $0 < k < p$, and an *(n, p)-preperiodic point* of f_c if $f_c^n(z)$ is a *p-periodic point* of f_c but $f_c^l(z)$ is not periodic for any $0 \leq l < n$.

Now, for any $n \geq 1, p \geq 1$, define

$$\begin{aligned}\check{\mathcal{X}}_{n,p} &= \{(c, z) \in \mathbb{C}^2 \mid z \text{ is a } (n, p)\text{-preperiodic point of } f_c\} \\ \mathcal{X}_{n,p} &:= \text{the closure of } \check{\mathcal{X}}_{n,p} \text{ in } \mathbb{C}^2.\end{aligned}$$

In fact, as we shall see below, all $\mathcal{X}_{n,p}$ are also affine algebraic curves, called the *preperiodic dynatomic curves*. Limited work has been done for this kind of curves. The special case $d = 2$ has been previously studied by Bousch [B], who established in this case that for any integers $n, p \geq 1$, the curve $\mathcal{X}_{n,p}$ is also smooth and irreducible (as the periodic dynatomic curves), and computed its associated Galois group.

The main purpose of this work is to extend these results to arbitrary $d \geq 2$. An obvious difference with the previous case is that, for $d > 2$, the curve $\mathcal{X}_{n,p}$ is no longer irreducible: it consists of $d - 1$ irreducible components. We may understand this by a simple observation. Consider the curve $\mathcal{X}_{1,p}$ of $(1, p)$ -preperiodic points, that is, the points z which are not p -periodic, but whose image $z_0 = f(z)$ is. The periodic point $z_{p-1} = f^{p-1}(z_0)$ is another preimage of z_0 . Because $f_c(z) = z^d + c$, we have $z = \omega z_{p-1}$, where ω is a d -th root of unity. According to the value of ω , we can partition the $(1, p)$ -preperiodic points into $d - 1$ classes, and this decomposition is of algebraic nature: it corresponds to a factorization of $f_c^{p+1}(z) - f_c(z)$.

We show that these $d - 1$ components are smooth and irreducible. Our approach to smoothness is by using elementary calculations on quadratic differentials and Thurston's contraction principle, following the method of Buff-Tan ([BT]). The approach to irreducibility is based on the connectedness of periodic dynatomic curves and then by an induction on the preperiodic index n . Moreover, we study the features of the singular points of $\mathcal{X}_{n,p}$.

Following Bousch, we compute the genus of each irreducible component and the associated Galois group of the curve $\mathcal{X}_{n,p}$.

Here is a list our main results. They are to be compared with results of periodic dynatomic curves.

Denote by $\{\nu_d(p)\}_{p \geq 1}$ the unique sequence of positive integers satisfying the recursive relation

$$d^p = \sum_{k|p} \nu_d(k), \quad \text{integer } d \geq 2 \tag{1.1}$$

and let $\varphi(m)$ be the Euler totient function (i.e. the number of positive integers less than m and co-prime to m). For $n, p \geq 1$, define the numbers

$$M_{n,p} := \nu_d(p) d^{n-2} (d-1) \left(n-1 - \sum_{t=1}^{\lfloor \frac{n-1}{p} \rfloor} d^{-tp} \right),$$

where $\lfloor x \rfloor$ denotes the maximal integer less than or equal to x , and

$$K_{n,p} := \nu_d(p) (d^{p-1} - 1) d^{n-1-p} \left(\sum_{t=1}^{\lfloor \frac{n-1}{p} \rfloor - 1} d^{-t(p-1)} - \sum_{t=1}^{\lfloor \frac{n-1}{p} \rfloor - 1} d^{-pt} \right) + (d^{\lfloor \frac{n-1}{p} \rfloor} - 1) \nu_d(p) d^{n-2 - \lfloor \frac{n-1}{p} \rfloor p}$$

(one can refer to (5.3) and (5.4) for the computation of them). For $n, p \geq 1$, set

$$g_p(d) = 1 + \frac{dp - d - p - 1}{2d} \nu_d(p) - \frac{d-1}{2d} \sum_{k|p, k < p} \varphi\left(\frac{p}{k}\right) k \cdot \nu_d(k),$$

$$g_{n,p} = 1 + \frac{1}{2} \nu_d(p) d^{n-2} (pd - d - p - 1) + \frac{1}{2} (M_{n,p} + K_{n,p}) - \frac{1}{2} d^{n-2} (d-1) \sum_{k|p, k < p} \varphi\left(\frac{p}{k}\right) k \cdot \nu_d(k).$$

Theorem 1.2. *For any $d \geq 2$, $n, p \geq 1$, the preperiodic dynatomic curve $\mathcal{X}_{n,p}$ has the following properties :*

1. *The set $\mathcal{X}_{n,p}$ is an affine algebraic curve. It has $d-1$ irreducible components and each one is smooth. Moreover, these components are pairwise intersecting at the singular points of $\mathcal{X}_{n,p}$. In particular, if $d = 2$, the curve $\mathcal{X}_{n,p}$ is smooth and irreducible.*
2. *The genus of every irreducible component of $\mathcal{X}_{n,p}$ (in some kind of compactification) is $g_{n,p}(d)$, and all irreducible components are mutually homeomorphic.*
3. *The Galois group associated with $\mathcal{X}_{n,p}$ is the same as that associated with $\mathcal{X}_{\leq n,p} := \cup_{l=0}^n \mathcal{X}_{l,p}$, which consists of all permutations on the roots of the defining polynomial of $\mathcal{X}_{\leq n,p}$ that commute with f_c and the rotation of argument $1/d$.*

Here is a tableau comparing these various curves, where \mathbf{S}_m denotes the group of permutations on $\{1, \dots, m\}$ and $G_{n,p}(d)$ is the associated Galois group of $\mathcal{X}_{n,p}$.

periodic $\mathcal{X}_{0,p}$	$d = 2$	$d > 2$
	irreducible	irreducible
	smooth	smooth
genus	$g_p(2)$	$g_p(d)$
Galois group	$\mathbf{S}_{\nu_2(p)/p} \times \mathbb{Z}_p^{\nu_2(p)/p}$	$\mathbf{S}_{\nu_d(p)/p} \times \mathbb{Z}_p^{\nu_d(p)/p}$

preperiodic $\mathcal{X}_{n,p}$, $n \geq 1$	$d = 2$	$d > 2$
	irreducible	$d - 1$ irreducible components
	smooth	not smooth, but each component is smooth
component-wise genus	$g_{n,p}(2)$	$g_{n,p}(d)$
Galois group	$G_{n,p}(2)$	$G_{n,p}(d)$
pairwise intersection	empty	$C_{n,p}(\text{singular})$: singularity set of $\mathcal{X}_{n,p}$

This manuscript is organized as follows:

In section 2, we summarize some preliminaries that will be used in this paper.

In section 3, we will prove that every $\mathcal{X}_{n,p}$ is an affine algebraic curve and find its defining polynomial.

In section 4, we give the irreducible factorization of $\mathcal{X}_{n,p}$, and prove that each irreducible factor is smooth and these irreducible components are pairwise intersecting at the singular points of $\mathcal{X}_{n,p}$.

In section 5, we calculate the genus of each irreducible component.

In section 6, we describe $\mathcal{X}_{n,p}$ from the algebraic point of view by calculating its associated Galois group.

Acknowledgement. I thank Tan Lei for helpful discussions and suggestions.

2 Preliminaries

1. Filled in Julia set and Multibrot set. These material can be found in [DH1, DH2] and [Eb].

For $c \in \mathbb{C}$, we denote by K_c the filled-in Julia set of f_c , that is the set of points $z \in \mathbb{C}$ whose orbit under f_c is bounded. We denote by M_d the *Multibrot set* in the parameter plane, that is the set of parameters $c \in \mathbb{C}$ for which the critical point 0 belongs to K_c . It is known that M_d is connected.

Assume $c \in M_d$. Then K_c is connected. There is a conformal isomorphism $\phi_c : \mathbb{C} \setminus \overline{K}_c \rightarrow \mathbb{C} \setminus \overline{\mathbb{D}}$ satisfying $\phi_c \circ f_c = (\phi_c)^d$ and $\phi_c'(\infty) = 1$ (i.e. $\frac{\phi_c(z)}{z} \rightarrow_{z \rightarrow \infty} 1$). The dynamical ray of angle $\theta \in \mathbb{T}$ is defined by

$$R_c(\theta) := \{z \in \mathbb{C} \setminus K_c \mid \arg(\phi_c(z)) = 2\pi\theta\}.$$

Assume $c \notin M_d$. Then K_c is a Cantor set and all periodic points of f_c are *repelling*, that is $|(f^p)'(z)| > 1$ for $p \geq 1$ and all p -periodic point z . There is a conformal isomorphism $\phi_c : U_c \rightarrow V_c$ between neighborhoods of ∞ in \mathbb{C} , which satisfies $\phi_c \circ f_c = (\phi_c)^d$ on U_c . We may choose U_c so that U_c contains the critical value c and V_c is the complement of a closed disk. For each $\theta \in \mathbb{T}$, there is an infimum $r_c(\theta) \geq 1$ such that ϕ_c^{-1} extends analytically along $R_0(\theta) \cap \{z \in \mathbb{C} \mid r_c(\theta) < |z|\}$. We denote by ψ_c this extension and by $R_c(\theta)$ the dynamical ray

$$R_c(\theta) := \psi_c \left(R_0(\theta) \cap \{z \in \mathbb{C} \mid r_c(\theta) < |z|\} \right).$$

As $|z| \searrow r_c(\theta)$, the point $\psi_c(re^{2\pi i\theta})$ converges to a point $x \in \mathbb{C}$ ([DH2, Pro.8.3]). If $r_c(\theta) > 1$, then $x \in \mathbb{C} \setminus K_c$ is an iterated preimage of 0 and we say that $R_c(\theta)$ bifurcates at x . If $r_c(\theta) = 1$, then x belongs to K_c and we say that $R_c(\theta)$ lands at x .

There are three kinds of important parameters in M_d : super-attracting, parabolic, and Misiurewicz parameters. Recall that a point z is said to be p -periodic if $f_c^p(z) = z$ but $f_c^k(z) \neq z$ for $0 < k < p$. We call $c \in \mathbb{C}$

- a p -super-attracting parameter if 0 is p -periodic by f_c ;
- a p -parabolic parameter if f_c has a p -periodic point z_0 with $(f_c^p)'(z_0) = 1$ or a m -periodic point z_0 such that $m \mid p$ and $(f_c^m)'(z_0)$ is a $\frac{p}{m}$ -th root of unity;
- a (n, p) -Misiurewicz parameter if 0 is a (n, p) -preperiodic point of f_c .

A well-known result in complex dynamics says that any parabolic cycle of a rational map has a critical point in its basin, whose orbit eventually converges to , but is disjoint with the cycle (see [Mil2, Thm.10.15]). So for the family of unicritical polynomials $\{f_c \mid c \in \mathbb{C}\}$, the three classes of parameters above are pairwise disjoint. We write this point as a lemma, since it will be repeatedly used throughout the paper.

Lemma 2.1. *If the critical point 0 is (pre)periodic for f_c , then c is not a parabolic parameter.*

2. Affine algebraic curve and singularity. These material can be found in [G].

A polynomial $f \in \mathbb{C}[x, y]$ is called *squarefree* if it is not divisible by $h(x, y)^2$ for any non-constant $h(x, y) \in \mathbb{C}[x, y]$. An *affine algebraic curve over \mathbb{C}* is defined as

$$\mathcal{C} = \{(x, y) \in \mathbb{C}^2 \mid f(x, y) = 0\}$$

where f is a non-constant squarefree polynomial in $\mathbb{C}[x, y]$, called the *defining polynomial* of \mathcal{C} . If $f = \prod_{i=1}^m f_i$, where f_i are the irreducible factor of f , we say that the affine curve defined by f_i is a *irreducible component* of \mathcal{C} .

Let $f \in \mathbb{C}[x, y]$. The *total degree* of $f(x, y)$ as a multivariate polynomial is the highest degree of its terms, denoted by $\text{Deg}(f)$. Correspondingly, we denote by $\text{deg}_x(f)$ and $\text{deg}_y(f)$ the degrees of f when considered as a polynomial in the variable x and y respectively. The following lemma is repeatedly used in this paper.

Lemma 2.2. (1) *If $f = f_1 f_2$ with $f_1, f_2 \in \mathbb{C}[x, y]$, then $\text{Deg}(f) = \text{Deg}(f_1) + \text{Deg}(f_2)$, $\text{deg}_x(f) = \text{deg}_x(f_1) + \text{deg}_x(f_2)$ and $\text{deg}_y(f) = \text{deg}_y(f_1) + \text{deg}_y(f_2)$.*

(2) *For $f_1, f_2 \in \mathbb{C}[x, y]$, if $f(x, y) = f_1(x, f_2(x, y))$, then $\text{deg}_y(f) = \text{deg}_y(f_1) \cdot \text{deg}_y(f_2)$.*

(3) *For $f_1, f_2 \in \mathbb{C}[x, y]$, if $f(x, y) = f_1(x, f_2(x, y))$ and $\text{Deg}(f_1) = \text{deg}_y(f_1) \geq 1$, $\text{Deg}(f_2) > 1$, then $\text{Deg}(f) = \text{Deg}(f_1) \cdot \text{Deg}(f_2)$.*

Proof. (1). Refer to [F, section 1.1].

(2). It is straightforward by a simple computation.

(3). Set $d_1 := \text{Deg}(f_1)$ and $d_2 := \text{Deg}(f_2)$. According to the conditions of the lemma, we have $\deg_y(f_1) = d_1 \geq 1$ and $d_2 > 1$. On one hand, since $\text{Deg}(f_1) = \deg_y(f_1) = d_1$, then there is a unique term in f_1 with the form $a_1 y^{d_1}$, where a_1 is a non-zero constant. So, by (1) and $d_1 \geq 1$, it follows that $\text{Deg}(a_1 f_2^{d_1}) = d_1 d_2$. On the other hand, any other term of f_1 has the form $ax^s y^t$, where a is a non-zero constant and either $s + t < d_1$ or $s + t = d_1$ and $s \geq 1$. According to point (1) and $d_2 > 1$,

$$\text{Deg}(x^s f_2^t(x, y)) = s + td_2 < d_1 d_2.$$

So we get $\text{Deg}(f) = d_1 d_2$. □

Let \mathcal{C} be an affine algebraic curve for \mathbb{C} defined by $f \in \mathbb{C}[x, y]$, and let $P = (a, b) \in \mathcal{C}$. The multiplicity of \mathcal{C} at P , denoted by $\text{mult}_P(\mathcal{C})$, is defined as the order s of the first non-vanishing term in the Taylor expansion of f at P , i.e.

$$f(x, y) = \sum_{s=0}^{\infty} \frac{1}{s!} \sum_{t=0}^s \binom{s}{t} (x-a)^t (y-b)^{s-t} \frac{\partial^s f}{\partial x^t \partial y^{s-t}}(a, b).$$

If $\text{mult}_P(\mathcal{C}) = 1$, the point P is called a *smooth point* of \mathcal{C} . If $\text{mult}_P(\mathcal{C}) = r > 1$, then we say that P is a *singular point of multiplicity r* . We say that \mathcal{C} or f is *smooth* if any point on \mathcal{C} is smooth. Note that the first non-vanishing term is a homogeneous polynomial about $x - a$ and $y - b$, so all its irreducible factors are linear and they are called the *tangents* of \mathcal{C} at P .

A singular point P of multiplicity r on an affine plane curve \mathcal{C} is called *ordinary* if the r tangents to \mathcal{C} at P are distinct.

The following result provides a topological interpretation of the irreducibility of polynomials.

Lemma 2.3. *A squarefree polynomial $f \in \mathbb{C}[x, y]$ is irreducible if and only if the set of smooth points of f is connected.*

3. Periodic dynatomic curves.

In this paper, some of proofs and statements rely on the work of the periodic curves $\mathcal{X}_{0,p}$. We list the related results in the following lemma. Its proof can be found in [B], [BT], [Eb], [GO], [LS], [Mil1], [S].

By abuse of notation, we will identify polynomials in $\mathbb{C}[c, z]$ as polynomials in $\mathbb{C}[z]$ with $\mathbf{C} = \mathbb{C}[c]$. Denote by \mathbf{K} a fixed algebraically closed field containing \mathbf{C} .

Let $f \in \mathbb{C}[c, z]$. By the *zeros of $f \in \mathbb{C}[c, z]$* , we mean the points $(c, z) \in \mathbb{C}^2$ with $f(c, z) = 0$. By the *roots of $f \in \mathbb{C}[z]$* , we mean the roots of f in \mathbf{K} when it is considered as a polynomial in $\mathbf{C}[z]$.

Recall that $\{\nu_d(p)\}_{p \geq 1}$ is a unique sequence of positive integers satisfying the recursive relation $d^p = \sum_{k|p} \nu_d(k)$, $\text{Deg}(f)$ denotes the total degree of f and $\deg_z(f)$ denotes the degree of f as a polynomial in $\mathbf{C}[z]$.

Lemma 2.4. *Let $\mathcal{X}_{0,p}$ be a periodic dynamomic curve. Then*

1. *[B, BT] There exists a unique sequence of monic polynomials $\{Q_{0,p} \in \mathbf{C}[z]\}_{p \geq 1}$ such that for all $p \geq 1$,*

$$\Phi_{0,p}(c, z) := f_c^{\circ p}(z) - z = \prod_{k|p} Q_{0,k}(c, z).$$

Moreover, we have $\text{Deg}(Q_{0,p}) = \text{deg}_z(Q_{0,p}) = \nu_d(p)$.

2. *[BT] Let c_0 be an arbitrary parameter. Then a point z_0 is a root of $Q_{0,p}(c_0, z) \in \mathbf{C}[z]$ if and only if one of the three exclusive conditions is satisfied:*

- (1) *z_0 is a p -periodic point of f_{c_0} and $[f_{c_0}^{\circ p}]'(z_0) \neq 1$,*
- (2) *z_0 is a p -periodic point of f_{c_0} and $[f_{c_0}^{\circ p}]'(z_0) = 1$,*
- (3) *z_0 is an m -periodic point of f_{c_0} , where m is a proper factor of p , and $[f_{c_0}^{\circ m}]'(z_0)$ is a primitive $\frac{p}{m}$ -th root of unity.*

3. *[B, BT, GO, LS, S] The polynomial $Q_{0,p}$ is smooth and irreducible for all $p \geq 1$ and*

$$\mathcal{X}_{0,p} = \{(c, z) \in \mathbf{C} \mid Q_{0,p}(c, z) = 0\}.$$

4. *[B, BT, GO] The projection $\pi_{0,p} : \mathcal{X}_{0,p} \rightarrow \mathbf{C}$, defined by $\pi_{0,p}(c, z) = c$, is a degree $\nu_d(p)$ (given in (1.1)) branched covering with two kinds of critical points:*

- (1) *$C_{0,p}(\text{primitive}) = \{(c, z) \in \mathcal{X}_{0,p} \mid (c, z) \text{ satisfies condition (2) of point 2}\}$. In this case, (c, z) is a simple critical point.*
- (2) *$C_{0,p}(\text{satellite}) = \{(c, z) \in \mathcal{X}_{0,p} \mid (c, z) \text{ satisfies condition (3) of point 2}\}$. In this case, the multiplicity of the critical point (c, z) is $\frac{p}{m} - 1$.*

The critical value set of $\pi_{0,p}$ consists of the parabolic parameters of period p .

5. *[Eb, Mil1] The projection $\varpi_{0,p} : \mathcal{X}_{0,p} \rightarrow \mathbf{C}$, defined by $\varpi_{0,p}(c, z) = z$, is a degree $\nu_d(p)/d$ branched covering, which is injective near each point $(c_0, 0) \in \mathcal{X}_{0,p}$.*
6. *[B] The Galois group $G_{0,p}$ for the polynomial $Q_{0,p} \in \mathbf{C}[z]$ consists of the permutations on roots of $Q_{0,p} \in \mathbf{C}[z]$ that commute with f_c .*

3 The defining polynomial for $\mathcal{X}_{n,p}$

The objective of this section is to show that $\mathcal{X}_{n,p}$ is an affine algebraic curve and find its defining polynomial.

Recall that \mathbf{C} denotes the ring $\mathbf{C}[c]$. For $n \geq 0, p \geq 1$, set $\Phi_{n,p}(c, z) = f_c^{\circ(n+p)}(z) - f_c^{\circ n}(z)$.

Lemma 3.1. *The polynomial $\Phi_{n,p} \in \mathbf{C}[z]$ has no repeated roots. Consequently, it is squarefree.*

Proof. To prove this lemma, it is enough to show that there exists $c_0 \in \mathbf{C}$ such that all roots of $\Phi_{n,p}(c_0, z)$ are simple. In fact, given $c_0 \in \mathbf{C} \setminus M_d$, a point z_0 is a root of $\Phi_{n,p}(c_0, z) \in \mathbf{C}[z]$ if and only if z_0 is a (l, k) -preperiodic point of f_{c_0} , where $0 \leq l \leq n$ and $k \mid p$. For such a c_0 , the critical point 0 goes to infinity and all periodic points of f_{c_0} are repelling. It follows that

$$(\partial\Phi_{n,p}/\partial z)(c_0, z_0) = [f_{c_0}^{on}]'(z_0)([f_{c_0}^{op}]'(z_0) - 1) \neq 0,$$

which completes the proof. \square

Lemma 3.2. *There exists a unique double indexed sequence of squarefree, monic polynomials $\{Q_{n,p} \in \mathbf{C}[z]\}_{n,p \geq 1}$, such that for all $n, p \geq 1$,*

$$\Phi_{n,p}(c, z) = \Phi_{n-1,p}(c, z) \prod_{k \mid p} Q_{n,k}(c, z). \quad (3.1)$$

Moreover, we have $\text{Deg}(Q_{n,p}) = \text{deg}_z(Q_{n,p}) = \nu_d(p)(d-1)d^{n-1}$.

Proof. The definition of $\{Q_{n,p}\}_{n,p \geq 1}$ is based on the polynomials $\{Q_{0,p}\}_{p \geq 1}$ which appear in part 1 of Lemma 2.4. We firstly show that $Q_{0,p}(c, z)$ divides $Q_{0,p}(c, f_c(z))$ for any $p \geq 1$. Since the polynomials $Q_{0,p}(c, f_c(z)) \in \mathbf{C}[z]$ are monic, we may perform a Euclidean division to find a monic quotient $Q \in \mathbf{C}[z]$ and a remainder $R \in \mathbf{C}[z]$ with $\text{degree}(R) < \text{degree}(Q_{0,p})$, such that $Q_{0,p}(c, f_c(z)) = Q_{0,p}Q + R$. We need to show that $R = 0$, which enable us to set $Q_{1,p}(c, z) := Q$.

Following Lemma 3.1 and part 1 of Lemma 2.4, the polynomial $Q_{0,p} \in \mathbf{C}[z]$ does not have repeated factors. So its discriminant $\Delta_{0,p} \in \mathbf{C}[c]$ does not identically vanish, and hence $\Delta_{0,p}(c) \neq 0$ outside a finite set. Fix $c_0 \in \mathbf{C}$ such that $\Delta_{0,p}(c_0) \neq 0$. Then any root z_0 of $Q_{0,p}(c_0, z)$ is simple. By part 2 of Lemma 2.4, the point z_0 is also a root of $Q_{0,p}(c_0, f_{c_0}(z))$. As a consequence, $R(c_0, z) = 0$ for all $z \in \mathbf{C}$. Since this is true for every c_0 outside a finite set, we have $R = 0$ as required.

For $n, p \geq 1$, we define $Q_{n,p}(c, z) := Q_{1,p}(c, f_c^{n-1}(z))$. It is clear that each $Q_{n,p} \in \mathbf{C}[z]$ is monic. Note that $\Phi_{n,p}(c, z) = \Phi_{0,p}(c, f_c^n(z))$ for any $n, p \geq 1$, then we have

$$\begin{aligned} \Phi_{n,p}(c, z) &= \Phi_{0,p}(c, f_c^n(z)) \stackrel{\text{Lem.2.4}}{=} \prod_{k \mid p} Q_{0,k}(c, f_c^n(z)) = \prod_{k \mid p} Q_{0,k}(c, f_c^{n-1}(z)) Q_{1,k}(c, f_c^{n-1}(z)) \\ &= \prod_{k \mid p} Q_{0,k}(c, f_c^{n-1}(z)) \prod_{k \mid p} Q_{1,k}(c, f_c^{n-1}(z)) = \Phi_{0,p}(c, f_c^{n-1}(z)) \prod_{k \mid p} Q_{n,k}(c, z) \\ &= \Phi_{n-1,p}(c, z) \prod_{k \mid p} Q_{n,k}(c, z). \end{aligned}$$

Since each $\Phi_{n,p}$ is squarefree (Lemma 3.1), so is each $Q_{n,p}$.

Repeatedly applying (2) and (3) of Lemma 2.2, we have $\text{Deg}(f_c^k(z)) = \deg_z(f_c^k(z)) = d^k$ for $k \geq 1$. It follows that $\text{Deg}(\Phi_{n,p}) = \deg_z(\Phi_{n,p}) = d^{n+p}$ for $n \geq 0, p \geq 1$. Then by the recursive formulas (3.1), (1.1) and (1) of Lemma 2.2, the degree conclusion in the lemma holds. \square

By the definition of $Q_{n,p}$, we get the inductive formulas

$$\begin{cases} Q_{n-1,p}(c, f_c(z)) = Q_{n,p}(c, z), & n \geq 2; \\ Q_{0,p}(c, f_c(z)) = Q_{0,p}(c, z)Q_{1,p}(c, z), & n = 1. \end{cases} \quad (3.2)$$

for each $p \geq 1$. This equation implies that we can obtain the properties of $Q_{n,p}$ by induction on n .

In fact, $Q_{n,p}(c, z)$ is the defining polynomial of $\mathcal{X}_{n,p}$. To see this, we will now study the properties of the roots of $Q_{n,p}(c_0, z) \in \mathbb{C}[z]$ for an arbitrary parameter $c_0 \in \mathbb{C}$.

Proposition 3.3. *Let $n \geq 1, p \geq 1$ be any pair of integers and $c_0 \in \mathbb{C}$ be any parameter. Then $z_0 \in \mathbb{C}$ is a root of $Q_{n,p}(c_0, z)$ if and only if one of the following 5 mutually exclusive conditions holds:*

- (1) z_0 is a (n, p) -preperiodic point of f_{c_0} such that $f_{c_0}^l(z_0) \neq 0$ for any $0 \leq l < n$ and $[f_{c_0}^p]'(f_{c_0}^n(z_0)) \neq 1$.
- (2) z_0 is a (n, p) -preperiodic point of f_{c_0} such that $f_{c_0}^l(z_0) \neq 0$ for any $0 \leq l < n$ and $[f_{c_0}^p]'(f_{c_0}^n(z_0)) = 1$.
- (3) z_0 is a (n, m) -preperiodic point of f_{c_0} such that $f_{c_0}^l(z_0) \neq 0$ for any $0 \leq l < n$ and m is a proper factor of p with $[f_{c_0}^m]'(f_{c_0}^n(z_0))$ a primitive $\frac{p}{m}$ -th root of unity.
- (4) z_0 is a (n, p) -preperiodic point of f_{c_0} such that $f_{c_0}^l(z_0) = 0$ for some $0 \leq l < n$.
- (5) $f_{c_0}^{(n-1)}(z_0) = 0$ and 0 is a p -periodic point of f_{c_0} .

We remark that in case (4), the case of $l = n - 1$ never occurs.

Proof. Fix $c_0 \in \mathbb{C}$. The proof goes by induction on n . As $n = 1$, $Q_{0,p}(c, f_c(z)) = Q_{0,p}(c, z) \cdot Q_{1,p}(c, z)$. We claim that z_0 is a common root of $Q_{0,p}(c_0, z)$ and $Q_{1,p}(c_0, z)$ if and only if $z_0 = 0$ is a p periodic point of f_{c_0} .

For sufficiency, we only need to note that, in this case, 0 is a multiple root of $Q_{0,p}(c_0, f_{c_0}(z))$, but a simple root of $Q_{0,p}(c_0, z)$ by part 2 of Lemma 2.4. For necessity, z_0 must be a multiple root of $Q_{0,p}(c_0, f_{c_0}(z))$. It follows that either $f_{c_0}(z_0)$ is a multiple root of $Q_{0,p}(c_0, z)$ or z_0 is a critical point of f_{c_0} . In the former case, by 4 of Lemma 2.4, c_0 is a parabolic parameter and $f_{c_0}(z_0)$ is a parabolic periodic point. It means that $Q_{0,p}(c_0, f_{c_0}(z))$ and $Q_{0,p}(c_0, z)$ have the same zero multiplicity at z_0 . Then $Q_{1,p}(c_0, z_0) \neq 0$. In the latter case, we have $z_0 = 0$, and by 2 of Lemma 2.4 0 is a p periodic point of f_{c_0} .

Such c_0, z_0 correspond to the condition (5). In any other case, z_0 is a root of $Q_{1,p}(c_0, z)$ if and only if $f_{c_0}(z_0)$ is a root of $Q_{0,p}(c_0, z)$ but z_0 is not periodic. In fact, if z_0 were periodic,

it would have the same period and multiplier as its first image. By part 2 of Lemma 2.4, we get that $Q_{0,p}(c_0, z_0)$ vanishes, which leads to a contradiction. Then part 2 of Theorem 2.4 implies that z_0 satisfies one of the conditions (1), (2), (3), (4) in Proposition 3.3.

Assume that the proposition is established for $1 \leq l < n$. At this time, $Q_{n,p}(c, z) = Q_{n-1,p}(c, f_c(z))$. So for any $c_0 \in \mathbb{C}$, z_0 is a root of $Q_{n,p}(c_0, z)$ if and only if $f_{c_0}(z_0)$ is a root of $Q_{n-1,p}(c_0, z)$. By Lemma 2.1, if $f_{c_0}(z_0)$ satisfies property (2) or (3), then the orbit of z_0 does not contain 0. Therefore by the inductive assumption, the point z_0 satisfies one of the 5 exclusive conditions in Proposition 3.3. \square

In Proposition 3.3, the zeros of $Q_{n,p}(c, z)$ are divided into 5 classes. We give some notations to denote the sets consisting of zeros of most classes in the following table.

The set	The points in the set
$C_{n,p}(\text{primitive})$	(c, z) satisfies the condition (2) in Proposition 3.3
$C_{n,p}(\text{satellite})$	(c, z) satisfies the condition (3) in Proposition 3.3
$C_{n,p}(\text{Misiurewicz})$	(c, z) satisfies the condition (4) in Proposition 3.3
$C_{n,p}(\text{singular})$	(c, z) satisfies the condition (5) in Proposition 3.3

Recall that for any $n, p \geq 1$, the sets $\check{\mathcal{X}}_{n,p}$ and $\mathcal{X}_{n,p}$ are defined by

$$\check{\mathcal{X}}_{n,p} = \{(c, z) \in \mathbb{C}^2 \mid z \text{ is a } (n, p)\text{-preperiodic point of } f_c\}$$

and

$$\mathcal{X}_{n,p} := \text{the closure of } \check{\mathcal{X}}_{n,p} \text{ in } \mathbb{C}^2 .$$

Proposition 3.4. *For $n, p \geq 1$, we have*

$$\mathcal{X}_{n,p} = \{(c, z) \mid Q_{n,p}(c, z) = 0\} \quad \text{and} \quad \mathcal{X}_{n,p} \setminus \check{\mathcal{X}}_{n,p} = C_{n,p}(\text{satellite}) \cup C_{n,p}(\text{singular})$$

Proof. Set $X := \{(c, z) \mid Q_{n,p}(c, z) = 0\}$. Then X is a closed, perfect set. By the definition of $\check{\mathcal{X}}_{n,p}$ and Proposition 3.3, we have

$$X \setminus (C_{n,p}(\text{satellite}) \cup C_{n,p}(\text{singular})) = \check{\mathcal{X}}_{n,p} \subset X \quad (3.3)$$

We claim that the sets $C_{n,p}(\text{satellite})$ and $C_{n,p}(\text{singular})$ are both finite. If so, we get

$$X = \overline{X \setminus (C_{n,p}(\text{satellite}) \cup C_{n,p}(\text{singular}))} = \overline{\check{\mathcal{X}}_{n,p}} = \mathcal{X}_{n,p} \subset X.$$

Hence it remains to check the claim.

If $(c_0, z_0) \in C_{n,p}(\text{satellite})$, it satisfies that $f_{c_0}^{n+p}(z_0) - f_{c_0}^n(z_0) = 0$ and $[f_{c_0}^p]'(f_{c_0}^n(z_0)) = 1$. Hence c_0 is a root of the resultant $R \in \mathbb{C}[c]$ of the equations $f_c^{n+p}(z) - f_c^n(z) = 0$ and $(f_c^p)'(f_c^n(z)) = 1$. For a parameter c outside the Multibrot set, all the periodic points of f_c are repelling, so the polynomials $f_c^{n+p}(z) - f_c^n(z)$ and $(f_c^p)'(f_c^n(z)) - 1$ do not have a common root. It follows that R is not identically zero, and hence, its roots form a finite set. If $(c_0, z_0) \in C_{n,p}(\text{singular})$, then $Q_{0,p}(c_0, 0) = 0$ by point (5) of Proposition 3.3 and point 2 of Lemma 2.4, whereas the roots of $Q_{0,p}(c, 0)$ form a finite set. \square

4 The irreducible factorization of $Q_{n,p}$

In this section, we will show that the curve $\mathcal{X}_{n,p}$, $n \geq 1$, has $d - 1$ smooth irreducible components and analyze the properties of its singular points. We always assume $n \geq 1$ without emphasizing.

4.1 Factorization of $Q_{n,p}$ and the features of its singular points

Recall that for $f \in \mathbb{C}[c, z]$, $\text{Deg}(f)$ denotes the total degree of f and $\text{deg}_z(f)$ denotes the degree of the variable z in f .

Lemma 4.1. (Algebraic version) *There exists a unique sequence of monic polynomials $\{q_{n,p}^j \in \mathbb{C}[z]\}_{1 \leq j \leq d-1}$ such that*

$$Q_{n,p}(c, z) = \prod_{j=1}^{d-1} q_{n,p}^j(c, z).$$

All points in $C_{n,p}(\text{singular})$ are zeros of $q_{n,p}^j \in \mathbb{C}[c, z]$, and there are no other common zeros for $q_{n,p}^i$ and $q_{n,p}^j$ with $i \neq j$. Moreover, we have $\text{Deg}(q_{n,p}^j) = \text{deg}_z(q_{n,p}^j) = \nu_d(p)d^{n-1}$.

(Topological version) *Let $\mathcal{V}_{n,p}^j = \{(c, z) \in \mathbb{C}^2 \mid q_{n,p}^j(c, z) = 0\}$ ($1 \leq j \leq d - 1$). Then $C_{n,p}(\text{singular}) \subset \mathcal{V}_{n,p}^j$ for each j and $\{\mathcal{V}_{n,p}^j \setminus C_{n,p}(\text{singular})\}_{1 \leq j \leq d-1}$ are pairwise disjoint.*

Proof. Recall that $\mathbf{C} = \mathbb{C}[c]$ and \mathbf{K} is a fixed algebraically closed field containing \mathbf{C} .

Let Δ be a root of $Q_{0,p} \in \mathbf{C}[z]$. Then by part 1 of Lemma 2.4,

$$\Phi_{0,p}(c, \Delta) = f_c^p(\Delta) - \Delta = 0.$$

We see from this equation that Δ is periodic under f_c and $\Delta, \dots, f_c^{p-1}(\Delta)$ are roots of $\Phi_{0,p}$. Note that $\Phi_{0,p}(c, 0) = f_c^p(0)$ is a polynomial in the variable c of the degree d^{p-1} , so $\Delta \neq 0$. Consequently, $\omega\Delta, \dots, \omega^{d-1}\Delta$ are not roots of $Q_{0,p}$, where $\omega = e^{\frac{2\pi i}{d}}$, because they are not periodic under f_c . Then by the equation $Q_{0,p}(c, f_c(z)) = Q_{0,p}(c, z)Q_{1,p}(c, z)$ (see (3.2)), we get that $\omega\Delta, \dots, \omega^{d-1}\Delta$ are roots of $Q_{1,p} \in \mathbf{C}[z]$.

Let us factorize $Q_{0,p}$ in \mathbf{K} by

$$Q_{0,p}(c, z) = \prod_{i=1}^{\nu_d(p)} (z - \Delta_i)$$

($\Delta_{s_1} \neq \Delta_{s_2}$ for $s_1 \neq s_2$, because all roots of $\Phi_{0,p} \in \mathbf{C}[z]$ are simple (Lemma 3.1), and so are $Q_{0,p}$ (part 1 of Lemma 2.4)). Then $Q_{1,p}$ can be expressed as

$$Q_{1,p} = \prod_{i=1}^{\nu_d(p)} (z - \omega\Delta_i) \cdots (z - \omega^{d-1}\Delta_i) = \prod_{j=1}^{d-1} \prod_{i=1}^{\nu_d(p)} (z - \omega^j\Delta_i) \quad (4.1)$$

To see this, we first note that for any $s, t \in [1, d-1]$ and $i_1 \neq i_2 \in [1, \nu_d(p)]$, $\omega^s \Delta_{i_1} \neq \omega^t \Delta_{i_2}$. But it is impossible because both Δ_{i_1} and Δ_{i_2} are periodic. Thus $\{\omega \Delta_i, \dots, \omega^{d-1} \Delta_i\}_{i=1}^{\nu_d(p)}$ are pairwise distinct roots of $Q_{1,p} \in \mathbf{C}[z]$ by the discussion above, then $\prod_{i=1}^{\nu_d(p)} (z - \omega \Delta_i) \cdots (z - \omega^{d-1} \Delta_i)$ is a divisor of $Q_{1,p}$. As its degree is $(d-1)\nu_d(p)$, equal to the degree of $Q_{1,p}$, and $Q_{1,p}$ is monic, we get (4.1). For $j \in [1, d-1]$, set

$$q_{1,p}^j(c, z) = \prod_{i=1}^{\nu_d(p)} (z - \omega^j \Delta_i) = (\omega^j)^{\nu_d(p)} \prod_{i=1}^{\nu_d(p)} (\omega^{-j} z - \Delta_i) = (\omega^j)^{\nu_d(p)} Q_{0,p}(c, \omega^{-j} z). \quad (4.2)$$

Note that $d \mid \nu_d(p)$, so $(\omega^j)^{\nu_d(p)} = 1$. Then $q_{1,p}^j(c, z)$ is a monic polynomial in $\mathbf{C}[z]$, satisfying

$$Q_{1,p}(c, z) = \prod_{j=1}^{d-1} q_{1,p}^j(c, z). \quad (4.3)$$

This gives a factorization of $Q_{1,p}$ in $\mathbf{C}[z]$. By formula (4.2) and the degree conclusion in point 1 of Lemma 2.4, the total degree $\text{Deg}(q_{1,p}^j)$ and $\text{deg}_z(q_{1,p}^j)$ are both $\nu_d(p)$.

For $n \geq 2$, we can define $q_{n,p}^j(c, z)$ inductively by $q_{n,p}^j(c, z) = q_{n-1,p}^j(c, f_c(z))$. Using induction, the degree conclusion in the lemma follows directly from (2) and (3) of Lemma 2.2. As $Q_{n,p}(c, z) = Q_{n-1,p}(c, f_c(z))$, we have

$$Q_{n,p}(c, z) = \prod_{j=1}^{d-1} q_{n,p}^j(c, z). \quad (4.4)$$

This is a factorization of $Q_{n,p}(c, z)$ in $\mathbf{C}[z]$.

We are left to prove that each $q_{n,p}^j(c, z)$ satisfies the remaining properties announced in the lemma. For $n = 1$, since $q_{1,p}^j(c, z) = Q_{0,p}(c, \omega^{-j} z)$, then (c_0, z_0) is a common root of $q_{1,p}^i(c, z)$ and $q_{1,p}^j(c, z)$ for some $1 \leq i \neq j \leq d-1 \iff$ both $(c_0, \omega^{-i} z_0)$ and $(c_0, \omega^{-j} z_0)$ are zeros of $Q_{0,p}(c, z)$. It follows that $\omega^{-i} z_0$ and $\omega^{-j} z_0$ are both periodic point of f_{c_0} , hence $z_0 = 0$. Note that, in case (3) of Lemma 2.4 item 2, the critical point 0 is never periodic (Lemma 2.1), so 0 has period p . It follows that $(c_0, z_0) \in C_{1,p}(\text{singular})$. On the other hand, if $(c_0, z_0) \in C_{1,p}(\text{singular})$, then $(c_0, \omega^{-i} z_0) = (c_0, \omega^{-j} z_0) = (c_0, 0)$ is a zero of $Q_{0,p}(c, z)$. For $n \geq 2$, the conclusion can be deduced from the case of $n = 1$ and the definition of $q_{n,p}^j(c, z)$. \square

For convenience, we summarize the definitions of $q_{1,p}^j$ in term of $Q_{0,p}$ and the inductive definitions of $q_{n,p}^j$ ($n \geq 2$) in terms of $q_{n-1,p}^j$ as a corollary.

Corollary 4.2. *For any $p \geq 1, 1 \leq j \leq d-1$, and $\omega = e^{\frac{2\pi i}{d}}$, we have*

$$\begin{cases} q_{1,p}^j(c, z) = Q_{0,p}(c, \omega^{-j} z), \\ q_{n,p}^j(c, z) = q_{n-1,p}^j(c, f_c(z)), \quad n \geq 2. \end{cases}$$

Example: Here are some examples of $Q_{n,p}$ and their decomposition. Let $d = 3$. Suppose $p = 1$, then we have $Q_{0,1}(c, z) = z^3 + c - z$,

$$\begin{aligned} Q_{1,1}(c, z) &= c^2 + cz + z^2 + 2cz^3 + z^4 + z^6 \\ &= (z^3 + c - e^{-\frac{2}{3}\pi i} z)(z^3 + c - e^{-\frac{4}{3}\pi i} z) \\ &= q_{1,1}^1(c, z) \cdot q_{1,1}^2(c, z). \end{aligned}$$

and

$$\begin{aligned} Q_{2,1}(c, z) &= 3c^2 + 3c^4 + (c^6 + 3c + 10c^3 + 6c^5)z^3 + (1 + 12c^2 + 15c^4)z^6 \\ &\quad + (6c + 20c^3)z^9 + (1 + 15c^2)z^{12} + 6cz^{15} + z^{18} \\ &= \left((1 - e^{-\frac{2}{3}\pi i})c + c^3 + (3c^2 - e^{-\frac{2}{3}\pi i})z^3 + 3cz^6 + z^9 \right) \times \\ &\quad \left((1 - e^{-\frac{4}{3}\pi i})c + c^3 + (3c^2 - e^{-\frac{4}{3}\pi i})z^3 + 3cz^6 + z^9 \right) \\ &= q_{2,1}^1(c, z) \cdot q_{2,1}^2(c, z). \end{aligned}$$

Suppose $p = 2$, then we have $Q_{0,2}(c, z) = 1 + c^2 + cz + z^2 + 2cz^3 + z^4 + z^6$ and

$$\begin{aligned} Q_{1,2}(c, z) &= 1 + 2c^2 + c^4 - (c + c^3) - z^2 + (3c + 4c^3)z^3 - 3c^2z^4 \\ &\quad + (1 + 6c^2)z^6 - 3cz^7 + 4cz^9 - z^{10} + z^{12} \\ &= \left(1 + c^2 + e^{-\frac{2}{3}\pi i} z + e^{-\frac{4}{3}\pi i} z^2 + 2cz^3 + e^{-\frac{2}{3}\pi i} z^4 + z^6 \right) \times \\ &\quad \left(1 + c^2 + e^{-\frac{4}{3}\pi i} z + e^{-\frac{2}{3}\pi i} z^2 + 2cz^3 + e^{-\frac{4}{3}\pi i} z^4 + z^6 \right) \\ &= q_{1,2}^1(c, z) \cdot q_{1,2}^2(c, z) \end{aligned}$$

From Lemma 4.1, we see that in the case $d \geq 3$, the polynomial $Q_{n,p}$ is both reducible and non-smooth, because $C_{n,p}(\text{singular})$, which is non-empty, belongs to the set of singular points of $Q_{n,p}$.

We now turn to the study of the components $q_{n,p}^j(c, z)$. The following theorem is the core of this section.

Theorem 4.3. *Given $d \geq 2$, for any $n, p \geq 1$, $j \in [1, d - 1]$, the polynomial $q_{n,p}^j(c, z)$ is smooth and irreducible.*

The proof of this theorem is postponed to § 4.2.

By this theorem, all components $\mathcal{V}_{n,p}^j$ are Riemann surfaces. Together with Lemma 4.1, this implies that the singularity set of $\mathcal{X}_{n,p}$ is equal to $C_{n,p}(\text{singular})$. The next proposition characterizes the features of these singularities.

Proposition 4.4. *Given $d \geq 2$, for $n, p \geq 1$, each singularity (c_0, z_0) of $\mathcal{X}_{n,p}$ has multiplicity $d - 1$. Furthermore, if $f_{c_0}^l(z_0) = 0$ for some $0 \leq l \leq n - 2$, then $\mathcal{X}_{n,p}$ has one tangent of multiplicity $d - 1$ at (c_0, z_0) ; otherwise, the singularity (c_0, z_0) is ordinary.*

Proof. Let (c_0, z_0) be a singular point of $\mathcal{X}_{n,p}$. Since each component of $\mathcal{X}_{n,p}$ is smooth and they are pairwise intersecting at (c_0, z_0) , then the first non-vanishing term of $Q_{n,p}(c, z)$ at (c_0, z_0) is $d - 1$. Hence the multiplicity of the singularity (c_0, z_0) is $d - 1$.

If $n = 1$, by (5) of Proposition 3.3, the fact that $(c_0, z_0) \in C_{1,p}(\text{singular})$ implies that $z_0 = 0$ and $(c_0, 0) \in \mathcal{X}_{0,p}$. According to Lemma 2.1 c_0 is not a parabolic parameter. Then it follows from part 4 of Lemma 2.4 that $(c_0, 0)$ is not a critical point of $\pi_{0,p}$, and hence $(\partial Q_{0,p}/\partial z)(c_0, 0) \neq 0$. Meanwhile, according to part 5 of Lemma 2.4, $(\partial Q_{0,p}/\partial c)(c_0, 0) \neq 0$. Thus $Q_{0,p}(c, z)$ has a local expression

$$Q_{0,p}(c, z) = a_{0,p}(c - c_0) + b_{0,p}z + \text{higher order terms}$$

around $(c_0, 0)$ with $a_{0,p}, b_{0,p} \neq 0$. It follows that

$$q_{1,p}^j(c, z) = Q_{0,p}(c, \omega^{-j}z) = a_{0,p}(c - c_0) + b_{0,p}\omega^{-j}z + \text{higher order terms}$$

Therefore the tangents of $\mathcal{V}_{1,p}^j$ ($1 \leq j \leq d-1$) at $(c_0, 0)$ are pairwise distinct.

For $n \geq 2$, we denote by $a_{n,p}^j(c - c_0) + b_{n,p}^j(z - z_0)$ the equation of the tangent of $\mathcal{V}_{n,p}^j$ at (c_0, z_0) . By the formula $q_{n,p}^j(c, z) = q_{1,p}(c, f_c^{n-1}(z))$ (Corollary 4.2), we have that

$$\begin{cases} a_{n,p}^j &= \frac{\partial q_{n,p}^j}{\partial c}(c_0, z_0) = \frac{\partial q_{1,p}^j}{\partial c}(c_0, 0) + \frac{\partial q_{1,p}^j}{\partial z}(c_0, 0) \frac{\partial f_c^{n-1}}{\partial c}(c_0, z_0) = a_{0,p} + b_{0,p}\omega^{-j} \frac{\partial f_c^{n-1}}{\partial c}(c_0, z_0) \\ b_{n,p}^j &= \frac{\partial q_{n,p}^j}{\partial z}(c_0, z_0) = \frac{\partial q_{1,p}^j}{\partial z}(c_0, 0)(f_{c_0}^{n-1})'(z_0) = b_{0,p}\omega^{-j}(f_{c_0}^{n-1})'(z_0) \end{cases}$$

If there exists $0 \leq l \leq n-2$ such that $f_{c_0}^l(z_0) = 0$, then $(f_{c_0}^{n-1})'(z_0) = 0$, and hence $b_{n,p}^j = 0$. It follows that the first non-vanishing term of $Q_{n,p}$ at (c_0, z_0) is $a(c - c_0)^{d-1}$ where a is a non-zero constant, i.e., $\mathcal{X}_{n,p}$ has the tangent $c = c_0$ of multiplicity $d-1$ at (c_0, z_0) . In the other cases, we get $(f_{c_0}^{n-1})'(z_0) \neq 0$. Combining this point and the fact that $a_{0,p}, b_{0,p} \neq 0$, it is not difficult to check that the pairs $(a_{n,p}^j, b_{n,p}^j)$ ($1 \leq j \leq d-1$) are pairwise non-colinear. Hence the tangents of $\mathcal{V}_{n,p}^j$ ($1 \leq j \leq d-1$) at (c_0, z_0) are pairwise distinct, that is, (c_0, z_0) is ordinary. \square

4.2 Proof of the smoothness and irreducibility of $q_{n,p}^j$

The objective here is to prove Theorem 4.3.

The approach to prove the smoothness is similar to that in [BT]. The idea is to prove that some partial derivative of $q_{n,p}^j$ is non vanishing. Following A. Epstein, we will express this derivative as the coefficient of a quadratic differential of the form $(f_c)_\star \mathcal{Q} - \mathcal{Q}$. Thurston's contraction principle gives $(f_c)_\star \mathcal{Q} - \mathcal{Q} \neq 0$, whence our partial derivative is non-zero.

The approach to the irreducibility is based on the connectedness of periodic curve $\mathcal{X}_{0,p}$. Then we will show the connectivity of $\mathcal{V}_{n,p}^j$ using a branched covering by induction on the preperiodic index n .

Here we list some definitions and results about quadratic differentials and Thurston's contraction principle. All their proofs can be found in [BT] and [Le].

We use $\mathcal{Q}(\mathbb{C})$ to denote the set of meromorphic quadratic differentials on \mathbb{C} whose poles (if any) are all simple. If $\mathcal{Q} \in \mathcal{Q}(\mathbb{C})$ and U is a bounded open subset of \mathbb{C} , the norm

$$\|\mathcal{Q}\|_U := \iint_U |q|$$

is well defined and finite.

For $f : \mathbb{C} \rightarrow \mathbb{C}$ a non-constant polynomial and $\mathcal{Q} = q dz^2$ a meromorphic quadratic differential on \mathbb{C} , the pushforward $f_*\mathcal{Q}$ is defined by the quadratic differential

$$f_*\mathcal{Q} := Tq dz^2 \quad \text{with} \quad Tq(z) := \sum_{f(w)=z} \frac{q(w)}{f'(w)^2}.$$

If $\mathcal{Q} \in \mathcal{Q}(\mathbb{C})$, then $f_*\mathcal{Q} \in \mathcal{Q}(\mathbb{C})$ also. The following lemma is a weak version of Thurston's contraction principle.

Lemma 4.5. *If $f : \mathbb{C} \rightarrow \mathbb{C}$ is a polynomial and if $\mathcal{Q} \in \mathcal{Q}(\mathbb{C})$, then $f_*\mathcal{Q} \neq \mathcal{Q}$.*

The formulas below appeared in [Le] chapter 2, we write them together as a lemma.

Lemma 4.6 (Levin). *For $f = f_c$, we have*

$$\begin{cases} f_*\left(\frac{dz^2}{z}\right) = 0 \\ f_*\left(\frac{dz^2}{z-a}\right) = \frac{1}{f'(a)}\left(\frac{dz^2}{z-f(a)} - \frac{dz^2}{z-c}\right) \quad \text{if } a \neq 0 \end{cases} \quad (4.5)$$

To prove the irreducibility of $q_{n,p}^j$, we need the following Lemma.

Lemma 4.7. *For each $n, p \geq 1$, $1 \leq j \leq d-1$, the polynomial $q_{n,p}^j(c, 0)$ (in the variable c) has degree $\nu_d(p)d^{n-2}$.*

Proof. For $n = 1$, we see that $q_{1,p}^j(c, 0) = Q_{0,p}(c, 0)$ from Corollary 4.2. Then the result follows directly from point 5 of Lemma 2.4.

For $n \geq 2$, $q_{n,p}^j(c, 0) = q_{1,p}^j(c, f_c^{n-1}(0))$. Since $\text{Deg}(q_{1,p}^j) = \deg_z(q_{1,p}^j) = \nu_d(p)$ (see Lemma 4.1) and $\text{Deg}(f_c^{n-1}(0)) = d^{n-2}$ (which is easily checked), we have

$$\text{Deg}(q_{n,p}^j(c, 0)) = \text{Deg}(q_{1,p}^j(c, z)) \cdot \text{Deg}(f_c^{n-1}(0)) = \nu_d(p)d^{n-2}$$

by (2) and (3) of Lemma 2.2. Then the proof is completed. \square

Proof of Theorem 4.3. The proof goes by induction on n .

For $n = 1$, as $q_{1,p}^j(c, z) = Q_{0,p}(c, \omega^{-j}z)$ and $Q_{0,p}(c, z)$ is smooth and irreducible, we know that $q_{1,p}^j(c, z)$ are smooth and irreducible. Assume that for $1 \leq l < n$, the polynomial

$q_{l,p}^j(c, z)$ are smooth and irreducible. Then we will show that $q_{n,p}^j(c, z)$ are smooth and irreducible. Now fix any $j_0 \in [1, d-1]$.

Smoothness of $q_{n,p}^{j_0}$: As $q_{n,p}^{j_0}(c, z) = q_{n-1,p}^{j_0}(c, f_c(z))$, for any (c_0, z_0) a zero of $q_{n,p}^{j_0}(c, z)$, we have

$$\begin{cases} \frac{\partial q_{n,p}^{j_0}}{\partial c}(c_0, z_0) = \frac{\partial q_{n-1,p}^{j_0}}{\partial c}(c_0, w_0) + \frac{\partial q_{n-1,p}^{j_0}}{\partial z}(c_0, w_0) \\ \frac{\partial q_{n,p}^{j_0}}{\partial z}(c_0, z_0) = \frac{\partial q_{n-1,p}^{j_0}}{\partial z}(c_0, w_0) \cdot f'_{c_0}(z_0) \end{cases} \quad (4.6)$$

where $w_0 = f_{c_0}(z_0)$. Then if $z_0 \neq 0$, by the smoothness of $\mathcal{V}_{n,p}^{j_0}$ (assumption of induction), $[\partial q_{n,p}^{j_0}/\partial c](c_0, z_0)$ and $[\partial q_{n,p}^{j_0}/\partial z](c_0, z_0)$ can not be equal to 0 simultaneously, it follows that $q_{n,p}^{j_0}(c, z)$ is smooth at (c_0, z_0) . So we are left to prove that $q_{n,p}^{j_0}(c, z)$ is smooth at $(c_0, 0) \in \mathcal{V}_{n,p}^{j_0}$. In this situation, c_0 is either a p -periodic super-attracting parameter or a (n, p) -Misiurewicz parameter, and $[\partial q_{n,p}^{j_0}/\partial z](c_0, 0) = 0$. So we have to show $[\partial q_{n,p}^{j_0}/\partial c](c_0, 0) \neq 0$.

In the former case $f_{c_0}^{n-1}(0) = 0$, then $p|n-1$. Since $q_{n,p}^{j_0}(c, z) = q_{1,p}^{j_0}(c, f_c^{n-1}(z))$, we have

$$\frac{\partial q_{n,p}^{j_0}}{\partial c}(c_0, 0) = \frac{\partial q_{1,p}^{j_0}}{\partial c}(c_0, 0) + \frac{\partial q_{1,p}^{j_0}}{\partial z}(c_0, 0) \frac{\partial f_c^{n-1}}{\partial c}(c_0, 0) \quad (4.7)$$

Note that $Q_{0,p}(c_0, 0) = 0$ and $p|n-1$, then differentiating both sides of the equation

$$f_c^{n-1}(z) - z = \prod_{k|n-1} Q_{0,k}(c, z),$$

which is raised in point 1 of Lemma 2.4, with respect to c and z respectively at the point $(c_0, 0)$, we have that

$$\begin{cases} \frac{\partial f_c^{n-1}}{\partial c}(c_0, 0) = \frac{\partial Q_{0,p}}{\partial c}(c_0, 0) \prod_{\substack{k|n-1 \\ k \neq p}} Q_{0,k}(c_0, 0) \\ -1 = \frac{\partial Q_{0,p}}{\partial z}(c_0, 0) \prod_{\substack{k|n-1 \\ k \neq p}} Q_{0,k}(c_0, 0) \end{cases} \quad (4.8)$$

Since $q_{1,p}^{j_0}(c, z) = Q_{0,p}(c, \omega^{-j_0}z)$, then

$$\frac{\partial q_{1,p}^{j_0}}{\partial c}(c_0, 0) = \frac{\partial Q_{0,p}}{\partial c}(c_0, 0), \quad \frac{\partial q_{1,p}^{j_0}}{\partial z}(c_0, 0) = \omega^{-j_0} \frac{\partial Q_{0,p}}{\partial z}(c_0, 0).$$

By substituting these two formulas into equation (4.7) and applying equation (4.8), we find

$$\begin{aligned} \frac{\partial q_{n,p}^{j_0}}{\partial c}(c_0, 0) &= \frac{\partial Q_{0,p}}{\partial c}(c_0, 0) + \omega^{-j_0} \frac{\partial Q_{0,p}}{\partial z}(c_0, 0) \frac{\partial Q_{0,p}}{\partial c}(c_0, 0) \prod_{\substack{k|n-1 \\ k \neq p}} Q_{0,k}(c_0, 0) \\ &= \frac{\partial Q_{0,p}}{\partial c}(c_0, 0) (1 + \omega^{-j_0} \frac{\partial Q_{0,p}}{\partial z}(c_0, 0) \prod_{\substack{k|n-1 \\ k \neq p}} Q_{0,k}(c_0, 0)) \\ &= \frac{\partial Q_{0,p}}{\partial c}(c_0, 0) (1 - \omega^{-j_0}). \end{aligned} \quad (4.9)$$

By point 5 of Lemma 2.4, $[\partial Q_{0,p}/\partial c](c_0, 0) \neq 0$, then so is $[\partial q_{n,p}^{j_0}/\partial c](c_0, 0)$.

In the latter case, since

$$\frac{\partial Q_{n,p}}{\partial c}(c_0, 0) = \prod_{1 \leq j \neq j_0 \leq d-1} q_{n,p}^j(c_0, 0) \cdot \frac{\partial q_{n,p}^{j_0}}{\partial c}(c_0, 0)$$

and the point $(c_0, 0)$ is not a zero of $\prod_{j \neq j_0} q_{n,p}^j(c, z)$ by Lemma 4.1, we only have to show $[\partial Q_{n,p}/\partial c](c_0, 0) \neq 0$. Furthermore, since

$$\frac{\partial \Phi_{n,p}}{\partial c}(c_0, 0) = \Phi_{n-1,p}(c_0, 0) \cdot \prod_{k|p, k < p} Q_{n,k}(c_0, 0) \cdot \frac{\partial Q_{n,p}}{\partial c}(c_0, 0)$$

and $\Phi_{n-1,p}(c_0, 0) \cdot \prod_{k|p, k < p} Q_{n,k}(c_0, 0) \neq 0$, it is equivalent to show $[\partial \Phi_{n,p}/\partial c](c_0, 0) \neq 0$. We shall choose a meromorphic quadratic differential with simple poles such that

$$(f_{c_0})_* \mathcal{Q} = \mathcal{Q} + \frac{\partial \Phi_{n,p}}{\partial c}(c_0, 0) \cdot \frac{dz^2}{z - c_0}.$$

Then by Lemma 4.5, we obtain $[\partial \Phi_{n,p}/\partial c](c_0, 0) \neq 0$.

We shall use the following notations:

$$\begin{aligned} z_k &:= f_{c_0}^{\circ(n+k)}(0), & \delta_k &:= f'_{c_0}(z_k) = dz_k^{d-1}, & 0 \leq k \leq p-1 \\ y_l &:= f_{c_0}^l(0), & \varepsilon_l &:= f'_l(y_l) = dy_l^{d-1}, & 1 \leq l \leq n-1 \end{aligned}$$

With these notations and a bit of calculations, we get

$$\begin{aligned} \frac{\partial \Phi_{n,p}}{\partial c}(c_0, 0) &= \frac{\partial f_c^{\circ(n+p)}}{\partial c}(c_0, 0) - \frac{\partial f_c^{\circ n}}{\partial c}(c_0, 0) \\ &= (\delta_0 \cdots \delta_{p-1} - 1)(\varepsilon_{n-1} \cdots \varepsilon_1 + \cdots + \varepsilon_{n-1} \varepsilon_{n-2} + \varepsilon_{n-1} + 1) \\ &\quad + \delta_{p-1} \cdots \delta_1 + \cdots + \delta_{p-1} + 1 \end{aligned}$$

Denote $(\delta_0 \cdots \delta_{p-1} - 1)(\varepsilon_{n-1} \cdots \varepsilon_1 + \cdots + \varepsilon_{n-1} \varepsilon_{n-2} + \varepsilon_{n-1} + 1)$ by α . Let

$$\mathcal{Q} = \sum_{k=0}^{p-1} \frac{\rho_k}{z - z_k} dz^2 + \sum_{l=1}^{n-1} \frac{\lambda_l}{z - y_l} dz^2$$

be a quadratic differential in $\mathcal{Q}(\mathbb{C})$. Here ρ_k ($0 \leq k \leq p-1$), λ_l ($1 \leq l \leq n-1$) are undetermined coefficients (note that $y_1 = c_0$). Applying Lemma 4.6 and writing f for f_{c_0} , we have

$$\begin{aligned} f_* \mathcal{Q} &= \sum_{k=0}^{p-1} \frac{\rho_k}{\delta_k} \left(\frac{dz^2}{z - z_{k+1}} - \frac{dz^2}{z - c_0} \right) + \sum_{l=1}^{n-2} \frac{\lambda_l}{\varepsilon_l} \left(\frac{dz^2}{z - y_{l+1}} - \frac{dz^2}{z - c_0} \right) + \frac{\lambda_{n-1}}{\varepsilon_{n-1}} \left(\frac{dz^2}{z - z_0} - \frac{dz^2}{z - c_0} \right) \\ &= \left(\frac{\rho_{p-1}}{\delta_{p-1}} + \frac{\lambda_{n-1}}{\varepsilon_{n-1}} \right) \frac{dz^2}{z - z_0} + \frac{\rho_0}{\delta_0} \frac{dz^2}{z - z_1} + \cdots + \frac{\rho_{p-2}}{\delta_{p-2}} \frac{dz^2}{z - z_{p-1}} \\ &\quad + \left(\alpha - \sum_{l=1}^{n-1} \frac{\lambda_l}{\varepsilon_l} \right) \frac{dz^2}{z - y_1} + \frac{\lambda_1}{\varepsilon_1} \frac{dz^2}{z - y_2} + \cdots + \frac{\lambda_{n-2}}{\varepsilon_{n-2}} \frac{dz^2}{z - y_{n-1}} - \left(\alpha + \sum_{k=0}^{p-1} \frac{\rho_k}{\delta_k} \right) \frac{dz^2}{z - c_0} \end{aligned}$$

Note that for $n = 1$, we can also define a map $\wp_{1,p}^j : \mathcal{V}_{1,p}^j \rightarrow \mathcal{X}_{0,p}$ by $\wp_{1,p}^j(c, z) = (c, f_c(z))$. By the smoothness of $\mathcal{V}_{n,p}^j$, we can check the following results.

- The map $\wp_{1,p}^j : \mathcal{V}_{1,p}^j \rightarrow \mathcal{X}_{0,p}$ is a homeomorphism. To see this, notice that $q_{1,p}^j(c, z) = Q_{0,p}(c, \omega^{-j}z)$ (Corollary 4.2), so we can define a map $\phi_{1,p}^j$ from $\mathcal{X}_{0,p}$ to $\mathcal{V}_{1,p}^j$ by mapping a point $(c_0, w_0) \in \mathcal{X}_{0,p}$ to $(c_0, \omega^j z_0) \in \mathcal{V}_{1,p}^j$, where z_0 is the point in the orbit of w_0 under f_{c_0} with $f_{c_0}(z_0) = w_0$. By a simple computation, we can see that $\phi_{1,p}^j \circ \wp_{1,p}^j = id_{\mathcal{V}_{1,p}^j}$ and $\wp_{1,p}^j \circ \phi_{1,p}^j = id_{\mathcal{X}_{0,p}}$. Hence $\wp_{1,p}^j$ is a homeomorphism.
- For $n \geq 2$, the map $\wp_{n,p}^j : \mathcal{V}_{n,p}^j \rightarrow \mathcal{V}_{n-1,p}^j$ is a degree d branched covering with critical set

$$D_{n,p}^j = \{(c, 0) \mid q_{n,p}^j(c, 0) = 0\}.$$

and each critical point has multiplicity $d - 1$.

In fact, a point $(c_0, w_0) \in \mathcal{V}_{n-1,p}^j \setminus \wp(D_{n,p}^j)$ has d preimages $(c_0, z_1), \dots, (c_0, z_d)$ under $\wp_{n,p}^j$, where z_1, \dots, z_d are preimages of w_0 under f_{c_0} . Fix $i \in [1, d]$. If $[\partial q_{n,p}^j / \partial z](c_0, z_i) \neq 0$, then by equation (4.6), $[\partial q_{n-1,p}^j / \partial z](c_0, w_0) \neq 0$. It implies that some neighborhoods of (c_0, z_i) and (c_0, w_0) can be parameterized by c respectively. Using such two local coordinates, the map $\wp_{n,p}^j$ has a local expression $c \mapsto c$ near (c_0, z_i) , which means that $\wp_{n,p}^j$ is a local homeomorphism near (c_0, z_i) . If $[\partial q_{n,p}^j / \partial z](c_0, z_i) = 0$, then by equation (4.6), the fact of $z_i \neq 0$ and the smoothness of $q_{n,p}^j$, we have that $[\partial q_{n-1,p}^j / \partial z](c_0, w_0) = 0$ and

$$\frac{\partial q_{n,p}^j}{\partial c}(c_0, z_i) = \frac{\partial q_{n-1,p}^j}{\partial c}(c_0, w_0) \neq 0$$

It implies that some neighborhoods of (c_0, z_i) and (c_0, w_0) can be parameterized by z respectively, and $c'(z_i) = 0$. Using such two local coordinates, the map $\wp_{n,p}^j$ has a local expression $z \mapsto f_{c(z)}(z)$ near (c_0, z_i) . Since $z_i \neq 0$, then $\left. \frac{df_{c(z)}(z)}{dz} \right|_{z=z_i} = dz_i \neq 0$, which still means that $\wp_{n,p}^j$ is a local homeomorphism near (c_0, z_i) .

By the discussion above, we can see that

$$\wp_{n,p}^j : \mathcal{V}_{n,p}^j \setminus (\wp_{n,p}^j)^{-1}(\wp(D_{n,p}^j)) \rightarrow \mathcal{V}_{n-1,p}^j \setminus \wp(D_{n,p}^j)$$

is a degree d covering. On the other hand, for any point in $\wp_{n,p}^j(D_{n,p}^j)$, it has only one preimage, which belongs to $D_{n,p}^j$. Hence we have that $\wp : \mathcal{V}_{n,p}^j \rightarrow \mathcal{V}_{n-1,p}^j$ is a degree d branched covering (because $(\wp_{n,p}^j)^{-1}(\wp(D_{n,p}^j)) = D_{n,p}^j$ and $D_{n,p}^j$ is finite) and the local degree of $\wp_{n,p}^j$ at each point of $D_{n,p}^j$ is d .

By the smoothness of $q_{n,p}^{j_0}(c, z)$ and the inductive assumption of irreducibility, we know that $\mathcal{V}_{n-1,p}^{j_0}$ and each connected component of $\mathcal{V}_{n,p}^{j_0}$ is a Riemann surface. Then the restriction of $\wp_{n,p}^{j_0}$ on any connected component of $\mathcal{V}_{n,p}^{j_0}$ is also a branched covering. Lemma 4.7 implies that the critical set $D_{n,p}^{j_0}$ of $\wp_{n,p}^{j_0}$ is non-empty. Since each critical point has multiplicity $d - 1$, the set $\mathcal{V}_{n,p}^{j_0}$ must be connected. By Lemma 2.3 and the smoothness of $q_{n,p}^{j_0}$, we conclude that $q_{n,p}^{j_0}(c, z)$ is irreducible in $\mathbb{C}[c, z]$.

□

5 Genus of the compactification of $\mathcal{V}_{n,p}^j$

In the previous section, we have seen that $\mathcal{X}_{n,p}$ consists of $d - 1$ Riemann surfaces $\mathcal{V}_{n,p}^j$ which are pairwise intersecting at the singular points of $\mathcal{X}_{n,p}$. In order to give a complete topological description of $\mathcal{X}_{n,p}$, we also need the topological characterization of each $\mathcal{V}_{n,p}^j$.

In fact, by adding an ideal boundary point at each end of $\mathcal{V}_{n,p}^j$, we obtain a compactification of $\mathcal{V}_{n,p}^j$, denoted by $\widehat{\mathcal{V}}_{n,p}^j$, such that $\widehat{\mathcal{V}}_{n,p}^j$ is a compact Riemann surface (in § 5.1). The genus of $\widehat{\mathcal{V}}_{n,p}^j$ is calculated (in § 5.2). Topologically, $\mathcal{X}_{n,p}$ is completely determined by the number of its singular points, the genus of $\widehat{\mathcal{V}}_{n,p}^j$ and the number of ideal boundary points added on $\mathcal{V}_{n,p}^j$ (or the number of ends of $\mathcal{V}_{n,p}^j$).

5.1 Compactification of $\mathcal{V}_{n,p}^j$

Denote by $\pi_{n,p}^j: \mathcal{V}_{n,p}^j \rightarrow \mathbb{C}$ the projection from $\mathcal{V}_{n,p}^j$ to the parameter plane, i.e., $\pi_{n,p}^j(c, z) = c$. It is easy to see

$$\pi_{n,p}^j = \pi_{0,p} \circ \wp_{1,p}^j \circ \cdots \circ \wp_{n-1,p}^j \circ \wp_{n,p}^j \quad (5.1)$$

where $\pi_{0,p}$ is the projection from $\mathcal{X}_{0,p}$ to the parameter plane and $\wp_{n,p}^j$ is defined in the proof of irreducibility. It follows that $\pi_{n,p}^j$ is a degree $\nu_d(p)d^{n-1}$ branched covering. To study the critical points of $\pi_{n,p}^j$, we define a subset $C_{n,p}^{\text{crit}}(\text{singular})$ of $C_{n,p}(\text{singular})$ by

$$C_{n,p}^{\text{crit}}(\text{singular}) = \{(c, z) \in C_{n,p}(\text{singular}) \mid f_c^l(z) = 0 \text{ for some } 0 \leq l \leq n - 2\} \quad (5.2)$$

Lemma 5.1. *For any $l, p \geq 1$, the critical set of $\pi_{l,p}^j$ is the union of $C_{l,p}^j(\text{primitive})$, $C_{l,p}^j(\text{satellite})$, $C_{l,p}^j(\text{Misiurewicz})$ and $C_{l,p}^{\text{crit}}(\text{singular})$, where $C_{l,p}^j(M) := C_{l,p}(M) \cap \mathcal{V}_{l,p}^j$ and M indicates different properties.*

Proof. We first note that (c_0, z_0) is a critical point of $\pi_{l,p}^j$ if and only if $[\partial q_{l,p}^j / \partial z](c_0, z_0) = 0$. By part 4 of Lemma 2.4 and the fact that $\wp_{1,p}^j$ is homeomorphic (which is shown in the proof of irreducibility of $q_{l,p}^j$), the critical set of $\pi_{1,p}^j$ is $C_{1,p}^j(\text{primitive}) \cup C_{1,p}^j(\text{satellite})$. In the case $l = 1$, $C_{l,p}(\text{Misiurewicz})$ and $C_{l,p}^{\text{crit}}(\text{singular})$ are empty.

For $l \geq 2$, by Corollary 4.2, we have $q_{l,p}^j(c, z) = q_{1,p}^j(c, f_c^{l-1}(z))$. Then a point (c_0, z_0) is critical for $\pi_{l,p}^j$ if and only if

$$\frac{\partial q_{l,p}^j}{\partial z}(c_0, z_0) = \frac{\partial q_{1,p}^j}{\partial z}(c_0, f_{c_0}^{l-1}(z_0)) \cdot (f_{c_0}^{l-1})'(z_0) = 0$$

It is equivalent that either $(c_0, f_{c_0}^{l-1}(z_0))$ is a critical point of $\wp_{1,p}^j$ or $f_{c_0}^l(z_0) = 0$ for some $0 \leq q \leq n - 2$. By Proposition 3.3, the former case happens if and only if $(c_0, z_0) \in C_{l,p}^j(\text{primitive}) \cup C_{l,p}^j(\text{satellite})$, and the latter case happens if and only if $(c_0, z_0) \in C_{l,p}^j(\text{Misiurewicz}) \cup C_{l,p}^{\text{crit}}(\text{singular})$. \square

From this Lemma, we see that the critical value set of $\pi_{n,p}^j$ is contained in the union of parabolic, super-attracting and Misiurewicz parameters. Hence $\mathbb{C} \setminus M_d$ contains no critical values. It follows that each connected component of $(\wp_{n,p}^j)^{-1}(\mathbb{C} \setminus M_d)$, called an *end* of $\mathcal{V}_{n,p}^j$, is conformal to $\mathbb{C} \setminus \overline{\mathbb{D}}$. By adding an ideal boundary point at the infinitely far boundary, each end of $\mathcal{V}_{n,p}^j$ is conformal to the unit disk, and then $\mathcal{V}_{n,p}^j$ becomes a compact Riemann surface. This gives a kind of compactification of $\mathcal{V}_{n,p}^j$ and we will calculate in the next subsection the genus of this compact Riemann surface.

More precisely, set $\{\mathcal{E}_{n,p,i}^j\}$ ($1 \leq i \leq m_{n,p}^j$) the ends of $\mathcal{V}_{n,p}^j$. Denote by $\infty_{n,p,i}^j$ the point added at the infinitely far boundary of $\mathcal{E}_{n,p,i}^j$. Then the surface $\widehat{\mathcal{V}}_{n,p}^j := \mathcal{V}_{n,p}^j \cup \{\infty_{n,p,i}^j\}_{i=1}^{m_{n,p}^j}$ is a compactification of $\mathcal{V}_{n,p}^j$ and $\widehat{\mathcal{E}}_{n,p,i}^j := \mathcal{E}_{n,p,i}^j \cup \{\infty_{n,p,i}^j\}$ is called an *end* of $\widehat{\mathcal{V}}_{n,p}^j$. In this case, the map $\pi_{n,p}^j$ can be extended to

$$\widehat{\pi}_{n,p}^j : \widehat{\mathcal{V}}_{n,p}^j \longrightarrow \widehat{\mathbb{C}}$$

by setting $\widehat{\pi}_{n,p}^j(\infty_{n,p,i}^j) = \infty$.

5.2 Calculation of the genus of $\widehat{\mathcal{V}}_{n,p}^j$

Now, for any $n, p \geq 1$, $j \in [1, d-1]$, we have obtained a branched covering $\widehat{\pi}_{n,p}^j : \widehat{\mathcal{V}}_{n,p}^j \rightarrow \widehat{\mathbb{C}}$ of degree $\nu_d(p)d^{n-1}$ between two compact Riemann surface. By the Riemann-Hurwitz formula, we have

$$2 - 2g_{n,p}^j + \text{total number of critical points of } \widehat{\pi}_{n,p}^j = 2\nu_d(p)d^{n-1}.$$

where $g_{n,p}^j$ denotes the genus of $\widehat{\mathcal{V}}_{n,p}^j$. So in order to calculate the genus of $\widehat{\mathcal{V}}_{n,p}^j$, we only need to count the number of critical points of $\widehat{\pi}_{n,p}^j$ counting with multiplicity. By Lemma 5.1, we know that the critical points of $\widehat{\pi}_{n,p}^j$ consists of the points of $C_{n,p}^j$ (primitive), $C_{n,p}^j$ (satellite), $C_{n,p}^j$ (Misiurewicz), $C_{n,p}^{\text{crit}}$ (singular) and maybe some added ideal boundary points. So we will count them class by class.

- Counting the points of $C_{n,p}^j$ (primitive) and $C_{n,p}^j$ (satellite).

In [B], Bousch counts the number of critical points in $C_{0,p}$ (primitive) and $C_{0,p}$ (satellite). His argument can be directly extended to our case (see also [Sil, Thm. 4.17]). so we only list the result without the counting process. The number of critical points counted with multiplicity of $\widehat{\pi}_{n,p}^j$ in $C_{n,p}^j$ (primitive) and $C_{n,p}^j$ (satellite) are

$$d^{n-1}p[(d-1)\nu_d(p)/d - \sum_{k|p, k < p} (\nu_d(k)/d)(d-1)\varphi(p/k)]$$

and

$$d^{n-1} \sum_{k|p, k < p} (\nu_d(k)/d)(d-1)\varphi(p/k)k(p/k - 1).$$

- Counting the points of $C_{n,p}^j$ (Misiurewicz).

Recall that $D_{s,p}^j = \{(c, 0) \in \mathbb{C}^2 \mid q_{s,p}^j(c, 0) = 0\}$, $s \geq 2$, is the set of critical points of $\wp_{s,p}^j$. By Proposition 3.3, if $(c, 0) \in D_{s,p}^j$, then c is either a (s, p) -Misiurewicz parameter or a p -super-attracting parameter. So we divide the set $D_{s,p}^j$ into two sets

$$D_{s,p}^j(\text{Misiurewicz}) = \{(c, 0) \in D_{s,p}^j \mid c \text{ is a Misiurewicz parameter}\}$$

and

$$D_{s,p}^j(\text{period}) = \{(c, 0) \in D_{s,p}^j \mid c \text{ is a super-attracting parameter}\}$$

By the definition of $C_{n,p}^j$ (Misiurewicz), we have

$$C_{n,p}^j(\text{Misiurewicz}) = \bigcup_{s=2}^n (h_{n,s,p}^j)^{-1}(D_{s,p}^j(\text{Misiurewicz})),$$

where $h_{n,s,p}^j := \wp_{s+1,p}^j \circ \cdots \circ \wp_{n,p}^j : \mathcal{V}_{n,p}^j \longrightarrow \mathcal{V}_{s,p}^j$

Fix any $s \in [2, n]$. Since the degree of $q_{s,p}^j(c, 0)$ is $\nu_d(p)d^{s-2}$ (Lemma 4.7) and $[\partial q_{s,p}^j / \partial c](c, 0) \neq 0$ at each $(c, 0) \in D_{s,p}^j$ (this point is shown in the proof of smoothness of $q_{s,p}^j(c, z)$), we get $\#D_{s,p}^j = \nu_d(p)d^{s-2}$. By point (5) of Proposition 3.3, the set $D_{s,p}^j(\text{period})$ is non-empty if and only if $p \mid s - 1$. In this case, we also see that $D_{s,p}^j(\text{period}) = \{(c, 0) \mid Q_{0,p}(c, 0) = 0\}$, then $\#D_{s,p}^j(\text{period})$ equals to $\nu_d(p)/d$ if $p \mid s - 1$ and 0 otherwise. It follows that

$$\#D_{s,p}^j(\text{Misiurewicz}) = \begin{cases} \nu_d(p)d^{s-2}, & \text{if } p \nmid s - 1; \\ \nu_d(p)d^{s-2} - \nu_d(p)/d, & \text{if } p \mid s - 1. \end{cases}$$

Note that the critical value set of $h_{n,s,p}^j$ is disjoint from $D_{s,p}^j(\text{Misiurewicz})$, so

$$\#(h_{n,s,p}^j)^{-1}(D_{s,p}^j(\text{Misiurewicz})) = \#D_{s,p}^j(\text{Misiurewicz}) \cdot d^{n-s}$$

and each point in $(h_{n,s,p}^j)^{-1}(D_{s,p}^j(\text{Misiurewicz}))$ is a critical point of $\widehat{\pi}_{n,p}^j$ with multiplicity $d - 1$. Therefore the number of critical points counting with multiplicity of $\widehat{\pi}_{n,p}^j$ in $C_{n,p}^j(\text{Misiurewicz})$, denoted by $M_{n,p}$, is equal to

$$\begin{aligned} M_{n,p} &: = \sum_{s=2}^n \#D_{s,p}^j(\text{Misiurewicz}) \cdot d^{n-s} \cdot (d - 1) \\ &= \nu_d(p)d^{n-2}(d - 1) \left(n - 1 - \sum_{t=1}^{\lfloor \frac{n-1}{p} \rfloor} d^{-tp} \right). \end{aligned} \quad (5.3)$$

- Counting the points of $C_{n,p}^{\text{crit}}$ (singular).

Recall that $C_{n,p}^{\text{crit}}$ (singular) consists of points of the form (c, z) with $f_c^{n-1}(z) = 0$ and such that there exists l between 0 and $n - 2$ (both included) with $f^l(z) = 0$ and s.t. 0 is p -periodic.

We divide the set $C_{n,p}(\text{singular})$ into several subsets $C_{n,p}^t(\text{singular})$ which consists of points $(c, z) \in C_{n,p}(\text{singular})$ such that

$$n - 1 - tp = \min\{ l \mid f_c^l(z) = 0 \}$$

The index t can take the values $0, \dots, [\frac{n-1}{p}]$, where $[x]$ denotes the maximal integer less than or equal to x , and the sets $C_{n,p}^t$ are pairwise disjoint and form a partition of $C_{n,p}(\text{singular})$. From (5.2), we see that $C_{n,p}^{\text{crit}}(\text{singular})$ is the union of $C_{n,p}^t(\text{singular}), t \geq 1$. Then we get $\#C_{n,p}^{\text{crit}}(\text{singular}) = 0$ if $n - 1 < p$. So in the following discussion, we only concern the case of $n - 1 \geq p$, i.e., $[\frac{n-1}{p}] \geq 1$.

Let $t \geq 1$. A point $(c, z) \in C_{n,p}^t(\text{singular})$ if and only if $(c, 0) \in D_{tp+1,p}^j(\text{period})$, $f_c^{n-1-tp}(z) = 0$ and $(f_c^{n-1-tp})'(z) \neq 0$. Hence

$$C_{n,p}^t(\text{singular}) = (h_{n,tp+1,p}^j)^{-1}(D_{tp+1,p}^j(\text{period})) \setminus (h_{n,(t+1)p+1,p}^j)^{-1}(D_{(t+1)p+1,p}^j(\text{period}))$$

if $(t+1)p+1 \leq n$, and $C_{n,p}^t(\text{singular}) = (h_{n,tp+1,p}^j)^{-1}(D_{tp+1,p}^j(\text{period}))$ otherwise. So we have

$$\#C_{n,p}^t(\text{singular}) = \begin{cases} d^{n-1-tp} \cdot \nu_d(p)/d, & \text{if } t = [\frac{n-1}{p}]; \\ d^{n-1-tp} \cdot \nu_d(p)/d - d \cdot d^{n-1-(t+1)p} \cdot \nu_d(p)/d, & \text{if } 1 \leq t < [\frac{n-1}{p}]. \end{cases}$$

On the other hand, the map $h_{n,tp+1,p}^j : \mathcal{V}_{n,p}^j \rightarrow \mathcal{V}_{tp+1,p}^j$ is injective in a neighborhood of any point $(c, z) \in C_{n,p}^t(\text{singular})$, and the map $\pi_{kp+1,p}^j : \mathcal{V}_{kp+1,p}^j \rightarrow \mathbb{C}$ has the local degree d^t at the point $(c, 0)$, so the number of critical points counting with multiplicity in $C_{n,p}^t(\text{singular})$ is $(d^t - 1)\#C_{n,p}^t(\text{singular})$. Then the total number of critical points counting with multiplicity in $C_{n,p}(\text{singular})$, in the case of $[\frac{n-1}{p}] \geq 1$, is

$$\begin{aligned} K_{n,p} &:= \sum_{t=1}^{[\frac{n-1}{p}]} (d^t - 1)\#C_{n,p}^t(\text{singular}) \\ &= \nu_d(p)(d^{p-1} - 1)d^{n-1-p}(\xi_{n,p} - \zeta_{n,p}) + (d^{[\frac{n-1}{p}]} - 1)\nu_d(p)d^{n-2-[\frac{n-1}{p}]p} \end{aligned} \quad (5.4)$$

where $\xi_{n,p} := \sum_{t=1}^{[\frac{n-1}{p}]-1} d^{-t(p-1)}$ and $\zeta_{n,p} := \sum_{t=1}^{[\frac{n-1}{p}]-1} d^{-pt}$.

Note that when $[\frac{n-1}{p}] = 0$, the number computed by formula (5.4) is 0, which is still equal to the number of $C_{n,p}^{\text{crit}}(\text{singular})$. So the number $K_{n,p}$, defined by (5.4), is equal to the number of critical points counting with multiplicity in $C_{n,p}^{\text{crit}}(\text{singular})$ in all cases.

- Counting the ideal boundary points.

In [B], [Mil1], Bousch and Milnor show that the local degree of $\pi_{0,p}$ at each ideal boundary point is 2 (in the case of $d = 2$) by analysing the asymptotic behavior of $f_c(z)$ as (c, z) goes to an ideal boundary point on $\mathcal{X}_{0,p}$. Their argument can be easily generalized to our case with degree $d \geq 2$. Just to be self-contained we give an alternative proof using the monodromy action (Lemma 5.3 below). By Lemma 5.3,

the local degree of $\widehat{\mathcal{V}}_{n,p}^j$ at each ideal boundary point is d . It follows that the number of ideal boundary points is $\nu_d(p)d^{n-2}$ because $\widehat{\pi}_{n,p}^j$ is a degree $\nu_d(p)d^{p-1}$ branched covering. So the number of critical points counting with multiplicity is equal to $\nu_d(p)d^{n-2}(d-1)$.

By the Riemann-Hurwitz formula, we have

$$g_{n,p}^j = 1 + \frac{1}{2}\nu_d(p)d^{n-2}(pd - p - 1 - d) + \frac{1}{2}(M_{n,p} + K_{n,p}) - \frac{1}{2}d^{n-2}(d-1) \sum_{k|p, k < p} k\nu_d(k)\varphi(p/k).$$

Here is a genus computation of some examples.

d	n	p	$\nu_d(p)$	$M_{n,p}$	$K_{n,p}$	$g_{n,p}$
3	1	1	3	0	0	0
3	2	1	3	4	2	1
2	2	2	2	2	0	0
2	3	2	2	7	1	1
2	2	3	6	6	0	2

Corollary 5.2. *Fix $n, p \geq 1$, the surfaces $\mathcal{V}_{n,p}^j$, $1 \leq j \leq d-1$ are pairwise homeomorphic.*

Proof. Topologically, the surface $\mathcal{V}_{n,p}^j$ is completely determined by the genus and the number of ideal boundary points of $\widehat{\mathcal{V}}_{n,p}^j$, whereas these two numbers are independent of j . \square

Lemma 5.3. *All ideal boundary points are critical points of $\widehat{\pi}_{n,p}^j$ with multiplicity $d-1$.*

Proof. To prove this lemma, we first give a symbolic description of the dynamics on the filled-in Julia set for a parameter outside the Multibrot set.

If $c \in \mathbb{C} \setminus M_d$, the Julia set of f_c is a Cantor set. If $c \in R_{M_d}(\theta)$ with $\theta \neq 0$ not necessarily periodic, the dynamical rays $R_c(\theta/d) \dots R_c((\theta + d - 1)/d)$ bifurcate on the critical point. The set $R_c(\theta/d) \cup \dots \cup R_c((\theta + d - 1)/d) \cup \{0\}$ decomposes the complex plane into d connected components. We denote by U_0 the component containing the dynamical ray $R_c(0)$ and by U_1, \dots, U_{d-1} the other components in counterclockwise order.

The orbit of a point $x \in K_c$ has an itinerary with respect to this partition. In other words, to each $x \in K_c$, we can associate a sequence in $\mathbb{Z}_d^{\mathbb{N}}$ whose j -th entry is equal to k if $f_c^{o_{j-1}}(x) \in U_k$. This gives a map $\iota_c : K_c \rightarrow \mathbb{Z}_d^{\mathbb{N}}$, which is bijective for any $c \in \mathbb{C} \setminus M_d$. Moreover, the dynamic of f_c on K_c is conjugate to shift on $\mathbb{Z}_d^{\mathbb{N}}$ via the map ι_c .

Now let $\pi := \pi_{n,p}^j|_{\mathcal{E}_{n,p,i}^j}$. The map $\pi : \mathcal{E}_{n,p,i}^j \rightarrow \mathbb{C} \setminus M_d$ is a covering of degree $d_{n,p,i}^j$. Fix $c_0 \in \mathbb{C} \setminus (M_d \cup R_{M_d}(0))$, $d_{n,p,i}^j = \#(\pi^{-1}(c_0))$. Since $\mathcal{E}_{n,p,i}^j$ is connected, the monodromy group induced by π , denoted by $\text{Mon}(\pi)$, acts on $\pi^{-1}(c_0)$ transitively. Then fixing any point $(c_0, z_0) \in \pi^{-1}(c_0)$, the set $\pi^{-1}(c_0)$ is exactly the orbit of (c_0, z_0) under $\text{Mon}(\pi)$.

Let $c_t : [0, 1] \rightarrow \mathbb{C} \setminus M_d$ be an oriented simple closed curve based at c_0 such that c_t intersects $R_{M_d}(0)$ at only one point c_{t_0} . Let z_t be the (n, p) -preperiodic point of f_{c_t}

obtained from the analytic continuation of z_0 along c_t . Note that as c varies in $\mathbb{C} \setminus (M_d \cup R_{M_d}(0))$, the (n, p) -preperiodic points of f_c , the dynamical rays $R_c(0)$ and $R_c((\theta_c + s)/d)$ ($s \in \mathbb{Z}_d$) move continuously. Consequently, we have

$$\begin{cases} \iota_{c_t}(z_t) = \iota_{c_0}(z_0) & \text{for } t \in [0, t_0) \\ \iota_{c_t}(z_t) = \iota_{c_0}(z_1) & \text{for } t \in (t_0, 1] \end{cases}$$

Furthermore, on one hand, z_t and $R_{c_t}(0)$ move continuously for $t \in [0, 1]$. On the other hand, when c_t passes through $R_{M_d}(0)$, the dynamical rays $R_{c_t}((\theta_t + s)/d)$ ($s \in \mathbb{Z}_d$) move discontinuously and jump from $R_{c_{t_-}}((\theta_{t_-} + s)/d)$ to $R_{c_{t_+}}((\theta_{t_+} + s + 1)/d)$, $t_- < t_0 < t_+$. So if $\iota_{c_0}(z_0) = \beta_n \dots \beta_1 \overline{\epsilon_1 \dots \epsilon_p}$, then

$$\iota_{c_0}(z_1) = (\beta_n + 1) \dots (\beta_1 + 1) \overline{(\epsilon_1 + 1) \dots (\epsilon_p + 1)} \quad (5.5)$$

Hence the map σ_{c_t} , an element of $\text{Mon}(\pi)$ induced by c_t , maps (c_0, z_0) to (c_0, z_1) with z_1 satisfying (5.5). Since $\pi_1(\mathbb{C} \setminus M_d, c_0) = \langle c_t \rangle$, then we have

$$\left(\pi_{n,p}^j \Big|_{\mathcal{E}_{n,p,i}^j} \right)^{-1} (c_0) = \{ (c_0, z) \mid \iota_{c_0}(z) = (\beta_n + s) \dots (\beta_1 + s) \overline{(\epsilon_1 + s) \dots (\epsilon_p + s)}, s \in \mathbb{Z}_d \}$$

As a consequence, $d_{n,p,i}^j = d$. □

6 The Galois group of $Q_{n,p}(c, z)$

The objective here is to study $\mathcal{X}_{n,p}$ from the algebraic point of view by calculating its associated Galois group.

Recall that \mathbf{C} denotes the ring $\mathbb{C}[c]$ and \mathbf{K} is a fixed algebraically closed field containing \mathbf{C} . Since the characteristic of $\mathbf{C}(c)$ is 0, any polynomial $f \in \mathbf{C}[z]$ induces a finite Galois extension $\mathbf{C}(c)(f)$ over $\mathbf{C}(c)$ (see [W, Thm. 3.2.6, 2.7.14]), where $\mathbf{C}(c)(f)$ is the splitting field of f , and hence a Galois group $G(f) := \text{Gal}(\mathbf{C}(c)(f)/\mathbf{C}(c))$. In particular, we denote the Galois group of $Q_{n,p}$ by $G_{n,p}$.

For each $n \geq 0, p \geq 1$, denote $\mathfrak{R}_{n,p}$ the set of roots of $Q_{n,p} \in \mathbf{C}[z]$. By (3.2), we have $f_c(\mathfrak{R}_{n,p}) = \mathfrak{R}_{n-1,p}$ if $n \geq 1$ and $f_c(\mathfrak{R}_{0,p}) = \mathfrak{R}_{0,p}$. Let us consider

$$\mathfrak{R}_{\leq n,p} := \bigcup_{0 \leq l \leq n} \mathfrak{R}_{l,p}.$$

Then $f_c(\mathfrak{R}_{\leq n,p}) \subset \mathfrak{R}_{\leq n,p}$ and the action of f_c induces a directed graph structure consisting of a certain number of disjoint cycles of order p , on each vertex of which is attached a tree of height n . More precisely, for each $0 \leq l \leq n$, we consider the roots in $\mathfrak{R}_{l,p}$ as the vertices of level l , and two vertices $\Delta_1, \Delta_2 \in \mathfrak{R}_{\leq n,p}$ are connected by an oriented edge from Δ_1 to Δ_2 if $f_c(\Delta_1) = \Delta_2$. Thus $\mathfrak{R}_{\leq n,p}$ has a graph structure, and we denote this graph by $\mathfrak{R}_{\leq n,p}^T$ (see Figure 1).

Example. For $d = 3, p = 4, n = 2$, the directed graph $\mathfrak{R}_{\leq 2,4}^T$ has 18 connected component, which are pairwise isomorphic. We draw one in the following.




Figure 1. A connected component of $\mathfrak{R}_{\leq 2,4}^T$.

On the other hand, note that $\mathfrak{R}_{\leq n,p}$ is the set of roots of

$$Q_{\leq n,p} := \prod_{l=0}^n Q_{l,p} \in \mathbf{C}[z].$$

So, correspondingly, we consider the Galois group $G_{\leq n,p}$ of $Q_{\leq n,p}$. Firstly, we have the following simple result.

Proposition 6.1. *For each $n \geq 0, p \geq 1$, we have $G_{n,p} = G_{\leq n,p}$.*

Proof. By (3.2), any root of $Q_{l,p} \in \mathbf{C}[z]$ ($0 \leq l \leq n$) can be written as a polynomial with coefficients in \mathbf{C} of roots of $Q_{n,p}$. It follows that the splitting field of $Q_{\leq n,p} = \prod_{l=0}^n Q_{l,p}$ over $\mathbf{C}(c)$ is the same as that of $Q_{n,p}$ over $\mathbf{C}(c)$. Hence $G_{\leq n,p} = G_{n,p}$. \square

By this proposition, computing the Galois group $G_{n,p}$ is equivalent to computing the group $G_{\leq n,p}$. Let σ be an element in $G_{\leq n,p}$. Since it fixes the base field $\mathbf{C}(c)$ pointwise, we have $\sigma(\mathfrak{R}_{l,p}) = \mathfrak{R}_{l,p}$ and $f_c \circ \sigma = \sigma \circ f_c$. Hence σ induces an *automorphism* of the graph $\mathfrak{R}_{\leq n,p}^T$, i.e., σ is a permutation on each l -level vertices of $\mathfrak{R}_{\leq n,p}^T$, and $\Delta_1, \Delta_2 \in \mathfrak{R}_{\leq n,p}$ are connected by an edge from Δ_1 to Δ_2 if and only if $\sigma(\Delta_1), \sigma(\Delta_2)$ are connected by an edge from $\sigma(\Delta_1)$ to $\sigma(\Delta_2)$. Clearly, different elements of $G_{\leq n,p}$ induce different automorphisms of $\mathfrak{R}_{\leq n,p}^T$. So $G_{\leq n,p}$ can be seen as a subgroup of $\text{Aut}(\mathfrak{R}_{\leq n,p}^T)$, the automorphic group of the graph $\mathfrak{R}_{\leq n,p}^T$.

In the case $d = 2$, Bousch [B] proved that

$$G_{\leq n,p} \simeq \text{Aut}(\mathfrak{R}_{\leq n,p}^T) \simeq H_{\leq n,p}(f_c),$$

where $H_{\leq n,p}(f_c)$ denotes the set of all permutations on $\mathfrak{R}_{\leq n,p}$ that commute with f_c . In the general case, the result is similar but needs a small modification. We will exhibit this point in the following.

Let σ be an element of the Galois group $G_{\leq n,p}$. As σ fixes the field $\mathbf{C}(c)$ pointwise, it must satisfy the following two conditions:

(P1) σ commutes with f_c , i.e., $\sigma \circ f_c = f_c \circ \sigma$.

(P2) σ commutes with the rotation of argument $1/d$. That is, if $\sigma(\Delta) = \tilde{\Delta}$ for $\Delta, \tilde{\Delta} \in \mathfrak{R}_{\leq n, p}$, then $\sigma(\omega^j \Delta) = \omega^j \tilde{\Delta}$, where $\omega = e^{\frac{2\pi i}{d}}$ and $1 \leq j \leq d-1$

Therefore, if a permutation on $\mathfrak{R}_{\leq n, p}$ wants to be a candidate of elements in the Galois group $G_{\leq n, p}$, it should satisfy the conditions (P1) and (P2).

In fact, in the case of $d = 2$, the condition (P1) implies (P2). To see this, let Δ_{n-1} be a root of $Q_{n-1, p}$ ($n \geq 1$) and $\Delta_n, -\Delta_n$ be the preimages of Δ_{n-1} under f_c . Let σ be an element of $G_{\leq n, p}$ and set $\tilde{\Delta}_n := \sigma(\Delta_n)$. By condition (P1) σ must map $-\Delta_n$ to $-\tilde{\Delta}_n$, then the condition (P2) holds. Therefore, it is possible for (P1) to be a sufficient condition for a permutation on $\mathfrak{R}_{\leq n, p}$ to be an element of $G_{\leq n, p}$, and Bousch [B] proved this point.

However, the situation has a little difference in the case of $d \geq 3$. Following the notations $\Delta_{n-1}, \Delta_n, \tilde{\Delta}_n$ and σ as above. In this case, Δ_{n-1} has at least 3 preimages, which are $\Delta_n, \omega \Delta_n, \dots, \omega^{d-1} \Delta_n$. By condition (P1), we only know that σ maps $\{\omega \Delta_n, \dots, \omega^{d-1} \Delta_n\}$ bijectively to $\{\omega \tilde{\Delta}_n, \dots, \omega^{d-1} \tilde{\Delta}_n\}$, but can not get $\sigma(\omega^j \Delta_n) = \omega^j (\tilde{\Delta}_n)$ for $1 \leq j \leq d-1$. So, in case of $d \geq 3$, the condition (P2) can not be omitted.

What we would like to prove is that, except (P1) and (P2), no other restrictions are imposed on $G_{\leq n, p}$. The proof is similar to that of Theorem 4 in Chapter III of [B].

Theorem 6.2. *The Galois group $G_{\leq n, p}$ consists of all permutations on $\mathfrak{R}_{\leq n, p}$ which commute with f_c and the rotation of argument $1/d$.*

Proof. We denote r_d the rotation of argument $1/d$, and $H_{\leq n, p}(f_c, r_d)$ the set of permutations on $\mathfrak{R}_{\leq n, p}$ which commute with f_c and r_d . By the definition, it is not difficult to check that $H_{\leq n, p}(f_c, r_d)$ leaves each $\mathfrak{R}_{l, p}$, and hence $\mathfrak{R}_{\leq l, p}$ invariant for $0 \leq l \leq n$.

Define a group homomorphism

$$\phi_n : G_{\leq n, p} \rightarrow H_{\leq n, p}(f_c, r_d)$$

such that $\phi_n(\sigma)$ is the restriction of σ to $\mathfrak{R}_{\leq n, p}$. According to the discussion above, we just need to prove the surjectivity of ϕ_n .

Note first that the result is true for $n = 0$ following 6 of Lemma 2.4.

For $n = 1$, since $H_{\leq 1, p}(f_c, r_d)$ leaves $\mathfrak{R}_{0, p}$ invariant, there is a natural homomorphism from $H_{\leq 1, p}(f_c, r_d)$ to $H_{\leq 0, p}(f_c, r_d)$ with $\tilde{\tau} \mapsto \tilde{\tau}|_{\mathfrak{R}_{0, p}}$. This homomorphism has an inversion which maps $\tau \in H_{\leq 0, p}(f_c, r_d)$ to $\tilde{\tau} \in H_{\leq 1, p}(f_c, r_d)$ such that $\tilde{\tau}|_{\mathfrak{R}_{0, p}} = \tau$ and $\tilde{\tau}(\omega^j \Delta) = \omega^j \tau(\Delta)$ for each $\Delta \in \mathfrak{R}_{0, p}, j \in [1, d-1]$. Thus $H_{\leq 1, p}(f_c, r_d) \simeq H_{\leq 0, p}(f_c, r_d)$. Note that $G_{1, p} = G_{0, p}$ (because the splitting fields of $Q_{0, p}$ and $Q_{1, p}$ over $\mathbb{C}(c)$ coincide), then the result is true for $n = 1$.

Now we argue by induction on n . Assume that $\phi_{n-1} : G_{\leq n-1, p} \rightarrow H_{\leq n-1, p}(f_c, r_d)$ is surjective ($n \geq 2$).

Let $\tau \in H_{\leq n, p}(f_c, r_d)$. As τ commutes with f_c , it leaves $\mathfrak{R}_{\leq n-1, p}$ invariant. Then $\tau|_{n-1}$, the restriction of τ on $\mathfrak{R}_{\leq n-1, p}$, belongs to $H_{\leq n-1, p}(f_c, r_d)$. By the assumption

of induction, there is an element σ_{n-1} of $G_{\leq n-1,p}$ with $\phi_{n-1}(\sigma_{n-1}) = \tau|_{n-1}$. According to the Galois theory (see [W, Thm. 2.88]), we can find an element σ of $G_{\leq n,p}$ such that its restriction on the splitting field of $Q_{\leq n-1,p}$ over $\mathbb{C}(c)$ coincides with σ_{n-1} . Set $\tau' := \tau \cdot \phi_n(\sigma)^{-1}$, then $\tau' \in H_{\leq n,p}(f_c, r_d)$ and it fixes $\mathfrak{R}_{\leq n-1,p}$ pointwise. Now it remains to prove that $G_{\leq n,p}$ contains τ' , i.e., there exists $\sigma' \in G_{\leq n,p}$ with $\phi_n(\sigma') = \tau'$, because if so, $\tau = \phi_n(\sigma')\phi_n(\sigma) = \phi_n(\sigma'\sigma)$, which implies ϕ_n is surjective.

Set $\kappa_l := \nu_d(p)(d-1)d^{l-1}$ for each $l \geq 1$ (which is the number of roots of $Q_{l,p}$), and denote

$$\mathfrak{R}_{n,p} = \{\Delta_n^i, \omega\Delta_n^i, \dots, \omega^{d-1}\Delta_n^i\}_{i=1}^{\kappa_{n-1}}$$

Since τ' fixes $\mathfrak{R}_{\leq n-1,p}$ pointwise and commutes with both f_c, r_d , it can be expressed as a product

$$\tau' = \prod_{i=1}^{\kappa_{n-1}} (s_i, s_i + 1, \dots, d-1, 0, \dots, s_i - 1),$$

where $(s_i, s_i + 1, \dots, d-1, 0, \dots, s_i - 1)$ is the cyclic permutation on $(\Delta_n^i, \dots, \omega^{d-1}\Delta_n^i)$ mapping Δ_n^i to $\omega^{s_i}\Delta_n^i$ and so on. Notice that all these cyclic permutations $(s_i, \dots, s_i - 1)$ are commutable.

The argument in this paragraph is a classical correspondence between the Galois theory and the covering theory; see [Z] for the detail. Let $V_{n,p}$ be the set of singular values of the projection $\pi : \mathcal{X}_{n,p} \rightarrow \mathbb{C}$. Then $\pi_{n,p}$ restricts to a cover from the complement of the preimage of $V_{n,p}$ in $\mathcal{X}_{n,p}$ to the complement of $V_{n,p}$ in \mathbb{C} . For all c_0 not in $V_{n,p}$, there is thus an action of $\pi_1(\mathbb{C} \setminus V_{n,p}, c_0)$ on the roots

$$Z_{n,p} = \{z_n^i, \dots, \omega^{d-1}z_n^i\}_{i=1}^{\kappa_{n-1}}$$

of $Q_{n,p}(c_0, z)$, which is seen as a polynomial in the variable z and with complex coefficients. By the correspondence between the Galois theory and the covering theory (see [Z, Thm. 1]), there is a (non-unique) choice of bijection between the roots of $Q_{n,p} \in \mathbb{C}[z]$ and the roots of $Q_{n,p}(c_0, z) \in \mathbb{C}[z]$ such that the set of permutations induced by the Galois group on the set $\mathfrak{R}_{\leq n,p}$ is conjugated by this bijection to the set of permutations on $Z_{n,p}$ induced by $\pi_1(\mathbb{C} \setminus V_{n,p}, c_0)$. Thus we get a surjective (not injective, usually) morphism from $\pi_1(\mathbb{C} \setminus V_{n,p}, c_0)$ to the Galois group. Moreover, this bijection is such that any polynomial relation between the Δ_n^i with coefficient in $\mathbb{C}(c)$ will give a relation between the z_n^i , with c_0 being substituted to c . It implies that the action of $\pi_1(\mathbb{C} \setminus V_{n,p}, c_0)$ on $Z_{n,p}$ preserves commutation with multiplication by ω .

Therefore, by the discussion above, to obtain the required permutation τ' , we only need to find a path in the basic group $\pi_1(\mathbb{C} \setminus V_{n,p}, c_0)$ such that its monodromy action on $\{(z_n^i, \dots, \omega^{d-1}z_n^i)\}_{i=1}^{\kappa_{n-1}}$ induces the same permutation as τ' . We now show the following result, which is sufficient: for any $i \in [1, \kappa_{n-1}]$, there exists a path in $\pi_1(\mathbb{C} \setminus V_{n,p}, c_0)$ such that its monodromy action induces the permutation $(s_i, s_i + 1, \dots, s_i - 1)$.

Fix any $i \in [1, \kappa_{n-1}]$. Suppose that $\{(c_0, z_n^i), (c_0, \omega z_n^i), \dots, (c_0, \omega^{d-1}z_n^i)\}$ belong to $\mathcal{V}_{n,p}^t$. Let \hat{c} be an (n, p) -Misiurewicz parameter with $(\hat{c}, 0) \in \mathcal{V}_{n,p}^t$. Such \hat{c} exists because the set $D_{n,p}^t(\text{Misiurewicz})$ is non-empty (see section 5.2 item 2). By (3.2), $(\hat{c}, \hat{c}) \in \mathcal{X}_{n-1,p}$.

Since \hat{c} is a Misiurewicz parameter and the orbit of \hat{c} does not contain 0, then the point (\hat{c}, \hat{c}) belongs to no sets in Lemma 5.1 in the case $l = n - 1$. Hence $w = \hat{c}$ is a simple root of the equation $Q_{n-1,p}(\hat{c}, w) = 0$ (in w). So by the Implicit Function Theorem, the equation $Q_{n-1,p}(c, w) = 0$ admits a unique solution $w = w(c)$ close to \hat{c} fulfilling $w(\hat{c}) = \hat{c}$. Thus, a neighborhood of $(\hat{c}, 0)$ in $\mathcal{X}_{n,p}$ can be written as

$$\{(c, z_c) \cup (c, \omega z_c) \cup \dots \cup (c, \omega^{d-1} z_c) \mid |c - \hat{c}| < \epsilon\}$$

where z_c is one of the preimages of $w(c)$ under f_c nearby 0.

When c makes a small turn around \hat{c} , the set $\{z_c, \dots, \omega^{d-1} z_c\}$ gets a cyclic permutation with $\omega^j z_c$ mapped to $\omega^{j+1} z_c$, because $\pi_{n,p}$ is a degree d covering in a punctured neighborhood of $(\hat{c}, 0)$ (which is shown in §5.2 item 2), and the other (n, p) -preperiodic points of f_c remain fixed, since $\pi_{n,p}$ is injective around each point (\hat{c}, ξ) with ξ a non-zero (n, p) -preperiodic point of $f_{\hat{c}}$. So if we choose a path $\gamma \in \pi_1(\mathbb{C} \setminus V_{n,p}, c_0)$ homotopic to \hat{c} , the induced permutation by its monodromy action gives the cyclic permutation $(2, \dots, d, 1)$ on $(z_n^*, \dots, \omega^{d-1} z_n^*)$ for a (n, p) -preperiodic point z_n^* of f_{c_0} fulfilling $(c_0, z_n^*) \in \mathcal{V}_{n,p}^t$, and keeps the other (n, p) -preperiodic points of f_{c_0} fixed. Now we join (c_0, z_n^i) and (c_0, z_n^*) by a curve from (c_0, z_n^i) to (c_0, z_n^*) in $\mathcal{V}_{n,p}^t \setminus \pi_{n,p}^{-1}(V_{n,p})$, and denote its projection under $\pi_{n,p}$ by β . Then $\beta \in \pi_1(\mathbb{C} \setminus V_{n,p}, c_0)$ and the path $\beta \cdot \gamma^{s_i} \cdot \beta^{-1}$ is what we expect. \square

Applying this theorem, we can also characterize the Galois group $G_{\leq n,p}$ by the automorphisms of the directed graph $\mathfrak{R}_{\leq n,p}^T$, as in the $d = 2$ case. For $d \geq 3$, denote by $\text{Aut}(\mathfrak{R}_{\leq n,p}^T, r_d)$ the set of automorphisms of $\mathfrak{R}_{\leq n,p}^T$ that commute with the rotation of argument $1/d$, and by $H_{\leq n,p}(f_c, r_d)$ the set of permutations on $\mathfrak{R}_{\leq n,p}$ that commute with f_c and the rotation of argument $1/d$.

Corollary 6.3. *For $n \geq 0, p \geq 1$, $G_{\leq n,p} \simeq \text{Aut}(\mathfrak{R}_{n,p}^T, r_d) \simeq H_{\leq n,p}(f_c, r_d)$.*

Following Bousch [B, Chap. 3, III] and Silverman ([Sil, § 3.9]), we express the Galois group $G_{n,p}$ ($n \geq 2$) as a wreath product.

Definition 6.4. *Let G be a group and Σ be a subgroup of \mathbf{S}_m , where \mathbf{S}_m denotes the set of permutations on $\{1, \dots, m\}$. Denote by $\Sigma \ltimes G^m$ the wreath product of G and Σ . As a set, it consists of $g = \sigma(g_1, \dots, g_m)$ where $g_i \in G$ and $\sigma \in \Sigma$. The multiplication is defined by*

$$g \cdot h = \sigma_g(g_1, \dots, g_m) \cdot \sigma_h(h_1, \dots, h_m) = \sigma_g \circ \sigma_h(g_{\sigma_h(1)} \cdot h_1, \dots, g_{\sigma_h(m)} \cdot h_m).$$

Under this multiplication law, $\Sigma \ltimes G^m$ is a group with $g^{-1} = \sigma_g^{-1}(g_{\sigma_g^{-1}(1)}^{-1}, \dots, g_{\sigma_g^{-1}(m)}^{-1})$ and unit element $(1, \dots, 1)$.

In [B], Bousch showed that the Galois group $G_{0,p}$ is isomorphic to the wreath product $\mathbf{S}_{\nu_d(p)/p} \ltimes (\mathbb{Z}/p\mathbb{Z})^{\nu_d(p)/p}$ (see also [Sil, § 3.9]). From the proof of Theorem 6.2, we have seen that $G_{1,p} = G_{0,p}$, so

$$G_{1,p} \simeq \mathbf{S}_{\nu_d(p)/p} \ltimes (\mathbb{Z}/p\mathbb{Z})^{\nu_d(p)/p}.$$

For $n \geq 2$, we can give inductively an isomorphic model of $G_{n,p}$ by a wreath product. Recall that $\kappa_n = \nu_d(p)(d-1)d^{n-1}$ ($n \geq 1$) is the number of roots of $Q_{n,p}$.

Proposition 6.5. For $n \geq 2$, we have $G_{n,p} \cong G_{n-1,p} \times (\mathbb{Z}/d\mathbb{Z})^{\kappa_{n-1}}$, where the action of $G_{n-1,p}$ on $(1, 2, \dots, \kappa_{n-1})$ comes from the action of $G_{n-1,p}$ on the roots of $Q_{n-1,p}$, of which there are exactly κ_{n-1} .

Proof. For $n \geq 2$, we denote $(\Delta_{n-1}^i)_{i=1}^{\kappa_{n-1}}$ the roots of $Q_{n-1,p} \in \mathbb{C}[z]$, and denote

$$\left(\{ \Delta_n^i, \dots, \omega^{d-1} \Delta_n^i \} \right)_{i=1}^{\kappa_{n-1}}$$

the roots of $Q_{n,p}$ such that $f_c(\Delta_n^i) = \Delta_{n-1}^i$.

$$\begin{array}{c|c|c|c} \Delta_{n-1}^1 & \Delta_{n-1}^2 & \dots & \Delta_{n-1}^{\kappa_{n-1}} \\ \hline \Delta_n^1 & \Delta_n^2 & \dots & \Delta_n^{\kappa_{n-1}} \\ \omega \Delta_n^1 & \omega \Delta_n^2 & \dots & \omega \Delta_n^{\kappa_{n-1}} \\ \vdots & \vdots & & \vdots \\ \omega^{d-1} \Delta_n^1 & \omega^{d-1} \Delta_n^2 & \dots & \omega^{d-1} \Delta_n^{\kappa_{n-1}} \end{array}$$

We define a group homomorphism

$$W : G_{n,p} \longrightarrow G_{n-1,p} \times (\mathbb{Z}/d\mathbb{Z})^{\kappa_{n-1}}$$

such that $W(\sigma) = \sigma|_{n-1}(s_1, \dots, s_{\kappa_{n-1}})$, where $\sigma|_{n-1}$ is the restriction of σ on the splitting field of $Q_{n-1,p}$ over $\mathbb{C}(c)$, and the i -th digit in $(s_1, \dots, s_{\kappa_{n-1}})$ is s_i if and only if once $\sigma(\Delta_{n-1}^i) = \Delta_{n-1}^t$ for some $1 \leq t \leq \kappa_{n-1}$, then $\sigma(\Delta_n^i) = \omega^{s_i} \Delta_n^t$. The injectivity of W is straightforward by the action of $G_{n,p}$ on $\mathfrak{R}_{\leq n,p}$ and the surjectivity of W is due to Theorem 6.2. \square

Before ending this section, we give a computation of $G_{n,p}$ for some small n, p . Note that although $G_{1,p}$ is isomorphic to a subgroup $\mathbf{S}_{\nu_d(p)/p} \times (\mathbb{Z}/p\mathbb{Z})^{\nu_d(p)/p}$ of $\mathbf{S}_{\nu_d(p)}$, it is indeed a subgroup of $\mathbf{S}_{\nu_d(p)(d-1)}$. So mimicking the action of $G_{1,p}$ on

$$\{ \omega \Delta_1^1, \dots, \omega \Delta_1^{\nu_d(p)}; \dots; \omega^{d-1} \Delta_1^1, \dots, \omega^{d-1} \Delta_1^{\nu_d(p)} \},$$

we define a subgroup $\mathbf{P}_{\nu_d(p)(d-1),d}$ of $\mathbf{S}_{\nu_d(p)(d-1)}$ such that $\tau \in \mathbf{P}_{\nu_d(p)(d-1),d}$ if and only if

$$\begin{aligned} & \tau(1, \dots, \nu_d(p); \dots; (d-2)\nu_d(p) + 1, \dots, (d-2)\nu_d(p) + \nu_d(p)) \\ = & (\sigma(1), \dots, \sigma(\nu_d(p)); \dots; (d-2)\nu_d(p) + \sigma(1), \dots, (d-2)\nu_d(p) + \sigma(\nu_d(p))) \end{aligned}$$

for a $\sigma \in \mathbf{S}_{\nu_d(p)}$. Then $\mathbf{P}_{\nu_d(p)(d-1),d} \simeq \mathbf{S}_{\nu_d(p)} \simeq G_{1,p}$, and $\mathbf{P}_{\nu_d(p)(d-1),d} = \mathbf{S}_{\nu_d(p)}$ in the case $d = 2$. The results of computation are listed in the following table.

d	n	p	$\nu_d(p)$	κ_{n-1}	$G_{n,p}(d)$
3	1	1	3	–	$\mathbf{S}_3 \simeq \mathbf{P}_{6,3}$
3	2	1	3	6	$\mathbf{P}_{6,3} \times (\mathbb{Z}/3\mathbb{Z})^6$
3	3	1	3	18	$(\mathbf{P}_{6,3} \times (\mathbb{Z}/3\mathbb{Z})^6) \times (\mathbb{Z}/3\mathbb{Z})^{18}$
2	3	2	2	4	$((\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})^2) \times (\mathbb{Z}/2\mathbb{Z})^4$
2	2	3	6	6	$((\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})^2) \times (\mathbb{Z}/2\mathbb{Z})^6$

References

- [B] T. Bousch, *Sur quelques problèmes de dynamique holomorphe*, Ph.D. thesis, Université de Paris-Sud, Orsay, 1992.
- [BT] X. Buff and L. Tan, *The quadratic dynatomic curves are smooth and irreducible*, *Frontiers in Complex Dynamics: In Celebration of John Milnor's 80th Birthday*, 2014, 49–72.
- [CG] L. Carleson and W. Gamelin, *Complex dynamics*, Springer-Verlag, 1993.
- [DH1] A. Douady, J. H. Hubbard, *Étude Dynamique Des Polynômes Complexes, I, II*, Orsay: Publ. Math. Orsay, 1984–1985.
- [DH2] A. Douady, J. H. Hubbard, *Exploring the Mandelbrot set: The Orsay Notes*, www.math.cornell.edu/~hubbard/OrsayEnglish.pdf
- [Eb] D. Eberlein, *Rational Parameter Rays of Multibrot Set*, Diploma Thesis, Technische Universität München, in preparation.
- [E] A. Epstein, *Infinitesimal Thurston rigidity and the Fatou-Shishikura inequality*, Stony Brook IMS Preprint 1999-1.
- [F] W. Fulton, *Algebraic curves. An introduction to algebraic geometry*, 3rd ed, 2008
- [G] P. A. Griffiths, *Introduction to algebraic curves*, *Translations of mathematical monographs*, Volume 76, AMS, 1989.
- [GO] Y. Gao and Y. F. Ou, *The dynatomic curves for polynomial $z \mapsto z^d + c$ are smooth and irreducible*, *SCIENCE CHINA Mathematics*, 2014, 57(6), 1175–1192
- [LS] E. Lau and D. Schleicher, *Internal addresses in the Mandelbrot set and irreducibility of polynomials*, *Stony Brook Preprint* **19**, 1994.
- [Le] G. Levin, *On explicit connections between dynamical and parameter spaces*, *Journal d'Analyse Mathématique*, 91 (2003), 297–327.
- [Mo] P. Morton, *On certain algebraic curves related to polynomial maps*, *Compositio Math*, 103 (1996), No. 3, 319-350.
- [Mil1] J. Milnor, *Periodic Orbits, External Rays and the Mandelbrot Set: An Expository Account*, *Astérisque* 261 (2000), 277–333.
- [Mil2] J. Milnor, *Dynamics in One Complex Variable*, Princeton University Press 2006.
- [S] D. Schleicher, *Internal addresses of the Mandelbrot set and Galois groups of polynomials*, arxiv:math/9411238v2, feb. 2008.
- [Sil] Silverman, *The Arithmetic of Dynamical Systems*, GTM 241, Springer, 2007.
- [W] S. H. Weintraub, *Galois Theory, 2nd edition (Universitext)*, Springer, 2006.

- [Z] H. Żołądek, *Note on the deck transformations group and the monodromy group*, Topological Methods In Nonlinear Analysis, Journal of the Juliusz Schauder Center, Volume 19, 2002, 237–256.